Attentional disengagement during external and internal distractions

reduces neural speech tracking in background noise

Yue Ren[1*], M. Eric Cui[1,2], Björn Herrmann[1,2,*]

[1] Rotman Research Institute, Baycrest Academy for Research and Education
North York, ON, Canada

[2] Department of Psychology, University of Toronto
Toronto, ON, Canada

*Correspondence concerning this article should be addressed to Yue Ren and Björn Herrmann,
Rotman Research Institute, Baycrest, 3560 Bathurst St, North York, ON, M6A 2E1, Canada. E-
mail: yren@research.baycrest.org and bherrmann@research.baycrest.org

Keywords: Electroencephalography (EEG), speech encoding, background noise, attention,
temporal response function

Conflict of interest statement: The author declares no competing interests.

Author contributions: YR: Methodology, data curation, formal analysis, visualization, writing -
original draft, writing - review and editing, project administration. **MEC**: Data curation, formal
analysis, writing - original draft, writing - review and editing, project administration. **BH**:
Conceptualization, methodology, formal analysis, writing - original draft, writing - review and
editing, supervision, funding acquisition.

Acknowledgements: We thank Tiffany Lao and Saba Junaid for their support with stimuli
creation and data collection. This study has been supported by BH's Canada Research Chair
program (CRC-2023-00383) and BH's Natural Sciences and Engineering Research Council of
Canada (Discovery Grant: RGPIN-2021-02602).

Word count: Abstract (240), Significance Statement (108), Introduction (650), Discussion (1498)

Abstract

Within-situation disengagement – the mental withdrawal during conversations in acoustically

challenging environments – is a common experience of older people with hearing difficulties. Yet,

most research on the neural mechanisms of attentional disengagement from speech listening has

focused on the distraction by one competing speaker, whereas within-situation disengagement is

often characterized by distraction towards external visual stimuli or internal thoughts and occurs

in   situations   with   ambient,   multi-talker   background   masking.   Across   three

electroencephalography (EEG) experiments in human participants of either sex, the current study

examined how disengagement due to external and internal distractions affect the neural tracking

of speech masked by different levels of multi-talker babble (speech in quiet, +6 dB, and -3 dB

SNR). We observed enhanced early neural responses (<0.2 s) to the speech envelope for speech

masked by background babble compared to speech in quiet (Experiments 1-3), suggesting

stochastic facilitation. Importantly, neural tracking of the speech envelope was reduced when

individuals were distracted by a visual-stimulus stream (Experiment 2) and by internal thought

and   imagination   (Experiment   3).   There   were   some   indices   suggesting   the   greatest

disengagement-related decline in neural speech tracking occurs for the most difficult speech-

masking condition, but this was not consistent across all measures. The current data show that

disengagement due to external and internal distractions yield decreases in neural speech

tracking, potentially suggesting converging neural pathways through which gain is downregulated

in auditory cortex. These results indicate that disengagement from listening can be identified

through non-invasive neural measures.

Significance

Many older adults with hearing difficulties mentally "tune out" during conversations in noisy
environments, yet the neural mechanisms underlying this within-situation disengagement remain
poorly understood. Across three electroencephalography experiments, we examined how
external (visual) and internal (thought-based) distractions influence neural tracking of speech
masked by multi-talker babble. We observed that attentional disengagement – whether induced
by external stimuli or internal thoughts – reduced the brain's tracking of the speech envelope.
These findings demonstrate that listening disengagement can be objectively identified through
neural measures and suggest a convergence in neural pathways through which both external and
internal distractions down-regulate auditory gain, providing new insight into attentional control in
challenging listening conditions.

Introduction

Conversations in cafés or restaurants often involves listeners extracting meaning from a speaker

amid background sounds. The cognitive resources required to comprehend under such conditions

makes listening effortful (Pichora-Fuller et al., 2016; Peelle, 2018). When the listening situation

becomes overly effortful, listeners may temporarily withdraw attention and "tune out" (Heffernan

et al., 2016; McNeice et al., 2024; Sarant et al., 2024). This within-situation disengagement is a

common response to sustained cognitive demand during listening (Heffernan et al., 2016;

Herrmann and Johnsrude, 2020) and may promote social withdrawal (Motala et al., 2024).

Characterizing markers of this disengagement could provide a window into how attention

fluctuates during listening and towards detecting disengagement in real-time.

Individuals may disengage from speech listening for multiple reasons. Externally, attention may

be captured by salient acoustic events, visual distractions, or competing speech inputs, disrupting

the processing of a target stimulus (Lavie, 2005; Shinn-Cunningham, 2008). Internally, attention

can drift away from the external environment as listeners engage in self-generated thought – often

characterized as mind wandering – during which cognitive resources are reallocated toward

memories, future plans, or imaginings (Eastwood et al., 2012; Ibaceta et al., 2024). This form of

internally directed disengagement becomes more frequent when individuals listen to speech in

background noise than when they listen in quiet (Steindorf et al., 2023). Thus, external distraction

and internal drift represent two pathways to the same functional outcome of temporarily

uncoupling attention from speech, but whether such disengagement leaves distinct neural

signatures and whether such signatures depend on listening demands is unclear.

One increasingly common approach to investigate the encoding of continuous speech in the brain

involves regressing relevant features of speech against signals recorded using electro- or

magnetoencephalography (EEG; MEG; Lalor and Foxe, 2010; Ding and Simon, 2012; Hertrich et

al., 2012). This approach, formalized as temporal response function (TRF) modeling (Crosse et

al., 2016), is often used to quantify how neural activity tracks the temporal dynamics of the speech

envelope, due to its importance for speech intelligibility (Ding et al., 2014; Lesenfants et al., 2019).

The neural tracking of the speech envelope can be decreased when speech is masked by

background sound (Ding and Simon, 2013; Synigal et al., 2023), although this seems to depend

on the specific TRF metric and type of background masker, sometimes even leading to noise-

related increases (Herrmann, 2025a, b).

Critically, a robust pattern across many studies is that the cortical tracking of the envelope of
attended speech is stronger than the tracking of the envelope of speech that is ignored, reflecting
selective amplification of behaviorally relevant neural signals (Kerlin et al., 2010; Presacco et al.,
2016; Tune et al., 2021; Orf et al., 2025). Yet, most prior research on the effects of attention on
speech tracking has relied on paradigms where a single talker competes with the target speech,
whereas communication also often occurs in contexts with diffuse multi-talker background babble.
Individuals may be particularly susceptible to attentional inward drifts under ambient background
noise (Steindorf et al., 2023). Distraction-related disengagement in multi-talker settings need not
resemble the explicit suppression of a competing speech stream used in selective-attention tasks.
Because the target speech remains behaviorally relevant, a competing speaker is absent, and
attentional decoupling may be intermittent, it is unclear whether external distraction or internal
drift will produce measurable changes in speech tracking under multi-talker noise conditions.

We investigate how external and internal distractions affect neural speech tracking across varying
levels of speech clarity using EEG and TRF analyses. After characterizing the dependence of
neural speech tracking on speech clarity (Experiment 1), we examine how disengagement – from
external visual distraction (Experiment 2) or internal thought (Experiment 3) – modulates neural
tracking. We hypothesize that both forms of distraction will reduce neural speech encoding,
particularly for noise-masked speech. Comparable effects across distraction types would point to
a shared neurocognitive mechanism – such as down-regulation of auditory-cortical gain – through
which attention decouples from sensory processing during effortful listening.

General Methods

Participants

Participants were aged 20-34 years, native English speakers or individuals who grew up in
English-speaking countries (mostly Canada) and had been speaking English since early
childhood (before age 6). All participants reported normal hearing abilities and no severe
psychological or neurological disorders. Specific participant demographics are detailed for each
experiment individually below.

Written informed consent was obtained prior to participation, and participants were compensated
at a rate of $15 per hour. The study adhered to the Declaration of Helsinki, the Canadian Tri-
Council Policy Statement on Ethical Conduct for Research Involving Humans (TCPS2-2014), and
received ethical approval from the Research Ethics Board of the Rotman Research Institute at
Baycrest Academy for Research and Education.

Story materials and stimuli

In each of the three experiments, participants listened to ~2-min stories. Stories were generated
using OpenAI's GPT-3.5 (OpenAI et al., 2023) based on researcher-provided prompts and
manually reviewed to ensure unique character names, absence of non-English terms, proper use
of punctuation, and avoidance of generic openings (e.g., "once upon a time"). Stories were revised
manually or rewritten using GPT-3.5 as needed. Each story was on a different topic (e.g., high
school reunion, friends on a trip; Panela et al., 2024; Herrmann, 2025a). For each story, GPT-3.5
was also used to generate five comprehension questions in a multiple-choice format, with four
options per question. Comprehension questions and answers were manually edited wherever
needed to ensure accuracy (Panela et al., 2024; Herrmann, 2025a).

Audio files were synthesized using Google's AI-based speech synthesizer using the male voice
en-US-Neural2-J (https://cloud.google.com/text-to-speech/docs/list-voices-and-types). Previous
research has demonstrated that AI-generated speech is perceived as highly naturalistic and yields
comparable speech-comprehension effects to human speech (Herrmann, 2023a, 2025c).

In each experiment, participants listened to 24 stories presented in 8 blocks (3 stories per block).
Stories were presented under one of three listening conditions: clear speech (i.e., speech in quiet)
or speech masked by 12-talker babble +6 dB SNR, or -3 dB SNR. Babble noise was derived from
the masker materials of the Revised Speech in Noise test (Bilger, 1984). 12-talker babble
simulates a crowded listening environment while preventing the identification of individual words
in the masker (Bilger, 1984; Bilger et al., 1984; Wilson et al., 2012). The +6 dB and -3 dB SNR
levels were chosen to create challenging listening conditions that correspond to approximately
90% and 55% speech intelligibility (Irsik et al., 2022). All stories and story-noise mixtures were
normalized to the same root-mean-square (RMS) amplitude and presented at approximately 65
dB SPL.

Acoustic presentation environment

148   Data collection took place in a sound-attenuating booth. Auditory stimuli were delivered via
149   Sennheiser HD 25-SP II headphones, with audio output controlled by an RME Fireface UCX
150   external sound card. Stimuli were presented using Psychtoolbox (v3.0.14) in MATLAB 2021b
151   (MathWorks Inc.) on a Lenovo T480 laptop running Microsoft Windows 7. Visual stimuli were
152   projected into the booth via screen mirroring.

Electroencephalography (EEG) recording and preprocessing

EEG data were recorded from 32 scalp electrodes and the left and right mastoids using a BioSemi system (Amsterdam, The Netherlands). The online sampling rate was set to 1024 Hz, with a low-pass filter of 208 Hz. Electrodes were referenced online using a monopolar reference feedback loop, which connected a driven passive sensor and a common-mode-sense (CMS) active sensor, both positioned posteriorly on the scalp.

Preprocessing was conducted in MATLAB (MathWorks, Inc.). Data were first notch-filtered to suppress 60-Hz power line noise using an elliptic filter. Signals were then re-referenced to the averaged mastoids to enhance the signal-to-noise ratio for auditory responses at fronto-central electrodes (Ruhnau et al., 2012; Herrmann, 2023b). Additional filtering included a 0.7 Hz high-pass filter (2449 samples, Hann window) and a 22 Hz low-pass filter (211 samples, Kaiser window with $\beta = 4$). EEG data were segmented into time series locked to story onset and downsampled to 512 Hz. Artifact removal was performed using Independent Component Analysis (ICA) in FieldTrip MATLAB toolbox to eliminate blinks, eye movements, and muscle artifacts (Makeig et al., 1995; Oostenveld et al., 2011). After ICA, further artifact rejection was applied by zeroing segments where EEG amplitude varied by more than 80 μV within 0.2 s in any channel (Dmochowski et al., 2012; Cohen and Parra, 2016). Finally, data were low-pass filtered at 10 Hz (251 points, Kaiser window) to focus on low-frequency neural signals most sensitive to acoustic features (Di Liberto et al., 2015; Zuk et al., 2021).

Calculation of amplitude-onset envelopes

For each clear speech signal, a cochleogram was generated using a simple auditory-periphery model consisting of 30 auditory filters (McDermott and Simoncelli, 2011). The amplitude envelope obtained for each auditory filter was compressed by a factor of 0.6 to simulate inner ear compression (McDermott and Simoncelli, 2011). The amplitude envelopes were then averaged across auditory filters and low-pass filtered using a 40-Hz Butterworth filter. The first derivative was calculated and all negative values were set to zero to obtain the amplitude-onset envelope (Hertrich et al., 2012; Tune et al., 2021; Yasmin et al., 2023; Panela et al., 2024). The amplitude-onset envelope was computed since it elicits strong neural speech tracking (Hertrich et al., 2012). The amplitude-onset envelope was down-sampled to match the EEG sampling rate.

Analysis of neural tracking of speech

To investigate neural speech tracking, we adopted Temporal Response Functions (TRFs) and
assessed EEG prediction accuracy (Crosse et al., 2016; Crosse et al., 2021). The TRF is a system
identification model that employs regularized linear regression to relate sensory stimulus features
with neural responses. Specifically, we utilized the Multivariate Temporal Response Function
(mTRF) Toolbox in MATLAB (Crosse et al., 2016; Crosse et al., 2021) to map the amplitude-onset
envelope of speech onto EEG signals (forward model). To mitigate overfitting, ridge regularization
was applied, with the ridge parameter (λ) set to 10, a value informed by prior research to avoid
extreme value selection during cross-validation and optimize computational efficiency (Fiedler et
al., 2017; Fiedler et al., 2019; Yasmin et al., 2023; Panela et al., 2024).

For each story, 100 randomly selected 25-s EEG snippets and their corresponding time series of
the amplitude-onset envelope were extracted for model training and testing using leave-one-
sample-out cross-validation. On each iteration, one snippet's EEG data and the corresponding
time series of the amplitude-onset envelope were reserved as the test set, while the remaining,
non-overlapping data were used for training. The model was trained using time lags ranging from
0 to 0.4 s, employing the *mTRFcrossval* and *mTRFtrain* functions. To assess prediction accuracy,
the acoustic feature time series of the test dataset was convolved with the TRF weights to
generate a predicted EEG time series, separately for each of the 100 iterations. EEG prediction
accuracy was calculated as the Pearson correlation between the predicted and actual EEG data,
computing using the *mTRFpredict* function. EEG prediction accuracy was averaged across the
leave-one-out iterations.

To investigate the neural-tracking response amplitude at a more nuanced timescale, TRFs for
each training dataset were calculated for lags ranging from -0.15–0.55 s. Baseline correction was
performed by subtracting the mean signal across -0.15–0 s from the TRF data at each time point.
The weights were averaged across the 100 training datasets.

For each participant, TRF weights and EEG prediction accuracy were averaged across stories,
separately for the three different speech-clarity conditions and averaged across the frontal-central
electrode cluster (F3, F4, Fz, FC1, FC2, Cz, FC5, FC6), which is sensitive to auditory cortical
activity (Näätänen and Picton, 1987; Picton et al., 2003).

Time courses of TRF weights exhibit positive and negative deflections (peaks), akin to traditional
event-related potentials (Luck, 2014; Crosse et al., 2016; Crosse et al., 2021). To examine
response magnitude differences across speech-clarity levels and attention conditions, we focused
on the P1-N1 (first major positive peak minus first major negative peak) and P2-N1 amplitude

differences. Using peak-to-peak measures helps reduce sensitivity to slow baseline shifts and

condition-related waveform morphology differences, providing a robust index of response

magnitude across conditions. The amplitudes of P1, N1, and P2 components were determined

by local extrema (maxima for P1 and P2, minima for N1) within a 0.1-s time window centered on

each component based on averaged data across all participants in each experiment. Latencies

for P1, N1, and P2 were identified as the time points corresponding to these extrema. For each

participant, amplitudes were calculated within a 0.03-s window (±0.015 s) centered on the

latencies of each component from grand-average data. This approach accommodates inter-

individual latency jitter and avoids unreliable participant-specific peak-picking in conditions where

some participants might not show a clear extremum, while keeping the P1, N1, and P2 windows

non-overlapping. The P1-N1 and P2-N1 amplitude differences were computed by subtracting N1

amplitudes from P1 and P2, respectively.

We also explored the extent to which semantic surprisal was tracked by the brain, but these

analyses were ambiguous and did not enable drawing clear conclusions about the semantic

processing in the current study, mirroring to some extent other debates in the field (Daube et al.,

2019; Gillis et al., 2021; Verschueren et al., 2022; Chalehchaleh et al., 2025). We provide these

explorations in the Supplementary Materials (Table S1, Figures S1 and S2).

Statistical analyses

All statistical analyses were carried out using MATLAB (MathWorks) and JASP software (version

0.19.2). Details about the specific tests used are provided in the relevant sections for each

experiment.

Experiment 1: Influences of speech clarity on neural speech tracking

Experiment 1 measured speech-evoked EEG activity under focused listening across speech-

clarity conditions to establish the baseline signal changes underlying distraction effects tested in

Experiments 2 and 3.

Method and materials

Participants

Twenty-four younger adults participated in Experiment 1 (mean = 25.2 years; SD = 3.85; 14

females, 10 males).

Stimuli and procedure

Twenty-four stories, each lasting between 109 and 133 seconds, were generated for Experiment

1. The stories were presented under one of three speech-clarity conditions: clear speech (no

noise), +6 dB SNR, or -3 dB SNR (12-talker babble). The experiment was divided into 8 blocks,

with each block containing 3 stories with the 3 speech-clarity conditions. Assignment of speech-

clarity conditions to specific stories was counter-balanced across participants. During each trial,

participants listened attentively to the story while their EEG signals were recorded. After each

story, participants answered five multiple-choice comprehension questions, each with four

response options (chance level = 25%). Additionally, participants provided a self-reported gist

rating on a scale from 1 to 9, with higher ratings indicating a better understanding of the story's

gist. Gist ratings correlate highly with speech intelligibility as measured using word report scores

(Davis and Johnsrude, 2003; Ritz et al., 2022). The exact wording for the gist rating was: "I

understood the gist of the story. Please rate this statement independently of how you felt about

the other stories". Gist ratings were linearly transformed to ranging from 0 to 1 in order to facilitate

interpretation similar to proportions (Mathiesen et al., 2024; Herrmann, 2025a). Participants

underwent a practice block prior to the main experimental procedures. The full experimental

session lasted approximately 2 hours per participant, with the experimental procedures –

including EEG recording – taking about 1 hour.

Analysis

The effect of Speech Clarity (clear, +6 dB, -3 dB SNR) on comprehension accuracy, gist rating,

EEG prediction accuracy, P1-N1 amplitude, and P2-N1 amplitude were evaluated in separate

one-way repeated-measures analysis of variance (rmANOVA). Post-hoc pairwise comparisons

were performed using the Holm correction for multiple comparisons (Holm, 1979).

Results

Both comprehension accuracy and gist ratings (Figure 1A) were affected by speech clarity

(comprehension: $F_{2, 46} = 11.986$, $p = 6.5·10^{-5}$, $\eta_p^2 = .343$; gist: $F_{2, 46} = 34.466$, $p = 7.1·10^{-10}$, $\eta_p^2 =$

.600). Performance dropped for the lowest speech clarity (-3 dB SNR): pair-wise comparisons

showed lower scores at -3 dB than at clear and at +6 dB for both behavioural measures (all p_{Holm}

< .01). Clear and +6 dB did not differ significantly on either measure (all p_{Holm} > .05).

--- Insert Figure 1 around here ---

EEG prediction accuracy (Figure 1C) showed a significant main effect of speech clarity ($F_{2, 46}$ =
9.376, p = $3.8 \cdot 10^{-4}$, η_p^2 = .269). The values were lower at -3 dB SNR compared to clear (t_{23} =
3.882, p_{Holm} = .002, d = .843) and +6 dB (t_{23} = 2.843, p_{Holm} = .018, d = .471), whereas clear and
+6 dB did not differ (t_{23} = 1.966, p_{Holm} = .062, d = .409). Scalp topographies revealed highest EEG
prediction accuracy at frontal-central electrodes that are sensitive to auditory cortex activity.

TRF time courses are displayed in Figure 1B, indicating that background noise affects the shape
and deflections of the response. To avoid that analyses of response amplitudes are affected by
overall morphological differences, analyses focused on the P1-N1 and P2-N1 amplitudes (Lanting
et al., 2013; de Boer and Krumbholz, 2018; Herrmann, 2025a, b). The speech-clarity effect for
the P1-N1 amplitude ($F_{2, 46}$ = 52.108, p = $1.5 \cdot 10^{-12}$, η_p^2 = .694) revealed a non-monotonic pattern:
responses were largest at +6 dB and remained elevated at -3 dB relative to clear (all p_{Holm} < .05).
For the P2-N1 amplitude the effect of speech clarity ($F_{2, 46}$ = 8.110, p = $9.6 \cdot 10^{-4}$, η_p^2 = .261) showed
larger amplitudes at +6 dB compared to -3 dB and clear (both p_{Holm} < .05), whereas -3 dB and
clear did not differ significantly (p_{Holm} = .110; Figures 1B, D).

In sum, the results of Experiment 1 show that background babble impairs speech comprehension,
gist rating, and overall EEG prediction accuracy, with declines observed only at the lowest speech
clarity (-3 dB). Early TRF responses (both P1-N1 and P2-N1 components) were selectively
enhanced under moderate noise – peaking at + 6 dB – consistent with recent work suggesting
stochastic resonance phenomena in neural speech tracking and for more simple sounds
(Parbery-Clark et al., 2011; Alain et al., 2014; Herrmann, 2025a, b). With the foundation of
Experiment 1, Experiment 2 examines how disengagement from listening due to visual distraction
affects neural speech tracking, and whether disengagement effects are affected by speech clarity.

Experiment 2: Influences of visual distraction on neural speech tracking

Method and materials

Participants

Twenty-four younger adults participated in Experiment 2 (mean = 26 years; SD = 3.95; 14
females, 10 males).

Stimuli and procedure

We generated a new set of 24 stories (102–141 s each) for Experiment 2. Similar to Experiment
1, the session comprised 8 blocks, with each block containing 3 stories, one for each speech-
clarity condition (clear, +6 dB, and -3 dB SNR). In four blocks, participants attended to the speech
('attend' condition), whereas in the other four blocks, participants ignored the speech ('ignore'
condition). Assignment of speech clarity and attention conditions to specific stories was counter-
balanced across participants. Blocks alternated in attention conditions, and the starting attention
condition was counter-balanced across participants.

In 'Ignore' blocks, participants were asked to ignore the stories and instead perform a visual 1-
back task. The visual 1-back task utilized images of white digits (0 to 9) on a black background,
sourced from the MNIST Handwritten Digit Classification Dataset (Herrmann 2024; Deng et al.
2012). Each digit image was displayed for 0.25 seconds, followed by a 0.25-s black screen before
the next image appeared. Visual stimuli thus repeat as a rate of 2 Hz. The digit stream included
a digit repetition – albeit a different image sample – every 6 to 12 digits. Participants were asked
to press a keyboard button as soon as they detected a repetition. Hit rates and response times
were recorded. Additionally, participants rated the degree to which they were able to ignore the
speech using a 9-point scale (1 = strongly disagree, 9 = strongly agree; linearly transformed to
range from 0-1). No comprehension questions or gist ratings were used in the 'Ignore' blocks to
avoid incentivizing participants to listen to the stories. In 'Attend' blocks, participants were
instructed to listen attentively to the stories, followed by comprehension questions and a gist rating
(similar to Experiment 1). Visual digit images at a 2-Hz rate were presented concurrently with the
stories to ensure similar visual and auditory stimulus presentation in 'attend' and 'ignore' blocks,
but participants were asked to pay attention only to the speech in 'attend' blocks and ignore the
visual stimuli.

Participants underwent a practice block prior to the main experimental procedures to familiarize
them with the tasks. The full experimental session lasted approximately 2 hours per participant,
with the experimental procedures – including EEG recording – taking about 1 hour.

Analysis

For each participant, comprehension accuracy and gist ratings from 'attend' blocks were averaged
across the four stories per speech-clarity condition (clear, +6 dB, -3 dB). Reaction times, visual
detection accuracy, and able-to-ignore ratings from 'ignore' blocks were also averaged across the
four stories per speech-clarity condition. For each behavioral measure, we calculated a one-way

rmANOVA with the within-participant factor Speech Clarity (clear, +6 dB, -3 dB). Note that

behavioral measures did not comprise the Attention factor.

Because Experiment 2 comprised a repetitive visual stream of digits occurring at 2 Hz, this

provided an opportunity to neurally investigate attention to the visual stream. To this end, EEG

data for each story / visual steam were divided into 7-s epochs (shifted by 0.5 s across the story

duration) that were time-locked to a visual stimulus. A fast Fourier transform (FFT) was calculated

for each epoch. Inter-trial phase coherence (ITPC; Lachaux et al., 1999) was then calculated

across epochs to obtain a measure of neural responses to the visual stream. ITPC was used here

instead of evoked amplitude because ITPC spectra do not show 1/f patterns that amplitude

spectra exhibit (Herrmann et al., 2019; Keitel et al., 2019; Irsik et al., 2021), but which can

complicate analyses (Donoghue et al., 2020; Donoghue et al., 2022). Mathematically, ITPC and

evoked amplitude are otherwise closely related. We calculated the average across ITPC at the 2-

Hz visual repetition rate and its harmonics up to 10 Hz (Irsik et al., 2021). Taking into account the

harmonics is needed because data are not sinusoidal in the time domain (as indicated by the

presence of harmonics). ITPC was averaged across 3 occipital channels (O1, Oz, O2) to focus

on visual-cortex activity. For the statistical analysis, ITPC was averaged across stories, separately

for each speech-clarity condition (clear, +6 dB, -3 dB) and attention condition (Attend, Ignore),

and a rmANOVA with the two factors was calculated.

For the neural speech tracking analysis, EEG prediction accuracy, P1-N1 amplitude, and P2-N1

amplitude were evaluated in separate rmANOVAs with the factors Speech Clarity (clear, +6 dB, -

3 dB) and Attention (Attend, Ignore). Post-hoc pairwise comparisons were performed using the

Holm correction for multiple comparisons (Holm, 1979).

Results

Behavioral performance during 'attend' speech blocks are shown in Figure 2A. Speech Clarity

affected both comprehension accuracy ($F_{2, 46}$ = 11.279, p = $1 \cdot 10^{-4}$, η_p^2 = .329) and gist ratings ($F_{2,}$

$_{46}$ = 47.964, p = $5.6 \cdot 10^{-12}$, η_p^2 = .676). Similar to Experiment 1, performance was lower in the -3

dB condition compared to the clear and +6 dB conditions for both measures (both p_{Holm} < .01).

For gist rating, +6 dB was also lower than clear (t_{23} = 3.174, p_{Holm} = .004, d = .444), whereas for

comprehension accuracy, clear and +6 dB did not differ (t_{23} = .554, p_{Holm} = .585, d = .103). During

'ignore' blocks (Figure 2B), speech clarity did not affect visual-detection accuracy ($F_{2, 46}$ = .506, p

= .606, η_p^2 = .022) or reaction times ($F_{2, 46}$ = 2.855, p = .068, η_p^2 = .110). In contrast, 'able-to-

ignore' ratings indicate that participants found the stories at -3 dB easier to ignore compared to

clear and +6 dB (both p_{Holm} < .05; effect of Speech Clarity: $F_{2, 46}$ = 7.828, p = .001, η_p^2 = .254).

Consistent with the periodic visual stream and isochronously occurring visual evoked potentials,

ITPC showed clear peaks at 2 Hz and harmonics (Figure 2C). Peak ITPC (2–10 Hz harmonics

averaged; Figure 2D) was higher when participants ignored than attended to the speech ($F_{1, 23}$ =

78.844, p = 6.8·10^{-9}, η_p^2 = .774). There was no effect of Speech Clarity ($F_{2, 46}$ = .882, p = .421, η_p^2

= .037), and no interaction ($F_{2, 46}$ = 1.676, p = .198, η_p^2 = .068). These data show that participants

indeed attended to the visual digit stream more when they ignored relative when they attended to

the speech.

EEG prediction accuracy was lower for -3 dB than at Clear (t_{23} = 4.137, p_{Holm} = 8·10^{-4}, d = .882)

and +6 dB (t_{23} = 4.665, p_{Holm} = 3.2·10^{-4}, d = .689), with no difference between Clear and +6 dB (t_{23}

= 1.001, p_{Holm} = .327, d = .193; effect of Speech Clarity: $F_{2, 46}$ = 12.368, p = 5·10^{-5}, η_p^2 = .350

Figure 2F). EEG prediction accuracy was greater for the attend than ignore conditions (effect of

Attention: $F_{1, 23}$ = 7.244, p = .013, η_p^2 = .240), although the significant interaction ($F_{2, 46}$ = 3.561, p

= .036, η_p^2 = .134) shows that this difference was specific for the -3 dB SNR condition (t_{23} = 3.346,

p_{Holm} = .031, d = .703).

380   --- Insert Figure 2 around here ---

TRF time courses are displayed in Figure 2E. The P1-N1 amplitude was smallest for clear speech,

largest for +6 dB SNR, and in-between for -3 dB (all pairwise p_{Holm} < .001; effect of Speech Clarity:

$F_{2, 46}$ = 33.186, p = 1.2·10^{-9}, η_p^2 = .591). There was no effect of Attention ($F_{2, 46}$ = 3.008, p = .096,

η_p^2 = .116), but the interaction between Speech Clarity and Attention was significant ($F_{2, 46}$ = 3.726,

p = .032, η_p^2 = .139). Although not surviving Holm correction, the interaction was driven by the

attention effect at -3 dB SNR as indicated by the uncorrected comparison (t_{23} = 2.471, p = .021,

p_{Holm} = .102, d = .504). Attention effects for other the two speech-clarity levels were not significant

(both effects, p_{Holm} and p > .50).

The P2-N1 amplitude was larger at +6 dB than at clear (t_{23} = 2.648, p_{Holm} = .014, d = .382) and -3

dB (t_{23} = 6.342, p_{Holm} = 5.4·10^{-6}, d = .947), and amplitudes for clear speech were larger than for -

3 dB (t_{23} = 3.450, p_{Holm} = .004, d = .565; effect of Speech Clarity: $F_{2, 46}$ = 19.481, p = 7.4·10^{-7}, η_p^2

= .459). P2-N1 amplitudes were larger when participants attended to the speech compared to

when they ignored it (effect of Attention: $F_{2, 46}$ = 4.426, p = .047, d = .194). There was no interaction

($F_{2, 46}$ = 0.089, p = .915, η_p^2 = .004).

In sum, disengagement from speech listening due to visual distraction reduced signatures of
neural speech tracking (EEG prediction accuracy and TRF component amplitudes), most
prominently for speech masked at -3 dB SNR. This result was mirrored by participants also
reporting greater ease ignoring speech at -3 dB than at +6 dB and clear speech. Experiment 2
demonstrates that neural speech tracking is sensitive to disengagement from listening due to
external distraction, such that the amplitude-envelope onset of the speech is encoded less
strongly. Critically, disengagement from listening in everyday life not only occurs because of
external distraction, but also because of internal distraction when a listener's mind is wandering
off. Experiment 3 was designed to examine whether disengagement due to internal distractions
affects neural speech tracking.

Experiment 3: Influences of internal imagination on neural speech tracking

Method and materials

Participants

Twenty-four younger adults were recruited for Experiment 3 (mean = 26.3 years, SD = 4.37; 13
females, 10 males, 1 non-binary).

Stimuli and procedure

A new set of 24 stories was generated, each lasting between 111 and 139 seconds. The
experiment was divided into 8 blocks, with each block containing 3 stories, one for each speech-
clarity condition (clear, +6 dB, and -3 dB SNR). In four blocks, participants attended to the speech
('attend' condition), whereas in the other four blocks, participants ignored the speech ('ignore'
condition). Assignment of speech clarity and attention conditions to specific stories was counter-
balanced across participants. Blocks with different attention conditions alternated in the
experimental session and starting attention condition in the session was counter-balanced across
participants. Participants underwent a practice block prior to the main experimental procedures
to familiarize them with the tasks.

In the 'attend' blocks, participants attentively listen to the stories, answered comprehension
questions, and rating gist understanding (similar to Experiments 1 and 2). In 'ignore' blocks,
participants were asked to disregard the spoken stories and engage in an 'imagination' task,
where they mentally wandered off. During the imagination task, participants were provided with a
verbal instruction outlining a daily scenario accompanied by three concrete cues (e.g., scenario:
public transport commute; cues: seats, tapping cards, and bus driver). After a brief preparation

period, participants were instructed to immerse themselves in the imagined scenario as soon as

the auditory stimulus began. They were encouraged to use internal narratives and mental imagery

to facilitate the task, and to mentally jump freely to any other scenarios that come to mind as much

as they wanted. This task aimed to mirror disengagement in daily life where people may think

about concrete daily activities or interactions (Smallwood and Andrews-Hanna, 2013; Ibaceta et

al., 2024), although here facilitated by task instructions. Following each story, participants

evaluated their experience using a 9-point scale (1 = strongly disagree, 9 = strongly agree) across

four statements: whether they were able to ignore the speech, mentally wander off, visualize the

given scenario, and would be able retell the story. No comprehension questions or gist ratings

were used in the 'ignore' blocks to avoid that participants attend to the stories. Ratings were

linearly transformed to range from 0-1 to facilitate interpretability (Mathiesen et al., 2024;

Herrmann, 2025a).

Participants underwent a practice block prior to the main experimental procedures to familiarize

them with the tasks. The full experimental session lasted approximately 2 hours per participant,

with the experimental procedures – including EEG recording – taking about 1 hour.

Analysis

For each behavioral measure, a one-way rmANOVA with the within-participant factor Speech

Clarity was calculated. Behavioral measures did not comprise the Attention factor to avoid that

participants are biased towards the irrelevant dimension. For each EEG measure (EEG prediction

accuracy, P1-N1 amplitude, and P2-N1 amplitude), a two-way rmANOVA was conducted using

the with-participant factors Speech Clarity (clear, +6 dB, and -3 dB SNR) and Attention (Attend,

Ignore).

Results

Speech clarity affected both comprehension accuracy ($F_{2, 46}$ = 20.647, p = $4 \cdot 10^{-7}$, η_p^2 = .473) and

gist ratings ($F_{2, 46}$ = 40.847, p = $7.5 \cdot 10^{-11}$, η_p^2 = .637). Both behavioral measures were lower at -3

dB compared to the clear (both p_{Holm} < .001) and +6 dB (both p_{Holm} < .001), with no difference

between Clear and +6 dB (both p_{Holm} > .05; Figure 3A).

Behavioral ratings related to the imagination task are shown in Figure 3B. Participants rated it to

be easier to ignore stories (effect of Speech Clarity: $F_{2, 46}$ = 18.591, p = $1.2 \cdot 10^{-6}$, η_p^2 = .447; p_{Holm}

< .05 for all pairwise comparisons) and to be less able to retell stories with decreasing speech

clarity (effect of Speech Clarity: $F_{2, 46}$ = 26.878, p = $1.9 \cdot 10^{-8}$, η_p^2 = .539; p_{Holm} < .05 for all

comparisons). The ability to mentally visualize was rated higher at -3 dB compared to Clear (t_{23} =

4.236, p_{Holm} = $9.4 \cdot 10^{-4}$, d = .552) and +6 dB (t_{23} = 3.636, p_{Holm} = .003, d = .364; effect of Speech

Clarity: $F_{2, 46}$ = 11.030, p = $1.2 \cdot 10^{-4}$, η_p^2 = .324), but speech clarity had no effect on participants'

ability to wander off mentally ($F_{2, 46}$ = .215, p = .807, η_p^2 = .009). Overall, the ratings suggest that

participants could well imagine mentally and ignore the speech, such that they felt relatively

unable to retell the story.

EEG prediction accuracy decreased with decreasing speech clarity ($F_{2, 46}$ = 26.774, p = $1.9 \cdot 10^{-8}$,

η_p^2 = .538; p_{Holm} < .05 for all pairwise comparisons; Figure 3D). Critically, EEG prediction accuracy

was lower during imagination (ignore) than attentive listening (effect of Attention: $F_{1, 23}$ = 7.405, p

= .012, η_p^2 = .244), with no Speech Clarity × Attention interaction ($F_{2, 46}$ =.091, p = .913, η_p^2 =

.004).

468   --- Insert Figure 3 around here ---

TRF time courses are depicted in Figure 3C. Speech Clarity modulated the P1-N1 amplitudes

(main effect of Speech Clarity: $F_{2, 46}$ = 40.196, p $8 \cdot 10^{-11}$, η_p^2 = .636), as amplitudes were largest

at +6 dB and smallest in the Clear condition (all pairwise p_{Holm} < .001). Mental imagination reduced

P1-N1 amplitudes compared to attentive speech listening (effect of Attention: $F_{1, 23}$ = 4.862, p =

.038, η_p^2 = .175). The interaction was not significant ($F_{2, 46}$ = .585, p = .561, η_p^2 = .025).

P2-N1 amplitudes also varied with Speech Clarity ($F_{2, 46}$ = 29.536, p = $5.6 \cdot 10^{-9}$, η_p^2 = .562), such

that amplitudes were largest at +6 dB and smallest in the Clear condition (all pairwise p_{Holm} < .05).

Mental imagination reduced P2-N1 amplitude compared to attentive speech listening (effect of

Attention: $F_{1, 23}$ = 5.837, p = .024, η_p^2 = .202). The interaction was not significant ($F_{2, 46}$ = 1.516, p

= .230, η_p^2 = .062).

In sum, Experiment 3 shows, consistent with the findings from Experiments 1 and 2, that

background babble can enhance early neural responses to the amplitude-envelope of speech

(see also Yasmin et al., 2023; Panela et al., 2024; Herrmann, 2025a). Experiment 3 results further

indicate that participants can ignore speech well while engaging in a mental imagination task.

Critically, Experiment 3 extends the external distraction effect in Experiment 2 by showing that

internal distraction (imagination) reduces the encoding of the amplitude-envelope of speech,

indicated by all three signatures of neural speech tracking. The main effect of attention in the

absence of an interaction indicates that internal distraction reduces speech encoding across

speech-clarity levels.

Comparison of attention effects between external and internal distraction

In order to directly compare the attention effects between external and internal distraction, we first

calculated an rmANOVA for the participants' ratings of their ability to ignore speech when they

performed the visual distraction or internal imagination task. Rating scores were used as

dependent measure, with the within-participant factor Speech Clarity (clear, +6 dB, -3 dB) and the

between-participants factor Distraction Type (external, internal; Experiments 2 and 3). The

rmANOVA showed higher speech ignoring ratings for -3 dB than for the clear (t_{47} = 6.125, p_{Holm} =

$5.6 \cdot 10^{-9}$, d = .892) and +6 dB conditions (t_{47} = 4.598, p_{Holm} = $6.7 \cdot 10^{-5}$, d = .600), and for +6 dB than

clear speech (t_{47} = 2.553, p_{Holm} = 0.014, d = .292; effect of Speech Clarity: $F_{2, 92}$ = 24.187, p =

$3.6 \cdot 10^{-9}$, η_p^2 = .345). However, there was no effect of Distraction Type ($F_{2, 92}$ = .735, p = .482, η_p^2

= .016), nor an interaction ($F_{1, 46}$ = 2.768, p = .103, η_p^2 = .057), indicating that participants were

able similarly ignore speech under external and internal distraction.

Next, we calculated an rmANOVA – separately for the three TRF metrics – using the within-

participants factors Speech Clarity (clear, +6 dB, -3 dB) and Attention (Attend, Ignore), and the

between-participants factor Distraction Type (external, internal; Experiments 2 and 3). If there

were any differences in the attention effect between distraction types, we would expect a

significant interaction involving Attention and Distraction Type. In no analysis did we observe a

significant interaction involving Attention and Distraction Type (for all p > 0.05), providing no

evidence for attention effects to differ between the type of distraction.

507   **Discussion**

The current EEG study investigated how disengagement from listening due to external and

internal distractions alters neural speech tracking under different levels of background noise. We

observed enhanced early neural responses (<0.2 s) to the speech envelope for speech masked

by background babble compared to speech in quiet (Experiments 1-3), suggesting stochastic

facilitation. Critically, neural speech tracking was reduced when individuals were distracted by a

visual-stimulus stream (Experiment 2) or by internal imagination (Experiment 3). Overall, our

study shows that disengagement from listening – towards external or internal spaces – that many

people experience in crowded, noisy situations is associated with reduced neural encoding of

speech.

Background babble altered neural speech tracking

Neural speech tracking was assessed using different TRF metrics (Crosse et al., 2016): EEG

prediction accuracy pools information across time (0-0.4 s), whereas P1-N1 and P2-N1

amplitudes indicate early sensory processing.

We observed a decline in EEG prediction accuracy with decreasing speech clarity (Figures 1-3),

with the most prominent decline for the -3 dB SNR condition, mirroring story-comprehension

accuracy and gist ratings (Figures 1-3). These results are consistent with previous research

showing a close link between EEG prediction accuracy and speech perception (Ding and Simon,

2013; Vanthornhout et al., 2018; Synigal et al., 2023).

Early TRF responses showed a different picture. P1-N1 and P2-N2 amplitudes for speech at

moderate and difficult levels of background noise were increased relative to clear speech. This

increase may be surprising. However, several recent works observed enhanced early speech-

tracking responses for speech presented in babble noise (Panela et al., 2024; Herrmann, 2025a,

b). The findings are also in line with research using tone bursts, syllables, and high-frequency

sound modulations, showing noise-enhanced early sensory responses (Alain et al., 2009;

Parbery-Clark et al., 2011; Alain et al., 2012; Shukla and Bidelman, 2021).

Recent research specifically examined the noise-related enhancement of early neural tracking

responses, showing that the enhancement is independent of attention, present at very high SNRs,

and strongest for multi-talker babble (Herrmann, 2025a, b). These findings argue against a

listening effort or attention mechanism, but instead suggest stochastic resonance/facilitation

(Stocks, 2000; Stein et al., 2005; McDonnell and Ward, 2011). Background noise can introduce

random fluctuations in membrane potential that transiently bring neurons closer to their firing

threshold (Ward et al., 2002; Moss et al., 2004; Ward et al., 2010), effectively priming them to

respond when speech-related input arrives. This stochastic resonance mechanism increases the

probability of stimulus-driven, time-locked firing without inducing independent noise-driven firing

(McDonnell and Abbott, 2009; McDonnell and Ward, 2011), thereby enhancing neural tracking of

the speech envelope rather than degrading it through masking. Regardless of the specific

mechanisms, the current study shows noise-related enhancements in neural speech tracking

across all three experiments.

In addition to amplitude changes, the response morphology also appeared to vary with

background noise, including a negative waveform shift (Figures 1B, 2E, 3C). Although

morphology was not the focus here, the pattern may reflect a more fundamental change in the

operating state of auditory system under noisy conditions. Prior work has shown that sustained
noise can elicit ultra-low-frequency auditory-cortical activity (Ross et al., 2002; Okamoto et al.,
2011; Barascud et al., 2016; Herrmann et al., 2022), though mechanisms remain incompletely
understood. Background noise may therefore modulate both transient speech-envelope
responses and the baseline state of auditory cortex.

External and internal distraction reduces neural speech tracking

The main purpose of the current study was to examine how disengagement from speech listening
due to external and internal distraction affects the neural encoding of speech as indicated by the
TRF metrics. EEG prediction accuracy, P1-N1 amplitude, and P2-N1 amplitude all decreased
when individuals attended to an external visual stream or internal thoughts and imagination.

Participants were able to disengage from listening in both the external and internal distraction
tasks. Enhanced inter-trial phase coherence when attending to the visual stream, and ignoring
speech, is consistent with previous work on attention and visual steady-state responses (Müller
and Hillyard, 2000; Joon Kim et al., 2007; Keitel et al., 2017; Keitel et al., 2019), and with
participants' subjective reports of being able to ignore the stories well. Similarly during the
imagination task, participants indicated being able to ignore and unable to retell the story. They
also stated that they could visualize well mentally. Subjective reports are common measures of
mental attentional drift (Smallwood and Schooler, 2006; Stawarczyk et al., 2011; Weinstein,
2018), because such internal changes resist objective assessments. The reported subjective
experiences point to mental disengagement from listening during both distraction tasks.

Previous work on attentional disengagement from speech listening has mainly focused on speech
in the presence of one competing speaker, finding that the tracking of the speech envelope of a
target speaker is reduced when listeners disengage from listening to it (Ding and Simon, 2012;
Power et al., 2012; Hambrook and Tata, 2019; Fuglsang et al., 2020; Kraus et al., 2021;
Straetmans et al., 2024). Fewer works have shown attention effects when individuals attend to
visual distractors while ignoring speech (Vanthornhout et al., 2019; for divided attention see Xie
et al., 2023), or under ambient speech-masking conditions, such as multi-talker babble. Such
conditions, however, reflect common scenarios in which listeners disengage in daily life, making
them prime targets for understanding altered speech encoding. Attention effects in single
competing-speaker paradigms have further been described as changes in TRF morphology, such
as a reduced/absent P2 component for ignored speech (Ding and Simon, 2012; Fiedler et al.,
2019; Kraus et al., 2021). By comparison, in the present babble-masking distraction paradigms,

attention effects were expressed mainly as reductions in TRF metrics, with limited evidence for

clear morphology-specific changes (Figures 2 and 3).

Importantly, the current work is first to show reduced speech encoding when individuals

disengage from speech listening and engage in internal thought and imagination. This is critical,

because internally oriented disengagement from listening is common among older people with

hearing difficulties when listening effort exceeds available resources and/or perceived

conversational benefit (Heffernan et al., 2016; Pichora-Fuller et al., 2016; Herrmann and

Johnsrude, 2020). We used an intentional imagination task over a spontaneous mind wandering

task as a proof on concept. The processes underlying intentional imagination and spontaneous

mind wandering may be similar (Villena-González and Cosmelli, 2020; Safati et al., 2024),

although others have argued that dissociating them may be beneficial (Forster and Lavie, 2009;

Seli et al., 2016a; Seli et al., 2016b). Critically, the reduced speech tracking due to internal thought

highlights neurophysiological feedback mechanisms, supressing auditory cortex activity in the

light of mental imagination.

Our data provide little evidence that external and internal distractions differ in their suppression

of neural speech tracking, suggesting that both may converge on a common pathway through

which attention decouples from sensory processing. Although visual distraction engages visual

sensory areas that internal thoughts may not, both likely recruit prefrontal and parietal networks

that feed back to auditory cortex (Corbetta and Shulman, 2002; Braga et al., 2013; Puschmann

et al., 2024), similarly reducing auditory-cortical gain and neural speech tracking. However,

although no difference in the self-reported ability to ignore speech was observed, the external and

internal distraction tasks were not explicitly equated for difficulty/cognitive load. Therefore, similar

TRF reductions across experiments do not by themselves establish identical pathways, but could

also reflect different mechanisms that produce a comparable degree of attentional withdrawal

from speech.

Nevertheless, there were a few surface-level differences between external and internal

distractions that may require additional research. Based on behavioral mind-wandering work

(Steindorf et al., 2023) and consistent with behavioral ratings in the current study (Figures 2B and

3B), we had expected the effects of attention on neural speech tracking to be greatest under the

challenging -3 dB SNR condition. This was observed for some TRF metrics during external visual

distraction (e.g., Figure 2F), but not during internal distraction, for which we only observed a main

effect of attention, but no interaction with speech clarity. This may potentially indicate that internal

thought and imagination can lead to a blanket form of disengagement from the acoustic

surroundings, whereas a distracting visual stream leaves more capacity for speech encoding in
quiet, but very little when speech is presented in noise. Nevertheless, direct comparisons between
distraction types did not show any significant differences. It is worth noting, however, that the
current experimental design was not specifically suited for such direct interpretations due to its
between-participant nature. This is thus an area for future research.

Conclusions

This study investigated how disengagement from speech listening due to distractions by external
visual stimuli or internal thought alters neural speech encoding under different degrees of
background masking babble. We show that background babble enhances early neural responses
to the speech envelope, consistent with recent reports of stochastic resonance. Critically, we
show that external and internal distractions both lead to reduced neural tracking of the amplitude-
onset envelope of speech, pointing to a converging neural pathway that supresses auditory
cortex. The current study provides a neurophysiological marker that indexes different ways of
disengagement from listening that are common among older adults with hearing challenges.

Declaration of conflict of interest

629 None.

Data availability

631 The data will be made available to others upon reasonable request.

Acknowledgements

633 We thank Tiffany Lao and Saba Junaid for their support with stimuli creation and data collection.
634 This study has been supported by BH's Canada Research Chair program (CRC-2023-00383) and
635 BH's Natural Sciences and Engineering Research Council of Canada (Discovery Grant: RGPIN-
636 2021-02602).

References

Alain C, McDonald K, Van Roon P (2012) Effects of age and background noise on processing a
mistuned harmonic in an otherwise periodic complex sound. Hearing Research 283:126-
135. doi: 10.1016/j.heares.2011.10.007.

Alain C, Roye A, Salloum C (2014) Effects of age-related hearing loss and background noise on
neuromagnetic activity from auditory cortex. Frontiers in Systems Neuroscience 8:Article
8. doi: 10.3389/fnsys.2014.00008.

Alain C, Quan J, McDonald K, Van Roon P (2009) Noise-induced increase in human auditory
evoked neuromagnetic fields. European Journal of Neuroscience 30:132-142. doi:
10.1111/j.1460-9568.2009.06792.x.

Barascud N, Pearce MT, Griffiths TD, Friston KJ, Chait M (2016) Brain responses in humans
reveal ideal observer-like sensitivity to complex acoustic patterns. Proceedings of the
National Academy of Sciences 113:E616-E625. doi.

Bilger RC (1984) Manual for the clinical use of the revised SPIN Test. Champaign, IL, USA: The
University of Illinois.

Bilger RC, Nuetzel JM, Rabinowitz WM, Rzeczkowski C (1984) Standardization of a test of speech
perception in noise. Journal of Speech, Language, and Hearing Research 27:32-48. doi:
10.1044/jshr.2701.32.

Braga RM, Wilson LR, Sharp DJ, Wise RJS, Leech R (2013) Separable networks for top-down
attention to auditory non-spatial and visuospatial modalities. NeuroImage 74:77-86. doi:
https://doi.org/10.1016/j.neuroimage.2013.02.023.

Chalehchaleh A, Winchester MM, Di Liberto GM (2025) Robust assessment of the cortical
encoding of word-level expectations using the temporal response function. Journal of
Neural Engineering 22:016004. doi: 10.1088/1741-2552/ada30a.

Cohen SS, Parra LC (2016) Memorable audiovisual narratives synchronize sensory and
supramodal neural responses. eNeuro 3:e0203. doi: 10.1523/ENEURO.0203-16.2016.

Corbetta M, Shulman GL (2002) Control of goal-directed and stimulus-driven attention in the
brain. Nature Reviews Neuroscience 3:201-215. doi: 10.1038/nrn755.

Crosse MJ, Di Liberto GM, Bednar A, Lalor EC (2016) The multivariate temporal response
function (mTRF) toolbox: A MATLAB toolbox for relating neural signals to continuous
stimuli. Frontiers in Human Neuroscience 10:604. doi: 10.3389/fnhum.2016.00604.

Crosse MJ, Zuk NJ, Di Liberto GM, Nidiffer AR, Molholm S, Lalor EC (2021) Linear modeling of
neurophysiological responses to speech and other continuous stimuli: Methodological
considerations for applied research. Frontiers in Neuroscience 15. doi:
10.3389/fnins.2021.705621.

Daube C, Ince RAA, Gross J (2019) Simple acoustic features can explain phoneme-based
predictions of cortical responses to speech. Current Biology 29:1924-1937. doi:
10.1016/j.cub.2019.04.067.

Davis MH, Johnsrude IS (2003) Hierarchical processing in spoken language comprehension. The
Journal of Neuroscience 23:3423-3431. doi: 10.1523/JNEUROSCI.23-08-03423.2003.

de Boer J, Krumbholz K (2018) Auditory Attention Causes Gain Enhancement and Frequency
Sharpening at Successive Stages of Cortical Processing—Evidence from Human
Electroencephalography. Journal of Cognitive Neuroscience 30:785-798. doi:
10.1162/jocn_a_01245.

Di Liberto Giovanni M, O'Sullivan James A, Lalor Edmund C (2015) Low-frequency cortical
entrainment to speech reflects phoneme-level processing. Current Biology 25:2457-2465.
doi: 10.1016/j.cub.2015.08.030.

Ding N, Simon JZ (2012) Emergence of neural encoding of auditory objects while listening to
competing speakers. Proceedings of the National Academy of Sciences 109:11854-
11859. doi: 10.1073/pnas.1205381109.

Ding N, Simon JZ (2013) Adaptive temporal encoding leads to a background-insensitive cortical
representation of speech. The Journal of Neuroscience 33:5728-5735. doi:
10.1523/JNEUROSCI.5297-12.2013.

Ding N, Chatterjee M, Simon JZ (2014) Robust cortical entrainment to the speech envelope relies
on the spectro-temporal fine structure. NeuroImage 88:41-46. doi:
10.1016/j.neuroimage.2013.10.054.

Dmochowski JP, Sajda P, Dias J, Parra LC (2012) Correlated components of ongoing EEG point
to emotionally laden attention – a possible marker of engagement? Frontiers in Human
Neuroscience 6:Article 112. doi: 10.3389/fnhum.2012.00112.

Donoghue T, Schaworonkow N, Voytek B (2022) Methodological considerations for studying
neural oscillations. European Journal of Neuroscience 55:3502-3527. doi:
https://doi.org/10.1111/ejn.15361.

Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, Noto T, Lara AH, Wallis JD,
Knight RT, Shestyuk A, Voytek B (2020) Parameterizing neural power spectra into periodic
and aperiodic components. Nature Neuroscience 23:1655-1665. doi: 10.1038/s41593-
020-00744-x.

Eastwood JD, Frischen A, Fenske MJ, Smilek D (2012) The Unengaged Mind: Defining Boredom
in Terms of Attention. Perspectives on Psychological Science 7:482-495. doi.

Fiedler L, Wöstmann M, Herbst SK, Obleser J (2019) Late cortical tracking of ignored speech
facilitates neural selectivity in acoustically challenging conditions. Neuroimage 186:33-42.
doi: 10.1016/j.neuroimage.2018.10.057.

Fiedler L, Wöstmann M, Graversen C, Brandmeyer A, Lunner T, Obleser J (2017) Single-channel
in-ear-EEG detects the focus of auditory attention to concurrent tone streams and mixed
speech. Journal of Neural Engineering 14:036020. doi: 10.1088/1741-2552/aa66dd.

Forster S, Lavie N (2009) Harnessing the wandering mind: The role of perceptual load. Cognition
111:345-355. doi: https://doi.org/10.1016/j.cognition.2009.02.006.

Fuglsang SA, Märcher-Rørsted J, Dau T, Hjortkjær J (2020) Effects of Sensorineural Hearing
Loss on Cortical Synchronization to Competing Speech during Selective Attention. The
Journal of Neuroscience 40:2562-2572. doi.

Gillis M, Vanthornhout J, Simon JZ, Francart T, Brodbeck C (2021) Neural Markers of Speech
Comprehension: Measuring EEG Tracking of Linguistic Speech Representations,
Controlling the Speech Acoustics. The Journal of Neuroscience 41:10316. doi:
10.1523/JNEUROSCI.0812-21.2021.

Hambrook DA, Tata MS (2019) The effects of distractor set-size on neural tracking of attended
speech. Brain and Language 190:1-9. doi: https://doi.org/10.1016/j.bandl.2018.12.005.

Heffernan E, Coulson NS, Henshaw H, Barry JG, Ferguson MA (2016) Understanding the
psychosocial experiences of adults with mild-moderate hearing loss: An application of
Leventhal's self-regulatory model. International Journal of Audiology 55:S3-S12. doi.

Herrmann B (2023a) The perception of artificial-intelligence (AI) based synthesized speech in
younger and older adults. International Journal of Speech Technology 26:395-415. doi:
10.1007/s10772-023-10027-y.

Herrmann B (2023b) Neural responses to repeated noise structure in sounds are invariant to
temporal interruptions. Auditory Perception & Cognition 6:330-352. doi.

Herrmann B (2025a) Enhanced neural speech tracking through noise indicates stochastic
resonance in humans. eLife 13:RP100830. doi: 10.7554/eLife.100830.3.

Herrmann B (2025b) Age-related differences in the impact of background noise on neural speech
tracking. Neurobiology of Aging 153:10-20. doi: 10.1101/2025.01.09.632269.

Herrmann B (2025c) Language-agnostic, automated assessment of listeners' speech recall using
large language models. Trends in Hearing 29:1-22. doi: 10.1177/23312165251347131.

Herrmann B, Johnsrude IS (2020) A model of listening engagement (MoLE). Hearing Research
397:108016. doi: 10.31234/osf.io/y3cr9.

Herrmann B, Buckland C, Johnsrude IS (2019) Neural signatures of temporal regularity
processing in sounds differ between younger and older adults. Neurobiology of Aging
83:73-85. doi: https://doi.org/10.1101/522375

Herrmann B, Maess B, Johnsrude IS (2022) A neural signature of regularity in sound is reduced
in older adults. Neurobiology of Aging 109:1-10. doi:
10.1016/j.neurobiolaging.2021.09.011.
Hertrich I, Dietrich S, Trouvain J, Moos A, Ackermann H (2012) Magnetic brain activity phase-
locked to the envelope, the syllable onsets, and the fundamental frequency of a perceived
speech signal. Psychophysiology 49:322-334. doi: 10.1111/j.1469-8986.2011.01314.x.
Holm S (1979) A Simple Sequentially Rejective Multiple Test Procedure. Scandinavian Journal of
Statistics 6:65-70. doi: http://www.jstor.org/stable/4615733.
Ibaceta M, Holman D, Niven K (2024) Mind-wandering contents and characteristics: an
exploratory study comparing between work and non-work contexts. European Journal of
Work and Organizational Psychology 33:503-516. doi: 10.1080/1359432X.2024.2328373.
Irsik VC, Johnsrude IS, Herrmann B (2022) Neural activity during story listening is synchronized
across individuals despite acoustic masking. Journal of Cognitive Neuroscience 34:933-
950. doi: 10.1162/jocn_a_01842.
Irsik VC, Almanaseer A, Johnsrude IS, Herrmann B (2021) Cortical responses to the amplitude
envelopes of sounds change with age. The Journal of Neuroscience 41:5045-5055. doi:
10.1523/JNEUROSCI.2715-20.2021.
Joon Kim Y, Grabowecky M, Paller KA, Muthu K, Suzuki S (2007) Attention induces
synchronization-based response gain in steady-state visual evoked potentials. Nature
Neuroscience 10:117-125. doi: 10.1038/nn1821.
Keitel C, Thut G, Gross J (2017) Visual cortex responses reflect temporal structure of continuous
quasi-rhythmic sensory stimulation. NeuroImage 146:58-70. doi.
Keitel C, Keitel A, Benwell CSY, Daube C, Thut G, Gross J (2019) Stimulus-Driven Brain Rhythms
within the Alpha Band: The Attentional-Modulation Conundrum. The Journal of
Neuroscience 39:3119. doi: 10.1523/JNEUROSCI.1633-18.2019.
Kerlin JR, Shahin AJ, Miller LM (2010) Attentional Gain Control of Ongoing Cortical Speech
Representations in a "Cocktail Party". The Journal of Neuroscience 30:620-628. doi.
Kraus F, Tune S, Ruhe A, Obleser J, Wöstmann M (2021) Unilateral acoustic degradation delays
attentional separation of competing speech. Trends in Hearing 25:23312165211013242.
doi: 10.1177/23312165211013242.
Lachaux J-P, Rodriguez E, Martinerie J, Varela FJ (1999) Measuring Phase Synchrony in Brain
Signals. Human Brain Mapping 8:194-208. doi.

Lalor EC, Foxe JJ (2010) Neural responses to uninterrupted natural speech can be extracted with
precise temporal resolution. European Journal of Neuroscience 31:189-193. doi:
10.1111/j.1460-9568.2009.07055.x.

Lanting CP, Briley PM, Summer CJ, Krumbholz K (2013) Mechanisms of adaptation in human
auditory cortex. Journal of Neurophysiology 110:973-983. doi.

Lavie N (2005) Distracted and confused?: Selective attention under load. Trends in Cognitive
Sciences 9:75-82. doi.

Lesenfants D, Vanthornhout J, Verschueren E, Decruy L, Francart T (2019) Predicting individual
speech intelligibility from the cortical tracking of acoustic- and phonetic-level speech
representations. Hearing Research 380:1-9. doi: 10.1016/j.heares.2019.05.006.

Luck SJ (2014) An Introduction to the Event-Related Potential Technique. USA: MIT Press.

Makeig S, Bell AJ, Jung T-P, Sejnowski TJ (1995) Independent component analysis of
electroencephalographic data. In: Advances in Neural Information Processing Systems
(Touretzky D, Mozer M, Hasselmo M, eds), pp 145-151. Cambridge, MA, USA: MIT Press.

Mathiesen SL, Van Hedger SC, Irsik VC, Bain MM, Johnsrude IS, Herrmann B (2024) Exploring
age differences in absorption and enjoyment during story listening. Psychology
International 6:667-684. doi: 10.3390/psycholint6020041.

McDermott Josh H, Simoncelli Eero P (2011) Sound texture perception via statistics of the
auditory periphery: Evidence from sound synthesis. Neuron 71:926-940. doi:
10.1016/j.neuron.2011.06.032.

McDonnell MD, Abbott D (2009) What Is stochastic resonance? Definitions, misconceptions,
debates, and its relevance to biology. PLOS Computational Biology 5:e1000348. doi:
10.1371/journal.pcbi.1000348.

McDonnell MD, Ward LM (2011) The benefits of noise in neural systems: bridging theory and
experiment. Nature Reviews Neuroscience 12:415-425. doi: 10.1038/nrn3061.

McNeice Z, Tomlin D, Timmer B, Short CE, Nixon G, Galvin K (2024) A scoping review exploring
how adults self-describe and communicate about the listening difficulties they experience.
International Journal of Audiology 63:163-170. doi: 10.1080/14992027.2023.2245136.

Moss F, Ward LM, Sannita WG (2004) Stochastic resonance and sensory information processing:
a tutorial and review of application. Clinical Neurophysiology 115:267-281. doi:
10.1016/j.clinph.2003.09.014.

Motala A, Johnsrude IS, Herrmann B (2024) A longitudinal framework to describe the relation
between age-related hearing loss and social isolation. Trends in Hearing 28:1-18. doi:
10.1177/23312165241236041.

Müller MM, Hillyard S (2000) Concurrent recording of steady-state and transient event-related
potentials as indices of visual-spatial selective attention. Clinical Neurophysiology
111:1544-1552. doi: https://doi.org/10.1016/S1388-2457(00)00371-0.

Näätänen R, Picton TW (1987) The N1 wave of the human electric and magnetic response to
sound: a review and an analysis of the component structure. Psychophysiology 24:375-
425. doi: 10.1111/j.1469-8986.1987.tb00311.x.

Okamoto H, Stracke H, Bermudez P, Pantev C (2011) Sound Processing Hierarchy within Human
Auditory Cortex. Journal of Cognitive Neuroscience 23:1855-1863. doi.

Oostenveld R, Fries P, Maris E, Schoffelen JM (2011) FieldTrip: Open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. Computational
Intelligence and Neuroscience 2011:Article ID 156869. doi: 10.1155/2011/156869.

OpenAI et al. (2023) GPT-4 Technical Report. arXiv. doi: 10.48550/arXiv.2303.08774.

Orf M, Hannemann R, Obleser J (2025) Does amplitude compression help or hinder attentional
neural speech tracking? The Journal of Neuroscience 45:e0238242024. doi:
10.1523/JNEUROSCI.0238-24.2024.

Panela RA, Copelli F, Herrmann B (2024) Reliability and generalizability of neural speech tracking
in younger and older adults. Neurobiology of Aging 134:165-180. doi:
10.1016/j.neurobiolaging.2023.11.007.

Parbery-Clark A, Marmel F, Bair J, Kraus N (2011) What subcortical–cortical relationships tell us
about processing speech in noise. European Journal of Neuroscience 33:549-557. doi:
10.1111/j.1460-9568.2010.07546.x.

Peelle JE (2018) Listening Effort: How the Cognitive Consequences of Acoustic Challenge Are
Reflected in Brain and Behavior. Ear & Hearing 39:204-214. doi:
10.1097/aud.0000000000000494.

Pichora-Fuller MK, Kramer SE, Eckert MA, Edwards B, Hornsby BWY, Humes LE, Lemke U,
Lunner T, Matthen M, Mackersie CL, Naylor G, Phillips NA, Richter M, Rudner M,
Sommers MS, Tremblay KL, Wingfield A (2016) Hearing impairment and cognitive energy:
The framework for understanding effortful listening (FUEL). Ear & Hearing 37 Suppl 1:5S-
27S. doi: 10.1097/aud.0000000000000312.

Picton TW, John SM, Dimitrijevic A, Purcell DW (2003) Human auditory steady-state responses.
International Journal of Audiology 42:177-219. doi: 10.3109/14992020309101316.

Power AJ, Foxe JJ, Forde E-J, Reilly RB, Lalor EC (2012) At what time is the cocktail party? A
late locus of selective attention to natural speech. European Journal of Neuroscience
35:1497-1503. doi: https://doi.org/10.1111/j.1460-9568.2012.08060.x.

Presacco A, Simon JZ, Anderson S (2016) Evidence of degraded representation of speech in
noise, in the aging midbrain and cortex. Journal of Neurophysiology 116:2346-2355. doi:
10.1152/jn.00372.2016.

Puschmann S, Regev M, Fakhar K, Zatorre RJ, Thiel CM (2024) Attention-Driven Modulation of
Auditory Cortex Activity during Selective Listening in a Multispeaker Setting. The Journal
of Neuroscience 44:e1157232023. doi: 10.1523/JNEUROSCI.1157-23.2023.

Ritz H, Wild CJ, Johnsrude IS (2022) Parametric cognitive load reveals hidden costs in the neural
processing of perfectly intelligible degraded speech. The Journal of Neuroscience
42:4619-4628. doi: 10.1523/JNEUROSCI.1777-21.2022.

Ross B, Picton TW, Pantev C (2002) Temporal integration in the human auditory cortex as
represented by the development of the steady-state magnetic field. Hearing Research
165:68-84. doi.

Ruhnau P, Herrmann B, Schröger E (2012) Finding the right control: The mismatch negativity
under investigation. Clinical Neurophysiology 123:507-512. doi:
10.1016/j.clinph.2011.07.035.

Safati AB, Carr TH, Lowe CJ, Smilek D (2024) Mind wandering on command. Frontiers in
Psychology Volume 15 - 2024. doi.

Sarant J, Lemke U, Giroud N, Scherpiet S, Weinstein B (2024) Promoting hearing and cognitive
health in audiologic rehabilitation for the well-being of older adults. International Journal
of Audiology 63:761-771. doi: 10.1080/14992027.2023.2260099.

Seli P, Risko EF, Smilek D (2016a) On the Necessity of Distinguishing Between Unintentional
and Intentional Mind Wandering. Psychological Science 27:685-691. doi:
10.1177/0956797616634068.

Seli P, Risko EF, Smilek D, Schacter DL (2016b) Mind-Wandering With and Without Intention.
Trends in Cognitive Sciences 20:605-617. doi: 10.1016/j.tics.2016.05.010.

Shinn-Cunningham BG (2008) Object-based auditory and visual attention. Trends in Cognitive
Sciences 12:182-186. doi.

Shukla B, Bidelman GM (2021) Enhanced brainstem phase-locking in low-level noise reveals
stochastic resonance in the frequency-following response (FFR). Brain Research
1771:147643. doi: 10.1016/j.brainres.2021.147643.

Smallwood J, Schooler JW (2006) The Restless Mind. Psychological Bulletin 132:946-958. doi.

Smallwood J, Andrews-Hanna J (2013) Not all minds that wander are lost: the importance of a
balanced perspective on the mind-wandering state. Frontiers in Psychology Volume 4 -
2013. doi.

Stawarczyk D, Majerus S, Maj M, Van der Linden M, D'Argembeau A (2011) Mind-wandering:
Phenomenology and function as assessed with a novel experience sampling method. Acta
Psychologica 136:370-381. doi: https://doi.org/10.1016/j.actpsy.2011.01.002.

Stein RB, Gossen ER, Jones KE (2005) Neuronal variability: noise or part of the signal? Nature
Reviews Neuroscience 6:389-397. doi: 10.1038/nrn1668.

Steindorf L, Pink S, Rummel J, Smallwood J (2023) When there is noise on Sherlock Holmes:
mind wandering increases with perceptual processing difficulty during reading and
listening. Cognitive Research: Principles and Implications 8:31. doi.

Stocks NG (2000) Suprathreshold stochastic resonance in multilevel threshold systems. Physical
Review Letters 84:2310-2313. doi: 10.1103/PhysRevLett.84.2310.

Straetmans L, Adiloglu K, Debener S (2024) Neural speech tracking and auditory attention
decoding in everyday life. Frontiers in Human Neuroscience Volume 18 - 2024. doi.

Synigal SR, Anderson AJ, Lalor EC (2023) Electrophysiological indices of hierarchical speech
processing differentially reflect the comprehension of speech in noise. BioRxiv. doi:
10.1101/2023.03.30.534927.

Tune S, Alavash M, Fiedler L, Obleser J (2021) Neural attentional-filter mechanisms of listening
success in middle-aged and older individuals. Nature Communications 12:4533. doi:
10.1038/s41467-021-24771-9.

Vanthornhout J, Decruy L, Francart T (2019) Effect of Task and Attention on Neural Tracking of
Speech. Frontiers in Neuroscience 13. doi: 10.3389/fnins.2019.00977.

Vanthornhout J, Decruy L, Wouters J, Simon JZ, Francart T (2018) Speech intelligibility predicted
from neural entrainment of the speech envelope. Journal of the Association for Research
in Otolaryngology 19:181-191. doi: 10.1007/s10162-018-0654-z.

Verschueren E, Gillis M, Decruy L, Vanthornhout J, Francart T (2022) Speech Understanding
Oppositely Affects Acoustic and Linguistic Neural Tracking in a Speech Rate Manipulation
Paradigm. The Journal of Neuroscience 42:7442. doi: 10.1523/JNEUROSCI.0259-
22.2022.

Villena-González M, Cosmelli D (2020) Chapter 5 - Imagination and mind wandering: two sides
of the same coin? A brain dynamics perspective. In: Creativity and the Wandering Mind
(Preiss DD, Cosmelli D, Kaufman JC, eds), pp 93-120: Academic Press.

Ward LM, Neiman A, Moss F (2002) Stochastic resonance in psychophysics and in animal
behavior. Biological Cybernetics 87:91-101. doi: 10.1007/s00422-002-0328-z.

Ward LM, MacLean SE, Kirschner A (2010) Stochastic resonance modulates neural
synchronization within and between cortical sources. PLoS ONE 5:e14371. doi:
10.1371/journal.pone.0014371.

Weinstein Y (2018) Mind-wandering, how do I measure thee with probes? Let me count the ways.
Behavior Research Methods 50:642-661. doi: 10.3758/s13428-017-0891-9.

Wilson RH, McArdle RA, Watts KL, Smith SL (2012) The revised speech perception in noise test
(R-SPIN) in a multiple signal-to-noise ratio paradigm. Journal of the American Academy
of Audiology 23:590-605. doi: 10.3766/jaaa.23.7.9.

Xie Z, Brodbeck C, Chandrasekaran B (2023) Cortical Tracking of Continuous Speech Under
Bimodal Divided Attention. Neurobiology of Language 4:318-343. doi:
10.1162/nol_a_00100.

Yasmin S, Irsik VC, Johnsrude IS, Herrmann B (2023) The effects of speech masking on neural
tracking of acoustic and semantic features of natural speech. Neuropsychologia
186:108584. doi: 10.1016/j.neuropsychologia.2023.108584.

Zuk NJ, Murphy JW, Reilly RB, Lalor EC (2021) Envelope reconstruction of speech and music
highlights stronger tracking of speech at low frequencies. PLOS Computational Biology
17:e1009358. doi: 10.1371/journal.pcbi.1009358.

Figure Captions

Figure 1: Results for Experiment 1. A: Comprehension scores (left), gist ratings (right) across different
speech-clarity conditions. **B:** Temporal response functions (TRF) waveforms. **C:** EEG prediction accuracy
and corresponding topographies. **D:** P1-N1 (left) and P2-N1 (right) amplitude and topographies for each
speech-clarity condition. Error bars denote the standard error of the mean. *p < .05, **p < .01, ***p < .001
(Holm-corrected where applicable).

Figure 2: Results for Experiment 2. A: Comprehension scores (left) and gist ratings (right) across three
speech-clarity levels during 'attend' blocks. **B:** Detection accuracy and reaction times (RT) for the visual 1-
back task, and ratings of the ability to ignore the spoken stories during visual distraction blocks. **C:** ITPC
showing peaks at 2 Hz and harmonics, reflecting responses to the visual stimuli occurring at a 2-Hz
stimulation rate. **D:** Peak ITPC (2-10 Hz averaged) for three speech-clarity conditions separated by
attention blocks. **E:** Temporal response functions (TRFs) for 'attend' (solid) and 'ignore' (dashed) conditions
for the different speech-clarity conditions. **F:** Mean EEG prediction accuracy and corresponding
topographies. **G:** Mean P1-N1 amplitudes and corresponding topographies. **H:** Mean P2-N1 amplitudes
and corresponding topographies. Error bars denote the standard error of the mean. *p < .05, **p < .01, ***p
< .001 (Holm-corrected where applicable).

Figure 3: Results for Experiment 3. A: Comprehension scores (left) and gist ratings (right) across three
speech-clarity conditions during attentive listening blocks. **B:** Self-evaluations of the ability 'to ignore the
spoken stories', 'to be able to wander off', 'to be able to visualize the given scenery', 'to be able to retell the
spoken stories' during imagination blocks. **C:** Temporal response functions (TRFs) during Attend (solid)
and Ignore (dashed) across Speech Clarity. **D:** EEG prediction accuracy and corresponding topographies.
E: P1-N1 amplitudes and corresponding topographies. **F:** P2-N1 amplitudes and corresponding
topographies. Error bars denote the standard error of the mean. *p < .05, **p < .01, ***p < .001 (Holm-
corrected where applicable).

35