Intersection of transient cell states with stable cell types in hippocampus

Jack A Olmstead[1,2,4]*, Lauren E King[3,4], Brenda L Bloodgood[3,4]*

[1] *Neurosciences Graduate Program*

[2] *Medical Scientist Training Program*

[3] *Department of Neurobiology, School of Biological Sciences*

[4] *UC San Diego School of Medicine, La Jolla, CA, United States*

* Corresponding Authors, jolmstea [at] health.ucsd.edu, blbloodgood [at] ucsd.edu

Abstract

The transcriptome of a brain cell encodes both its stable identity and its dynamic responses to environmental stimuli. While significant progress has been made in categorizing cell types within the brain, deciphering to what extent transcriptional identity and transcriptional state are related remains a major technical and conceptual challenge. Here, we present a single-nucleus RNA-sequencing atlas of the mouse hippocampus spanning physiological and pathological stimuli and multiple circadian phases, enabling unified analysis of activity-, circadian-, and cell-type-dependent transcriptional programs. Taxonomically assigned cell types are largely stable despite the induction of different activity states, with a notable exception in the dentate gyrus. Activity and circadian rhythm each drive robust, largely nonoverlapping transcriptional responses, with convergent regulation on genes involved in specific pathways, including endocannabinoid signaling, excitability, and chromatin remodeling. These results underscore the necessity of integrating cell-type taxonomy with transcriptional state to capture how diverse cell types respond to experience.

Introduction

Activity-dependent gene expression is a fundamental mechanism that neural cell types use to respond to experience and coordinate adaptive cellular responses. For canonical immediate-early genes (IEGs) such as *Fos*, *Egr1*, and *Npas4*, transcription begins to rise within minutes and peaks within an hour, highlighting the transient kinetics of these programs[1–6]. Large-scale transcriptomic atlases have provided precise molecular definitions of cell types in the brain[7,8], but interpretation of IEG expression within these foundational datasets has been fraught with known artifacts produced during cell capture, a process which itself induces IEGs[9,10]. A concerted effort is therefore necessary to characterize the intersection of activity states with a standardized taxonomy of cell types.

The hippocampus is a central structure for learning and episodic memory[11], and displays strong cell-type-specific network activity patterns during exploration and rest[12–17]. Behavioral states that drive plasticity—such as exploration during arousal and sharp-wave ripples during rest or sleep—predominate at different points in the circadian cycle, positioning the hippocampus as a model system for developing conceptual approaches to the integration of transcriptomic cell types and multidimensional cell states. Prior bulk RNA-seq studies in mice have shown circadian cycling of clock genes in the hippocampus[18,19], but no single-cell data exist to determine how neurons and glia engage these programs. Circadian phase influences excitability[20–23], seizure susceptibility[24,25], and memory performance[18,26], making the interaction between activity-dependent and circadian-dependent transcription biologically important, yet poorly characterized. To date, transcriptional rhythms across the circadian cycle have not been described in the hippocampus using single-cell RNA-seq approaches, leaving open how different cell types entrain their gene expression to this important biological variable.

In this study, we leverage state-of-the-art cell-type resources and a multifactorial experimental design to create an atlas of hippocampal cell states across activity conditions and circadian timepoints, which we call the Activity and Circadian

Transcriptomes Dependent on Physiological and Pathological activity (ACT-DEPP)

dataset. With this resource, we show that enriched-environment (EE)- and seizure-

induced neuronal activity engage qualitatively different transcriptomic programs within

all cell types of the hippocampus. We find that within the Allen Brain Cell taxonomy of

cell types, there is a collapse of cell type and activity state into a unitary identity among

a subset of dentate granule cells. We describe a novel suite of marker genes that are

transiently induced by activity in subpopulations of excitatory, inhibitory, and glial cells

in the hippocampus. We report a gradient of IEG induction along the dorsal-ventral axis

of CA1 in response to EE exposure. Finally, we show here that circadian rhythm

coordinates widespread transcriptomic programs in both hippocampal neurons and glia

at baseline. By comparing EE exposure to baseline states across the diurnal cycle, we

systematically describe how activity- and circadian-dependent transcriptional programs

intersect within hippocampal cell types and show that these variables influence

endocannabinoid signaling, cell excitability, and chromatin remodeling.

By integrating transcriptional signatures of activity and circadian time into a taxonomy

of cell types within the ACT-DEPP dataset, we provide both a conceptual framework for

understanding state-dependent gene regulation and a resource for probing how activity

patterns shape hippocampal plasticity across the circadian cycle.

71 Results

72 Generation of the ACT-DEPP dataset

Single-nucleus RNA sequencing (snRNA-seq) using SPLiT-seq[27] was performed on

hippocampi from animals exposed to a standard environment (SE); an enriched

environment (EE) for 30 minutes or 6 hours; or kainic acid-induced seizures (KA) for 30

minutes or 6 hours. The resulting dataset includes 93,066 nuclei from 48 biological

replicates, capturing gene expression in baseline, physiological, and pathological activity

conditions across circadian timepoints (**Fig. 1A-B, Supp. Fig. 1**).

Quality-controlled nuclei were annotated using MapMyCells and integrated with the Allen

Brain Cell (ABC) taxonomy, assigning each nucleus to hierarchical ranks of Class >

Subclass > Supertype > Cluster (**Fig. 1C inset;** ref [7]). Note: the Subclass rank generally

corresponds to conventionally understood cell types within the hippocampus, e.g.,

dentate granule cells, CA1, PV-expressing basket cells, etc. The UMAP embedding

recapitulated relations of canonical hippocampal cell populations at the Subclass level

(**Fig. 1C**), with additional clear transcriptional distinctions for both KA and EE conditions

within neuronal and glial Subclasses (**Fig. 1D**).

To confirm that cell-type annotation was independent of activity-induced transcriptional

states, a set of known activity-regulated genes[2] (ARGs) were removed, and annotations

were remapped. Cell-type re-annotations at the Class and Subclass rank were highly

stable (**Fig. 1E**-**F**), with minimal reassignment even at the lower ABC taxonomic ranks of

Subclass and Cluster (**Fig. 1G-H**). The ACT-DEPP dataset thus robustly registers

hippocampal cell types to the ABC taxonomy and reliably distinguishes transcriptional

states driven by physiological and pathological stimuli.

Figure 1

Figure 1. Generation of the ACT-DEPP dataset.
(A) Experimental schematic of generation of the ACT-DEPP dataset.
(B) Distribution of nuclei number, experimental conditions, and sample metadata among major
subclasses. Some neocortical and subiculum Subclasses appearing in Figure 1D have been
omitted from this panel, as they were not analyzed in any further downstream analyses.
(C) UMAP embedding of ACT-DEPP nuclei colored by subclass. Inset: Single-nucleus
transcriptomes were mapped to the Allen Brain Cell (ABC) whole-brain atlas, which assigns cells
in a hierarchical taxonomy to one class, subclass, supertype, and cluster.
(D) As in (C), but colored by activity condition.
(E-H) Sankey flow diagrams showing changes in ABC-taxon identity for nuclei before (+) and
after (-) removing ARGs from single-nucleus transcriptomes. (G) and (H) show the effect at lower
ABC taxa for the two largest neuronal subclasses, CA1 and DG.
CA1-ProS, cornu ammonis 1-prosubiculum; L2/3 PPP, layer 2/3 pre-, pro-, parasubiculum; NP
Sub, near-projecting subiculum; CT Sub, corticothalamic subiculum; L6 Ent, layer 6 entorhinal;
DG, dentate gyrus granule cells; Sub-ProS, Subiculum-prosubiculum; Micro, microglia; Astro,
astroglia; Oligo, oligodendrocyte; OPC, oligodendrocyte precursor cells; Epen, ependymal cell;
NGF, neurogliaform cell; IMN, DG immature neuron; SST, GABAergic somatostatin-expressing
interneuron; PV, GABAergic parvalbumin-expressing interneuron; GABAergic vasopressin
internal peptide-expressing interneuron; Sncg, GABAergic Sncg-expressing interneuron; Lhx6,
GABAergic Lhx6-expressing interneuron

The activity-dependent transcriptome is stimulus-specific and obscures cell
identity in DG

Seizures have been used extensively to study ARG expression in the past, but direct

comparisons to naturalistic stimuli are limited[28,29]. Moreover, the near-complete

segregation of KA populations within some Subclasses in our uniform manifold

approximation and projection (UMAP) plots led us to hypothesize that seizures constitute

a categorically different transcriptional state from physiological manipulations. We

therefore analyzed differentially expressed genes (DEGs) in CA1 and DG, the most

abundant excitatory neuron Subclasses in our samples, at 30 minutes and 6 hours. KA

induced an order of magnitude more DEGs than EE at both time points. Approximately

one-third of EE-induced DEGs at 30 minutes were distinct from those induced by KA

(**Fig. 2A**). At 6 hours, CA1 responses were fully shared between EE and KA, while DG

maintained a unique DEG set specific to EE (**Fig. 2B**). Thus, novel experiences and

seizures trigger categorically distinct transcriptional responses—with respect to both

magnitude and specific genes—in these hippocampal subregions.

To look more closely at the effects of activity on different excitatory-neuron populations,

we re-embedded UMAPs and color-coded nuclei by condition or Subclass/Supertype

(**Fig. 2C**). Supertypes were clearly segregated within Subclass clusters, and strikingly,

distinct Supertypes emerged as seizure-specific populations. This effect was particularly

pronounced in DG, identified by its dramatically altered Supertype distributions between

activity conditions (**Fig. 2D**). In DG, nearly all KA cells clustered within Supertype 138-3,

with notable dissociation into Clusters 508-3 (KA 30m; green) and 509-4 (KA 6hr; purple,

Fig. 2E). Analysis of Cluster-defining genes revealed six marker genes strongly

associated with seizure timing and cluster identity (**Fig. 2F-G**): *Egr2, Egr4, Cenpa,*

Bhlhe41, Lct, Npy, the latter four not characterized as canonical ARGs. Thus, Supertype

DG 138-3 labels recently active granule cells, with additional discrimination of early and

late transcriptional responses to activity at the Cluster level. This would likely not be

detectable without both EE and KA conditions, given the sparsity of DG activation during

naturalistic activity[30–34].

Figure 2

Figure 2. The activity-dependent transcriptome is stimulus-specific and obscures cell identity in DG.

(A-B) Venn diagrams of shared DEGs between KA or EE vs SE at 30 min (A) or 6 hr (B) in CA1 (n_{EE30m} = 3329 nuclei, n_{SE} = 1907 nuclei, n_{KA30m} = 354 nuclei) and DG (n_{EE30m} = 7119 nuclei, n_{SE} = 6754 nuclei, n_{KA30m} = 961 nuclei). DEGs were defined as genes with absolute log2(FC) > 1.5 and FDR < 0.05 using pseudobulk expression (Methods).

(C) Re-embedded UMAPs of excitatory neuronal nuclei colored by Subclass (left, top), activity condition (left, bottom), or Supertype (right). Cross-reference of the UMAPs reveals KA-specific populations marked by red arrows.

(D) Distribution of Supertypes within each subclass across different conditions. Colors within each column represent proportions of different Supertypes within the labeled Subclass.

(E) As in (D), but for DG Clusters only.

(F-G) Single-nucleus, SC-transformed expression (Hafemeister & Satija 2019) of marker genes for DG Cluster identity, plotted by condition (F) or Cluster (G)

Discovery of cell-type-specific ERGs induced by novel experience

To understand how different cell populations respond to naturalistic activity, we compared early-response gene (ERG) expression across major groups of hippocampal Subclasses (DEGs were categorized as either ERGs or LRGs based on their peak-expression timing; see Methods and ref [10]). Both excitatory and inhibitory neurons share broad upregulation of canonical ERGs such as *AP1-*, *Egr-*, *Nr4a-*, and *Dusp*-family genes, *Arc*, and *Npas4* across nearly all Subclasses examined (**Fig. 3A, E**); these genes unsurprisingly exhibit some of the strongest induction genome-wide across nearly all Subclasses, including astrocytes and microglia (**Fig. 3B, F, I**).

Next, we were interested in determining which upregulated ERGs exhibited specific activity-dependent induction in one or a few Subclasses. We calculated a tau specificity score previously developed for bulk RNA-seq count data[35] to filter for specific changes in expression within EE 30 min vs SE across Subclasses (Methods). When doing so, we observe that canonical ERGs have some of the lowest cell-type specificity (**Fig. 3C, G, J**). Meanwhile, screening for high-tau ERGs reveals genes whose induction is specific to one or a few related cell types. For example, *Hs3st2* and *Rcan2* are both genes whose expression is (1) specific to spatially restricted subiculum Subclasses, (2) acutely upregulated following activity, and (3) decaying to baseline expression at 6 hours (**Fig. 3D**). Likewise, *Tac1* is an exemplar activity-dependent marker gene among PV-

expressing GABAergic cells and *Crhbp* in medial-ganglionic-eminence-derived

interneurons broadly (**Fig. 3H**). Finally, we make special note of the astrocytes, surprising

in their robust Subclass-specific response to acute EE exposure (**Fig. 3I-J**). Among the

DEGs, *Map3k19* and *Slco1c1* stand out as genes highly specific to activated astrocytes

(**Fig. 3K**). Thus, comparison of activity-dependent ERG expression following novel

experience yields novel marker genes with highly cell-type-specific induction.

Figure 3

Figure 3. Discovery of cell-type-specific ERGs induced by novel experience.
(A) Change in expression in EE30m vs SE for all ERGs in excitatory neuronal subclasses; gene columns are ordered by hierarchical clustering, and genes encoding transcription factors are annotated with black ticks above heatmaps. For every Subclass, we performed pseudobulk DEG analysis at EE 30 min vs SE and EE 6 hr vs SE. The heatmap contains all DEGs—ascertained from any Subclass listed—that were classified as ERGs. ERGs are defined as DEGs whose fold-change in EE vs SE was higher after 30 min instead of 6 hr (Methods).
(B) Volcano plot of DEGs in CA1-ProS nuclei at EE30m (n = 3329 nuclei) vs SE (n = 1907 nuclei).
(C) Heatmap of upregulated ERGs in excitatory neuronal Subclasses, ordered by Subclass-specific induction (tau score; Methods). Canonical ERGs are annotated in black text, while novel example Subclass-specific genes are annotated in red text.
(D) Violin plots of SC-transformed expression for example genes with high tau scores.
(E-H) as in (A-D) but for inhibitory neuronal subclasses.
(I-K) as in (A-D) but for immature neurons and glial subclasses.

Distribution of IEG expression across CA1-ProS Supertypes

Because CA1-ProS neurons were the largest Subclass in our dataset, we used the opportunity to explore experience-induced activity responses within sub-populations of these principal cells. First, we characterized activity in the entire population by adapting a previously used method[10] to classify individual neurons as "active" or "inactive" based on thresholded induction of immediate early genes (**Fig. 4A**; Methods). Crucially, our library preparation captured only nuclear transcripts before they were exported to the cytoplasm, increasing the likelihood that active cells are classified as such specifically in response to the stimulus[3,9]. At baseline (SE), only 8% of cells were active. After 30 minutes of EE, activation increased to 33% of CA1 cells, then declined to 20% by 6 hours. KA seizure robustly activated nearly all cells at 30 minutes, decreasing to about half at 6 hours, with a bimodal distribution possibly reflecting ongoing versus resolved seizures[36] (**Fig. 4B**).

Transcriptionally defined Supertypes correspond to gradations of anatomically distinct sub-populations of CA1 pyramidal neurons, evidenced by visualization within the ABC MERFISH atlas[7] (**Fig. 4C**). We therefore hypothesized that we would observe differences between CA1 Supertypes that may underlie functional specialization within the hippocampus along anatomical axes[37,38]. We find that EE strongly activates Supertypes 69-1 and 70-2 compared to others—notable given the dorsal distribution of these

Supertypes within hippocampus (**Fig. 4C**) and the engagement of this part of the

hippocampus during exploration[39–41]. In contrast, KA seizure manipulations demonstrate

that all Supertypes are competent at inducing IEGs following 30 minutes of global

activity, although variation in active cells at 6 hours suggests differences in ongoing

neural activity, ongoing transcription, or mRNA export among CA1 Supertypes (**Fig. 4D-**

E).

Prior single-cell studies have analyzed IEG expression in different neuronal cell types,

including in the hippocampus, to discern whether there are IEG "modules," i.e. genes

that are co-induced together[8,10,29]. No such analyses have revealed any interpretable

structure of IEG activation. We sought to determine if the unprecedented cell-type

resolution among ABC taxa in our dataset would reveal patterns of IEG expression in

CA1 subpopulations that were aggregated in prior analyses. If a pattern to stimulus-

dependent IEG induction existed, it would emerge in recently active cells, appearing as

highly correlated expression modules among IEG subsets, with anti-correlation between

subsets. When we tested this hypothesis among active cells in our most abundant

Supertypes after 30 minutes of EE, we observed patterns of modest positive correlation

among nearly all canonical IEGs across all Supertypes (**Fig. 4F**). We therefore conclude

there is minimal evidence for structured or modular expression of IEGs within subsets of

CA1-ProS neurons.

Figure 4

A
→ *Fos* induced
SE
EE 30m
KA 30m

→ *Arc* induced | 90th percentile in SE
SE
EE 30m
KA 30m

→ *Nr4a1* induced
SE
EE 30m
KA 30m

Expression

B 016 CA1-ProS Glut

Subclass

8% active
33% active
20% active
99% active
56% active

Num. IEGs induced

SE | EE 30m | EE 6h | KA 30m | KA 6h

C CA1 Supertypes, MERFISH (Yao 2023)

coronal view — D / V
diagonal view — A / P

D Percent of active CA1 cells by Supertype

	Total n	SE	EE30m	EE6h	KA30m	KA6h
CA1 69-1	14,030	7%	41%	24%	100%	56%
CA1 70-2	5,110	7%	34%	22%	99%	53%
CA1 71-3	495	3%	12%	4%	100%	14%
CA1 72-4	2,919	2%	8%	3%	100%	76%
CA1 73-5	133	14%	16%	3%	100%	14%
CA1 74-6	692	2%	11%	3%	100%	50%

E CA1 Supertypes across activity conditions

SE | EE 30m | EE 6h | KA 30m | KA 6h

Num. IEGs induced

69-1 70-2 71-3 72-4 73-5 74-6

F
Active cells in Supertype CA1 69-1
Fosb, Fosl2, Nr4a3, Btg2, Nr4a1, Dusp1, Npas4, Fos, Junb, Egr4, Egr3, Arc, Egr2, Egr1, Dusp5
n = 1768

Active cells in Supertype CA1 70-2
Fosb, Npas4, Btg2, Fos, Nr4a3, Fosl2, Nr4a1, Egr3, Egr4, Egr2, Junb, Arc, Egr1, Dusp1, Dusp5
n = 481

Active cells in Supertype CA1 72-4
Npas4, Fosb, Fos, Btg2, Egr3, Nr4a3, Fosl2, Nr4a1, Junb, Arc, Egr1, Egr2, Egr4, Dusp1, Dusp5
n = 73

Pearson's *r*
1
0.5
0
-0.5
-1

Figure 4. Distribution of IEG expression across CA1-ProS Supertypes.
(A) Example plots demonstrating IEG induction thresholds using *Fos*, *Arc*, and *Nr4a1*. For a given gene, the induction threshold was defined as the 90th percentile of expression among SE nuclei; thresholds are represented as dotted lines.
(B) Activation of single cells across activity conditions within CA1-ProS neurons. Active cells were defined as those that induced 3 or more IEGs, similar to the approach in Hrvatin et al. 2018 (Methods).
(C) 3-D plots of ABC MERFISH brain slices (published data from Yao et al. 2023) with only CA1-ProS Supertypes plotted.
(D-E) Activation percentages as in (B), de-aggregated by Supertype.
(F) Gene-gene correlation matrices for IEG expression within active cells at EE 30 min. Dendrograms represent hierarchical clustering of genes.

Circadian-dependent transcription among major hippocampal cell types in SE

Biological rhythms entrained to the diurnal light cycle are ancient cellular processes driven by transcription-translation feedback lopps that occur in many tissues, including the mouse hippocampus[23,42]. Yet, circadian cycling of transcription remains poorly understood outside the suprachiasmatic nucleus, with no single-cell studies characterizing the extent to which this occurs among different cell types in the hippocampus. We therefore sought to characterize transcriptional changes occurring at four Zeitgeber time points (ZTs) throughout the diurnal cycle under baseline conditions using our snRNA-seq approach, focusing on the four largest Subclasses to reduce noise. Canonical clock genes *Per1*, *Per2*, and *Cry2* peaked at ZT12 (start of the dark cycle), whereas *Clock* and *Bmal1* had opposite periodicity at ZT0 (**Fig. 5A**), aligning with previous hippocampal studies[18,19,23,43–47].

We next sought to characterize transcriptional changes taking place genome-wide using DGE analysis. When doing so, we capture the differential expression of canonical clock genes in every subclass tested (**Fig. 5B-E**, yellow annotations). We note again that *Per*-family expression peaks at ZT12 in every case but one. In addition to the *Per* family, its heterodimer partners *Cry1*, *Cry2*—as well as transcriptionally correlated Clock-gene regulators *Nr1d1*, and *Nr1d2*—peak at ZT12 in Subclasses where they are differentially expressed (**Fig. 5B-D**). Conversely, although we do not capture differential expression of *Clock* and *Bmal1* in every Subclass, their peak expression in such cases is not

concurrent with *Per-*, *Cry-*, or *Nr1d*-family peaks. These phase-shifted expression peaks

are functionally consistent with the canonical transcription-translation feedback loop

that undergirds circadian rhythms[48]. Thus, the ACT-DEPP dataset captures circadian

oscillations of transcription with high fidelity in multiple neuronal and glial cell types.

We next created UMAP embeddings of single-nucleus transcriptomes in circadian-DEG

space (2725 genes across all Subclasses) to determine whether there is a transcriptomic

structure tracking circadian time. We observe a clear separation of nuclei collected at

ZT12 in every Subclass examined, with additional gradations of other timepoints in some

cell types (**Fig. 5B-D**). High tau-specificity analysis confirms statistically what is visually

apparent: In all Subclasses, ZT12 is a unique circadian-transcriptomic state with broad

upregulation of many genes specifically at that time. Interestingly, in CA1-ProS cells, 12

of 17 of high-tau genes encoded lncRNAs, in contrast to DG granule cells (0/1),

astrocytes (4/13) and oligodendrocytes (4/40). This points to a unique recruitment of the

non-coding transcriptome in CA1-ProS neurons during this diurnal transition, the

functional significance of which is unclear. Finally, among previously characterized

ARGs[2] (ref [49] for *Adcy* genes), several of these varied throughout the circadian cycle,

suggesting that there may be an interaction between ZT and activity within the

hippocampus.

Figure 5

Figure 5. Circadian-dependent gene expression among major hippocampal cell types.
(A) Pseudobulk expression of canonical clock genes across the circadian period in SE. Error bars represent s.e.m.
(B-E) Above: UMAP embeddings of nuclei in circadian-DEG space, colored by ZT of collection (Methods). Below: Pseudobulk expression for circadian DEGs across the diurnal cycle. Clock genes (yellow), high-tau genes (red), and ARGs (black) are highlighted.

Interactions between activity- and circadian-dependent transcription in hippocampus

Previous studies have shown differences in canonical ARG expression at different times of day, although these studies could not make direct claims about stimulus-specific ARG induction due to their experimental design[18,18,19]. To directly test whether circadian time influences activity-dependent gene expression, we performed EE exposure at different ZTs. We hypothesized, based on these prior studies, that ARG induction would vary in at least some cell types across circadian time; that is, fold-change in ARG expression in EE vs SE would not be constant across the circadian period. We restricted our initial analyses to IEG and clock gene sets used in **Figures 4** and **5**, respectively. Surprisingly, there were no significant ZT:Activity interactions for IEGs in any of the Subclasses tested (**Fig. 6A-D**). This was unexpected, given previous reports of circadian-dependent IEG expression at baseline and upstream signaling cascades using non-RNA-seq methods[43,49,50]. We nevertheless note strong IEG induction in CA1, irrespective of ZT, unlike other Subclasses.

Having established that the expression of clock genes varies substantially in baseline conditions (**Fig. 5A**), we hypothesized that, if present, activity-dependent induction of these genes would not be uniform across ZTs. Indeed, we find that not only are *Per1* and *Per2* inducible in response to EE exposure, but also that *Per1* exhibits a significant ZT:Activity interaction in all Subclasses except DG, with inducibility restricted to times excluding ZT12—notable because ZT12 corresponds to peak expression in SE (**Fig. 5A**). This observation can be interpreted as a "ceiling effect" of EE-induced expression mimicking the peak circadian expression.

Finally, we tested for ZT:Activity interaction genome-wide. Surprisingly, we only found a

small number of genes whose expression was significantly modulated by ZT:Activity

interaction, and these genes only appeared in the neuronal Subclasses (**Fig. 6I-J**).

Among the cell types tested, we observe 13 genes that have a statistically significant

interaction between ZT:EE exposure, none of which are canonical IEGs. Functional

annotation of these genes reveals that two are involved in endocannabinoid synthesis

(*Daglb, Napepld*), three in cell signaling or excitability (*Fcho2, Itpkb, Kcnn2*), and four in

chromatin remodeling or gene expression (*Cecr2, Zfp420, Zpf523*). We therefore

conclude that changes in gene expression following novel experience are relatively

constant across the circadian cycle, although special mechanisms may be in place to

selectively change induction mechanisms for genes which modulate endocannabinoid

synthesis and cell excitability within neurons of the hippocampus.

Figure 6

Figure 6. Interactions between activity- and circadian-dependent transcription in the
hippocampus.
(A-D) Expression change between EE30m vs SE for IEGs across the circadian period. Baseline
pseudobulk expression for each gene at each ZT is plotted on the right.
(E-H) As in (A-D) but for canonical clock genes.
(I) Expression change between EE30m vs SE for genes with significant ZT:activity_condition
interaction (Methods). Baseline pseudobulk expression for each gene at each ZT is plotted on
the right.
(J) As in (I) but for DG. Note: astrocytes and oligodendrocytes had no significantly interacting
genes between ZT and activity condition.

Discussion

The ACT-DEPP dataset is an atlas of single-nucleus transcriptomes from the

hippocampus, capturing transcriptional responses to both physiological and

pathological brain activity. The creation of this atlas accomplishes two goals: (1) mapping

of experience-dependent gene expression in a high-resolution cell-type taxonomy, and

(2) characterization of circadian-dependent transcriptional oscillations in both neurons

and glia of the hippocampus. All cell types are registered to a gold-standard brain-wide

taxonomy, providing rich annotation that we leverage to demonstrate that stimulus-

responsive cell states are embedded within broader cell-type identities, such as

hippocampal sub-region (**Fig. 3**) and anatomical location (**Fig. 4**), with DG serving as an

exception to this rule (**Fig. 2**). We identify the light-to-dark transition as a key inflection

point for diurnal gene regulation in the hippocampus (**Fig. 5**) and show that activity-

dependent induction does not vary widely across the circadian period (**Fig. 6**). We

envision this resource as a roadmap for analyzing cell-type-specific, stimulus-responsive

states in the brain using modern transcriptome-registration tools. More broadly, the

ACT-DEPP dataset is a resource for understanding how physiological and pathological

activity patterns drive experience-dependent transcriptional programs to support neural

plasticity across the circadian cycle.

Limitations

We take care to point out a potential conceptual limitation of our study: Dissociating

neuronal gene expression driven by circadian phase from that driven by acute

locomotion is inherently challenging because the two are tightly coupled in behavior and

physiology. Locomotor activity co-occurs with changes in arousal, hormones, and

neuromodulator release, many of which independently influence transcription, while the

propensity for locomotor activity varies across the circadian cycle[23,51]. One argument that

favors the distinction we make between activity- and circadian-dependent expression is

that canonical IEGs do not overlap with circadian DEGs (**Fig. 5**), suggesting orthogonal

regulatory logic between activity programs and circadian programs. Second, in contrast

to neurons, glial transcription is less sensitive to rapid fluctuations in circuit activity and

more influenced by systemic cues, making their rhythmic gene expression less

confounded by locomotor events. Taken together, these observations support the

validity of interpreting our findings as reflecting both activity- and circadian-driven

transcriptional programs, rather than a single merged signal.

We also disclaim notable variance in cell number and read depth among a small number

of our biological replicates (**Supp. Fig. 2 and 3**). Although our pseudobulk approach

accounts for both of these factors by computing size factors for gene-count

normalization before performing DGE analysis, it is nevertheless likely that estimates of

variance are noisier in these conditions. However, for all of our DGE analyses, we set

DESeq2 gene-filtering parameters that would bias our analysis toward excluding genes

from DEG lists based on unacceptably high levels of within-condition variance for that

gene model. Pseudobulk DGE analysis has been shown to be the most robust against

Type I errors for single-cell data while performing the best in balancing Type II error

rate[52,53]. Thus, the results reported in this paper should be protected from unacceptably

high false-positive rates.

Cell states and cell types

Beyond our investigation of circadian transcription, our study has three main contributions to our understanding of activity-dependent gene expression. First, we extensively leverage registration of our dataset with the ABC atlas—a state-of-the-art ontology of cell types in the mouse brain[7]—to query activity-dependent states among high-resolution cell types. The ability to observe the effects of activity manipulations at the Supertype and Cluster level allows us to infer activity distributions along functionally relevant anatomical axes in the hippocampus (**Fig. 4**). Second, we distinguish several candidate genes that are highly specific not only to cell types, but also activity states (**Fig. 3**). These genes may serve as guideposts for the development of simpler next-generation tools to study activity-dependent processes with cell-type granularity going beyond IEG-based permanent labeling systems using *Fos, Arc, or Npas4*[54–57]—all of which are broadly upregulated across multiple cell types in our data, including non-neuronal cell types (**Fig. 3**). Such tools would have the advantage of being able to permanently access task-specific ensembles within specific cell types without the need for complex intersectional genetic strategies. Third, our descriptions of activity in baseline and "ceiling" conditions such as seizure provide strong evidence for the annotation of specific Supertypes and Clusters as highly temporally specific signatures of recent activity (**Fig. 2**).

Within DG granule cells, we observed a collapse of cell type and activity state at the supertype level (**Fig. 2**), and we also demonstrate that the Cluster rank in this population further resolves activity states into both acute and subacute activity histories, on the order of 30 minutes to 6 hours within our dataset (**Fig. 2E**). This example case in DG begs the question: Does the annotation of activity state—or any other condition-specific state, e.g. cellular manipulation or disease—at the Supertype or Cluster rank merit the collapse of these taxa with closely related inactive populations within the ABC taxonomy? More broadly, to what extent should reference atlases balance heterogeneity of cell states with the classification of cell types? On the one hand, the development of genetic tools to interrogate a particular Supertype or Cluster may produce unexpected

results if that taxon represents a distinct cell state not present, or overrepresented, within

an experimental dataset. On the other hand, it is easy to imagine how the annotation of

active DG cells, for example, through a simple compositional analysis of Supertypes or

Clusters would be useful for high-throughput readout of activity. It is our opinion that the

correlation of specific cell states with lower ranks in the ABC atlas is acceptable, as long

as these cell states are ascertained using robust experimental methods whose results

are clearly disclosed to the community.

Circadian rhythm changes baseline gene expression in the hippocampus

It has long been established that cells in the suprachiasmatic nucleus (SCN) use

transcription-translation feedback loops entrained to ambient-light sensation to keep a

"molecular clock" that tracks the diurnal cycle (Patton & Hastings 2018; Patton 2023).

However, there are few investigations of hippocampal circadian-dependent transcription

in the omics era, and no snRNA-seq studies that we are aware of. Therefore, we sought

to characterize how transcription changes in the hippocampus across the circadian

cycle using snRNA-seq. Our results extend previous work by confirming the presence of

clock-gene cycling in hippocampal neurons and glia across the circadian cycle. The

expression patterns of canonical clock genes we observe are consistent with earlier bulk

hippocampal studies[18,19,23,43], lending confidence that our sampling strategy captures

genuine circadian processes rather than stochastic variation. Importantly, we find that

ZT12 represents a distinct transcriptional inflection point across multiple cell types,

coinciding with the transition from the light to dark period. This time is behaviorally salient

for nocturnal animals such as mice, as locomotor activity begins to rise sharply before

and during the onset of darkness[23,46,51,58]. Our data therefore support the interpretation

that circadian gene expression in hippocampal cells is tightly coupled to this

environmental and behavioral transition.

The similarity of transcriptomic profiles at ZT16 and ZT0 further suggests that circadian-

driven transcription is concentrated within a narrow temporal window. Rather than a

gradual progression across the cycle, many genes appear to undergo sharp, temporally

restricted bursts of expression near the light–dark transition. This "bursty" model of circadian regulation is consistent with prior reports of pulse-like transcription in single cells[59–62] and may reflect a strategy by which hippocampal cells track salient transitions without maintaining high levels of rhythmic transcription throughout the entire dark period.

Among the cell types tested, we observed a striking specificity in circadian-dependent transcription: oligodendrocytes, which exhibit relatively few activity-dependent DEGs (**Fig. 3**), display the largest number of circadian DEGs. This dissociation underscores that activity-dependent and circadian-dependent transcription represent partially non-overlapping regulatory axes. For oligodendrocytes, the high number of circadian DEGs suggests that diurnal regulation may play a dominant role in processes such as myelin turnover or sheath maintenance, at least during the developmental period tested (approximately P28 mice). In contrast, neuronal populations such as CA1 exhibit fewer circadian DEGs overall, with enrichment of lncRNAs at ZT12 compared to other cell types, hinting at separate mechanisms of regulatory control. Together, these results suggest that circadian transcription exerts heterogeneous but highly structured influence across hippocampal cell types.

Circadian cycle has limited effect on activity-dependent transcription

We found no circadian differences in IEG induction (**Fig. 6A–D**), despite expecting higher expression in the dark period due to increased locomotor activity[51], state-dependent hippocampal network events such as sharp-wave ripples during rest[14], and prior reports of circadian oscillations in cAMP, MAPK, and CREB signaling[49,51,63,64]. Instead, IEGs were induced uniformly across the diurnal cycle in both neuronal and glial populations, exemplified by CA1 cells (**Fig. 6**). Prior work reported dark-phase upregulation of IEGs in rats under chronic EE[65] or in cortex at ZT20[50], but those studies differ in species, brain region, EE paradigm, stimulus, timing, and bulk vs. snRNA-seq, making direct comparison to our results inappropriate.

It has been previously shown that *Per* genes are inducible in the hippocampus in
response to circuit activity[18,19,47], a finding that we recapitulate after 30 minutes of EE.
This finding, in conjunction with *Per*'s circadian oscillations in the absence of activity,
raises important conceptual questions. What does it mean for there to be circadian-
dependent expression when these genes are also induced by activity? We observe that
at the highest peaks of EE-responsive induction, *Per* expression is comparable to its
circadian peak. One speculation that could explain this phenomenon is that in the
absence of activity, daily peaks in *Per* induction "guarantee" a dose of permissive
regulatory-element access for PER-CRY target genes which would otherwise not be
transcribed but are nevertheless necessary for cell function; activity-dependent *Per*
expression would act as an adjuvant to predictable circadian bursts, possibly at genes
important for cellular adaptations to circuit activity. Clearly, additional studies are
necessary to understand the multifactorial causes and functional consequences of *Per*
expression.

When testing for ZT:activity interaction genes, only 13 DEGs were found, all in neuronal
subclasses. The functional classes of these DEGs suggest that there is a special role for
fluctuations in intercellular endocannabinoid signaling throughout the circadian cycle, an
enticing idea that may be relevant for neurological phenotypes. For example, despite the
untold number of molecular etiologies for different epilepsies, circadian rhythms to
seizure attacks are common in drug-resistant epilepsy, occurring in approximately 90%
of patients[25]. This observation implies that there are circadian changes taking place in
the brain that are not themselves pathogenic, but may affect seizure threshold across
the diurnal cycle. In summary, our examination of the intersection of activity- and
circadian-dependent gene expression is an important first step toward understanding
the circadian transcriptome in the hippocampus, as it may canalize certain disease
phenotypes.

Author contributions

Jack A Olmstead: Conceptualization, Investigation, Methodology, Data curation, Formal analysis, Software, Visualization, Writing – original draft, Writing – review and editing

Lauren E King: Investigation, Writing – review and editing

Brenda L Bloodgood: Conceptualization, Supervision, Project administration, Funding acquisition, Writing – original draft, Writing – review and editing

Acknowledgements

We thank members of the Bloodgood lab and Dhanajay Bambah-Mukku for continuous discussion and feedback about experimental results and data analysis. We are indebted to Eric Halgren for providing computing resources. We thank Elizabeth Chamiec-Case and Lauren Valdez for their assistance with experimental troubleshooting. We kindly acknowledge the detailed experimental and analytic support provided by Brian Davis, Sam You, and Daniel Diaz of Parse Biosciences. This work was supported by funding awarded to BLB by the National Institute of Neurological Disorders and Stroke (R01 NS111162) and to the UC San Diego Medical Scientist Training Program (T32 GM154642). This publication includes data generated at the UC San Diego IGM Genomics Center utilizing an Illumina NovaSeq X Plus that was purchased with funding from a National Institutes of Health SIG grant (#S10 OD026929).

The authors declare no competing interests.

Materials and Methods

Animals

Animal experiments were approved by the University of California San Diego Institutional Animal Care and Use Committee and followed guidelines according to the National Institutes of Health's *Guide for the Care and Use of Laboratory Animals.* All experiments used wild-type C57BL6/J mice from The Jackson Laboratory (JAX#000664). Mice were

516 approximately four weeks old, with balanced proportions of males and females
517 distributed across control and experimental groups (**Supp. Table 1**).

518 **Experimental procedures**

519 *Standard environment and circadian timepoints*
520 Mice were group-housed in a vivarium on a 12-hour light/dark cycle and provided food
521 *ad libitum*. To avoid non-experimental stimulation artifacts resulting from cage handling
522 or transport, mice were transported from the vivarium to the standard environment (SE)
523 at least 18 hours before experiments. The SE consisted of a plywood box paneled with
524 sound-dampening foam, air-circulation fans, and LED lights whose power was controlled
525 by a timer mimicking the vivarium's light/dark cycle. Circadian timepoints were
526 measured and reported on the 24-hour Zeitgeber time scale, with ZT0 marked as "lights-
527 on" (at 06:00 PST) and ZT12 as "lights-off" (at 18:00 PST).

528 *Enriched environment*
529 For enriched-environment (EE) experimental conditions, animals were transferred from
530 their home cage in the standard environment to a larger cage in a different room
531 containing plastic and metal toys, tunnels, vertical scaffolds, and a running wheel. Mice
532 were placed in the EE with at least one cagemate and left to explore continuously for the
533 durations reported (i.e., for 30 minutes or 6 hours).

534 *Kainic-acid seizures*
535 Pharmacological seizures were induced using peritoneal injection of 15 mg/kg of kainic
536 acid (KA; Santa Cruz, #SC-200454) as described previously[36,66]. Seizure onset was
537 measured as the onset of rearing and accompanying bilateral forelimb clonus,
538 approximately corresponding to Racine stage 4[67]. Mice were sacrificed either 30 minutes
539 or 6 hours after seizure onset.

540 *snRNA-seq sample collection*
541 For sacrifice and sample collection, mice were deeply and rapidly anesthetized with
542 isoflurane in an induction chamber. Following induction, the brain was removed from the
543 skull into a dissection dish containing ice-cold dPBS (Gibco Cat. No. 14040133).
544 Hippocampus was rapidly extracted from one hemisphere, and the approximate CA1
545 area was microdissected using a microblade. Although the purpose of this
546 microdissection was to de-enrich samples of dentate granule cells, we still note that

these cells nevertheless comprise the 4th largest subclass in our dataset (**Fig. 1B**). The
dissected tissue sample was then placed in a microtube and flash-frozen. The other
hemisphere was placed in a cryomold as a whole, embedded in OCT compound, and
rapidly flash-frozen in isopentane chilled with dry ice. Left and right hemispheres were
alternated for the collection of CA1 and whole-hemisphere collection, across sexes and
conditions, to avoid any effects of lateralization (**Supp. Table 1**). Both samples from each
animal were stored at -70°C until use.

Nucleus isolation and fixation

Flash-frozen hippocampal dissectates were stored at -70°C until all experimental
samples had been collected. Before nucleus isolation, all buffers, tubes, and instruments
were pre-cooled on ice for 15-20 minutes before tissue was removed from -70°C.
Reusable instruments such as the Dounce tubes and pestles (Millipore cat. #D8938-
1SET) were rinsed with RNAse-free PBS and then with RNAse Away detergent before
being chilled on ice. All pipette tips (filtered tips only) and disposable tubes contacting
samples were RNAse-free. All buffers used during nucleus isolation contained
recombinant RNAsin to preserve RNA integrity (Promega, cat. #N2111), and actinomycin
D to prevent artifactual transcription stimulated by the isolation protocol (Tocris, cat.
#1229). Two main buffers were used for nucleus isolation: nucleus-isolation medium
(NIM) and NIM-Douce-pestle buffer. NIM buffer consisted of 250 mM sucrose, 25 mM
KCl, 5 mM $MgCl_2$, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.2 U/uL RNAsin, 25 uM
actinomycin D, and 1 cOmplete protease inhibitor mini tablet (Roche cat.
#11836170001), dissolved in 10 mL of buffer. NIM-DP buffer consisted of 0.1% triton-X
100 detergent in NIM buffer.

To begin nucleus isolation, tissue samples were removed from -70°C and placed directly
on ice to thaw. No more than five samples underwent isolation at any one time in order
to minimize sample-handling time before fixation. First, 1 mL of NIM-DP buffer was
added to the sample storage microtube, and the sample was transferred to a 2 mL
Dounce tube using a 1000 uL pipette tip. Samples were homogenized using a Douce
pestle in 1 mL of NIM-DP buffer, first with 10 strokes of the loose pestle, and then 17
strokes of the tight pestle. Two sets of dounce tubes and pestles were used at a time,
and each was washed five times with PBS in-between samples. Dounced tissue
homogenates were then spun in a fixed-angle benchtop centrifuge at 800 x g for 8
minutes at 4°C. The supernatant was removed, and the nuclear pellet, clearly visible,

was resuspended thoroughly in 500 uL of NIM buffer using 30 strokes with a P200
pipette tip. Suspensions were then filtered through a 70 um filter before being spun at
200 x g for 10 min at 4°C in a swinging-bucket centrifuge. The supernatant was removed,
and nuclei pellets were resuspended in 187.5 uL of Parse Biosciences's "Resuspension
Buffer" and strained through a 20 um filter to begin Parse's Evercode Nuclei Fixation
protocol (version 1.1, kit chemistry v3). In total, samples/nuclei were suspended in
actinomycin D and kept on ice or in the pre-chilled centrifuge for the entirety of the
isolation protocol, which took approximately 1 hour before proceeding directly to the
Parse fixation protocol.

snRNA-seq barcoding and library construction

snRNA-seq library construction was performed using Parse Biosciences's Whole
Transcriptome v3 kit according to the manufacturer's protocol. Briefly, nuclei originating
from different samples are multiplexed and barcoded according to the SPLiT-pool
combinatorial barcoding method[27], wherein individual nuclei go through serial rounds of
PCR barcoding, pooling, and splitting amongst a 96-well plate containing well-specific
oligo barcodes. Split-pooling occurs in four serial barcoding rounds to ensure that each
nucleus has a unique combination of barcodes ligated. All samples were loaded and
barcoded together to avoid batch effects from separate library preparations. Libraries
were sequenced at the UCSD Institute for Genomic Medicine on an Illumina NovaSeq X
Plus using paired-end 100-bp reads. Sequencing was targeted at approximately 90,000
reads per nucleus, with a final average read depth of 66,702 reads per nucleus.

Quantification and statistical analysis

Data preprocessing and quality control

Nuclei quality control was performed as outlined in **Supplementary Figure 1**. Raw
nucleus x gene count matrices from demultiplexed samples were generated from the
FASTQ files using Parse Bioscience's Trailmaker software pipeline, version 1.4.1. A
"raw" Seurat object created by Trailmaker containing 110,728 nuclei, along with the
digital gene-count matrix, associated sample metadata, and DoubletFinder scores[68] was
downloaded and used to begin nuclei QC. Percent of reads mapping to mitochondrial
reads was calculated for each nucleus. Nuclei that did not meet the following criteria
were removed: transcript count (UMI) >= 1500; gene count >= 1500; and percent-
mitochondrial-read count <= 5%. After this filtering step, 108,748 nuclei remained. Allen
Brain Cell (ABC) taxonomic labels were assigned using the MapMyCells online tool. Next,

single-nucleus raw gene counts were transformed using Seurat's SCTransform function
(v0.4.1), with the vars.to.regress argument set as the percent of reads mapping to
mitochondrial genes (percent.mt) and SPLiT-Seq sublibrary. SCTransformation was
computed across nuclei from all samples. (Note: Across-sample SCT-normalization,
rather than within-sample normalization followed by integration was determined
appropriate after normalizing, clustering, and visualizing UMAPs on a pilot sequencing
run from this dataset, where we observed no batch effects due to sample or SPLiT-seq
sublibrary). SCT-normalized counts were then used to perform principal component
analysis (PCA), nearest-neighbor graph construction, initial Leiden cluster assignment,
and uniform manifold approximation and projection (UMAP) embedding. This resulted in
the segmentation of 108,748 nuclei into 47 Leiden clusters.

At this step, manual curation of 47 detected Leiden clusters was performed by
experimenter J.O. cross-referencing plots and tables of UMI count, gene count, percent
mitochondrial reads, doublet scores, and putative ABC cell type within each Leiden
cluster. Based on these judgements, nuclei in 6 small Leiden clusters were removed
because of low counts, high mitochondrial reads, high doublet scores, or the collision of
many different putative ABC cell types. Finally, any nuclei with a DoubletFinder score <
0.02 were removed. After these quality-control steps, 93,066 nuclei remained. These
again went through final PCA, Leiden clustering, and UMAP embedding to create the
UMAP in **Figure 1**.

ABC taxon assignment stability
Stability of ABC taxonomy assignment for single nuclei was performed by comparing
two separate runs of the curated data through the ABC atlas MapMyCells tool. The first
run was performed using the native mouse transcriptome. The second run was
performed after removing ARGs that had been ascertained in a published study of
activity-dependent gene expression[2]. The stability of assignment was then determined
by comparing taxon equality within the same barcode and visualized with Sankey flow
diagrams (**Fig. 1E-H**).

Log2(FC) methods

Average log2(FC) expression values in **Figures 3** and **6** represent pseudobulk fold-
change expression values calculated using DESeq2.

Differential gene expression

To identify differentially expressed genes (DEGs) across activity conditions, we
performed pseudobulk DGE analysis independently for each Subclass using DESeq2[69].
For each Subclass, nuclei were subset and grouped by their sample of origin.
Pseudobulk counts were aggregated at the sample level using Seurat's
AggregateExpression() function. Genes expressed in fewer than 1% of nuclei were
excluded. DESeq2 was then used to construct a DESeqDataSet with sample-level
counts and a design formula including activity condition (expression ~
activity_condition). Only condition contrasts meeting a minimum per-group nucleus
count threshold (n = 30 nuclei per condition) were retained for testing. Wald tests were
performed, and log2 fold changes were shrunken using the apeglm method. Genes with
absolute log2 fold change > 0.585 (corresponding to 1.5-fold change) and fdr < 0.05
were classified as differentially expressed. For analysis discovering DEGs between ZTs,
log2 fold change values were not shrunken. This was done for ZT analyses but not
activity analyses to minimize type II error rate; there are fewer biological replicates at
each ZT (**Supplementary Table 1**). Results were saved separately for each Subclass
and contrast.

Hierarchical clustering

For heatmap plots accompanied by a dendrogram, genes were ordered according to
hierarchical clustering of the Euclidean gene distances using the package
ComplexHeatmap[70]. In the heatmaps of **Figure 5**, genes were first clustered into 4 k-
means clusters, and only after were organized according to hierarchical methods within
each k-means cluster.

ZT:Activity interaction testing

To test for statistical interactions between the effect of ZT and the effect of activity on
gene expression, we used the linear formula

$$expression \sim ZT + activity + ZT{:}activity$$

to test for a significant ZT:Activity interaction term with DESeq2. Pseudobulk gene

counts were computed for each subclass separately. log2 fold changes were not

shrunken in this analysis in order to increase sensitivity of this test. For *a priori* gene sets,

multiple-hypothesis correction was performed by calculating the FDR among the *a priori*

gene set. For example, the FDR rates reported in **Figure 6A-D** were corrected for 15

multiple comparisons for the set of 15 IEGs. Only genes with an FDR < 0.05 were labeled

as significant DEGs.

ERG vs LRG classification

We used a similar method to a prior single-cell study to classify DEGs as mutually

exclusive early-response genes (ERGs) or late-response genes (LRGs)[10]. Classification

was performed separately for excitatory neurons, inhibitory neurons, and glial

Subclasses. First, DEGs for all Subclasses and contrasts (EE30m vs SE, EE6h vs SE)

were aggregated. Genes were provisionally labeled as ERG, LRG, or both based on

differentially expressed in EE 30 min or EE 6 hr. Genes initially labeled as "both" were

reclassified based on their peak fold-change timepoint. In cases where gene

classification differed across Subclasses, the modal classification was assigned. Final

classifications were saved for downstream analysis (**Figure 3**).

Tau calculation

We calculated a tau score for each gene, adapted from ref. [35], to discover and quantify

highly cell-type specific gene expression. For example, a gene with a tau score of 0

would have promiscuous expression among the cell types examined, whereas a gene

with a tau score approaching or greater than 1 would have highly cell-type specific

expression. For analyses on non-fold-change expression data among different cell types

(e.g. **Figure 5**), we directly applied the tau formula, in which tau was bounded by (0, 1):

$$\tau = \frac{1}{n-1} \sum_{i=1}^{n} \left(\frac{expression_i}{max(expression)} \right),$$

where *n* is the number of subclasses, and *expression$_i$* represents the expression of the

gene in subclass *i*; in these analyses, our tau threshold was set at 0.75 due to the index

being bounded by (0, 1). For analyses on fold-change expression where expression

values can be negative, the tau score was bounded by (0, ∞); for these analyses, our tau

threshold was set higher at 0.8. In practice, tau scores above 0.75 were observed to be

highly specific to one or a small number of subclasses.

Defining active cells

In order to determine the number of cells activated within a given condition, it was

determined whether cells expressed the IEGs above a certain threshold: The IEG list was

defined *a priori* based on the following 15 canonical IEGs: *Arc, Btg2, Dusp1, Dusp5,*

Egr1, Egr2, Egr3, Egr4, Fos, Fosb, Fosl2, Junb, Npas4, Nr4a1, and *Nr4a3*. The induction

threshold was determined for each gene individually, and was defined as the 90th

percentile of expression in SE cells of the CA1 subclass. Thus, for each gene in the gene

list, a binary classification was assigned as either induced or not-induced based on

whether or not it surpassed the expression threshold (**Fig. 4A**). Finally, a cell was defined

as "active" if it induced at least 20% of the genes (i.e., 3 or more) in the activity-regulated-

gene list.

Data and code availability

Fastq files for sequenced reads will be publicly deposited in the GEO database.

Metadata for biological replicates can be downloaded from the GEO database or

https://github.com/Bloodgood-Lab/Olmstead-2025-preprint. We used the Trailmaker

pipeline running split-pipe v1.4.1 through Parse Biosciences' web portal for creation of

the cell-count matrix and unprocessed Seurat object, which will be publicly deposited in

the GEO database. All the code and package versions to reproduce this manuscript are

publicly available at https://github.com/Bloodgood-Lab/Olmstead-2025-preprint.

References

1. Greenberg, M. E. & Ziff, E. B. Stimulation of 3T3 cells induces transcription of the c-fos proto-oncogene. *Nature* **311**, 433–438 (1984).

2. Tyssowski, K. M. *et al.* Different Neuronal Activity Patterns Induce Different Gene Expression Programs. *Neuron* **98**, 530-546.e11 (2018).

3. Guzowski, J. F., McNaughton, B. L., Barnes, C. A. & Worley, P. F. Environment-specific expression of the immediate-early gene Arc in hippocampal neuronal ensembles. *Nat Neurosci* **2**, 1120–1124 (1999).

4. Tullai, J. W. *et al.* Immediate-Early and Delayed Primary Response Genes Are Distinct in Function and Genomic Architecture*. *Journal of Biological Chemistry* **282**, 23981–23995 (2007).

5. Lituma, P. J., Singer, R. H., Das, S. & Castillo, P. E. Real-time imaging of Arc/Arg3.1 transcription ex vivo reveals input-specific immediate early gene dynamics. *Proceedings of the National Academy of Sciences* **119**, e2123373119 (2022).

6. Akiki, R. M. *et al.* A long noncoding eRNA forms R-loops to shape emotional experience–induced behavioral adaptation. *Science* **386**, 1282–1289 (2024).

7. Yao, Z. *et al.* A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse brain. *Nature* **624**, 317–332 (2023).

8. Langlieb, J. *et al.* The molecular cytoarchitecture of the adult mouse brain. *Nature* **624**, 333–342 (2023).

9. Lacar, B. *et al.* Nuclear RNA-seq of single neurons reveals molecular signatures of activation. *Nat Commun* **7**, 11022 (2016).

10. Hrvatin, S. *et al.* Single-cell analysis of experience-dependent transcriptomic states in the mouse visual cortex. *Nat Neurosci* **21**, 120–129 (2018).

11. Jarrard, L. E. On the role of the hippocampus in learning and memory in the rat. *Behavioral and Neural Biology* **60**, 9–26 (1993).

12. O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Research* **34**, 171–175 (1971).

13. Nádasdy, Z., Hirase, H., Czurkó, A., Csicsvari, J. & Buzsáki, G. Replay and Time Compression of Recurring Spike Sequences in the Hippocampus. *J. Neurosci.* **19**, 9497–9507 (1999).

14. Buzsáki, G. Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. *Hippocampus* **25**, 1073–1188 (2015).

15. Joo, H. R. & Frank, L. M. The hippocampal sharp wave–ripple in memory retrieval for immediate use and consolidation. *Nat Rev Neurosci* **19**, 744–757 (2018).

16.   Carr, M. F., Jadhav, S. P. & Frank, L. M. Hippocampal replay in the awake state: a
potential substrate for memory consolidation and retrieval. *Nat Neurosci* **14**, 147–
153 (2011).

17.   Pfeiffer, B. E. The content of hippocampal "replay". *Hippocampus* **30**, 6–18
(2020).

18.   Bellfy, L. *et al.* The clock gene Per1 may exert diurnal control over hippocampal
memory consolidation. *Neuropsychopharmacol.* **48**, 1789–1797 (2023).

19.   Kwapis, J. L. *et al.* Epigenetic regulation of the circadian gene Per1 contributes to
age-related changes in hippocampal memory. *Nat Commun* **9**, 3323 (2018).

20.   Harvey, J. R. M., Plante, A. E. & Meredith, A. L. Ion Channels Controlling Circadian
Rhythms in Suprachiasmatic Nucleus Excitability. *Physiological Reviews* **100**,
1415–1454 (2020).

21.   Gonzalez, J. C. *et al.* Circadian regulation of dentate gyrus excitability mediated
by G-protein signaling. *Cell Reports* **42**, 112039 (2023).

22.   Flourakis, M. *et al.* A Conserved Bicycle Model for Circadian Clock Control of
Membrane Excitability. *Cell* **162**, 836–848 (2015).

23.   Birnie, M. T. *et al.* Circadian regulation of hippocampal function is disrupted with
corticosteroid treatment. *Proceedings of the National Academy of Sciences* **120**,
e2211996120 (2023).

24.   Gerstner, J. R. *et al.* BMAL1 controls the diurnal rhythm and set point for electrical
seizure threshold in mice. *Front. Syst. Neurosci.* **8**, (2014).

25.   Karoly, P. J. *et al.* Circadian and circaseptan rhythms in human epilepsy: a
retrospective cohort study. *The Lancet Neurology* **17**, 977–985 (2018).

26.   Rawashdeh, O. *et al.* PERIOD1 coordinates hippocampal rhythms and memory
processing with daytime. *Hippocampus* **24**, 712–723 (2014).

27.   Rosenberg, A. B. *et al.* Single-cell profiling of the developing mouse brain and
spinal cord with split-pool barcoding. *Science* **360**, 176–182 (2018).

28.   Fernandez-Albert, J. *et al.* Immediate and deferred epigenomic signatures of in
vivo neuronal activation in mouse hippocampus. *Nat Neurosci* **22**, 1718–1730
(2019).

29.   Traunmüller, L. *et al.* Novel environment exposure drives temporally defined and
region-specific chromatin accessibility and gene expression changes in the
hippocampus. *Nat Commun* **16**, 7787 (2025).

30.   Johnson, A., Varberg, Z., Benhardus, J., Maahs, A. & Schrater, P. The
hippocampus and exploration: dynamically evolving behavior and neural
representations. *Front. Hum. Neurosci.* **6**, (2012).

31. Diamantaki, M., Frey, M., Berens, P., Preston-Ferrer, P. & Burgalossi, A. Sparse
activity of identified dentate granule cells during spatial exploration. *eLife* **5**,
e20252 (2016).

32. Chavlis, S., Petrantonakis, P. C. & Poirazi, P. Dendrites of dentate gyrus granule
cells contribute to pattern separation by controlling sparsity: DENDRITIC ROLE IN
PATTERN SEPARATION. *Hippocampus* **27**, 89–110 (2017).

33. Lee, J. Q. *et al.* Sparsity of Population Activity in the Hippocampus Is Task-
Invariant Across the Trisynaptic Circuit and Dorsoventral Axis. *Hippocampus* **35**,
e23651 (2025).

34. Borzello, M. *et al.* Assessments of dentate gyrus function: discoveries and
debates. *Nat. Rev. Neurosci.* **24**, 502–517 (2023).

35. Yanai, I. *et al.* Genome-wide midrange transcription profiles reveal expression
level relationships in human tissue specification. *Bioinformatics* **21**, 650–659
(2005).

36. Pollina, E. A. *et al.* A NPAS4–NuA4 complex couples synaptic activity to DNA
repair. *Nature* 1–10 (2023) doi:10.1038/s41586-023-05711-7.

37. Yao, Z. *et al.* A transcriptomic and epigenomic cell atlas of the mouse primary
motor cortex. *Nature* **598**, 103–110 (2021).

38. van Strien, N. M., Cappaert, N. L. M. & Witter, M. P. The anatomy of memory: an
interactive overview of the parahippocampal–hippocampal network. *Nat Rev
Neurosci* **10**, 272–282 (2009).

39. Rolotti, S. V., Blockus, H., Sparks, F. T., Priestley, J. B. & Losonczy, A.
Reorganization of CA1 dendritic dynamics by hippocampal sharp-wave ripples
during learning. *Neuron* **110**, 977-991.e4 (2022).

40. Foster, D. j., Morris, R. g. m. & Dayan, P. A model of hippocampally dependent
navigation, using the temporal difference learning rule. *Hippocampus* **10**, 1–16
(2000).

41. Dragoi, G. & Tonegawa, S. Preplay of future place cell sequences by hippocampal
cellular assemblies. *Nature* **469**, 397–401 (2011).

42. Hor, C. N. *et al.* Sleep–wake-driven and circadian contributions to daily rhythms in
gene expression and chromatin accessibility in the murine cortex. *Proc. Natl.
Acad. Sci. U.S.A.* **116**, 25773–25783 (2019).

43. Reick, M., Garcia, J. A., Dudley, C. & McKnight, S. L. NPAS2: An Analog of Clock
Operative in the Mammalian Forebrain. *Science* **293**, 506–509 (2001).

44. Wang, L. M. C. *et al.* Expression of the Circadian Clock Gene Period2 in the
Hippocampus: Possible Implications for Synaptic Plasticity and Learned
Behaviour. *ASN Neuro* **1**, AN20090020 (2009).

45. Jilg, A. *et al.* Temporal dynamics of mouse hippocampal clock gene expression
support memory processing. *Hippocampus* **20**, 377–388 (2010).

46.   Moriya, S., Tahara, Y., Sasaki, H., Ishigooka, J. & Shibata, S. Housing under
abnormal light–dark cycles attenuates day/night expression rhythms of the clock
genes Per1, Per2, and Bmal1 in the amygdala and hippocampus of mice.
Neuroscience Research **99**, 16–21 (2015).

47.   Bering, T., Blancas-Velazquez, A. S. & Rath, M. F. Circadian Clock Genes Are
Regulated by Rhythmic Corticosterone at Physiological Levels in the Rat
Hippocampus. *Neuroendocrinology* **113**, 1076–1090 (2023).

48.   Chiou, Y.-Y. *et al.* Mammalian Period represses and de-represses transcription by
displacing CLOCK–BMAL1 from promoters in a Cryptochrome-dependent
manner. *Proc. Natl. Acad. Sci. U.S.A.* **113**, (2016).

49.   Eckel-Mahan, K. L. *et al.* Circadian oscillation of hippocampal MAPK activity and
cAMP: implications for memory persistence. *Nat Neurosci* **11**, 1074–1082 (2008).

50.   Cirelli, C. & Tononi, G. Gene expression in the brain across the sleep–waking
cycle1. *Brain Research* **885**, 303–321 (2000).

51.   Phan, T. H., Chan, G. C.-K., Sindreu, C. B., Eckel-Mahan, K. L. & Storm, D. R. The
Diurnal Oscillation of MAP (Mitogen-Activated Protein) Kinase and Adenylyl
Cyclase Activities in the Hippocampus Depends on the Suprachiasmatic Nucleus.
J. Neurosci. **31**, 10640–10647 (2011).

52.   Squair, J. W. *et al.* Confronting false discoveries in single-cell differential
expression. *Nat Commun* **12**, 5692 (2021).

53.   Zimmerman, K. D., Espeland, M. A. & Langefeld, C. D. A practical solution to
pseudoreplication bias in single-cell studies. *Nat Commun* **12**, 738 (2021).

54.   Denny, C. A. *et al.* Hippocampal Memory Traces Are Differentially Modulated by
Experience, Time, and Adult Neurogenesis. *Neuron* **83**, 189–201 (2014).

55.   Sun, X. *et al.* Functionally Distinct Neuronal Ensembles within the Memory
Engram. *Cell* **181**, 410-423.e17 (2020).

56.   Guenthner, C. J., Miyamichi, K., Yang, H. H., Heller, H. C. & Luo, L. Permanent
Genetic Access to Transiently Active Neurons via TRAP: Targeted Recombination
in Active Populations. *Neuron* **78**, 773–784 (2013).

57.   Sørensen, A. T. *et al.* A robust activity marking system for exploring active
neuronal ensembles. *eLife* **5**, e13918 (2016).

58.   Riggle, J. P. *et al.* Spontaneous Recovery of Circadian Organization in Mice
Lacking a Core Component of the Molecular Clockwork. *J Biol Rhythms* **37**, 94–
109 (2022).

59.   Tunnacliffe, E. & Chubb, J. R. What Is a Transcriptional Burst? *Trends in Genetics*
36, 288–297 (2020).

60.   Corrigan, A. M. & Chubb, J. R. Regulation of Transcriptional Bursting by a
Naturally Oscillating Signal. *Current Biology* **24**, 205–211 (2014).

61. Coulon, A., Chow, C. C., Singer, R. H. & Larson, D. R. Eukaryotic transcriptional
dynamics: from single molecules to cell populations. *Nat Rev Genet* **14**,
10.1038/nrg3484 (2013).

62. Chen, L.-F. *et al.* Enhancer Histone Acetylation Modulates Transcriptional
Bursting Dynamics of Neuronal Activity-Inducible Genes. *Cell Reports* **26**, 1174-
1188.e5 (2019).

63. Luo, J., Phan, T. X., Yang, Y., Garelick, M. G. & Storm, D. R. Increases in cAMP,
MAPK Activity, and CREB Phosphorylation during REM Sleep: Implications for
REM Sleep and Memory Consolidation. *J. Neurosci.* **33**, 6460–6468 (2013).

64. Saraf, A., Luo, J., Morris, D. R. & Storm, D. R. Phosphorylation of Eukaryotic
Translation Initiation Factor 4E and Eukaryotic Translation Initiation Factor 4E-
binding Protein (4EBP) and Their Upstream Signaling Components Undergo
Diurnal Oscillation in the Mouse Hippocampus. *Journal of Biological Chemistry*
289, 20129–20138 (2014).

65. Rönnbäck, A., Dahlqvist, P., Bergström, S.-A. & Olsson, T. Diurnal effects of
enriched environment on immediate early gene expression in the rat brain. *Brain*
Research **1046**, 137–144 (2005).

66. Bosco, D. B. *et al.* RNAseq analysis of hippocampal microglia after kainic acid-
induced seizures. *Mol Brain* **11**, 34 (2018).

67. Racine, R. J. Modification of seizure activity by electrical stimulation: II. Motor
seizure. *Electroencephalography and Clinical Neurophysiology* **32**, 281–294
(1972).

68. McGinnis, C. S., Murrow, L. M. & Gartner, Z. J. DoubletFinder: Doublet Detection
in Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. *Cell*
Systems **8**, 329-337.e4 (2019).

69. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

70. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. *Bioinformatics* **32**, 2847–2849
(2016).

Supplementary Figure 1

Supplementary Figure 1. Quality control of single-nuclei dataset.
The initial unfiltered Seurat object was downloaded from Parse Biosciences' Trailmaker pipeline. Nuclei underwent QC thresholding, doublet rejection, and preliminary clustering to arrive at the final dataset of 93,066 nuclei.

Supplementary Figure 2

Supplementary Figure 2. UMI counts across major cell types.
Nuclei (left) and unique molecular identifier (UMI) counts (right) across major cell types in all biological replicates of the ACT-DEPP dataset. Colored dots on the left correspond to ZT times tested.

Supplementary Figure 3

Supplementary Figure 3. Gene counts across major cell types.
Nuclei (left) and gene counts (right) across major cell types in all biological replicates of the ACT-DEPP dataset. Colored dots on the left correspond to ZT times tested.

Factor design in ACT-DEPP dataset

		Age ± sd	Replicates	Sex		Hemisphere		Nuclei	
				M	F	R	L	Total	Average
SE	ZT0	30.0 ± 0.0	4	2	2	2	2	6,912	1,728
	ZT4	28.5 ± 0.6	4	2	2	2	2	8,424	2,106
	ZT12	30.0 ± 0.0	3	2	1	2	1	3,656	1,219
	ZT16	28.0 ± 0.0	4	2	2	2	2	8,016	2,004
	n_{total}		15	8	7	8	7	27,008	7,057
EE30m	ZT0	30.0 ± 0.0	4	2	2	2	2	8,091	2,023
	ZT4	28.5 ± 0.6	4	2	2	2	2	7,596	1,899
	ZT12	32.0 ± 0.0	3	2	1	2	1	4,497	1,499
	ZT16	28.5 ± 0.6	4	2	2	2	2	9,179	2,295
	n_{total}		15	8	7	8	7	29,363	7,716
EE6h	ZT0	29.0 ± 0.0	3	1	2	3	0	5,983	1,994
	ZT4	28.0 ± 0.0	3	1	2	2	1	6,114	2,038
	ZT12	31.0 ± 0.0	3	1	2	1	2	6,685	2,228
	ZT16	32.0 ± 0.0	3	1	2	2	1	5,946	1,982
	n_{total}		12	4	8	8	4	24,728	8,243
KA30m	—	29.0 ± 0.0	3	1	2	1	2	3,744	1,248
KA6h	—	29.0 ± 0.0	3	1	2	1	2	8,223	2,741
$n_{ACT\text{-}DEPP}$			48	22	26	26	22	93,066	

Supplementary Table 1. Factor design in the ACT-DEPP dataset.