Conservation units for anadromous Arctic Char (*Salvelinus alpinus*) in the Canadian Arctic informed by genetic structure, population connectivity and adaptive genomic variation

Abstract

Intraspecific genetic diversity is a crucial aspect of biodiversity conservation as it preserves evolutionary potential and enhances resilience to environmental change. Genomic-informed delineation of Conservation Units (CUs) offers ways of subdividing species into groups based on historical isolation and adaptive differentiation, to develop biologically relevant conservation and management policies. CUs have been defined in many species of harvested anadromous salmonids, but broad scale data remains lacking in the Canadian Arctic, where anadromous Arctic Char (*Salvelinus alpinus*) dominates catches in Indigenous-led subsistence and commercial fisheries. In this study, we use low-coverage whole-genome data from 30 Canadian populations of Arctic Char to define CUs based on population structure and connectivity, as well as adaptive genetic variation. We highlight two main genetic groups, each of which comprises three subgroups, or candidate CUs: the North (above the 67th parallel), including the North Baffin Island, Kitikmeot, and Inuvialuit Settlement Region CUs; and the South (below the 67th parallel), including the South Baffin Island, Ungava Bay, and Hudson Bay CUs. This delimitation is supported by areas of low effective migration between candidate CUs, as well as isolation-by-environment, which suggests adaptive differentiation. Finally, we discuss opportunities and caveats relating to linkage when identifying adaptive genetic variation from whole genome sequencing data through genome scans and Gene-Environment Associations.

Introduction

Given the numerous threats to global biodiversity, there has been a call for conservation at the levels of ecosystems, species, and within-species genetic

diversity (Laikre et al., 2020). Of these three levels, species conservation has
gathered much of the attention, suggesting a simple categorization of nature and
offering a common metric of biodiversity. In practice, the delimitation and
definition of species is rarely straightforward (Mace, 2004; Frankham et al., 2012;
Garnett & Christidis, 2017), and speciation proceeds via gradual genomic
divergence among isolated populations driven by the combined effects of
genetic drift and natural selection in contrasted habitats (Roux et al., 2016). Both
these evolutionary forces are subject to fluctuations over long timescales, as
populations reconnect in secondary contact zones, or changes in the
environment shift selective pressures (Seehausen et al., 2008; Rosenblum et al.,
2012). The resulting intraspecific genetic diversity is a key component of
biodiversity that warrants protection, as it underpins evolutionary potential and
enhances resilience to environmental change and anthropogenic pressures
(Forest et al., 2007; Schindler et al., 2010).

Biological systems are inherently complex, and speciation is best
understood as a continuous process. Nevertheless, discrete intraspecific
classifications remain necessary to inform conservation priorities within limited
budgets and to guide the management of harvested species. There exists a vast
body of literature on the delineation of conservation units (CU), which generally
relies on the identification of historical isolation (leading to genomic divergence)
and adaptive differentiation (Waples, 1991; Moritz, 2002; Coates et al., 2018).
Among the numerous CU models developed over the years, the Evolutionary
Significant Unit (ESU; first proposed in Ryder, 1986) aims to provide a framework
for the definition of subspecific groups that represent both unique genetic
variation and the historical or ongoing connectivity among populations within a
species. Depending on its definition, it can either be based exclusively on genetic
characters or incorporate ecological data (Fraser & Bernatchez, 2001). As an
additional concept, the Management Unit (MU) can be defined within ESUs as
are demographically independent populations identified based on contemporary
gene flow and population dynamics rather than deep evolutionary history. This
subdivision allows managers to implement local actions, such as

supplementation, restoration, or monitoring, that maintain genetic diversity,
reduce inbreeding, buffer against environmental fluctuations, and support the
long-term viability of the broader ESU (Moritz, 2002).

The concept of the ESU has been translated into legislation in multiple
countries (e.g., the USA 'Endangered Species Act' and the Australian
'Endangered Species Protection Act'). In Canada, the 'Species at Risk Act'
applies a similar concept of Designatable Units (DU, Green, 2005), which are
considered discrete if there is evidence of limited transmission of heritable traits
or markers between them, or if they are separated by a natural disjunction in their
distribution. A discrete unit is then considered significant if its evolutionary
trajectory dates back long enough to create an evolutionary history not found
elsewhere in Canada (e.g. units originating from distinct glacial refugia) or if it
possesses unique and verifiable adaptive traits.

Genetic data has been central to work related to CUs, as their delineation
often revolves around a two-step process: 1) describing the structure of
populations, which is often hierarchical within species, and 2) choosing the level
of distinctiveness appropriate for defining units, which implies some degree of
subjectivity (Waples, 1995). Cytoplasmic (both mitochondrial and chloroplastic)
DNA has long been used to characterize intraspecific lineages (Avise et al.,
1987), while microsatellites have contributed to investigating population structure
and connectivity (Ellegren, 2004). However, genomic data, i.e. genome-wide
datasets produced by genotyping-by-sequencing (GBS, thousands of SNPs) or
whole-genome sequencing (WGS, millions of SNPs), have the potential to
improve delineation of CUs by quantifying adaptive variation in addition to neutral
variation (Funk et al., 2012). This can be done by scanning the genome for loci
displaying higher differentiation than expected under neutral evolution,
suggesting the imprint of natural selection, or assessing correlations between
allele frequencies and biologically relevant environmental variables (hereafter
gene-environment associations, GEA, Rellstab et al., 2015; Dauphin et al.,
2023).

Anadromous salmonids (i.e., those migrating from marine to freshwater
habitats to spawn) represent a diverse group of fishes with significant economic
and cultural importance worldwide. Their strong homing behavior (i.e., returning
to natal sites for spawning, thus limiting gene flow among populations) plays a
crucial role in shaping their population structure and is a factor long recognized
as key for their conservation and management (Moulton 1939; Thompson 1959).
Many salmonid populations are facing strong harvesting pressures and are
actively managed within CUs to ensure sustainability and long-term population
persistence. For instance, Atlantic Salmon (*Salmo salar*) along Canada's eastern
coast are currently managed within 15 extant DUs (COSEWIC, 2010; Moore et
al., 2014). However, further subdivisions into 19 DUs was recently suggested
based on genomic data (Lehnert et al., 2023). On the western coast, Pacific
Salmons (*Oncorhynchus spp.*) are divided into dozens of CUs per species, with
larger watersheds, such as the Fraser River in British Columbia, supporting
multiple units (Xuereb et al., 2022; DFO, 2024). Overall, CUs have proven
essential for guiding effective management and sustaining salmonid populations
across diverse systems.

Northern Canada is home to populations of Arctic Char (*Salvelinus
alpinus*), a salmonid whose anadromous form has long been the most harvested
fish across Inuit Nunangat (i.e. the homeland of Inuit; Friesen, 2004; Priest &
Usher, 2004). Arctic Char fisheries are predominantly small-scale and harvested
for subsistence purposes (Roux et al., 2019; Tallman et al., 2019), but
collectively amount to a high replacement value, i.e., the estimated cost of
substituting the harvested fish with equivalent grocery-bought protein (Brubacher,
2004). While subsistence stocks are locally managed, some regions harbour
commercial fisheries that are co-managed between the Indigenous land claim
body and federal government (Fisheries and Oceans Canada). Continent-wide
genetic variation in Arctic Char has been investigated using mtDNA and
microsatellites. These studies suggest that most North American Arctic
populations derive from a single High Arctic glacial refugium, the Arctic lineage
(Moore et al., 2015), with the exception of Labrador (Salisbury et al., 2019) and

Greenland (Jacobsen et al., 2022), where an additional Atlantic lineage is also present. However, recent whole-genome data revealed that admixture between the Arctic and Atlantic lineage was a prominent feature of Arctic Char genomes in all Canadian populations south of the 67[th] parallel, redefining the extent of the second contact between lineages (Dallaire et al., 2025). Other recent genomic datasets for the species were focused on spatial scales from local to regional and have been used to investigate questions relating to population structure and connectivity (Moore et al., 2017; Li et al., 2021), morphotype differentiation (Kess et al., 2021), and local adaptation (Madsen et al., 2019; Dallaire et al., 2021; Layton et al., 2021).

In this study, we revisit the whole-genome resequencing dataset introduced in Dallaire et al. (2025), including more than 1,000 anadromous Arctic Char whole-genomes in 30 sampling sites across the Canadian Arctic, to define candidate CUs at a national scale. To inform these units, we 1) characterize in depth the hierarchical genetic structure of populations in relation to administrative regions in the Inuvialuit Settlement Region (ISR, Northwest Territories), Nunavut, and Nunavik (Québec), and 2) identify areas of lower gene flow. Then, we aim to 3) disentangle patterns of i) isolation-by-distance, ii) isolation-by-environment, and iii) isolation-by-colonization (related to post-glacial recolonization) that characterize this structure. Finally, we 4) detect signals of natural selection and explore patterns of population structure driven by putatively adaptive genetic variation using genome scans and Gene-Environment Associations. We discuss those results through the lens of the Designatable Unit framework and explore and discuss the added value of whole-genome data to conservation genomics.

Methodology

Sequencing and SNP calling

We reanalysed sequencing data from 1016 Arctic Char from 30 Canadian populations, published in Dallaire et al. (2025). Briefly, whole-genome sequences were obtained following Therkildsen & Palumbi (2017) on DNA extracted from fin

clips from fish harvested at the mouth of rivers in the Inuvialuit Settlement Region (Northwest Territories), Nunavut, and Nunavik (Québec) between 2007 and 2023 (Fig.1a). As in Dallaire et al. (2025), data were processed following the pipeline described at https://github.com/enormandeau/wgs_sample_preparation: sequences were trimmed and aligned on the ASM291031v2 reference genome (likely from *Salvelinus malma* or a *S. alpinus* x *S. malma* hybrid; Christensen et al., 2021), duplicate reads were removed, indels were realigned and overlapping ends of paired reads were clipped.

Mapped and cleaned reads yielded an individual depth of coverage around 2X, and we estimated Genotype Likelihoods (GL) at polymorphic positions identified in Dallaire et al. (2025). Those positions were distributed on all assembled linkage groups, but SNPs in sex-linked regions identified in Beemelmanns et al. (2024) were removed from all analyses. The SNP list was filtered for paralogs and other deviations from expected patterns due to transposable elements and ancestral autoploidization in salmonids, using *ngsparalog* (Linderoth, 2018) following Dallaire et al. (2023). We kept SNPs with global minor allele frequency above 1% and with a minimal depth of coverage of 1X in at least 75% of individuals. We used *ngsLD* (Fox et al., 2019) to estimate linkage between SNPs within 500 kb using a random subset of half the samples to create a LD-pruned dataset by iteratively removing SNPs with the graph method until no SNP pairs within 200kb had an r^2 above 0.1.

Extraction of environmental data

We used the ArcGIS software v10.4 (ESRI, 2011) to extract environmental data. Marine variables represent sea-surface values for factors of potential biological importance for Arctic Char available in the BIO-Oracle v3.0 dataset (Assis et al., 2024) and were averaged in a 20 km radius around each sampled river mouth and within 5 km from the coast. This standardized approach was used to characterize the local coastal environment in accordance with current knowledge of Arctic Char marine habitat use in Arctic Canada (Spares et al., 2015; Moore et al., 2016; Harris et al., 2020). We selected annual coastal averages for five

marine variables (temperature, salinity, primary productivity, dissolved O_2, and
ice cover) as they represented good predictors for values in summer, when Arctic
Char occupies coastal habitats.

Freshwater variables were extracted from the RiverATLAS dataset by
selecting the linear segment closest to the mouth of each sampled river, as these
segments include values averaged over the entire upstream watershed. We
initially considered seven variables: % of the watershed covered by lake or tree,
average slope, sand fraction in the soil, and annual averages of discharge, air
temperature, and precipitation. These variables are linked to both migratory
challenges (e.g., slope, discharge, temperature) and spawning habitats (e.g.,
lake cover, sand in soil), two factors that may play an important role in local
adaptation of Arctic Char (Moore et al., 2017; Dubos et al., 2023). Precipitation
was later excluded because it was highly correlated with air temperature
(Pearson r = 0.84), resulting in a final set of 11 environmental predictors (Table
S1, Table S2, Fig. S1).

Population structure

When describing population structure in the dataset, we used a priori
administrative regions to categorize sampling sites across the study area, namely
the Inuvialuit Settlement Region (Northwest Territories); Kitikmeot, Kivalliq and
Baffin (Nunavut); and Nunavik (Québec). We used the term Baffin instead of
Qikiqtaaluk, as this administrative region also includes the High Arctic
Archipelago, which was not sampled as part of this study. Note that anadromous
populations of Arctic Char are less common at those latitudes where resident and
landlocked Arctic Char dominate (Reist et al., 2013). North and South Baffin were
further split around the 67[th] parallel. Note that the Pamiurluk Lake sampling site
(PAM) was grouped within the Kivalliq region, despite technically being in
Qikiqtaaluk, as this fish population is traditionally harvested by the people of
Naujaat, a community in the Kivalliq region.

The overall distribution of genetic variation and population structure was
initially evaluated through a principal component analysis (PCA) conducted on
the LD-pruned dataset using PCAngsd 1.10 (Meisner & Albrechtsen, 2018).
Subsequently, we examined the hierarchical population structure by applying
NGSadmix (Skotte et al., 2013), estimating individual admixture proportions while
varying the number of ancestral populations (K) from 1 to 30 (i.e. the number of
sampling sites). For each value of K, we repeated the analysis over 50
independent runs, then used the CLUMPAK algorithm (Kopelman et al., 2015) to
cluster runs with a similarity score over a fixed threshold of 0.85. We interpreted
the cluster with the most runs (hereafter the major mode) for each value of K.
Ancestry proportions were averaged over runs in the major mode and visualized
using a stacked bar plot. To compare the fit of the model at different values of K,
we 1) compared the proportion of runs in the major mode and 2) computed the
ΔK (Evanno et al., 2005) by dividing the absolute value of the second order rate
of change in the mean likelihood by the standard deviation of likelihoods for all
runs of a given K.

Next, we used ngsDist (Vieira et al., 2016) on the unpruned dataset to
compute genetic distances between every individual pair. We chose the
uncorrected p-distance, i.e. the proportion of nucleotide sites (polymorphic in the
dataset) at which they differ. SNPs with missing data in either individual of the
pair were ignored. To express p-distances as the proportion of genome-wide
differences between two individuals, we multiplied them by the ratio of the
number of SNPs (4,044,588) to the total number of sites in the reference genome
(1,516,033,356). We then built an unrooted neighbour-joining tree on the
distance matrix using the BIONJ algorithm (Gascuel, 1997) in the ape v5.7 R
package (Paradis & Schliep, 2019). We visualized the tree with the R packages
ggtree (Yu et al., 2017) and *ggtreeExtra* (Xu et al., 2021), while collapsing
monophyletic groups of individuals from a single sampling site, then aligned
individual ancestry proportions estimated with ngsAdmix for K = 2, 7, and 30
genetic clusters with the corresponding tree tips. These values of K were chosen
as they represented Arctic vs Atlantic ancestry as detailed in Dallaire et al. (2025)

(K = 2), roughly corresponded to *a priori* geographic regions (K = 7), or matched
the number of sampling sites (K =30).

Connectivity, Isolation-by-distance, Isolation-by-environment

Dallaire et al. (2025) revealed strong patterns of isolation-by-distance that were
distinct in groups of populations above (hereafter northern) and below (southern)
the 67[th] parallel, with southern population pairs being more genetically distant at
equal marine distances. Here, we revisited the relationship between genetic
distance, as measured by the pairwise fixation index (F_{ST}) and allele frequency
difference (AFD), and either marine, environmental, and ancestry distance to
investigate the interplay between i) isolation-by-distance (IBD), ii) isolation-by-
environment (IBE), and iii) isolation-by-colonisation (IBC; defined as the
proportion of genetic distance explained by the difference in genetic background
contributions from two glacial lineages that recolonized the Canadian Arctic),
respectively.

F_{ST}, AFD, and marine distances were estimated as in Dallaire et al.
(2025). Environmental distance between sites was calculated by performing a
PCA on all environmental variables and measuring the Euclidian distance
between population coordinates on the first seven axes, which encompassed
92.2% of the variation. Ancestry distance was measured as the difference in the
average ancestry proportion for a sampling site in the NGSadmix analysis for K =
2, to represent a proxy of the proportion of ancestry from the Arctic and Atlantic
lineages.

The correlations between each measure of genetic distance and either
marine, environmental, or ancestry distances was evaluated separately using
simple Mantel tests as implemented in the R package *vegan* (Oksanen et al.,
2022). We then used partial Mantel tests to test whether each correlation was
robust to controlling for either other type of distance (e.g. AFD ~ marine distance
while controlling for ancestry distance, then for environmental distance, etc.). To
control for the effect of two distance matrices conjointly, we first built a linear

model between the genetic distance and the admixture model while accounting for the category of the population pair (i.e. two northern populations, two southern populations, or one northern and one southern population), and used the residuals as a substitute for genetic distance in a partial Mantel test with marine or environmental distance. To further investigate which component of the environment could contribute to isolation-by-environment, we repeated partial Mantel tests using the difference in the value for each environmental variable as environmental distance. We compared Mantel r statistics and their associated p-values estimated from 999 permutations for tests including all populations pairs, then only northern or southern pairs. We excluded GEO from all Mantel tests due to its high genetic distance from other sites. These diverged from all patterns of isolation, likely due to heavy introgression from the Atlantic lineage in genomes from this population (Dallaire et al., 2025).

To characterize geographic areas of limited connectivity, we conducted an Estimated Effective Migration Surfaces (EEMS) analysis (Petkova et al., 2016) using FEEMS (Marcus et al., 2021). Briefly, the effective migration rate is comparable to the concept of effective population size (N_e), in that it represents the rate of migration at which an idealized stepping stone model at equilibrium would produce genetic dissimilarities equivalent to values observed in the data. In FEEMS, we modified the python scripts *spatial_graph* and *run_cv* to skip the allele frequency estimation from genotypes and used the population MAF estimated in ANGSD as input since we did not call individual genotypes. We used the R package *dggridR* (Barnes & Sahr, 2024) to create a triangular grid at resolution 7 (~ 55km between cells) over the study area. To select the value for the smoothing parameter λ, we followed the leave-one-out cross-validation method suggested in Marcus et al. (2021): we repeated the EEMS analysis after removing one population at a time with 20 values of λ varying from 0.1 to 10,000 and compared the migration surfaces for λ displaying local minima in cross-validation errors.

Loci under putative selection and gene-environment association

SNPs with a global minor allele frequency over 0.05 were investigated for signatures of selection using *pcadapt* (Luu et al., 2017) and the XtX statistic in *Baypass v2.1* (Gautier, 2015), and for Gene-Environment Associations (GEA) using the Bayes Factor statistic in *Baypass* and redundancy analyses (RDA).

We applied the *pcadapt* procedure in PCAngsd, which allows for individual genotype likelihoods as input. Briefly, *pcadapt* scans the genome using significant components of a PCA, here the first 15 PCs, as selected by PCAngsd. The multi-dimensional z-scores in output were converted to Mahalanobis distances using the orthogonalized Gnanadesikan–Kettenring method, corrected for genomic inflation following recommendations (Luu et al., 2017), then converted into p-values via a χ^2 test with 14 degrees of freedom. To build the input for *Baypass*, we converted population allele frequencies estimated in ANGSD (-domaf 1) to allele counts by multiplying (then rounding) the frequencies by the SNP-specific number of individuals to account for missing data. We first sampled 10,000 random SNPs to compute a covariance matrix with the core model in *Baypass*, then used this matrix in the standard covariate model with every SNP (maf > 0.05) and all 11 environmental variables. We ran five replicate analyses and used the median XtX as an index of selection and the median Bayes Factor (BF) as a measure of the association between the allele frequency of a SNP and each environmental variable.

As a second method to detect Gene-Environment Associations, we used redundancy analyses, which are constrained ordinations that assess the relationship between a multivariate response variable (here allele frequencies in sampling sites) and explanatory variables (the environmental factors). To account for collinearity between variables, we ran a first RDA with all variables in the R package *vegan*, then iteratively removed the variable with the highest Variance Inflation Factor (VIF) until all VIF were under 10 (we only removed Dissolved O_2 this way). We then ran a backward selection of model with the function *ordistep*, and iteratively pruned variables from the model until all

variables contribute to the model according to an ANOVA-like permutation test (p
< 0.1). Finally, we assessed the significance of RDA axes through an ANOVA-
like permutation test on the selected model and kept axes with p < 0.05.

Since the ancestry proportion from the Arctic and Atlantic lineages (as
estimated by the NGSadmix analysis for K = 2; see also Dallaire et al. 2025) was
a major determinant of the genetic variation in our study area and strongly
correlated with some of the environmental variables, we also conducted a partial
RDA (pRDA) using the average ancestry proportion as a covariate. We followed
the same variable and axes selection steps as described above. We use the
varpart function in the R package *vegan* to estimate the portions of the genetic
variation that was explained conjunctly and distinctively by environment and
lineage ancestry in the final pRDA.

Every selection scan and GEA methods described above produced
summary statistics by SNP that were aggregated by gene using a windowed Z
analysis (WZA; Booker et al., 2024). WZA combines the information of multiple
SNPs linked in a genomic region to assess if this region is in GEA or displays
signs of selection. Since linkage varies across the genome, and in the absence
of a robust recombination map, we opted to use genes annotated on the
reference genome instead of windows of an arbitrarily defined length. We ran
separate WZAs by using as summary statistics i) the *p*-values from pcadapt, ii)
the XtX and iii) BF (for each environmental variable) from Baypass, and iv) the
squared loading on each significant axis of the RDA and pRDA. Briefly, those
statistics were ranked, ordered and converted to empirical p-values, then to z-
scores that were combined for SNPs inside each gene (i.e. between the start and
end position based on the annotated reference genome) to compute a *p*-value
corrected for the number of SNPs in the gene (Booker et al., 2024).

For each WZA, we listed top candidate genes (i.e. the genes with p-values
< 0.001), thus showing the strongest signal of selection or association with the
environment. To explore the relationship between our top candidate genes and

regions of low recombination as inferred by local PCA in Dallaire et al. (2025), we

conducted a series of χ^2 tests to check if top candidates identified by each

analysis were more likely than by chance to be found in long putative local

ancestry tracts, then in other local PCA outlier regions not correlated with the

admixture gradient between the Arctic and Atlantic (as listed in Table S3).

For each selection scan or GEA analysis, we performed a gene ontology

(GO) enrichment test with goatools 1.2.3 (Klopfenstein et al., 2018) on all top

candidate genes, and then on genes identified as top candidates by more than

one method. We used a custom annotation table, obtained by mapping Arctic

Char transcripts (accession GCF_002910315.2; ASM291031v2) against the

Swissprot 1.2 database (UniProt Consortium, 2024) with blast 2.12.0 (Camacho

et al., 2009), according to the pipeline GAWN v0.3.5

(https://github.com/enormandeau/gawn). As we lacked power to detect selection

in WZA for genes with few SNPs, only genes on which we considered at least 5

SNPs were kept as the background set. Terms for biological processes (BP) with

corrected p-values (Benjamini & Hochberg, 1995) under 0.05 were considered

significantly enriched.

Results

Genetic population structure

Out of the 5.98 million polymorphic SNPs in Canadian and Greenlandic Arctic

Char populations identified in Dallaire et al. (2025), 4.04 million passed the

coverage filters and displayed MAF above 0.01 in Canadian populations alone.

Subsequent LD-pruning reduced this number to 277,570 SNPs. To facilitate

knowledge transfer to local policy makers, populations were first categorized *a*

priori according to administrative regions of Inuit Nunangat: the Inuvialuit

Settlement Region (ISR, Northwest Territories); Nunavik (Québec); and

Kitikmeot, Kivalliq, and Baffin (Nunavut) (Fig. 1a, circles and labels).

A principal component analysis revealed a population structure consistent

with geography on the first two axes (comprising 32.9% of the variation; Fig. 1b).

However, most sites in ISR and Kitikmeot were densely clustered on these axes
and while HRN and KJA were distinct on the fourth axis (3.6%), the rest of north-
western sites remained indistinguishable near the middle of the plot for the first
10 axes (totaling 55.8% of the variation; Fig. S2). A neighbour-joining tree based
on p-distances displayed a similar geographic pattern, with north-western
individuals on shorter branches than south-eastern individuals (Fig. 1c). Many
sampling sites clustered into exclusive groups, with the notable exception of
closely located pairs of sites: EKA and HAL (Kitikmeot; 39 km, F_{ST} = 0.010, Table
S4), AKL and CRB (Kivalliq; 64 km, F_{ST} = 0.011), and SAT and KUL (North
Baffin; 65 km, F_{ST} = 0.014). Interestingly, PAY and CHR were only marginally
further apart (69 km), but exhibited much stronger structure (F_{ST} = 0.078). Finally,
some sampling sites displayed considerable substructure, as their clade was split
into subgroups (e.g., SVG, KUL-SAT) or had a few individuals grouping into
smaller distant clades (e.g., NTS, KGJ).

Figure 1: a) Arctic Char sampling sites with colored labels showing their administrative region of origin. Pie charts show the site average in ancestry proportion for K = 7 genetic clusters, as detailed in panel d). b) The first two axes of a principal component analysis for 277,570 independent SNPs. *95% confidence interval ellipses are drawn around each site and the percentage of variance explained by each axis is noted in parentheses. c) Unrooted neighbor-joining tree built on genome-wide uncorrected p-distances between Arctic Char samples. Monophyletic clades where at least 80% of samples originate from a single site (or a pair of neighbouring sites) are collapsed as triangles and truncated at the minimal length. Sampling sites are delimited by dashed lines and arrows note the origin of samples in smaller clades. Arrows accompanied by fractions denote clades where a portion of individuals were captured at a different site. d) Individual proportion of ancestry for K = 2, 7, and 30 genetics clusters, CLUMPAK-averaged over replicate runs in the major mode (out of 50 runs). Individuals are aligned to their position in the tree in c). Note that this figure reuses the same color palette for two codes: administrative regions in panel a) (labels, circles), b), and c); and their most concordant genetic cluster (K = 7) in panel a) (pie charts) and d). Discordance between administration regions and genetic clusters are noted by an asterisk. Additional colors for the K = 30 plot were assigned randomly.*

The *NGSadmix* analysis of ancestry proportions with *K* ranging from 2 to 30 ancestral populations revealed a highly hierarchical structure (Fig.1d, Fig. S3, Fig. S4). At K = 2, eastern Nunavik and northern sites displayed pure ancestry for different groups, and sites in between followed a cline in admixture closely matching the pattern attributed to Arctic vs. Atlantic lineage ancestry in Dallaire et al. (2025). At K = 30, the analysis discriminated most sampling sites, in line with clades identified in the neighbour-joining tree. Interestingly, K = 7 groups produced a structure parallel to geographic regions defined *a priori* (Fig. 1a, pie charts), with a few key discordances. For example, PAM clustered more closely with Kitikmeot than Kivalliq populations, and ABL aligned with North Baffin rather than Kitikmeot. Most notably, at this level of structure, populations along Hudson Bay and Ungava Bay clustered into different genetic groups, rather than aligning with the Kivalliq and Nunavik regions defined *a priori*. We hereafter refer to groups of sampling sites sharing a dominant genetic cluster at K = 7 as our proposed CUs, with the exception of HRN and KJA who were grouped into the Inuvialuit Settlement Region CUs, as discussed later.

Connectivity and isolation

Pairwise F_{ST} ranged from 0.011 (HAL-EKA) to 0.407 (IQG-GEO), while average allele frequency difference (AFD) in SNPs with MAF > 0.01 ranged from 0.033 (HAL-EKA) to 0.231 (HRN-GEO) (Table S4). The strongest predictor of genetic distance was lineage ancestry difference, with Mantel correlations (r) of 0.92 for F_{ST} and 0.90 for AFD (Fig.2a, Table S5). In comparison, marine geographical distance showed lower correlations (r = 0.65 for F_{ST}, 0.58 for AFD), and environmental distance (summarized by PCA of 13 factors) had even weaker, but still significant (p < 0.01), correlations (r = 0.49 for F_{ST}, 0.51 for AFD) (Table S5. At equal ancestry, marine, or environmental distances, pairs of populations from the southern group were more genetically distant than northern pairs, and the George River (GEO) was strikingly more distant from all other populations than other southern sites.

Figure 2: *a) Linear regression of allele frequency differences (AFD) between a pair of populations and their ancestry distance (i.e. the difference in their average ancestry proportion, NGSadmix, K=2). Pairs were categorized depending on if they include two northern (blue), two southern (red), or one northern and one southern population (green). Pairs including the George River (GEO) are displayed in grey and excluded from panel b-c. Partial Mantel tests were conducted between the residuals of the model from panel a and either b) marine or c) environmental distance for all populations (top), or either northern (middle) or southern ones (below). The remaining distance, i.e. b) environmental or c) marine, was used as covariable and is displayed in color (yellow to blue). Mantel r statistics are noted in bold when the associated p-value was under 0.01.*

Partial Mantel tests indicated no signs of isolation-by-distance in the global

dataset when correcting for both ancestry and environmental distance (Fig.2b, r =

-0.02, p = 0.972), but signs of weak isolation-by-environment when correcting for

ancestry and marine distance (Fig. 2c, r = 0.23, p = 0.025). When repeating

these analyses exclusively for northern or southern pairs, we revealed apparent

isolation-by-distance in the North (Fig. 2b, r = 0.52, p < 0.001), and isolation-by-

environment in the South (Fig. 2c, r = 0.57, p < 0.001). Partial Mantel tests with

specific environmental variables indicated that difference in discharge (r = 0.33)

and primary productivity (r = 0.46) explained a significant portion (p < 0.01) of the

variation in study-wide AFD, while differences in primary productivity (r = 0.47)

and salinity (r = 0.42) explained a significant portion of the variation in AFD in
southern pairs of sites (Table S5, Fig. S5).

We estimated effective migration rates across the study area from allele
frequencies. The resulting migration map at different values of the smoothing
parameter lambda mainly highlights an area of low effective migration between
the previously described northern and southern regions (Fig. 3, Fig. S6). Smaller
areas of low effective migration are observed over the Boothia Peninsula
(between NTS and ABL), Amundsen Gulf (between HRN, KJA), Hudson Strait
(between QNB and FRM), and inland Nunavik (between Hudson Bay and
Ungava Bay). Sampling sites within CUs were generally connected by areas of
effective migration rates higher than 1.

Figure 3: *Edge-specific effective migration rates (w) estimated from sampling allele frequencies in FEEMS with a smoothing parameter λ = 21.5. Sampling sites were repositioned to the nearest node on a triangular grid (cell spacing ≈ 55 km) and colored with the proposed conservation unit, following the most dominant genetic cluster in an NGSadmix analysis (K = 7), except for Inuvialuit S. R. (see Discussion for details).*

452 Selection scan and Gene-environment association

453 We investigated 2.6 million SNPs with MAF over 0.05 for signatures of selection,
454 via pcadapt and the XtX statistic in Baypass (details below); or their association
455 with environmental variables, with redundancy analyses without (RDA) and with

(pRDA) correction for lineage ancestry, and the Bayes Factor statistic in
Baypass.

A global RDA with all 11 selected environmental factors and Arctic-Atlantic
lineage ancestry (as estimated by the average ancestry proportion inferred with
NGSadmix at K=2) explained 62.4% of the variation in MAF between sampling
sites (Fig. 4a). Environmental factors explained 53.3% of the variation, but a
large part (38.4%) was conjointly explained by lineage ancestry, as further
explored by the following RDA and pRDA.

Figure 4: a) Proportion of the variation in allele frequency explained independently and conjointly by a set of 11 environmental variables and glacial lineage ancestry, as estimated by NGSadmix (K = 2). Biplots of b) a redundancy analysis (RDA, 2 significant axes) and c-d) a partial RDA accounting for the variation in ancestry (pRDA, 3 significant

After backward variable selection with ordistep, the first two axes of the

RDA (adjusted R^2 = 0.36) were significant (p < 0.05) and were largely correlated

with 1) air temperature and ice cover; and 2) sea surface temperature (Fig. 4b).

The first three axes of the pRDA (adjusted R^2 = 0.12) were significant and were

correlated with 1) sea-surface temperature, lake cover, fraction of sand in soil,

and slope; 2) discharge and primary productivity; and 3) primary productivity and

ice cover (Fig 4c-d). RDA1 was highly correlated with latitude and lineage

ancestry, while RDA2 and pRDA1 discriminated Baffin Island sites from the rest

of the study area. Both RDA2 and pRDA2 also highlighted the genetic and

environmental distinction between western (Hudson Bay: TJL, KOA) and eastern

(Ungava Bay: PAY, CHR, GEO) Nunavik.

With a windowed Z analysis (WZA), we combined SNP-specific statistics

from the 5 methods (a total of 18 tests: pcadapt, XtX, 2 RDA axes, 3 pRDA axes,

and 11 univariate tests for the BF statistics in Baypass) for 21,379 genes (with at

least 5 SNPs) to provide rank-ordered support for selection or GEA, in the form

of empirical p-values (Fig. 5a). The genome scan methods (pcadapt and XtX)

identified respectively 151 and 188 top candidate genes (p < 0.001), with

significant overlap (81, or 31.2% of genes in common, Fig. 5b). The GEA

methods (BF, RDA, and pRDA) identified 2,079, 381, and 544 top genes. BF had

notable overlap with the other methods: 164 (7.1%) of top genes were in

common with the RDA and 250 (10.5%) with the pRDA (Fig. 5b). The RDA and

pRDA had a similar level of overlap, with 57 top genes in common (6.6%). Top

candidate genes from the XtX, pcadapt method or associated to pRDA1 or slope

had more chance to be found in putative local ancestry tracts identified in Dallaire

et al. (2025), while other local PCA outlier regions were enriched in pcadapt,

RDA2, pRDA1, Lake cover, and Sea T° top candidate genes (χ^2 test, adjusted p

< 0.05; Table S6).

Figure 5: *a) Support for selection scans or gene-environment associations on genes with at least 10 SNPs on 4 linkage groups, in 5 panels regrouping methods: 1) XtX (Baypass) or pcadapt; 2) redundancy analysis axes (RDA); 3) partial RDA axes; and Bayes Factors (BF, Baypass) for 4) freshwater and 5) marine variables. Support at the gene level (empirical p-values) was derived from combined per-SNP summary statistics in a windowed Z analysis (WZA). Top candidate genes (p < 0.001) for more than one type of test were highlighted with bigger diamonds than the other genes (small dots). b) Venn diagrams presenting the number of top candidate genes (p < 0.001) for selection (top) or gene environment association (bottom) and the overlap in genes between methods. c) Karyotype frequencies per sampling site for a 290kb-long putative inversion on LG13, as inferred by local PCA in Dallaire et al. (2025). The genomic position of the putative inversion is noted by an arrow in panel a.*

The higher number of top genes identified by the BF method is linked to
the univariate nature of the method, as top-ranking candidate genes were
separately identified based on their correlation with each of the 11 environmental
variables. Individual variables had on average 262.1 (sd = 14.7) associated top
genes (empirical p < 0.001 in the WZA). Out of 2,079 unique BF top genes, 571
(27.4%) were associated with more than one variable (404 with two variables,
120 with three, 47 with four or more). Salinity and dissolved O_2 (55 top genes in
common), followed by lake cover and slope (19 genes), were the most frequent
pairs.

Across all genome scans and GEA methods, 549 out of 2,634 unique top
candidate genes (20.8 %) were identified by more than one method (Table S7,
Fig. S7). We tested for enrichment in gene ontology (GO) terms in lists of outlier
genes from each individual test, as well as the list of genes that were outliers
according to at least two methods and identified 89 unique GO terms with a p-
value for enrichment under 0.001 and associated with at least three top
candidate genes (Table S8). However, after correction for multiple testing, no GO
terms presented corrected p-values under 0.1.

Discussion

In this study, we aimed to define conservation units for anadromous populations
of Arctic Char across its continuous distribution in the Inuvialuit Settlement
Region, Nunavut, and Nunavik in Arctic Canada. Through analysis of population
structure and connectivity, we highlight two major population groupings,
respectively above and below the 67[th] parallel, which we interpret as having
distinct evolutionary trajectories and reduced gene flow. These northern and
southern groups were further subdivided in six candidate CUs (see Fig. 3) based
on genetic similarity and connectivity. While we observed interconnected patterns
of isolation-by-distance and isolation-by-environment across the landscape,
glacial lineage ancestry was the strongest predictor of genetic distance among
populations. Using genome scans and GEA, we detected a polygenic and

complex signal of selection in the face of multiple selective pressures,
highlighting putatively adaptive variation between the six candidate CUs.

Hierarchical population structure informs conservation units

As a first step toward delimiting conservation units for anadromous Arctic Char,
we used a combination of common methods to assess the genetic population
structure of our 30 sampling sites. This approach revealed a complex but
geographically coherent and hierarchical picture of genetic structure.

At the broadest level, visible in the PCA (PC1) and NGSadmix (K = 2)
results, the system is defined by an admixture cline from the Arctic lineage in the
northwest to the Atlantic lineage in the southeast, as explored in detail in Dallaire
et al. (2025). According to mitochondrial data, the divergence time between those
lineages could reach up to 400,000 years, predating many glacial cycles
(Jacobsen et al., 2022). Regardless of whether admixture occurred during
previous interglacial periods, a prolonged phase of allopatry allowed these
lineages to diverge significantly, which explains why much of the genetic
variation observed in our current dataset reflects this secondary contact. This
pattern provides an initial framework for delineating conservation units, as it
reveals two distinct evolutionary trajectories: Northern sites are dominated by the
Arctic lineage, while Southern sites exhibit varying degrees of admixture between
the Arctic and Atlantic lineages. The distinctiveness between these two
population segments is also supported by a low effective migration surface
between the Kitikmeot and Kivalliq region, as well as Northern and Southern
Baffin Island. Although the Northern and Southern populations do not represent
entirely pure lineages, we argue that they should be managed as separate
broad-scale conservation units. The persistence of this historical signal in
contemporary patterns of variation suggests that secondary contact has not
erased the imprint of prolonged allopatry. This dominance of ancient divergence
over more recent demographic processes is particularly striking in the Arctic,
where deglaciation occurred only a few thousand years ago (Dalton et al., 2020),
and resembles findings in other post-glacial populations of fishes such as

salmonids (Lehnert et al., 2019) and sticklebacks (Kirch et al., 2021), where deep
lineage histories continue to structure modern diversity.

However, our analysis of structure revealed additional hierarchical levels
of population structure within the North and South groups, where most individuals
showed pure ancestry to a genetic cluster specific to its sampling site. While the
ΔK summary statistic suggested 2 groups, studies have shown that this statistic
is strongly biased toward K = 2 (Janes et al., 2017). In our study, this is not
surprising given the overwhelming signal of lineage ancestry that this level of
structure represents. Considering that ΔK was stable from K = 4 to 30 and that
significant structure between rivers is expected according to the philopatric
nature of Arctic Char reproductive migrations (Gyselman, 1994; Moore et al.,
2013; Dallaire et al., 2021), we argue that all 30 clusters identified by NGSadmix
could represent populations with some degree of demographic independence.
These results highlight the strong hierarchical nature of genetic structure in Arctic
Char, as observed in other anadromous salmonids (e.g. Beacham et al., 2004;
Habicht et al., 2007; Dionne et al., 2008; Gilbert-Horvath et al., 2016; Östergren
et al., 2021). Importantly, this fine-scale, river-specific structure aligns with
current fisheries management practices, which often implement river-specific
quotas and management measures to reflect the biological reality of Arctic Char
populations.

Our sampling strategy focused on broad coverage across the Canadian
range of anadromous Arctic Char rather than fine-scale resolution. As such, the
six candidate CUs proposed here each cover coastal areas ranging for hundreds
to thousands of km, which is much larger than current CUs in other salmonids in
southern Canada (e.g., Xuereb et al., 2022; Lehnert et al., 2023). However, our
analyses still uncovered patterns that may hold significance for local
management. For example, we identified evidence of straying between most
pairs of rivers inside 100 km from each other, and in one instance, up to 400 km
(one fish sampled at EKA was genetically closer to fish in COP). Despite their
philopatric behavior, Arctic Chars are known to show less fidelity to their natal

site during the upstream migration in non-reproductive years (Gyselman, 1994; Klemetsen et al., 2003; Moore et al., 2013). When migrating strictly for overwintering purposes, fish might select watersheds with easily accessible winter habitats, for example the Ekalluk system in the Kitikmeot Region (EKA; Moore et al., 2017). In the case of AKL and CRB (Kivalliq), fish from either site were indistinguishable using whole-genome data, which suggests either panmixia between the two systems or that sampling at these sites is not targeting individuals during the upstream reproductive migration, but rather a mixed-stock of poorly differentiated source populations. Proper delimitation of MUs in Arctic Char would warrant additional local studies of genetic structure and migration, as was implemented in areas sustaining commercial and subsistence fisheries, such as in Cambridge Bay (Harris, Moore, et al., 2016; Moore et al., 2017), Cumberland Sound (Harris et al., 2014), Paulatuk (Harris, Boguski, et al., 2016), and Ulukhaktok (Lea et al., 2023). Going forward, whole-genome sequencing is likely not the most cost-effective method to tackle such local-scale questions. Alternatively, SNP panels can be developed to assign fish caught in mixed-stock fisheries to their population of origin, which holds promise for the accurate monitoring of fish stocks and the conciliation of commercial and subsistence fishing (Beemelmanns et al., 2025).

We chose K = 7 as an intermediate level of structure that could be used to delimit candidate CUs in Canadian anadromous Arctic Char. This level of structure subdivides our top-level Northern and Southern CUs, described earlier, and creates genetic groupings that conveniently, but somewhat unexpectedly, closely match the administrative regions of the Canadian Arctic (Fig. 1). A notable exception, however, is the union of both the Kivalliq (western) and Nunavik (eastern) coasts of the Hudson Bay in a single cluster (both coasts display signs of common ancestry until K = 17), while the rest of the Nunavik populations, around Ungava Bay, appeared as a distinct unit. Considering that K = 6 and K = 7 discriminated HRN and KJA (Fig. S4), our two westernmost sites, we suggest that these two could be paired to form a sixth CU, until additional sampling is performed in the Inuvialuit Settlement Region.

The choice of a level of genetic structure to define CUs implies subjectivity
to some degree (Waples, 1995), as genetic differentiation typically occurs along
a continuum rather than in clear-cut divisions. In the case of Arctic Char, many
populations at the edge of candidate CUs display admixed ancestry to multiple
clusters. However, the borders of the proposed CUs were supported by shifts in
the dominant cluster that were concordant with areas of low effective migration in
our EEMS analyses, as was for example observed in multiple marine species in
Atlantic Canada (Wilcox et al., 2023). In the present study, candidate boundaries
included the Boothia Peninsula between the Kitikmeot and the North Baffin CUs,
and the Foxe Basin and Hudson Strait between the South Baffin and Hudson
Bay/Ungava Bay CUs. While EEMS areas of low effective migration do not
necessarily imply physical barriers to gene flow (Petkova et al., 2016), they
should coincide with shifts in allele frequency that are relevant for informing
conservation actions.

Detecting local adaptation using whole-genome data
We further used our genomic dataset to detect signals of local adaptation in
Arctic Char populations to support the delineation of CUs that also reflect
patterns of local adaptation to divergent environments (sensu Funk et al., 2012).
By using multiple analyses on millions of genetic markers, we obtained an
overwhelmingly polygenic signal at the SNP-level, with numerous peaks of
differentiation or associations with the environment across all linkage groups. To
summarise this amount of noise and account for the high rate of false positives in
the detection of adaptation, we 1) opted for a Windowed Z Analysis (WZA) to
identify top candidate genes while accounting for linkage, and 2) focused on
genes that were top candidates in multiple analyses.

As such, we were interested in the relationship between signals of local
adaptation and known genomic regions of low recombination, as these might
either be the result of hard selective sweeps or genomic features such as
inversions or centromeres (Lotterhos, 2019). Our previous work inferred the
position of many long haplotype blocks (i.e. sequences containing many

polymorphisms inherited as a unit), suggesting a very low recombination rate

within those segments (Dallaire et al. 2025). The haplotype frequencies of many

of those putative haploblocks were heavily correlated with the admixture gradient

between the Arctic and Atlantic glacial lineages. This implies that they act as

local ancestry tracts and are much more constrained by post-glacial demography

than spatial variation in selection, though the two are not mutually exclusive. In

the partial RDA, we included each population's average ancestry proportion as a

covariate in the model, so that the analysis only tests for associations between

genotypes and environment once the effect of lineage admixture has been

factored out. Surprisingly, both putative local ancestry tracts and other local PCA

outliers (identified in Dallaire et al. 2025) were enriched in top candidates from

pRDA1, despite the partial RDA specifically controlling for the effect of Arctic-

Atlantic ancestry.

However, RDA2 and pRDA1 identified common top candidates in a region

of the LG13 identified as a potential inversion in Dallaire et al. (2025) (Fig 5a,c).

This 290 kb region overlaps seven genes, including the insulin-like 5a and a

protein regulating mitochondrial genes, hinting at their importance in metabolism

or energy production. The alternative karyotype for this putative inversion is

present in higher frequencies in both North and South Baffin, which present

contrasted habitats compared to the rest of the Canadian Arctic, with fjord-like

topography (watersheds with a higher slope and lower lake cover) and colder

sea-surface temperatures. It is thus possible that this putative inversion could

play a role in adaptation to these environments. Chromosomal inversions

suppress recombination and are known to promote co-adaptation of

neighbouring genes in the face of gene flow (Dobzhansky & Sturtevant, 1938;

Kirkpatrick & Barton, 2006; Wellenreuther et al., 2019). This has previously been

documented in rainbow trout (*Onchorhyncus mykiss*, Pearse et al., 2019),

Atlantic Cod (*Gadus morhua*, Matschiner et al., 2022), Atlantic Herring (*Clupea*

harengus, Jamsandekar et al., 2024), and Atlantic Silverside (*Menidia menidia*,

Tigano et al., 2021). Another larger (1.2 Mb) candidate inversion on LG12 was

found to be polymorphic in Arctic Char populations from Kitikmeot (Hale et al.,

2021), but we did not find well-supported top candidate genes in this genomic
region.

When using multiple genome scans and GEA methods with their
respective biases, giving greater weight to markers identified by more than one
method can help to reduce false positives (Forester et al., 2018). By doing so, we
reduced our list of top candidates from 2,635 to 541 genes widely distributed
across the genome. Despite low statistical support for GO enrichment after
correction for multiple testing, we identified key gene functions related to lipid
homeostasis and catabolic processes, which ranked among the best-supported
GO categories across multiple sets of candidate genes. Lipid metabolism plays a
critical role in the adaptation to Arctic climates, supporting membrane fluidity in
cold-water ectotherms such as fishes (Ernst et al., 2016; Wang et al., 2021) and
energy reserves during extreme dietary challenges such as prolonged fasting
and hibernation (Olsen et al., 2021). The importance of fatty acid homeostasis in
local adaptation was also observed in subarctic Atlantic Salmon populations,
where genes regulating lipid metabolism were identified as drivers of adaptation
(Lehnert et al., 2023).

In summary, identifying the biological processes and molecular functions
involved in polygenic adaptation to multiple selective pressures remains
challenging (Pritchard and Di Rienzo 2010; Le Corre and Kremer 2012; Rees et
al. 2020). This is particularly true when using indirect inference of selection in the
absence of phenotypic data on species with complex life cycles. Here, genome-
wide, high-density SNP data provided unmatched resolution in the detection of
signals of selection but required considerable data aggregation (e.g., through the
WZA, as recently used in plants, Battlay et al., 2023; and *Drosophila*, Nunez et
al., 2024) to make the results interpretable. This supported the importance of
considering the relation between selection and recombination in our data, as
highly linked regions of the genome could either be a consequence of selective
sweeps or could have biased the detection of adaptation.

Putatively adaptive variation reinforces candidate CUs

Despite challenges in detecting specific targets for selection, we used the overall putatively adaptive genomic variation to highlight groups of populations that could show local adaptation to similar habitats, thus supporting the candidate Conservation Units described earlier. For example, the distribution of allele frequencies in relation to environmental variation, as described by the pRDA (Fig. 4c-d), emphasizes one of our main findings—that a distinction based on water bodies in the Ungava and Hudson Bay groups was more significant than one based on landmasses (i.e., Nunavik vs. Kivalliq). These waterbodies mainly differed in coastal sea-surface temperature and salinity, and watersheds flowing in them are topographically distinct: Nunavik watersheds are generally less steep and have higher lake cover compared with those in South Baffin, our third recommended Conservation Unit in the South Arctic region.

In the North, environmental and geographical distances between populations were too confounded to support isolation-by-environment, but putatively adaptive genetic variation supported the distinction between the Kivalliq and North Baffin CUs, as well as the inclusion of the Abernathy Lake (ABL) population, east of the Boothia Peninsula, in the North Baffin CU. Similar to our results on genome-wide variation, we observed a weaker shift in allele frequencies associated with the environment between Kitikmeot and Inuvialuit Settlement Region populations, putting into question the distinction between those two candidate CUs. However, following a precautionary approach, we recommend maintaining these population segments as separate CUs, with boundaries that could be refined through additional sampling at the western limit of our study area.

Across our study area, conclusions based on genome-wide, neutral, and putatively adaptive genetic variation were largely consistent, with the adaptive data reinforcing the neutral patterns rather than providing novel insights. While one of the promises of population genomics was to reveal previously undetected adaptive groups that could be considered for conservation (Funk et al., 2012),

there is growing evidence that genome-wide variation effectively predicts
adaptive variation in many systems (Chhina et al., 2024). This was observed in
other anadromous salmonids like Atlantic Salmon (Moore et al., 2014; Lehnert et
al., 2023) and Coho Salmon (Xuereb et al., 2022), and this observation could be
influenced by a few key factors. First, polygenic traits are expected to create
weak selection signals (as identified in this study) that do not necessarily differ
from neutral population structure (Pritchard et al., 2010). In contrast, large-effect
loci such as GREB1L in steelhead (anadromous *O. mykiss*) and Chinook Salmon
(*O. tshawytscha*), linked to migration timing, and vgll3 in Atlantic Salmon (*Salmo*
salar), associated with age at maturity, underlie key phenotypes that are now
explicitly considered in conservation and management (Waples et al., 2022).

Second, isolation-by-distance patterns suggest limited dispersal over large
geographic scales (Aguillon et al., 2017) and straying is likely more frequent
among watersheds sharing environmental characteristics. This could be either
because of migratory cues such as water chemistry that differ among regional
river systems (Dittman et al., 1996) or because of selection against maladapted
migrants (Peterson et al., 2014) at local scales. There is in fact a long-standing
hypothesis that homing evolved in anadromous salmonids as it promotes
advantageous local adaptations (Quinn, 1993; McDowall, 2001; Keefer & Caudill,
2014). In contrast to the long marine migrations of other anadromous salmonids
to offshore feeding areas, Arctic Char feed in coastal waters close to their natal
rivers in summer (Dempson & Kristofferson, 1987; Spares et al., 2015; Moore et
al., 2016), extending the potential for local adaptation to marine habitats in the
species (Dallaire et al., 2021). As both freshwater and marine environments tend
to be spatially autocorrelated (Legendre, 1993), it is not surprising that a
significant part of the putatively adaptive genetic variation would be concordant
with the neutral structure.

In this study, we reanalyzed genomic data from Arctic Char across the
Canadian Arctic and provided a detailed assessment of population structure and
connectivity. From this analysis, we identified six candidate Conservation Units

(CUs) within two major groups displaying distinct evolutionary trajectories. This framework offers an important baseline for understanding intraspecific diversity at a national level, which can be further refined as additional data become available. Incorporating environmental, ecological, and life-history information alongside genomic insights will be essential for improving the resolution of these units and ensuring their alignment with meaningful biological boundaries. In that regard, Inuit ecological traditional knowledge will be invaluable as it includes detailed and fine-scale observations on distinct traits between population, e.g. size, taste and color, migratory behavior, and phenology.

Ultimately, this refined understanding of Arctic Char population structure can guide resource management strategies, ensuring the protection of locally adapted populations and their associated roles in the ecology of northern freshwater ecosystems, their importance for food security, and their contribution to the development of a sustainable economy in the Canadian Arctic. Future analyses that integrate genomic baselines with monitoring programs, harvest data, and indigenous traditional knowledge could improve the practical implementation of CUs. Such an approach would allow managers to better anticipate shifts in population dynamics, adjust harvest regulations proactively, and strengthen co-management frameworks with northern communities.

Data Archiving Statement

All *Salvelinus alpinus* raw sequencing data analysed in this manuscript is available on Short Read Archive as part of project PRJNA1031558. A bioinformatical pipeline presenting the main lcWGS analyses is available at: https://github.com/xav9536/angsd_pipeline.

Reference

Aguillon, S. M., Fitzpatrick, J. W., Bowman, R., Schoech, S. J., Clark, A. G., Coop, G., & Chen, N. (2017). Deconstructing isolation-by-distance: The genomic consequences of limited dispersal. *PLoS Genetics*, *13*(8), 1–27. https://doi.org/10.1371/journal.pgen.1006911

Assis, J., Fernández Bejarano, S. J., Salazar, V. W., Schepers, L., Gouvêa, L.,
Fragkopoulou, E., Leclercq, F., Vanhoorne, B., Tyberghein, L., Serrão, E.
A., Verbruggen, H., & De Clerck, O. (2024). Bio-ORACLE v3.0. Pushing
marine data layers to the CMIP6 Earth System Models of climate change
research. *Global Ecology and Biogeography*, *33*(4), e13813.
https://doi.org/10.1111/geb.13813
Avise, J. C., Arnold, J., Ball, R. M., Bermingham, E., Lamb, T., Neigel, J. E.,
Reeb, C. A., & Saunders, N. C. (1987). Intraspecific Phylogeography: The
Mitochondrial DNA Bridge Between Population Genetics and Systematics.
Annual Review of Ecology and Systematics, *18*, 489–522.
Barnes, R., & Sahr, K. (2024). *dggridR: Discrete Global Grids*. https://CRAN.R-
project.org/package=dggridR
Battlay, P., Wilson, J., Bieker, V. C., Lee, C., Prapas, D., Petersen, B., Craig, S.,
Van Boheemen, L., Scalone, R., De Silva, N. P., Sharma, A.,
Konstantinović, B., Nurkowski, K. A., Rieseberg, L. H., Connallon, T.,
Martin, M. D., & Hodgins, K. A. (2023). Large haploblocks underlie rapid
adaptation in the invasive weed Ambrosia artemisiifolia. *Nature
Communications*, *14*(1), 1717. https://doi.org/10.1038/s41467-023-37303-
4
Beacham, T. D., Lapointe, M., Candy, J. R., McIntosh, B., MacConnachie, C.,
Tabata, A., Kaukinen, K., Deng, L., Miller, K. M., & Withler, R. E. (2004).
Stock Identification of Fraser River Sockeye Salmon Using Microsatellites
and Major Histocompatibility Complex Variation. *Transactions of the
American Fisheries Society*, *133*(5), 1117–1137.
https://doi.org/10.1577/T04-001.1
Beemelmanns, A., Bouchard, R., Michaelides, S., Normandeau, E., Jeon, H.-B.,
Chamlian, B., Babin, C., Hénault, P., Perrot, O., Harris, L. N., Zhu, X.,
Fraser, D., Bernatchez, L., & Moore, J.-S. (2025). Development of SNP
Panels from Low-Coverage Whole Genome Sequencing (lcWGS) to
Support Indigenous Fisheries for Three Salmonid Species in Northern
Canada. *Molecular Ecology Resources*, e14040.
https://doi.org/10.1111/1755-0998.14040
Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A
Practical and Powerful Approach to Multiple Testing. *Journal of the Royal
Statistical Society: Series B (Methodological)*, *57*(1), 289–300.
https://doi.org/10.1111/j.2517-6161.1995.tb02031.x
Booker, T. R., Yeaman, S., Whiting, J. R., & Whitlock, M. C. (2024). The WZA□:
A window□based method for characterizing genotype–environment
associations. *Molecular Ecology Resources*, *24*(2), e13768.
https://doi.org/10.1111/1755-0998.13768
Brubacher. (2004). *An overview of Nunavut fisheries* (p. 93) [Report].
Government of Nunavut.
Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K.,
& Madden, T. L. (2009). BLAST+: architecture and applications. *BMC
Bioinformatics*, *10*(1), 421. https://doi.org/10.1186/1471-2105-10-421

Chhina, A. K., Fernandez-Fournier, P., Lewthwaite, J., Booker, T., & Mooers, A.
(2024). *Data from: Does genome-wide variation and putatively adaptive*
variation identify the same set of distinct populations? (Version 4, p.
271770344 bytes) [Dataset]. Dryad.
https://doi.org/10.5061/DRYAD.NVX0K6F1J
Christensen, K. A., Rondeau, E. B., Minkley, D. R., Leong, J. S., Nugent, C. M.,
Danzmann, R. G., Ferguson, M. M., Stadnik, A., Devlin, R. H., Muzzerall,
R., Edwards, M., Davidson, W. S., & Koop, B. F. (2021). Retraction: The
Arctic charr (Salvelinus alpinus) genome and transcriptome assembly.
PLOS ONE, *16*(2), e0247083.
https://doi.org/10.1371/journal.pone.0247083
Coates, D. J., Byrne, M., & Moritz, C. (2018). Genetic Diversity and Conservation
Units: Dealing With the Species-Population Continuum in the Age of
Genomics. *Frontiers in Ecology and Evolution*, *6*.
https://www.frontiersin.org/articles/10.3389/fevo.2018.00165
COSEWIC. (2010). *COSEWIC assessment and status report on the Atlantic*
Salmon Salmo salar (Nunavik population, Labrador population, Northeast
Newfoundland population, South Newfoundland population, Southwest
Newfoundland population, Northwest Newfoundland population, Quebec
Eastern North Shore population, Quebec Western North Shore population,
Anticosti Island population, Inner St. Lawrence population, Lake Ontario
population, Gaspé-Southern Gulf of St. Lawrence population, Eastern
Cape Breton population, Nova Scotia Southern Upland population, Inner
Bay of Fundy population, Outer Bay of Fundy population) in Canada (p.
xlvii + 136). Committee on the Status of Endangered Wildlife in Canada.
Dallaire, X., Bouchard, R., Hénault, P., Ulmo-Diaz, G., Normandeau, E., Mérot,
C., Bernatchez, L., & Moore, J.-S. (2023). Widespread Deviant Patterns of
Heterozygosity in Whole-Genome Sequencing Due to Autopolyploidy,
Repeated Elements, and Duplication. *Genome Biology and Evolution*,
15(12), evad229. https://doi.org/10.1093/gbe/evad229
Dallaire, X., Normandeau, E., Brazier, T., Harris, L., Michael, M. M. H., Mérot, C.,
& Moore, J.-S. (2025). Leveraging whole genomes, mitochondrial DNA,
and haploblocks to decipher complex demographic histories: an example
from a broadly admixed arctic fish. *Molecular Ecology*.
https://doi.org/10.1101/2024.12.11.628006
Dallaire, X., Normandeau, É., Mainguy, J., Tremblay, J.-É., Bernatchez, L., &
Moore, J.-S. (2021). Genomic data support management of anadromous
Arctic Char fisheries in Nunavik by highlighting neutral and putatively
adaptive genetic variation. *Evolutionary Applications*, *n/a*(n/a).
https://doi.org/10.1111/eva.13248
Dalton, A. S., Margold, M., Stokes, C. R., Tarasov, L., Dyke, A. S., Adams, R. S.,
Allard, S., Arends, H. E., Atkinson, N., Attig, J. W., Barnett, P. J., Barnett,
R. L., Batterson, M., Bernatchez, P., Borns, H. W., Breckenridge, A.,
Briner, J. P., Brouard, E., Campbell, J. E., … Wright, H. E. (2020). An
updated radiocarbon-based ice margin chronology for the last deglaciation

of the North American Ice Sheet Complex. *Quaternary Science Reviews*,
234. https://doi.org/10.1016/j.quascirev.2020.106223
Dauphin, B., Rellstab, C., Wüest, R. O., Karger, D. N., Holderegger, R., Gugerli,
F., & Manel, S. (2023). Re-thinking the environment in landscape
genomics. *Trends in Ecology & Evolution*, *38*(3), 261–274.
https://doi.org/10.1016/j.tree.2022.10.010
Dempson, J. B., & Kristofferson, A. H. (1987). Spatial and Temporal Aspects of
the Ocean Migration of Anadromous Arctic Char. *American Fisheries*
Society Symposium, *1*, 340–357.
DFO. (2024). *Pacific Salmon Outlook: Pacific Region*. Fisheries and Oceans
Canada. https://waves-vagues.dfo-mpo.gc.ca/library-
bibliotheque/41263625.pdf
Dionne, M., Caron, F., Dodson, J. J., & Bernatchez, L. (2008). Landscape
genetics and hierarchical genetic structure in Atlantic salmon: The
interaction of gene flow and local adaptation. *Molecular Ecology*, *17*(10),
2382–2396. https://doi.org/10.1111/j.1365-294X.2008.03771.x
Dittman, A. H., Quinn, T. P., & Nevitt, G. A. (1996). Timing of imprinting to natural
and artificial odors by coho salmon (*Oncorhynchus kisutch*). *Canadian*
Journal of Fisheries and Aquatic Sciences, *53*(2), 434–442.
https://doi.org/10.1139/f95-185
Dobzhansky, T., & Sturtevant, A. H. (1938). Inversions in the Chromosomes of
Drosophila Pseudoobscura. *Genetics*, *23*(1), 28–64.
Dubos, V., St-Hilaire, A., & Bergeron, N. E. (2023). Fuzzy logic modelling of
anadromous Arctic char spawning habitat from Nunavik Inuit knowledge.
Ecological Modelling, *477*, 110262.
https://doi.org/10.1016/j.ecolmodel.2022.110262
Ellegren, H. (2004). Microsatellites: simple sequences with complex evolution.
Nature Reviews Genetics, *5*(6), 435–445. https://doi.org/10.1038/nrg1348
Ernst, R., Ejsing, C. S., & Antonny, B. (2016). Homeoviscous Adaptation and the
Regulation of Membrane Lipids. *Journal of Molecular Biology*, *428*(24,
Part A), 4776–4791. https://doi.org/10.1016/j.jmb.2016.08.013
Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters
of individuals using the software STRUCTURE□: a simulation study.
Molecular Ecology, *14*(8), 2611–2620. https://doi.org/10.1111/j.1365-
294X.2005.02553.x
Forest, F., Grenyer, R., Rouget, M., Davies, T. J., Cowling, R. M., Faith, D. P.,
Balmford, A., Manning, J. C., Procheş, Ş., van der Bank, M., Reeves, G.,
Hedderson, T. A. J., & Savolainen, V. (2007). Preserving the evolutionary
potential of floras in biodiversity hotspots. *Nature*, *445*(7129), 757–760.
https://doi.org/10.1038/nature05587
Forester, B. R., Lasky, J. R., Wagner, H. H., & Urban, D. L. (2018). Comparing
methods for detecting multilocus adaptation with multivariate genotype–
environment associations. *Molecular Ecology*, *27*(9), 2215–2233.
https://doi.org/10.1111/mec.14584

Fox, E. A., Wright, A. E., Fumagalli, M., & Vieira, F. G. (2019). ngsLD: evaluating
linkage disequilibrium using genotype likelihoods. *Bioinformatics*, *35*(19),
3855–3856. https://doi.org/10.1093/bioinformatics/btz200
Frankham, R., Ballou, J. D., Dudash, M. R., Eldridge, M. D. B., Fenster, C. B.,
Lacy, R. C., Mendelson, J. R., Porton, I. J., Ralls, K., & Ryder, O. A.
(2012). Implications of different species concepts for conserving
biodiversity. *Biological Conservation*, *153*, 25–31.
https://doi.org/10.1016/j.biocon.2012.04.034
Fraser, D. J., & Bernatchez, L. (2001). Adaptive evolutionary conservation:
Towards a unified concept for defining conservation units. *Molecular*
Ecology, *10*(12), 2741–2752. https://doi.org/10.1046/j.1365-
294X.2001.t01-1-01411.x
Friesen, T. M. (2004). Contemporaneity of Dorset and Thule Cultures in the
North   American Arctic: New Radiocarbon Dates from
Victoria Island,   Nunavut. *Current Anthropology*, *45*(5), 685–
691. https://doi.org/10.1086/425635
Funk, W. C., McKay, J. K., Hohenlohe, P. A., & Allendorf, F. W. (2012).
Harnessing genomics for delineating conservation units. *Trends in*
Ecology and Evolution, *27*(9), 489–496.
https://doi.org/10.1016/j.tree.2012.05.012
Garnett, S. T., & Christidis, L. (2017). Taxonomy anarchy hampers conservation.
Nature, *546*(7656), 25–27. https://doi.org/10.1038/546025a
Gascuel, O. (1997). BIONJ: an improved version of the NJ algorithm based on a
simple model of sequence data. *Molecular Biology and Evolution*, *14*(7),
685–695. https://doi.org/10.1093/oxfordjournals.molbev.a025808
Gautier, M. (2015). Genome-wide scan for adaptive divergence and association
with population-specific covariates. *Genetics*, *201*(4), 1555–1579.
https://doi.org/10.1534/genetics.115.181453
Gilbert-Horvath, E. A., Pipal, K. A., Spence, B. C., Williams, T. H., & Garza, J. C.
(2016). Hierarchical Phylogeographic Structure of Coho Salmon in
California. *Transactions of the American Fisheries Society*, *145*(5), 1122–
1138. https://doi.org/10.1080/00028487.2016.1201003
Green, D. M. (2005). Designatable Units for Status Assessment of Endangered
Species. *Conservation Biology*, *19*(6), 1813–1820.
https://doi.org/10.1111/j.1523-1739.2005.00284.x
Gyselman, E. C. (1994). Fidelity of Anadromous Arctic Char. *Canadian Journal of*
Fisheries and Aquatic Sciences, *51*, 1927–1934.
Habicht, C., Seeb, L. W., & Seeb, J. E. (2007). Genetic and Ecological
Divergence Defines Population Structure of Sockeye Salmon Populations
Returning to Bristol Bay, Alaska, and Provides a Tool for Admixture
Analysis. *Transactions of the American Fisheries Society*, *136*(1), 82–94.
https://doi.org/10.1577/T06-001.1
Hale, M. C., Campbell, M. A., & McKinney, G. J. (2021). A candidate
chromosome inversion in Arctic charr (*Salvelinus alpinus*) identified by
population genetic analysis techniques. *G3 Genes|Genomes|Genetics*,
jkab267. https://doi.org/10.1093/g3journal/jkab267

Harris, L. N., Boguski, D. A., Gallagher, C. P., & Howland, K. L. (2016). Genetic
Stock Identification and Relative Contribution of Arctic Char (*Salvelinus*
alpinus) from the Hornaday and Brock Rivers to Subsistence Fisheries in
Darnley Bay, Northwest Territories + Supplementary Appendix Tables S1
to S4 (See Article Tools). *ARCTIC*, *69*(3), 231–245.
https://doi.org/10.14430/arctic4578
Harris, L. N., Moore, J.-S., Bajno, R., & Tallman, R. F. (2016). Genetic Stock
Structure of Anadromous Arctic Char in Canada's Central Arctic: Potential
Implications for the Management of Canada's Largest Arctic Char
Commercial Fishery. *North American Journal of Fisheries Management*,
36(6), 1473–1488. https://doi.org/10.1080/02755947.2016.1227399
Harris, L. N., Moore, J.-S., Galpern, P., Tallman, R. F., & Taylor, E. B. (2014).
Geographic influences on fine-scale, hierarchical population structure in
northern Canadian populations of anadromous Arctic Char (Salvelinus
alpinus). *Environmental Biology of Fishes*, *97*(11), 1233–1252.
https://doi.org/10.1007/s10641-013-0210-y
Harris, L. N., Yurkowski, D. J., Gilbert, M. J. H., Else, B., Duke, P. J., Ahmed, M.,
Tallman, R. F., Fisk, A. T., & Moore, J. S. (2020). Depth and temperature
preference of anadromous Arctic Char, Salvelinus alpinus, in the Kitikmeot
Sea: a shallow and low salinity area of the Canadian Arctic. *Marine*
Ecology Progress Series, *634*, 175–197.
https://doi.org/10.3354/meps13195
Jacobsen, M. W., Jensen, N. W., Nygaard, R., Præbel, K., Jónsson, B., Nielsen,
N. H., Pujolar, J. M., Fraser, D. J., Bernatchez, L., & Hansen, M. M.
(2022). A melting pot in the Arctic: Analysis of mitogenome variation in
Arctic char (*Salvelinus alpinus*) reveals a 1000 km contact zone
between highly divergent lineages. *Ecology of Freshwater Fish*, *31*(2),
330–346. https://doi.org/10.1111/eff.12633
Jamsandekar, M., Ferreira, M. S., Pettersson, M. E., Farrell, E. D., Davis, B. W.,
& Andersson, L. (2024). The origin and maintenance of supergenes
contributing to ecological adaptation in Atlantic herring. *Nature*
Communications, *15*(1), 9136. https://doi.org/10.1038/s41467-024-53079-
7
Janes, J. K., Miller, J. M., Dupuis, J. R., Malenfant, R. M., Gorrell, J. C.,
Cullingham, C. I., & Andrew, R. L. (2017). The K = 2 conundrum.
Molecular Ecology, 26(14), 3594–3602. https://doi.org/10.1111/mec.14187
Keefer, M. L., & Caudill, C. C. (2014). Homing and straying by anadromous
salmonids: A review of mechanisms and rates. *Reviews in Fish Biology*
and Fisheries, *24*(1), 333–368. https://doi.org/10.1007/s11160-013-9334-6
Kess, T., Dempson, J. B., Lehnert, S. J., Layton, K., Einfeldt, A., Bentzen, P.,
Salisbury, S. J., Messmer, A. M., Duffy, S., Ruzzante, D. E., Nugent, C.
M., Ferguson, M. M., Leong, J. S., Koop, B. F., O'Connell, M. F., &
Bradbury, I. R. (2021). Genomic basis of deep water adaptation in Arctic
Charr (*Salvelinus alpinus*) morphs. *Molecular Ecology*, mec.16033.
https://doi.org/10.1111/mec.16033

Kirch, M., Romundset, A., Gilbert, M. T. P., Jones, F. C., & Foote, A. D. (2021). Ancient and modern stickleback genomes reveal the demographic constraints on adaptation. *Current Biology*, *31*(9), 2027-2036.e8. https://doi.org/10.1016/j.cub.2021.02.027

Kirkpatrick, M., & Barton, N. (2006). Chromosome inversions, local adaptation and speciation. *Genetics*, *173*(1), 419–434. https://doi.org/10.1534/genetics.105.047985

Klemetsen, A., Amundsen, P.-A., Dempson, J. B., Jonsson, B., Jonsson, N., O'Connell, M. F., & Mortensen, E. (2003). Atlantic salmon Salmo salar L., brown trout Salmo trutta L. and Arctic charr Salvelinus alpinus (L.): a review of aspects of their life histories. *Ecology of Freshwater Fish*, *12*(1), 1–59. https://doi.org/10.1034/j.1600-0633.2003.00010.x

Klopfenstein, D. V., Zhang, L., Pedersen, B. S., Ramírez, F., Warwick Vesztrocy, A., Naldi, A., Mungall, C. J., Yunes, J. M., Botvinnik, O., Weigel, M., Dampier, W., Dessimoz, C., Flick, P., & Tang, H. (2018). GOATOOLS: A Python library for Gene Ontology analyses. *Scientific Reports*, *8*, 10872. https://doi.org/10.1038/s41598-018-28948-z

Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak: a program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources*, *15*(5), 1179–1191. https://doi.org/10.1111/1755-0998.12387

Laikre, L., Hoban, S., Bruford, M. W., Segelbacher, G., Allendorf, F. W., Gajardo, G., Rodríguez, A. G., Hedrick, P. W., Heuertz, M., Hohenlohe, P. A., Jaffé, R., Johannesson, K., Liggins, L., MacDonald, A. J., OrozcoterWengel, P., Reusch, T. B. H., Rodríguez-Correa, H., Russo, I.-R. M., Ryman, N., & Vernesi, C. (2020). Post-2020 goals overlook genetic diversity. *Science*, *367*(6482), 1083–1085. https://doi.org/10.1126/science.abb2748

Layton, K. K. S., Snelgrove, P. V. R., Dempson, J. B., Kess, T., Lehnert, S. J., Bentzen, P., Duffy, S. J., Messmer, A. M., Stanley, R. R. E., DiBacco, C., Salisbury, S. J., Ruzzante, D. E., Nugent, C. M., Ferguson, M. M., Leong, J. S., Koop, B. F., & Bradbury, I. R. (2021). Genomic evidence of past and future climate-linked loss in a migratory Arctic fish. *Nature Climate Change*, *11*(2), 158–165. https://doi.org/10.1038/s41558-020-00959-7

Lea, E. V., Olokhaktomiut Hunters and Trappers Committee, & Harwood. (2023). *Fish and Marine Mammals Harvested near Ulukhaktok, Northwest Territories, with a focus on Anadromous Arctic Char (Salvelinus alpinus)* (No. 2023/014; p. 23). Fisheries and Oceans Canada.

Legendre, P. (1993). Spatial Autocorrelation: Trouble or New Paradigm? *Ecology*, *74*(6), 1659–1673. https://doi.org/10.2307/1939924

Lehnert, S. J., Bentzen, P., Kess, T., Lien, S., Horne, J. B., Clément, M., & Bradbury, I. R. (2019). Chromosome polymorphisms track trans-Atlantic divergence and secondary contact in Atlantic salmon. *Molecular Ecology*, *28*(8), 2074–2087. https://doi.org/10.1111/mec.15065

Lehnert, S. J., Bradbury, I. R., Wringe, B. F., Van Wyngaarden, M., & Bentzen, P. (2023). Multifaceted framework for defining conservation units: An

example from Atlantic salmon (Salmo salar) in Canada. *Evolutionary Applications*, *16*(9), 1568–1585. https://doi.org/10.1111/eva.13587

Li, P., van Coeverden de Groot, P., Clemente-Carvalho, R. B. G., & Lougheed, S. C. (2021). ddRAD genotyping reveals hierarchical genetic population structure in anadromous Arctic char (Salvelinus alpinus) in the Lower Northwest Passage, Nunavut. *Canadian Journal of Fisheries and Aquatic Sciences*, *78*(4), 457–471. https://doi.org/10.1139/cjfas-2020-0069

Linderoth, T. (2018). *Identifying Population Histories, Adaptive Genes, and Genetic Duplication from Population-Scale Next Generation Sequencing*. University of California, Berkeley.

Lotterhos, K. E. (2019). The Effect of Neutral Recombination Variation on Genome Scans for Selection. *G3 Genes|Genomes|Genetics*, *9*(6), 1851–1867. https://doi.org/10.1534/g3.119.400088

Luu, K., Bazin, E., & Blum, M. G. B. (2017). pcadapt: an R package to perform genome scans for selection based on principal component analysis. *Molecular Ecology Resources*, *17*(1), 67–77. https://doi.org/10.1111/1755-0998.12592

Mace, G. M. (2004). The role of taxonomy in species conservation. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, *359*(1444), 711–719. https://doi.org/10.1098/rstb.2003.1454

Madsen, R. P. A., Jacobsen, M. W., O'Malley, K. G., Nygaard, R., Præbel, K., Jónsson, B., Pujolar, J. M., Fraser, D. J., Bernatchez, L., & Hansen, M. M. (2019). Genetic population structure and variation at phenology-related loci in anadromous Arctic char (Salvelinus alpinus). *Ecology of Freshwater Fish*, *00*, 1–14. https://doi.org/10.1111/eff.12504

Marcus, J., Ha, W., Barber, R. F., & Novembre, J. (2021). Fast and flexible estimation of effective migration surfaces. *eLife*, *10*, e61927. https://doi.org/10.7554/eLife.61927

Matschiner, M., Barth, J. M. I., Tørresen, O. K., Star, B., Baalsrud, H. T., Brieuc, M. S. O., Pampoulie, C., Bradbury, I., Jakobsen, K. S., & Jentoft, S. (2022). Supergene origin and maintenance in Atlantic cod. *Nature Ecology & Evolution*, *6*(4), 469–481. https://doi.org/10.1038/s41559-022-01661-x

McDowall, R. M. (2001). Anadromy and homing: two life□history traits with adaptive synergies in salmonid fishes? *Fish and Fisheries*, *2*(1), 78–85. https://doi.org/10.1046/j.1467-2979.2001.00036.x

Meisner, J., & Albrechtsen, A. (2018). Inferring Population Structure and Admixture Proportions in Low-Depth NGS Data. *Genetics*, *210*(2), 719–731. https://doi.org/10.1534/genetics.118.301336

Moore, J.-S., Bajno, R., Reist, J. D., & Taylor, E. B. (2015). Post-glacial recolonization of the North American Arctic by Arctic char (Salvelinus alpinus): Genetic evidence of multiple northern refugia and hybridization between glacial lineages. *Journal of Biogeography*, *42*(11), 2089–2100. https://doi.org/10.1111/jbi.12600

Moore, J.-S., Bourret, V., Dionne, L., Bradbury, I. A. N., Reilly, P. O., & Kent, M. (2014). Conservation genomics of anadromous Atlantic salmon across its North American range□: outlier loci identify the same patterns of

population structure as neutral loci. *Molecular Ecology*, *23*, 5680–5697. https://doi.org/10.1111/mec.12972

Moore, J.-S., Harris, L. N., Kessel, S. T., Bernatchez, L., Tallman, R. F., & Fisk, A. T. (2016). Preference for near-shore and estuarine habitats in anadromous Arctic char (Salvelinus alpinus) from the Canadian high Arctic (Victoria Island, NU) revealed by acoustic telemetry. *Canadian Journal of Fisheries and Aquatic Sciences*, *53*(9), 1689–1699. https://doi.org/10.1017/CBO9781107415324.004

Moore, J.-S., Harris, L. N., Luyer, J. L., Sutherland, B., Rougemont, Q., Tallman, R. F., Fisk, A. T., & Bernatchez, L. (2017). Genomics and telemetry suggest a role for migration harshness in determining overwintering habitat choice, but not gene flow, in anadromous Arctic Char. *Molecular Ecology*, 1–14. https://doi.org/10.1111/mec.14393

Moore, J.-S., Harris, L. N., Tallman, R. F., & B, T. E. (2013). The interplay between dispersal and gene flow in anadromous Arctic char (Salvelinus alpinus): implications for potential for local adaptation. *Canadian Journal of Fisheries and Aquatic Sciences*, *1338*(July), 1327–1338.

Moritz, C. (2002). Strategies to Protect Biological Diversity and the Evolutionary Processes That Sustain It. *Systematic Biology*, *51*(2), 238–254. https://doi.org/10.1080/10635150252899752

Nunez, J. C. B., Lenhart, B. A., Bangerter, A., Murray, C. S., Mazzeo, G. R., Yu, Y., Nystrom, T. L., Tern, C., Erickson, P. A., & Bergland, A. O. (2024). A cosmopolitan inversion facilitates seasonal adaptation in overwintering *Drosophila*. *GENETICS*, *226*(2), iyad207. https://doi.org/10.1093/genetics/iyad207

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., Caceres, M. D., Durand, S., … Weedon, J. (2022). *vegan: Community Ecology Package*. https://CRAN.R-project.org/package=vegan

Olsen, L., Thum, E., & Rohner, N. (2021). Lipid metabolism in adaptation to extreme nutritional challenges. *Developmental Cell*, *56*(10), 1417–1429. https://doi.org/10.1016/j.devcel.2021.02.024

Östergren, J., Palm, S., Gilbey, J., Spong, G., Dannewitz, J., Königsson, H., Persson, J., & Vasemägi, A. (2021). A century of genetic homogenization in Baltic salmon—evidence from archival DNA. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1949), rspb.2020.3147, 20203147. https://doi.org/10.1098/rspb.2020.3147

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, *35*(3), 526–528. https://doi.org/10.1093/bioinformatics/bty633

Pearse, D. E., Barson, N. J., Nome, T., Gao, G., Campbell, M. A., Abadía-Cardoso, A., Anderson, E. C., Rundio, D. E., Williams, T. H., Naish, K. A., Moen, T., Liu, S., Kent, M., Moser, M., Minkley, D. R., Rondeau, E. B., Brieuc, M. S. O., Sandve, S. R., Miller, M. R., … Lien, S. (2019). Sex-dependent dominance maintains migration supergene in rainbow trout.

1156   *Nature Ecology & Evolution*, *3*(12), 1731–1742.

1157   https://doi.org/10.1038/s41559-019-1044-6

1158 Peterson, D. A., Hilborn, R., & Hauser, L. (2014). Local adaptation limits lifetime

1159   reproductive success of dispersers in a wild salmon metapopulation.

1160   *Nature Communications*, *5*(1), 3696. https://doi.org/10.1038/ncomms4696

1161 Petkova, D., Novembre, J., & Stephens, M. (2016). Visualizing spatial population

1162   structure with estimated effective migration surfaces. *Nature Genetics*,

1163   *48*(1), 94–100. https://doi.org/10.1038/ng.3464

1164 Priest, H., & Usher, P. J. (2004). *Nunavut Wildlife Harvest Study* (p. 822).

1165   Nunavut Wildlife Management Board.

1166 Pritchard, J. K., Pickrell, J. K., & Coop, G. (2010). The Genetics of Human

1167   Adaptation: Hard Sweeps, Soft Sweeps, and Polygenic Adaptation.

1168   *Current Biology*, *20*(4), R208–R215.

1169   https://doi.org/10.1016/j.cub.2009.11.055

1170 Quinn, T. P. (1993). A review of homing and straying of wild and hatchery-

1171   produced salmon. *Fisheries Research*, *18*(1–2), 29–44.

1172   https://doi.org/10.1016/0165-7836(93)90038-9

1173 Reist, J. D., Power, M., & Dempson, J. B. (2013). Arctic charr (Salvelinus

1174   alpinus): a case study of the importance of understanding biodiversity and

1175   taxonomic issues in northern fishes. *Biodiversity*, *14*(1), 45–56.

1176   https://doi.org/10.1080/14888386.2012.725338

1177 Rellstab, C., Gugerli, F., Eckert, A. J., Hancock, A. M., & Holderegger, R. (2015).

1178   A practical guide to environmental association analysis in landscape

1179   genomics. *Molecular Ecology*, *24*(17), 4348–4370.

1180   https://doi.org/10.1111/mec.13322

1181 Rosenblum, E. B., Sarver, B. A. J., Brown, J. W., Des Roches, S., Hardwick, K.

1182   M., Hether, T. D., Eastman, J. M., Pennell, M. W., & Harmon, L. J. (2012).

1183   Goldilocks Meets Santa Rosalia: An Ephemeral Speciation Model

1184   Explains Patterns of Diversification Across Time Scales. *Evolutionary*

1185   *Biology*, *39*(2), 255–261. https://doi.org/10.1007/s11692-012-9171-x

1186 Roux, C., Fraïsse, C., Romiguier, J., Anciaux, Y., Galtier, N., & Bierne, N. (2016).

1187   Shedding Light on the Grey Zone of Speciation along a Continuum of

1188   Genomic Divergence. *PLOS Biology*, *14*(12), e2000234.

1189   https://doi.org/10.1371/journal.pbio.2000234

1190 Roux, M.-J., Tallman, R. F., & Martin, Z. A. (2019). Small-scale fisheries in

1191   Canada's Arctic: Combining science and fishers knowledge towards

1192   sustainable management. *Marine Policy*, *101*, 177–186.

1193   https://doi.org/10.1016/j.marpol.2018.01.016

1194 Ryder, O. (1986). Conservation and systematic: The dilemma of subspecies.

1195   *Trends Ecol Evol*, *1*, 9–10.

1196 Salisbury, S. J., McCracken, G. R., Keefe, D., Perry, R., & Ruzzante, D. E.

1197   (2019). Extensive secondary contact among three glacial lineages of

1198   Arctic Char (Salvelinus alpinus) in Labrador and Newfoundland. *Ecology*

1199   *and Evolution*, *9*(4), 2031–2045. https://doi.org/10.1002/ece3.4893

1200 Schindler, D. E., Hilborn, R., Chasco, B., Boatright, C. P., Quinn, T. P., Rogers,

1201   L. A., & Webster, M. S. (2010). Population diversity and the portfolio effect

in an exploited species. *Nature*, *465*(7298), 609–612.
https://doi.org/10.1038/nature09060
Seehausen, O., Takimoto, G., Roy, D., & Jokela, J. (2008). Speciation reversal
and biodiversity dynamics with hybridization in changing environments.
Molecular Ecology, *17*(1), 30–44. https://doi.org/10.1111/j.1365-
294X.2007.03529.x
Skotte, L., Korneliussen, T. S., & Albrechtsen, A. (2013). Estimating Individual
Admixture Proportions from Next Generation Sequencing Data. *Genetics*,
195(3), 693–702. https://doi.org/10.1534/genetics.113.154138
Spares, A. D., Stokesbury, M. J. W., Dadswell, M. J., O'Dor, R. K., & Dick, T. A.
(2015). Residency and movement patterns of Arctic charr Salvelinus
alpinus relative to major estuaries. *Journal of Fish Biology*, *86*(6), 1754–
1780. https://doi.org/10.1111/jfb.12683
Tallman, R. F., Roux, M.-J., & Martin, Z. A. (2019). Governance and assessment
of small-scale data-limited Arctic Charr fisheries using productivity-
susceptibility analysis coupled with life history invariant models. *Marine*
Policy, *101*, 187–197. https://doi.org/10.1016/j.marpol.2017.11.032
Therkildsen, N. O., & Palumbi, S. R. (2017). Practical low-coverage genomewide
sequencing of hundreds of individually barcoded samples for population
and evolutionary genomics in nonmodel species. *Molecular Ecology*
Resources, *17*(2), 194–208. https://doi.org/10.1111/1755-0998.12593
Tigano, A., Jacobs, A., Wilder, A. P., Nand, A., Zhan, Y., Dekker, J., &
Therkildsen, N. O. (2021). Chromosome-Level Assembly of the Atlantic
Silverside Genome Reveals Extreme Levels of Sequence Diversity and
Structural Genetic Variation. *Genome Biology and Evolution*, *13*(6),
evab098. https://doi.org/10.1093/gbe/evab098
UniProt Consortium. (2024). *Swiss-Prot: A manually annotated and reviewed*
protein sequence database. (Version 1.2) [Dataset].
Vieira, F. G., Lassalle, F., Korneliussen, T. S., & Fumagalli, M. (2016). Improving
the estimation of genetic distances from Next-Generation Sequencing
data. *Biological Journal of the Linnean Society*, *117*(1), 139–149.
https://doi.org/10.1111/bij.12511
Wang, C., Gong, Y., Deng, F., Ding, E., Tang, J., Codling, G., Challis, J. K.,
Green, D., Wang, J., Chen, Q., Xie, Y., Su, S., Yang, Z., Raine, J., Jones,
P. D., Tang, S., & Giesy, J. P. (2021). Remodeling of Arctic char (
Salvelinus alpinus) lipidome under a stimulated scenario of Arctic
warming. *Global Change Biology*, *27*(14), 3282–3298.
https://doi.org/10.1111/gcb.15638
Waples, R. S. (1991). Pacific salmon, Oncorhynchus spp., and the definition of
"species" under the Endangered Species Act. *Marine Fisheries Review*,
53(3), 11–22.
Waples, R. S. (1995). Evolutionarily significant units and the conservation of
biological diversity under the Endangered Species Act. *American*
Fisheries Society Symposium, *17*, 8–27.
Waples, R. S., Ford, M. J., Nichols, K., Kardos, M., Myers, J., Thompson, T. Q.,
Anderson, E. C., Koch, I. J., McKinney, G., Miller, M. R., Naish, K., Narum,

S. R., O'Malley, K. G., Pearse, D. E., Pess, G. R., Quinn, T. P., Seamons,
T. R., Spidle, A., Warheit, K. I., & Willis, S. C. (2022). Implications of
Large-Effect Loci for Conservation: A Review and Case Study with Pacific
Salmon. *Journal of Heredity*, *113*(2), 121–144.
https://doi.org/10.1093/jhered/esab069
Wellenreuther, M., Mérot, C., Berdan, E., & Bernatchez, L. (2019). Going beyond
SNPs: The role of structural genomic variants in adaptive evolution and
species diversification. *Molecular Ecology*, *28*(6), 1203–1209.
https://doi.org/10.1111/mec.15066
Wilcox, M. A., Jeffery, N. W., DiBacco, C., Bradbury, I. R., Lowen, B., Wang, Z.,
Beiko, R. G., & Stanley, R. R. E. (2023). Integrating seascape resistances
and gene flow to produce area-based metrics of functional connectivity for
marine conservation planning. *Landscape Ecology*, *38*(9), 2189–2205.
https://doi.org/10.1007/s10980-023-01690-2
Xu, S., Dai, Z., Guo, P., Fu, X., Liu, S., Zhou, L., Tang, W., Feng, T., Chen, M.,
Zhan, L., Wu, T., Hu, E., Jiang, Y., Bo, X., & Yu, G. (2021). ggtreeExtra:
Compact Visualization of Richly Annotated Phylogenetic Data. *Molecular*
Biology and Evolution, *38*(9), 4039–4042.
https://doi.org/10.1093/molbev/msab166
Xuereb, A., Rougemont, Q., Dallaire, X., Moore, J.-S., Normandeau, E., Bougas,
B., Perreault-Payette, A., Koop, B. F., Withler, R., Beacham, T., &
Bernatchez, L. (2022). Re-evaluating Coho salmon (Oncorhynchus
kisutch) conservation units in Canada using genomic data. *Evolutionary*
Applications, *15*(11), 1925–1944. https://doi.org/10.1111/eva.13489
Yu, G., Smith, D. K., Zhu, H., Guan, Y., & Lam, T. T.-Y. (2017). ggtree: an r
package for visualization and annotation of phylogenetic trees with their
covariates and other associated data. *Methods in Ecology and Evolution*,
8(1), 28–36. https://doi.org/10.1111/2041-210X.12628
1276