Long-Term Intestinal Epithelial Remodeling Induced by Acute Protein-Energy

Malnutrition

Fenja A. Schuran[1,*], Neha Mishra[1,*], Víctor A. López-Agudelo[1], Nina Sommer[1], Joana P.

Bernardes[1], Alesia Walker[2], Finn Hinrichsen[1,3], Ting Gong[1], Felix Gilbert[1,3], Lena Schröder[1],

Archana Bhardwaj[1], Sven Künzel[4], Saskia Weber-Stiehl[1], Go Ito[5], Florian Tran[1,3], Matthieu

Groussin[1,6], Christoph Röcken[7], Juan Matute[,8,9], Stefan Schreiber[3], Josef M. Penninger[10,11,12],

Richard S. Blumberg[8], Philippe Schmitt-Kopplin[2], John F. Baines[4,13], Felix Sommer[1,#] & Philip

Rosenstiel[1,3#]

[1] Institute of Clinical Molecular Biology, University of Kiel and University Hospital Schleswig-

Holstein, Rosalind-Franklin-Straße 12, 24105 Kiel, Germany

[2] Research Unit Analytical BioGeoChemistry, Helmholtz Zentrum München, German

Research Center for Environmental Health, Ingolstädter Landstrasse 1, 85764 Neuherberg,

Germany.

[3] Dep. of Internal Medicine I, University Hospital Schleswig-Holstein, Campus Kiel, Arnold-

Heller-Straße 3, 24105 Kiel, Germany

[4] Max-Planck-Institute for Evolutionary Biology, August-Thienemann-Straße 2, 24306 Plön,

Germany

[5] Department of Gastroenterology and Hepatology, Institute of Science Tokyo, 2 Chome-12-

1 Ookayama, 152-8550, Tokyo, Japan

[6] The Global Microbiome Conservancy, Massachusetts Institute of Technology, Cambridge,

Massachusetts, USA.

[7] Department of Pathology, Christian-Albrechts-University, University Hospital Schleswig-

Holstein, 24105, Kiel, Germany.

[8] Gastroenterology Division, Department of Medicine, Brigham and Women's Hospital,

Harvard Medical School, 77 Avenue Louis Pasteur, Boston MA 02115, USA

[9]current address: Division of Neonatology, Department of Pediatrics, Children's Hospital of

Philadelphia, Perelman School of Medicine at the University of Pennsylvania, 3401 Civic

Center Blvd, Philadelphia PA 19004, USA

[10]Department of Laboratory Medicine, Medical University of Vienna, Währinger Gürtel 18-20,

1090 Vienna, Austria

[11] Helmholtz Centre for Infection Research, Inhoffenstraße 7, 38124 Braunschweig, Germany

[12]Department of Medical Genetics, Life Sciences Institute, University of British Columbia,

Vancouver V6T 1Z3, Canada

[13]Institute of Experimental Medicine, University of Kiel, Michaelisstr. 5, 24105 Kiel, Germany

[*] Authors contributed equally with shared first authorship.

[#] Shared senior authorship.

Keywords: Protein energy malnutrition / intestinal inflammation/ Paneth cell / microbiome

/ GDF15 / intestinal epithelium

Correspondence:

Philip Rosenstiel (p.rosenstiel@mucosa.de) and Felix Sommer (f.sommer@ikmb.uni-kiel.de)

Abstract

Protein-energy malnutrition (PEM) is a global health burden with lasting effects that extend

well beyond the initial nutrient deficiency. To systematically investigate the long-term effects

of a single episode of PEM on the structure and function of the intestinal epithelium and its

associated microbiota, we employed a comprehensive multi-omics approach, including

(spatial) transcriptomics, DNA methylation analysis, fecal metagenomics, and metabolomics.

Our findings show that PEM persistently alters the intestinal epithelium by depleting Paneth

cells and suppressing antimicrobial gene expression - changes linked to DNA methylation that

persist despite dietary recovery. In germ-free mice, the sustained epithelial phenotype after

was absent. We identified the microbial lipid metabolite 9-HODE and epigenetically

deregulated PPAR-driven GDF15 expression as key molecular drivers of the persistent PEM-

induced Paneth cell dysfunction. Targeting microbial lipid production and its link to the host

GDF15 pathway could offer novel therapeutic strategies for long-term consequences of

malnutrition and other Paneth cell-associated diseases.

Introduction

Undernutrition is a major global health burden and is responsible for 45% of child deaths world-wide [1,2]. It is defined as a disordered nutritional state resulting from a combination of inflammation and a negative nutrient balance, leading to changes in body composition, function and outcome and associated disorders [3]. Clinically, undernutrition has several facets, ranging from a mild caloric and/or micronutrient deficit to more severe forms resulting from sustained inadequate uptake of protein (protein energy malnutrition, termed PEM hereafter). The latter is the main form of global undernutrition and combines an energy deficit with lack of essential amino acids, which are necessary building blocks for new proteins and regulate cellular energy homeostasis. PEM is by far the most important causative factor of morbidity and mortality in developing countries and crisis regions, but also in industrialized countries the growing socioeconomic gap and specifically increasing poverty at young and old age, are linked to a significant burden of this disease state [4]. The gut microbiota has been convincingly shown to be involved in the etiology of PEM [5–8] and microbiota-directed nutritional interventions have been utilized to overcome acute periods of malnutrition [9–12].

Whereas the pathophysiology of acute PEM has been intensely studied [7], long-term consequences of PEM and the molecular factors that contribute to such effects are not sufficiently understood. These consequences are characterized by a "memory effect" on specific body functions, which can be established after just one episode of malnutrition and may even be inherited by future generations[13]. Malnutrition during fetal development or early life can lead to ongoing poor weight gain and growth (known as stunting), which often persists even when a normal diet is introduced during childhood. Other clinical features associated with previous episodes of PEM include delayed motor [14] and intellectual development [15–17] and predisposition to obesity [18] and Type 2 diabetes in adulthood [19–21]. Importantly, individuals who suffered from chronic malnutrition in the past are also more vulnerable to intestinal infections and have a higher mortality rate due to infection-related complications [22,23]. Previous PEM may extend the duration of infectious diarrhea [24], potentially due to a decrease of intestinal barrier function, facilitating translocation of commensal bacteria and pathogens. We had previously shown that a lack of ACE2/BOAT1-dependent absorption of tryptophan in intestinal epithelial cells of mice may be a pivotal link between malnutrition, commensal bacteria and intestinal inflammation [25], but the duration

and the exact cellular mechanism of this effect remained unclear. Early studies have shown that PEM may sustainably hinder the growth of epithelial cells in the small intestinal crypts, causing delayed cellular migration along the crypt-villus axis [26]. This may result in a delay in the restoration of the epithelial structure and the recovery of the enzymatic and absorptive capacity in malnourished individuals [27]. Moreover, serum anti-antibodies to intestinal microbial antigens (e.g. flagellin, LPS) are persistently elevated after PEM, indicating an insufficient barrier function and sustained immune responses against the gut microbiota [28–30]. Despite this wealth of observations pointing towards fundamental long-term effects of PEM on the intestinal mucosa, the effector mechanisms, e.g. microbe-derived metabolites acting on cell fate decisions, remains largely unexplained. Here, we employed a murine model of adult PEM to systematically understand long-term effects of severe episode of protein undernutrition on the structure and functionality of the intestinal mucosa and its associated microbiota.

Results

Persistent consequences of experimental protein energy malnutrition in the small intestine of adult mice.

Protein-restricted isocaloric diets in mice have been suggested to resemble the acute phenotype of protein energy malnutrition [23,31,33]. To establish a dietary PEM model in mice, we first conducted a pilot study to assess the ability of isocaloric experimental diets with different protein levels (0.7%, 2%, 5% protein, hereafter termed "PEM diet") to cause a PEM phenotype, as determined by the maximum weight loss within a four-week treatment period, compared to a standard diet with 20% protein, which is termed "control or ctrl diet" hereafter (Fig S1A). To investigate the long-term physiological effects of an episode of protein-energy malnutrition (PEM) on the intestinal tract, we conducted a dietary study using 10-week-old adult male C57 BL/6J mice. The mice were first subjected to a 4-week isocaloric PEM diet containing 0.7% protein, followed by a 6-week recovery phase on a regular control diet with 20% protein. (Figure 1A). After 4 weeks of 0.7% protein diet, mice lost approx. 25% of body weight compared to their littermates fed with control diet. After switching the recovery group to the control diet following 4 weeks of PEM, the mice experienced rapid weight gain during the first 2 weeks. However, by the end of the 6-week follow-up period, they had not fully regained the weight levels of the mice that had been maintained on the control diet for the entire duration of the study. Notably, this weight difference emerged after the mice had completed their axial growth, suggesting a lasting, "memory-like" effect of the prior PEM period in adult mice (Figure 1B). In a separate experiment, we confirmed that the dietary regimen successfully induced stunting in 4-week-old mice. However, we opted to proceed with the adult model to avoid potential secondary effects of perturbed developmental processes (Figure S2). No significant difference in food consumption was detected between the study groups (Figure 1C). As expected, PEM induced transient effects in several organ systems. We noticed decreased liver, epididymal white adipose tissue (EWAT) and spleen weight after 4 weeks of PEM diet (Figure S1B), which were however completely restored to the levels of their littermate controls after the shift to the control diet. Strikingly, exposure to PEM also led to several persistent changes in the intestinal tract. We observed a significantly reduced length of the small intestine in acute PEM and after recovery. While only subtle changes of crypt-villous architecture (crypt depth and villous length) were detectable (Figure

S1C), persistently reduced Ki-67 and TUNEL-positive epithelial cells indicated a slower

proliferation and turnover of the small intestinal epithelial lining as sequelae of PEM as shown

previously in humans [26,27]. This effect was limited to the small intestine, the primary site for

protein digestion and absorption. In contrast, the colon showed reduced length (Figure 1D)

and impaired epithelial proliferation (data not shown) only during the acute phase of PEM,

both of which fully normalized after 6 weeks on the control diet.

Acute PEM induces a complex transcriptional dysregulation and leads to sustained

downregulation of anti-microbial defense modules in intestinal epithelial cells

We next aimed to delineate the dynamics of the transcriptional response of intestinal

epithelial cells during PEM and after recovery. We performed RNA sequencing of purified

intestinal epithelial cells from the small intestine at acute PEM (4 weeks) and after 6 weeks

recovery (10 weeks). Principal component (PC) analysis showed a strong separation between

the acute PEM samples and all other conditions on the first PC (Figure S3A) indicating a severe

alteration of global transcriptome signatures of the epithelium under acute PEM. A total of

8,197 genes were differentially expressed as a result of PEM, with 4,190 downregulated and

4,007 upregulated, either during the acute phase or after recovery, compared to controls.

4,125 of these differentially expressed genes (DEGs) were unique to the acute PEM phase

only (2,007 down- and 2,118 upregulated genes). The upregulated PEM-specific DEGs

included genes involved in key metabolic processes, such as acyl-CoA dehydrogenases (Acadl,

Acadm, Acadvl, Acads, Acad11), phosphokinases (Khk, Tkfc), and glutathione transferases

(Gsta1, Gsta2, Gsta3, Gsta4, Gstm1, Gstm2, Gstm3, Gstm4, Gstm6, Gstp1, Gstp2, etc.). Gene

set enrichment analyses identified an over-representation of metabolic processes, including

beta-oxidation, glyceraldehyde-3-phosphate metabolism, and glutathione homeostasis, likely

reflecting a significant metabolic shift and heightened oxidative stress in response to PEM.

(Figures 2A, S3D). Among the downregulated DEGs, changes comprised decreased innate

immune and TNF production, such as *Cd14*, *Cd2*, *Ccr2*, *Ccr5*, *Ccr7*, *Il17a*, *Stat3*, *Tlr2* among

others (Figures 2A, S3D).

Following recovery, 3,791 genes were uniquely differentially expressed in post-PEM animals

compared to age-matched controls, with 2,011 genes downregulated and 1,781 upregulated.

(Figure S3B). Several DEGs followed an inverse pattern at acute PEM (4w) and recovery (10w)

suggesting compensatory regulatory processes after recovery from a period of protein

restriction, e.g. downregulation of fatty acid beta-oxidation and glutathione metabolism-

related DEGs relative to age-matched controls in gene set enrichment analyses (Figure S3D).

A comparison of the two timepoints to identify potential transcriptional "PEM-memory"

effects revealed an overlap of 281 DEGs (172 downregulated and 109 upregulated; Figure 2

and S3B). These genes exhibited consistent transcriptional regulation in the same direction

during both the acute phase of PEM (4 weeks) and after recovery (10 weeks), compared to

age-matched controls. Upregulated shared DEGs were enriched for differentiation-related

transcripts including *Dvl1* and *Dvl3* (Dishevelled Segment Polarity Protein 1 and 3), *Notch1*

(Neurogenic locus notch homolog protein 1), *Neurod1* (Neuronal Differentiation 1), which

pointed towards sustained alteration of epithelial lineage decisions (Figure 2B). Persistently

downregulated DEGs comprised a large body of antimicrobial effector molecules such as

several alpha-defensins (*Defa3, 17, 21, 22, 24, 26*), epithelial lysozyme (*Lyz1*) and intelectin

(*Itln1*), indicating an ongoing reduction of epithelial defense mechanisms after PEM (Figure

2D). To further explore the topographical distribution of epithelial alterations, we performed

spatial transcriptomics on intestinal tissues isolated from mice under acute PEM (4 weeks)

and age-matched controls (Figure 3C). Spots containing greater than 60% epithelial tissue

within the capture zone were manually selected and classified according to their anatomical

position, namely "Crypt", "Villus-Intermediate" (including both bottom and middle sections)

and "Villus-Tip", using high-resolution images of H&E-stained intestinal Swiss roles (Figure 3C,

S3F). When comparing these regions between control and PEM mice, we found that the

upregulated metabolic processes in PEM were primarily localized to the villi (Figure 3D),

whereas downregulation of antimicrobial effector molecules (*Defa3, 5, 17, 23, 26*) was

confined to the crypt region (Figure 3E).

As many of the genes involved in the transcriptional memory effect of PEM, such as Defa3,

Defa17, Defa21, Defa22, Defa24, Defa26, as well as Ang4, Hspb8, Itln1, and Lyz1 (Figure 2D,

3A), are recognized as Paneth cell marker genes, we next enumerated small intestinal Paneth

cells by lysozyme immunohistochemistry. Indeed, numbers of lysozyme-positive Paneth cells

were significantly reduced in the acute phase of PEM and remained at a much lower level

after recovery (Figure 3B). Proportions of Goblet cells, another cell type of the secretory

lineage, were unaffected (Figure S3C).

To test the long-term stability of PEM-dependent gene signatures, we isolated intestinal
organoids from mice during the PEM-recovery diet experiment at acute PEM (4 weeks) and
after recovery (10 weeks) including from CTRL diet-fed mice. These organoids were cultured
for at least five passages in regular organoid medium (no nutritional restriction), before RNA
was isolated and subjected to RNA sequencing. Comparing the general expression patterns
between intestinal epithelial cells isolated directly from the mice and cultured organoids
revealed a largely similar expression pattern both under acute PEM (Figure 4A) and after
recovery (Figure 4B). The persistence of these changes in the *ex vivo* setting suggests that a
single episode of PEM induced fixed regulatory alterations, leading to the observed long-
lasting transcriptional changes.

Integrative Analysis of Genome-Wide DNA Methylation and Differential Gene Expression in
Small Intestinal Epithelial Cells During Acute PEM and Post-Recovery

DNA methylation is an epigenetic mechanism that plays a key role in the long-term regulation
of gene expression. In our previous work, we demonstrated its involvement in the
differentiation processes of intestinal epithelial cells [32,34]. To find a potential mechanism of
the observed long-lasting transcriptional memory effect after PEM, we next analyzed
genome-wide DNA methylation signatures by bead-based arrays using purified intestinal
epithelial cells from the small intestine at acute PEM (4 weeks) and after recovery (10 weeks)
and respective controls. In the acute PEM samples, we identified 33,013 differentially
methylated positions (DMPs), of which 18,214 were hyper- and 14,799 hypomethylated
(Figure 4C). Whereas after recovery, hypomethylation was the prevailing pattern compared
to age-matched controls (21,753 hypo- vs. 9550 hypermethylated sites). Similar to the
intersection of transcriptome signatures shared by acute PEM and recovery samples, we
found that a notable proportion of DMPs from acute PEM overlapped with DMPs detected in
the recovery group, with 1026 hyper- and 1007 hypomethylated sites being persistently
deregulated (Figure 4D). Differentially methylated sites significantly overlapped with binding
motifs for several transcription factors, for example the cell division proteins and PPAR targets
RAD21 and SMC3 [35,36], as well as the mRNA transporter ZC3H11A and transcriptional
repressor CTCF which were enriched in hypomethylated sites shared between acute PEM and
recovery samples (Figure 4E).

To delineate regulatory DNA methylation changes linked with persistently altered gene

expression *in cis*, we integrated transcriptome and methylome data using a hierarchical

approach as described [32,37] (Figure 4F). First, we located DMPs situated within a range of 5kb

before or after the transcription start site of any DEG. Next, we computed the correlation

between the expression level of each DEG and the intensity of methylation at the

corresponding DMPs. Among 56,271 deregulated DMPs analyzed, 6,554 were located within

the defined window of at least one DEG. Out of these, 6,197 DMP-DEG pairs representing

3,394 genes exhibited significant correlation, with 51% displaying a reciprocal correlation (i.e.,

high methylation linked to low transcript levels or vice versa) and 49% exhibiting a

cooperative correlation (i.e., high methylation linked to high transcript levels or vice versa).

31 DMP-DEG with a persistent pattern in acute PEM and after recovery were identified (Figure

4G). We were able to identify persistent links between DNA methylation and transcriptome

signatures, both after acute PEM and in the recovery phase. The cis-linked DNA methylation-

DEG pairs point towards potential long-term alteration of cellular states through epigenetic

regulation after a PEM episode. Notably, GDF15 (Growth/Differentiation Factor 15) emerged

as a promising candidate for mediating some of the long-term PEM memory effects, given its

established role as an endocrine signal of nutritional stress and body mass regulation and as

a mediator of cancer cachexia [38–40.]

Intestinal microbial community dynamics in acute PEM and after recovery indicate a high

level of resilience to PEM-induced perturbation

To explore the potential involvement of gut microbial dysbiosis for the observed sustained

phenotypes, we performed 16S rRNA gene sequencing on sequential stool samples collected

from mice during the PEM-recovery experiment (Figure 5A). During PEM (up to week 4),

bacterial α-diversity as measured by the Shannon Index declined significantly compared to

control-fed mice, but fully normalized again after switching to control diet during recovery

(from week 5 to 10) (Figure 5B). Composition of the fecal microbiota from mice of under acute

PEM differed profoundly from baseline (before dietary intervention) or control-fed

littermates (both at 4 weeks and 10 weeks "), and PEM-recovery mice at 10 weeks

(PERMANOVA on between-sample Bray-Curtis dissimilarities, F-statistics = 25.5, R^2 = 0.54, p =

1e-4, Figure 5C). Specifically, in mice subjected to PEM, *Enterobacteriaceae*, a family of the

class Proteobacteria that includes opportunistic pathogenic microorganisms such as *Escherichia* and *Shigella*, which have been linked with infection-related pathologies in malnourished individuals [41,42], were expanded along with *Bacteroides*, *Parabacteroides* or *Lachnospiraceae* UCG-006. In contrast, the relative abundance of *Bifidobacterium*, *Bilophila*, *Lachnospiraceae* NK4A136, *Lachnoclostridium*, *Alistipes*, *Parasutterella*, *Faecalibaculum* and *Muribaculaceae* were significantly decreased (Wilcoxon rank sum test, $p < 0.001$) compared to control-fed mice (Figures 5D and S4A, S4B). No significant differences were observed at the taxonomical level between PEM-recovery mice and their littermate controls. Notably, the intraclass correlation coefficient (ICC), which measures the similarity among samples within the same group, was significantly higher in fecal samples from mice with acute PEM compared to all other experimental groups (Figure S4D). This suggests that acute PEM induces uniform changes in the microbiota, with minimal influence from individual baseline condition or other environmental factors. Through shotgun metagenomic sequencing at weeks 4 and 10, we also explored changes of the functional potential of the microbiome. At 4 weeks, we found that a wide range of metabolic functions related to fatty acid and amino acid metabolism were enriched in the microbiome of mice under acute PEM compared to control-fed mice. (Figure 5E and S4C). Again, after recovery at week 10, no significant changes compared to control mice were observed, suggesting that adult microbial communities exhibit strong resilience to this type of perturbation.

Lack of an intestinal microbiome ameliorates phenotype severity and alters fecal metabolite patterns upon PEM perturbation

We next explored the role of the microbiome in shaping PEM-related phenotypes in our mouse model. We hypothesized that, similar to other stressors like DSS or radiation-induced enteritis [43], the absence of intestinal microbes—which induces an energy-restricted state, particularly in the intestinal mucosa—would exacerbate PEM symptoms. We thus performed the same PEM-recovery diet intervention in germfree (GF) mice of the same genetic background. Contrary to our expectation, GF mice exhibited significantly less body weight loss than conventionally raised (CONV-R) mice with a normal microbiome (Figure 6A). PEM-induced body weight loss could be fully normalized during recovery after switching to a control diet in GF, but not in CONV-R mice (Figure 6A). Importantly, in contrast to CONV-R

mice, numbers of Paneth cells were not affected by PEM in the absence of a microbiota (Figure 6B). Organ measures and abundance of other cell types were also unaltered by PEM in GF mice (Figure S5). Untargeted metabolomics of fecal samples from GF and CONV-R mice during acute PEM (day 28, week 4) revealed that lipid metabolites were significantly impacted by the nutritional challenge in a microbiota-dependent manner (Figure 6C). Specifically, conjugated linoleic acids and oxylipins such as 12,13-diHOME and 9-HODE were strongly induced under acute PEM in CONV-R mice, but were virtually absent in GF mice as individually validated by targeted quantitative metabolomics (Figure 6C, D). Supporting this metabolomic data, abundances of the respective functional genes in the microbiome, namely of lipoxygenases (LOX) and epoxide hydroxylases (EH), which generate 9-HODE and 12,13-diHOME from linoleic acid, respectively, were increased under acute PEM only in CONV-R mice as revealed by shotgun metagenomics (Figure 6E). Importantly, using publicly available metagenomic data of malnourished children undergoing a nutritional intervention with either a Microbiota Directed Complementary Food (MDCF-2) or standard Ready-to-Use Supplementary Food (RUSF) [10], we found that abundances of microbial LOX and EH genes declined during the therapeutic nutritional intervention (Figure 6F). These findings collectively demonstrate that fecal oxylipins, such as 9-HODE and 12,13-diHOME, along with the microbial enzymes that produce them, are highly sensitive to protein-energy malnutrition (PEM).

Activation of the PPAR-GDF15 pathway disrupts Paneth cell development during PEM

Oxylipins like 12,13-diHOME and 9-HODE are recognized as peroxisome proliferator-activated receptor (PPAR) agonists, and Paneth cell differentiation is regulated by PPAR signaling. We hypothesized that these microbially-derived metabolites may serve as a mechanistic link between the gut microbiota and disruptions in Paneth cell function [44–47]. PPARs are nuclear lipid-activable receptors controlling a variety of genes regulating metabolism, including GDF15 [48,49]. Strikingly, we observed that the expression patterns of both *Gdf15* and *Pparg* were co-dependent on PEM and the microbiome, showing upregulation in response to PEM and during recovery in CONVR mice, but not in GF mice (Figure 7A). We next assessed the functional role of PPAR activation and GDF15 for Paneth cell differentiation. Paneth cell differentiation was induced by a switch to standard [50] ENR-CD medium containing CHIR99021

(a GSK3 inhibitor/WNT activator) and DAPT (a γ-secretase inhibitor) (Figure 7B). Successful

induction was reflected by increased expression of Paneth cell markers over time, such as

Lyz1 (Figures 7C, D, E) and Defa5 (Figure 7C). We first tested the effects of GDF15 on Paneth

cells *in vitro*. Strikingly, stimulation of intestinal organoids with recombinant and TGFβ-free

GDF15 (rGDF15) [51] resulted in downregulation of Paneth cell markers (Figure 7C). To further

explore the link between PPAR signaling and GDF15, we established an *in vitro* organoid PEM

model. Intestinal organoids were cultured in either normal or protein-reduced ENR medium

(see methods), while simultaneously inducing Paneth cell differentiation by CHIR99021 and

DAPT addition. Of note, when organoid cultures were treated with Lanifibranor, a pan-PPAR

agonist, under PEM conditions, Paneth cell marker expression was significantly repressed

(Figure 7D), while Lanifibranor in the absence of protein restriction had no discernible effect.

Additionally, Lanifibranor increased *Gdf15* expression, confirming it as a direct downstream

target of PPAR under PEM (Figure 7D). Similar effects were observed with the microbial PPAR

agonist 9-HODE *in vitro* under PEM conditions. After 96 hours, 9-HODE decreased *Lyz1*

expression and upregulated *Gdf15* after 48 hours (Figure 7E). The timing of events, with an

initial upregulation of *Gdf15* expression followed 48 hours later by a decrease in *Lyz1* marker

expression in response to 9-HODE stimulation, collectively still supports a mechanistic link.

The interaction of PEM and PPAR-GDF15 activation was required to induce defective Paneth

cell differentiation as stimulation with Lanifibranor or 9-HODE in the absence of PEM did not

reduce *Lyz1* or *Gdf15* expression (Figure S6). These findings therefore suggest the presence

of a host-intrinsic PPAR-GDF15 pathway in the intestinal epithelium which can be activated

by microbially-derived oxylipin metabolites during PEM and mediates sustained effects on

Paneth cell differentiation and function.

Discussion

PEM remains a major global health challenge, impacting millions of individuals worldwide. In recent decades most studies have focused on understanding acute protein undernutrition. Major achievements are based on using locally available ingredients and co-targeting the intestinal microbiome [5,8,10,11,25,52–57]. However, the exact nature of the interactions between the intestinal microbiota and host tissues during nutritional stress are still largely unknown. In this study, we utilized a murine model of adult PEM to systematically investigate the long-term effects of a single severe episode of protein deficiency on both the structure and function of the intestinal mucosa, as well as its associated microbiota. Using multi-omics profiling, we uncovered extensive transcriptional, epigenetic, metabolic, and cellular effects of acute malnutrition. Our findings provide a resource of mucosa-microbiota alterations in the acute phase of PEM, which significantly extends previous insights and can be leveraged to mine for novel targeted interventions [5,8,9,11,58–61]. Importantly, we show that a single episode of PEM leads to a prolonged reduction in body weight and induces significant structural changes in the intestinal epithelium. These include macroscopic alterations, such as a shortened small intestine, and microscopic changes, such as a depletion of Paneth cells, which is a common feature of environmental enteric dysfunction (EED) in malnourished children [62,63]. Paneth cells are key regulators of intestinal physiology and antimicrobial responses, thus, our findings of permanently impaired Paneth cell function due to PEM could potentially explain the increased susceptibility towards gastrointestinal pathogens, i.e. *Enterobacteriaceae*, accounting for severe diarrhea, transmigration peritonitis and sepsis observed in severely malnourished humans [22,23]. The structural changes were accompanied by "memory-like" transcriptional alterations in the intestinal epithelium, notably a prolonged reduction in the expression of key Paneth cell genes, including those encoding antibacterial factors such as defensins and lysozyme, during and after acute PEM. These gene expression changes were sustainably present even in intestinal organoids grown from intestinal stem cells of mice harvested during acute PEM or recovery, when cultured for multiple passages *in vitro* under full-nutrient conditions. This suggests the involvement of long-term regulatory mechanisms, such as alterations in DNA methylation [64,65], which control the lasting effects of PEM. Indeed, we could show a strong remodeling of DNA methylation patterns upon PEM in purified intestinal epithelial cells. We conducted an interaction analysis between gene

expression and differentially methylated positions [32,37] and identified, besides other

differentiation factors (e.g. Neurod1 [66]), a link between cis-linked methylation changes and

upregulation of mRNA levels of the pleiotropic stress-induced cytokine GDF15 as a potential

mediator of PEM on intestinal function and metabolic processes [67–69].

To assess the dynamics of gut microbial community shifts during and after PEM we conducted

16S rDNA and metagenomic shotgun sequencing. In line with previous studies [5,7–9,11,58,61] we

detected a significant shift of gut microbiota composition and functional potential during

acute PEM. Strikingly, PEM-dependent changes completely reverted back to their original

configuration during recovery indicating a high level of resilience of fecal microbial

communities[70]. We initially hypothesized that lack of microbiome would increase PEM

susceptibility, e.g. due to an energetic restriction in the absence of bacterial fermentation [71,72]

or lack of TLR-dependent protective signals[43] . Surprisingly, when subjecting GF mice to PEM,

we found a significantly milder gastrointestinal phenotype with a less pronounced

bodyweight loss, and neither a long-lasting reduction of bodyweight nor significant depletion

of Paneth cells. These findings indicate that a second microbially-dependent hit is required to

trigger a strong PEM phenotype. Using metabolomics in CONV-R and GF mice, we identified

the fecal oxylipin metabolites 12,13-diHOME and 9-HODE as potential sources of the PEM-

and microbiome-dependent phenotype. By analyzing metagenomic data, we found increased

abundances of LOC and EH genes encoding enzymes that generate 9-HODE and 12,13-

diHOME from linoleic acid under acute PEM in CONV-R mice and in malnourished children,

which suggests that the same mechanism functions in mice and humans. Both, 9-HODE and

12,13-diHOME, have been implicated as immunomodulatory lipids, which may act through

PPARs [73–75]. Using an *ex vivo* malnutrition model in intestinal organoids, we then found that

protein deprivation alone did not impair Paneth cell differentiation, but a second (microbial)

hit targeting the PPAR/GDF15-axis was required.

Our data indicate that the sustained loss of Paneth cells could be linked to an epigenetically

fixed upregulation of GDF15 in intestinal epithelial cells. Whereas the intestinal epithelium

has been shown to be the major site of inducible GDF15 [76] upon nutritional stressors, there

are conflicting results on the expression distribution of its cognate receptor glial cell line-

derived neurotrophic factor family receptor α-like (GFRAL) as well as on biological effects of

yet unidentified non-GFRAL GDF15 receptors. Although the specific receptor system involved

was not identified in our study, we used a recombinant GDF15 source, which ruled out the

possibility of co-purified TGF being responsible. This suggests that the observed effects are

directly attributable to GDF15 itself. Our findings are of particular clinical relevance as biologic

treatments targeting GDF15 have not only proven effective in mice [77], but they are currently

entering phase 2 clinical trials for tumor cachexia in patients [40,78]. While the main effect of

this new drug class is to increase bodyweight via enhanced appetite, muscle function and

physical performance, these drugs could also prove effective in restoring Paneth cell numbers

and function in malnourished patients, thereby alleviating gastrointestinal symptoms and

reducing PEM lethality. A major obstacle in a global use of biologicals for PEM will be the

expected high cost of therapy. Instead, targeting the microbial metabolites 12,13-diHOME or

9-HODE could provide a cost-effective alternative significantly extending the scope of

successful nutritional interventions targeting the intestinal microbiome [10].

Potential limitations of the study

This study was conducted using a single episode of PEM in an adult murine model. Even

though there is evidence that these findings can be translated to the human situation (e.g.

Paneth cell disruption, elevated GDF15 levels, elevated abundance of LOX/EH genes), our

comprehensive molecular profiling results pointing towards a long-term memory-like effect

of PEM remain to be validated in a human setting. Moreover, studies on the relevance of our

findings for the childhood situation, e.g. for developmental phenotypes such as stunting [2], are

warranted. In addition, like most controlled PEM studies we used an isocaloric set-up, which

does not reflect the usual undernutrition scenario in crisis regions, where both available

protein and overall calories are severely reduced. It must be emphasized that the alteration

of lipid metabolism, which we see both in the mucosal transcriptome and bacterial

metagenome, is not linked to increased lipid content of the synthetic diet, as protein calories

are replaced by carbohydrates. Since lipid metabolism has been shown to be altered in human

malnutrition cohorts through both microbiome and serum analyses [79–82], we are confident

that the observed changes are primarily driven by metabolic shifts resulting from limited

protein availability. Importantly, due to our experimental setup, we do neither address the

effects of multiple PEM periods nor the duration of the observed memory phenotype in the

intestinal epithelium. From earlier regeneration kinetics, it seems however unlikely that the

effect is simply caused by the physiological lag time of differentiation, as acute chemical

ablation of Paneth cells was followed by re-appearance of mature cells already after 24h [83].

An interesting explanation for the disappearance of Paneth cells could be a permanent trans-

differentiation to stem-like dividing cells, which could partially replace the original stem cells

[84,85]. Further follow up studies including longer observation periods and/or multiple PEM

episodes and single cell-based analyses are required to comprehensively study the effects on

epithelial lineage decisions, but also long-term effects in other cell types and tissue

compartments.

It is increasingly evident that phenotypic responses to PEM vary between the sexes [86]. In our

study, most analyses have been done in males. In the GF condition, we included both sexes

and observed no significant differences between them. Further research should therefore

investigate sex-dependent effects during PEM, especially since woman during pregnancy and

breast-feeding are a very vulnerable group of patients prone to suffering from PEM due to an

increased protein demand [87]. Moreover, in this study we utilized a PEM model characterized

by a general protein deficiency, rather than a shortage of specific amino acids or their

combinations. Certain amino acids, such as tryptophan, undergo extensive metabolism by

both the host and the microbiota, generating metabolites with diverse signaling roles in

immune regulation and differentiation processes. These metabolites are known to influence

the expression of antimicrobial peptides [25,88]. It will be crucial to investigate whether the

acute and memory-like effects of PEM can be reversed through supplementation of these

amino acids or their metabolites, such as those involved in the kynurenine/nicotinamide

pathway.

Conclusions

In this study we present a comprehensive multi-omics resource of a dietary PEM model in

mice, revealing that a single episode of PEM induces long-lasting phenotypic changes, with a

particular focus on the intestinal mucosa and microbiota. Key findings include sustained

weight loss, shortening of the small intestine, impaired proliferation of intestinal epithelial

cells, a reduction in Paneth cells, and diminished expression of genes encoding antimicrobial

agents. Notably, germ-free (GF) mice exhibited alleviated acute PEM symptoms and did not

manifest any of the aforementioned long-term PEM-associated phenotypic changes. Through

an integrative multi-omics analysis and *ex vivo* intestinal organoid stimulations, we identified

microbial lipid metabolites, such as 12,13-diHOME and 9-HODE, along with the host PPAR-

GDF15 axis, as potential molecular drivers of the enduring PEM phenotype, particularly the

dysfunction of Paneth cells. These findings suggest that targeting microbial lipid production

or the host GDF15 pathway could provide novel therapeutic strategies, not only for

malnutrition, but also for other conditions associated with Paneth cell dysfunction, such as

Crohn's disease [89,90].

Methods

Animals. All animal experiments were approved by the local animal safety review board of the federal ministry of Schleswig Holstein and conducted according to national and international laws and policies (V242-30405/2016(55-5/16), V242-6900/2019(28-3/19) and V242-60285/2019(105-9/19)). C57Bl6/J mice were purchased from Charles River Laboratories and housed in the Central Animal Facility (ZTH) of the University Hospital Schleswig Holstein (UKSH, Kiel, Germany) under specific-pathogen free (SPF) conditions in individually ventilated cages (Green Line, Techniplast) or under sterile conditions in gnotobiotic flexible film isolators at the Max Planck Institute for Evolutionary Biology in Plön. All mice were kept under a 12-h light cycle and fed a regular gamma-irradiated chow diet *ad libitum*.

Dietary PEM intervention experiment. All custom PEM and CTRL diets were purchased from Ssniff (Table S1). To induce PEM in adult mice, 10-week-old CONV-R or 10-16-week-old GF C57Bl6/J mice (n=40) were either given a control diet (CTRL, 20% protein) or an isocaloric diet with reduced protein content (PEM, 0.7% protein) for 4 weeks to induce an acute PEM. To induce PEM in young mice, 4-week-old C57Bl6/J mice (n=40) were either given a control diet (CTRL, 20% protein) or an isocaloric diet with reduced protein content (PEM, 2% protein due to a greater PEM susceptibility of young mice) for 4 weeks to induce an acute PEM. One half of each experimental group was killed for blood & organ harvest at 4 weeks to assess the acute PEM phenotype. The other half of the experimental groups was either kept on or switched to a diet with regular protein content for an additional six weeks to assess recovery from PEM. Up to five recipient mice were co-housed. Mice were weighed every week and fecal pellets were collected, immediately frozen on dry ice and stored at –80°C. For final analyses, blood was collected from the vena cava, mice were killed by cervical dislocation and tissues were removed for histological and molecular analyses.

Histology and immunostainings. Tissue specimen were fixed in methacarn (60% dry methanol, 30% dry chloroform, 10% glacial acetic acid) for 1-2 weeks at room temperature prior to paraffin embedding and sectioning. 5 µm thick sections were cut and deparaffinized in Roti-Histol and rehydrated in ethanol followed by hematoxylin and eosin (H&E) or periodic acid-Schiff (PAS, TH Geyer, 3952016) staining. Alternatively, sections were boiled in citrate buffer for antigen retrieval and subjected to immunostaining. For this purpose, we used the Vectastain ABC kit (Vector Laboratories), peroxidase-substrate solution from DAB Peroxidase

(HRP) Substrate Kit (Vector Laboratories) and primary antibodies for detecting proliferating
cells were incubated in a 1:500 diluted mouse anti-Ki67 antibody (BD Biosciences, cat.no.
556003) and for Paneth cells in a 1:500 diluted mouse anti-LYZ1 antibody (Santa Cruz, cat.no.
sc-27956) overnight at 4°C. TUNEL assay was performed with the *in situ* cell death detection
Kit, TMR red (Roche, 12156792910) according to the manufacturer's instructions, while nuclei
were counterstained with DAPI (4',6-diamidino-2-phenylindole, dihydrochloride) nuclear dye
(Thermo Fisher, 62247). Slides were visualized using a Zeiss Imager Z1 microscope (Zeiss) and
pictures were taken using ZEN pro (Zeiss) software.

Isolation of intestinal epithelial cells (IECs). IECs were isolated from small intestinal tissue
using the Lamina Propria Dissociation Kit (Miltenyi BioTech, Bergisch Gladbach, Germany)
according to the manufacturer's protocol. In brief, intestinal epithelial cells were isolated by
disruption of the structural integrity of the epithelium using ethylenediaminetetraacetic acid
(EDTA) and dithiothreitol (DTT). Purity of individual IEC fractions was analyzed by flow
cytometry on a FACS Calibur flow cytometer (B&D, Heidelberg, Germany) with Cellquest
analysis software from Becton Dickinson. We used the Anti-EpCam-PE (Clone: G8.8,
Biolegend, San Diego, USA) antibody for analysis of IEC purity.

Noggin-conditioned medium. HEK Noggin cells, previously described by Miyoshi and
Stappenbeck (2013) and provided by Prof. Zeissig's group at CRTD, Dresden University,
Germany, were used in this study. In summary, HEK cells were modified to secrete Noggin
protein into the culture medium. The cells were thawed in DMEM + 10%(v/v) FBS purchased
from Biochrom. Once confluent, the cells were cultured for 3 cycles in DMEM with 10% (v/v)
FBS and 10 µg/mL Puromycin (Sigma-Aldrich). After the 3rd cycle, the cells were diluted 1:20
and cultured in DMEM with 10% (v/v) FBS without Puromycin. Four days later, the medium
was collected and centrifuged to remove excess cells. Fresh DMEM with 10% (v/v) FBS was
added and after another four days, the second batch of medium was collected and
centrifuged. The medium from the first batch was then combined with the second batch.

R-Spondin-conditioned medium. HEK R-Spondin-1-producing cells, previously described by
Miyoshi and Stappenbeck (2013) and provided by Prof. Zeissig´s group at CRTD, Dresden
University, Germany, were used in this study. In summary, HEK cells were modified to secrete
R-Spondin-1 protein into the culture medium. The cells were thawed in DMEM with 10% (v/v)
FBS (Biochrom). Once confluent, the cells were cultured for 3 cycles using DMEM with 10%

(v/v) FBS+ 300 µg/mL Zeocin (Thermo Fisher). After the 3rd cycle cells were diluted 1:20 and

cultured in DMEM with 10% (v/v) FBS without Zeocin. Four days later, the medium was

collected and centrifuged to remove excess cells. Fresh DMEM with 10% (v/v) FBS was added

and after another four days, the second batch of medium was collected and centrifuged. The

medium from the first batch was then combined with the second batch.

Isolation and culture of intestinal organoids. Organoids were generated from small intestinal

crypts of male C57Bl6/J mice receiving either a control or PEM diet following procedures

described earlier by Sato et al., 2009 [91]. Organoids were cultivated in 24 well plates at 37°C

with 5% CO_2 atmosphere in Matrigel (BD) with ENR-conditioned medium supplemented with

0.1% human recombinant EGF (50 µg/mL) as described by Sato et al., 2011 [50]. ENR-

conditioned medium consisted of 70% (*v/v*) 2 × basal medium (Advanced DMEM/F12

supplemented with HEPES [1M], Glutamax [100×], Penicillin/streptomycin 10,000 U/mL

[1:50] and N-Acetylcysteine [500 mM]), 10% (*v/v*) Noggin-conditioned medium and 20% R-

Spondin-conditioned medium.

Organoid PEM model. Murine intestinal organoids were initially cultured for 48 h in ENR-CV

medium, an ENR-conditioned medium containing 3.3 µM CHIR99021 (Miltenyi Biotech, #130-

103-926) and 1 mM valproic acid (Sigma, PHR1061-1G) promoting differentiation into the

stem cell niche. These stem cell organoids were then transitioned to either ENR-CD or PEM-

CD medium containing 3.3 µM CHIR99021 and 10 µM DAPT (Tebubio, 25704-0041) to further

drive differentiation into the Paneth cell niche.

The PEM culture medium (1.25% amino acids) consisted of 68.75% (*v/v*) Advanced DMEM/F12

(ADF) medium without amino acids or glucose [Biozol/US Biological Life Sciences, #USB-

D9800-27-10] supplemented with 10 mM HEPES, 1× Glutamax, 100 U/mL

Penicillin/streptomycin and 1 mM N-Acetylcysteine), 1.25% (*v/v*) regular ADF medium with

full amino acid concentrations, 10% (*v/v*) dialysed Noggin-conditioned medium and 20%

dialysed R-Spondin-conditioned medium. The Noggin- and 20% R-Spondin-conditioned media

were dialyzed using using cellulose-hydrate tubing with a cutoff of 15 kDa (Reichelt

Chemietechnik GmbH, catalogue number 84605)) to prevent the introduction of nutrients

from the added conditioned media while retaining Noggin and R-Spondin required for

organoid growth. for the in-vitro PEM model the conditioned Medium was dialyzed. Dialysis

success was assessed using Multistix 10 SG testing strips (Siemens).

Organoid stimulations with GDF15, Lanifibranor or 9-HODE. Murine intestinal organoids
were initially cultured for 48 h in ENR-CV medium, an ENR-conditioned medium containing
3.3 µM CHIR99021 (Miltenyi Biotech, #130-103-926) and 1 mM valproic acid (Sigma,
PHR1061-1G) promoting differentiation into the stem cell niche. These stem cell organoids
were then transitioned to either ENR-CD or PEM-CD medium containing 3.3 µM CHIR99021
and 10 µM DAPT (Tebubio, 25704-0041) to further drive differentiation into the Paneth cell
niche. Organoids were divided into experimental groups and stimulated with 1 µg *E. coli*
derived recombinant mouse GDF15 (R&D Systems, 8944-GD) per 500 µl organoid medium for
up to 120h, 20 µM of the pan-PPAR agonist Lanifibranor (MedChemExpress, HY-104049) for
24h to 96h or 1 µM 9(S)-HODE (Sigma Aldrich, SML0503) for 24h to 96h. Corresponding
control groups received vehicle treatments identical to those used for the stimuli. All
organoids were then cultured as described above.

RNA isolation and qPCR. RNA was extracted using the RNeasy Mini Kit (QIAGEN) according to
the manufacturer's protocol. Total RNA concentration was determined with a NanoDrop ND-
1000 spectrophotometer (PeqLab Biotechnologie). Total RNA was reverse transcribed to
cDNA using the High-Capacity cDNA Reverse Transcription Kit (ThermoFisher Scientific,
4368814). qPCR was carried out using TaqMan® Gene Expression Assays (ThermoFisher
Scientific) (Table S2) according to the manufacturer's instructions on a Viia 7 Real-Time PCR
System (ThermoFisher Scientific). Relative gene expression was normalized to housekeeper
gene *Actb* (β-actin) and calculated by the $2^{-\Delta Ct}$ method.

Transcriptional profiling by RNA sequencing. Total RNA from IECs and intestinal organoids
was extracted using the RNeasy Mini Kit (Qiagen) according to the manufacturer's protocol.
RNA concentration and integrity were analysed using a TapeStation 4200 System (Agilent)
and a Qubit 4 fluorometer (ThermoFisher Scientific). RNA-sequencing libraries were prepared
using the TruSeq® RNA seq Library Prep Kit v2 according to the Illumina TruSeq® messenger
(mRNA) sequencing protocol. The RNA-seq libraries were sequenced on an Illumina HiSeq
4000 sequencer (Illumina, San Diego,CA) with an average of 23 million reads (1x 50 bp) at
IKMB NGS core facilities. The nf-core rnaseq pipeline (version 1.2) (https://github.com/nf-
core/rnaseq) [92] was used to pre-process the sequencing data. Adapter and low-quality bases
from the sequencing reads were trimmed using Trim Galore (0.5.0)
(http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) and reads shorter than

35bp after trimming were removed. The filtered genes were mapped to the mouse genome

(GRCm38) using STAR aligner (version 2.6.1a) [93]. The expression counts of the genes were

estimated using featureCounts (version 1.6.2) [94] and were normalized across samples using

the DESeq normalization method. Differentially expressed genes (DEGs) between Ctrl_4w and

PEM_4w samples as well as Ctrl_10w and Recovery 10w samples were identified using the

Bioconductor package DESeq2 (version 1.20.0) [95]. Genes with FDR adjusted p-value of 0.05

were regarded as differentially expressed (DEGs). Gene ontology enrichment analyses were

conducted using the Bioconductor package topGO (version 2.46.0) [96] with genes with similar

expression level as the universe set. In the topGO analysis, the Fisher.elim p-value, calculated

using the weight algorithm, of 0.05 was used as the significance threshold.

Transcription factor binding sites (TFBS) enriched in the promoter regions of DEGs were

identified by conducting enrichment analysis using the Bioconductor package LOLA (version

1.24.0) [97]. Promoter regions were defined as the region between 1,500 bp upstream to 500

bp downstream of the transcription start site.

Spatial transcriptomics. Small intestinal tissue was prepared for spatial transcriptomics using

the Visium Spatial Fresh Frozen Protocol CG000240 (RevE, 10x Genomics). Briefly, small

intestinal tissue was rolled into Swiss rolls, embedded in O.C.T. medium (Tissue-Tek) within

Cryomolds (Tissue-Tek), and then rapidly immersed in pre-cooled isopentane on liquid

nitrogen until the O.C.T. medium solidified. The samples were subsequently cryopreserved at

-80°C. To determine the optimal permeabilization time of 15 minutes, serial 10μm sections

were cut and mounted on a Visium Tissue Optimization Slide (10x Genomics). For spatial

transcriptomics, additional 10μm-thick sections were mounted on a Visium Gene Expression

Slide for Fresh Frozen (10x Genomics), followed by methanol fixation and H&E staining

following protocol CG000160 (RevD, 10x Genomics). Whole slide imaging was conducted

using a NanoZoomer S60 Digital Slide Scanner (Hamamatsu, C13210-04) with 40x

magnification. Spatial library construction was carried out according to the manufacturer's

protocol CG000239 (RevH, 10x Genomics), and the libraries were then sequenced on a

NovaSeq 6000 sequencer (Illumina) with 2x 100bp read length. Sequencing data was mapped

to the mouse reference genome (mm10) using 10x spaceranger v.2.1. Feature-barcode

matrices for each sample were imported into the R package 'Seurat' v.4.2.0 for quality control,

normalization, dimensionality reduction, and Louvain clustering [98]. As quality control, spots

were filtered if they had <200 features (genes) or >25% of mitochondrial gene expression.

After spot filtration, sample level normalization was performed using SCTransform [99] function

in Seurat, which uses a regularized negative binomial regression to transform the UMI count

data. Thereafter, sample integration was performed with the CCA algorithm followed by a

Shared Nearest Neighbor (SNN) construct graph and clustering using default settings. Using

high-resolution images of H&E-stained intestinal Swiss roles, multiple regions were identified,

and spots classified as belonging to "Crypt", "Muscle", "Villus-Intermediate" (including both

bottom and middle sections), or "Villus-Tip" regions. The dominant clusters corresponding to

each category were then used for differential gene analysis. The "FindMarkers" function in

Seurat v4.2.0 was used to identify differentially expressed genes (DEGs) within each cluster

corresponding to Crypts, Villus-Tips, and Villus-Intermediate under both CT and PEM

conditions. Genes were classified as upregulated if they had log2 fold changes >0 and as

downregulated if they had log2 fold changes <0 with an adjusted p-value <0.05. Gene

ontology enrichment analysis of these DEGs was performed using clusterProfiler v4.12.1 [100]

with thresholds of adjusted p-value <0.05 (hypergeometric test).

DNA methylation profiling. DNA was isolated from IECs using the DNeasy Blood and Tissue

Kit (Qiagen) according to the manufacturer's protocol. DNA methylation levels were

measured using the Infinium Mouse Methylation BeadChip (Illumina), which covers > 285k

methylation sites spanning over CpG islands, transcription start sites, enhancers, and gene

regions. DNA methylation data was analysed using the Bioconductor package RnBeads

(version 2.12.1) [101]. Sites that overlapped with SNPs and had unreliable measurements were

filtered out. Context-specific probes, probes on the sex chromosomes, and probes with

missing values were also removed resulting in the removal of 54,378 out of 261,600 probes.

The signal intensity values were normalized using the dasen method. Differentially

methylated positions (DMPs) and regions (DMRs) between Ctrl_4w and PEM_4w samples as

well as Ctrl_10w and Recovery 10w samples were identified using the automatically selected

rank cutoff of RnBeads. Predicted transcription factor binding sites (TFBS) enriched in DMPs

were identified by conducting enrichment analysis using the Bioconductor package LOLA

(version 1.24.0) [97].

Integration of DNA methylation and transcriptome data. To integrate gene expression data

with DNA methylation data, we used a hierarchical approach as described before [32,102]. We

first identified DMPs located 5,000 bp upstream and downstream of the transcription start
sites of DEGs. Spearman's rank correlation coefficient between the normalized expression
count of each DEG and the methylation intensity (β-values) of its corresponding DMPs were
calculated. To test the statistical significance of the correlations, we calculated the false
discovery rate (FDR) using a permutation approach.

16S rRNA gene amplicon sequencing. DNA was isolated from fecal material using the DNeasy
PowerSoil Kit (Qiagen) following the manufacturer's protocol. Extracted DNA was eluted from
the spin filter silica membrane with 100 µl of elution buffer and stored at -80 °C. 16S profiling
and MiSeq sequencing was performed as described earlier [103,104], with the following
modifications: the V3-V4 region of the 16S rRNA gene was amplified using the dual barcoded
primers 319F (ACT CCT ACG GGA GGC AGC AG) and 806R (GGA CTA CHV GGG TWT CTA AT)
[105]. Each primer contained additional sequences for a 12 base Golay barcode, Illumina adaptor
and a linker sequence [106]. PCR was performed using the Phusion Hot Start Flex 2X Master Mix
(NEB) in a GeneAmp PCR system 9700 (Applied Biosystems) and the following program (98°C
for 3 min, 25-30x (98°C for 20 s, 55°C for 30 s, 72°C for 45 s), 72°C for 10 min, hold at 4°C).
Performance of the PCR reactions was checked using agarose gel electrophoresis.
Normalization was performed using the SequalPrep Normalization Plate Kit (Thermo Fisher
Scientific, Darmstadt, Germany) following the manufacturer's instructions. Equal volumes of
SequalPrep-normalized amplicons were pooled and sequenced on an Illumina MiSeq (2 x 300
nt). Quality control of the MiSeq 16S amplicon sequence data was performed in MacQIIME
v1.9.1 (http://www.wernerlab.org/software/macqiime) [107]. Sequencing reads were trimmed
keeping only nucleotides with a Phred quality score of at least 20, then paired-end assembled
and mapped onto the samples using the barcode information. Microbiome phylogenetic
analyses were conducted using QIIME2 [108] (release 2022.8). The paired-end sequences were
filtered, denoised, and chimera removed using DADA2 [109] plugin resulting in abundance tables
of amplicon sequence variants (ASVs) with on average 7.7K ± 4.2K (mean ± sd) reads per
sample. Taxonomy was assigned using the Bayesian classifier provided in DADA2 and using
the Silva rRNA database v.138 [110]. ASVs not assigned to "Bacteria" or annotated as
"Mitochondria", "Chloroplast", or "Eukaryote" were removed. QIIME2 outputs (feature table,
taxonomy assignation, and rooted tree), were combined into a single object using the qiime2R
(0.99.6) and Bioconductor phyloseq R (1.40.0) [111] packages.

Metagenomic sequencing. A total of 32 fecal DNA samples (8 Ctrl_4w, 8 Ctrl_10w, 8 PEM_4w, and 8 Recovery_10w), which were already included in the 16S profiling, were also subjected to shotgun metagenomic sequencing performed at the Competence Centre for Genomic Analysis (Kiel). DNA libraries were generated using the Illumina DNA Prep kit following the manufacturer's instructions. Libraries were then pooled and sequenced on an Illumina NovaSeq 6000 with 2 x 150 nt for 19.6 ± 4.14 (mean ± sd) million reads per sample. Shot-gun reads were processed using an in-house established workflow from our institute (https://github.com/ikmb/TOFU-MAaPO) that relies on BBtools and bowtie2 for mapping and host decontamination, and bioBakery tools (MetaPhlAn 4.0 and HUMaN 3.6) for taxonomic and functional potential profiling. Raw reads were quality trimmed and mapped to the mouse genome (GRCm39) to discern between sequenced reads from the host and the gut microbiome bacteria. Only samples containing >1 Gbp after trimming and host decontamination were kept in the analysis. Taxonomic features and microbial relative abundances were determined using MetaPhlAn 4.0 [112] with default parameters and its SGB database containing unique marker genes of 26,970 species-level genome bins (SGBs) derived from a collection curated 1.01M prokaryotic and metagenome-assembled genomes (MAGs). Functional profiles (stratified pathways, gene families and enzyme categories) were computed using HUMAnN 3.6 [113] with default parameters. All abundances were transformed to CPM (count per million) before any statistical analysis.

Microbiome phylogenetic analyses of 16S and metagenomics data. Uni- and multivariate analyses were performed in R (v.4.2.1) under phyloseq [111] (v.1.40.0), vegan [114] (v.2.6-2) and MAasLin2 [115] (v.1.10.0). All samples for diversity analyses were normalized by rarefaction to the minimum shared read count to account for differential sequencing depth among samples. Relative abundance was calculated by dividing the number of reads for an ASV by the total number of sequences in the sample. Within sample diversity (alpha diversity) was explored using various diversity indexes (Chao1, Shannon, Simpson, Observed ASVs) on ASV abundance data coupled to significance testing (Wilcoxon signed rank test or Kruskal-Wallis tests). Between-sample diversity (beta diversity) as well as differences in diet groups and time were explored and visualized using a Principal Coordinate Analysis (PCoA) plot based on Bray-Curtis distances. Associations of microbiome composition to specified covariates were tested with the implementation of PERMANOVA models (using the adonis2 function from the vegan

package). The P and R^2 values were determined by 10,000 permutations using time. To

explore variation between and within individuals per diet phase at the genus level, the

intraclass correlation coefficient (ICC) was computed by the *ICCest* function of the ICC R

package, using as input the longitudinal rarefied abundances, defining each individual mouse

as a distinct class. ICC estimation uses the variance components from a one-way ANOVA

(between-group variance and within-group variance). To detect differences in the abundance

of microbial features (16S rRNA, metagenomics) or pathways (metagenomics) between the

experimental groups over time, we built linear mixed models using the MaAslin2 [116] package

that included time, diet group and individual mice as a random variable (*microbial*

abundances ~ (1|mouse) + diet switch + time). P values were corrected for multiple hypothesis

testing using the Benjamin-Hochberg procedure, and a false discovery rate <0.05 was defined

as the significant threshold. Features (taxa, or pathways) appearing in at least 20% of the

samples were included in the analysis.

Identification of genes involved in the bacterial transformation of Linoleic acid-derived

oxylipins. Databases of known bacterial oxylipin transformation enzyme genes such as

Epoxide hydrolases (~66.000) and Lipooxigenases (~7800) were generated using the NCBI

protein database. All genes tagged with the following search criteria were included in the

database. (i) Epoxide hydrolases (10th of March, 2023): (epoxide hydrolase[Protein Name] OR

(epoxide[All Fields] AND hydrolase[All Fields])) AND (bacteria[filter] AND refseq[filter]). (ii)

Lipooxygenases (24th of March, 2023): Lipoxygenase[All Fields] AND bacteria[filter]. The

specific enzyme database (Epoxide hydrolases and Lipooxigenases) and the UniRef90

database [117] were input into the ShortBRED identify pipeline (v.0.9.5) and used to generate

protein-specific markers [118]. These markers along with the quality-controlled metagenomics

samples were used as input for ShortBRED quantify to get the presence of copy numbers of

the target protein markers.

Non-targeted metabolomics using HILIC UHPLC-MS/MS. Fecal pellets (~25mg) were weighed

in sterile ceramic bead tubes (NucleoSpin® Bead Tubes, Macherey-Nagel, Dueren, Germany)

and 1mL of methanol was added (−20°C; LiChrosolv, Supelco, Merck, Darmstadt, Germany).

Metabolite extraction was performed with Precellys® Evolution Homogenizer with following

settings (4,500 rpm, 40 seconds of 3 cycles with 2 seconds pause time, Bertin Corp., Rockville,

Maryland, USA). Samples were centrifuged for 10 minutes at 21,000 × g, cooled at 4°C and

900 µL of supernatant was transferred into new tubes. Fifty microliters of supernatant were

evaporated (40°C, SpeedVac Concentrator, Savant SPD121P, ThermoFisher Scientific,

Waltham, MA, United States) and reconstituted in 75% acetonitrile (ACN; LiChrosolv,

hypergrade for LC-MS; Merck KGaA). Samples were analyzed using an UHPLC system (ExionLC,

Sciex LLC, Framingham, MA, USA) coupled to a quadrupole time-of-flight mass spectrometer

(X500R QTOF MS, Sciex LLC, Framingham, MA, USA). Parameters of the MS for positive and

negative electrospray ionization mode are summarized in Table S3 and hydrophilic interaction

liquid chromatography (HILIC) was performed as described previously [119] to analyze small

polar metabolites.

Targeted metabolomics identification of 9-HODE and 12,13-diHOME. Identification and

semi-targeted peak picking of 12,13-diHOME ((9Z,12S,13S)-12,13-dihydroxyoctadec-9-enoic

acid) and 9-HODE was conducted with the same LC-MS method as described previously [120].

Here, we used an UHPLC system (ExionLC, Sciex LLC, Framingham, MA, USA) coupled to a

quadrupole time-of-flight mass spectrometer (X500R QTOF MS, Sciex LLC, Framingham, MA,

USA). We analyzed 12,13-diHOME standard, present on the GUTMLS01-1EA plate (Sigma-

Aldrich, Plate 1, Row A, final concentration of around 0.0025 µg/µL, dissolved and diluted in

methanol) and fecal extracts of CONVR and GF mice. The identity of 9-HODE was verified by

matching against spectral library of MassBank of North America. The X500R QTOF MS was set

to negative electrospray ionization mode with parameters summarized in Table S3.

Metabolomics data processing and analysis. Raw LC-MS (.wiff2) data were post-processed

using GeneData Expressionist Refiner MS 15.0 (GeneData GmbH, Basel, Switzerland) including

chemical noise subtraction (signal with intensities <200 were removed), chromatographic

peak picking, chromatogram isotope clustering, valid feature filter (features present in 10%

of samples and at least 500 maximum intensity), retention time range restriction (0.4-9

minutes), annotate known peaks (mass-to-charge tolerance of 0.005 Dalton and retention

time tolerance of 0.1 minutes), MS/MS consolidation (keep signals with highest total ion

count for one cluster in one sample), MS/MS peak detection and export to MASCOT generic

files. Processing resulted in a data matrix containing clusters (m/z and retention time values)

and observed maximum intensity values for each sample. Maximum intensity values were

normalized to the wet fecal weight. For each cluster we kept MS/MS with highest total ion

counts within one sample. MS/MS of clusters across samples were combined with the

consensusSpectrum() function of MSnbase (), using mzd of 0.005, minProp of 0.1 and sum for

intensityFun argument. Ions of MS/MS were matched with 0.005 Dalton, and intensities of

aggregated peaks were summed, if ions were present in minimum 10% of matched spectra.

Ions with intensities ≤500 and m/z values ≥ precursor m/z+2 were discarded from consensus

spectra. Identification was done by matching experimental MS/MS spectra against spectral

libraries, downloaded from MassBank of North America by using MS PepSearch (0.01 Dalton

mass tolerance for precursor and fragment Search). Furthermore, GNPS molecular

networking and GNPS Library Search was engaged for metabolite identification of

experimental MS/MS. Within MS PepSearch output, matches with dot product over 500 were

kept and multiple matches for one cluster were filtered for the best match. We kept clusters

that are present in at least 4 samples and clusters that have valid MS/MS information or

metabolite identities. With this approach, we kept 793 and 833 clusters for positive and

negative mode, respectively. Furthermore, SIRIUS was used for metabolite identification and

classification by an in-silico approach using MS/MS [121]. Multivariate statistical analysis was

then performed using Wilcoxon signed rank test implemented in R.

Statistical analysis. Biostatistical analyses were performed using GraphPad Prism (version 8)

software (GraphPad, Inc, La Jolla, CA) or R (v 3.2.5). Specific comparisons and analyses are

described in the individual method sections. All pairwise comparisons were performed using

Mann-Whitney U test and time-series multiple comparisons were done using Holm-Šídák's

multiple comparisons test. Data are shown as mean ± standard error of the mean (SEM).

References

1. World Health Organization (2024). World health statistics 2024: monitoring health for the SDGs, Sustainable Development Goals.

2. UNICEF, WHO, and WORLD BANK (2023). Level and trend in child malnutrition. World Health Organization, 4.

3. Soeters, P., Bozzetti, F., Cynober, L., Forbes, A., Shenkin, A., and Sobotka, L. (2017). Defining malnutrition: A plea to rethink. Clinical Nutrition 36, 896–901. https://doi.org/10.1016/J.CLNU.2016.09.032.

4. World Health Organization (2000). Turning the tide of malnutrition : responding to the challenge of the 21st century. Preprint at World Health Organization.

5. Smith, M.I., Yatsunenko, T., Manary, M.J., Trehan, I., Mkakosya, R., Cheng, J., Kau, A.L., Rich, S.S., Concannon, P., Mychaleckyj, J.C., et al. (2013). Gut microbiomes of Malawian twin pairs discordant for kwashiorkor. Science (1979) 339, 548–554. https://doi.org/10.1126/science.1229000.

6. Gordon, J.I., Dewey, K.G., Mills, D.A., and Medzhitov, R.M. (2012). The Human Gut Microbiota and Undernutrition. Sci Transl Med 4. https://doi.org/10.1126/SCITRANSLMED.3004347.

7. Barratt, M.J., Ahmed, T., and Gordon, J.I. (2022). Gut microbiome development and childhood undernutrition. Cell Host Microbe 30, 617–626. https://doi.org/10.1016/J.CHOM.2022.04.002.

8. Subramanian, S., Huq, S., Yatsunenko, T., Haque, R., Mahfuz, M., Alam, M.A., Benezra, A., DeStefano, J., Meier, M.F., Muegge, B.D., et al. (2014). Persistent gut microbiota immaturity in malnourished Bangladeshi children. Nature 510, 417–421. https://doi.org/10.1038/nature13421.

9. Gehrig, J.L., Venkatesh, S., Chang, H.W., Hibberd, M.C., Kung, V.L., Cheng, J., Chen, R.Y., Subramanian, S., Cowardin, C.A., Meier, M.F., et al. (2019). Effects of microbiota-directed foods in gnotobiotic animals and undernourished children. Science (1979) 365. https://doi.org/10.1126/SCIENCE.AAU4732/SUPPL_FILE/AAU4732_GEHRIG_TABLES-S1-S23.XLSX.

10. Chen, R.Y., Mostafa, I., Hibberd, M.C., Das, S., Mahfuz, M., Naila, N.N., Islam, M.M., Huq, S., Alam, M.A., Zaman, M.U., et al. (2021). A Microbiota-Directed Food Intervention for Undernourished Children. New England Journal of Medicine 384, 1517–1528. https://doi.org/10.1056/NEJMOA2023294/SUPPL_FILE/NEJMOA2023294_DATA-SHARING.PDF.

11. Barratt, M.J., Nuzhat, S., Ahsan, K., Frese, S.A., Arzamasov, A.A., Sarker, S.A., Munirul Islam, M., Palit, P., Islam, M.R., Hibberd, M.C., et al. (2022). Bifidobacterium infantis treatment promotes weight gain in Bangladeshi infants with severe acute malnutrition. Sci Transl Med 14, 1107. https://doi.org/10.1126/SCITRANSLMED.ABK1107/SUPPL_FILE/SCITRANSLMED.ABK1107_MDAR_REPRODUCIBILITY_CHECKLIST.PDF.

12. Mostafa, I., Fahim, S.M., Das, S., Gazi, M.A., Hasan, M.M., Saqeeb, K.N., Mahfuz, M., Lynn, H.B., Barratt, M.J., Gordon, J.I., et al. (2022). Developing shelf-stable Microbiota Directed Complementary Food (MDCF) prototypes for malnourished children: study

protocol for a randomized, single-blinded, clinical study. BMC Pediatr *22*, 1–9.
https://doi.org/10.1186/S12887-022-03436-6/TABLES/3.

13.   Heijmans, B.T., Tobi, E.W., Stein, A.D., Putter, H., Blauw, G.J., Susser, E.S., Slagboom,
P.E., and Lumey, L.H. (2008). Persistent epigenetic differences associated with
prenatal exposure to famine in humans. Proc Natl Acad Sci U S A *105*, 17046–17049.
https://doi.org/10.1073/PNAS.0806560105/ASSET/E45FA062-D76D-424C-9EBE-
A06956CB0B4C/ASSETS/GRAPHIC/ZPQ9990854020001.JPEG.

14.   Upadhyay, S.K., Agarwal, K.N., and Agarwal, D.K. (1989). Influence of malnutrition on
social maturity, visual motor coordination & memory in rural school children. Indian J
Med Res *90*, 320–327.

15.   Chopra, J.S., and Sharma, A. (1992). Protein energy malnutrition and the nervous
system. J Neurol Sci *110*, 8–20. https://doi.org/10.1016/0022-510X(92)90003-4.

16.   Stoch, M.B., Moodie, A.D., and Bradshaw, D. (1982). Psychosocial Outcome and CT
Findings after Gross Undernourishment during Infancy: a 20-year Developmental
Study. Dev Med Child Neurol *24*, 419–436. https://doi.org/10.1111/J.1469-
8749.1982.TB13647.X.

17.   Kar, B.R., Rao, S.L., and Chandramouli, B.A. (2008). Cognitive development in children
with chronic protein energy malnutrition. Behavioral and Brain Functions *4*, 1–12.
https://doi.org/10.1186/1744-9081-4-31/TABLES/3.

18.   Victora, C.G., Adair, L., Fall, C., Hallal, P.C., Martorell, R., Richter, L., and Sachdev, H.S.
(2008). Maternal and child undernutrition: consequences for adult health and human
capital. The Lancet *371*, 340–357. https://doi.org/10.1016/S0140-6736(07)61692-
4/ATTACHMENT/B220BF22-B4C5-4C79-8345-C6091D8A67AC/MMC10.PDF.

19.   Chattopadhyay, P.S., Gupta, S.K., Chattopadhyay, R., Kundu, P.K., and Chakraborti, R.
(1995). Malnutrition-Related Diabetes Mellitus (MRDM), Not Diabetes-Related
Malnutrition: A report on genuine MRDM. Diabetes Care *18*, 276–277.
https://doi.org/10.2337/DIACARE.18.2.276.

20.   TRIPATHY, B.B., and KAR, B.C. (1965). Observations on Clinical Patterns of Diabetes
Mellitus in India. Diabetes *14*, 404–412. https://doi.org/10.2337/DIAB.14.7.404.

21.   Hugh-Jones, P. (1955). DIABETES IN JAMAICA. The Lancet *266*, 891–897.
https://doi.org/10.1016/S0140-6736(55)92530-7.

22.   Taylor, A.K., Cao, W., Vora, K.P., De La Cruz, J., Shieh, W.J., Zaki, S.R., Katz, J.M.,
Sambhara, S., and Gangappa, S. (2013). Protein energy malnutrition decreases
immunity and increases susceptibility to influenza infection in mice. J Infect Dis *207*,
501–510. https://doi.org/10.1093/infdis/jis527.

23.   Chatraw, J.H., Wherry, E.J., Ahmed, R., and Kapasi, Z.F. (2008). Diminished primary
CD8 T cell response to viral infection during protein energy malnutrition in mice is
due to changes in microenvironment and low numbers of viral-specific CD8 T cell
precursors. J Nutr *138*, 806–812.

24.   Chandra, R.K. (1979). Nutritional deficiency and susceptibility to infection. Bull World
Health Organ *57*, 167.

25.   Hashimoto, T., Perlot, T., Rehman, A., Trichereau, J., Ishiguro, H., Paolino, M., Sigl, V.,
Hanada, T., Hanada, R., Lipinski, S., et al. (2012). ACE2 links amino acid malnutrition to
microbial ecology and intestinal inflammation. Nature *487*, 477–481.
https://doi.org/10.1038/nature11228.

26.   Guiraldes, E., and Hamilton, J.R. (1981). Effect of chronic malnutrition on intestinal
895   structure, epithelial renewal, and enzymes in suckling rats. Pediatr Res *15*, 930–934.
896   https://doi.org/10.1203/00006450-198106000-00010.

897   27.   Butzner, J.D., Butler, D.G., Miniats, O.P., and Hamilton, J.R. (1985). Impact of Chronic
898   Protein-Calorie Malnutrition on Small Intestinal Repair after Acute Viral Enteritis: A
899   Study in Gnotobiotic Piglets. Pediatric Research 1985 19:5 *19*, 476–481.
900   https://doi.org/10.1203/00006450-198505000-00014.

901   28.   Lauer, J., Ghosh, S., Ausman, L., Webb, P., Bashaasha, B., Agaba, E.,
902   Turyashemererwa, F., Tran, H., Gewirtz, A., Erhardt, J., et al. (2020). Markers of
903   Environmental Enteric Dysfunction Are Associated with Poor Growth and Iron Status
904   in Rural Ugandan infants. Curr Dev Nutr *4*, nzaa053_064.
905   https://doi.org/10.1093/CDN/NZAA053_064.

906   29.   Syed, S., Iqbal, N.T., Sadiq, K., Ma, J.Z., Akhund, T., Xin, W., Moore, S.R., Liu, E.,
907   Qureshi, S., Gosselin, K., et al. (2018). Serum anti-flagellin and anti-lipopolysaccharide
908   immunoglobulins as predictors of linear growth faltering in Pakistani infants at risk for
909   environmental enteric dysfunction. PLoS One *13*, e0193768.
910   https://doi.org/10.1371/JOURNAL.PONE.0193768.

911   30.   Phiri, T.N., Mutasa, K., Rukobo, S., Govha, M., Mushayanembwa, P., Mwakamui, S.,
912   Haider, T., Zyambo, K., Dumbura, C., Tome, J., et al. (2023). Severe acute malnutrition
913   promotes bacterial binding over proinflammatory cytokine secretion by circulating
914   innate immune cells. Sci Adv *9*.
915   https://doi.org/10.1126/SCIADV.ADH2284/SUPPL_FILE/SCIADV.ADH2284_SM.PDF.

916   31.   Iyer, S.S., Chatraw, J.H., Tan, W.G., Wherry, E.J., Becker, T.C., Ahmed, R., and Kapasi,
917   Z.F. (2012). Protein energy malnutrition impairs homeostatic proliferation of memory
918   CD8 T cells. J Immunol *188*, 77–84. https://doi.org/10.4049/jimmunol.1004027.

919   32.   Pan, W.H., Sommer, F., Falk-Paulsen, M., Ulas, T., Best, P., Fazio, A., Kachroo, P.,
920   Luzius, A., Jentzsch, M., Rehman, A., et al. (2018). Exposure to the gut microbiota
921   drives distinct methylome and transcriptome changes in intestinal epithelial cells
922   during postnatal development. Genome Med *10*, 27. https://doi.org/10.1186/s13073-
923   018-0534-5.

924   33.   McGee, D.W., and McMurray, D.N. (1988). The effect of protein malnutrition on the
925   IgA immune response in mice. Immunology *63*, 25–29.

926   34.   Fazio, A., Bordoni, D., Kuiper, J.W.P., Weber-Stiehl, S., Stengel, S.T., Arnold, P.,
927   Ellinghaus, D., Ito, G., Tran, F., Messner, B., et al. (2022). DNA methyltransferase 3A
928   controls intestinal epithelial barrier function and regeneration in the colon. Nat
929   Commun *13*. https://doi.org/10.1038/s41467-022-33844-2.

930   35.   Gizard, F., Amant, C., Barbier, O., Bellosta, S., Robillard, R., Percevault, F., Sevestre, H.,
931   Krimpenfort, P., Corsini, A., Rochette, J., et al. (2005). PPAR alpha inhibits vascular
932   smooth muscle cell proliferation underlying intimal hyperplasia by inducing the tumor
933   suppressor p16INK4a. J Clin Invest *115*, 3228–3238.
934   https://doi.org/10.1172/JCI22756.

935   36.   Daniel, B., Nagy, G., Czimmerer, Z., Horvath, A., Hammers, D.W., Cuaranta-Monroy, I.,
936   Poliska, S., Tzerpos, P., Kolostyak, Z., Hays, T.T., et al. (2018). The Nuclear Receptor
937   PPARγ Controls Progressive Macrophage Polarization as a Ligand-Insensitive
938   Epigenomic Ratchet of Transcriptional Memory. Immunity *49*, 615-626.e6.
939   https://doi.org/10.1016/J.IMMUNI.2018.09.005.

37. Mishra, N., Aden, K., Blase, J.I., Baran, N., Bordoni, D., Tran, F., Conrad, C., Avalos, D.,
Jaeckel, C., Scherer, M., et al. (2022). Longitudinal multi-omics analysis identifies early
blood-based predictors of anti-TNF therapy response in inflammatory bowel disease.
Genome Med *14*. https://doi.org/10.1186/S13073-022-01112-Z.

38. Wang, D., Day, E.A., Townsend, L.K., Djordjevic, D., Jørgensen, S.B., and Steinberg,
G.R. (2021). GDF15: emerging biology and therapeutic applications for obesity and
cardiometabolic disease. Nature Reviews Endocrinology 2021 17:10 *17*, 592–607.
https://doi.org/10.1038/s41574-021-00529-7.

39. Patel, S., Alvarez-Guaita, A., Melvin, A., Rimmington, D., Dattilo, A., Miedzybrodzka,
E.L., Cimino, I., Maurin, A.C., Roberts, G.P., Meek, C.L., et al. (2019). GDF15 Provides
an Endocrine Signal of Nutritional Stress in Mice and Humans. Cell Metab *29*, 707-
718.e8. https://doi.org/10.1016/J.CMET.2018.12.016/ATTACHMENT/0A532AF4-
416C-4CD4-BFB9-C8F898B06E4E/MMC2.PDF.

40. Groarke, J.D., Crawford, J., Collins, S.M., Lubaczewski, S., Roeland, E.J., Naito, T.,
Hendifar, A.E., Fallon, M., Takayama, K., Asmis, T., et al. (2024). Ponsegromab for the
Treatment of Cancer Cachexia. N Engl J Med.
https://doi.org/10.1056/NEJMOA2409515.

41. Bona, M., Medeiros, P.H., Santos, A.K., Freitas, T., Prata, M., Veras, H., Amaral, M.,
Oliveira, D., Havt, A., and Lima, A.Â. (2019). Virulence-related genes are associated
with clinical and nutritional outcomes of Shigella/Enteroinvasive Escherichia coli
pathotype infection in children from Brazilian semiarid region: A community case-
control study. Int J Med Microbiol *309*, 151–158.
https://doi.org/10.1016/J.IJMM.2019.01.003.

42. Gazi, M.A., Alam, M.A., Fahim, S.M., Wahid, B.Z., Khan, S.S., Islam, M.O., Hasan,
M.M., Hasan, S.M.T., Das, S., Mahfuz, M., et al. (2022). Infection With Escherichia Coli
Pathotypes Is Associated With Biomarkers of Gut Enteropathy and Nutritional Status
Among Malnourished Children in Bangladesh. Front Cell Infect Microbiol *12*.
https://doi.org/10.3389/FCIMB.2022.901324.

43. Rakoff-Nahoum, S., Paglino, J., Eslami-Varzaneh, F., Edberg, S., and Medzhitov, R.
(2004). Recognition of commensal microflora by toll-like receptors is required for
intestinal homeostasis. Cell *118*, 229–241. https://doi.org/10.1016/j.cell.2004.07.002
S0092867404006610 [pii].

44. Quaranta, A., Revol-Cavalier, J., and Wheelock, C.E. (2022). The octadecanoids: an
emerging class of lipid mediators. Biochem Soc Trans *50*, 1569–1582.
https://doi.org/10.1042/BST20210644.

45. Levan, S.R., Stamnes, K.A., Lin, D.L., Panzer, A.R., Fukui, E., McCauley, K., Fujimura,
K.E., McKean, M., Ownby, D.R., Zoratti, E.M., et al. (2019). Elevated faecal 12,13-
diHOME concentration in neonates at high risk for asthma is produced by gut bacteria
and impedes immune tolerance. Nat Microbiol *4*, 1851–1861.
https://doi.org/10.1038/S41564-019-0498-2.

46. Cunha, M.C.R., Lima, F. da S., Vinolo, M.A.R., Hastreiter, A., Curi, R., Borelli, P., and
Fock, R.A. (2013). Protein Malnutrition Induces Bone Marrow Mesenchymal Stem
Cells Commitment to Adipogenic Differentiation Leading to Hematopoietic Failure.
PLoS One *8*, e58872. https://doi.org/10.1371/JOURNAL.PONE.0058872.

47. Varnat, F., Heggeler, B.B. Ten, Grisel, P., Boucard, N., Corthésy-Theulaz, I., Wahli, W.,
and Desvergne, B. (2006). PPARβ/δ Regulates Paneth Cell Differentiation Via

Controlling the Hedgehog Signaling Pathway. Gastroenterology *131*, 538–553.
https://doi.org/10.1053/J.GASTRO.2006.05.004.

48. Aguilar-Recarte, D., Barroso, E., Gumà, A., Pizarro-Delgado, J., Peña, L., Ruart, M.,
Palomer, X., Wahli, W., and Vázquez-Carrera, M. (2021). GDF15 mediates the
metabolic effects of PPARβ/δ by activating AMPK. Cell Rep *36*.
https://doi.org/10.1016/J.CELREP.2021.109501.

49. Wan, Y., and Fu, J. (2023). GDF15 as a key disease target and biomarker: linking
chronic lung diseases and ageing. Molecular and Cellular Biochemistry 2023 479:3
479, 453–466. https://doi.org/10.1007/S11010-023-04743-X.

50. Sato, T., Stange, D.E., Ferrante, M., Vries, R.G.J., Van Es, J.H., Van Den Brink, S., Van
Houdt, W.J., Pronk, A., Van Gorp, J., Siersema, P.D., et al. (2011). Long-term
expansion of epithelial organoids from human colon, adenoma, adenocarcinoma, and
Barrett's epithelium. Gastroenterology *141*, 1762–1772.
https://doi.org/10.1053/J.GASTRO.2011.07.050.

51. Olsen, O.E., Skjærvik, A., Størdal, B.F., Sundan, A., and Holien, T. (2017). TGF-β
contamination of purified recombinant GDF15. PLoS One *12*, e0187349.
https://doi.org/10.1371/JOURNAL.PONE.0187349.

52. Mostafa, I., Nahar, N.N., Islam, M.M., Huq, S., Mustafa, M., Barratt, M., Gordon, J.I.,
and Ahmed, T. (2020). Proof-of-concept study of the efficacy of a microbiota-directed
complementary food formulation (MDCF) for treating moderate acute malnutrition.
BMC Public Health *20*, 242. https://doi.org/10.1186/s12889-020-8330-8.

53. Reyes, A., Blanton, L. V, Cao, S., Zhao, G., Manary, M., Trehan, I., Smith, M.I., Wang,
D., Virgin, H.W., Rohwer, F., et al. (2015). Gut DNA viromes of Malawian twins
discordant for severe acute malnutrition. Proc Natl Acad Sci U S A *112*, 11941–11946.
https://doi.org/10.1073/pnas.1514285112.

54. Schwarzer, M., Makki, K., Storelli, G., Machuca-Gayet, I., Srutkova, D., Hermanova, P.,
Martino, M.E., Balmand, S., Hudcovic, T., Heddi, A., et al. (2016). Lactobacillus
plantarum strain maintains growth of infant mice during chronic undernutrition.
Science (1979) *351*, 854–857. https://doi.org/10.1126/science.aad8588.

55. Storelli, G., Defaye, A., Erkosar, B., Hols, P., Royet, J., and Leulier, F. (2011).
Lactobacillus plantarum promotes Drosophila systemic growth by modulating
hormonal signals through TOR-dependent nutrient sensing. Cell Metab *14*, 403–414.
https://doi.org/10.1016/j.cmet.2011.07.012.

56. Kau, A.L., Planer, J.D., Liu, J., Rao, S., Yatsunenko, T., Trehan, I., Manary, M.J., Liu, T.C.,
Stappenbeck, T.S., Maleta, K.M., et al. (2015). Functional characterization of IgA-
targeted bacterial taxa from undernourished Malawian children that produce diet-
dependent enteropathy. Sci Transl Med *7*, 276ra24.
https://doi.org/10.1126/scitranslmed.aaa4877.

57. Charbonneau, M.R., O'Donnell, D., Blanton, L. V, Totten, S.M., Davis, J.C., Barratt,
M.J., Cheng, J., Guruge, J., Talcott, M., Bain, J.R., et al. (2016). Sialylated Milk
Oligosaccharides Promote Microbiota-Dependent Growth in Models of Infant
Undernutrition. Cell *164*, 859–871. https://doi.org/10.1016/j.cell.2016.01.024.

58. Hartman, S.J., Hibberd, M.C., Mostafa, I., Naila, N.N., Islam, Md.M., Zaman, M.U.,
Huq, S., Mahfuz, M., Islam, Md.T., Mukherji, K., et al. (2024). A microbiome-directed
therapeutic food for children recovering from severe acute malnutrition. Sci Transl
Med *16*. https://doi.org/10.1126/SCITRANSLMED.ADN2366.

59.   Chama, M., Amadi, B.C., Chandwe, K., Zyambo, K., Besa, E., Shaikh, N., Ndao, I.M.,
1033   Tarr, P.I., Storer, C., Head, R., et al. (2019). Transcriptomic analysis of enteropathy in
1034   Zambian children with severe acute malnutrition. EBioMedicine *45*, 456–463.
1035   https://doi.org/10.1016/J.EBIOM.2019.06.015.
1036   60.   Schulze, K. V., Swaminathan, S., Howell, S., Jajoo, A., Lie, N.C., Brown, O., Sadat, R.,
1037   Hall, N., Zhao, L., Marshall, K., et al. (2019). Edematous severe acute malnutrition is
1038   characterized by hypomethylation of DNA. Nature Communications 2019 10:1 *10*, 1–
1039   13. https://doi.org/10.1038/s41467-019-13433-6.
1040   61.   Blanton, L. V, Charbonneau, M.R., Salih, T., Barratt, M.J., Venkatesh, S., Ilkaveya, O.,
1041   Subramanian, S., Manary, M.J., Trehan, I., Jorgensen, J.M., et al. (2016). Gut bacteria
1042   that prevent growth impairments transmitted by microbiota from malnourished
1043   children. Science (1979) *351*. https://doi.org/10.1126/science.aad3311.
1044   62.   Mulenga, C., Sviben, S., Chandwe, K., Amadi, B., Kayamba, V., Fitzpatrick, J.A.J.,
1045   Mudenda, V., and Kelly, P. (2022). Epithelial Abnormalities in the Small Intestine of
1046   Zambian Children With Stunting. Front Med (Lausanne) *9*.
1047   https://doi.org/10.3389/FMED.2022.849677.
1048   63.   Kelly, P., Amadi, B., Chandwe, K., Besa, E., Zyambo, K., Chama, M., Tarr, P.I., Shaikh,
1049   N., Ndao, I.M., Storer, C., et al. (2021). Gene expression profiles compared in
1050   environmental and malnutrition enteropathy in Zambian children and adults.
1051   EBioMedicine *70*. https://doi.org/10.1016/j.ebiom.2021.103509.
1052   64.   Howell, K.J., Kraiczy, J., Nayak, K.M., Gasparetto, M., Ross, A., Lee, C., Mak, T.N., Koo,
1053   B.K., Kumar, N., Lawley, T., et al. (2018). DNA Methylation and Transcription Patterns
1054   in Intestinal Epithelial Cells From Pediatric Patients With Inflammatory Bowel
1055   Diseases Differentiate Disease Subtypes and Associate With Outcome.
1056   Gastroenterology *154*, 585–598. https://doi.org/10.1053/J.GASTRO.2017.10.007.
1057   65.   Kraiczy, J., Nayak, K.M., Howell, K.J., Ross, A., Forbester, J., Salvestrini, C., Mustata, R.,
1058   Perkins, S., Andersson-Rolf, A., Leenen, E., et al. (2019). DNA methylation defines
1059   regional identity of human intestinal epithelial organoids and undergoes dynamic
1060   changes during development. Gut *68*, 49–61. https://doi.org/10.1136/GUTJNL-2017-
1061   314817.
1062   66.   Li, H.J., Ray, S.K., Pan, N., Haigh, J., Fritzsch, B., and Leiter, A.B. (2019). Intestinal
1063   Neurod1 expression impairs paneth cell differentiation and promotes
1064   enteroendocrine lineage specification. Sci Rep *9*. https://doi.org/10.1038/S41598-
1065   019-55292-7.
1066   67.   Gross, B., Pawlak, M., Lefebvre, P., and Staels, B. (2017). PPARs in obesity-induced
1067   T2DM, dyslipidaemia and NAFLD. Nat Rev Endocrinol *13*, 36–49.
1068   https://doi.org/10.1038/NRENDO.2016.135.
1069   68.   Wang, D., Townsend, L.K., DesOrmeaux, G.J., Frangos, S.M., Batchuluun, B., Dumont,
1070   L., Kuhre, R.E., Ahmadi, E., Hu, S., Rebalka, I.A., et al. (2023). GDF15 promotes weight
1071   loss by enhancing energy expenditure in muscle. Nature *619*, 143–150.
1072   https://doi.org/10.1038/S41586-023-06249-4.
1073   69.   Coll, A.P., Chen, M., Taskar, P., Rimmington, D., Patel, S., Tadross, J.A., Cimino, I.,
1074   Yang, M., Welsh, P., Virtue, S., et al. (2020). GDF15 mediates the effects of metformin
1075   on body weight and energy balance. Nature *578*, 444–448.
1076   https://doi.org/10.1038/S41586-019-1911-Y.

70. Sommer, F., Anderson, J.M., Bharti, R., Raes, J., and Rosenstiel, P. (2017). The
resilience of the intestinal microbiota influences health and disease. Nat Rev
Microbiol. https://doi.org/10.1038/nrmicro.2017.58.
71. Bäckhed, F., Ding, H., Wang, T., Hooper, L. V, Koh, G.Y., Nagy, A., Semenkovich, C.F.,
and Gordon, J.I. (2004). The gut microbiota as an environmental factor that regulates
fat storage. Proc Natl Acad Sci U S A *101*, 15718–15723.
https://doi.org/10.1073/pnas.0407076101.
72. Turnbaugh, P.J., Ley, R.E., Mahowald, M.A., Magrini, V., Mardis, E.R., and Gordon, J.I.
(2006). An obesity-associated gut microbiome with increased capacity for energy
harvest. Nature *444*, 1027–1031. https://doi.org/nature05414 [pii]
10.1038/nature05414.
73. Itoh, T., Fairall, L., Amin, K., Inaba, Y., Szanto, A., Balint, B.L., Nagy, L., Yamamoto, K.,
and Schwabe, J.W.R. (2008). Structural basis for the activation of PPARγ by oxidized
fatty acids. Nature Structural & Molecular Biology 2008 15:9 *15*, 924–931.
https://doi.org/10.1038/nsmb.1474.
74. Levan, S.R., Stamnes, K.A., Lin, D.L., Panzer, A.R., Fukui, E., McCauley, K., Fujimura,
K.E., McKean, M., Ownby, D.R., Zoratti, E.M., et al. (2019). Elevated faecal 12,13-
diHOME concentration in neonates at high risk for asthma is produced by gut bacteria
and impedes immune tolerance. Nature Microbiology 2019 4:11 *4*, 1851–1861.
https://doi.org/10.1038/s41564-019-0498-2.
75. Nagy, L., Tontonoz, P., Alvarez, J.G.A., Chen, H., and Evans, R.M. (1998). Oxidized LDL
regulates macrophage gene expression through ligand activation of PPARgamma. Cell
93, 229–240. https://doi.org/10.1016/S0092-8674(00)81574-3.
76. Kincaid, J.W.R., Rimmington, D., Tadross, J.A., Cimino, I., Zvetkova, I., Kaser, A.,
Richards, P., Patel, S., O'Rahilly, S., and Coll, A.P. (2024). The gastrointestinal tract is a
major source of the acute metformin-stimulated rise in GDF15. Sci Rep *14*.
https://doi.org/10.1038/S41598-024-51866-2.
77. Suriben, R., Chen, M., Higbee, J., Oeffinger, J., Ventura, R., Li, B., Mondal, K., Gao, Z.,
Ayupova, D., Taskar, P., et al. (2020). Antibody-mediated inhibition of GDF15–GFRAL
activity reverses cancer cachexia in mice. Nature Medicine 2020 26:8 *26*, 1264–1270.
https://doi.org/10.1038/s41591-020-0945-x.
78. Groarke, J.D., Crawford, J., Collins, S.M., Lubaczewski, S.L., Breen, D.M., Harrington,
M.A., Jacobs, I., Qiu, R., Revkin, J., Rossulek, M.I., et al. (2024). Phase 2 study of the
efficacy and safety of ponsegromab in patients with cancer cachexia: PROACC-1 study
design. J Cachexia Sarcopenia Muscle *15*, 1054–1061.
https://doi.org/10.1002/JCSM.13435.
79. Badaloo, A. V., Forrester, T., Reid, M., and Jahoor, F. (2006). Lipid kinetic differences
between children with kwashiorkor and those with marasmus. Am J Clin Nutr *83*,
1283–1288. https://doi.org/10.1093/AJCN/83.6.1283.
80. Bartz, S., Mody, A., Hornik, C., Bain, J., Muehlbauer, M., Kiyimba, T., Kiboneka, E.,
Stevens, R., Bartlett, J., St Peter, J. V., et al. (2014). Severe Acute Malnutrition in
Childhood: Hormonal and Metabolic Status at Presentation, Response to Treatment,
and Predictors of Mortality. J Clin Endocrinol Metab *99*, 2128–2137.
https://doi.org/10.1210/JC.2013-4018.
81. Mahmud, I., Kabir, M., Haque, R., and Garrett, T.J. (2019). Decoding the Metabolome
and Lipidome of Child Malnutrition by Mass Spectrometric Techniques: Present Status

and Future Perspectives. Anal Chem *91*, 14784.
https://doi.org/10.1021/ACS.ANALCHEM.9B03338.

82. Oxidation of free fatty acids of plasma in infants with protein-calorie malnutrition |
South African Medical Journal
https://journals.co.za/doi/10.10520/AJA20785135_34805.

83. Sawada, M., Takahashi, K., Sawada, S., and Midorikawa, O. (1991). Selective killing of
Paneth cells by intravenous administration of dithizone in rats. Int J Exp Pathol *72*,
407.

84. Yilmaz, O.H., Katajisto, P., Lamming, D.W., Gultekin, Y., Bauer-Rowe, K.E., Sengupta,
S., Birsoy, K., Dursun, A., Yilmaz, V.O., Selig, M., et al. (2012). mTORC1 in the Paneth
cell niche couples intestinal stem-cell function to calorie intake. Nature *486*, 490–495.
https://doi.org/10.1038/nature11163.

85. Schmitt, M., Schewe, M., Sacchetti, A., Feijtel, D., van de Geer, W.S., Teeuwssen, M.,
Sleddens, H.F., Joosten, R., van Royen, M.E., van de Werken, H.J.G., et al. (2018).
Paneth Cells Respond to Inflammation and Contribute to Tissue Regeneration by
Acquiring Stem-like Features through SCF/c-Kit Signaling. Cell Rep *24*, 2312-2328.e7.
https://doi.org/10.1016/J.CELREP.2018.07.085.

86. Thurstans, S., Opondo, C., Seal, A., Wells, J., Khara, T., Dolan, C., Briend, A., Myatt, M.,
Garenne, M., Sear, R., et al. (2020). Boys are more likely to be undernourished than
girls: a systematic review and meta-analysis of sex differences in undernutrition. BMJ
Glob Health *5*, e004030. https://doi.org/10.1136/BMJGH-2020-004030.

87. WHO (2016). WHO recommendations on antenatal care. WHO Library Cataloguing,
172.

88. Agus, A., Planchais, J., and Sokol, H. (2018). Gut Microbiota Regulation of Tryptophan
Metabolism in Health and Disease. Cell Host Microbe *23*, 716–724.
https://doi.org/10.1016/J.CHOM.2018.05.003.

89. Cadwell, K., Liu, J.Y., Brown, S.L., Miyoshi, H., Loh, J., Lennerz, J.K., Kishi, C., Kc, W.,
Carrero, J.A., Hunt, S., et al. (2008). A key role for autophagy and the autophagy gene
Atg16l1 in mouse and human intestinal Paneth cells. Nature *456*, 259–263.
https://doi.org/10.1038/NATURE07416.

90. Adolph, T.E., Tomczak, M.F., Niederreiter, L., Ko, H.J., Böck, J., Martinez-Naves, E.,
Glickman, J.N., Tschurtschenthaler, M., Hartwig, J., Hosomi, S., et al. (2013). Paneth
cells as a site of origin for intestinal inflammation. Nature *503*, 272–276.
https://doi.org/10.1038/NATURE12599.

91. Sato, T., Vries, R.G., Snippert, H.J., van de Wetering, M., Barker, N., Stange, D.E., van
Es, J.H., Abo, A., Kujala, P., Peters, P.J., et al. (2009). Single Lgr5 stem cells build crypt-
villus structures in vitro without a mesenchymal niche. Nature *459*, 262–265.
https://doi.org/10.1038/nature07935.

92. nf-core/rnaseq: nf-core/rnaseq v3.14.0 - Hassium Honey Badger
https://doi.org/10.5281/ZENODO.10471647.

93. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15–21. https://doi.org/10.1093/BIOINFORMATICS/BTS635.

94. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose
program for assigning sequence reads to genomic features. Bioinformatics *30*, 923–
930. https://doi.org/10.1093/BIOINFORMATICS/BTT656.

95.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change
1170   and dispersion for RNA-seq data with DESeq2. Genome Biol *15*.
1171   https://doi.org/10.1186/S13059-014-0550-8.

1172   96.   Alexa, A., Rahnenfuhrer, J., and Lengauer, T. (2006). Improved scoring of functional
1173   groups from gene expression data by decorrelating GO graph structure.
1174   Bioinformatics *22*, 1600–1607. https://doi.org/10.1093/bioinformatics/btl140.

1175   97.   Sheffield, N.C., and Bock, C. (2016). LOLA: enrichment analysis for genomic region
1176   sets and regulatory elements in R and Bioconductor. Bioinformatics *32*, 587–589.
1177   https://doi.org/10.1093/BIOINFORMATICS/BTV612.

1178   98.   Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J.,
1179   Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal single-
1180   cell data. Cell *184*, 3573-3587.e29. https://doi.org/10.1016/J.CELL.2021.04.048.

1181   99.   Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of
1182   single-cell RNA-seq data using regularized negative binomial regression. Genome Biol
1183   *20*. https://doi.org/10.1186/S13059-019-1874-1.

1184   100.   Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for
1185   comparing biological themes among gene clusters. OMICS *16*, 284–287.
1186   https://doi.org/10.1089/OMI.2011.0118.

1187   101.   Assenov, Y., Müller, F., Lutsik, P., Walter, J., Lengauer, T., and Bock, C. (2014).
1188   Comprehensive analysis of DNA methylation data with RnBeads. Nat Methods *11*,
1189   1138–1140. https://doi.org/10.1038/NMETH.3115.

1190   102.   Bernardes, J.P., Mishra, N., Tran, F., Bahmer, T., Best, L., Blase, J.I., Bordoni, D.,
1191   Franzenburg, J., Geisen, U., Josephs-Spaulding, J., et al. (2020). Longitudinal Multi-
1192   omics Analyses Identify Responses of Megakaryocytes, Erythroid Cells, and
1193   Plasmablasts as Hallmarks of Severe COVID-19. Immunity *53*, 1296-1314 e9.
1194   https://doi.org/10.1016/j.immuni.2020.11.017.

1195   103.   Varga, A., Makszin, L., Bufa, A., Sipos, D., Kása, P., Pál, S., Rosenstiel, P., Sommer, F.,
1196   Kocsis, B., and Péterfi, Z. (2023). Efficacy of lyophilised bacteria-rich faecal sediment
1197   and supernatant with reduced bacterial count for treating patients with Clostridioides
1198   difficile Infection – A novel method for capsule faecal microbiota transfer. Front Cell
1199   Infect Microbiol *13*. https://doi.org/10.3389/fcimb.2023.1041384.

1200   104.   Castrillón-Betancur, J.C., López-Agudelo, V.A., Sommer, N., Cleeves, S., Bernardes,
1201   J.P., Weber-Stiehl, S., Rosenstiel, P., and Sommer, F. (2023). Epithelial Dual Oxidase 2
1202   Shapes the Mucosal Microbiome and Contributes to Inflammatory Susceptibility.
1203   Antioxidants (Basel) *12*. https://doi.org/10.3390/ANTIOX12101889.

1204   105.   Fadrosh, D.W., Ma, B., Gajer, P., Sengamalay, N., Ott, S., Brotman, R.M., and Ravel, J.
1205   (2014). An improved dual-indexing approach for multiplexed 16S rRNA gene
1206   sequencing on the Illumina MiSeq platform. Microbiome *2*.
1207   https://doi.org/10.1186/2049-2618-2-6.

1208   106.   Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N.,
1209   Owens, S.M., Betley, J., Fraser, L., Bauer, M., et al. (2012). Ultra-high-throughput
1210   microbial community analysis on the Illumina HiSeq and MiSeq platforms. ISME J *6*,
1211   1621–1624. https://doi.org/10.1038/ismej.2012.8.

1212   107.   Caporaso, J.G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F.D., Costello, E.K.,
1213   Fierer, N., Pena, A.G., Goodrich, J.K., Gordon, J.I., et al. (2010). QIIME allows analysis
1214   of high-throughput community sequencing data. Nat Methods *7*, 335–336.
1215   https://doi.org/10.1038/nmeth.f.303.

108. Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A.,
Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., et al. (2019). Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2. Nat
Biotechnol *37*, 852–857. https://doi.org/10.1038/S41587-019-0209-9.

109. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes,
S.P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data.
Nat Methods *13*, 581–583. https://doi.org/10.1038/NMETH.3869.

110. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and
Glöckner, F.O. (2013). The SILVA ribosomal RNA gene database project: improved
data processing and web-based tools. Nucleic Acids Res *41*.
https://doi.org/10.1093/NAR/GKS1219.

111. McMurdie, P.J., and Holmes, S. (2013). phyloseq: An R Package for Reproducible
Interactive Analysis and Graphics of Microbiome Census Data. PLoS One *8*, e61217.
https://doi.org/10.1371/journal.pone.0061217.

112. Blanco-Míguez, A., Beghini, F., Cumbo, F., McIver, L.J., Thompson, K.N., Zolfo, M.,
Manghi, P., Dubois, L., Huang, K.D., Thomas, A.M., et al. (2023). Extending and
improving metagenomic taxonomic profiling with uncharacterized species using
MetaPhlAn 4. Nature Biotechnology 2023 41:11 *41*, 1633–1644.
https://doi.org/10.1038/s41587-023-01688-w.

113. Beghini, F., McIver, L.J., Blanco-Míguez, A., Dubois, L., Asnicar, F., Maharjan, S.,
Mailyan, A., Manghi, P., Scholz, M., Thomas, A.M., et al. (2021). Integrating
taxonomic, functional, and strain-level profiling of diverse microbial communities
with biobakery 3. Elife *10*. https://doi.org/10.7554/ELIFE.65088.

114. Dixon, P. (2003). VEGAN, a package of R functions for community ecology. Journal of
Vegetation Science *14*, 927–930. https://doi.org/10.1111/j.1654-
1103.2003.tb02228.x.

115. Mallick, H., Rahnavard, A., McIver, L.J., Ma, S., Zhang, Y., Nguyen, L.H., Tickle, T.L.,
Weingart, G., Ren, B., Schwager, E.H., et al. (2021). Multivariable association
discovery in population-scale meta-omics studies. PLoS Comput Biol *17*, e1009442.
https://doi.org/10.1371/journal.pcbi.1009442.

116. Mallick, H., Rahnavard, A., McIver, L.J., Ma, S., Zhang, Y., Nguyen, L.H., Tickle, T.L.,
Weingart, G., Ren, B., Schwager, E.H., et al. (2021). Multivariable association
discovery in population-scale meta-omics studies. PLoS Comput Biol *17*, e1009442.
https://doi.org/10.1371/journal.pcbi.1009442.

117. Suzek, B.E., Wang, Y., Huang, H., McGarvey, P.B., and Wu, C.H. (2015). UniRef
clusters: a comprehensive and scalable alternative for improving sequence similarity
searches. Bioinformatics *31*, 926–932.
https://doi.org/10.1093/BIOINFORMATICS/BTU739.

118. Kaminski, J., Gibson, M.K., Franzosa, E.A., Segata, N., Dantas, G., and Huttenhower, C.
(2015). High-Specificity Targeted Functional Profiling in Microbial Communities with
ShortBRED. PLoS Comput Biol *11*. https://doi.org/10.1371/JOURNAL.PCBI.1004557.

119. Sillner, N., Walker, A., Harrieder, E.M., Schmitt-Kopplin, P., and Witting, M. (2019).
Development and application of a HILIC UHPLC-MS method for polar fecal
metabolome profiling. Journal of Chromatography B *1109*, 142–148.
https://doi.org/10.1016/J.JCHROMB.2019.01.016.

120.   Sillner, N., Walker, A., Koch, W., Witting, M., and Schmitt-Kopplin, P. (2018).
1262   Metformin impacts cecal bile acid profiles in mice. Journal of Chromatography B
1263   *1083*, 35–43. https://doi.org/10.1016/J.JCHROMB.2018.02.029.
1264   121.   Dührkop, K., Fleischauer, M., Ludwig, M., Aksenov, A.A., Melnik, A. V., Meusel, M.,
1265   Dorrestein, P.C., Rousu, J., and Böcker, S. (2019). SIRIUS 4: a rapid tool for turning
1266   tandem mass spectra into metabolite structure information. Nature Methods 2019
1267   16:4 *16*, 299–302. https://doi.org/10.1038/s41592-019-0344-8.
1268

1269

Declarations

Funding

This work was supported by the German Research Foundation (DFG) through the Collaborative Research Centre CRC1182 "Origin and Function of Metaorganisms" (project C2, A2 and Z3), the Excellence Cluster EXS2167 "Precision Medicine in Chronic Inflammation", the individual grant SO1141/10-1 and the Research Unit FOR5042 "miTarget - The Microbiome as a Target in Inflammatory Bowel Diseases" (projects P3, P4 and P7). RB is supported by the NIH grant DK088199. JMP received funding from the T. von Zastrow foundation and the German Federal Ministry of Education and Research (BMBF) under the project "Microbial Stargazing - Erforschung von Resilienzmechanismen von Mikroben und Menschen" (Ref. 01KX2324).

Acknowledgments

The authors thank Sabine Kock, Stefanie Baumgarten, Maren Reffelmann, Vivian Wegner, Tanja Klostermeier, Dorina Ölsner, Meike Hansen, Ronja Möhring and Sophie Reiher for excellent technical assistance. We further thank Hans-Christian Behrens for providing access to the Hamamatsu NanoZoomer S60 Digital Slide Scanner, Jacob Hamm for experimental support and Robert Häsler, Louise Thingholm, Eike Wacker and David Ellinghaus for valuable input regarding the RNAseq and metagenomic analyses. We thank Anthony Coll, John Tadross and Stephen O´ Rahilly, University Cambridge for insightful critical discussions and provision of reagents.

Author contributions

FS, NM, NS, FS and PR designed research. FS, NM, VL, NS, JPB, AW, FH, TG, FG, LS, AB, SK, SW, GI, FT, FS and PR performed experiments and analyzed the data. SS, JMP, RSB, PSK, JFB provided critical resources. FS, NM, VL, JPB, FS and PR prepared the figures. FS and PR obtained funding. FS, NM, VL, JPB, FH, FS and PR co-wrote the manuscript with critical input from all authors. All authors read and approved the final manuscript.

Declaration of Interests

1299   The authors declare no competing interests.

1300

1301   **Figures and Tables**

1302

1303 **Figure 1: PEM induces long-lasting changes in intestinal physiology independent of caloric**

1304 **intake. A)** Scheme of the PEM regimen in mice (n=10 mice per group). 12-week-old C57BL6

1305 mice were given either isocaloric control or protein-reduced food for 28 days to induce an

1306 acute PEM, and the PEM-mice were then switched to control diet as well for an additional 42

1307 days to monitor recovery from PEM. **B)** Body weight curve shows drastic weight loss during

PEM and incomplete recovery after switch to control diet. *p < 0.05, ***p < 0.001 and ****p

< 0.0001 using two-way ANOVA. **C)** Food intake of the experimental groups. ns = not

significant using Mann-Whitney U test. **D)** Lengths of the small and large intestine during PEM

and recovery. PEM shortened both the small and large intestine and the small intestinal

length could not be restored during recovery. **p < 0.01, ***p < 0.001 and ns = not significant

using Mann-Whitney U test. **E-F)** Fewer proliferating cells per crypt **(E)** and apoptotic cells per

villus **(F)** in PEM mice as determined by Ki-67 immunostaining and TUNEL assay including

representative images. The scale bar represents 50 μm. *p < 0.05, **p < 0.01 and ns = not

significant using Mann-Whitney U test.

1317

1318

Figure 2: Transcriptome analysis of purified intestinal epithelial cells reveals lasting antimicrobial and metabolic alterations. A, C. Gene ontology (GO) terms enriched in upregulated (A) and downregulated (C) genes only at 4 weeks (PEM only), 10 weeks (Recovery only) and at both 4 and 10 weeks (Shared). Dot size is proportional to the gene ratio and color corresponds to the p-value of enrichment. Top selected terms are visualized. B, D. Heatmaps showing the expression of DEGs corresponding to selected GO terms enriched in upregulated

1325   (**B**) and downregulated (**D**) shared genes. Scaled normalized gene expression counts across all

1326   samples are plotted. Note that due to the GO setup, genes can belong to multiple GO

1327   categories.

1328

Figure 3: PEM impairs Paneth cell differentiation and leads to spatially confined gene expression changes along the crypt-villus axis. A) Log2-fold change (y axis) of differentially expressed genes (DEGs) between mice under Ctrl and PEM diet at 4 weeks and 10 weeks (x axis) with n=10 per group. Color discriminates genes that are upregulated or downregulated only at 4 weeks (PEM only), 10 weeks (Recovery only) and at both 4 and 10 weeks (Shared), and point size represents statistical significance (adjusted p value). Paneth cell related genes are highlighted in blue. **B)** Boxplots showing average number of paneth cells (Lyz+) per crypt in duodenum and ileo-jejunal regions of the small intestine of mice under Ctrl and PEM diets along with representative histological images. ***: $p < 0.001$ Ctrl_4w versus PEM_4w and Ctrl_10w versus Rec_10w using Mann-Whitney U test. **C)** Classification of spots into multiple regions: crypts represented in green, intermediate villi in purple, and villi tips in orange. Identified spots were used for DEGs analysis across PEM vs Ctrl conditions. **D-E)** Top enriched

1341   gene ontology terms of significant upregulated **(D)** and downregulated **(E)** DEGs discovered

1342   in crypts are presented. DEGs were sorted into "up" (log2foldchange > 0) and "down"

1343   (log2foldchange < 0) categories with an adjusted p-value (padj) < 0.05. Similar steps of gene

1344   ontology analysis were then applied to the Villi-Intermediate and Villi-Tip regions.

Figure 4: PEM-dependent persistent DNA methylation changes and *ex vivo* stability of gene expression changes. A-B) Comparison of transcriptional PEM effects in intestinal epithelial cells (IEC, blue) and organoids (yellow) isolated of mice from two separate dietary PEM-recovery intervention experiments during **(A)** acute PEM and **(B)** after recovery. The dot plot

depicts the Log$_2$[fold-changes PEM_4w/10w vs Ctrl_4w/10w] whereas the Venn diagram

depicts the overlap in up- and downregulated DEGs. **C)** Number of differentially (hypo- and

hyper-) methylated positions (DMPs) detected by BeadCHiP array in small intestinal epithelial

cells from mice during acute PEM (4 weeks) and after recovery (10 weeks) and respective

controls. **D)** Shared DMPs between acute PEM (4 weeks) and recovery (10 weeks). **E)**

Predicted transcription factor binding sites enriched in observed DMPs. Dot size is

proportional to the odds ratio and color corresponds to the p-value of enrichment. Top

selected transcription factors are visualized. **F)** Schematic depiction of gene expression - DNA

methylation integration analysis. Spearman's rank correlation coefficient between the

methylation intensity and gene expression values was calculated between the DMPs located

within a range of 5kb before or after the transcription start site of the respective DEG. **G)**

Venn Diagram depicting the overlaps between shared DEGs between PEM_4w and Rec_10w

and DEGs correlated with their nearby DMPs, identifying 31 candidate DEGs linked with DMPs

including GDF15 that has been linked to body mass regulation.

1364

Figure 5: Acute PEM drastically alters microbiome composition and function. A) Stool

sample collection scheme for microbiome analyses. Stool samples were collected weekly

during the 10 weeks of diet experiment (normal or reduced protein diet) and used for 16S

amplicon sequencing analysis and taxonomy characterization. Shot-gun metagenomic sequencing and functional potential analysis was performed only on stool samples from weeks 4 and 10 (Ctrl_4w, PEM_4w, Ctrl_10w and Rec_10w, each n=10 per group). **B)** Longitudinal alpha diversity of stool samples from mice under normal and reduced protein chow. Shannon index was computed at ASV level. Longitudinal cross-sectional comparisons were performed using Wilcoxon signed rank test. ns: not significant, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$. **C)** Principal Coordinate Analysis based on Bray-Curtis's distances of stool fecal microbiomes of mice during PEM and recovery to explore further differences in microbiome composition. Differences in the centroids of groups were tested with PERMANOVA with 10000 permutations. **D)** Linear discriminant analysis (LDA) effect size (Lefse) of the fecal microbiomes at peak acute PEM (day 28, PEM_4w versus Ctrl_4w). The cladogram depicts the phylogenetic distribution of differential taxa and the bar plot depicts differential taxa ranked by LDA. Differences are shown by the color of the most abundant class, with red indicating PEM_4w and blue indicating Ctrl_4w. The diameter of each circle corresponds to the abundance of the taxon. This analysis uses the SILVA v138 taxonomy. The bar plot represents the top 30 taxa based on LDA scores. The letters in brackets denote the abbreviation of the taxon shown in the cladogram. **E)** Differential amino acid and fatty acid metabolic microbiome pathways (PEM_4w vs Ctrl_4w and Rec_10w vs Ctrl_10w). The triangles show the hues and direction of the effect size (b coefficient). Color intensity and size represent the magnitude of the effect size and FDR significance of the specific pathway, respectively.

1389

Figure 6: Absence of microbial signals attenuates the Paneth-cell phenotype and fecal metabolic alterations upon PEM. A) Body weight curves of GF mice without a microbiome under 4-week PEM regimen followed by 6-week recovery with regular diet compared to CONV-R mice with a regular microbiome. Note that GF mice displayed milder disease symptoms with significantly less body weight loss during PEM and a full recovery (n=10-11 mice per group). **B)** Paneth (LYZ1-positive) cell count in small intestinal sections of GF mice. **C)** Top 20 most significant lipid metabolites identified in feces of GF and CONV-R mice fed a CTRL or PEM for 28 days. **D)** Levels of 9-HODE and 12,13diHOME in feces of GF and CONV-R mice during the PEM & recovery experiment. **E)** Bacterial lipoxygenase and epoxide hydrolase (EH) genes identified in stool metagenomics of GF and CONV-R mice during PEM & recovery.

F) Lipoxygenase and EH genes in feces collected on day 0, 90 and 120 of malnourished children

undergoing a nutritional intervention with either a Microbiota Directed Complementary Food

(MDCF-2) or standard Ready-to-Use Supplementary Food (RUSF) using published

metagenomics data [10]. Data in (A) were compared by two-way ANOVA and in (B) and (D-F)

using Wilcoxon signed rank test. ns: not significant, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$,

****: $p < 0.0001$.

1406

Figure 7: Microbiome and PEM dependent factors block Paneth cell differentiation. A)
Expression of *Gdf15*, *Pparg* and *Pparbd* were assessed in small intestinal tissue of both CONVR
and GF mice subjected to PEM and after recovery. *p < 0.05, ***p < 0.001 and ****p < 0.0001,
ns = non-significant. Significance testing was performed using Wilcoxon-Mann-Whitney-Test.
B) Schematic drawing of organoid intervention experiment. Murine intestinal organoids were
first cultured in ENR-CV stem cell organoid medium and then Paneth cell differentiation was
induced using ENR-CD or PEM-CD medium. Stimulants (recombinant GDF15 [1 µg/ml], pan-

PPAR-agonist Lanifibranor [20 μM] or the lipid 9-HODE [1 μM] were added during initial ENR-
CD and PEM-CD culture and kept throughout the experimental duration. All experiments were
performed independently at least twice in triplicates. **C-E)** Relative *Lyz1*, *Defa5* and *Gdf15*
mRNA expression in organoids during Paneth cell differentiation and stimulated with **(C)**
recombinant GDF15, **(D)** the Pan-PPAR-agonist Lanifibranor or the lipid **(E)** 9-HODE. Note that
PPAR-activation boosts GDF15 levels thereby blocking Paneth cell differentiation and the
bacterial PEM metabolite 9-HODE also suppresses Paneth cell differentiation. *: $p < 0.05$, **:
$p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$ and ns = not significant using Mann-Whitney U test.

Figure S1: PEM drastically alters intestinal architecture and organ measures. A) Body weight development of adult mice fed either a control diet (20% protein) or PEM diets with various protein contents (0.7, 2 and 5%) for 3 weeks. B) Cecum, EWAT (epididymal white adipose tissue), liver and spleen weights relative to body weight of mice during the PEM diet and recovery intervention. C) Crypt depth, villus height and villus width from small intestine of mice during the PEM diet and recovery intervention.

1429

Figure S2: PEM during postnatal development aggravates the disease phenotype. A) Body

weight curve of 4-week-old mice weaned onto PEM diet for 4 weeks and then switched to

regular control diet for recovery. **p < 0.01, ***p < 0.001 and ****p < 0.0001 using two-way

ANOVA. **B**) Lengths of the small and large intestine during PEM and recovery. *p < 0.05 and

ns = not significant using Mann-Whitney U test. **C)** Cecum, EWAT (epididymal white adipose

tissue), liver and spleen weights relative to body weight of mice during the PEM diet and

recovery intervention. *p < 0.05, **p < 0.01 and ns = not significant using Mann-Whitney U

test.

1438

Figure S3: Differential regulation of genes in intestinal epithelial cells. A) Principal component analysis (PCA) for RNA sequencing samples. **B)** Number of significantly upregulated and downregulated genes in small intestinal epithelial cells from mice during acute PEM (4 weeks) and after recovery (10 weeks) and respective controls. **C)** Boxplots showing average number of goblet cells (PAS+) per villus and crypt in duodenum and ileo-jejunal regions of the small intestine of mice under Ctrl and PEM diets along with representative histological images. ns: $p > 0.05$ Ctrl_4w versus PEM_4w and Ctrl_10w versus Rec_10w using Mann-Whitney U test. **D)** Heatmap showing the expression of DEGs corresponding to selected GO terms enriched in upregulated and downregulated PEM only DEGs. Scaled normalized gene expression counts across all samples are plotted. **E)** Overlay of

1449   spatial transcriptomic spots classified by different regions on high-quality H&E stainings from

1450   PEM_4w and Ctrl_4w mice.

1451

Figure S4: Longitudinal microbiome functions affected by PEM and recovery. A) Heatmap of relative abundance of bacterial species (ASV) and families with significant changes under acute PEM (PEM_4w vs Ctrl_4w) and after recovery (Rec_10w vs Ctrl_10w). For each cell, colors indicate the row-wise z-score of relative abundances, asterisks denote the FDR significance at each cross-sectional comparison and prevalence represents the percentage of

non-zero features. Row-wise clusters represent features that belong to the same Phylum. **B)**

Longitudinal area-stacked relative abundances of the top 20 significant genera identified by

LefSe analysis. Unclassified genera and those with low relative abundance are grouped in

"Others". Relative abundances comparisons of two representative genera, *Bifidobacterium*

and *Escherichia-Shigella*. Wilcoxon signed rank test was used to test significance. **C)** Heatmap

of abundances of microbiome metabolic pathways (obtained from HUMAnN 3.6) with

significant changes under acute PEM (PEM_4w vs Ctrl_4w) and after recovery (Rec_10w vs

Ctrl_10w). For each cell, colors indicate the row-wise z-score of relative abundances, asterisks

denote the FDR significance at each cross-sectional comparison and prevalence represents

the percentage of non-zero features. Row-wise clusters represent features that belong to the

same higher-level pathway. **D)** Intraclass correlation coefficient (ICC) of individual genera

across diet phases. Statistical differences were assessed by Wilcoxon signed-rank test.

1469

Figure S5: Ameliorated PEM phenotype in GF mice. GF mice (n=10-11 per group) were

subjected to a dietary PEM-recovery intervention experiment and analyzed for organ

measures and cell counts. **A)** Length of the small and large intestines. **B)** Number of PAS-

positive goblet cells per villus-crypt-unit in intestinal small sections. **C)** Number of Ki67-

positive proliferative epithelial cells per crypt. **D)** Number of TUNEL-positive apoptotic

epithelial cells per crypt-villus unit. Significance testing was performed using Wilcoxon-Mann-

Whitney-Test.

1477

Figure S6: PPAR-activation or the microbial metabolite 9-HODE alone do with Microbiome

status and PEM drive *Ppar / Gdf15* expression and Paneth cell differentiation. Expression of

Lyz1 and *Gdf15* were assessed in intestinal organoids stimulated with the **(A)** pan-PPAR-

agonist Lanifibranor [20 µM] or the **(B)** lipid 9-HODE [1 µM] during Paneth cell differentiation.

Note that stimulation with Lanifibranor or 9-HODE did not alter *Lyz1* or *Gdf15* expression in

contrast to stimulation during PEM as shown in Figure 6.

 Table S1: Formulation of PEM diets.

Ingredient	U	0.7 % Protein	2.0 % Protein	5.0 % Protein	20.0 % Protein
Product No. (Ssniff)		*S0071-E710*	*S0071-E711*	*S0071-E712*	*S0071-E714*
Casein	%	**0.680**	**2.200**	**5.600**	**21.500**
L-Cystine	%	0.100	0.100	0.100	0.300
Maltodextrin	%	32.020	30.500	27.100	11.000
Corn starch, pre-gelatinized	%	32.400	32.400	32.400	32.400
Dextrose	%	12.000	12.000	12.000	12.000
Sucrose	%	0.200	0.200	0.200	0.200
Cellulose powder	%	7.000	7.000	7.000	7.000
Inulin powder	%	2.300	2.300	2.300	2.300
Vitamin premix	%	1.000	1.000	1.000	1.000
Mineral premix AIN	%	5.000	5.000	5.000	5.000
Choline Cl (50 %)	%	0.250	0.250	0.250	0.250
Dye [Y, G, B, V, R]	%	0.050	0.050	0.050	0.050
Soybean oil	%	7.000	7.000	7.000	7.000
Proximate contents					
Crude protein	%	**0.7**	**2.0**	**5.0**	**19.0**
Crude fat	%	7.0	7.0	7.0	7.0
Crude fibre	%	8.4	8.4	8.4	8.4
Crude ash (minerals)	%	4.2	4.2	4.2	4.3
Carbohydrates	%	76.5	75.1	71.7	56.1
Energy (Atwater)	MJ/kg	15.6	15.6	15.6	15.5
kcal% Protein		0.7	2	5	21
kcal% Fat		17	17	17	17
kcal% Carbohydrates		82	81	78	62

1486   **Table S2: List of** TaqMan® Gene Expression Assays **used in this study.**

Gene Name	Gene Symbol	Assay
Actin, beta	*Actb*	Mm00607939_s1
Lysozyme 1	*Lyz1*	Mm00657323_m1
Defensin, alpha, 5	*Defa5*	Mm00651548_g1
Growth differentiation factor 15	*Gdf15*	Mm00442228_m1
Peroxisome proliferator-activated receptor gamma	*Pparg*	Mm01184322_m1
Peroxisome proliferator-activated receptor beta/delta	*Pparbd*	Mm00803184_m1

1487

 Table S3: X500R MS parameters for non-targeted metabolomics and targeted identification

 of fatty acids (12,13-diHOME and 9-HODE)

	Non-targeted metabolomics		Targeted identification	Unit
Method duration (minutes)	13.5	13.5	25	min
Total Scan Time	0.412	0.412	0.412	s
Curtain gas	30	30	30	psi
Ion source gas 1	45	45	45	psi
Temperature	500	500	500	°C
Ion source gas 2	45	45	45	psi
CAD gas	7	7	7	-
Polarity	**Positive**	**Negative**	**Negative**	-
Spray voltage	5500	-4500	-4500	V
TOF start mass	50	65	65	Dalton
TOF stop mass	1000	1000	1000	Dalton
Declustering potential	50	-50	-50	V
DP spread	0	0	0	V
Accumulation time	0.1	0.1	0.1	s
Collision energy	5	-5	-5	V
CE spread	0	0	0	V
IDA Criteria	Small molecule	Small molecule	Small molecule	-
Maximum candidate ions	10	10	10	-
Intensity threshold	1000	1000	1000	cps
TOF start mass	50	50	50	Dalton
TOF stop mass	1000	1000	1000	Dalton
Declustering potential	80	-80	-80	V
DP spread	0	0	0	V
Accumulation time	0.025	0.025	0.025	s
Collision energy	35	-35	-35	V
CE spread	15	15	15	V

1490