1

Sensory Neural Noise as a Limiting Factor in Visual Working Memory

Precision in Neurotypicals and Individuals with Schizophrenia

Woon Ju Park[1*], Megan Ichinose[2], Gengshi Hu[2], Geoffrey F. Woodman[2,3,4], Duje
Tadin[5#], & Sohee Park[2,3#*]

[1]School of Psychology, Georgia Institute of Technology, Atlanta, GA, USA
[2]Department of Psychology, Vanderbilt University, Nashville, TN, USA
[3]Center for Integrative and Cognitive Neuroscience, Vanderbilt University, TN, USA
[4]Vanderbilt Vision Research Center, Vanderbilt University, Nashville, TN, USA
[5]Department of Brain and Cognitive Sciences, Department of Neuroscience,
Department of Ophthalmology and Center for Visual Science, University of Rochester,
Rochester, NY, USA

[#] Equally contributing senior authors

*Correspondence to:
Woon Ju Park
School of Psychology
Georgia Institute of Technology
654 Cherry St NW, Atlanta, GA 30332
E-mail: woonju@gatech.edu

or

Sohee Park
Department of Psychology
Vanderbilt University
111 21st Ave S., Nashville, TN 37240-7817
E-mail: sohee.park@vanderbilt.edu

ABSTRACT

The neural mechanisms that determine the capacity limits of working memory (WM) are not well understood, with traditional views identifying prefrontal circuitry as the main source of WM limits. Sensory noise, however, remains an underexplored explanation for WM limits despite its unavoidable influence on brain systems that encode and relay sensory information. Here, we show that individual differences in internal sensory noise during visual processing predict visual WM capacity. We further demonstrate that the well-documented WM deficits observed in schizophrenia can be associated with atypically high levels of visual sensory noise, offering a new explanatory framework for these deficits. By experimentally manipulating sensory noise via transcranial direct current stimulation applied to the visual cortex, we also show that changes in sensory noise led to corresponding changes in visual WM precision in individuals with schizophrenia. Finally, a computational model demonstrates that the same sensory noise measured during visual perception can limit WM precision during WM maintenance. These findings show that sensory noise may explain a significant amount of variance in WM function in both neurotypical adults and individuals with schizophrenia.

MAIN TEXT

INTRODUCTION

Working memory (WM) is central to flexible and adaptive human cognition, allowing intelligent guidance of behavior [1]. The efficacy of WM is determined by its capacity, with capacity limitations leading to impaired information processing [2–4]. WM deficits are present in a wide range of neuropsychiatric conditions, including schizophrenia, anxiety disorders, and dementias [5–9], and have a substantial impact on real-world functioning [10]. Thus, elucidating mechanisms that determine the capacity of WM is of broad scientific and clinical importance. WM storage involves coordinating the functioning of multiple sensory, attentional, and cognitive control mechanisms [11–13]. Traditional theoretical approaches have focused on prefrontal access and control as the primary constraint on WM effectiveness [14–16], and it remains controversial whether sensory processing plays a key role in WM capacity limitations [17–19].

Limitations in sensory processing have been overlooked as a critical determinant of WM capacity largely because the fidelity of sensory processing far exceeds the WM capacity for sensory information [20]. At first glance, this makes sensory processing an unlikely culprit for WM limitations. The argument, however, fails to consider how variations in noise levels within sensory processing could cascade and affect brain functions that rely on sensory processing [21]. Noise is ubiquitous in neural processing [22]: It is present in sensory transduction, cellular processes of neurons, neural responses, and motor outputs [23], limiting the efficiency of the brain's communication channels [24]. Noise present at earlier stages of sensory processing is also subject to amplification during typical gain processes [21]. Thus, atypical increases in sensory noise may have unavoidable cascading effects on a wide range of brain functions that rely on sensory inputs, including WM.

Electrophysiological, neuroimaging, and brain stimulation studies demonstrate that sensory areas remain engaged during the maintenance of visual WM representations [25–27], forming the basis of the *sensory-recruitment hypothesis* [17,21]. Building on this view,

we test the *sensory-noise hypothesis*, focusing on how sensory noise may limit WM

capacity. We reasoned that visual WM may be particularly and uniquely vulnerable to

sensory noise because it depends on the fidelity of sensory signals both for its initial

encoding and for the maintenance of stored sensory information. In this framework,

visual WM processing may be affected by sensory noise both in the feed-forward

processing of visual information to be stored in WM and in the active maintenance of

that information.

This framework makes several important predictions. First, it predicts that WM capacity

limits will be systematically related to sensory noise levels in the general population.

Second, it suggests that atypically high sensory noise could contribute to WM deficits

observed in neuropsychiatric populations, such as schizophrenia, where limitations are

often attributed to higher-level cognitive dysfunctions [28]. Third, the framework also

predicts that these limits should be changeable if we could manipulate noise levels in

the visual cortex itself. Finally, it predicts that sensory noise estimated during perception

should propagate and be amplified during the maintenance stage, thereby limiting WM

precision.

We tested these predictions using converging methods of psychophysics and

computational modeling, brain stimulation, and clinical science. Psychophysics

quantified sensory noise during visual perception, computational modeling formalized

how this noise limits WM precision, and brain stimulation provided a causal means to

manipulate sensory noise in visual cortex. In our clinical work, we focused on

schizophrenia where WM deficits have been proposed as an endophenotypic marker

and a core feature of this condition [8,9]. WM deficits in schizophrenia have been

characterized as a failure of high-level executive control [29]. However, schizophrenia is

also associated with broadly atypical sensory processing [30–33], with implications for

increased neural noise [34–36]. Together with our sensory noise hypothesis of WM, these

suggest a yet untested possibility that WM deficits in schizophrenia may derive from

impaired sensory mechanisms.

RESULTS

To test whether sensory noise limits visual WM, we asked both neurotypical adults (NT)

and individuals with schizophrenia or schizoaffective disorder (SZ) to complete visual

WM and orientation discrimination tasks (Study 1; 57 participants, 30 NT, 27 SZ, Table

1). The visual WM task required participants to remember locations and orientations of

one, two, or four gratings and reproduce the orientation of a probed item after a 1

second delay with an input dial (Figure 1A; Methods). WM recall variance was the

primary performance metric of interest. Less precise WM performance is reflected by

higher recall variance, as demonstrated by the characteristic increase in recall variance

(i.e., decline in precision of the reported orientation) as the number of items

remembered increases [37,38]. Given that internal noise is commonly assumed to add

variability to the internal representation [21,39], we hypothesized that elevated sensory

noise would lead to higher WM recall variance.

Table 1. Demographic and Clinical Information for Studies 1 and 2.

	Study 1		Study 2
	SZ (n = 27)	**NT** (n = 30)	**SZ** (n = 20)
M/F	14/13	16/14	11/9
Age	45.04 (8.10)	44.96 (7.55)	48.60 (9.84)
Years of education	13.00 (1.80)	15.65 (2.61)	13.00 (2.08)
IQ[a]	102.30 (9.03)	106.14 (8.41)	102.25 (8.11)
Handedness[b]	+60.18 (13)	+72.59 (2.61)	+66.00 (53.87)
Ethnicity/Race:			
% Minority	62.96	36.67	70
BPRS	17.07 (6.61)	-	20.7 (10.95)
SAPS	18.37 (14.52)	-	28.25 (15.92)
SANS	28.81 (11.96)	-	29.85 (17.90)
CPZE	380.46 (460.90)		394.94 (271.43)

Mean values are shown with *SD* in parentheses. BPRS, Brief Psychiatric Rating Scale
(Overall & Gorman, 1962); SANS, Scale for the Assessment of Negative Symptoms
(Andreasen, 1983); SAPS, Scale for the Assessment of Positive Symptoms
(Andreasen, 1984).

[a] National Adult Reading Test (NART; Nelson, 1982).
[b] Edinburgh Handedness Inventory (Oldfield, 1971).
[c] Chlorpromazine Equivalent Doses (mg/kg per day) (Woods, 2003).
147

Figure 1. **A. & B.** Procedures for the visual WM task and the orientation discrimination task.

See main text and Methods for details. **C.** Results for the visual WM task. Both NT (red) and SZ

(blue) showed worse WM performance with increasing set size, with SZ performing significantly

worse than NT. **D**. Results for the orientation discrimination task. Both NT (red) and SZ (blue)

showed the characteristic non-linear increase in contrast thresholds with increasing external

noise. SZ performed overall worse than NT, best explained by higher baseline internal noise

and unfiltered external noise in SZ (relevant statistics are in the main text).

To quantitatively estimate sensory noise in visual processing, we used the equivalent

noise paradigm—a well-established psychophysical approach that has been extensively

validated in both basic vision science [22,40,41] and clinical populations [42,43]. In this

paradigm, participants discriminated coarse orientation differences (±45° tilt from

vertical) of centrally presented gratings (1 cycle/°) embedded in varying levels of

external noise (Figure 1B; Methods). Performance across these external noise levels

was modeled with the Perceptual Template Model (PTM) [22]. The equivalent noise

approach leverages a systematic manipulation of external noise to dissociate different

sources of noise that limit perception. The method provides quantitative estimates of (i)

baseline additive internal noise, (ii) stimulus-dependent internal noise that scales with

stimulus strength, and (iii) the ability of the visual system to filter out external noise in

the stimulus (unfiltered external noise), which is commonly associated with tuning of

sensory neurons. A key insight behind the approach is that, at low levels of external

noise, performance is primarily limited by baseline internal noise, and external noise

only begins to affect performance at high noise levels that exceed baseline internal

noise.

We estimated sensory noise associated with visual orientation perception at both group

and individual levels. This allowed us to test whether individual differences in sensory

noise predict individual variability in WM precision, and whether there is a group

difference in sensory noise in SZ and NT. Importantly, we designed WM and noise

estimation tasks to involve the same type of visual information, oriented visual stimuli,

but to maximize differences in (a) stimulus salience, (b) task demands, and (c) how

subjects report their responses. The orientation discrimination task focused on stimulus

visibility, estimated contrast thresholds in noise (i.e., stimuli shown at low signal-to-noise

levels), and required only a coarse categorical response with a keystroke (±45° left vs

right tilt). In contrast, the WM task used highly visible noise-free grating stimuli (i.e., high

signal-to-noise) and required a precise report of the remembered orientation with a dial.

If we had measured perceptual noise using one feature, like orientation, and memory

using another feature, like color, then differences in sensory noise between these

features across individuals would be confounded with our measurements.

High baseline internal noise is associated with worse WM performance in NT

We began by examining whether sensory noise measured from the orientation

discrimination task was linked to WM performance in NT. In the visual WM task, NT

exhibited the expected pattern of greater WM variance with increasing set size (Figure

1C, red symbols; $F(2, 58) = 200.42$, $p < .001$, $\eta p^2 = .87$). In the orientation discrimination

task, the same participants showed a highly characteristic nonlinear result, with contrast

thresholds remaining fairly constant at low levels of external noise but increasing

linearly (in log space) at higher levels of external noise (Figure 1D, red) [22]. These data

were modeled with the PTM to estimate the baseline internal noise, stimulus-dependent

internal noise, and the unfiltered external noise for each participant (see Methods).

To test our hypothesis that sensory noise predicts WM capacity in NT adults, we used a

multiple linear regression with the noise estimates from PTM as predictors of the

average WM recall variance (Figure 2A & 2B, red). The model significantly accounted

for 31.5% of the variance in WM precision ($F(3, 26) = 3.99$, $p = .02$, $R^2 = .32$, R^2_{Adj}

$= .24$). However, only the baseline internal noise and unfiltered external noise were

significant predictors of WM recall variance on their own (baseline internal noise: $B =$

1.02, $SE = .44$, $t(29) = 2.31$, $p = .03$, $\eta p^2 = .17$; unfiltered external noise: $B = .21$, SE

$= .09$, $t(29) = 2.31$, $p = .03$, $\eta p^2 = .17$). The stimulus-dependent internal noise ($B = -$

11.75, $SE = 8.83$, $t(29) = -1.33$, $p = .20$, $\eta p^2 = .06$) did not significantly predict WM

variance. This suggests that sensory noise associated with perceiving orientations is

significantly associated with WM performance in NT adults.

Figure 2. Sensory noise limiting orientation discrimination was associated with WM

performance in both SZ (blue) and NT (red). **A**. In NT, baseline internal noise and unfiltered

external noise were significant predictors of WM performance. **B**. In SZ, unfiltered external noise

was a significant predictor of WM deficit. **C**. The 'larger' noise between the two (baseline internal

noise vs. unfiltered external noise; z-scored per group) for each individual was significantly

correlated with WM variance in both groups. This suggests that WM precision may be primarily

constrained by the dominant source of sensory noise within an individual.

Elevated sensory noise is associated with impaired WM performance in SZ

If sensory noise is a key limiting factor of visual WM precision, then a deficit in WM may

be, at least in part, due to a deficit in sensory processing. We tested this hypothesis in

SZ a disorder with a well-established WM deficit [8]. In the WM task, individuals with SZ

showed a marked impairment: the SZ group exhibited higher WM recall variance across

all set sizes relative to NT (Figure 1C, blue; $F(1,55) = 18.04$, $p < .001$, $\eta p^2 = .25$). There

was also a significant group by set size interaction ($F(2,110) = 5.05$, $p = .008$, $\eta p^2 = .08$)

with individuals with SZ showing greater deficits at larger set sizes. Because the WM

task relies on reporting remembered orientation with fine motor movements, we also

conducted a control motor precision task where participants reproduced the orientation

of visible gratings (see Methods). However, the results showed no significant

differences between NT and SZ ($t(55) = 1.65$, $p = 0.11$, $\eta p^2 = .05$), indicating that the

observed WM impairment in SZ was not due to differences in motor precision.

In contrast to the well-established WM deficit in SZ, much less is known about visual

orientation processing in SZ, particularly limitations related to sensory noise. Our results

from the orientation discrimination task reveal a previously unknown perceptual deficit in

SZ, with individuals with SZ exhibiting overall elevated contrast thresholds across a full

range of external noise levels (Figure 1D, blue). These group differences were best

explained by a model in which SZ and NT differed in the levels of both the baseline

internal noise and unfiltered external noise. Specifically, the best-fitting model indicated

a 61% increase in baseline internal noise along with a 21% increase in unfiltered

external noise in the SZ group relative to the noise levels of the NT ($R^2 = .98$). The

model was significantly different from the null model which assumed that the two groups

did not differ in their noise ($F(2, 26) = 29.95$, $p < .001$, $\eta p^2 = .70$, $\Delta R^2 = .03$), and was

not significantly different from the full model where all three types of noise varied

between the two groups ($F(1,25) = .69$, $p = .41$, $\eta p^2 = .03$, $\Delta R^2 = .0004$). These findings

together suggest that SZ is associated with elevated levels of sensory noise.

Critically, similar to NT, atypically elevated levels of visual noise in the SZ group

predicted their WM performance (Figure 2A & 2B, blue). To examine this relationship, a

multiple regression analysis was carried out within SZ with individually estimated

baseline internal noise and unfiltered external noise, shown to be different in SZ, as

predictors. The analysis revealed that the baseline internal noise and unfiltered external

noise together accounted for 34% of the variance in WM recall performance ($F(2, 24) =$

6.06, $p = .007$, $R^2 = .34$, $R^2_{Adj} = .28$), with only unfiltered external noise significantly

predicted WM recall variance on its own in SZ ($B = .19$, $SE = .06$, $t(26) = 3.12$, $p = .005$,

$\eta p^2 = .29$; baseline internal noise B = .29, SE = .44, $t(26) = .67$, $p = .51$, $\eta p^2 = .02$).

Our results are consistent with the hypothesis that the noise associated with sensory

processing limits visual WM. In NT, baseline internal noise and unfiltered external noise

predicted individual differences in WM performance. In SZ, both baseline internal noise

and unfiltered external noise were significantly elevated, with variations in the latter

predicting variability in WM. One possibility is that WM precision is primarily constrained

by the 'larger' noise (baseline internal noise vs. unfiltered external noise) that overrides

the other within an individual, similar to the rationale for the equivalent noise paradigm

explained above. To explore this further, we computed z-scores for both baseline

internal and unfiltered external noise separately for the NT and SZ groups. Then, for

each subject, we kept only the larger of the two z-scores. This simple max operation is

intended to capture noise that is more prominent for that subject relative to that subject's

peer group (NT or SZ). We then correlated the result with WM performance and found

that, for both NT and SZ, variability in sensory noise predicted visual WM precision

(Figure 2C; NT: $r(28) = .62$, $p = .0002$; SZ: $r(25) = .49$, $p = .009$).

Direct manipulation of sensory noise impacts WM performance in SZ

Next, we tested the observed link between sensory noise and WM in a causal design

using transcranial direct current stimulation (tDCS) to affect the noise in the sensory

processing (Study 2). The goal was to causally manipulate sensory noise and measure

its influence on WM performance within SZ. Using tDCS involves the application of low-

amplitude direct current to the scalp, thereby modulating the underlying excitability of

cortical neurons in a polarity-specific manner. We focused on anodal stimulation

because, in contrast to cathodal stimulation, it increases cortical excitability during and

after stimulation, a modulation that is associated with increasing neural noise [44–46].

Twenty participants diagnosed with schizophrenia or schizoaffective disorder (SZ; Table

1) completed three stimulation conditions: anodal stimulation over occipitoparietal

cortex, anodal stimulation over medial-frontal cortex, and sham (the location of which

was randomly assigned to either occipitoparietal or medial-frontal cortex; Figure 3A).

Stimulation over the occipitoparietal cortex served as our test condition based on prior

work indicating that anodal stimulation over the visual cortex can heighten neuronal

excitability underlying visual perception [47]. Active stimulation over the medial-frontal

cortex served as a spatial active control condition due to its distance from the target

sensory regions. Following 20 minutes of either active 2.0 mA stimulation or 20 minutes

of sham stimulation, participants immediately completed the same WM task and

orientation discrimination task described in the experiments above.

Across all analyses, the "change" following stimulation was defined as the difference

between the active (Medial-frontal or Occipitoparietal) and sham stimulation conditions

in either the WM or PTM noise metric.

Figure 3. **A**. Modeled current distribution for the medial-frontal and occipitoparietal stimulation sites. **B**. Visual WM performance in SZ following tDCS across medial-frontal (Fr, darker blue), occipitoparietal (Occ, blue), and Sham conditions. (Sh, lighter blue), **C**. Visual orientation discrimination performance following tDCS. **D**. Changes in sensory noise estimated using PTM following tDCS. The parameter estimates from the hierarchical Bayesian model (Methods) are depicted for each stimulation conditions (left column) and for changes between active and sham conditions (right column; FrSh: medial-frontal vs. sham; OccSh: occipitoparietal vs. sham).

We first tested whether stimulation over the occipitoparietal cortex would increase sensory noise, as measured by PTM noise parameters derived from the orientation

discrimination task. Participants' performance across the stimulation conditions again
yielded the characteristic nonlinear dependency on external noise (Figure 3C). To
characterize the effects of tDCS, the performances in the medial-frontal and
occipitoparietal stimulation conditions were each compared with that in the sham
condition separately using the PTM.

As hypothesized, visual orientation perception during the occipitoparietal stimulation
was characterized by a significantly higher level of baseline internal noise compared to
the sham condition (Figure 3D). Specifically, the effect of stimulation was best explained
by a model in which active occipitoparietal and sham differed in the level of baseline
internal noise. This model specified a 61% increase in internal noise with
occipitoparietal stimulation relative to sham ($R^2 = .96$). The model fit was significantly
better than a null model in which all noise parameters were constrained to be equal
($F(1, 27) = 4.36$, $p = .046$, $\eta p^2 = .14$, $\Delta R^2 = .01$), but did not differ from a full model in
which all noise sources were allowed to differ from the sham condition ($F(1, 27) = .42$, p
$= .66$, $\eta p^2 = .02$, $\Delta R^2 = .001$).

The increase in sensory noise was only observed following the occipitoparietal
stimulation. Compared to the sham condition, we found no evidence that medial-frontal
stimulation significantly impacted noise levels (Figure 3D). No model with any
combination of varying noise parameters was significantly different from the null model
in which all the noise parameters were held constant across active medial-frontal and
sham conditions (all Fs < 1.87, ps > .18; null model $R^2 = .97$).

The observation that the anodal tDCS over the visual cortex increases sensory noise
allowed us to test the hypothesized causal relationship between sensory processing
and WM capacity. On average, similar to Study 1, SZ showed worsening WM
performance with increasing set size (Figure 3B, $F(1,19) = 61.52$, $p < .001$, $\eta p^2 = .76$),
but, neither the medial-frontal nor the occipitoparietal tDCS stimulations significantly
influenced the overall WM recall variance at the group level (main effect of stimulation
condition: $F(2,38) = 1.26$, $p = .29$, $\eta p^2 = .06$), with no significant interaction between set
size and stimulation condition ($F(2,38) = .17$, $p = .84$, $\eta p^2 = .009$).

Figure 4. Relationship between changes in sensory noise and WM performance following tDCS stimulation in SZ. **A**. Changes in sensory noise following the medial-frontals stimulation did not significantly predict changes in WM performance as compared to sham. **B**. However, changes in baseline internal noise (left) following the occipitoparietal stimulation significantly predicted changes in WM performance.

However, the *concurrent changes* in sensory noise and WM performance *within individuals* were significantly correlated with each other following only the occipitoparietal stimulation (Figure 4B). In other words, although stimulation did not alter WM performance at the group level, individuals who showed a greater active-stimulation-related increase in baseline internal noise exhibited greater impairment in WM performance, but only following occipital stimulation. A multiple linear regression model with changes in all noise parameters (stimulation – sham) as predictors

explained changes in WM performance (stimulation – sham), accounting for 45% of

variance ($F(3,14) = 3.79$, $p = .04$, $R^2 = .45$, $R^2_{Adj} = .33$), and only the changes in

baseline internal noise on its own significantly predicted the changes in WM variance in

these respective conditions ($B = 12.56$, SE $= 5.31$, $t(17) = 2.36$, $p = .03$, $\eta p^2 = .29$;

stimulus-dependent noise: B $= -.30$, SE $= .15$, $t(17) = -2.05$, $p = .06$, $\eta p^2 = .23$; unfiltered

external noise: $B = .05$, SE $= .16$, $t(27) = .29$, $p = .78$, $\eta p^2 = .006$). In contrast, changes

in the noise parameters following medial-frontal stimulation did not significantly explain

changes in WM variance (Figure 4A; $F(3,13) = 1.72$, $p = .21$, $R^2 = .28$, $R^2_{Adj} = .12$).

Stated differently, increases in baseline internal noise specifically after occipitoparietal

stimulation predicted the degree to which WM became less precise and more variable.

Modeling the effects of sensory noise on WM

To understand the mechanistic ways in which the sensory noise contributes to WM

precision, we constructed a two-stage computational model of WM with encoding and

maintenance stages (see Methods for details). The model draws from prior observations

using delayed-estimation studies showing that WM precision degrades with set size and

delay [48]. A key assumption we make is that this reduction WM precision can arise from

stochastic drift in memory representations during maintenance [49], with its rate

determined by sensory noise.

Briefly, during encoding, stimulus orientations were represented by von Mises–tuned

orientation channels with Poisson-like variability and decoded using a population-vector

readout. Encoding precision decreased with set size. Sensory noise entered at this

stage in two forms: unfiltered external noise broadened channel tuning, while baseline

internal noise was modeled as circular Gaussian noise added to the decoded

orientations.

During the maintenance stage (1s delay), memory precision degraded through "drift" in

representations, implemented as diffusion of the encoded orientations over time [49] with

faster diffusion at larger set sizes. Critically, we assumed that the same sensory noise

that limits perception also adds variability to memory representations. Accordingly, the

diffusion rate was multiplicatively scaled by a subject-specific sensory-noise factor,

defined as the larger of the two z-scored noise sources estimated from the PTM. As a

result, sensory noise reduced WM precision both by limiting encoding fidelity and by

amplifying drift during maintenance.

We first simulated a full model in which PTM-derived baseline internal noise and

unfiltered external noise could limit WM precision at three distinct stages as explained

above (encoding tuning, post-decoding additive noise, and maintenance via diffusion

during the delay period). Consistent with our observed empirical data (Figure 1C), the

full model predicted reduced WM precision (higher WM variance) in the SZ group

relative to the NT group (Figure 5A).

Figure 5. **A**. Group-averaged predicted WM variance as a function of set size for NT and SZ. **B**.

Observed versus model-predicted WM variance for the full model across participants, shown

separately for NT and SZ groups, with dots colored by set size and regression lines derived

from mixed-effects models. **C**. Model comparisons. Left: bootstrapped Pearson correlation (r)

between observed and predicted variance for models in which each noise component was

separately simulated, with error bars showing 68% CI; the gray band indicates the 95% CI of the

full model. Right: paired-bootstrap difference in correlation (Δr) relative to the full model,

showing the relative contribution of baseline encoding noise, tuning-related encoding noise, and

maintenance drift to explaining empirical observed performance. The error bars show 95% CI.

To quantify the relationship between empirical and model-predicted WM variance, we fit

linear mixed-effects models in which the model-predicted WM variance was regressed

on observed empirical WM variance, while accounting for repeated measurements

across subjects and set sizes by including random intercepts for subject and for set

size. Separate models were fit for the NT and SZ groups. In both groups, empirical WM

variance significantly predicted model-predicted variance (Figure 5B; NT: β = .33, SE

= .05, t(58) = 7.15, p < .001; SZ: β = .22, SE = .08, t(55) = 2.75, p = .008), indicating

that the model explains inter-individual variability in WM precision.

To test whether this relationship between predicted and observed WM variance indeed

depended on individual-specific sensory noise estimates, we conducted a permutation

analysis in which the estimated PTM parameters were randomly shuffled across

individuals within each group. Model goodness of fit was quantified as the summed

partial R^2 of the empirical WM variance predictor across groups, derived from the

mixed-effects models similar to those above. The goodness-of-fit score obtained using

the true subject-specific PTM estimates significantly exceeded the shuffled null

distribution (permutation test, N = 1000; p = .01; true partial R^2 = .59; shuffled mean

partial R^2 = .12, SD = .13), demonstrating that preserving individual sensory noise

estimates significantly improved the model's ability to explain inter-individual WM

variability.

We then assessed which noise stage contributed most strongly to inter-individual

variability in WM variance. For this, we ran reduced "ablation" simulations in which only

one of the three noise stages was retained at a time (encoding tuning, post-decoding

baseline noise, or maintenance drift). For each reduced model, we performed a

bootstrap analysis with 1,000 iterations. Within each bootstrap iteration, subjects were

resampled with replacement within group, and empirical and predicted WM variance

were pooled across set sizes. Overall model goodness of fit was quantified as the

Fisher-z-aggregated Pearson correlation between empirical and predicted variance

across set sizes and groups.

The analysis revealed that sensory noise influencing the drift rate during the

maintenance stage accounted for the largest share of inter-individual variability in WM

variance (Figure 5C, left). To further quantify the relative contribution of each stage, we

computed the difference in correlation coefficient (Δr) between the full model and each

reduced model. Consistent with the ablation results, the model retaining only drift-

related noise showed the most similar explanatory power to the full model (Δr closest to

zero), suggesting that the sensory noise affecting the maintenance stage captured

much of the variance explained by the full model (Figure 5C, right).

Finally, we asked whether the model could account for within-subject changes in WM

variance following tDCS in the SZ group. Based on the tDCS and the above model

results, we focused on a reduced model in which baseline internal noise modulates WM

precision during the delay period by scaling the drift in the maintenance stage.

Accordingly, we simulated WM performance using a reduced model where only the

maintenance-stage diffusion term was noise-dependent and made subject-specific

predictions for sham and active stimulation conditions. For each participant, we

obtained the mean WM variance across set sizes and computed the condition

difference (stimulation − sham) for both empirical and predicted WM variance.

Consistent with our stimulation results, for occipitoparietal stimulation, changes in

predicted WM variance were positively correlated with changes in empirical WM

variance ($r(16) = .50$, $p = .03$). In contrast, the analogous relationship was not significant

for medial-frontal stimulation ($r(15) = .36$, $p = .15$). Together, these results support the

idea that the same sensory noise linked to the perception of visual orientations can

constrain visual WM precision during maintenance.

DISCUSSION

Our study provides a set of observations supporting the role of sensory noise in

determining WM limits. The findings suggest that noise present in early stages of visual

processing can limit the precision of WM at recall. Using a model of visual perception

[22,50], we show that individual differences in sensory noise can both explain individual

variability in WM capacity and account for the well-established WM deficit in individuals

with schizophrenia. Our manipulations of sensory noise with tDCS over the visual cortex

provide a causal link between sensory processing and WM, where changes in internal

noise with stimulation resulted in corresponding changes in WM precision. Our

computational model further suggests that the same sensory noise measured from

visual perception can limit WM precision during maintenance by affecting the drift of

WM representations. These findings indicate that lower-level sensory mechanisms

contribute to capacity limitations typically attributed to high-level cognitive processes.

We conjecture that visual WM is particularly vulnerable to sensory noise not only

because it relies on sensory information as its input, but more importantly, because it

recruits sensory mechanisms for the maintenance of stored information [17,21]. One

putative advantage of involving sensory mechanisms for information maintenance in

WM is the high fidelity of sensory processing. We posit that this is also what makes the

precision of visual WM particularly sensitive to noise levels within the visual process.

Because effective WM performance depends on multiple stages, including encoding

and maintenance [51,52], it is also susceptible to processing inefficiencies, including noise,

at each of those stages [4]. Both the sensory recruitment hypothesis and the sensory

noise hypothesis suggest that sensory areas are actively recruited in these stages,

where the fidelity of sensory representations directly influences WM precision. Our

model suggests that sensory noise can also influence maintenance. Sensory noise

likely is a potent factor even for maintenance because neural activity for memory

retention in the sensory cortex is generally weaker than when a physical stimulus is

present [53], leaving the representations more vulnerable to existing internal noise.

Our support for the idea that sensory noise affects WM storage does not preclude a role

of the prefrontal circuitry in determining WM capacity [18]. Instead, our findings add a

complementary view from the sensory processing perspective: in tasks that rely on

sensory details, such as orientation, color, or location, the fidelity of sensory cortex

representations is crucial. Studies have shown that WM capacity limits can be observed

even when visual stimuli remain in view [54]. This suggests that WM capacity results not

simply from remembering and maintaining representations but from performing

simultaneous tasks on multiple representations. In this context, sensory noise is critical

as it constrains the precision with which representations can be manipulated and

compared. At the same time, cognitive processes also shape operations to be

performed on these representations [55]. Thus, WM capacity is a joint outcome of

limitations in representational fidelity and the efficiency of higher-order control

processes.

Our results also show that the WM deficits in SZ, traditionally associated with

impairment in prefrontal processing, can be explained by elevated levels of sensory

noise. Our observation is consistent with a recent finding that implied faster WM decay

in SZ [56], which could result from noisy sensory representations. At the group level, the

WM precision in SZ was best explained by their ability to filter out external noise. In the

PTM, external noise filtering occurs at the earliest stages of visual processing, where

'templates' selectively enhance relevant signals and suppress irrelevant external noise.

The effectiveness of this filtering depends on the tuning properties of these templates.

Notably, prior work has demonstrated broader orientation tuning in SZ [57], which could

explain elevated levels of unfiltered external noise observed in our study. It is possible

that noisy representations due to broader orientation tuning in SZ are having a larger

effect on WM capacity than elevated baseline internal noise.

We also show that direct manipulation of baseline internal noise through brain

stimulation affects WM capacity in SZ. Anodal tDCS is known to increase cortical

excitability [58,59], which can modulate the spontaneous fluctuation in neural activity.

Given the lack of temporal precision and task-specificity in tDCS (stimulation is

continuously applied for 20 minutes prior to completing the tasks), it is less likely that

the stimulation selectively enhances task-related activity. If the tDCS rather amplifies

random neural variability, it can reduce the overall signal-to-noise ratio in neural

responses by increasing baseline neural noise. Our findings are not at odds with the

previous observation of increased visual sensitivity to low-contrast stimuli following

anodal tDCS [60], as in our task, increased excitability would amplify the responses for not

only the target signal (oriented grating) but also the external noise added to the stimuli.

It is also worth noting that the effects we observed were individual-specific: individuals

with SZ who showed an increase in baseline internal noise following tDCS to the

occipitoparietal cortex also exhibited a corresponding decline in WM precision.

Recognizing this variability provides critical considerations for designing individualized

intervention strategies.

Our findings identify the early visual system as a significant WM bottleneck in SZ.

Elevated sensory noise in visual cortex has far-reaching consequences in addition to

contributing to WM deficits. Other aspects of visual information processing may also be

impacted by increased sensory noise in SZ, including motion perception [35] and visual

context processing [61]. Furthermore, face discrimination deficits associated with

increased internal noise in SZ [36] could potentially cascade to social cognition

impairments. Increased sensory noise has also been observed in individuals with

autism spectrum disorders (ASD) [43,62], a population similarly noted for their WM deficits

[63,64]. In both SZ and ASD, elevated sensory noise may reflect or contribute to an

excitation/inhibition imbalance in cortical circuits [34], a mechanism that has been widely

proposed in both populations [65–67]. In other words, increased sensory noise may play a

mechanistic role in shaping higher-level cognitive outcomes such as WM in these

populations. Understanding how sensory noise interacts with developmental and

neuroanatomical factors, as well as its potential interactions with medications, could

provide key insights into the shared and distinct pathways through which WM deficits

emerge in a wide range of neuropsychiatric disorders [6,51,68–71].

In conclusion, we report both the correlational and causal evidence for the effects of

sensory noise on WM capacity. We conclude that the noise that limits the efficiency in

perceptual processing plays a critical role in constraining WM recall precision. The

results offer new insights into the sensory recruitment hypothesis of WM and highlight

sensory noise as a possible target for intervention (e.g., through brain stimulation or

perceptual training) that could enhance WM capacity in populations whose WM deficits

have been primarily associated with cognitive dysfunctions.

METHODS

PARTICIPANTS

Demographic and clinical information for participants who completed Studies 1 and 2 is summarized in Table 1. Twenty-seven (48% women; Study 1) and 20 (45% women; Study 2) medicated and clinically stable outpatients with chronic schizophrenia or schizoaffective disorder (SZ) were recruited from outpatient facilities in Nashville for participation. Diagnoses were made or confirmed according to the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision (DSM-IV-TR) criteria using the Structured Clinical Interview for DSM-IV (SCID-I/P) [72]. SCIDs were administered by masters-level clinical psychology graduate students.

In Study 1, 30 (47% women) neurotypical controls (NT) were recruited from the same metropolitan area through advertisements. NT had no history of DSM-IV Axis I disorders or family history of psychosis. We estimated premorbid IQ using the North American Adult Reading Test (NART) [73]. SZ and NT in Study 1 were matched on age, estimated IQ, and handedness but not on years of education ($t(55) = 4.18$, $p < .01$). Exclusion criteria included a history of head injury, neurological disorder, or substance abuse in the 6 months preceding the study.

All SZ participants were taking medications at the time of the study; in Study 1, 26 were taking atypical antipsychotics, 1 was taking a typical antipsychotic, 4 were taking mood stabilizers, and 10 were taking SSRIs, and in Study 2, 19 were taking atypical antipsychotics, 1 was taking a typical antipsychotic, 1 was taking a mood stabilizer, and 10 were taking SSRIs. Antipsychotic doses were converted to chlorpromazine equivalents for comparative analyses. Symptom severity in SZ was assessed with the Brief Psychiatric Rating Scale (BPRS; Overall and Gorman, 1962), the Scale for the Assessment of Negative Symptoms (SANS) [74], and the Scale for the Assessment of Positive Symptoms (SAPS) [75]. All participants were screened for normal or corrected-to-normal vision with the Snellen test of visual acuity. For the Snellen test, participants read through the chart using one eye at a time, viewed the letters from a distance of 20 feet, and read letters from the top of the chart aloud to an experimenter who scored their performance. Based on this viewing distance and the reference standard (20/20),

the 8th row of letters from the top of the chart consisted of letters subtending an angle of

5° with each letter part subtending 1°.

All participants provided written informed consent to study procedures approved by the

Vanderbilt University Institutional Review Board and were compensated for their

participation at a rate of $20 per hour.

STIMULI, TASKS, AND PROCEDURE

All tasks and stimuli in Studies 1 and 2 were programmed in MATLAB and Psychtoolbox

[76,77] and presented on a linearized monitor (20-inch Sony CRT; 1024 x 640 resolution;

120 Hz) on a gray background. A chin rest was used to maintain viewing distance at 77

cm with each pixel subtending 0.036°. An experimenter was present in the room during

both visual discrimination and working memory tasks to encourage task engagement.

Task order was counterbalanced across participants.

Visual Working Memory Precision Task

To assess visual WM, participants completed a WM task in which they were instructed

to remember locations and orientations of sinewave gratings (1 cycle/deg) in a two-

dimensional raised cosine envelope (radius = 1°). All gratings were presented at

maximum contrast to ensure visibility. The trial procedure was as follows: first, a

memory array consisting of one, two, or four oriented gratings was presented for 2 s.

Orientations were randomly selected from all possible options [-90°, 90°] with the

constraint that orientations within a given memory array were unique. Grating locations

for each trial were randomly chosen from eight positions around an invisible circle

(radius 4°) centered on a fixation point. After a short delay (1 s), a new, randomly

oriented [-90°, 90°] grating appeared at one of the prior locations (test probe).

Participants used an input dial (PowerMate USB Mulitmedia Controller; Griffin

Technology) to adjust the test probe's orientation so that it matched the orientation of

the grating presented at that same location. Participants were asked to be as precise as

possible when matching orientations and responses were untimed. Participants

indicated their final response by pressing down on the dial, leading to a 3 s inter-trial

interval. Trials began with a centrally-presented dynamic fixation sequence [78].

Participants completed a total of 450 trials, split equally into 10 blocks with rest periods

lasting a minimum of 30 s.

Because group differences in ability to use the manual dial may confound hypothesized

group differences in WM precision, participants also completed a short motor precision

task that did not tax memory. Participants completed the motor precision task prior to

the WM task in order to become acclimated to using the manual dial. Trials involved

presentation of a randomly oriented grating located at one of eight positions around an

invisible circle (radius 4°) that remained on the display. A randomly oriented grating was

then presented at the center of the display, and participants were instructed to use the

manual dial to adjust the orientation of this central grating until it was aligned with the

orientation of the first grating, which remained on the display until the participant's

response was submitted. Consistent with instructions on the WM task, participants were

asked to be as precise as possible when matching orientations and responses were

untimed. Participants completed a total of 24 trials such that each of the 8 possible

grating locations was sampled three times.

This manual recall method has been used with both healthy adults and psychiatric

populations to obtain a more detailed, continuous measure of visual WM precision

compared to typical binary match/non-match response paradigms [21,79]. Importantly, this

method enables WM performance to be analyzed as a distribution of recall errors (the

difference between input response and target orientation) with accompanying

descriptive properties of the distribution. As the number of visual features or items to be

remembered increases, the decline in recall precision is captured by the increasing

variance of the error distribution.

Orientation Discrimination Task

Participants performed a coarse orientation discrimination task [43] where they judged the

tilt of centrally presented gratings with the same spatial frequency and size as those in

the WM task (tilted ±45° from vertical). Gratings were temporally embedded in two

independent external noise samples. The grey-level pixel values for external noise were

sampled from a Gaussian distribution with a mean of 0 and SD that varied across trials,

such that its root mean square contrast defined the external noise level to be one of 8

set levels ranging from 0-21% (0, 0.3, 0.61, 1.24, 2.51, 5.1, 10.3, or 21%). The rapid

temporal succession of noise, stimulus, and noise frames (16.7 ms duration per frame)

results in an intentional perceptual merging of the grating and noise frames that is

consistent with PTM experiments, as it allows for finer adjustments of stimulus contrast

[50]. Participants indicated their response by a left or right arrow key press, and brief

auditory feedback (50 ms tone) was provided after correct trials to maintain task

engagement. As in WM task, trials started with a dynamic, centrally-presented fixation

sequence. Participants completed 680 trials, split equally into 8 blocks with rest periods

lasting a minimum of 30 s.

The stimulus contrast in the visual discrimination task was adjusted every trial using the

Functional Adaptive Sequential Testing (FAST toolbox [80]) procedure, which determines

the most informative contrast levels to be sampled for estimating participants' contrast

thresholds. Two FAST structures were interleaved to estimate psychophysical functions

reflecting the contrast level required to elicit reliable orientation discrimination at two

different sensitivity levels (d'=1.089 and 1.634), corresponding to 70.71% and 79.37%

accuracy. Of note, data analyses were completed separately from the within-task FAST

procedure in order to limit biased threshold estimations from potential increased

accidental error responses in SZ. See Park et al. (2017) [43] for a complete description of

the psychophysical procedure.

TRANSCRANIAL DIRECT CURRENT STIMULATION

The tDCS was administered (Study 2) using a battery driven, constant current stimulator

(MindAlive Inc., Alberta, Canada) and a pair of conductive rubber electrodes (active:

19.25 cm2, reference: 52 cm2). The electrodes were placed over saline-soaked

sponges and each held in place with elastic cloth headbands. In the active anodal

conditions, current is applied for 20 minutes. Stimulation intensity is fixed at 2.0 mA, as

this intensity demonstrated improvement in visual acuity in a prior study [47]. For medial

frontal stimulation, the anodal electrode was placed over the medial frontal lobe at

midline (site FCz of the International 10-20 System) and the cathodal electrode as

placed over the right cheek to avoid confounding effects from other brain regions. For

occipitoparietal stimulation, the anodal electrode was placed over occipitoparietal cortex

at midline (site Pz of the International 10-20 System), with the cathodal electrode again

placed over the right cheek. Consistent with tDCS montages in related work, the

placement of the cheek electrode is on a diagonal, parallel to the jaw line, 3 cm from the

lip corner (cheilion) above the jaw [47,81]. Location of anodal electrode placement over

FCz or Pz was determined with individual scalp measurements. We modeled the

current flow using COMETS and our previously described methods for estimating the

fields generated by the stimulator [59,82].

The tDCS sham condition involves similar administration as that for the active condition,

except the stimulation only lasts 30 seconds, which ramps up and down at the

beginning and end of the 20-minute period. This sham procedure results in the same

physical sensations (e.g., tingling, itching) that are reported with active tDCS [47].

Following each session, participants completed a questionnaire and visual analog scale

that included questions regarding attention, concentration, mood, vision, headache,

fatigue, and skin sensations during tDCS. Participants were additionally asked to rate

whether they had received active versus sham stimulation and the degree of confidence

associated with those ratings. Participants tended to overestimate the presence of

stimulation, leading to an accuracy rate of detecting the correct stimulation condition of

57%. However, the hit rate for detecting sham stimulation was 15%, well below chance,

suggesting that participants were blind to the sham versus active conditions.

Additionally, confidence ratings did not differ across stimulus condition ($p > .23$). Scores

across all questionnaire ratings did not significantly differ between stimulation conditions

(ps > .092).

The present study followed a within-subjects design, such that each participant

completed the three stimulation conditions (anodal active over medial-frontal, anodal

active over occipitoparietal, and sham) on three separate days. Session order was

counterbalanced across participants, and the location of sham (medial

frontal/occipitoparietal) was randomly assigned. The time interval between session visits

was at least 48 hours to minimize potential task practice effects or carryover effects

related to the previous session's brain stimulation (M=6.60 days between sessions,

SD=4.11 days) [83].

All study sessions began with 20 minutes of active or sham tDCS applied over medial

frontal or occipitoparietal cortex. Immediately after tDCS, participants completed the

visual WM precision task (20-30 minutes), followed by the visual discrimination task (40-

60 minutes). For the WM task, participants completed a shortened version of the visual

WM task described above in order to reduce task duration and cognitive burden on

participants. Participants completed a total of 60 trials, split equally into 3 blocks with

rest periods lasting a minimum of 30 s. While the trial procedure was identical, memory

arrays only consisted of two of four oriented gratings. In total, the two tasks lasted

approximately 90 minutes, which is within the time window in which tDCS effects have

been shown to last in comparable stimulation procedures [81].

ANALYSES

Visual Working Memory Task Analysis

Grating orientations were recorded and reported in the parameter space of all

possible line orientation values [-90°, 90°) and converted to the circular space [-π, π)

radians. Recall error on each trial was calculated as the difference between the

orientation of the target grating and the orientation of the test probe that was reported

(input with the manual dial) by the participant. Given that internal noise is hypothesized

to add variability to, or destabilize, the internal representation [21,39], the primary

performance metric of interest on the visual WM task was participants' recall variance.

Poorer WM performance is thus reflected by higher recall variance, as demonstrated by

the characteristic increase in recall variance (decline in precision) as the number of

items remembered increases [37,38]. The Von Mises probability density function (circular

normal distribution) with mean $\theta = 0$ (no error, such that recall is centered at the target

memory location) and σ was used to fit every participant's distribution of recall errors at

each memory array set size. The Von Mises probability density function for a given

angle x, centered at θ with a concentration κ where $I_0(\kappa)$ is the modified Bessel function

of order 0:

$f(x|\theta,\kappa) = \dfrac{e^{\kappa \cos (x-\theta)}}{2\pi I_0(\kappa)}$.

The variance of the Von Mises probability density function (σ^2), which reflects the

dispersion of the distribution, is analogous to the inverse of the distribution's

concentration ($1/\kappa$). The concentration parameter (κ) was estimated from each

participant's recall errors at every memory array set size using a maximum likelihood

method. Estimates of distribution concentration were converted to variances by taking

the inverse for subsequent between-group and individual differences analyses.

Orientation Discrimination Task

Data from the visual discrimination task were analyzed with two different approaches.

The first involved a conventional PTM analysis to estimate model parameters at a group

level, thus allowing us to test for relative groups differences in levels of noise in the

three noise parameters between SZ and NT. The second approach utilized hierarchical

Bayesian model fitting to estimate each individual's model parameters. The latter thus

allowed for a better understanding of individual differences in noise estimates. It also

enabled us to examine how noise estimates related to WM performance.

Conventional PTM Analysis. To examine group-level differences in noise estimates,

visual discrimination performance was first analyzed using a conventional PTM fitting

method. First, psychophysical contrast thresholds were estimated for each participant at

each of the 8 external noise levels with data pooled across the two FAST structures.

Individuals' contrast thresholds for orientation discrimination were estimated with

Weibull functions at every external noise level using the following equation:

$$P(c) = 1 - (1 - 0.5) \times 2^{-(\frac{\log (c)}{a})^{\eta}}$$

where P represents accuracy (percent correct), c is stimulus contrast, α is threshold at

75% accuracy, and η is the slope. The two free parameters, α and η, were estimated

with a Bayesian model fitting method implementing a Markov Chain Monte Carlo

(MCMC) technique. Group averages of individuals' estimated thresholds were used to fit

the conventional PTM. Coefficient indices for each noise source (unfiltered external

noise (A_e), baseline internal noise (A_a), stimulus-dependent internal noise (A_m)), were

introduced into the standard PTM to account for differences in noise levels between

groups:

749
$$c_\tau = \frac{1}{\beta} \left[\frac{(1 + (A_m N_m)^2)(A_e N_e)^{2\gamma} + (A_a N_a)^2}{(\frac{1}{d'^2} - (A_m N_m)^2)} \right]^{\frac{1}{2\gamma}}$$

During PTM fitting, these three SZ coefficients (A_e, A_a, A_m) could freely vary or be fixed

at 1, with the latter constraining a given noise estimate to be equal across groups

(Study 1) or stimulation conditions (Study 2) and the former allowing for relative

differences between groups or stimulation conditions in a given noise estimate. To

determine if noise sources differed between groups or conditions, all possible candidate

models accounting for each combination of free and fixed noise estimates were run: the

null model, which assumed no differences in any of the noise sources, the full model,

which assumed differences in all three noise sources, and six remaining models

reflecting the different possible combinations of the three noise estimates. In addition to

which ever noise coefficients were specified to freely vary, all candidate models

included the following four free parameters: stimulus-dependent internal noise (N_m),

internal additive noise (N_a), signal gain from the perceptual template (β), and the

exponent of the nonlinear transducer function (γ). Thus, the full model included 7 free

parameters, the null model included 4 free parameters, and the other six models

included between 5 and 6 free parameters. The eight candidate models were each fitted

to average thresholds using a least-squares method. Candidate models were compared

by evaluating differences in goodness-of-fit using r^2. The best fitting model was

determined with the F-test for nested models. Specifically, the best fitting model was

identified as the model with the fewest free parameters that was not significantly

different from the full model, in which all parameters could freely vary.

Hierarchical Bayesian Model Analysis. A hierarchical Bayesian modeling technique was

used to fit each participant's data with the PTM in order to obtain individual noise

parameter estimates and examine relations between noise estimates and visual WM

performance [43]. Unlike in the conventional PTM analysis approach described above, in

which model parameters are estimated from the contrast thresholds at each level of

external noise, the hierarchical Bayesian approach allows for model parameters to be

estimated for each participant. The following PTM equation was used:

777
$$c_{\tau_{ij}} = \frac{1}{\beta} \left[\frac{(1 + N_m{}^2)(A_f N_{e_{ij}})^{2\gamma} + N_a{}^2}{(\frac{1}{d'^2{}_i} - N_m^2)} \right]^{\frac{1}{2\gamma}}$$

where the contrast threshold is estimated at a given accuracy criterion, i (either 70.71%

or 79.37%), and external noise level, j. Each participant's response on a given trial (with

a given external noise level and at a given difficulty level) was assumed to be drawn

from a Bernoulli distribution to capture the set of possible outcomes from the 2-

alternative forced-choice response in the visual discrimination task. PTM fitting was

constrained such that the gain parameter β and nonlinear power parameter γ were held

constant to minimize the number of free model parameters while enabling estimation of

all three noise parameters of interest (N_a, N_m, A_f) and the slope of the psychometric

function (η). Values were set at β = 1 and γ = 2.64 for Study 1, and β = 1 and γ = 1.5 for

Study 2, for all individuals, based on estimates generated from the conventional PTM

analysis. Importantly, modifications in the values of the gain parameter β and nonlinear

power parameter γ did not change the results regarding relations between noise

estimates and WM recall variance. For Study 1, based on results from the conventional

PTM analysis which showed a group difference in baseline internal noise and unfiltered

external noise, only the N_a and A_f noise parameters were set to vary across individuals.

Free model parameters were estimated for each individual with the MCMC method for

sampling from posterior probability distributions. A given individual's model parameters

were constrained by hierarchical priors in that parameters were assumed to be drawn

from a group's (SZ or NT) population distribution with their own means and SDs. Priors

for the two groups were set to broad uniform distributions.

Statistics

Repeated measures ANOVAs were conducted to determine changes in recall variance

with increasing WM set size and to probe group differences (Study1) and effects of

tDCS locations (Study 2). To assess the relationship between the sensory noise

estimates and WM variance, multiple linear regressions were completed within each

group. Nested F-tests were used to compare different models of PTM with varying

numbers of free parameters. Statistical analyses were performed at the significance

level of .05.

COMPUTATIONAL MODELING OF WORKING MEMORY PRECISION

Model

The model consists of two stages: encoding of stimulus orientations by a population

code and maintenance of the encoded representation via diffusion during a delay.

Subject-specific sensory noise parameters estimated from the PTM were used to

modulate encoding tuning, post-decoding noise, and memory drift.

Encoding stage. On each trial, a stimulus orientation θ is encoded by a population of M

orientation-tuned neurons with preferred orientations $\{\phi_i\}_{i=1}^{M}$. The mean response of a

neuron i is given by a von Mises tuning function:

$\mu_i(\theta) = r_0 + g_N \exp(\kappa_{enc} \cos(\theta - \phi_i))$,

where r_0 is the baseline firing rate, $g_N = g_0/N$ is the gain scaled by set size N, and

κ_{enc} is the orientation tuning. Unfiltered external noise n_{ext} estimated from the PTM

modulates encoding precision through setting the sharpness of the tuning curves:

$\kappa_{enc} = \dfrac{\kappa_0}{1+\gamma n_{ext}}$,

where κ_0 is the baseline tuning sharpness, and γ controls the influence of the unfiltered

external noise. Neural responses included Poisson-like variability, approximated as

Gaussian noise with variance proportional to the mean response:

$r_i = \mu_i(\theta) + \epsilon_i, \epsilon_i \sim Normal(0, \sigma_i^2), \sigma_i^2 = \alpha\mu_i(\theta)$,

where α controls the strength of the variability. Following orientation decoding, baseline

internal noise was added as a form of additive circular Gaussian noise applied to the

decoded orientation:

$\hat{\theta}_{post} = \hat{\theta}_{enc} + \eta, \; \eta \sim WrappedNormal(0, n_{int}^2),$

where n_{int} is mapped from the PTM baseline internal noise estimate.

Maintenance stage. During the delay period of duration T, the memory representation

undergoes diffusion on the circular orientation space. Memory noise is modeled as a

Gaussian random walk with variance increasing linearly over time:

$\hat{\theta}_{mem} = \hat{\theta}_{post} + \xi, \; \xi \sim WrappedNormal(0, \sigma_{diff}^2), \; \sigma_{diff}^2 = 2D_N T.$

The diffusion constant D_N depends on set size and subject-specific sensory noise:

$D_N = D_0(1 + \beta_{set}(N-1))(1 + \beta_{noise}z_\sigma),$

Where D_0 is the baseline diffusion rate, β_{set} controls the effect of set size, and z_σ is the

dominant internal noise defined as the larger of the z-scored baseline internal noise and

unfiltered external noise.

Behavioral error and precision. Memory error was defined as the circular difference

between the remembered and true orientations. Memory precision was quantified by the

concentration of the error distribution across trials, converted to an equivalent von Mises

concentration parameter. WM variance was reported as the inverse of this

concentration (1/κ).

Subject-specific PTM estimates included baseline internal noise and unfiltered external

noise. These parameters were z-scored across subjects and mapped linearly onto

model parameters governing three noise stages: (i) orientation tuning during encoding,

(ii) post-decoding noise, and (iii) diffusion strength during maintenance. Mapping

coefficients were calibrated by fitting the model to match the mean empirical WM

variance across set sizes. All parameters unrelated to sensory noise or diffusion were

held constant across simulations and serve only to define the scale of the population

code.

Analyses

<u>Relationship between predicted vs empirical WM variance</u>

For each subject and model variant, we simulated 1,000 trials per condition to predict
WM precision using a full model in which PTM-specific sensory noise affected WM
precision at all three noise stages. Model predictions were generated separately for
each subject and set size.

To assess whether the model captured individual differences in WM performance, we
related empirical and model-predicted WM variance using linear mixed-effects models.
Model-predicted WM variance was regressed on empirical WM variance while
accounting for repeated measurements across subjects and set sizes via random
intercepts for subject and set size. Models were fit separately for NT and SZ.

<u>Permutation test of individual specificity</u>

To test whether individual-specific sensory noise estimates were critical for model
performance, we conducted a permutation analysis in which PTM parameters were
randomly shuffled across individuals within each group. For each permutation, model
predictions were recomputed and related to empirical WM variance using the same
mixed-effects framework. Model goodness of fit was quantified as the summed partial
R^2 of the empirical predictor across groups, derived from the mixed-effects models. The
observed goodness-of-fit score was compared against the null distribution to obtain a
permutation p-value.

<u>Bootstrap ablation analysis</u>

To determine which noise stage contributed most to inter-individual variability in WM
precision, we performed a bootstrap analysis on the reduced model variants. In the
reduced models, sensory noise was allowed to affect only one stage at a time, with
other noise parameters fixed to group-median values. In each bootstrap replicate (1,000
iterations), subjects were resampled with replacement within group, and empirical and
predicted WM variance were pooled across set sizes. Overall model goodness of fit was
quantified as the Fisher z–aggregated Pearson correlation between empirical and

predicted WM variance. Confidence intervals were obtained from the bootstrap

distribution, and differences in correlation relative to the full model (Δr) were computed

to assess similarity in explanatory power.

<u>tDCS analysis</u>

To examine whether the model could account for changes in WM performance following

tDCS in SZ participants, we focused on a reduced model in which baseline internal

noise modulated diffusion during the maintenance stage. WM variance was simulated

for sham and stimulation conditions, and condition differences (stimulation − sham)

were computed for both empirical and predicted WM variance, averaged across set

sizes. Pearson correlations were used to assess the relationship between changes in

predicted and empirical WM variance for occipito-parietal and frontal stimulation sites.

DATA AVAILABILITY

The data generated and/or analyzed during the current study are not publicly available

due to participant confidentiality concerns. Although the data do not contain direct

identifiers, they include information that, in combination (e.g., location, patient group),

may pose a risk of re-identification. De-identified data will be made available from the

corresponding authors on request, subject to institutional review and approval.

CODE AVAILABILITY

Code to run experiments and analyze data (MATLAB and Python) are available on

Github repository: https://github.com/gt-core-lab/wm-noise-sz.

ACKNOWLEDGMENTS

Supported by the NIH (R00-EY034546 to WJP; T32-EY007135, P30-EY08126 to GFW;

R01-MH110378 to GFW and SP, R01-MH073028, R01-MH128967 to SP), Smithgall

Watts Early Career Award to WJP, Brain & Behavior Research Foundation (NARSAD

Independent Investigator Grant to DT; Distinguished Investigator Grant to SP), NSF

(BCS-2147064 to GFW), the E. Bronson Ingram Family (to GFW), American

Psychological Foundation (Alexander Gralnick Research Investigator Award to SP), and

Gertrude Conaway Vanderbilt Endowment (to SP).

AUTHOR CONTRIBUTIONS

Conceptualization: WJP, MI, GFW, DT, and SP; Methodology: WJP, MI, GFW, DT, and

SP; Software: WJP, and MI; Formal analysis: WJP, GH, and MI; Investigation: WJP,

and MI.; Writing—Original draft: MI and WJP; Writing—Review & Editing: WJP, GFW,

DT, and SP; Funding acquisition: GFW, DT, and SP.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Postle, B. R. Cognitive neuroscience of visual working memory. in *Working memory:*

State of the science 333–357 (2021).

2. Brady, T. F., Konkle, T. & Alvarez, G. A. A review of visual memory capacity:

Beyond individual items and toward structured representations. *Journal of Vision* **11**,

4–4 (2011).

3. Luck, S. J. Visual short-term memory. in *Visual memory* (2008).

4. Ma, W. J., Husain, M. & Bays, P. M. Changing concepts of working memory. *Nature*

Neuroscience **17**, 347–356 (2014).

5. Barnes-Scheufler, C. V. *et al.* Transdiagnostic comparison of visual working memory

capacity in bipolar disorder and schizophrenia. *Int J Bipolar Disord* **9**, 12 (2021).

6. Kensinger, E. A., Shearer, D. K., Locascio, J. J., Growdon, J. H. & Corkin, S.

Working memory in mild Alzheimer's disease and early Parkinson's disease.

Neuropsychology **17**, 230–239 (2003).

7. Moran, T. P. Anxiety and Working Memory Capacity: A Meta-Analysis and Narrative

Review. (2016).

8. Park, S. & Gooding, D. C. Working memory impairment as an endophenotypic

marker of a schizophrenia diathesis. *Schizophrenia Research: Cognition* **1**, 127–136

(2014).

9. Park, S. & Holzman, P. S. Schizophrenics show spatial working memory deficits.

Archives of General Psychiatry **49**, 975–982 (1992).

10. Xie, W., Campbell, S. & Zhang, W. Working memory capacity predicts individual

differences in social-distancing compliance during the COVID-19 pandemic in the

United States. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 17667–17674 (2020).

11. Eriksson, J., Vogel, E. K., Lansner, A., Bergström, F. & Nyberg, L. Neurocognitive

Architecture of Working Memory. *Neuron* **88**, 33–46 (2015).

12. Hakim, N., Adam, K. C. S., Gunseli, E., Awh, E. & Vogel, E. K. Dissecting the Neural

Focus of Attention Reveals Distinct Processes for Spatial Attention and Object-

Based Storage in Visual Working Memory. *Psychological Science* **30**, 526–540

(2019).

13. Lin, H.-Y. & Oberauer, K. An interference model for visual working memory:

Applications to the change detection task. *Cognitive Psychology* **133**, 101463

(2022).

14. Buschman, T. J., Siegel, M., Roy, J. E. & Miller, E. K. Neural substrates of cognitive

capacity limitations. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 11252–11255 (2011).

15. Gazzaley, A. & Nobre, A. C. Top-down modulation: bridging selective attention and

working memory. *Trends in Cognitive Sciences* **16**, 129–135 (2012).

16. Goldman-Rakic, P. S. & Selemon, L. D. Functional and Anatomical Aspects of

Prefrontal Pathology in Schizophrenia. *Schizophrenia Bulletin* **23**, 437–458 (1997).

17. Scimeca, J. M., Kiyonaga, A. & D'Esposito, M. Reaffirming the Sensory Recruitment

Account of Working Memory. *Trends in Cognitive Sciences* **22**, 190–192 (2018).

18. Xu, Y. Reevaluating the Sensory Account of Visual Working Memory Storage.

Trends in Cognitive Sciences **21**, 794–815 (2017).

19. Zhao, Y.-J., Kay, K. N., Tian, Y. & Ku, Y. Sensory Recruitment Revisited: Ipsilateral

V1 Involved in Visual Working Memory. *Cerebral Cortex* **32**, 1470–1479 (2022).

20. Luck, S. J. & Vogel, E. K. Visual working memory capacity: From psychophysics and

neurobiology to individual differences. *Trends in Cognitive Sciences* **17**, 391–400

(2013).

21. Bays, P. M. Noise in Neural Populations Accounts for Errors in Working Memory.

Journal of Neuroscience **34**, 3632–3645 (2014).

22. Dosher, B. A. & Lu, Z. L. Perceptual learning reflects external noise filtering and

internal noise reduction through channel reweighting. *Proceedings of the National*

Academy of Sciences **95**, 13988–13993 (1998).

23. Faisal, A. A., Selen, L. P. J. & Wolpert, D. Noise in the nervous system. *Nature*

Reviews Neuroscience **9**, 292–303 (2008).

24. Shannon, C. E. A Mathematical Theory of Communication. *Bell System Technical*

Journal **27**, 379–423 (1948).

25. Harrison, S. A. & Tong, F. Decoding reveals the contents of visual working memory

in early visual areas. *Nature* **458**, 632–635 (2009).

26. Pasternak, T. & Greenlee, M. W. Working memory in primate sensory systems.

Nature Reviews Neuroscience **6**, 97–107 (2005).

27. Vogel, E. K. & Machizawa, M. G. Neural activity predicts individual differences in

visual working memory capacity. *Nature* **428**, 748–751 (2004).

28. Castner, StacyA., Goldman-Rakic, PatriciaS. & Williams, GrahamV. Animal models

of working memory: insights for targeting cognitive dysfunction in schizophrenia.

Psychopharmacology **174**, (2004).

29. Barch, D. M. & Ceaser, A. Cognition in schizophrenia: core psychological and neural

mechanisms. *Trends in Cognitive Sciences* **16**, 27–34 (2012).

30. Javitt, D. C. When Doors of Perception Close: Bottom-up Models of Disrupted

Cognition in Schizophrenia. *Annu. Rev. Clin. Psychol.* **5**, 249–275 (2009).

31. Kim, J., Park, S. & Blake, R. Perception of biological motion in Schizophrenia and

healthy individuals: A behavioral and fMRI study. *PLoS ONE* **6**, (2011).

32. Tadin, D. *et al.* Weakened center-surround interactions in visual motion processing

in schizophrenia. *Journal of Neuroscience* **26**, 11403–11412 (2006).

33. Yang, E. *et al.* Visual context processing in schizophrenia. *Clinical psychological*

science **1**, 5–15 (2013).

34. Abbasi, S., Wolff, A., Çatal, Y. & Northoff, G. Increased noise relates to abnormal

excitation-inhibition balance in schizophrenia: a combined empirical and

computational study. *Cerebral Cortex* **33**, 10477–10491 (2023).

35. Chen, Y., Norton, D. & McBain, R. Effects of domain-specific noise on visual motion

processing in schizophrenia. *PLoS ONE* **9**, (2014).

36. Christensen, B. K., Spencer, J. M. Y., King, J. P., Sekuler, A. B. & Bennett, P. J.

Noise as a mechanism of anomalous face processing among persons with

Schizophrenia. *Front. Psychol.* **4**, (2013).

37. Bays, P. M. & Husain, M. Dynamic Shifts of Limited Working Memory Resources in

Human Vision. *Science* **321**, 851–854 (2008).

38. Wilken, P. & Ma, W. J. A detection theory account of change detection. *Journal of*

vision **4**, 1120–35 (2004).

39. Starc, M. *et al.* Schizophrenia is associated with a pattern of spatial working memory

deficits consistent with cortical disinhibition. *Schizophrenia Research* **181**, 107–116

(2017).

40. Pelli, D. G. & Farell, B. Why use noise? *Journal of the Optical Society of America. A,*

Optics, image science, and vision **16**, 647–653 (1999).

41. Dosher, B. A. & Lu, Z. Noise exclusion in spatial attention. *Psychological Science*

11, 139–146 (2000).

42. Manning, C., Dakin, S. C., Tibber, M. S. & Pellicano, E. Averaging, not internal

noise, limits the development of coherent motion processing. *Developmental*

cognitive neuroscience **10**, 1–13 (2014).

43. Park, W. J., Schauder, K. B., Zhang, R., Bennetto, L. & Tadin, D. High internal noise

and poor external noise filtering characterize perception in autism spectrum

disorder. *Scientific Reports* 1–12 (2017).

44. Nitsche, M. A. & Paulus, W. Excitability changes induced in the human motor cortex

by weak transcranial direct current stimulation. *The Journal of Physiology* **527**, 633–

639 (2000).

45. Nitsche, M. A. & Paulus, W. Sustained excitability elevations induced by transcranial

DC motor cortex stimulation in humans. *Neurology* **57**, 1899–1901 (2001).

46. Wagner, T., Valero-Cabre, A. & Pascual-Leone, A. Noninvasive Human Brain

Stimulation. *Annu. Rev. Biomed. Eng.* **9**, 527–565 (2007).

47. Reinhart, R. M. G., Xiao, W., McClenahan, L. J. & Woodman, G. F. Electrical

Stimulation of Visual Cortex Can Immediately Improve Spatial Vision. *Current*

Biology **26**, 1867–1872 (2016).

48. Zhang, W. & Luck, S. J. Sudden Death and Gradual Decay in Visual Working

Memory. *Psychol Sci* **20**, 423–428 (2009).

49. Schneegans, S. & Bays, P. M. Drift in Neural Population Activity Causes Working

Memory to Deteriorate Over Time. *J. Neurosci.* **38**, 4859–4869 (2018).

50. Lu, Z.-L. & Dosher, B. A. Perceptual learning retunes the perceptual template in

foveal orientation identification. *Journal of vision* **4**, 44–56 (2004).

51. Mayer, J. S. & Park, S. Working memory encoding and false memory in

schizophrenia and bipolar disorder in a spatial delayed response task. *Journal of*

Abnormal Psychology **121**, 784–794 (2012).

52. Thakkar, K. N. & Park, S. Impaired Passive Maintenance and Spared Manipulation

of Internal Representations in Patients With Schizophrenia. *Schizophrenia Bulletin*

38, 787–795 (2012).

53. D'Esposito, M. From cognitive to neural models of working memory. *Phil. Trans. R.*

Soc. B **362**, 761–772 (2007).

54. Tsubomi, H., Fukuda, K., Watanabe, K. & Vogel, E. K. Neural Limits to Representing

Objects Still within View. *J. Neurosci.* **33**, 8257–8263 (2013).

55. Adam, K. C. S., Mance, I., Fukuda, K. & Vogel, E. K. The Contribution of Attentional

Lapses to Individual Differences in Visual Working Memory Capacity. *Journal of*

Cognitive Neuroscience **27**, 1601–1616 (2015).

56. Wang, J.-K. *et al.* Enhancing visual working memory in schizophrenia: effects of

frontoparietal theta tACS in low-performing patients. *Schizophr* **10**, 97 (2024).

57. Rokem, A. *et al.* Broader Visual Orientation Tuning in Patients with Schizophrenia.

Frontiers in Human Neuroscience **5**, 127 (2011).

58. Boroda, E., Sponheim, S. R., Fiecas, M. & Lim, K. O. Transcranial direct current

stimulation (tDCS) elicits stimulus-specific enhancement of cortical plasticity.

NeuroImage **211**, 116598 (2020).

59. Reinhart, R. M. G., Cosman, J. D., Fukuda, K. & Woodman, G. F. Using transcranial

direct-current stimulation (tDCS) to understand cognitive processing. *Atten Percept*

Psychophys **79**, 3–23 (2017).

60. Antal, A., Kincses, T. Z., Nitsche, M. A., Bartfai, O. & Paulus, W. Excitability

Changes Induced in the Human Primary Visual Cortex by Transcranial Direct

Current Stimulation: Direct Electrophysiological Evidence. *Invest. Ophthalmol. Vis.*

Sci. **45**, 702 (2004).

61. Uhlhaas, P. J., Phillips, W. A. & Silverstein, S. M. The course and clinical correlates

of dysfunctions in visual perceptual organization in schizophrenia during the

remission of psychotic symptoms. *Schizophrenia Research* **75**, 183–192 (2005).

62. Dinstein, I. *et al.* Unreliable evoked responses in autism. *Neuron* **75**, 981–991

(2012).

63. Habib, A., Harris, L., Pollick, F. & Melville, C. A meta-analysis of working memory in

individuals with autism spectrum disorders. *PLoS ONE* **14**, e0216198 (2019).

64. Wang, Y. *et al.* A Meta-Analysis of Working Memory Impairments in Autism

Spectrum Disorders. *Neuropsychol Rev* **27**, 46–61 (2017).

65. Calvin, O. L. & Redish, A. D. Global disruption in excitation-inhibition balance can

cause localized network dysfunction and Schizophrenia-like context-integration

deficits. *PLoS Comput Biol* **17**, e1008985 (2021).

66. Gao, R. & Penzes, P. Common Mechanisms of Excitatory and Inhibitory Imbalance

in Schizophrenia and Autism Spectrum Disorders. *CMM* **15**, 146–167 (2015).

67. Rubenstein, J. L. R. & Merzenich, M. M. Model of autism: Increased ratio of

excitation/inhibition in key neural systems. *Genes, brain, and behavior* **2**, 255–267

(2003).

68. Fried, R. *et al.* Clinical correlates of working memory deficits in youth with and

without ADHD: A controlled study. *Journal of Clinical and Experimental*

Neuropsychology **38**, 487–496 (2016).

69. Jeffries, S. & Everatt, J. Working memory: Its role in dyslexia and other specific

learning difficulties. *Dyslexia* **10**, 196–214 (2004).

70. Kessels, R. P. C., Molleman, P. W. & Oosterman, J. M. Assessment of working-

memory deficits in patients with mild cognitive impairment and Alzheimer's dementia

using Wechsler's Working Memory Index. *Aging Clin Exp Res* **23**, 487–490 (2011).

71. Lee, J. & Park, S. Working Memory Impairments in Schizophrenia: A Meta-Analysis.

Journal of Abnormal Psychology **114**, 599–611 (2005).

72. First, M. B., Spitzer, R. L., Gibbon, M. & Williams, J. B. W. First, M. B., Spitzer, R. L.,

Gibbon, M., & Williams, J. B. W. (2002). Structured clinical interview for DSM-IV-TR

axis I disorders, research version, patient edition. (SCID-I/P). *Biometrics Research*

(2002).

73. Nelson, H. E. National Adult Reading Test (NART): For the Assessment of

Premorbid Intelligence in Patients with Dementia: Test Manual. (1982).

74. Andreasen, N. C. Scale for the assessment of negative symptoms (SANS). *Iowa*

City: University of Iowa. (1983).

75. Andreasen, N. C. Scale for the assessment of positive symptoms (SAPS). *University*

of Iowa: Iowa City. (1984).

76. Brainard, D. H. The Psychophysics Toolbox. *Spatial Vision* **10**, 433–436 (1997).

77. Pelli, D. G. The VideoToolbox software for visual psychophysics: Transforming

numbers into movies. *Spatial Vision* **10**, 437–442 (1997).

78. Foss-Feig, J., Tadin, D., Schauder, K. B. & Cascio, C. J. A substantial and

unexpected enhancement of motion perception in autism. *Journal of Neuroscience*

33, 8243–8249 (2013).

79. Gold, J. M. *et al.* Reduced Capacity but Spared Precision and Maintenance of

Working Memory Representations in Schizophrenia. *Arch Gen Psychiatry* **67**, 570

(2010).

80. Vul, E., Bergsma, J. & MacLeod, D. I. Functional Adaptive Sequential Testing.

Seeing and Perceiving **23**, 483–515 (2010).

81. Reinhart, R. M. G. & Woodman, G. F. Causal Control of Medial–Frontal Cortex

Governs Electrophysiological and Behavioral Indices of Performance Monitoring and

Learning. *J. Neurosci.* **34**, 4214–4227 (2014).

82. Jung, Y.-J., Kim, J.-H. & Im, C.-H. COMETS: A MATLAB toolbox for simulating local

electric fields generated by transcranial direct current stimulation (tDCS). *Biomed.*

Eng. Lett. **3**, 39–46 (2013).

83. Monte-Silva, K. *et al.* Induction of Late LTP-Like Plasticity in the Human Motor

Cortex by Repeated Non-Invasive Brain Stimulation. *Brain Stimulation* **6**, 424–432

(2013).

1122

1123