TDP-43 dysfunction leads to impaired proteostasis and predisposes mice to worse neurological outcomes after brain injury

Melissa S. Rotunno [1,2], Megan Fowler-Magaw [1,3,#], Jianjun Zhong [1,4,#], Kennedy O'Hara [1,3], Elenore A. Wiggin [1,5], Debra Cameron [1], Karly Stallworth [1,2], James Bouley [1], Holly McEachern [6,7], Mina N. Anadolu [8], Jeffrey A. Nickerson [9], Andrew R. Tapper [10], Susanna Molas [10], Francesca Massi [2,11], Nils Henninger [1,3,12], Oliver D. King [1,3], Daryl A. Bosco [1,2,3,5,11,13]*

[1]Department of Neurology, University of Massachusetts Chan Medical School, Worcester, Massachusetts, MA, 01605

[2]Department of Biochemistry and Molecular Biotechnology, University of Massachusetts Chan Medical School, Worcester, MA, 01605

[3]Neuroscience Program, Morningside Graduate School of Biomedical Sciences, University of Massachusetts Chan Medical School, Worcester, MA, 01605

[4]Department of Neurosurgery, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China, 400016.

[5]Interdisciplinary Graduate Program, Morningside Graduate School of Biomedical Sciences, University of Massachusetts Chan Medical School, Worcester, MA, 01605

[6]RNA Therapeutics Institute, UMass Chan Medical School, Worcester, MA, 01605

[7]RNA Therapeutics and Biology Program, Morningside Graduate School of Biomedical Sciences, University of Massachusetts Chan Medical School, Worcester, MA, 01605

[8]Program in Molecular Medicine, University of Massachusetts Chan Medical School, Worcester, MA, 01605

[9]Department of Pediatrics, University of Massachusetts Medical School, Worcester, Worcester, MA, 01605

[10]Department of Neurobiology, Brudnick Neuropsychiatric Research Institute, University of

Massachusetts Chan Medical School, Worcester, MA, 01605

[11]Biochemistry and Molecular Biotechnology Program, Morningside Graduate School of

Biomedical Sciences, University of Massachusetts Chan Medical School, Worcester, MA, 01605

[12]Department of Psychiatry, University of Massachusetts Chan Medical School, Worcester, MA

[13]Lead Contact

[#]These authors contributed equally

*To whom correspondence should be addressed: Daryl A. Bosco: Department of Neurology,

University of Massachusetts Chan Medical School, Worcester, MA 01605;

Daryl.Bosco@umassmed.edu; Tel. (774) 455-3745; Fax. (508) 856-6750

ABSTRACT

Background: Pathological TAR DNA-binding protein 43 (TDP-43) dysfunction is associated with

multiple neurodegenerative disorders. However, the mechanistic link between TDP-43

dysfunction and neurodegeneration is poorly understood and likely involves a combination of

genetic and environmental risk factors. A major risk factor for neurodegenerative disease is

exposure to traumatic brain injury (TBI). Here, we investigated the synergistic interplay between

TDP-43 dysfunction and TBI in a murine model of amyotrophic lateral sclerosis (ALS)/

frontotemporal dementia (FTD).

Methods: A model of TDP-43 dysfunction caused by a knock-in Q331K mutation in *Tardbp* was

combined with a mild model of TBI. Control conditions included both WT mice and mice with sham

surgery. Animals were evaluated for behavioral deficits at timepoints pre- and post-surgery.

Additionally, post-mortem brain tissues were examined using RNA sequencing and mass

spectrometry-based quantitative proteomics together with histological and biochemical analyses.

Results: Expression of dysfunctional TDP-43 *in vivo* caused deficits in multiple branches of the

proteostasis network, including protein folding, protein synthesis, and protein turnover. Examples

include mis-expression of chaperones and genes within the ubiquitin-proteosome pathway in

mutant TDP-43 versus WT mice. Further, mutant TDP-43 expression correlated with reduced

thermostability of proteins associated with the ribosome and the chaperonin containing TCP-1

complex. In response to TBI, mutant TDP-43 mice exhibited significantly worse neurological

outcomes relative to WT animals. Heightened neurological deficits in mutant TDP-43 mice

following TBI coincided with a robust upregulation of proteostasis- and stress-related genes at

the transcript level. However, this upregulation was not detected at the protein level.

Conclusions: Our data demonstrate that expression of dysfunctional TDP-43 leads to deficits

within the proteostasis network in vivo at baseline. Despite an upregulation of proteostasis-related

genes at the transcript level in mutant TDP-43 mice after TBI, mutant TDP-43 mice exhibit an

impaired response to, and recovery from, brain trauma relative to their WT counterparts.

Restoring proteostasis is expected to protect against the detrimental effects of TDP-43

dysfunction, especially under stress conditions that promote neurodegenerative disease.

Keywords: TDP-43, traumatic brain injury, proteostasis, ribosome, Proteome Integral Solubility

Alteration (PISA), multistep hypothesis, amyotrophic lateral sclerosis, frontotemporal dementia,

mass spectrometry, transcriptomics.

BACKGROUND

TAR DNA-binding protein 43 (TDP-43) is an essential RNA/DNA-binding protein that performs

critical functions in the context of RNA splicing, transport, and expression.(1) TDP-43 shuttles

between the nucleus and cytoplasm while fulfilling these functions, although the protein is

predominately expressed in the nucleus under physiological conditions. A significant cytoplasmic

shift in TDP-43 localization has been observed in cultured cells exposed to various external stimuli

or "stressors" that inhibit proteasomal and transcriptional activity, or that induce hyperosmolarity,

hyperexcitability or excitotoxicity.(2-6) While the mechanisms underlying stress-induced

cytoplasmic accumulation of TDP-43 have not been fully elucidated, recent reports have shown

that both RNA binding capacity and post-translational modification of TDP-43 affect TDP-43

subcellular localization.(3, 7-10) TDP-43 has also been identified as a component of cytoplasmic

condensates called stress granules, which form in response to acute cellular stress.(11-13)

Autosomal dominant mutations in TDP-43 that cause familial amyotrophic lateral sclerosis (ALS)

alter the properties of stress granules as well as other TDP-43-associated ribonucleoprotein

(RNP) granules,(10-12, 14-16) possibly through altered phase separation behavior of mutant

TDP-43 with itself, other RNA-binding proteins and/or RNA.(10, 17-20) Collectively, these

observations indicate that subcellular TDP-43 localization can serve as a proxy for cellular stress.

Further, the effects of TDP-43 on stress response pathways can be perturbed by disease-causing

mutations.

In addition to TDP-43 mutations in familial ALS, TDP-43 has emerged as a hallmark pathological

feature across multiple neurodegenerative disorders termed TDP-43 proteinopathies,(21)

including ALS, frontotemporal lobar degeneration (FTLD),(22, 23) Alzheimer's disease (AD),(24)

limbic-predominant age-related TDP-43 encephalopathy (LATE)(25) and chronic traumatic

encephalopathy (CTE).(26). Cells with cytoplasmic accumulation and/or nuclear depletion of

TDP-43 have been shown to exhibit splicing defects that are consistent with TDP-43 loss-of-

function in human post-mortem CNS tissues from ALS/FTLD and AD cases.(4, 27-31) Most cases

of human TDP-43 proteinopathy occur sporadically with poorly defined etiologies, and likely

develop through a combination of genetic susceptibility and environmental risk factors.(32, 33)

Indeed, emerging evidence supports that ALS pathogenesis occurs through a multistep process,

even in cases with penetrant genetic mutations in *TARDBP* encoding TDP-43.(34) The multistep

hypothesis purports that individual ALS-linked mutations cannot fully account for the disease

process, which must include other factors such as additional genetic mutations, environmental

exposures and/or physical stress.(34, 35)

Traumatic brain injury (TBI) is a form of physical stress that also represents a major risk factor for

developing neurodegenerative diseases, including those with TDP-43 pathology.(36) Cytoplasmic

shifts in TDP-43 localization have been reported in laboratory animal models of TBI.(13, 37-41)

Further, recurrent TBI leads to CTE with cytoplasmic TDP-43 pathology in humans.(26, 42) Here,

we sought to investigate the relationship between stress and TDP-43 dysfunction in the context

of a multistep model of neurodegenerative disease. To this end, we combined a murine model of

TDP-43 dysfunction caused by a genetic knock-in mutation in *Tardbp* together with a mild model

of TBI.(43, 44) Our results revealed that mutant TDP-43 mice exhibit significantly worse

neurological outcomes during the acute phase post-TBI relative to their wild-type (WT)

counterparts. Corresponding transcriptomics and RT-PCR analyses revealed synergistic effects

of TDP-43 mutation and TBI on gene expression and processing. For example, a significant

upregulation of proteostasis- and stress-related genes was detected specifically in brain tissue

from mutant TDP-43 mice after TBI. Many of the pathways that became upregulated in mutant

TDP-43 mice following TBI were also highlighted in a recent transcriptomics analysis of post-

mortem ALS and FTLD-TDP CNS tissues.(45, 46) While this TBI model caused a cytoplasmic

shift in TDP-43 localization, this change in TDP-43 localization did not appear to correlate with

mis-splicing events caused by TDP-43 loss-of-function. Rather, we found that mutant TDP-43

expression correlated with reduced thermostability of cytoplasmic ribosomal proteins. In fact,

diminished thermostability of ribosomal proteins correlated with mutant TDP-43 expression,

irrespective of whether mutant TDP-43 mice had been exposed to brain trauma or not. Notably,

TDP-43 has been shown to associate with ribosomes and to modulate both global and local

translation.(47-50)

The outcomes of this study are consistent with a model in which expression of mutant TDP-43

leads to deficits throughout the proteostasis network. Upon CNS challenge, compensatory

mechanisms appear to become activated in mutant TDP-43 mice, yet these mice fail to mount

the physiological response to brain injury exhibited by their WT counterparts. Therefore, our data

demonstrate that expression of dysfunctional TDP-43 leads to impaired response to, and recovery

from, brain trauma and provide mechanistic insight into TBI as a risk factor for TDP-43

proteinopathies. Restoring proteostasis may therefore protect against the effects of TDP-43

dysfunction, especially under conditions of CNS challenge that promote neurodegenerative

disease.

METHODS

Animals

All animal related protocols were conducted in accordance with the Guide for the Care and Use

of Laboratory Animals published by the National Research Council (US) Committee. The

protocols were approved by the Institutional Animal Care and Use Committee (IACUC) of the

University of Massachusetts Chan Medical School. Mice with the human-equivalent TDP-43

Q331K mutation on a pure C57BL/6J (WT) background were generated and characterized in a

previous study.(44) All experiments herein were performed with aged-matched, male mice with

either the homozygous TDP-43 Q331K or TDP-43 WT genotype. Mice were generated using

standard breeding approaches or via *in vitro* fertilization at the UMass Chan Medical School

Transgenic Animal Modeling Core (TAMC; Worcester, MA). All mice were maintained under

standard conditions with no more than 5 mice per cage, on a standard 12h light/dark cycle (white

light ON at 7:00am and OFF at 7pm), and were provided *ad libitum* access to food and water.

Traumatic brain injury (TBI) surgery

Mice at 16-weeks of age were exposed to TBI or sham surgery. TBI surgeries were based on a

previously published protocol.(43) Details of the TBI surgery used herein were published by us

elsewhere.(51) Briefly, mice were anesthetized with 3% isoflurane (Pivetal, NDC 46066-755-03)

in carrier gas at 1.5 L/min flow. Prior to surgery, buprenorphine (PAR Pharmaceutical, NDC

42023-179-05) was administered subcutaneously at 1 mg/kg. Hair was removed from the top of

the head and the scalp skin was disinfected with 75% ethanol and betadine. A ~1.2-1.5 cm

incision was made along the midline starting approximately 3 mm posterior from the eyes to

expose the right parietal skull. The skin over the left and right hemispheres of the skull was

excised. The periosteum was removed and the exposed skull was positioned on a cushion

beneath the weight-drop. A transducer rod tip was aligned to the skull-top location: 2.5 mm

posterior to Bregma and 2 mm lateral to the midline. From a height of 15 cm, a 50-gram

weight was released that came into contact with a transducer, which in turn delivered an impact

onto the mouse's intact skull (see diagram in **Fig. 1A**). After the impact, mice were placed onto a

heating blanket in a supine position for the time needed to wake from the anesthesia (i.e., time

required to right themselves to a prone position), which took on the order of minutes for each

mouse. Mice were then returned to the anesthesia chamber in preparation for a subsequent

impact; this process was repeated a total of ten times within a single surgical session. As the

impacts were delivered within the same surgical session and not spaced by days or weeks, this

model is not considered a repetitive TBI model that we described previously.(40, 52) After the

surgery was completed, cefazolin (HIKMA Pharmaceutical, NDC 0143-9924-90) was

administered intramuscularly at 500 mg/kg and meloxicam (Norbrook, NDC 55529-040-10) was

administered subcutaneously at 5 mg/kg. Mice were placed on a heating blanket and monitored

until they were ambulatory (approximately 15 minutes), and then transferred to their home cage.

Buprenorphine was administered as described above at 8 h, 16 h and 24 h after the first dose.

For the sham surgery, mice underwent the same procedure as the TBI mice but without the

weight-drop impact.

Neurological Severity Score (NSS) Test

The NSS test was performed at 12 time points: pre-surgery and 2h, 24h, 48h, 1 week, 2 weeks,

3 weeks, 4 weeks, 8 weeks, 12 weeks, 16 weeks and 20 weeks post-surgery, as previously

described in detail.(43, 53) Body weight was measured prior to every NSS testing session. Briefly,

the NSS consists of 10 individual tasks designed to assess motor function, alertness and

physiological behavior. Each task was completed regardless of whether the mouse passed or

failed previous tasks. For each task, mice were assigned a score of 1 to indicate failure of the

task or 0 if the task was completed, resulting scores ranging between 0 (no deficit) and 10

(maximum deficit). The NSS test was performed by a lab technician who was blinded to the

genotype and surgery type of the mice. For statistical analyses, between-group comparisons of

the NSS over repeated measurements (time) were conducted using longitudinal mixed models.

Time was treated as a categorical variable. The models included group (each group representing

both a genotype and surgery type) and time as fixed covariates, as well as the two-way interaction.

When the main effects were significant, pairwise comparisons were conducted to identify

significance by group and time point. A two-sided $p < 0.05$ was considered statistically significant.

Statistical analyses were performed using SPSS Statistics Version 29 (IBM-Armonk, NY).

Cued and Contextual Fear Conditioning (CCFC) Test

Animals were generated through two in vitro fertilization procedures at the Transgenic Animal

Modeling Core (TAMC) at UMass Chan Medical School using heterozygous Q331K male and

female mice. Homozygous Q331K and WT mice were randomly assigned to group 1 or group 2,

and both groups were subjected to the CCFC test before surgery. The CCFC test was conducted

at the UMass Chan Mouse Behavioral Core facility using the EthoVision XT system (Noldus,

BOM-EV-0001). Mice were placed in a Ugo Basile Fear Conditioning Chamber (Noldus, BOM-

XUGO-F101) made of optical transparent plastic 26 cm long, 26 cm wide and 38 cm tall. The top

of the chamber was equipped with a monochrome GigE camera (Noldus, BOM-XCAM-GIGE)

used to track animal movement in real-time during the testing session. The bottom of the chamber

contained electricity-conductive metal strips to deliver foot shocks. The chamber was nested in a cabinet equipped with a speaker to deliver white noise and high-frequency tones per the experimental design detailed below. The cabinet wall was insulated to reduce environmental noise and a built-in ventilator allowed for air-exchange. All the inputs, including white noise, tone, and foot shock, were controlled with a computer-based operation console (Ugo Basile, version 1.1.8.9). EthoVision XT 15 software was used for animal tracking and data analysis. The live tracking started immediately when the animal entered the chamber, after which there was a 60s delay prior to the start of an input. The full test session was conducted over four consecutive days as follows. Note that the white noise and the tone input were delivered randomly, with random intervals (ranging from 5s to 30s) between these two inputs for all tests on each day.

Day 1:

White noise: 50% intensity, 30s duration, and repeated 5 times.

Tone: 1000 Hz, 50% intensity, 30s duration, and repeated 5 times.

Day 2:

White noise: 50% intensity, 30s duration, and repeated 5 times.

Tone: 1000 Hz, 50% intensity, 30s duration, and repeated 5 times.

Electrical foot shock: 0.75 mA intensity that occurred at the last second of each tone. Each mouse received five foot shocks in total on Day 2 with each shock lasting for 1s.

Day 3:

White noise: 50% intensity, 30s duration, and repeated 5 times.

Tone: 1000 Hz, 50% intensity, 30s duration, and repeated 5 times.

Novel context: To create a novel context, a new scent and visual component were introduced to the chamber as follows. One drop of vanilla extract (novel scent) was applied to a piece of tissue paper, which was placed in the feces tray underneath the chamber. Paper boards with white and black stripes (novel visual) were attached to the outer side of the chamber.

Day 4:

224   The context was reverted as per Days 1 and 2 (i.e., the paper boards were removed and the feces

225   tray cleaned before Day 4) for testing on Day 4.

226   Neither white noise nor tone were implemented.

227   Below is a table summarizing the inputs and contexts for the CCFC test:

	Day 1	Day 2	Day 3	Day 4
White noise	+	+	+	-
Tone	+	+	+	-
Foot shock	-	+	-	-
Novel context	-	-	+	-

For analysis of Days 1-3, the initial 29s of each input was extracted to avoid potential interference

from the last second when the foot shock was delivered. Therefore, the total duration for white

noise and tone inputs is 29s x 5 = 145s. For Day 4, the consecutive 145s after the initial 60s delay

was extracted for the analysis. Mouse movement was detected by the software at the pixel level

for the entire chamber from one video frame to the next. Thresholds based on animal activity

within the chamber were set as follows: immobility state (<0.1% pixel change in the chamber),

movement between immobility and walking (between 0.1% and 0.5%) and walking (>0.5%). The

amount of time animals spent in an immobility state was interpreted as freezing behavior and

plotted for each day (Day 1-4). The statistical analysis was performed as described above for

NSS, where between-group comparisons over repeated measurements (time) were conducted

using longitudinal mixed models. Time was treated as a categorical variable. The models included

group and time as fixed covariates, as well as the two-way interaction. When the main effects

were significant, pairwise comparisons were conducted to identify significance by group and time

point. A two-sided $p<0.05$ was considered statistically significant. Statistical analyses were

performed using SPSS Statistics Version 29 (IBM-Armonk, NY).

RNA isolation

Mouse brain tissue samples (~40 mg and 4 mm of diameter) were collected from the impact site

(**Fig. 1A**) and the corresponding site on the contralateral side using a sterile disposable biopsy

punch (Integra Miltex, 33-32) that excised tissue from the brain surface (i.e., cortex) to the brain

bottom (i.e., hypothalamus). Total RNA was isolated from mouse brain tissues using Qiagen's

RNeasy Lipid Tissue Mini Kit (Cat. No. 74804) with DNase treatment following the manufacturer's

protocols. RNA concentration and absorbance were measured by a Nanodrop device (ND1000).

Samples with A260/A280 between 1.8 to 2.2 were considered suitable for downstream

applications.

Transcriptome analysis by RNA-sequencing (RNAseq)

For the 24h-post TBI timepoint, RNAseq analysis was performed on RNA isolated from brain

tissue as described above for six groups of mice: homozygous Q331K and WT genotypes, each

under naïve, sham, and TBI conditions. For the 7mo-post TBI timepoint, only sham and TBI

conditions for homozygous Q331K and WT genotypes were included. Sham and TBI conditions

are described under TBI surgery. Naïve mice were not subjected to surgery and are therefore

treated a separate control group than sham surgery mice. Between 3-6 mice were included per

group for a total of 46 samples. RNA quality assessment, library preparation and sequencing were

conducted by Novogene Co., LTD (Beijing, China). Poly-T oligo-attached magnetic beads were

used for poly-A capture, random hexamer primers for first strand cDNA synthesis, and dTTP for

second strand cDNA synthesis of non-directional libraries. Libraries were sequenced to a depth

of 50M-125M raw read pairs (2x150 PE) per sample using an Illumina NovaSeq 6000 sequencer.

After trimming adapter sequences and low-quality bases, reads were mapped with HISAT2

(2.0.5)(54) to the mouse mm39 reference genome with Ensembl release 107 gene annotations.

Two samples (one Q331K sham and one Q331K TBI) from the 24h post-TBI timepoint were

excluded from further analyses, as they were strong outliers in Principal Component Analysis

(PCA) of expression values (counts-per-million on $\log_2$ scale with prior.count=1; **Fig. S1A**); see

Additional File 1 for all supplemental figures. These samples were also outliers for several

mapping quality control (QC) metrics, with higher-than-usual fractions of multi-mapping, exonic,

and unspliced reads. Two samples (one Q331K sham and one WT TBI) were also excluded from

the 7mo-post TBI dataset because they were determined to by outliers in the PCA analysis (**Fig.**

S1B).

Differential gene expression analyses were performed in R (4.3.2) using quasi-likelihood methods

in the edgeR package (4.0.12).(55) Data from 24hr-post TBI and 7mo-post TBI experiments were

analyzed separately. The TMM method was used for computing normalized library sizes and

genes with low counts were filtered out in preprocessing. For both datasets, ~25,000 genes were

retained with at least 5 scaled counts in at least 3 samples, where the scaling adjusts for

differences in total library sizes (read counts across all genes) between samples while keeping

the median library size unchanged. The function glmQLFit (with non-default options robust=T

and prior.count=1) was used to fit the expression data with a design matrix that included factors

for injury (INJ, with levels Naïve, Sham, and TBI), genotype (GT, with levels Q331K and WT) and

their interaction (model formula ~0 + GT:INJ). Differential expression was assessed using the

function glmQLFTest for contrasts that included nine simple between-group comparisons

(WT.TBI vs WT.Sham, etc.); main effects of GT (Q331K vs WT) and INJ (TBI vs Sham), though

in both cases excluding the groups with INJ=Naïve; and a test for an interaction between GT and

INJ, again excluding the groups with INJ=Naïve. For the 7mo-post TBI data the naïve mice were

not included, and the model also included an additive factor for IVF batch (with two levels, 162

and 167). Supplemental Table I includes the $\log_2$(fold-change) [LFC], p-value, and FDR for each

gene for each contrast discussed in the results section.

Gene Set Enrichment Analysis (GSEA, 4.3.3) was performed using the Hallmark

(mh.all.v2024.1.Mm.symbols.gmt) and Gene Ontology Biological Process (GOBP,

m5.go.bp.v2024.1.Mm.symbols.gmt) gene set collections from the Molecular Signatures

Database (MSigDB).(56-60) Cutoffs on Normalized Enrichment Score (NES) and adjusted p-

value (FDR) were used to determine significantly altered gene sets (|NES| > 1.5 and FDR<0.05).

NES is a score that reflects the overrepresentation of a gene set at the upper (positive value) or

lower (negative value) end of a ranked gene list, after normalization to account for variation in

gene set size. The gene list was pre-ranked based on directional p-values for the indicated

comparison ($-\log_{10}$(p-value)*sign of $\log_2$(fold change)). A weighted enrichment statistic was used

in the analysis. The 10 top gene sets based on highest |NES| are displayed for all enrichment

analyses. If fewer than 10 are displayed then all gene sets that meet the cutoff criteria are shown.

All gene set hits that meet the cutoff criteria can be found in **Supplemental Table II**.

Quantitative Proteomics

Mice were perfused with PBS prior to tissue isolation for all proteomic analyses. Mouse brain

tissue samples were collected as described above under RNA isolation and cut into ~1 mm

diameter pieces. For whole brain proteomics, the tissue was transferred to a pre-cooled glass

homogenizer on ice with 900 µl of RIPA buffer (Sigma-Aldrich R0278-50ML) containing 1x

protease inhibitor (Roche, 11873580001) and 1x phosphatase inhibitor (Roche, 04906837001).

Complete tissue homogenization was performed with a WHEATON Power Homogenizer and

Overhead Stirrer (catalog number 903475, serial number G082210) by setting the stirring speed

on level 2 for 1 minute. The sample was transferred to a plastic tube together with 300ul of RIPA

buffer that was added to rinse the homogenizer. The sample was left on ice for 30 min and then

centrifuged at 15,899 xg for 10 min at 4°C in a table-top centrifuge (AXYGEN, serial number

LRMPL1-1710018, lot number 04619100). The supernatant was transferred into a fresh tube and

the bicinchoninic acid (BCA) assay (Thermo Scientific, 23223) was used to measure the protein

concentration. For cytoplasmic samples depleted of nuclei and mitochondria for PISA (below),

samples were homogenized in 1.1 mL of KPBS (130 mM KCl, 10 mM KH_2PO_4, pH 7.3). Minced

tissue was homogenized twice on level 2 for 1 min each with 1 min on ice in between.

Homogenate was transferred to a 1.5 ml Eppendorf tube. The glass tube was rinsed with 0.3 ml

of KPBS +PI and added to a 1.5 ml Eppendorf tube. Samples were spun twice at 600 xg at 4°C

for 5 min with supernatant moved to a fresh Eppendorf tube each time to remove the nuclear

fraction and debris. Samples were then centrifuged at 7,000 xg at 4°C for 10 min to remove

mitochondria.

For PISA analysis, cytoplasmic samples depleted of nuclei and mitochondria were diluted in half

to 1 mg/ml with PISA lysis buffer (2 mM MgCl2, 1% CA630, PBS (Corning 21-031-CV)). Lysates

were incubated on ice for 30 min and subdivided into 4 aliquots in PCR strip tubes. One aliquot

was kept on ice to use for the total protein level quantitative proteomics. The remaining three

aliquots were exposed to a single temperature (52°C, 53°C, 54°C) for 3 min followed by

equilibration to 25°C for 5 min. The three temperatures (52°C, 53°C, 54°C) for each sample were

combined and centrifuged at 20,817 xg for 99 min at 4°C to remove aggregated protein.

Supernatant (soluble fraction) was moved to a new tube. All samples were frozen on dry ice and

dried down in speedvac (**Fig. S10A**).

For tandem mass tag (TMT) proteomics, 50 µg of protein (whole brain) or 12 µg (organelle

depleted cytoplasm) was dried down in a speedvac (Thermo Scientific, SPD111V), resuspended

in 23 µl of 1x lysis buffer (5% SDS, 50 mM TEAB, pH8.5) and digested with Trypsin/LysC (Thermo

Scientific, A41007) using S-Trap micro columns (ProtiFi, C02-micro-80) as described by the

manufacturer. Eluted peptides were dried in a speedvac and resuspended in 20 µl (whole brain)

or 5 µl (organelle depleted cytoplasm) of 100 mM TEAB pH 8.5. The resuspended peptides were

added to a tube containing TMT tag for a final protein:TMT ratio of 1:2 (ThermoFisher, #A44521).

The reaction was quenched with 0.5 µl of 5% hydroxylamine and incubated for 15 min at 25°C

with shaking (300 rpm). Quenched samples were combined and dried down in aliquots with 50

μg set aside as "pre-fractionation" sample. For fractionation (whole brain only), 125 μg aliquot

was resuspended in 300 μl of 0.1% TFA and sonicated for 5 min. The sample was subjected to a

high pH reversed phase fractionation procedure as described by manufacturer (Pierce,

#84868). Fractions 2 and 8 were combined and dried down. Fractions 3-7 were kept separate

and dried down. Fractions and input samples were reconstituted with 20 μl of 5% acetonitrile,

0.1% formic acid and 3.8 μl was injected per replicate.

Liquid chromatography tandem mass spectrometry (LC–MS/MS) experiments were performed in

duplicate for fractionated samples and in triplicate for unfractionated samples. Samples were run

in a 145 min gradient from 10% of acetonitrile (0.1% formic acid) to 70% acetonitrile, 0.1% formic

acid in data dependent mode on a nanoACQUITY UltraPerformance LC (UPLC; Waters, Milford,

MA, USA) coupled to an Orbitrap Fusion Lumos Tribid mass spectrometer. MS data acquisition

was collected from 400 to 1400 m/z with a resolution of 120,000. MS2 Data Dependent Acquisition

was acquired with a resolution of 50000 at m/z 200 with a mass injection time of 120 ms. Other

acquisition methods and column setup were identical to those described previously.[61] Raw

data files were processed with MaxQuant version 2.4.11.0. with reviewed entries only of mouse

uniprotKB FASTA file (downloaded 08/09/2023) to identify peptide-spectrum matches (PSM) and

to consolidate replicate injections.[62] Search parameters included Trypsin/P enzyme, fixed

modification of carbamidomethylation on cysteine (+57 on C) and variable modifications of

oxidized methionine (+16 on M), phosphorylation of serine (S), threonine (T), and tyrosine (Y)

(+80 on STY) and N-terminal acetylation (+42). The files were processed with Reporter MS2 type

set to TMT 16-plex (+304 on peptide N-Terminal and Lysine). Assignments were made using a

20-ppm mass tolerance for first search and 4.5 for main search for the precursor and 0.5 Da mass

tolerance for the fragments with a 1% FDR set for PSM and Protein. The proteinGroups output

text file was imported into Perseus (2.0.10.0) for log_2 transformation, imputation using values

based on normal distribution, and quantile normalization.[63] Additional filters included numerical

value in at least 4 of the 16 samples and 2 unique peptides for protein identification. The exported

table was then processed in RStudio (R version 4.3.3) using limma (3.58.1) for differential

expression (or stability) testing with a linear model that had a factor for group, whose levels were

the four combinations of GT (Q331K or WT) and INJ (TBI or Sham). Empirical Bayes moderated

t-statistics were computed for contrasts that included four between-group comparisons (WT.TBI

vs WT.Sham, Q331K.TBI vs Q331K.Sham, WT.TBI vs Q331K.TBI, and WT.Sham vs

Q331K.Sham), main effects of GT (Q331K vs WT) and INJ (TBI vs Sham), and the interaction

between GT and INJ. Proteins with an FDR < 0.1 were considered significantly different for these

contrasts. For the PISA data processing, one Q331K sham sample was identified as an outlier

based on total intensity and excluded from further analyses. Data are available via

ProteomeXchange with identifier PXD068825 and 10.6019/PXD068825.

Western blotting

For Western blot analyses, protein samples were mixed with loading buffer (Boston BioProducts,

BP-111R) and boiled at 100 °C for 5 min. Gel electrophoresis was performed with the BIO-RAD

PowerPac HC followed by a transfer onto polyvinylidene fluoride (PVDF; Merck Millipore,

IPFL00010) membrane. The PVDF membrane was blocked for 1h at ambient temperature using

Intercept Blocking Buffer (LI-COR, 927-70003) and then primary antibody (see table below) was

incubated at 4°C overnight (>12 h) in PBST and secondary antibody was incubated at ambient

temperature for 1 h in blocking buffer. The PVDF membrane was imaged using the LI-COR

ODYSSEY infrared imager system (Serial No. ODY-2215). The target protein intensity was

quantified using Image Studio software (Lite Ver 5.2).

Antibody	Manufacturer	Catalog #	Dilution
H3	Proteintech	17168-1-AP	1:1000
Calr	Enzo life sciences	ADI-SPA-602	1:1000
Cox4	Abcam	AB14744-1005	1:1500
GAPDH	Sigma-Aldrich	G8795	1:2000
Rpl4	Proteintech	11302-1-AP	1:2000
Rps6	Cell Signaling	2217	1:500
TDP43	Encor	MCA3H8	1:5000
Anti-Mouse IgG 680	LICORbio	926-68072	1:10,000
Anti-Rabbit IgG 800	LICORbio	926-32211	1:10,000

Immunostaining

Mice were euthanized and transcardially perfused with PBS and 4% paraformaldehyde (PFA;

Fisher Scientific, AAA1131336) in 1X PBS, pH 7.4. Mice were decapitated, and the scalp and

skull carefully removed to harvest the intact brain. All brain samples were immersed in 4% PFA

solution for 24h at 4°C for post-fixation. A brain matrix (CellPoint Scientific, Alto Acrylic 1mm

Mouse Brain Coronal) was used to pre-section the brains into anterior, medial, posterior and

cerebellar sections, which were embedded in paraffin by the Morphology Core (UMass Chan

Medical School). Coronal sections of 7 μm thickness were prepared on a Leica RM2125 RTS and

mounted onto charged microscope slides (Globe Scientific Cat#1354W). Prior to immunostaining,

tissues were deparaffinized with xylenes (Fisher Scientific, X3P-1GAL) and rehydrated in 100%

and 70% ethanol (Fisher Scientific, 04355223EA). Tissues were subjected to antigen retrieval

using 10mM Sodium Citrate (Fisher Scientific, S25545) containing 0.05% Tween 20 (Fisher

Scientific, BP337-500) at pH 6.0, heated to boiling, and then maintained at 95°C in a steamer

(Black&Decker Cat# HS1050) for 30 minutes. Tissues were blocked using 1% bovine serum

albumin (BSA; Fisher Scientific, BP1600-100) and 3% horse serum (Gibco, 26050-070) in PBS

for 1h at ambient temperature. Tissues were probed with rabbit anti-TDP-43 (1:250 Proteintech,

10782-2-AP) and chicken anti-NeuN (Millipore Sigma, ABN91) antibodies at 4°C overnight.

Tissues were incubated at ambient temperature for 1h with fluorophore-conjugated secondary

antibodies (Jackson ImmunoResearch Laboratories) at 1:500 and counterstained with DAPI.

Prolong Gold antifade reagent (Invitrogen, P36930) with a refractive index of 1.46 was used to

mount coverslips onto the tissues.

Confocal Microscopy

TDP-43/NeuN/DAPI immunofluorescence images were obtained using a Leica TCS SP5II (S/N:

5100001537, 11888906 BZ:01) confocal microscope. A 40x objective lens with oil was used to

acquire images. Leica Application Suite Advanced Fluorescence (LAS AF) software was used for

image acquisition as follows: imaging mode: XYZ; Sequential imaging: on; Format: 512x512;

Pinhole: on; Speed: 400 Hz; Line average: 1; Frame average: 3, Accu for Line average: 1; Accu

for Frame average: 1; Z-stack mode: Z-wide; Sequential scan mode: between frames, number of

steps: 3; Z-step size: ~0.5 μm. Using these settings, images were collected in the following order:

ipsilateral cortex, ipsilateral amygdala, ipsilateral hypothalamus, contralateral hypothalamus,

contralateral amygdala and contralateral cortex.

TDP-43 localization analyses

TDP-43 nuclear-to-cytoplasmic (N/C) ratio analysis was performed with 40x confocal images

described above on a per image basis with a custom macro generated with Fiji (**Additional file**

2). (64) DAPI signal was used to define the area of nuclear TDP-43 fluorescence and the

cytoplasm was defined as any area outside of the DAPI-defined region. Mean TDP-43

fluorescence intensity was measured for the nuclear and cytoplasmic regions, and the N/C ratio

was calculated by dividing the mean nuclear TDP-43 signal by the mean cytoplasmic TDP-43

signal.

TDP-43 N/C ratio analysis was also performed on a per-cell basis using a customized pipeline

that was developed in CellProfiler(65) for calculation of the percentage of cells with increased

cytoplasmic TDP-43 (**Additional file 3**). DAPI signal was used to identify the nuclear region,

which was then expanded by 8 pixels to capture a contiguous cytoplasmic region. The mean TDP-

43 fluorescence intensity was measured for the defined nuclear and cytoplasmic regions and the

N/C ratio was calculated as described above for the per image analysis. Cells with an N/C ratio

<5.72 were classified as having cytoplasmic TDP-43. This threshold was determined using the

average N/C ratio of each field of view (FOV) in the WT sham condition (representing a condition

that did not exhibit robust cytoplasmic TDP-43) minus one standard deviation. Within a FOV, the

number of cells with an N/C ratio <5.72 was then divided by the total number of cells, multiplied

by 100 to obtain the percentage cells with cytoplasmic TDP-43 for each condition.

The CellProfiler pipeline above was also used to measure the distribution of TDP-43 signal within

the nucleus and to quantify the number of cells with a "ring-like" TDP-43 staining appearance.

The nucleus, as defined by DAPI, was divided into three concentric regions using the Measure

Object Intensity Distribution module in CellProfiler and setting the number of bins to three. The

rings are therefore proportional to the size of each nucleus and nuclear area is accounted for in

the pipeline. For example, the outer ring represented the outer 1/3 of the nucleus, regardless of

nuclear size. TDP-43 fluorescent signal was measured within each of the three nuclear ring-

regions. Cells exhibiting ring-like TDP-43 staining were defined as those with >56% of the total

nuclear TDP-43 signal in the outer region (i.e., at the boundary of the nuclear envelope). This

threshold was determined by calculating the average percent of total nuclear TDP-43 signal

present in the outer region of the nucleus (the outermost concentric ring) in all WT sham cells

plus one standard deviation. Within a FOV, the percentage of cells with >56% nuclear TDP-43

signal in the outer nucleus out of the total cells per each condition was determined and used for

analysis.

The outcomes of the TDP-43 localization analyses above were combined to generate a measure

of cells with altered TDP-43 localization, which included cells that exhibited TDP-43 ring-like

and/or cytoplasmic TDP-43 according to the following parameters:

TDP-43 localization	N/C Ratio Value	% TDP-43 in outer ring of nucleus
Nuclear	>5.72	<56%
Ring-like, no cytoplasmic	>5.72	>56%
Ring-like with cytoplasmic	<5.72	>56%
Cytoplasmic, without ring-like	<5.72	<56%

The total number of cells that demonstrated each of the four TDP-43 distributions from the above

table (nuclear, ring-like +/- cytoplasmic and cytoplasmic) was summed then divided by the total

cells per FOV in each condition and multiplied by 100 to calculate the percent of cells with altered

TDP-43 localization per mouse. To visualize the overall effect of genotype and TBI on altered

TDP-43 localization, the percent of cells with each TDP-43 distribution was averaged across all

mice in each condition by FOV and plotted in a stacked bar graph in **Fig. 4**.

Reverse transcription polymerase chain reaction (RT-PCR)

RNA was isolated as described above and cDNA was synthesized using iScript Reverse

Transcription Supermix (Bio-Rad, 9170-8840) according to the manufacturer's instructions. RT-

PCR was performed on mouse tissue using 2x Taq RED Master Mix (APExBio Cat. No. K1034)

with the primers listed (**Supp. Table V**) with a touchdown PCR according to the manufacturer's

instructions. Products were separated on 3.5% agarose gels using the SYBR Safe DNA gel stain

(Invitrogen) and imaging was performed using the Bio-Rad Chemidoc software. Quantification of

the different isoforms was performed with ImageStudio software (LICORbio), using manual

settings to analyze the densitometry of each band. The intensity ratios between products with

included and excluded exons were analyzed for samples from a minimum of four individual mice

per group.

Structural analysis

The mm39 reference genome in Integrative Genomics Viewer, with its default RefSeq Curated gene annotation track, was used to identify exons in *Ube3c*. The splice variant, with the corresponding exon removed from the amino acid sequence, was modeled in AlphaFold 3.(66) Structural alignments of the models were performed in PyMOL (Molecular Graphics System, Version 3.1.3.1, Schrödinger) by minimizing the root-mean-square deviation (RMSD) between superimposed structures. The resulting aligned structures were visualized in PyMOL.

Ribosomal Profiling of cytoplasmic fractions

Ribosomal profiling was performed as previously described (67) on cytoplasmic fractions. A 20% to 50% sucrose gradient was prepared using a Gradient Master 107 (BioComp) in 14 x 89 mm polypropylene ultracentrifuge tubes (Beckman 331372). A total of 6 mL of 20% sucrose solution in 10 mM HEPES pH7.4, 150 mM NaCl, 5 mM MgCl2 with 100 μg/ml of cycloheximide, was added and overlayed with 50% sucrose solution made using the same buffer conditions. The top of the sucrose gradient (300 μl) was removed and replaced with 300 μl of sample (800 μg) containing 100 μg/ml of cycloheximide and 40 U/ml of RNaseIN. Samples were centrifuged at 197,000 x g for 2h at 4°C using a Beckman SW41 Ti rotor. Samples were loaded on a Density Gradient Fractionation system (Brandel), pushed with 60% sucrose and 0.72 ml fractions were collected with real-time UV monitoring at 254 nm using an ISCO UA-6 UV detector. Protein was precipitated from fractions using cold acetone. Protein pellets were collected by centrifugation at 13,000 x g for 10 min at 4°C, resuspend in 100 μl of SDS loading buffer, and heated for 5 min at 90°C.

RESULTS

Mutant TDP-43 mice exhibit heightened neurological deficits and delayed recovery following TBI. To examine the effects of TDP-43 dysfunction in the context of CNS challenge,

we administered a mild TBI using a weight-drop device (**Fig. 1A**) to mice harboring a knock-in

ALS-linked Q331K mutation in TDP-43.(44) TDP-43 Q331K mice (hereon referred to as Q331K

mice) exhibit phenotypes consistent with frontotemporal dementia (FTD), but without motor

deficits or motor neuron degeneration that is associated with ALS.(44) This mild TBI model has

been described by us previously.(39, 40, 43, 51, 52, 68) Here, a free-falling 50g weight was

dropped from a height of 15cm and used to deliver multiple impacts to the mouse's exposed but

intact skull within a single surgical session (**Fig. 1A,B**). Sham animals underwent the same

surgery as TBI animals to retract the skin from the skull but without any impact.

TBI studies included four main groups of mice at 16-weeks of age: Q331K and WT mice,

each with sham or TBI surgery. Neurological evaluations of motor function, alertness and

physiological behavior pre- and post-TBI were accomplished using a set of 10 tasks.(53) The

results of these tests are reported as a Neurological Severity Score (NSS), which is used to

assess potential neurological deficits following TBI in rodent models, where a higher score reflects

worse neurological deficits.(53). Consistent with our previous reports,(39, 43, 68) WT mice

exhibited worse NSS scores after TBI compared to their sham counterparts at 2 h post-TBI.

Notably, Q331K mice exhibited significantly worse NSS scores during the acute phase post-TBI

(i.e., 2h, 24h, and 48h post-TBI) compared to their WT counterparts, demonstrating heightened

vulnerability and delayed recovery to brain trauma **(Fig. 1C)**. All groups returned to baseline by

one-week post-TBI, as expected for a mild-TBI paradigm.(43)

Mice were also evaluated for long-term deficits resulting from TBI using the cued and contextual

fear conditioning (CCFC) test, an assessment of learning and memory.(69, 70) The CCFC test

consisted of four days, with habituation on day 1 (**Fig. 1D**), conditioning with a paired tone and

foot shock on day 2 (**Fig. 1E**), testing recall of the foot shock in response to the tone (i.e., cue) on

day 3 (**Fig. 1F**), and testing recall of the CCFC chamber (i.e., context) on day 4 (**Fig. 1G**). An

animal's freezing behavior on days 3 and 4 reflects its capacity for learning and memory, where

Figure 1. Neurological and cognitive deficits in mice with and without brain trauma. (A) Diagram of the weight drop device used to induce multiple closed-skull impacts with a 50g weight dropped from a 15cm height within a single surgical session. Location of impact site (red circle) is shown on the right. Created with biorender.com. **(B)** Illustration depicting the location of the impact site (red lightning bolt and oval). Several brain regions of interest are highlighted as follows: cortex (Ctx), amygdala (Amg), and hypothalamus (Hypo) on both ipsilateral and contralateral sides of the brain. Created with biorender.com. **(C)** Neurological severity score (NSS) plotted as a function of time post-surgery and cohort (see key). Statistically significant comparisons shown on the graph include Q331K TBI versus Q331K sham at 2h (****p=2.98e-6), 24h (***p=0.00058) and 48h (***p=0.00018) post-surgery, WT TBI versus WT sham at 2h post-surgery

(###p=0.00070) and Q331K TBI versus WT TBI at 2h (△△△p=0.00026), 24h (△△△p=0.00022) and 48h (△△p=0.0058) post-surgery. Additional significant comparisons include Q331K TBI versus WT sham at 2h (p=7.0e-10), 24h (p=2.54e-6) and 48h (p=0.00014) post-surgery. **(D-G)** Mice were randomly assigned to Group 1 (sham) or 2 (TBI). Results of the cued and contextual fear conditioning (CCFC) test. **(E)** The Q331K-2 cohort exhibited altered freezing behavior during conditioning compared to Q331K-1 (*p=0.028) and WT-2 (*p=0.017). **(F)** Both Q331K cohorts exhibited attenuated cue-induced freezing relative to WT animals (*p=0.012 for WT-2 vs. Q331K-1; **p=0.009 for WT-2 vs. Q331K-2) pre-surgery. WT animals exhibited attenuated cue-induced 5-months after sham (#p=0.048) and TBI (###p<0.001) surgeries.

attenuated freezing is indicative of deficits in learning and memory.(69, 70) WT and Q331K mice

were randomly assigned to group 1 or group 2, and both groups were subjected to the CCFC test

before surgery. Group 1 and group 2 animals were then, respectively, administered sham or TBI

surgery and assessed using the CCFC test again 5-months post-surgery. Linear mixed effect

analyses were then used to determine potential differences between groups and over time (i.e.,

between pre- and post-surgery) for the CCFC test. This uncovered significant effects for the

conditioning (Day 2) and cue-induced freezing response (Day 3). For conditioning, significant

group (P=0.010; numerator degrees of freedom or df=3, denominator df=107.770, F=3.991)

effects were detected, but not significant time (P=0.139; df=1, denominator df=107.773, F=2.220)

or group x time interaction (P = 0.724; numerator df=3, denominator df=107.770, F=0.441) effects.

On pair-wise testing, one of the Q331K cohorts was found to exhibit altered freezing behavior

during conditioning compared to one of the WT cohorts and also to the other Q33K1 cohort (**Fig.**

1E). As mice were randomly assigned to groups 1 and 2, the reason for the latter difference in

pre-surgery mice is unclear. For cue-induced freezing response, significant group (P = 0.016;

numerator df=3, denominator df=107.938, F=3.595) and time (P <0.001; numerator df=1,

denominator df=107.955, F=14.651) effects were detected, but not significant group x time

interaction (P = 0.298; numerator df=3, denominator df=107.938, F=1.243) effects. On pair-wise

testing, both Q331K cohorts exhibited attenuated cue-induced freezing on day 3 relative to WT

animals, where comparisons with the WT-2 cohort reached significance (**Fig. 1F**). These

outcomes are consistent with Q331K mice exhibiting cognitive deficits prior to surgery, as shown

for other cognition tests described by White et al.(44)

Both WT sham and WT TBI groups showed differences in the cue-induced freezing response at

5 months post-surgery versus the pre-surgery timepoint (**Fig. 1F**). The reduced freezing behavior

in WT sham animals 5 months post-surgery may stem from aging and/or adverse effects related

to the sham surgery. Conversely, Q331K mice showed no difference in the cue-induced freezing

response at 5 months when compared to pre-surgery, when they already exhibited cognitive

deficits. Therefore, there may be a ceiling effect with respect to cognitive dysfunction in Q331K

mice, given that Q331K mice exhibit similar cue-induced freezing behavior pre- and post-surgery,

regardless of surgery-type or age (**Fig. 1F**). Finally, there were no significant differences in

context-induced fear behavior due to genotype or surgery (**Fig. 1G**). Taken together, this mild-

TBI paradigm appears to exert minimal long-term effects on learning and memory, whereas

mutant TDP-43 expression may be sufficient to cause deficits in the amygdala, the brain region

that influences cue-induced behavior more so than others such as the hippocampus.(69, 70)

Proteostasis-related genes become upregulated in mutant TDP-43 mouse brain after TBI.

To investigate the gene expression changes associated with enhanced neurological severity

scores and delayed recovery in Q331K mice (**Fig. 1C**), bulk RNA sequencing (RNAseq) was

pursued using mouse whole brain lysate near the impact site (**Fig. 1A**) at the 24h post-TBI

timepoint. This analysis focused on four groups: Q331K and WT mice, each with sham or TBI

surgery. Of the ~25,000 genes that remained after filtering out genes with very low expression,

our analysis identified 228 that were differentially expressed between the Q331K TBI and WT TBI

mice (**Fig. 2A, Supp. Table I**).

Gene Set Enrichment Analysis(59, 60) was performed using the hallmark and Gene Ontology

Biological Process (GOBP) gene set collections from the Molecular Signatures Database

(MSigDB).(56-58) The hallmark collection includes 50 gene sets, with clearly defined gene

expression, as a high-level representation of a compilation of other gene sets in MSigDB, whereas

GOBP represents >7000 gene sets capturing biological processes. A normalized enrichment

score ($|NES|>1.5$) and an adjusted p-value cutoff ($FDR<0.05$) were used to determine significantly

altered pathways between Q331K TBI and WT TBI mice. NES reflects the overrepresentation of

a gene set at the upper (positive value) or lower (negative value) end of a ranked gene list. GSEA

analysis of the hallmark collection identified "mammalian target of rapamycin complex 1

(MTORC1) signaling" as the top upregulated gene set (**Fig. 2B**; **Supp. Table II**). MTORC1

signaling is involved with multiple cellular processes, including protein synthesis and

autophagy.(71) Protein folding was associated with the top 3 gene sets from GOBP identified in

the GSEA analysis (**Fig. 2C**). The relative expression for the top 30 ranked genes in the "MTORC1

signaling" (**Fig. 2D**) and "protein folding" (**Fig. 2E**) pathways are shown in a heatmap

representation, highlighting striking differences in gene expression between Q331K TBI mice

relative to the other three cohorts. Seven genes overlap between the top ranked list for MTORC1

signaling and protein folding, with *Hspa5*, encoding the endoplasmic reticulum (ER)-associated

chaperone heat shock 70 kDa protein 5, as the top ranked gene in both pathways (**Fig. 2D-E**).

To further examine differentially expressed genes between Q331K and WT mice as a function of

TBI, we compared significant transcriptional changes in Q331K TBI versus Q331K sham mouse

brain lysates (**Fig. S2A and Supp. Table I**) with those in WT TBI versus WT sham (**Fig. S2B and**

Supp. Table I) by plotting directional p-values (**Fig. 3A**), in which -$\log_{10}$(p-value) – a measure of

significance – is multiplied by the sign of the $\log_2$(fold-change), so that negative scores reflect a

decrease in gene expression for a specific comparison and positive scores reflect an increase in

expression. In this plot, genes that significantly change in expression in the Q331K TBI versus

Q331K sham comparison (sections shaded in pink) can be visualized along the x-axis, whereas

the y-axis highlights genes that significantly change in expression in WT TBI versus WT sham

mice (sections shaded in blue). Sections shaded in grey highlight genes that are differentially

expressed as a function of TBI, irrespective of genotype (**Fig. 3A**), and include 63 genes with

FDR<0.1 in both comparisons (**Fig. 3B, Supp. Table I**). GSEA hallmark pathway analysis

revealed "TNFa signaling via NFkB" as the top ranked, upregulated pathway in both Q331K (**Fig.**

3C and Fig. S2C) and WT (**Fig. 3D and Fig. S2D**) brain lysates post-TBI relative to their sham

counterparts (**Supp Table II**). This is not surprising, given that inflammation and changes in innate

immunity-related processes are well-documented following brain trauma.(72) Pathway changes

Figure 2. Upregulation of MTORC1 and protein folding genes in Q331K versus WT mouse brain after TBI. (A) Volcano plot of RNAseq data comparing Q331K and WT mice, each with either TBI or sham surgery. Differentially expressed genes (DEGs) with FDR <0.1 are shown in red. Top up- and down-regulated annotated genes are labeled. (B,C) Top GSEA gene set hits (both up and down regulated, as applicable) for Q331K TBI relative to WT TBI using the Molecular Signatures Database (MSigDB) hallmark (B) and Gene Ontology Biological Process (C) collections. Purple bars denote gene sets associated with proteostasis. (D,E) Heatmaps showing the top 30 ranked genes in the hallmark gene set "MTORC1 signaling" (D) and the Gene Ontology Biological Process gene set "protein folding" (E). The $\log_2$(fold change) relative to the mean expression is shown for each sample. All pathway hits meet cutoff criteria of

650   FDR <0.05 and |NES|>1.5. Genes in both "MTORC1 signaling" and "protein folding" gene sets are indicated

651   in bold.

that were specific to the WT TBI versus WT sham comparison indicate a downregulation of

metabolic and bioenergetic processes that occur in the mitochondria, such as "oxidative

phosphorylation" and "NADH dehydrogenase complex assembly" (**Fig. 3D,E and Fig. S2E,F**).

TBI is known to affect mitochondrial bioenergetics, where TBI induces mitochondrial damage and

results in reduced energy production.(73) Although these pathways were not significantly altered

between Q331K TBI and Q331K sham brain lysates, expression of genes in these pathways were

variable and potentially dysregulated in Q331K mice under both TBI and sham conditions, an

observation that may reflect transient and adverse effects of mutant TDP-43 on mitochondria

reported by others.(74)

Strikingly, nine out of the top ten GOBP gene sets emerging from the Q331K TBI versus Q331K

sham comparison were proteostasis-related, including protein folding and response to ER-stress

(**Fig. 3F**), which were not enriched gene sets in the WT TBI versus WT sham comparison (**Fig.**

3D, Supp Table II). Further, proteostasis-related genes *Hspa5*, mesencephalic astrocyte-derived

neurotrophic factor (*Manf*), calreticulin (*Calr*) and protein disulfide isomerase family A, members

3 (*Pdia3*) and 6 (*Pdia6*) were significantly increased in Q331K TBI versus Q331K sham mice (**Fig.**

3A; pink-shaded region). Proteostasis-related gene expression changes did not persist long-term

post-TBI, as determined by a separate RNA-seq analysis on whole brain mouse lysates 7-months

post-TBI or sham surgery, for both WT and Q331K mice. In fact, there were no significant gene

expression changes (at cutoff FDR < 0.1) due to TBI for either genotype (i.e., Q331K TBI versus

Q331K sham and WT TBI versus WT sham; **Supp Table I**).

Considering the upregulation of transcripts involved in protein folding and ER-stress in Q331K

mice 24h post-TBI compared to the other three cohorts, we assessed protein level changes with

quantitative proteomics analysis of mouse brain lysates labeled with tandem mass tags (TMTs)

from the four cohorts. Other than TDP-43, no significantly upregulated proteins were detected

(**Fig. S3A, Supp Table III**). In fact, TDP-43 upregulation was observed in Q331K versus WT

Figure 3. Differentially expressed genes in WT and Q331K mouse brain as function of surgery. (A)
Plot of directional p-values ($-\log_{10}$(p-value)*sign of $\log_2$(fold change)) from transcriptomics datasets for the
WT TBI versus WT sham comparison plotted against the Q331K TBI versus Q331K sham comparison. The
sign of the score reflects the direction of the fold change. Orange dots denote genes with expression
changes specifically due to TBI, irrespective of genotype. Blue and green dots denote genes with
expression changes specifically in WT and Q331K mice, respectively, after TBI. Color-shaded sections
indicate unique and overlapping gene changes, as illustrated in B. Dotted lines indicate FDR=0.1 **(B)** Venn
diagram illustrating minimal gene overlap between Q331K and WT mice subjected to TBI using FDR <0.1.
(C-F) Top GSEA gene set hits from the RNAseq analysis for Q331K TBI versus Q331K sham conditions

705   using the hallmark collection **(C)**, WT TBI versus WT sham conditions using the hallmark collection **(D)**, WT

706   TBI versus WT sham conditions using the Gene Ontology Biological Process collection **(E)** and Q331K TBI

707   versus Q331K sham conditions using the Gene Ontology Biological Process collection **(F)**. All gene set hits

708   meet cutoff criteria of FDR <0.05 and |NES|>1.5. Purple bars denote gene sets that are associated with

709   proteostasis.

comparisons irrespective of surgery type, at both mRNA and protein levels (**Fig. S3C,D**),

consistent with a loss of TDP-43 autoregulation due to the Q331K mutation.(44) Despite a

significant upregulation of *Hspa5* at the mRNA level in Q331K TBI mice, differential expression of

Hspa5 protein was not observed across the four mouse cohorts (**Fig. S3B**). We additionally

probed for isoforms of active X-box binding protein 1 (Xbp1; **Fig. 2D**), which becomes spliced

(Xbp1s) and activated under conditions of ER-stress (**Fig. 3F**),(75) by Western blot analysis.

Xbp1s was not detected in samples from any mouse cohort and levels of unspliced Xbp1 (Xbp1u)

were unchanged across conditions (**Fig. S3E**). The apparent lack of upregulation at the protein

level may be due to deficits in protein synthesis or because 24h post-TBI is too early a timepoint

for detection of newly translated protein. Nonetheless, the transcriptomics data clearly indicate a

robust upregulation of proteostasis-related pathways at the mRNA level that is dependent on both

TBI and expression of mutant TDP-43.

To gain insight into the relevance of our TBI model to neurodegenerative diseases such as ALS

and FTLD-TDP, we compared the GSEA hallmark gene set analyses of the TBI versus sham

conditions for both WT and Q331K mice with recent transcriptomic analyses of human ALS post-

mortem spinal cord and FTLD-TDP brain tissues.(45, 46) Significantly upregulated gene sets in

FTLD-TDP and/or ALS CNS tissues, such as "TNFa signaling via NFkB" and "hypoxia", were also

significantly upregulated in our TBI model, irrespective of genotype (**Fig. S4**). Notably, several

significantly upregulated gene sets in FTLD-TDP and/or ALS CNS post-mortem tissues were only

observed in Q331K mice after TBI, but not WT mice after TBI, including "MTORC1 signaling",

"unfolded protein response" and "inflammatory response" (**Fig. S4**). Similarly, GO terms related

to cytoplasmic translation and innate immunity emerged from a single nucleus sequencing

analysis of neurons from a separate study of human post-mortem ALS and FTLD brain tissue.(76)

Therefore, there appears to be a synergistic effect of mutant TDP-43 dysfunction and acute brain

trauma on biological processes in the CNS that, at least partially, recapitulates neurodegenerative

disease processes on the ALS/FTLD spectrum.

Proteostasis-related factors and pathways are downregulated in mutant TDP-43 mice at

baseline. We performed similar omics analyses on cohorts of WT and Q331K mice without TBI,

including RNAseq on mice without surgery (i.e., naïve mice). In our view, the sham condition

represents the most relevant control for TBI, as it accounts for gene expression changes that

occur from the surgery itself, thereby revealing expression changes due to the impact. Including

naïve mice here serves as an additional control for identifying differentially expressed genes due

to expression of mutant TDP-43. For the Q331K versus WT sham comparison, only two

downregulated gene sets emerged from our GSEA analysis, "protein folding" and "aerobic

electron transport chain" (**Supp. Table II**). Similarly, "MTORC1 signaling", which includes

proteostasis-related genes, such as heat shock protein family chaperones *Hspd1*, *Hspe1*, and

Hspa4, was a downregulated pathway for the Q331K versus WT naïve comparison (**Supp. Table**

II). Plotting transcriptomics data with directional p-values for the Q331K versus WT sham groups

against Q331K versus WT naïve groups revealed a significant downregulation of genes involved

in the ubiquitin-proteosome pathway, including glucose-induced degradation protein 4 homolog

(*Gid4*) and speckle-type POZ protein (*Spop*), the latter a downregulated gene that was also

observed by White et al. in this Q331K model (**Fig. S5B** and **Supp. Table IV**).(44) Of note, both

Gid4 and *Spop* remain downregulated in Q331K sham and TBI mice at 7 months post-TBI relative

to WT groups (**Supp. Table I, Fig. S5C,D**). Further, our quantitative proteomics analysis of Q331K

versus WT sham groups uncovered a modest yet significant decrease in the large ribosomal

subunit protein P1 (Rplp1), which plays a critical role in embryonic and brain development (**Fig.**

S3F, **Supp. Table III**).(77) Collectively, these omics data point toward a deficit in proteostasis-

related pathways in Q331K mice at baseline.

TDP-43 exhibits altered nuclear and cytoplasmic localization following TBI. Different TBI models reportedly induce subcellular redistribution of TDP-43 from the nucleus, where the protein is predominately expressed under homeostatic conditions, to the cytoplasm.(13, 37-41) Enhanced cytoplasmic localization of TDP-43 is also well-documented in other stress paradigms in vivo and in vitro, indicating that subcellular redistribution of TDP-43 represents a physiological response to stress.(2-6, 78) Given the disparate responses between Q331K and WT mice to TBI (**Figs. 1-3**), we examined the cellular localization of TDP-43 in situ as a function of surgery and TDP-43 genotype. Mouse brain tissue sections from TBI and sham cohorts were stained with anti-TDP-43 antibody and counterstained with DAPI. Obvious changes in subcellular TDP-43 distribution were observed in the cortex, amygdala, and hypothalamus (regions defined in **Fig. 1B**) of both Q331K and WT mice (**Fig. 4**), whereas changes in TDP-43 subcellular distribution were minimal or not detected in other regions of the tissue section (not shown). Sham animals also exhibited some degree of cytoplasmic TDP-43 redistribution in these brain regions, possibly indicative of cellular "stress" caused by the sham surgery, and therefore the naïve condition was included here for further analysis.

To quantify TDP-43 cellular localization, confocal microscopy images were acquired within the cortex, amygdala, and hypothalamus of Q331K and WT mice in the absence of surgery (i.e., naïve) or 24h after receiving a sham or TBI surgery, for a total of six groups (**Fig. 4A,B**). First, TDP-43 nuclear-to-cytoplasmic (N/C) ratios were determined for all cells across a field of view (FOV) on a per image basis. Reductions in TDP-43 N/C in TBI versus naïve conditions were generally observed in all three brain regions for both WT and Q331K mice, although some comparisons did not reach statistical significance (**Fig. S6A**). Notably, TBI-induced reductions in TDP-43 N/C ratios occurred in both ipsilateral (i.e., TBI side) and contralateral hemispheres of the brain (**Fig. S6A**) and were prominent in the amygdala and hypothalamus. Therefore, the effects of TBI on subcellular TDP-43 localization extend beyond the site of injury in the cortex,

Figure 4. Alterations in TDP-43 localization in WT and Q331K mice as a function of surgery. (A-B)
Confocal immunofluorescence images from WT and Q331K mice under native (i.e., no surgery) conditions
or 24h post-surgery (sham and TBI). Tissues were stained for TDP-43 (green) and counterstained with
DAPI (blue). Images from the hypothalamus (A) and cortex (B) are shown. Cropped insets highlight cells
with cytoplasmic (asterisks) and ring-like distribution (triangles) of TDP-43 under conditions of TBI. Scale
bar=50 um in the low magnification image and =20 um in the insets. **(C)** Percent of cells that exhibit altered
TDP-43 localization (ring-like, reduced nuclear-to-cytoplasmic (N/C) ratio, or both) in the cortex, amygdala,
or hypothalamus, either ipsilateral (left) or contralateral (right) to the impact site 24h post-surgery. **(D)**
Stacked bar graphs representing the average percent of cells that exhibit nuclear TDP-43 localization (grey)
or a type of altered TDP-43 localization (defined in C and color-coded according to the key) within the
indicated brain region. **(C-D)** Quantification of TDP-43 localization was performed using tissues from n=4-
6 mice, where each dot represents a different mouse. Statistical analysis was performed using two-way
ANOVA followed by Tukey's multiple comparisons test: *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

likely because the weight-drop induces contact between the skull and brain tissue at the base of

the skull (**Fig. 1A,B**). TBI-induced changes in TDP-43 subcellular localization represented an

acute response to TBI, as differences in TDP-43 N/C ratios were no longer observed at 5-months

post-surgery (**Fig. S7**).

As changes in TDP-43 subcellular localization were not observed in every cell, the percentage of

cells exhibiting reduced TDP-43 N/C ratio relative to the naïve condition was determined from the

FOVs above. Q331K TBI mice exhibited a significantly higher percentage of cells with reduced

TDP-43 N/C ratio in all three brain regions relative to Q331K naïve mice for both ipsilateral and

contralateral hemispheres, which was collectively a more robust change than for their WT

counterparts (**Fig. S6B**). During this analysis, we also noticed a "ring-like" TDP-43 staining

pattern, which appeared as dense TDP-43 signal near the outer edge of (but still within) the

nucleus (**Fig. 4A**, Q331K inset) and that also occurred in cells with cytoplasmic TDP-43 staining.

We developed a CellProfiler pipeline (see Methods) to quantify the percentage of cells with this

phenotype and found significant ring-like TDP-43 staining in TBI versus control comparisons in

all brain regions for at least one genotype (**Fig. S6C**). Ring-like TDP-43 staining was most

prominent in the hypothalamus and amygdala for both Q331K TBI and WT TBI mice versus their

respective controls, on both sides of the brain (**Fig. S6C**).

To assess the full extent of altered TDP-43 subcellular localization after TBI, we compiled the

data above (**Fig. S6**) and determined the percentage of cells exhibiting reduced TDP-43 N/C ratio

and/or ring-like TDP-43 (**Fig. 4C,D**). TBI induced more pronounced changes in TDP-43

localization in the cortical region below the impact site in Q331K versus WT mouse brain,

particularly on the contralateral side (**Fig. 4C,D**). These analyses also highlighted that Q331K

sham animals exhibited significant TDP-43 subcellular localization changes that were more

similar to the Q331K TBI condition and were significantly different than their Q331K naïve

counterparts, especially in the hypothalamus (**Fig. S6B,C**). Taken together, these data

demonstrate an acute response involving the TDP-43 protein to brain injury that occurred

irrespective of genotype. That Q331K mice appeared to be more sensitive to both sham and TBI

surgery with respect to changes in subcellular TDP-43 localization further supports the notion that

TDP-43 dysfunction leads to altered stress responses in vivo.

Effects of TDP-43 mutation and TBI on select TDP-43 splice targets. An important nuclear

function of TDP-43 involves the alternative splicing of conserved exons for numerous RNA

targets,(1) which can become mis-spliced as a result of TDP-43 knock-down, overexpression

and/or mutation.(79, 80) We therefore considered whether TBI-induced subcellular redistribution

of TDP-43 (**Fig. 4** and **Fig. S6**) would manifest in altered splicing of TDP-43 RNA targets. To

address this possibility, we probed for potential changes in splicing patterns for eight known RNA

targets of TDP-43 within mouse brain lysates derived from near the TBI site (**Fig. 1A**) using RT-

PCR and primer sequences that cover the exon junctions of interest (**Supp. Table V**). TDP-43

RNA targets included several that become mis-spliced due to TDP-43 misexpression, including

DnaJ heat shock protein family (Hsp40) member 5 (*Dnajc5*), potassium voltage-gated channel

interacting protein 2 (*Kcnip2*), protein phosphatase 3 catalytic subunit alpha (*Ppp3ca*), and

translin (*Tsn*).(81-84) Splicing patterns were assessed by gel electrophoresis (**Fig. 5A**) and

reported as the fraction of the mis-spliced variant versus the common or more abundant variant

expressed under normal conditions (**Fig. 5B**). For *Dnajc5*, no differences in splicing of exon 5

were observed between the mouse groups studied herein. In contrast, enhanced exon exclusion

most consistent with TDP-43 loss-of-function was detected for Q331K mice, irrespective of

surgery, for *Kcnip2* (exon 3 only, indicated by an increase in S1), *Ppp3ca* and *Tsn*.(84) Consistent

with TDP-43 gain-of-function reported in this and other Q331K mouse models,(44, 82) enhanced

exclusion of exon 17b in Sortilin-1 (*Sort1*) was observed in both Q331K TBI and sham mice; in

contrast, inclusion of exon 17b correlates with TDP-43 loss-of-function.(29, 81-83)

Figure 5. **Alternative splicing patterns for TDP-43 targets in WT and Q331K mouse brain after TBI.**

(A) Splice variants of known TDP-43 RNA targets (i-vi) detected by RT-PCR analysis in WT and Q331K

mice 24h after either TBI or sham surgery. The PCR primers, exons of interest and references for the

primers used in this study are shown in **Supp Table VI**. Splicing patterns that are consistent with TDP-43

loss of function or gain of function are indicated. Common and mis-spliced variants are denoted by,

respectively, closed and open arrowheads with the corresponding size (bp=base pairs) of the PCR product.

Gels are representative of at least n=3 separate RT-PCR experiments. **(B)** Densitometry analysis of the

gels in (A) was used to determine the ratio of the mis-spliced/common PCR product for each gene of

interest. Each dot in the graph represents a separate mouse.

We examined additional genes with known skiptic exons caused by mutant TDP-43 gain-of-

function in vivo.(80, 82) These included parkin coregulated like (*Pacrgl*), solute carrier family 6

member 6 (*Slc6a6*) and ubiquitin protein ligase E3C (*Ube3c*).(82) Skiptic exons were detected in

Q331K samples for all three genes. Intriguingly, the skiptic exon in *Ube3c* was attenuated in

Q331K mice after TBI, thus representing a gene-splicing event that was modulated by TBI,

specifically in Q331K mice (**Fig. 5A,B**). *Ube3c* encodes an E3 ubiquitin ligase that promotes

proteasomal degradation of misfolded proteins.(85) TDP-43-mediated skipping of exon 6 in

Ube3c causes the removal of 54 amino acid residues between the ubiquitin-binding domain and

the catalytic Homologous to the E6-AP Carboxyl Terminus (HECT) domain (**Fig. S8A**). While the

impact of exon 6 removal on Ube3c function is unknown,(82) modeling the 3-dimensional protein

structure of Ube3c in AlphaFold(66) suggests that omission of exon 6 could induce conformational

perturbations in the protein N-terminal to the HECT domain (**Fig. S8B**). The effects of TBI on

Ube3c exon 6 skipping in Q331K mice were acute, as exon skipping was no longer observed at

7-months post-TBI. Additionally, the splicing patterns for the other seven candidate TDP-43

mRNA targets were unchanged at 7-months post-TBI compared to 24h post-TBI (**Fig. S9**). Taken

together, these RT-PCR results indicate that TBI-induced changes in subcellular TDP-43

localization do not strongly correlate with a loss of TDP-43 splicing function, as most of the

observed splicing differences were due to TDP-43 mutation and not TBI. An exception is TDP-43

Q331K-induced skipping of exon 6 in *Ube3c*, which is attenuated during the acute phase post-

TBI.

Ribosomal proteins exhibit thermal instability in Q331K mice. A cytoplasmic shift in TDP-43

localization appears to represent a physiological response to brain trauma in mice, irrespective of

TDP-43 genotype (**Fig. 4**). We hypothesized that mutations in TDP-43 could alter the nature of

this response by perturbing cytoplasmic cellular processes involved in stress recovery. For

example, disease-linked mutations in TDP-43 have been shown to affect the interaction of TDP-

43 with cytoplasmic RNA granules as well as the biophysical properties of these granules.(11, 12,

14-16) Here, we examined protein content and protein thermostability of mouse brain lysates in

both WT and Q331K mice after sham or TBI surgery using the Proteome Integral Solubility

Alteration (PISA) assay (**Fig. S10A**) coupled with TMT quantitative proteomics using mass

spectrometry.(86, 87) PISA capitalizes on the principle that proteins become insoluble when

denatured. By extension, how a protein partitions between the soluble and insoluble fraction as a

function of temperature can be used to detect changes in protein thermostability under various

conditions. PISA and similar assays were historically used to elucidate protein targets of small

molecules, which generally induce an increase in thermostability of their protein targets upon

binding.(86, 88) We reasoned that PISA could be repurposed here for identifying thermostability

shifts of cytosolic proteins and protein complexes in brain lysates as a function of TBI and TDP-

43 mutation. To simplify sample processing and analysis, lysates were first depleted of large

organelles. A Western blot analysis with histone H3 and cytochrome c oxidase subunit IV (CoxIV)

antibodies confirmed that this protocol depleted nuclei and mitochondria, respectively, whereas

anti-calreticulin confirmed that ER was retained (**Fig. S10B**).

For the PISA assay, lysates were subjected to a narrow temperature gradient (52-54°C) followed

by centrifugation to remove the insoluble protein (**Fig. S10A**). The resulting soluble fraction was

subjected to quantitative proteomics (**Fig. S10C**). In parallel, quantitative proteomics was

performed on cytosolic lysates without heat exposure to determine total protein levels (**Fig.**

S10D). Only modest differences in total protein levels between Q331K TBI and WT TBI cytosolic

lysates were detected by the proteomic analysis (**Fig. S10D and Supp. Table VI)**, for which no

gene sets were significantly enriched by GSEA (**Supp. Table VII**). In striking contrast, GSEA

analysis of the PISA proteomics for the Q331K TBI versus WT TBI comparison revealed multiple

downregulated processes, with seven of the top ten GOBP hits including terms for translation and

ribosome (**Fig. 6A**). As shown in the heatmap for "cytoplasmic translation" (**Fig. 6B**), ribosomal

proteins account for most of the top 30 proteins in this pathway. Proteins in the "cytoplasmic

translation" pathway become elevated within the soluble fraction from WT animals after TBI compared to their sham counterparts, as demonstrated in the heatmap (**Fig. 6B**) and a box plot for the top five ranked proteins in the heatmap, all of which are ribosomal proteins (**Fig. 6C**). Conversely, these ribosomal proteins are diminished within the soluble fraction from Q331K sham mice relative to WT mice, and while they do become enhanced after TBI, their levels remain significantly below that of WT TBI lysates (**Fig. 6B,C**).

Considering that decreased protein solubility in the PISA assay is indicative of reduced protein thermostability, our interpretation of the PISA outcomes is that ribosomal proteins and potentially ribosome complexes themselves exhibit reduced thermostability in Q331K mice relative to WT mice. Polysome profiling of cytosolic lysates confirmed that intact 80S ribosomes were indeed present for both WT and Q331K conditions (**Fig. S11A-C**). RNA fragment analysis of isolated RNA confirmed the presence of both 28S and 18S ribosomal RNA (**Fig. S11D**). Given that the effect of TDP-43 Q331K expression on ribosome stability occurred irrespective of surgery, and that moreover there were no proteins for which the genotype x injury interaction contrast was significant (FDR < 0.1) in the limma linear model, we focused on the main effect of genotype for additional proteomic analyses. This approach further verified that total levels of ribosomes in the cytosolic lysates were unchanged between TDP-43 genotypes (**Fig. 6D**), but that their thermostability was reduced in Q331K mice relative to WT mice (**Fig. 6E**). We note that Rplp1, which was reduced in the proteomics analysis of total brain lysates (**Fig. S3F**), was not quantifiable in the cytosolic proteomics because only one unique peptide was identified whereas ≥2 were required in our pipeline for protein level quantification (**Supp. Table VI**). A subsequent STRING analysis using Markov Clustering (MCL) for the 50 destabilized proteins (FDR < 0.1) revealed a major cluster of 27 proteins (Cluster 1) associated with "cytoplasmic translation" and "ribosome" (**Fig. 6F**).(89, 90) Importantly, most of the ribosomal proteins in this cluster were shown to immunoprecipitate with TDP-43 from cultured cell and/or mouse brain lysates, raising

A PISA: Q331K TBI vs WT TBI
GO Biological Process gene sets

B Q331K TBI vs WT TBI
Cytoplasmic Translation

C

D Q331K All vs WT All
Quantitative proteomics

E Q331K All vs WT All
PISA proteomics

F PISA: Q331K all vs WT all
destabilized proteins

Figure 6. PISA analysis identifies melting temperature shifts in ribosomal and chaperonin proteins

in Q331K mouse brain lysates. (A) Top 10 gene set hits with GSEA enrichment of thermostability changes

in Q331K TBI versus WT TBI brain lysates using GOBP. Purple bars denote gene sets involved in ribosome

biology. **(B)** Heatmap showing levels of soluble protein detected by PISA for the top proteins in the pathway

"Cytoplasmic Translation". The $\log_2$(fold change) relative to the mean expression is shown for each sample.

(C) Box plot of top-ranking proteins in the "Cytoplasmic Translation" gene set for the indicated condition.

(D,E) Volcano plot for the quantitative proteomics (**D**) and PISA (**E**) results for Q331K ("Q331K All") versus

WT ("WT All"), where TBI and sham conditions are combined for each genotype to assess the main effects

of genotype while controlling for baseline differences related to injury. Ribosomal proteins are highlighted

as purple dots. (**F**) Results of a STRING analysis using the Markov Cluster Algorithm showing significantly

destabilized proteins (FDR < 0.1) in the "Q331K All" versus "WT All" comparison. Key terms included for

each cluster are as follows. Cluster 1: cytoplasmic translation, Signal Recognition Particle (SRP)-dependent

co-translational protein targeting to membrane, ribosome; Cluster 2: positive regulation of establishment of

protein localization to telomere, association of TriC/CCT with target proteins during biosynthesis, ATP-

dependent protein folding chaperone; Cluster 3, tRNA-splicing ligase complex; Cluster 4: Myosin II filament,

cell shape; Cluster 5: Nisch, Ube2i; Cluster 6: synthesis of active ubiquitin: roles of E1 and E2 enzymes;

Cluster 7: SHC-transforming protein 1 (SHC1) events in Erythroblastic Blast 2 (ERBB2) signaling, calcium-

dependent protein kinase C activity, presynaptic cystosol. String protein-protein interaction enrichment p-

value < $1.0e^{-16}$. Ribosomal proteins found to interact with TDP-43 in other studies are in bold within Cluster

1.(18, 91, 92)

the possibility that association of mutant TDP-43 with ribosomes and/or ribosomal proteins could

impact ribosome stability.(18, 91, 92) For the polysome profiling study, we noticed that the

absorbance at 254nm was consistently lower for Q331K cytosolic lysates relative to that of WT

within the peak where medium polysomes elute from the gradient (**Fig. S11B**). Additionally, we

observed a subpopulation of TDP-43 associating with the 80S ribosome (**Fig. S11C**). These

observations suggest that translation efficiency may be reduced in Q331K mouse brain relative

to WT mice. "Cluster 2", the second largest cluster revealed by STRING, includes multiple proteins

involved in protein folding such as heat shock 70 kDa protein 4 (Hspa4) and four of the eight

subunits that comprise the eukaryotic cytosolic T-complex protein 1 (TCP-1) ring complex or

chaperonin containing TCP-1 (TRiC/CCT), including Tcp1/Cct1, Cct6a, Cct7 and Cct8.

Interestingly, Cheng et al. identified these as putative TDP-43 interactors.(91) TRiC/CCT plays

important roles in maintaining proteostasis and ensuring the proper folding of critical cellular

proteins, lending further support to the notion that proteostasis-related processes are vulnerable

and dysregulated in mutant TDP-43 mice.(93)

There were 40 proteins that exhibited enhanced stabilization in Q331K versus WT cytosolic

lysates, although no cluster larger than 5 proteins was observed with STRING analyses and MCL

clustering. Cluster 1 and 2 each contain 5 proteins associated with smooth muscle contraction

and endocytosis, respectively (**Fig. S10G**). Interestingly, PISA detected four peptides derived

from ubiquitin (**Fig. 6E**) with sequence homology shared amongst ubiquitin C, ubiquitin B,

ubiquitin A-52 (Uba52) and Rps27a (**Supp. Table VI**). Uba52 and Rps27a are both ubiquitin-

ribosomal fusion proteins.(94) Therefore, despite downregulation of proteostasis-related

processes in Q331K mice at baseline, some compensatory mechanisms may be at play to

overcome these adverse effects.

1007

1008

DISCUSSION

Herein, we show that expression of dysfunctional TDP-43 leads to deficits in multiple branches of the proteostasis network (**Fig. 7**). Proteostasis deficits occur independently of overt TDP-43 overexpression, aggregation or mislocalization, which were not observed here or in prior reports describing this TDP-43 Q331K knock-in mouse model.(6, 44, 95) Analysis of our RNAseq datasets comparing Q331K and WT mice without brain trauma (i.e., naïve or sham conditions) revealed downregulation of genes involved in protein folding, including multiple heat shock proteins and chaperones, as well as genes involved in the ubiquitin proteosome pathway, such as *Gid4* and *Spop*. Exon skipping of *Ube3c*, an RNA target of TDP-43 that encodes an E3 ubiquitin ligase, was also detected in brain lysates from Q331K mice. This gain of TDP-43 mis-splicing function was initially reported in TDP-43 mouse strains with endogenous mutations in the low complexity domain, including a Q331K variant.(82) Although changes in *Ube3c* expression levels were not detected by RNAseq, AlphaFold and PyMol analyses predict that mis-splicing of *Ube3c* could alter the structure of Ube3c near the ubiquitin-binding domain. Further, Fratta et al. reported that about 15% of the exon skipping events identified in mutant TDP-43 mice occurred in genes within the ubiquitin proteosome pathway.(82) A decrease in proteosome-encoding transcripts early in life is a predictor of shortened lifespan in killifish, a vertebrate model used for aging studies.(96) Together, these observations support the notion that dysfunctional TDP-43 adversely affects the quality control arm of the proteostasis network, which in turn correlates with a decline in health and life-expectancy.

PISA, a novel approach for identifying alterations in biological complexes through thermostability measurements,(86, 87) further highlighted protein folding as a vulnerable component of the proteostasis network in Q331K mice, where multiple proteins within the essential molecular chaperone TRiC/CCT were found to be destabilized. Further, 20 out of 79 core ribosomal proteins were found to exhibit reduced thermal stability through our PISA analysis.(97) These data suggest

Figure 7. Proposed model for how the proteostasis network is affected by TDP-43 dysfunction. The proteostasis network is illustrated in the center of the diagram. Under baseline conditions (top), genes involved in proteostasis and the ubiquitin-proteasome pathway (UPP) are downregulated and mis-spliced in Q331K mouse brain relative to WT mice. In Q331K mouse brain lysates, ribosomal proteins are destabilized, which may compromise protein synthesis and thus contribute to the cognitive deficits observed in Q331K mice. In response to CNS challenge in form of a TBI (bottom), compensatory mechanisms appear to be activated in Q331K mice, including an upregulation of proteostasis-related genes at the RNA level. However, Q331K mice fail to mount a physiological response to TBI, as demonstrated by worse neurological deficits and delayed recovery after brain injury compared to their WT counterparts. We posit that destabilization of ribosomal proteins and chaperone complexes involved in proteostasis underlies the enhanced neurological deficits exhibited by Q331K mice after TBI. Although Q331K mice ultimately recover from TBI, we predict that repeated CNS challenge as a function of aging will exacerbate the phenotypes uncovered herein, and could lead to neurodegeneration and pathological TDP-43 aggregation. Model figure generated with BioRender.

that complexes formed by ribosomal proteins, namely ribosomes, could be affected by expression

of dysfunctional TDP-43 in a manner that compromises this macromolecular complex (**Fig. 7**).

TDP-43 has been shown to associate with ribosomal proteins through co-immunoprecipitation

and proximity labeling experiments,(18, 48, 91, 92) and with actively translating ribosomes in

cultured cells.(47, 48) Ribosomal proteins have also been detected within pathological TDP-43

aggregates(48, 98) and in amyloid-positive aggregates within the aging brain.(96) In various

contexts, TDP-43 expression can either repress or activate protein synthesis.(49, 50)

Interestingly, the association of TDP-43 with translation-related factors can be modulated by the

phase-separation behavior of TDP-43.(18) Proper phase separation of TDP-43 in vitro and in vivo

requires residues 321-340, referred to as the conserved region (CR).(17-19) Deletion of the CR

resulted in enhanced interactions between TDP-43 and translation factors together with

heightened rates of global protein synthesis in vivo.(18) While the Q331K variant studied herein

also reportedly disrupts TDP-43 phase separation,(17, 19) our polysome profiling data suggest

that translation efficiency is reduced in Q331K mouse brain relative to WT brain. Therefore, there

may be differential effects of the CR on translational regulation depending on how the TDP-43

sequence is modified (i.e., deletion versus single point mutation). We note that PISA was

performed on organelle-depleted brain lysates in this study, such that changes to nuclear and

mitochondrial proteins would not have been captured in this analysis. Indeed, it will be interesting

to extend PISA for analysis of other cellular compartments and organelles in future studies.

1075

It remains to be determined precisely how expression of mutant Q331K in mice leads to reduced

thermostability of ribosomal proteins. Disease-linked mutations may perturb interactions between

TDP-43 and ribosomal proteins,(91) which could induce conformational changes within the

ribosome (**Fig. 7**), other complexes that include ribosomal proteins and/or free ribosomal

proteins.(99) Ribosomal protein stability could also be indirectly affected by mutant TDP-43

expression. For example, altered proteostasis, activation of stress-response pathways and other

stimuli can affect the post-translational modification status of ribosomal proteins, which may

impact ribosome conformation and activity.(100) Intriguingly, ribosomal proteins underwent a shift

toward enhanced stability in WT animals after TBI, possibly representing a mechanism that

protects these proteins during cellular stress.(101) Although we do not observe changes in overall

levels of ribosomal proteins in WT or Q331K mice, expression of mutant TDP-43 could influence

the stoichiometry of ribosomal proteins, if not at the age of mice studied here (i.e., 16 weeks),

potentially in older mice. The stoichiometry of ribosomes is reportedly compromised during aging,

possibly a side-effect of impaired proteosome activity, which may lead to an accumulation of

"orphan" ribosomal proteins that are prone to aggregate.(96, 101) TDP-43 dysfunction has also

been linked to deficits in transporting mRNAs that encode ribosomal proteins(102) and in the

stability of ribosome-encoding transcripts,(103) both of which could also impact ribosome integrity

and protein translation.(49, 50) Deficiencies in protein synthesis may contribute to some cognitive

deficits observed in Q331K mice,(44) including attenuated cue-induced freezing in the CCFC test

in Q331K mice prior to surgery.(69) Although TBI is known to induce translational repression,

upregulated expression of select proteins is thought to be important for mitigating the effects of

trauma.(104) Our data support a model whereby Q331K mice are unable to mount a physiological

response to TBI due to multiple deficits within the proteostasis network that likely include deficits

in protein synthesis and folding (**Fig. 7**).

1100

In addition to TBI,(13, 37-41) TDP-43 has been linked to other stress-related pathways in vitro(2-

6, 11, 12) and in vivo.(15, 78). That TDP-43 undergoes a change in subcellular localization in

response to a broad range of stressors suggests TDP-43 may serve as a general stress-sensor,

a notion supported by our observation that TDP-43 re-distributes to the cytoplasm under

conditions of sham surgery. Stress-induced subcellular re-distribution of TDP-43 is modulated by

TDP-43 RNA-binding capacity and TDP-43 post-translational modification status, and this

redistribution can coincide with mis-splicing signatures corresponding to TDP-43 loss of

function.(3, 4, 7-9, 105) Here, we observe robust TBI-induced changes in subcellular TDP-43

localization in both WT and Q331K mouse brain tissue, including a ring-like TDP-43 staining

pattern that has not been widely reported. Changes in TDP-43 localization were most prominent

in the hypothalamus and the amygdala, the latter region having been linked to behavioral

dysfunction in human sporadic ALS.(106) *A priori*, we expected TBI-induced changes in TDP-43

localization to coincide with splicing defects associated with TDP-43 loss-of-function.(4, 9, 82,

105) However, the splicing patterns of known TDP-43 targets were primarily influenced by TDP-

43 mutation and largely unaffected by TBI. It is possible that emerging reporters of TDP-43-

mediated cryptic splicing could be more sensitive for detection of TDP-43 loss-of-function in cells

with altered TDP-43 localization in our TBI model.(107)

1118

The outcomes of our study lend support to the hypothesis that ALS pathogenesis occurs through

a multistep process, where steps may represent genetic, environmental and/or behavioral risk

factors.(34, 35) Notably, we detected a synergistic effect of TDP-43 mutation and TBI on gene

expression profiles that are relevant to ALS pathogenesis. Indeed, there is substantial overlap in

GSEA hallmark gene sets in CNS tissues between Q331K TBI mice 24h post-TBI and human

post-mortem ALS/FTLD-TDP cases,(45, 46) even though the Q331K TBI mice eventually recover,

and the human cases represent end-stage disease. Our interpretation is that expression of mutant

TDP-43 causes mice to become vulnerable to CNS challenge, resulting in dysregulation of key

biological pathways that are also dysregulated during ALS/FTD pathogenesis. In Q331K mice,

TBI-induced phenotypes that were observed 24h post-TBI, such as exaggerated neurological

deficits (i.e., higher NSS), proteostasis-related gene upregulation, changes in *Ube3c* splicing and

subcellular TDP-43 re-distribution, were largely restored to baseline levels 5-7 months post-TBI,

consistent with this being a mild model of TBI.(43). Further, compensatory mechanisms, such as

an upregulation of proteostasis-related genes, appear to be activated in Q331K mice (**Fig. 7**).

Despite these protective measures, Q331K mice exhibited worse neurological outcomes in the

acute phase post-TBI, indicating that stress-related pathways were vulnerable to dysfunctional

TDP-43. This notion is further supported by a recent study demonstrating reduced stress granule

formation in mutant TDP-43 mice exposed to hyperthermia.(15) We posit that repeated incidence

of TBI and/or exposure to additional risk factors would lead to pathological and chronic

degenerative phenotypes that persist months after the injury occurs (**Fig. 7**). Mathematical

modeling of incidence data for human ALS cases with TDP-43 mutation indicated that disease

pathogenesis may require up to five steps, where only one step is attributed to TDP-43

mutation.(34, 35) In the case of sporadic ALS, six steps are predicted for disease pathogenesis.

Accordingly, environmental and behavioral risk factors likely play a significant role in ALS

pathogenesis.(32, 33) While the TDP-43 Q331K mutation represents one step in the current

study, TDP-43 dysfunction can also manifest in the absence of genetic mutations. For example,

altered post-translational and transcriptional forms of TDP-43 that have been detected in sporadic

ALS cases have also been linked to TDP-43 dysfunction.(108, 109)

1147

Conclusions

Outcomes of this study demonstrate that expression of dysfunctional TDP-43 predisposes mice

to worse neurological outcomes following brain trauma. Deficiencies in multiple branches of the

proteostasis network likely contribute to this vulnerability of mutant TDP-43 mice to TBI (**Fig. 8**).

Therefore, targeting the proteostasis network should be a viable therapeutic direction for treating

TDP-43 proteinopathies. For example, proteostasis could be enhanced by mitigating protein

misfolding and stress through activation of heat-shock proteins and chaperones.(110) TDP-43

dysfunction can manifest through both genetic and non-genetic causes, where mutant TDP-43

expression altered the physiological response to TBI herein. We predict that TDP-43 dysfunction

will similarly affect cellular response and recovery pathways associated with other forms of CNS

challenge.(15) Further, repeated and/or additional incidences of CNS challenge are expected to

increase the likelihood of developing a TDP-43 proteinopathy such as ALS (**Fig. 8**).

1160   List of abbreviations

Abbreviation	Description
AD	Alzheimer's disease
ALS	amyotrophic lateral sclerosis
BCA	bicinchoninic acid
BSA	bovine serum albumin
Calr	calreticulin
CCFC	cued and contextual fear conditioning
CR	conserved region
CTE	chronic traumatic encephalopathy
Dnajc5	DnaJ heat shock protein family (Hsp40) member 5
ER	endoplasmic reticulum
ES	enrichment score
FOV	field of view
FTD	frontotemporal dementia
FTLD	frontotemporal lobar degeneration
Gid4	glucose-induced degradation protein 4 homolog
GO	gene ontology
GOBP	Gene Ontology Biological Process
GSEA	Gene Set Enrichment Analysis
GT	genotype
HECT	homologous to the E6-AP Carboxyl Terminus
Hspa4	heat shock 70 kDa protein 4
IACUC	Institutional Animal Care and Use Committee
IDR	intrinsically disordered region
INJ	Injury (i.e. sham or TBI)
Kcnip2	potassium voltage-gated channel interacting protein 2
LATE	limbic-predominant age-related TDP-43 encephalopathy
LC-MS/MS	liquid chromatography tandem mass spectrometry
Manf	mesencephalic astrocyte-derived neurotrophic factor
MCL	Markov Clustering
MSigDB	molecular signatures database
MTOR1	mammalian target of rapamycin complex 1
N/C	nuclear-to-cytoplasmic
NES	normalized enrichment score
NMDA	N-methyl-D-aspartate
NSS	neurological severity score
Pacrgl	parkin coregulated like

PCA	principal component analysis
Pdia3	protein disulfide isomerase family A, member 3
Pdia6	protein disulfide isomerase family A, member 6
PFA	paraformaldehyde
PISA	proteome integral solubility alteration
Ppp3ca	protein phosphatase 3 catalytic subunit alpha
pre-mRNA	pre-messenger RNA
PSM	peptide-spectrum match
PVDF	polyvinylidene fluoride
RNA-seq	bulk RNA-sequencing
RNP	ribonucleoprotein
Rplp1	large ribosomal subunit protein P1
Rps27a	ribosomal protein S27a
RT-PCR	Reverse transcription polymerase chain reaction
Slc6a6	solute carrier family 6 member 6
Sort1	Sortilin-1
Spop	speckle-type POZ protein
Ssr1	signal sequence receptor subunit 1
TAMC	Transgenic Animal Modeling Core
TBI	traumatic brain injury
TCP-1	cytosolic T-complex protein 1
TDP-43	TAR DNA-binding protein 43
TMT	tandem mass tag
TRiC/CCT	eukaryotic TCP-1 ring complex or chaperonin containing TCP-1
Tsn	translin
Uba52	ubiquitin A-52
Ube3c	ubiquitin protein ligase E3C
UPLC	UltraPerformance LC
WT	wild-type
Xbp1	active X-box binding protein 1
Xbp1s	spliced active X-box binding protein 1
Xbp1u	unspliced active X-box binding protein 1

1161

1162

1163

1164

1165

Declarations

Ethics approval and consent to participate. Not applicable.

Consent for publication. Not applicable.

Availability of data and materials. Mass spectrometry proteomics data are available via ProteomeXchange with identifier PXD068825 and 10.6019/PXD068825.Submission of RNAseq dataset to GEO are in progress at the time of this manuscript submission.

Competing interests. The authors declare that they have no competing interests

Funding. This work was supported by CDMRP/DoD DOD/W81XWH-20-1-0271 (D.A.B and N.H), the Angel Fund for ALS Research (D.A.B and N.H.), NINDS R21NS139100 (D.A.B.), and the First Affiliated Hospital of Chongqing Medical University (J.Z.).

Authors' contributions. MSR performed all the mass spectrometry experiments and analyzed the resulting data, performed most of the RNAseq analyses and was a major contributor in writing and preparing the manuscript; **MFM** performed the TDP-43 localization analyses, contributed to the RNA-related analyses, and created multiple figures in the manuscript; **JZ** performed animal surgeries, performed the CCFC analyses, and acquired the confocal microscopy images; **KO** performed the RT-PCR analyses; **EAW** isolated RNA from mouse tissue and coordinated the RNA sequencing and provided some assistance in the analysis; **DC** maintained the mouse colony and assisted in the animal studies; **KS** assisted in the animal studies and performed immunofluorescence; **JB** performed the NSS tests and helped with surgeries; **HM** performed the AlphaFold analyses and contributed to the RT-PCR analyses; **MNA** performed the polysome profiling studies together with MSR; **FM** performed the PyMOL analyses and interpreted the results; **ART** and **SM** assisted in the CCFC study; **JAN** assisted with confocal microscopy; **NH** contributed to the experimental design for the TBI surgeries and analyzed the animal behavior data; **ODK** performed the tests of differential gene expression for the RNAseq data and was a

major contributor to the omics analyses. **DAB** conceived and directed the project, analyzed data and wrote the manuscript with contributions from co-authors.

Acknowledgements. We are grateful for services provided by UMass Chan core facilities including the Transgenic Animal Modeling Core, Mouse Behavioral Core, Mass Spectrometry Facility and Molecular Biology Core Labs. We thank Dr. Joel Richter for use of equipment and advice related to polysome profiling; Drs. Patrick Sullivan, Hemendra Vekaria, Myriam Heiman, Jack Humphrey, Sami Barmada, Christine Vande Velde and Clotilde Lagier-Tourenne and members of their labs for valuable advice; and current and former members of Bosco lab, especially Dr. Ozge Yildiz and Jonathan Jung for help on this project.

Authors' information. Not applicable.

 Additional file 1. Supplemental Figures (Fig. S1 thru S11)

1217
1218 **Figure S1. Principal component analysis (PCA) identifies outliers in RNAseq datasets.** PCA analysis using 1000

1219 genes with the highest variance for the 24 h post-TBI **(A)** and 7 mo post-TBI (**B**) datasets. Genes were selected based

1220 on their variance (sigma^2) after adjusting for the mean-variance trend by fitting the expression data with the limma

1221 functions voom and lmFit, both using a design with a single group.

1222

1223

1224

1225

1226

1227

1228

Figure S2. Overlapping and differential gene expression profiles in WT versus Q331K mice after TBI. (A) Volcano plot of transcriptional changes from RNAseq analyses for Q331K mice under conditions of TBI versus sham surgery. **(B)** Volcano plot of transcriptional changes from RNAseq analyses for WT mice under conditions of TBI versus sham surgery. **(A,B)** Hits with FDR <0.1 are shown in red. Orange dots highlight top gene hits (*Klf2*, *Klf4*, *Mog*, *Clk1*) resulting from TBI, irrespective of genotype. Blue dots denote top gene hits specifically in the WT TBI condition whereas green dots are top gene expression changes specifically in the Q331K TBI condition. **(C,D)** Heatmap showing the top 30 ranked genes in the top up-regulated gene set for hallmark "TNFalpha signaling via NFKb" for Q331K TBI versus Q331K sham (C) and WT TBI versus WT sham (D). Fourteen genes (in bold) are significantly altered under "TNFalpha signaling via NFKb" for both the Q331K TBI and WT TBI conditions relative to their respective sham group. **(E,F)** Heatmap showing the top 30 ranked genes in the top down-regulated gene set in hallmark collection, "Oxidative Phosphorylation" (E), and Gene Ontology Biological Process collection, "NADH Dehydrogenase Complex Assembly" (F), for the WT TBI versus WT sham comparison. The log_2(fold change) relative to the mean expression is shown for each sample.

1258

Figure S3. Proteomic analyses identify protein changes altered by TBI and mutant TDP-43 genotype. (A)

Volcano plot showing relative protein level changes between Q331K TBI and WT TBI cohorts. Hits with FDR <0.1 are

highlighted in red. **(B)** Hspa5 is unchanged at the protein level by quantitative proteomics. **(C)** Volcano plot showing

relative protein level changes between Q331K sham and WT sham cohorts. Hits with FDR <0.1 are highlighted in red.

(D) Tardbp is significantly elevated in Q331K mouse brain by transcriptomics and proteomics analyses. **(E)** Total levels

of Xbp1 are unchanged and there is no detection of spliced Xbp1 by Western analysis. The expected molecular weight

of spliced Xbp (Xbp1s) is 54 kDa and of unspliced Xbp1 (Xbp1u) is 29 kDa. Note, the band located at ~35 kDa is

nonspecific. **(F)** Protein levels of Rplp1 as determined by quantitative proteomics. Nominal p-value = * <0.05; **<0.01,

****<0.0001.

1268

1269

1270

1271

Figure S4. Comparison of published human ALS and FTLD post-mortem datasets with current TBI mouse study. Directional p-values were calculated from published datasets reported in Humphrey et al,(46) and Hasan et al (45) and subjected to GSEA using the hallmark database as described in methods for RNAseq. Asterisks to the right of the bars denote FDR<0.05 for the indicated comparison defined in the key. The dotted line indicates NES of 1.5. Gene sets with a pink asterisk highlight those that are significant for the Q331K TBI versus Q331K sham comparison (pink bars), and at least one of the human datasets (grey and black bars). There are 16/21 gene sets that meet this criterion, and only 5/21 gene sets that are significant for the WT TBI versus WT sham comparison (blue bars) and at least one of the human datasets. Gene set names in bold are discussed in the main text.

1280

1281

1282

1283

1284

1285

1286

Figure S5. Omics analyses identify Q331K-specific changes to the transcriptome (A) Volcano plot with the significant up- and down-regulated transcripts in red for the Q331K sham versus WT sham comparison. **(B)** Directional p-value plot overlaying the Q331K sham versus WT sham datasets with the Q331K naïve versus WT naïve datasets to identify reproducible mutant TDP-43-specific changes, which are highlighted in red. Directional p-value = -log₁₀(p-value)*sign of log₂(fold change) **(C-D)** Plots of the most significant Q331K-specific down-regulated genes, *Spop* and *Gid4*, which are both involved in the ubiquitin-proteosome pathway. **(E)** Elevated *Tardbp* is observed in all Q331K vs WT comparisons. For C-E, time post-surgery is indicated below. Nominal p-value = **<0.01, ****<0.0001.

Figure S6. Changes in TDP-43 subcellular localization in mouse brain tissue 24h post-surgery. (A) Nuclear-to-cytoplasmic (N/C) ratio of TDP-43 signal in naïve mice or 24h post-surgery (sham or TBI) mice determined from immunofluorescence analysis of tissue sections from the cortex (left), amygdala (middle), and hypothalamus (right), either ipsilateral (top) or contralateral (bottom) to the TBI site. **(B)** Percent of cells that exhibit reduced TDP-43 N/C ratios in the tissues from (A). **(C)** Percent of cells that exhibit "ring-like" TDP-43 distribution in the tissues from (A). Each dot represents data from an individual mouse, with n=4-6 mice per cohort. Statistics was performed using two-way ANOVA followed by Tukey's multiple comparisons test between conditions. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

1317

Figure S7. Lack of significant TDP-43 subcellular alteration in mouse brain tissue 5 months post-surgery. (A)

Confocal immunofluorescence images of hypothalamus brain sections from WT and Q331K mice, either without surgery

(naïve) or 5-months post-surgery (sham or TBI). Sections are stained for TDP-43 (green) and counterstained with DAPI

(blue). Scale bars=20 um. **(B,C)** Quantification of the nuclear-to-cytoplasmic (N/C) ratio of TDP-43 signal in WT and

Q331K mice 5-months post-surgery in the cortex, amygdala and hypothalamus, either ipsilateral (B) or contralateral

with respect to the impact site. Each dot represents data from an individual mouse, with n=4-9 mice per cohort. Statistics

was performed using two-way ANOVA followed by Tukey's multiple comparisons test between conditions.

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

1335

Figure S8. Structural analysis of TDP-43-relevant splice variants using AlphaFold and PyMOL. (A) Schematic

representation of the ubiquitin-protein ligase E3C (Ube3c) protein highlighting distinct domains. Residue numbers are

indicated at the start of each domain. The ubiquitin-binding domain (residues 1–132) is shown in purple, the second

domain (residues 133–743) in green, and the catalytic Homologous to the E6-AP Carboxyl Terminus (HECT) domain

(residues 744–1083) in blue. Residues corresponding to exon 6 (residues 153–206) are highlighted in red. **(B)** Overlay

of AlphaFold-predicted structures for Ube3c containing exon 6 (domains in purple, green, and blue) with the predicted

structures for Ube3c lacking exon 6 (domains in light purple, light green, and light blue). Structural alignments were

performed in PyMOL, yielding an RMSD of 2.3 Å for alignment of only the ubiquitin-binding domain and an RMSD of

8.5 Å for the entire protein.

1345

1346

1347

1348

1349

1350

Figure S9. Alternative splicing targets of TDP-43 in WT and Q331K mice at 7 months post-injury. (A) Splice

variants of known TDP-43 RNA targets (i-vi) detected by RT-PCR analysis in WT and Q331K mice 24h after either TBI

or sham surgery. The PCR primers, exons of interest and references for the primers used in this study are shown in

Supp Table VI. Splicing patterns that are consistent with TDP-43 loss of function or gain of function are indicated.

Common and mis-spliced variants are denoted by, respectively, closed and open arrowheads with the corresponding

size (bp=base pairs) of the PCR product. **(B)** Densitometry analysis of the gels in (A) was used to determine the ratio

of the mis-spliced/common PCR product for each gene of interest. Each dot in the graph represents a separate mouse.

1358

1359

1360

1361

1362

Figure S10. Quantitative proteomics of organelle-depleted cytoplasmic brain lysates. (**A**) PISA protocol for identifying proteins with altered thermostability between conditions: (1) Lysates are generated with mild, non-denaturing conditions and homogenization. Lysates are cleared by centrifugation. (2) Each sample is divided into four samples, where one is placed on ice for traditional quantitative proteomics and three are subjected to heat for protein denaturation. (3A) Aggregated protein is removed by centrifugation. (3B) The sample for traditional TMT quantitative proteomics remains on ice while samples for PISA are heated. (4) The soluble protein is digested with trypsin and labeled (5) with isobaric TMTpro-18plex tags. (6) Tagged samples are pooled, generating two sample sets (one from PISA and one from traditional quantitative proteomics) and subjected to mass spectrometry for quantification (7). Created with biorender.com. (**B**) Western analyses of organelle depleted supernatant (supernatant) compared to the whole brain sample input (input). (**C-D**) Volcano plot of protein changes identified by traditional quantitative proteomics (**C**) and PISA (**D**) for Q331K TBI versus WT TBI conditions. (**E**) Overlapping hits between quantitative proteomics and PISA are assessed to distinguish protein level changes from thermostability changes. (**F**) Directional p-value plot of Q331K TBI versus WT TBI for PISA and QUANT (quantitative proteomics). Purple dots indicate ribosomal proteins.

Orange dots are highlighting top hits found in both PISA and QUANT. (**G**) Clustering using the Markov Cluster Algorithm

for "Q331K All" versus "WT All" (defined in Figure 7 legend) for the stabilized proteins passing FDR adjustment (FDR

<0.1). Only proteins identified as part of a cluster are shown. Cluster 1: Smooth Muscle Contraction; Cluster 2: Synaptic

vesicle endocytosis, Synaptobrevin 2-SNAP-25-syntaxin-1a-complexin II complex; Cluster 3: Inactivation of

Granulocyte colony-stimulating factor (G-CSF) signaling; Cluster 4: Protein-disulfide reductase nicotinamide adenine

dinucleotide phosphate (NAD(P)) activity; Cluster 5: S-100/Intestinal calcium-binding protein (ICaBP) type calcium

binding domain; Cluster 6: ER membrane insertion complex; Cluster 7: Epoxide hydrolase 2 (Ephx2), reactive

intermediate imine deaminase A (Rida); Cluster 8: Negative regulation of calcium ion export across plasma membrane.

Protein-protein interaction enrichment p-value = $2.94e^{-06}$.

1385

1386

1387

Figure S11. Polysome profiling of cytoplasmic fractions confirm the presence of intact ribosomes. **(A)** Graph of sucrose gradient fractions post-ultracentrifugation. The prominent peaks are at fractions 9/10, fractions 11/12, and fractions 15/16. **(B)** The area under the curve was calculated for all samples at 400 sec to 500 sec to estimate relative amount of medium polysomes, as this was the only ribosomal peak consistently observed in all samples. * = p-value <0.05. **(C)** Western analyses of the protein precipitated fractions of the polysome profile shown in A. Numbers indicate specific fractions analyzed. **(D)** Fragment analysis of RNA isolated from cytoplasmic fractions for each sample subjected to polysome profiling using the Agilent Fragment Analyzer.

Additional file 2. FIJI Macro for Per FOV N/C Analysis. Macro script to be used in FIJI/ImageJ to complete TDP-43 localization analysis on a per FOV level. DAPI signal is used to define and separate the nuclear and non-nuclear regions where TDP-43 fluorescent signal is measured to determine the N/C ratio.

Additional file 3. CellProfiler Pipeline for Per Cell N/C Ratio and Ring-like Analysis. CellProfiler Pipeline used for per cell analyses of TDP-43 fluorescent signal including N/C ratio and ring-like TDP-43 signal. DAPI signal is used to identify individual cells, as well as define nuclear and cytoplasmic regions for N/C ratio analyses. For ring-like analysis, the nuclear region is binned into 3 rings to report on distribution of TDP-43 signal.

Additional file 4, Supp. Table I. Statistical output of RNAseq analyses. For all comparisons, logFC refers to log2(fold change), PValue refers to nominal p-value, and FDR refers to adjusted p-value. Nominal p-values <0.05 are indicated in orange and FDR < 0.1 in red. (**IA**) Statistical analyses for the 4 mo old mouse cohorts including the 24 h post-surgery (TBI and sham) and naïve groups. (**IB**) Statistical analyses for the 7 mo post-surgery cohorts.

Additional file 5, Supp.Table II. GSEA analyses for all comparisons from the RNAseq analysis. All gene set hits that passed the cutoff criteria (NES > 1.5, FDR > 0.05) are listed. Each tab is a unique group comparison. In each tab, the top upregulated gene sets are beneath the purple-shaded header on the left, while the top downregulated gene sets are beneath the blue-shaded header on the right. Results of hallmark and GOBP are shown. (**IIA**) List of abbreviations used in the file. (**IIB-E**) Gene set hits for the 24h-post surgery (TBI and sham). (**IIF**) Gene set hits for Q331K naïve mice relative to WT naïve mice at 4 months of age. (**IIG-IIJ**) Detailed gene lists for the top pathway hits for the 24h group comparisons. (**IIK-IIN**) Gene set hits for each group comparison at 7 mo post-surgery.

Additional file 6, Supp. Table III. Statistical analyses of whole brain quantitative proteomics from Q331K and WT mice with surgery. Unadjusted p-values $\leq$ 0.05 are highlighted red (columns H-K). Adjusted p-values $\leq$ 0.1 are highlighted in red (columns L-O). Coef. refers to Log_2(fold change); P.value indicates nominal p-value; P.value.adj. represents adjusted p-value (FDR).

Additional file 7, Supp. Table IV. Overlap of published omics datasets with the current study. Includes all identified proteome and transcriptome level changes in Q331K relative to WT mice from the current study. Columns B-D refer to datasets in the present study. RNAseq_Sham refers to the Q331K sham versus WT sham comparison. RNAseq_Naive refers to Q331K naïve versus WT naïve at 4 months of age. Proteomics_Sham refers to Q331K sham versus WT sham whole brain proteomics. Columns E/F refer to results published by White et al.(44).

Additional file 8, Supp. Table V. RT-PCR primers. Sequence information for primers that were used to identify alternative splicing for targets of TDP-43 in **Figures 5** and **S8**.

Additional file 9, Supp. Table VI. Significant outcomes from quantitative proteomics on organelle-depleted cytoplasmic brain lysates. (**VIIA**) PISA quantitative proteomics results. (**VIIB**) Traditional quantitative proteomics results (i.e. not exposed to heat). Unadjusted p-values $\leq$ 0.05 are highlighted red (columns K-Q). Adjusted p-values $\leq$ 0.1 are highlighted in red (columns R-X). Coef. refers to Log_2(fold change); P.value indicates nominal p-value; P.value.adj. represents adjusted p-value (FDR).

Additional file 10, Supp. Table VII. GSEA analyses of quantitative proteomics outcomes for
organelle-depleted cytoplasmic brain lysates. All gene set hits that passed the cutoff criteria
(NES > 1.5, FDR > 0.05) are listed. Each tab is a unique group comparison. In each tab, the top
upregulated gene sets are beneath the purple-shaded header on the left, while the top
downregulated gene sets are beneath the blue-shaded header on the right. Results of hallmark
and GOBP are shown. (**VIIIA-D**) Gene set hits for samples processed using PISA. (**VIIIE-H**) Gene
set hits from the traditional quantitative proteomics analyses.
1457

References
1.   Buratti E, Baralle FE. Multiple roles of TDP-43 in gene expression, splicing regulation,
and human disease. Front Biosci. 2008;13:867-78.
2.   Dewey CM, Cenik B, Sephton CF, Dries DR, Mayer P, 3rd, Good SK, et al. TDP-43 is
directed to stress granules by sorbitol, a novel physiological osmotic and oxidative stressor.
Molecular and cellular biology. 2010.
3.   Duan L, Zaepfel BL, Aksenova V, Dasso M, Rothstein JD, Kalab P, et al. Nuclear RNA
binding regulates TDP-43 nuclear localization and passive nuclear export. Cell Rep.
2022;40(3):111106.
4.   Klim JR, Williams LA, Limone F, Guerra San Juan I, Davis-Dusenbery BN, Mordes DA,
et al. ALS-implicated protein TDP-43 sustains levels of STMN2, a mediator of motor neuron
growth and repair. Nat Neurosci. 2019;22(2):167-79.
5.   Tischbein M, Baron DM, Lin YC, Gall KV, Landers JE, Fallini C, et al. The RNA-binding
protein FUS/TLS undergoes calcium-mediated nuclear egress during excitotoxic stress and is
required for GRIA2 mRNA processing. J Biol Chem. 2019;294(26):10194-210.
6.   Weskamp K, Tank EM, Miguez R, McBride JP, Gomez NB, White M, et al. Shortened
TDP43 isoforms upregulated by neuronal hyperactivity drive TDP43 pathology in ALS. J Clin
Invest. 2020;130(3):1139-55.
7.   Perez-Berlanga M, Wiersma VI, Zbinden A, De Vos L, Wagner U, Foglieni C, et al. Loss
of TDP-43 oligomerization or RNA binding elicits distinct aggregation patterns. EMBO J.
2023;42(17):e111719.
8.   Zhang S, Vazquez-Sanchez S, Lu S, Baughn MW, Lim J, Oung S, et al. Acetylation of
lysine 82 initiates TDP-43 nuclear loss of function by disrupting its nuclear import. bioRxiv.
2024:2024.09.04.611121.
9.   Necarsulmer JC, Simon JM, Evangelista BA, Chen Y, Tian X, Nafees S, et al. RNA-
binding deficient TDP-43 drives cognitive decline in a mouse model of TDP-43 proteinopathy.
Elife. 2023;12.
10.   Dos Passos PM, Hemamali EH, Mamede LD, Hayes LR, Ayala YM. RNA-mediated
ribonucleoprotein assembly controls TDP-43 nuclear retention. PLoS Biol.
2024;22(2):e3002527.
11.   Liu-Yesucevitz L, Bilgutay A, Zhang YJ, Vanderwyde T, Citro A, Mehta T, et al. Tar DNA
binding protein-43 (TDP-43) associates with stress granules: analysis of cultured cells and
pathological brain tissue. PloS one. 2010;5(10):e13250.
12.   McDonald KK, Aulas A, Destroismaisons L, Pickles S, Beleac E, Camu W, et al. TAR
DNA-binding protein 43 (TDP-43) regulates stress granule dynamics via differential regulation of
G3BP and TIA-1. Human molecular genetics. 2011.
13.   Anderson EN, Gochenaur L, Singh A, Grant R, Patel K, Watkins S, et al. Traumatic
injury induces Stress Granule Formation and enhances Motor Dysfunctions in ALS/FTD Models.
Human molecular genetics. 2018.

14.   Alami NH, Smith RB, Carrasco MA, Williams LA, Winborn CS, Han SSW, et al. Axonal
transport of TDP-43 mRNA granules is impaired by ALS-causing mutations. Neuron.
2014;81(3):536-43.

15.   Dubinski A, Gagne M, Peyrard S, Gordon D, Talbot K, Vande Velde C. Stress granule
assembly in vivo is deficient in the CNS of mutant TDP-43 ALS mice. Human molecular
genetics. 2023;32(2):319-32.

16.   Gopal PP, Nirschl JJ, Klinman E, Holzbaur EL. Amyotrophic lateral sclerosis-linked
mutations increase the viscosity of liquid-like TDP-43 RNP granules in neurons. Proc Natl Acad
Sci U S A. 2017;114(12):E2466-E75.

17.   Conicella AE, Zerze GH, Mittal J, Fawzi NL. ALS Mutations Disrupt Phase Separation
Mediated by alpha-Helical Structure in the TDP-43 Low-Complexity C-Terminal Domain.
Structure. 2016;24(9):1537-49.

18.   Gao J, Wang L, Ren X, Dunn JR, Peters A, Miyagi M, et al. Translational regulation in
the brain by TDP-43 phase separation. J Cell Biol. 2021;220(10).

19.   Hallegger M, Chakrabarti AM, Lee FCY, Lee BL, Amalietti AG, Odeh HM, et al. TDP-43
condensation properties specify its RNA-binding and regulatory repertoire. Cell.
2021;184(18):4680-96 e22.

20.   Grese ZR, Bastos AC, Mamede LD, French RL, Miller TM, Ayala YM. Specific RNA
interactions promote TDP-43 multivalent phase separation and maintain liquid properties.
EMBO Rep. 2021;22(12):e53632.

21.   de Boer EMJ, Orie VK, Williams T, Baker MR, De Oliveira HM, Polvikoski T, et al. TDP-
43 proteinopathies: a new wave of neurodegenerative diseases. J Neurol Neurosurg Psychiatry.
2020;92(1):86-95.

22.   Arai T, Hasegawa M, Akiyama H, Ikeda K, Nonaka T, Mori H, et al. TDP-43 is a
component of ubiquitin-positive tau-negative inclusions in frontotemporal lobar degeneration
and amyotrophic lateral sclerosis. Biochem Biophys Res Commun. 2006;351(3):602-11.

23.   Neumann M, Sampathu DM, Kwong LK, Truax AC, Micsenyi MC, Chou TT, et al.
Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis.
Science. 2006;314(5796):130-3.

24.   Amador-Ortiz C, Lin WL, Ahmed Z, Personett D, Davies P, Duara R, et al. TDP-43
immunoreactivity in hippocampal sclerosis and Alzheimer's disease. Ann Neurol.
2007;61(5):435-45.

25.   Nelson PT, Dickson DW, Trojanowski JQ, Jack CR, Boyle PA, Arfanakis K, et al. Limbic-
predominant age-related TDP-43 encephalopathy (LATE): consensus working group report.
Brain. 2019;142(6):1503-27.

26.   McKee AC, Gavett BE, Stern RA, Nowinski CJ, Cantu RC, Kowall NW, et al. TDP-43
proteinopathy and motor neuron disease in chronic traumatic encephalopathy. J Neuropathol
Exp Neurol. 2010;69(9):918-29.

27.   Brown AL, Wilkins OG, Keuss MJ, Hill SE, Zanovello M, Lee WC, et al. TDP-43 loss and
ALS-risk SNPs drive mis-splicing and depletion of UNC13A. Nature. 2022;603(7899):131-7.

28.   Estades Ayuso V, Pickles S, Todd T, Yue M, Jansen-West K, Song Y, et al. TDP-43-
regulated cryptic RNAs accumulate in Alzheimer's disease brains. Mol Neurodegener.
2023;18(1):57.

29.   Ling JP, Pletnikova O, Troncoso JC, Wong PC. TDP-43 repression of nonconserved
cryptic exons is compromised in ALS-FTD. Science. 2015;349(6248):650-5.

30.   Ma XR, Prudencio M, Koike Y, Vatsavayai SC, Kim G, Harbinski F, et al. TDP-43
represses cryptic exon inclusion in the FTD-ALS gene UNC13A. Nature. 2022;603(7899):124-
30.

31.   Melamed Z, Lopez-Erauskin J, Baughn MW, Zhang O, Drenner K, Sun Y, et al.
Premature polyadenylation-mediated loss of stathmin-2 is a hallmark of TDP-43-dependent
neurodegeneration. Nat Neurosci. 2019;22(2):180-90.

32.   Al-Chalabi A, Hardiman O. The epidemiology of ALS: a conspiracy of genes,
environment and time. Nat Rev Neurol. 2013;9(11):617-28.

33.   Goutman SA, Savelieff MG, Jang DG, Hur J, Feldman EL. The amyotrophic lateral
sclerosis exposome: recent advances and future directions. Nat Rev Neurol. 2023;19(10):617-
34.

34.   Chio A, Mazzini L, D'Alfonso S, Corrado L, Canosa A, Moglia C, et al. The multistep
hypothesis of ALS revisited: The role of genetic mutations. Neurology. 2018;91(7):e635-e42.

35.   Al-Chalabi A, Calvo A, Chio A, Colville S, Ellis CM, Hardiman O, et al. Analysis of
amyotrophic lateral sclerosis as a multistep process: a population-based modelling study.
Lancet Neurol. 2014;13(11):1108-13.

36.   Barker S, Paul BD, Pieper AA. Increased Risk of Aging-Related Neurodegenerative
Disease after Traumatic Brain Injury. Biomedicines. 2023;11(4).

37.   Heyburn L, Sajja V, Long JB. The Role of TDP-43 in Military-Relevant TBI and Chronic
Neurodegeneration. Front Neurol. 2019;10:680.

38.   Jankovic T, Dolenec P, Rajic Bumber J, Grzeta N, Kriz J, Zupan G, et al. Differential
Expression Patterns of TDP-43 in Single Moderate versus Repetitive Mild Traumatic Brain Injury
in Mice. Int J Mol Sci. 2021;22(22).

39.   Kahriman A, Bouley J, Smith TW, Bosco DA, Woerman AL, Henninger N. Mouse closed
head traumatic brain injury replicates the histological tau pathology pattern of human disease:
characterization of a novel model and systematic review of the literature. Acta Neuropathol
Commun. 2021;9(1):118.

40.   Kahriman A, Bouley J, Tuncali I, Dogan EO, Pereira M, Luu T, et al. Repeated mild
traumatic brain injury triggers pathology in asymptomatic C9ORF72 transgenic mice. Brain.
2023;146(12):5139-52.

41.   Wiesner D, Tar L, Linkus B, Chandrasekar A, Olde Heuvel F, Dupuis L, et al. Reversible
induction of TDP-43 granules in cortical neurons after traumatic injury. Exp Neurol. 2018;299(Pt
A):15-25.

42.   Stein TD, Alvarez VE, McKee AC. Chronic traumatic encephalopathy: a spectrum of
neuropathological changes following repetitive brain trauma in athletes and military personnel.
Alzheimers Res Ther. 2014;6(1):4.

43.   Henninger N, Bouley J, Sikoglu EM, An J, Moore CM, King JA, et al. Attenuated
traumatic axonal injury and improved functional outcome after traumatic brain injury in mice
lacking Sarm1. Brain. 2016;139(Pt 4):1094-105.

44.   White MA, Kim E, Duffy A, Adalbert R, Phillips BU, Peters OM, et al. TDP-43 gains
function due to perturbed autoregulation in a Tardbp knock-in mouse model of ALS-FTD. Nat
Neurosci. 2018;21(4):552-63.

45.   Hasan R, Humphrey J, Bettencourt C, Newcombe J, Consortium NA, Lashley T, et al.
Transcriptomic analysis of frontotemporal lobar degeneration with TDP-43 pathology reveals
cellular alterations across multiple brain regions. Acta Neuropathol. 2022;143(3):383-401.

46.   Humphrey J, Venkatesh S, Hasan R, Herb JT, de Paiva Lopes K, Kucukali F, et al.
Integrative transcriptomic analysis of the amyotrophic lateral sclerosis spinal cord implicates
glial activation and suggests new risk genes. Nat Neurosci. 2023;26(1):150-62.

47.   Neelagandan N, Gonnella G, Dang S, Janiesch PC, Miller KK, Kuchler K, et al. TDP-43
enhances translation of specific mRNAs linked to neurodegenerative disease. Nucleic Acids
Res. 2019;47(1):341-61.

48.   Russo A, Scardigli R, La Regina F, Murray ME, Romano N, Dickson DW, et al.
Increased cytoplasmic TDP-43 reduces global protein synthesis by interacting with RACK1 on
polyribosomes. Human molecular genetics. 2017;26(8):1407-18.

49.   Bjork RT, Mortimore NP, Loganathan S, Zarnescu DC. Dysregulation of Translation in
TDP-43 Proteinopathies: Deficits in the RNA Supply Chain and Local Protein Production. Front
Neurosci. 2022;16:840357.

50.   Piol D, Robberechts T, Da Cruz S. Lost in local translation: TDP-43 and FUS in
axonal/neuromuscular junction maintenance and dysregulation in amyotrophic lateral sclerosis.
Neuron. 2023;111(9):1355-80.

51.   Zhong J, Gunner G, Henninger N, Schafer DP, Bosco DA. Intravital Imaging of
Fluorescent Protein Expression in Mice with a Closed-Skull Traumatic Brain Injury and Cranial
Window Using a Two-Photon Microscope. J Vis Exp. 2023(194).

52.   Dogan EO, Bouley J, Zhong J, Harkins AL, Keeler AM, Bosco DA, et al. Genetic ablation
of Sarm1 attenuates expression and mislocalization of phosphorylated TDP-43 after mouse
repetitive traumatic brain injury. Acta Neuropathol Commun. 2023;11(1):206.

53.   Flierl MA, Stahel PF, Beauchamp KM, Morgan SJ, Smith WR, Shohami E. Mouse closed
head injury model induced by a weight-drop device. Nat Protoc. 2009;4(9):1328-37.

54.   Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 2019;37(8):907-15.

55.   Chen Y, Lun AT, Smyth GK. From reads to genes to pathways: differential expression
analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline.
F1000Res. 2016;5:1438.

56.   Castanza AS, Recla JM, Eby D, Thorvaldsdottir H, Bult CJ, Mesirov JP. Extending
support for mouse data in the Molecular Signatures Database (MSigDB). Nat Methods.
2023;20(11):1619-20.

57.   Liberzon A, Birger C, Thorvaldsdottir H, Ghandi M, Mesirov JP, Tamayo P. The
Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst.
2015;1(6):417-25.

58.   Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP.
Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011;27(12):1739-40.

59.   Mootha VK, Lindgren CM, Eriksson KF, Subramanian A, Sihag S, Lehar J, et al. PGC-
1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated
in human diabetes. Nat Genet. 2003;34(3):267-73.

60.   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene
set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. Proc Natl Acad Sci U S A. 2005;102(43):15545-50.

61.   Funes S, Jung J, Gadd DH, Mosqueda M, Zhong J, Shankaracharya, et al. Expression
of ALS-PFN1 impairs vesicular degradation in iPSC-derived microglia. Nat Commun.
2024;15(1):2497.

62.   Tyanova S, Temu T, Cox J. The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. Nat Protoc. 2016;11(12):2301-19.

63. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, et al. The Perseus
computational platform for comprehensive analysis of (prote)omics data. Nat Methods.
2016;13(9):731-40.

64. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.

65. Kamentsky L, Jones TR, Fraser A, Bray MA, Logan DJ, Madden KL, et al. Improved
structure, function and compatibility for CellProfiler: modular high-throughput image analysis
software. Bioinformatics. 2011;27(8):1179-80.

66. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly
accurate protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9.

67. Darnell JC, Van Driesche SJ, Zhang C, Hung KY, Mele A, Fraser CE, et al. FMRP stalls
ribosomal translocation on mRNAs linked to synaptic function and autism. Cell.
2011;146(2):247-61.

68. Dogan EO, Simonini SR, Bouley J, Weiss A, Brown RH, Jr., Henninger N. Genetic
Ablation of Sarm1 Mitigates Disease Acceleration after Traumatic Brain Injury in the
SOD1(G93A) Transgenic Mouse Model of Amyotrophic Lateral Sclerosis. Ann Neurol. 2025.

69. Curzon P, Rustay NR, Browman KE. Cued and Contextual Fear Conditioning for
Rodents. 2nd ed: CRC Press/Taylor & Francis, Boca Raton (FL); 2009 2009.

70. Maren S, Quirk GJ. Neuronal signalling of fear memory. Nat Rev Neurosci.
2004;5(11):844-52.

71. Querfurth H, Lee HK. Mammalian/mechanistic target of rapamycin (mTOR) complexes in
neurodegeneration. Mol Neurodegener. 2021;16(1):44.

72. Simon DW, McGeachy MJ, Bayir H, Clark RS, Loane DJ, Kochanek PM. The far-
reaching scope of neuroinflammation after traumatic brain injury. Nat Rev Neurol.
2017;13(3):171-91.

73. Hubbard WB, Joseph B, Spry M, Vekaria HJ, Saatman KE, Sullivan PG. Acute
Mitochondrial Impairment Underlies Prolonged Cellular Dysfunction after Repeated Mild
Traumatic Brain Injuries. J Neurotrauma. 2019;36(8):1252-63.

74. Wang W, Wang L, Lu J, Siedlak SL, Fujioka H, Liang J, et al. The inhibition of TDP-43
mitochondrial localization blocks its neuronal toxicity. Nat Med. 2016;22(8):869-78.

75. Yoshida H, Matsui T, Yamamoto A, Okada T, Mori K. XBP1 mRNA is induced by ATF6
and spliced by IRE1 in response to ER stress to produce a highly active transcription factor.
Cell. 2001;107(7):881-91.

76. Pineda SS, Lee H, Ulloa-Navas MJ, Linville RM, Garcia FJ, Galani K, et al. Single-cell
dissection of the human motor and prefrontal cortices in ALS and FTLD. Cell.
2024;187(8):1971-89 e16.

77. Perucho L, Artero-Castro A, Guerrero S, Ramon y Cajal S, ME LL, Wang ZQ. RPLP1, a
crucial ribosomal protein for embryonic development of the nervous system. PLoS One.
2014;9(6):e99956.

78. Moisse K, Volkening K, Leystra-Lantz C, Welch I, Hill T, Strong MJ. Divergent patterns
of cytosolic TDP-43 and neuronal progranulin expression following axotomy: implications for
TDP-43 in the physiological response to neuronal injury. Brain Res. 2009;1249:202-11.

79. Mehta PR, Brown AL, Ward ME, Fratta P. The era of cryptic exons: implications for ALS-
FTD. Mol Neurodegener. 2023;18(1):16.

80. Rouaux C, Gonzalez De Aguilar JL, Dupuis L. Unmasking the skiptic task of TDP-43.
EMBO J. 2018;37(11).

81. Arnold ES, Ling SC, Huelga SC, Lagier-Tourenne C, Polymenidou M, Ditsworth D, et al.
ALS-linked TDP-43 mutations produce aberrant RNA splicing and adult-onset motor neuron
disease without aggregation or loss of nuclear TDP-43. Proc Natl Acad Sci U S A.
2013;110(8):E736-45.

82. Fratta P, Sivakumar P, Humphrey J, Lo K, Ricketts T, Oliveira H, et al. Mice with
endogenous TDP-43 mutations exhibit gain of splicing function and characteristics of
amyotrophic lateral sclerosis. EMBO J. 2018;37(11).

83. Polymenidou M, Lagier-Tourenne C, Hutt KR, Huelga SC, Moran J, Liang TY, et al. Long
pre-mRNA depletion and RNA missplicing contribute to neuronal vulnerability from loss of TDP-
43. Nat Neurosci. 2011;14(4):459-68.

84. Yang C, Qiao T, Yu J, Wang H, Guo Y, Salameh J, et al. Low-level overexpression of
wild type TDP-43 causes late-onset, progressive neurodegeneration and paralysis in mice.
PLoS One. 2022;17(2):e0255710.

85. Chu BW, Kovary KM, Guillaume J, Chen LC, Teruel MN, Wandless TJ. The E3 ubiquitin
ligase UBE3C enhances proteasome processivity by ubiquitinating partially proteolyzed
substrates. J Biol Chem. 2013;288(48):34575-87.

86. Gaetani M, Zubarev RA. Proteome Integral Solubility Alteration (PISA) for High-
Throughput Ligand Target Deconvolution with Increased Statistical Significance and Reduced
Sample Amount. Methods Mol Biol. 2023;2554:91-106.

87. Li J, Van Vranken JG, Paulo JA, Huttlin EL, Gygi SP. Selection of Heating Temperatures
Improves the Sensitivity of the Proteome Integral Solubility Alteration Assay. J Proteome Res.
2020;19(5):2159-66.

88. Savitski MM, Reinhard FB, Franken H, Werner T, Savitski MF, Eberhard D, et al.
Tracking cancer drugs in living cells by thermal profiling of the proteome. Science.
2014;346(6205):1255784.

89. Brohee S, van Helden J. Evaluation of clustering algorithms for protein-protein
interaction networks. BMC Bioinformatics. 2006;7:488.

90. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The
STRING database in 2023: protein-protein association networks and functional enrichment
analyses for any sequenced genome of interest. Nucleic Acids Res. 2023;51(D1):D638-D46.

91. Cheng F, Chapman T, Venturato J, Davidson JM, Polido SA, Rosa-Fernandes L, et al.
Proteomics Analysis of the TDP-43 Interactome in Cellular Models of ALS Pathogenesis. J
Neurochem. 2025;169(5):e70079.

92. Freibaum BD, Chitta RK, High AA, Taylor JP. Global analysis of TDP-43 interacting
proteins reveals strong association with RNA splicing and translation machinery. J Proteome
Res. 2010;9(2):1104-20.

93. Zeng C, Han S, Pan Y, Huang Z, Zhang B, Zhang B. Revisiting the chaperonin T-
complex protein-1 ring complex in human health and disease: A proteostasis modulator and
beyond. Clin Transl Med. 2024;14(2):e1592.

94. Luo J, Zhao H, Chen L, Liu M. Multifaceted functions of RPS27a: An unconventional
ribosomal protein. J Cell Physiol. 2023;238(3):485-97.

95. Lin Z, Kim E, Ahmed M, Han G, Simmons C, Redhead Y, et al. MRI-guided histology of
TDP-43 knock-in mice implicates parvalbumin interneuron loss, impaired neurogenesis and
aberrant neurodevelopment in amyotrophic lateral sclerosis-frontotemporal dementia. Brain
Commun. 2021;3(2):fcab114.

96. Kelmer Sacramento E, Kirkpatrick JM, Mazzetto M, Baumgart M, Bartolome A, Di Sanzo
S, et al. Reduced proteasome activity in the aging brain results in ribosome stoichiometry loss
and aggregation. Mol Syst Biol. 2020;16(6):e9596.

97. Amirbeigiarab S, Kiani P, Velazquez Sanchez A, Krisp C, Kazantsev A, Fester L, et al.
Invariable stoichiometry of ribosomal proteins in mouse brain tissues with aging. Proc Natl Acad
Sci U S A. 2019;116(45):22567-72.

98. Zhao B, Cowan CM, Coutts JA, Christy DD, Saraph A, Hsueh SCC, et al. Targeting
RACK1 to alleviate TDP-43 and FUS proteinopathy-mediated suppression of protein translation
and neurodegeneration. Acta Neuropathol Commun. 2023;11(1):200.

99. Bhavsar RB, Makley LN, Tsonis PA. The other lives of ribosomal proteins. Hum
Genomics. 2010;4(5):327-44.

100. Simsek D, Barna M. An emerging role for the ribosome as a nexus for post-translational
modifications. Curr Opin Cell Biol. 2017;45:92-101.

101. Ali A, Garde R, Schaffer OC, Bard JAM, Husain K, Kik SK, et al. Adaptive preservation
of orphan ribosomal proteins in chaperone-dispersed condensates. Nat Cell Biol.
2023;25(11):1691-703.

102. Nagano S, Jinno J, Abdelhamid RF, Jin Y, Shibata M, Watanabe S, et al. TDP-43
transports ribosomal protein mRNA to regulate axonal local translation in neuronal axons. Acta
Neuropathol. 2020;140(5):695-713.

103. Tank EM, Figueroa-Romero C, Hinder LM, Bedi K, Archbold HC, Li X, et al. Abnormal
RNA stability in amyotrophic lateral sclerosis. Nat Commun. 2018;9(1):2845.

104. Chou A, Krukowski K, Jopson T, Zhu PJ, Costa-Mattioli M, Walter P, et al. Inhibition of
the integrated stress response reverses cognitive deficits after traumatic brain injury. Proc Natl
Acad Sci U S A. 2017;114(31):E6420-E6.

105. Huang WP, Ellis BCS, Hodgson RE, Sanchez Avila A, Kumar V, Rayment J, et al.
Stress-induced TDP-43 nuclear condensation causes splicing loss of function and STMN2
depletion. Cell Rep. 2024;43(7):114421.

106. Rifai OM, Waldron FM, O'Shaughnessy J, Read FL, Gilodi M, Pastore A, et al. Amygdala
TDP-43 pathology is associated with behavioural dysfunction and ferritin accumulation in
amyotrophic lateral sclerosis. bioRxiv. 2024.

107. Wilkins OG, Chien M, Wlaschin JJ, Barattucci S, Harley P, Mattedi F, et al. Creation of
de novo cryptic splicing for ALS and FTD precision medicine. Science. 2024;386(6717):61-9.

108. Francois-Moutal L, Perez-Miller S, Scott DD, Miranda VG, Mollasalehi N, Khanna M.
Structural Insights Into TDP-43 and Effects of Post-translational Modifications. Front Mol
Neurosci. 2019;12:301.

109. Shenouda M, Xiao S, MacNair L, Lau A, Robertson J. A C-Terminally Truncated TDP-43
Splice Isoform Exhibits Neuronal Specific Cytoplasmic Aggregation and Contributes to TDP-43
Pathology in ALS. Front Neurosci. 2022;16:868556.

110. Elliott E, Bailey O, Waldron FM, Hardingham GE, Chandran S, Gregory JM. Therapeutic
Targeting of Proteostasis in Amyotrophic Lateral Sclerosis-a Systematic Review and Meta-
Analysis of Preclinical Research. Front Neurosci. 2020;14:511.

1766