Core ideas:

1) Morphological nut characteristics are under highly polygenic control in American and

American-European interspecific hazelnuts.

2) Best linear unbiased predictors allow for accurate prediction of morphological nut

characteristics.

3) Marker density and training population design must be tailored to the sample population for

which predictions are being made.

Title: COMPOSITE INTERVAL MAPPING AND GENOMIC PREDICTION OF NUT

QUALITY TRAITS IN AMERICAN HAZELNUTS AND AMERICAN-EUROPEAN

INTERSPECIFIC HYBRIDS

Authors: Scott H. Brainard[1*], Julie C. Dawson[1]

Affiliations

[1] Department of Plant and Agroecosystem Sciences, University of Wisconsin-Madison, Madison,

WI 53706, USA

[*]corresponding author: shbrainard@wisc.edu

ABSTRACT

The native, perennial shrub American hazelnut (*Corylus americana*) is cultivated in the

Midwestern U.S. for its significant ecological benefits, as well as its high-value nut crop.

Perennial crops are a long-term breeding effort, and would benefit from the use of genetic data in

selection to reduce breeding cycle time. In addition, high-throughput phenotyping methods are

essentially to the efficient and accurate screening of large breeding populations. This study

reports novel advances in both of these domains, for American (*C. americana*) and interspecific

hybrids between European (*C. avellana*) and American hazelnuts. Two populations of hazelnuts,

one composed of *C. americana* and one composed of *C. americana* x *C. avellana* hybrids, were

phenotyped over the course of two years in two locations using a digital imagery-based method

for quantifying morphological nut and kernel traits. This data was used to perform composite

interval mapping (CIM) using a recently released genetic map, and genomic prediction using a

newly-available chromosome-scale reference genome for *C. americana*. Multiple QTL were

detected for all traits analyzed, with an average total R^2 of 52%. Genomic prediction exhibited

high accuracy, with an average correlation coefficient between genotypic values and phenotypic

observations of 0.78 across both environments. These results suggest that incorporating genetic

data in selection is a tenable method for improving genetic gain for highly-polygenic traits in

hazelnut breeding programs.

Keywords

Corylus americana, *Corylus avellana*, hazelnut, quantitative genetics, genomic prediction,

linkage mapping, composite interval mapping, digital phenotyping

INTRODUCTION

Hazelnuts (*Corylus* spp.) are a globally significant nut crop, with annual production of over 1.3

million tons, produced across 34 countries (FAOSTAT, 2023). Currently, this cultivation is

limited to areas with climatic conditions approximating the Mediterranean region in which the

dominant cultivars of *C. avellana* were bred (Mehlenbacher and Molnar, 2021). The narrow

range of cultivars has limited U.S. production of hazelnuts to the Willamette Valley in Oregon,

where over 95% of U.S. acreage is situated (USDA Economic Research Service, 2025). In this

context, the native species *C. americana*, which is endemic and well-adapted to much of the

eastern U.S., represents a valuable source of genetic diversity for expanding hazelnut production.

American hazelnut can exhibit vigorous growth in coldenvironments, and robust resistance to

endemic pathogens such as the eastern filbert blight (EFB, *Anisogramma anomala*) (Molnar et

al., 2018; Revord et al., 2020).

The genetic improvement of hazelnut is limited by its long generation time. A seedling hazelnut

will not typically flower under field conditions for at least three years, and many of the most

agronomically important traits (e.g., total kernel yield and bush architecture) require additional

time to accurately phenotype. Mature plant yields will often not be fully apparent for a decade,

and multi-year harvest data are necessary to precisely quantify such traits. Methods that can

make parental and progeny selection decisions more efficient are therefore extremely valuable

for a perennial species such as hazelnut, by accelerating this traditional breeding process.

Quantitative genetic tools are essential in this regard. The identification of QTL is essential to

the implementation of marker-assisted selection, while genomic prediction can dramatically

accelerate genetic gain for polygenic traits (Würschum, 2012; Heslot et al., 2015). The recent

development of genomic resources for *C. americana* have made both of these approaches

accessible to breeding programs. First, a genetic map constructed using an F_1 progeny family

descended from a cross between Oregon and Midwestern hazelnut varieties 'Jefferson' and

'Eric4-21', respectively, now permits linkage map-based approaches to detecting QTL (Brainard

et al., 2023). Such interspecific populations are highly relevant to Midwest breeding efforts, as

C. avellana remains a valuable source of diversity for nut quality-related traits. The use of

linkage mapping populations is well-suited to such populations, where the contrasting

phenotypes of parental varieties frequently produces high degrees of segregation in progeny

families. In addition, chromosome-scale genome assemblies for the *C. americana* selections

'Rush' and 'Winkler' also now allow for the efficient identification of polymorphisms for use in

genomic selection models (Brainard et al., 2024).   In this study, a diverse collection of wild *C.*

americana was utilized to test such a method. Diversity panels such as this are useful for

assessing highly polygenic traits, and developing methods for American hazelnut are

appropriate, as this native species not only has potential as a crop for the Upper Midwest, but is

also a useful source of traits such as resistance to Eastern Filbert Blight, and cold hardiness.

This study uses both of these tools, in combination with a novel digital imagery-based

phenotyping pipeline for precisely quantifying morphological characteristics of inshell hazelnuts

and kernels. Recent developments in contour analysis of digital images has led to numerous

methods for quantifying the sizes and shapes of plant structures (Hameed et al., 2018). In this

study, we present the design and demonstration of such a platform that quantifies the size and

shape features of hazelnuts, and validate its performance using two experimental populations: a

diversity panel composed of a wild population of *C. americana* sourced from the Wisconsin

Department of Natural Resources, and three F_1 interspecific biparental progeny families

generated by the University of Minnesota hazelnut breeding program.

MATERIALS AND METHODS

Plant Materials

Two populations of seedling hazelnut plants were used in this study. One is composed of 473

wild *C. americana* seedlings sourced from the Wisconsin Department of Natural Resources and

planted in 2016 at a farm outside of Barneveld, Wisconsin (43.043266 N, -89.926068 W). This

population was treated as a "diversity panel", and used to fit and validate genomic prediction

models that could be used in introgressing valuable genetic diversity from *C. americana* into

Midwest breeding programs. The second is 258 plants which comprise three F_1 families

descended from crosses between clonal varieties sourced from the University of Minnesota and

Oregon State University, specifically: Eric4-21 x Jefferson ("Eric\Jeff"), Gibs5-15 x OSU-919-

031 ("Gibs\OSU"), and Gibs5-15 x York ("Gibs\York"). These were planted at a University of

Minnesota research farm in Rosemount, MN (44.695784 N, -93.079196 W) in 2015 and 2016.

These represented the biparental families that were used to test both genomic prediction in

breeding populations that include *C. avellana*, as well as perform F_1 linkage mapping.

Phenotyping

Morphological characteristics of in-shell hazelnuts and shelled kernels were used as the trait data

for this study. This was collected primarily using an adapted version of the digital imagery

acquisition and analysis pipeline reported by (Brainard et al., 2021) (Fig. 1). In brief, bushes

114 were completely harvested by hand in August, 2020 and August 2021. Harvested clusters were

115 dried in the greenhouse and husked. A sample of 30 in-shell nuts were randomly sampled per

116 bush, and weighed in bulk. These nuts were arranged on a 6x5 grid with a QR code, and a Nikon

117 5600 DSLR camera tethered to a desktop computer was used to acquire a single image. The

118 OpenCV Python library was used to isolate each in-shell nut, and produce a binary mask by

119 applying a fixed HSV threshold to each pixel. An ellipse was then fit to each binary mask, and

120 the length of its major and minor axis was calculated by converting pixel length to physical

121 distance using a scale bar embedded in each image. Circularity of the nut was calculated as the

122 ratio of these two lengths. Each nut was individually cracked, and the kernel was then returned

123 to the grid, preserving the original arrangement of the nuts. A second photo was acquired, and

124 the same traits were calculated. Finally, the kernels were weighed in bulk. This allowed for both

125 a volumetric and gravimetric estimation of the percent kernel for each nut sampled. Python

126 scripts for the image acquisition, processing, phenotyping, and file management are available at:

127 https://github.com/shbrainard/hazelnut-phenotyping. Phenotypic measurements were averaged

128 for each individual plant. The prcomp R function was used to perform PCA on this phenotypic

129 data, and the ggplot2 function autoplot was used to visualize the relationship between these

130 traits.

132 **Sequencing**

133 Roughly 1 cm^2 of leaf tissue was sampled from each bush in May, 2021, immediately following

134 budbreak. Tissue was sampled into 96-well Qiagen Collection Microtubes (Qiagen N.V., Venlo,

135 The Netherlands), and lyophilized using a Labconco 18 L freeze dryer set to 0.004mBar for 72

136 hours. Freeze-dried tissue was then macerated. DNA extraction, quantification, library

preparation, and sequencing was performed at the University of Wisconsin-Madison

Biotechnology Center. Libraries were prepared for genotyping-by-sequencing using a double

digestion with the restriction enzymes *NsiI* and *BfaI* following the methodology described by

(Elshire et al., 2011). This combination was pre-selected based on an analysis of k-mer

distributions of various enzyme digestions, where *NsiI/BfaI* was observed to maximize the k-mer

diversity of the library. Illumina adapters and sample-specific barcodes were then annealed.

Samples were multiplexed, and paired-end 150-bp sequence data was generated using an

Illumina NovaSeq 6000, with an average of 10 million reads per sample. Trimming and

demultiplexing of raw Illumina reads was performed using a custom Java application

https://github.com/shbrainard/gbsTools. Reads were aligned to the *C. americana* genome for

'Winkler' (Brainard et al., 2024).

SNP calling

Haplotype-based markers were called using Stacks 2 (Rochette et al., 2019), which can identify

multi-allelic markers using the phased nature of multiple indels or SNPs that appear within a

single 150-bp paired-end read. Because such markers cannot be directly filtered for depth, the

parameter 'gt-alpha' was increased to 0.01 as a method for ensuring genotype quality. Markers

were then filtered for linkage disequilibrium using bcftools (prune plugin with a window size of

10kb, and r^2 filter of 0.95, retaining one site per window based on highest allele frequency). This

generated a set of 78,079 markers, with an average of ~7,000 per chromosome.

Biallelic SNPs were called using the TASSEL GBSv2 pipeline (Bradbury et al., 2007).

1,161,144 called SNPs were first filtered to exclude sites with minor allele frequency < 0.05,

more than one alternate allele, or greater than 10% missingness across all samples, which

reduced the number of SNPs to 878,306. This set was then further reduced to include only SNPs

where the 80^{th} percentile of allele depth across all samples was > 8, leaving 102,096 sites. Next,

filtering was performed for linkage disequilibrium ($r^2 < 0.75$, with the same selection criteria

above), which retained 53,696 SNPs. Finally, markers were then subset to only include those

which were retained in both the *C. americana* diversity panel in Wisconsin and interspecific

biparental populations in Minnesota, leaving 44,961 SNPs. All filtering was performed using

bcftools (Danecek et al., 2021).

Linkage map construction and composite interval mapping

Since the interspecific biparental populations in Minnesota were constructed from controlled

crosses between known hazelnut varieties, it was possible to build a genetic map by using the R

package onemap (Margarido et al., 2007) (https://github.com/augusto-garcia/onemap). This map

was previously reported in (Brainard et al., 2023). Briefly, markers called using Stacks 2 were

first filtered to include only those of segregation types A1, A2 and B3.7 (following the notation

of (Wu et al., 2002)), such that only markers with either three or four alleles remained. In these

segregation types, the four possible alleles present in a biparental cross between two outbred

diploid parents are represented as *a, b, c,* and *d*. A1 segregation types are loci for which the

parental genotypes are *ab* x *cd*, A2 represents *ab* x *ac* crosses, and B3.7 represents *ab* x *ab*.

Next, markers for which more than 5% of all samples had no called genotype were removed and

two-point recombination frequencies were calculated for all possible phase configurations

between all remaining markers using maximum likelihood. Maximum likelihood estimates were

able to fully resolve phase between pairs of markers, due to the fully-informative segregation

types that were used. A hierarchical clustering algorithm was utilized to construct linkage

groups, and markers were ordered and phased within these groups by using a Hidden Markov

Model with an error rate of 0.05. Recombination frequencies were finally converted to genetic

distances using the Kosambi mapping function (Kosambi, 1943). Finally, this map was imported

into the R package fullsibQTL (Gazaffi et al., 2020) (https://github.com/augusto-

garcia/fullsibQTL) which was used to perform composite interval mapping. Cofactors were

selected to minimize the Akaike Information Criteria, and composite interval mapping was

employed to perform a single QTL scan for each of the phenotypes measured using the digital

image analysis protocol described above. Permutation tests were calculated to empirically

estimate a LOD threshold representing a 0.05 genome-wide p-value. Additive and dominance

effects were estimated for QTL exceeding this threshold using the function cim_char(). Least

squares estimation was used to calculate the percent variance explained both for both these QTL

individually, and collectively, using the function $r2\_ls$. This R^2 is calculated using the residual

sums of squares (RSS) for the null and full models:

$$R^2 = \frac{RSS.null - RSS.full}{RSS.null}$$

Calculation of GEBVs

In order calculate genomic-estimated breeding values from the biallelic SNP dataset described

above, the R package StageWise (Endelman, 2023) (https://github.com/jendelman/StageWise)

was used to compute variance components and best linear unbiased predictors (BLUPs) of

additive genetic value. This software is designed to perform two-stage analysis, by first

computing best linear unbiased estimators (BLUEs) for each genotype using a specified

experimental design. Since the genotypes in both populations were comprised of unreplicated

seedlings, a fixed-effects linear model was used to first compute BLUEs for each genotype

where:

$$Y_{ij} = \mu + G_i + Y_j + \varepsilon_{ij}$$

G_i represents the i^{th} genotype effect, Y_j the j^{th} year effect, and ε_{ij} the residual variance (which

included the $G * Y$ effect), with $\varepsilon_{ij} \sim N(0, \sigma_\varepsilon^2)$). Models were fit for each trait independently,

and then passed to the function *Stage2* in order to estimate genotypic values for each trait

individually. Additive genetic values are here assumed to follow a multivariate normal

distribution with a variance-covariance matrix proportional to the G matrix (VanRaden, 2008),

where a centered marker of allele dosages **W** is normalized by the product of the major and

minor allele frequencies (*p* and *q*) for the k^{th} marker:

$$G = \frac{WW^T}{2 \sum_k p_k q_k}$$

A baseline for the reliability of genomic estimated genetic values was obtained by predicting the

genetic value for all samples, using the full phenotypic and genotypic datasets available. This

form of marker-assisted prediction provides a point of comparison for testing different

parameters of the prediction model.

Reliabilities were calculated as both the mean r^2 returned by StageWise (here denoted as

"Stagewise- r^2"), which is the squared correlation between the true and predicted values

assuming the model is correct. This is a model parameter that is proportional to the prediction

error variance (PEV) of the BLUP. Correlation coefficient between the predicted genotypic

values and BLUEs for each genotype (calculated on the basis of the observed phenotypes) are

also reported (here denoted as "Pearson-r^2").

Marker density evaluation

In order to assess the impact of number of SNPs used in the estimation of the variance-

covariance "G" matrix by StageWise, the input marker matrix was randomly downsampled to

simulate densities ranging from 500 to 45,000 markers. The resulting G matrix was then

subtracted from G matrix calculated using the complete set of markers in order to estimate the

ability of smaller marker sets to accurately capture the degree of relatedness between all

individuals in each population.

Validation analyses were performed to assess the impact of marker density on the reliability of

the predictions. Marker density was varied from 500 to 45,000, in 100 marker increments, in

order to model the impact of the number of SNPs on prediction accuracy. This was assessed

directly by calculating the difference between the G matrix calculated with the full set of

markers, and the reduced set, as well as by modeling trait prediction accuracy at different marker

densities. At each marker density, twenty random subsets of the marker matrix were produced,

with even distribution of markers across the 11 chromosomes.

Population relatedness

The impact of the degree of relatedness between the training and validation population when

performing marker-based selection was also assessed, using a leave-one-out cross-validation

approach. In the *C. americana* population, this was done by first sorting the population by the

degree of relatedness of each individual in the population to the validation individual, then using

a 100-indivdual subset of the whole population as the training population, which was

progressively designed to be less related to the validation individual (i.e., the 1st-100th most

related individuals, the 2^{nd}-101^{st} most related individuals, the 3^{rd}-102^{nd} most related individuals,

etc.). Once an asymptotically-minimal prediction accuracy was identified, a secondary analysis

was performed, by progressively expanding the training population starting from the 20^{th}-121^{st}

most-related individual, adding in increasing numbers of less-related individuals. This allowed

for an assessment of whether training population size could compensate for a lack of training

population relatedness.

In the interspecific biparental populations, the known pedigrees of the families provided an

alternative method for assessing the impact of relatedness on accuracy. These analyses were

performed for each of the three families individually, assessing prediction accuracy by predicting

the genotypic value each member of each of the three full-sib families, using each of the three

families as the training population.

RESULTS

Morphological inshell and kernel characteristics

Histograms for six traits, inshell height and width, kernel height and width, and percent kernel,

measured both volumetrically and gravimetrically, are shown in Fig. 2. In general, the

interspecific biparental populations, being composed of crosses with *C. avellana* pollen parents

from the Oregon State University breeding program, have larger inshell and kernel dimensions.

However, the distribution of percent kernel was observed to be either nearly identical (when

measured gravimetrically) or larger in the *C. americana* diversity panel (when measured

volumetrically from the digital images).

Fig. 3 presents PCA biplots for the *C. americana* population (Fig. 3A), interspecific biparental

populations (Fig. 3B) and the combined set of all individuals (Fig. 3C), illustrating the close

correlation between inshell and kernel traits, and their relative independence from both measures

of percent kernel.

Composite interval mapping

The use of multialleic markers in the onemap pipeline produced a high-quality genetic map.

Map quality was confirmed by assessing inflation. Given the relatively small size of hazelnut

chromosomes (with an average physical size of 29.78 Mb), a given linkage group should not be

longer than 100 cM (under normal biological assumptions and a standard mapping function).

Inflated map lengths in excess of this upper bound are nevertheless frequently observed by not

controlling for genotyping errors, which increase estimates of recombination frequency. Using

multi-point regression when converting genetic distances to the final genetic map counteracts

this tendency. The linkage groups in our final map are no longer than 89.9 cM, with an average

length of 66.3 cM, and a total length of 729.6 cM.

Single QTL scans were performed for each of six phenotypic traits. LOD profiles across the

genetic map are shown in Fig 4. Table 1 summarizes the results of the QTL identified via

composite interval mapping for each of the six traits. For the two inshell traits, four QTL were

identified (with only one shared in common between both traits), with an average total R^2 of

52%. For the two kernel-specific measures, three QTL were detected for height, while four were

detected for length (with no overlap between the two). In both cases, lower total R^2 was

observed, with an average of 28.6%. The two methods for measuring percent kernel similarly

were associated with three (gravimetric) and four (volumetric) unique QTL, with a higher total

average R^2 of 41.1%

Genomic prediction accuracy

Table 1 presents the heritabilities calculated by StageWise when BLUPs are calculated with all

phenotypic and genotypic data for each population. This is equivalent to the proportion of

variance attributable to the genotypes.

Two measures of predication accuracy are presented in Table 2. These are the average

StageWise-r^2 across all individuals in each population, and the Pearson-r^2, calculated as the

correlation between predicted genetic values and phenotypes for all individuals in each

population. In the *C. americana* diversity panel, the average StageWise-r^2 across all traits was

0.56, while the average Pearson-r^2 was 0.88. In the interspecific biparental populations, these

two measures were closer to one another: average StageWise-r^2 was 0.68, and the average

Pearson-r^2 was 0.67.

Effect of marker density on prediction accuracy

The results of the marker density analysis are shown in Fig 5., where the y-axis represents the

absolute value of the sum of the matrix resulting from the subtraction of the downsampling-

derived G matrix. In both populations, an asymptotic minimum in the difference between the

matrices is reached between 5,000 and 10,000 markers.

A similar pattern was observed for average StageWise-r^2 values, where there was a rapid

increase in mean StageWise-r^2 as marker density increased, until an asymptotic maximum

StageWise-r^2 was attained between 5,000 and 10,000 markers (Supp. Fig. 1) , further suggesting

that a marker density in this range will most efficiently estimate relatedness between individuals.

When all individuals were used in the calculation of BLUPs, the difference was relatively minor,

representing an increase of only ~0.04 in the case of the *C. americana* diversity population. In

the interspecific biparental populations, the difference is almost entirely negligible: while the

exponential relationship is evident, the difference between the minimum and maximum

reliability is ~0.01.

Impact of relatedness on prediction accuracy

Results from the comparison of variably-related training populations are shown in Fig 6.

Prediction accuracy quickly drops as the most closely related individuals are removed from the

training population, leading to asymptotically-minimal accuracies when the twenty most closely

related individuals are removed. This reduction in accuracy represented a nearly 55% reduction

in relatedness (measured by the average of the coefficients of the corresponding elements of the

G matrix). However, even at maximal relatedness, prediction accuracy in the *C. americana*

diversity population was relatively low, with maximum Pearson-r^2 values of ~0.45, and an

average of 0.3.

The ability to compensate for reduced prediction accuracy by increasing the size of the training

population was subsequently assessed. The results of this analysis are shown in Fig. 6. For the

two measures of percent kernel (which also exhibited the lowest prediction accuracy in general),

this increase in population size had no impact on prediction accuracy, while for inshell height,

the reduction in relatedness associated with increasing population size led to a reduction in

prediction accuracy. Finally, for kernel height, kernel length, and inshell length, prediction

accuracy slightly increased, however all of these changes in prediction accuracy represented less

than 0.25% in total variation. This indicates that if the 20 most-closely related individuals are

absent from the training population design, this cannot be substantially offset by simply

including more weakly-related individuals. Importantly, the average degree of relatedness in the

C. americana panel is relatively low, and thus this observation is not necessarily generalizable to

all other training population designs.

In the interspecific biparental populations, similar results were observed when reducing the

relatedness of the training population, and keeping the total size of the population constant at 100

individuals Supp. Fig. 2. However, there are limitations to this approach for these populations.

Because of the structured nature of the full-sib populations, correlations between the genotypic

values and phenotypic data across all individuals primarily reflects this structure. These results

are shown in Table 3. Interestingly, marker-based prediction led to low (near-zero or negative)

Pearson-r^2 values, when the training population represented the same interspecific biparental

cross from which the sample being validated was drawn. For each of the families, however,

there was one other biparental population which led to positive prediction accuracies, although

these maximum Pearson-r^2 values were significantly lower than the maximum values observed in

the diversity panel.

DISCUSSION

Morphological nut traits exhibit high degree of polygenicity

The composite interval mapping results reported above demonstrate clearly that the traits

measured here related to physical characteristics of inshell nuts and kernels are highly polygenic.

On average, the identified QTL individually explained only 11% of the phenotypic variation for

a given trait. While for certain traits the total explained variance was much higher, the average

total R^2 for all QTL associated with a trait was only 40.5%, with the unexplained variance

ranging from 44.5% - 80.2%. A low total R^2 was particularly apparent for kernel traits, which

was also reflected in the prediction accuracy for these traits when evaluating genomic prediction

models using the interspecific biparental populations in Minnesota .

To our knowledge, this is the first study to ever map kernel and nut quality traits in an

interspecific *C. americana* x *C. avellana* family. Several studies have performed association

analysis in *C. avellana* populations of similar size. (Baytar et al., 2024) used similar GBS

marker data, and evaluated both kernel and nut size, as well as percent kernel. However, they

found only a single QTL for percent kernel. Another GWAS utilized a low density SSR map and

only found one QTL for kernel weight and none for percent kernel (Ozturk et al., 2017). Most

comparable to these results was a linkage mapping study performed using GBS markers and a

relatively large F_1 population derived from Italian *C. avellana* cultivars (Torello Marinoni et al.,

2018). This analysis included phenotypic data on both nut and kernel size, identifying one QTL

for kernel size (on LG 5) and four for nut size (on LGs 1, 2, 4, and 6). However, none of these

QTL explained more than 10% of the total phenotypic variance.

In light of these results, the CIM analyses reported here represent a significant advance in our

understanding of the genetic control of these physical nut characteristics, which is likely

attributable both to the significant variation observed in these interspecific crosses, the high

resolution of the genetic map used, and the precision and scale of the phenotyping method

utilized. Nevertheless, it appears likely that this study was underpowered to detect the numerous

remaining QTL associated with these traits. This is likely not due to a limitation in the number

of markers used, since the resolution of the map was greater than the number of expected

recombination events present in the population. Indeed, it is likely that the relatively small size

of the mapping population used in this study is primarily responsible for reduced statistical

power. Building larger F_1 populations would therefore be useful in attempting to more

accurately assess the true genetic architecture of these traits, however, the size of F_1 progeny

families are limited by *Corylus* biology. Multiparent QTL mapping would, in particular, be a

valuable approach (e.g., via a diallel mating design), as this would enable more precise

estimation of dominance effects. However, the present results point strongly towards many QTL

influencing these traits, and the absence of any single large effect QTL. As will be discussed

below, this motivates relying upon genomic prediction models as the preferred method for

utilizing marker information in applied breeding programs.

Discrepancies between C. americana diversity panel and interspecific biparental

populations

While the correlation coefficient-based estimates of prediction accuracy in the interspecific

biparental populations were nearly identical (but slightly less than) the mean Pearson-r^2 values,

correlation coefficients in the *C. americana* diversity population were dramatically higher than

the mean Pearson-r^2 values. This difference is so great that while mean Pearson-r^2 values were

higher in the interspecific biparental populations than in the *C. americana* diversity population

(higher for every trait other the kernel-specific measurements), correlation coefficients were

much higher in the *C. americana* diversity population than in the interspecific biparental

populations. Possible interpretations of these discrepancies, and how they should inform the

interpretation of these two metrics of reliability, are provided below. It should be noted at the

outset that because best linear predictors are designed to maximize the correlation between the

true and predicted genotypic values (and the covariance of the true and predicted values is equal

to the variance of the true values, reliability of the model is equivalent to broad sense heritability

as computed in a completely randomized design (Endelman, 2023). Correlation coefficients

derived from a leave-one-out cross-validation approach can introduce bias by assuming the

observed phenotypic data represents the "true" value of an individual (Zhou et al., 2017).

However, in general, it is clear that marker-assisted selection generated strikingly high

reliabilities, particularly for populations composed of unreplicated seedling genotypes. Except

for the kernel-specific traits in the interspecific biparental populations, both measures of

reliability were higher than 0.5. To our knowledge, this study represents the first reported

evaluation of genomic selection in any *Corylus* species, but these results are comparable to

assessments of prediction accuracy for morphological characteristics in other tree nut crops. For

example, (Nishio et al., 2018) reported prediction accuracies of 0.4 for the weight of chestnuts

using a GWAS model (BayesB).

There are numerous differences between the *C. americana* diversity population and the

interspecific biparental populations in Minnesota which likely underlie the differences in trait

heritabilities observed for these two locations. While spatial variation within the field locations

was not apparently greater in either the *C. americana* diversity population or the interspecific

biparental populations (Supp. Fig. 3), the fact that both trials were composed of unreplicated

seedling genotypes certainly lowered heritabilities, and unavoidably increased the relative size of

environmental variance components, potentially to differing degrees.

In addition, the populations themselves were obviously markedly different. While the

interspecific biparental populations were substantially smaller, they also exhibited much higher

degrees of average relatedness both within and across biparental populations. The *C. americana*

diversity population location was composed of wild-collected accessions with no known shared

pedigree. The interspecific biparental populations, on the other hand, was composed of full-sib

crosses in which maternal parents were both drawn from the Minnesota breeding program, and

paternal parents were selected form the Oregon breeding program. As a result, in most training

populations designed to test prediction accuracy in the interspecific biparental populations, the

average coefficients of the respective G matrix were roughly an order of magnitude higher than

in the *C. americana* diversity panel.

Excessive shrinkage within the interspecific biparental populations

The cross-validation results analyzing the impact of relatedness on prediction accuracy in the

interspecific biparental populations are striking, and are consistent across traits. It is not

immediately clear why prediction accuracies were consistently negative when the training

population was composed of the same full sib family as the individual for which the prediction

was being made. This is particularly surprising, given the relatively high accuracies (which for

certain traits, exceeded the asymptotic maximum prediction accuracy seen in the *C. americana*

diversity population) when the prediction population selected is **not** the full sib family from

which the given sample is drawn. Two important features of this analysis are likely impacting

these results. First, there is substantial shrinkage, much greater than that seen in the *C.*

americana diversity population, apparent across traits when performing these cross-validation

analyses Supp. Fig. 4. With small population sizes, and unreplicated genotypes, it has been

previously reported that shrinkage-based estimators can lower prediction accuracy for

unphenotyped individuals (Endelman and Jannink, 2012; Zhao et al., 2013).

Second, and relatedly, the population sizes for these comparisons across families is small,

ranging from 75-81 individuals being used in the given prediction populations. This is much

smaller than is typically considered in studies assessing factors influencing genomic prediction

accuracy, where significant drop-offs in accuracy are frequently observed when population size

drops below 100 (Tayeh et al., 2015; Zhang et al., 2017; Sverrisdóttir et al., 2018; Edwards et al.,

2019). As such, these were also smaller training populations than those used to analyze the

impact of relatedness in the *C. americana* diversity population. Generating larger biparental

families is biologically challenging in hazelnut, so breeding programs should adopt a strategy of

ensuring biparental families are interconnected through shared parentage, such that higher

effective numbers can be achieved in training populations.

Mean r^2 values vs. correlation coefficients as measures of reliability

The average of the r^2 values returned by the StageWise package and the Pearson correlation

coefficients between the genotypic values estimated by StageWise and the BLUEs represent two

intuitive methods for estimating reliability. The high values for reliability reported here (>0.5 in

nearly all cases) suggest relatively large broad-sense heritabilities i.e., large additive genetic

variance components, for morphological nut traits. This suggests that applying genomic

prediction methods within selection programs is realistic and tenable. There are, however,

several caveats to be aware of in evaluating these measures of reliability.

Any correlation-based measure of reliability will be inflated to a degree that is proportion to the

correlation between population structure and the traits of interest (Werner et al., 2020). The

mean r^2 of the BLUPs returned by StageWise should in principle be less sensitive to this artificial

inflation, as they are individual measures of correlation across replicates of each specific

genotype (Endelman, 2023). At the same time, these correlation coefficients will only ever be

accurate measures of true reliability to the degree that the BLUEs are taken as a form of

"ground-truthed" data (Rincent et al., 2012). In the case of unreplicated seedling genotypes,

where environmental variance components cannot be explicitly included in the model, this

assumption is clearly not entirely justified (Waldmann, 2019).

Application of genomic prediction within hazelnut breeding programs

There are numerous practical considerations that will impact the use of genomic prediction in the

various stages of a specific hazelnut breeding program. In particular, the per sample cost of

genotyping will be a determining factor influencing the number of plants for which predictions

can be made. In addition, important traits may express themselves relatively quickly (e.g., for

disease screens which can be performed in a greenhouse environment), or take over a decade

before a phenotype is available (e.g., for mature plant architecture, or multi-year cumulative

yield). This will significantly impact the benefit of genomic-based selection, relative to

phenotypic selection.

In general, however, several conclusions are evident from the data presented here. First,

prediction accuracies that are obtained when phenotypic data is not masked for individuals in the

training population are clearly high enough to warrant including genetic marker information in

the selection of parents. The set of parental candidates is typically relatively small, and breeders

frequently have limited access to replicated trial data for all individuals with which crosses might

realistically be made. At the same time, some phenotypic data will be available before a cross

can physiologically be performed, and thus the inclusion of genotypic information would in this

context be used to perform genomic-assisted selection (Jannink et al., 2010). In addition, in

mature breeding programs, close relatives of potential parents may have substantial phenotypic

information available, which can be used to improve the training population. The improvements

in trait prediction offered by such an approach will generally be warranted, given the

progressively declining cost of sequencing.

Second, in many applied contexts, genomic prediction is not simply used in a marker-assisted

fashion, but also to perform marker-based prediction, where only genotypic data is available for

some individuals in the population. In these cases, a so-called "training population" is

phenotyped and genotyped, allowing genotypic values to be predicted for un-phenotyped

individuals. The use of purely genomic-based prediction at the seedling stage, performed to

determine which subset of progeny families to advance into field trials, will necessarily be more

limited. Phenotypic data will be largely absent for such individuals. While there may be cases

in which phenotypic data exists for half-sib or full-sib families created from crosses using the

same parents, any training population will be to some degree removed from the individuals for

which predictions are being made. In addition, while the size of any specific progeny family will

be limited, a breeding program as a whole could easily generate thousands of seedlings each

year, genotyping all of which would likely represent a prohibitive expense. Instead, a prudent

balancing act should be made between resources dedicated to genotyping, and field trials used to

grow out new progeny each year, such that the number of progeny generated each year can in

fact be genotyped. Even in the absence of phenotypic data, and thus reduced prediction

accuracies, genotyping at the seedling stage will improve genetic gain, and allow for a more

informed preservation of genetic variance within the breeding program (Endelman, 2025).

In sum, these results provide a promising basis for further developing genomic selection models

for use in hazelnut breeding programs. This study represents the first such model developed for

Corylus, and despite the limited degree of replication, the prediction accuracies obtained for nut

quality traits suggest this method can substantially improve the rates of genetic gain for

polygenic traits. We hope future studies will expand upon this work, validating models with

more robust experimental designs to minimize better minimize environmental variances, and

validate predication accuracies for a greater diversity of traits.

ACKNOWLEDGEMENTS

The Dawson Lab assisted greatly in the collection of phenotypic and genotypic data, specifically,

Marissa Nix, Peyton Higgins, Martha Barta, Ava Gorius, Ava Glaser, Tressa Peskar, Malachi

Persche, Matt Mirkes, Shea Tillotson, Raeann Rich, Maya Giordano, Calliana Wickus, Thomas

Hickey, and Brent Johnson. We are thankful to the support of Chuck Zinda, for allowing us

access to the *C. americana* diversity panel in Wisconsin, as well as Mark Hamann, Lois Braun,

and Les Everett, for providing the interspecific biparental families in Minnesota.

ORCID

S. H. Brainard: 0000-0001-7678-3716

J. C. Dawson: 0000-0002-9907-8611

SUPPLMENTAL MATERIAL

Supplementary Figure 1. The impact of marker dilution is illustrated in terms of its impact on

prediction accuracy. **A:** results from the *C. americana* diversity panel; **B:** results from the

interspecific biparental populations. As in Figure 5, asymptotic prediction accuracy is reached

between 5,000 and 10,000 markers.

Supplementary Figure 2. Progressive dilution of the training population is shown for the

interspecific biparental populations in Minnesota. The analysis is similar to what is shown in

Fig. 6, however in this case, prediction accuracy is largely a function of correctly prediction F_1

population identity, and not the genetic value of individual samples, due to the significant degree

of population structure present when all three biparental families are combined.

Supplementary Figure 3. Spatial variation across the field sites for **A:** kernel length in the

Wisconsin *C. americana* diversity panel; **B:** kernel height in the Wisconsin *C. americana*

diversity panel; **C:** kernel length in the Minnesota interspecific biparental populations; **D:** kernel

height in the Minnesota interspecific biparental populations. No clear structured spatial variation

is observed within sites, or any degree of relatively greater or lesser variation between sites.

Supplementary Figure 4. Violin plots comparing observed phenotypic values with the genetic

values predicted using BLUPs. **A:** Shrinkage is observed in the *C. americana* diversity panel,

and this effect increases as the training population because less related to the sample. **B:** Much

more extreme shrinkage is observed in the biparental populations across all traits, despite the

similar training population size.

FUNDING

Funding for this project was provided by USDA NIFA-SCRI Award H007913501, and the

Savanna Institute.

CONFLICT OF INTERETS

The authors declare no conflict of interest.

DATA AVAILABILITY

Digital imagery (both raw and binary masks) which were used for phenotyping, genetic map

using multialleic makers, ordered linkage groups for the "Eric\Jeff" F_1 population, input marker

and phenotype files for use in fullsibQTL, as well as input marker and phenotype files for use in

StageWise, are available via DataDryad [temporary link for peer review:

http://datadryad.org/share/DdbiZHYdCypYeb2d7gCfkZC8edWZ4jpWjfmRZcyitpw].

References

Baytar, A.A., E.G. Yanar, A. Frary, and S. Doğanlar. 2024. Association mapping and candidate
gene identification for yield traits in European hazelnut (L.). Plant Direct 8(8): e625. doi:
10.1002/pld3.625.

Bradbury, P.J., Z. Zhang, D.E. Kroon, T.M. Casstevens, Y. Ramdoss, et al. 2007. TASSEL:
software for association mapping of complex traits in diverse samples. Bioinformatics
23(19): 2633–2635. doi: 10.1093/bioinformatics/btm308.

Brainard, S.H., J.A. Bustamante, J.C. Dawson, E.P. Spalding, and I.L. Goldman. 2021. A Digital
Image-Based Phenotyping Platform for Analyzing Root Shape Attributes in Carrot.
Frontiers in Plant Science 12.
https://www.frontiersin.org/articles/10.3389/fpls.2021.690031 (accessed 20 August
2022).

Brainard, S.H., J.C. Dawson, J.A. Fischbach, and L.C. Braun. 2023. Improving selection
efficiency in C. americana × C. avellana interspecific hybrids through the development of
an indel-based genetic map. Acta Hort 1379: 135–140. doi:
https://doi.org/10.17660/ActaHortic.2023.1379.20.

Brainard, S.H., D.M. Sanders, T. Bruna, S. Shu, and J.C. Dawson. 2024. The first two
chromosome-scale genome assemblies of American hazelnut enable comparative
genomic analysis of the genus Corylus. Plant Biotechnology Journal 22(2): 472–483. doi:
10.1111/pbi.14199.

Danecek, P., J.K. Bonfield, J. Liddle, J. Marshall, V. Ohan, et al. 2021. Twelve years of
SAMtools and BCFtools. GigaScience 10(2): giab008. doi: 10.1093/gigascience/giab008.

Edwards, S.M., J.B. Buntjer, R. Jackson, A.R. Bentley, J. Lage, et al. 2019. The effects of
training population design on genomic prediction accuracy in wheat. Theoretical and
Applied Genetics 132(7): 1943–1952. doi: 10.1007/s00122-019-03327-y.

Elshire, R.J., J.C. Glaubitz, Q. Sun, J.A. Poland, K. Kawamoto, et al. 2011. A Robust, Simple
Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. PLOS ONE
6(5): e19379. doi: 10.1371/journal.pone.0019379.

Endelman, J.B. 2023. Fully efficient, two-stage analysis of multi-environment trials with
directional dominance and multi-trait genomic selection. Theor Appl Genet 136(4): 65.
doi: 10.1007/s00122-023-04298-x.

Endelman, J.B. 2025. Genomic prediction of heterosis, inbreeding control, and mate allocation in
outbred diploid and tetraploid populations. Genetics 229(2): iyae193. doi:
10.1093/genetics/iyae193.

Endelman, J.B., and J.-L. Jannink. 2012. Shrinkage estimation of the realized relationship
matrix. G3: Genes, Genomes, Genetics 2(11): 1405–1413. doi: 10.1534/g3.112.004259.

FAOSTAT. 2022. Value of Agricultural Production Database. Food and Agricultural
Organization of the United Nations. https://www.fao.org/faostat/en/#data/QV (accessed
29 August 2022).

Gazaffi, R., R.R. Amadeu, M. Mollinari, J.R.B.F. Rosa, C.H. Taniguti, et al. 2020. fullsibQTL:

an R package for QTL mapping in biparental populations of outcrossing species. doi:

10.1101/2020.12.04.412262.

Hameed, K., D. Chai, and A. Rassau. 2018. A comprehensive review of fruit and vegetable

classification techniques. Image and Vision Computing 80: 24–44. doi:

10.1016/j.imavis.2018.09.016.

Heslot, N., J.-L. Jannink, and M.E. Sorrells. 2015. Perspectives for Genomic Selection

Applications and Research in Plants. Crop Science 55(1): 1–12. doi:

https://doi.org/10.2135/cropsci2014.03.0249.

Jannink, J.-L., A.J. Lorenz, and H. Iwata. 2010. Genomic selection in plant breeding: from

theory to practice. Briefings in Functional Genomics 9(2): 166–177. doi:

10.1093/bfgp/elq001.

Kosambi, D.D. 1943. The Estimation of Map Distances from Recombination Values. Annals of

Eugenics 12(1): 172–175. doi: 10.1111/j.1469-1809.1943.tb02321.x.

Margarido, G.R.A., A.P. Souza, and A. a. F. Garcia. 2007. OneMap: software for genetic

mapping in outcrossing species. Hereditas 144(3): 78–79. doi: 10.1111/j.2007.0018-

0661.02000.x.

Mehlenbacher, S.A., and T.J. Molnar. 2021. Hazelnut Breeding. In: Goldman, I., editor, Plant

Breeding Reviews. 1st ed. Wiley. p. 9–141

Molnar, T.J., J.A. Honig, A. Mayberry, R.S. Revord, S.T. Lovell, et al. 2018. Corylus americana:

a valuable genetic resource for developing hazelnuts adapted to the eastern United States.

Acta Hortic. 1226: 115–122. doi: 10.17660/ActaHortic.2018.1226.16.

Nishio, S., T. Hayashi, T. Yamamoto, S. Terakami, H. Iwata, et al. 2018. Bayesian genome-wide

association study of nut traits in Japanese chestnut. Mol Breeding 38(8): 99. doi:

10.1007/s11032-018-0857-3.

Ozturk, S.C., S.E. Ozturk, I. Celik, F. Stampar, R. Veberic, et al. 2017. Molecular genetic

diversity and association mapping of nut and kernel traits in Slovenian hazelnut (Corylus

avellana) germplasm. Tree Genetics & Genomes 13(1): 16. doi: 10.1007/s11295-016-

1098-4.

Revord, R.S., S.T. Lovell, J.M. Capik, S.A. Mehlenbacher, and T.J. Molnar. 2020. Eastern

Filbert Blight Resistance in American and Interspecific Hybrid Hazelnuts. J. Amer. Soc.

Hort. Sci. 145(3): 162–173. doi: 10.21273/JASHS04732-19.

Rincent, R., D. Laloë, S. Nicolas, T. Altmann, D. Brunel, et al. 2012. Maximizing the Reliability

of Genomic Selection by Optimizing the Calibration Set of Reference Individuals:

Comparison of Methods in Two Diverse Groups of Maize Inbreds (Zea mays L.).

Genetics 192(2): 715–728. doi: 10.1534/genetics.112.141473.

Rochette, N.C., A.G. Rivera-Colón, and J.M. Catchen. 2019. Stacks 2: Analytical methods for

paired-end sequencing improve RADseq-based population genomics. Molecular Ecology

28(21): 4737–4754. doi: 10.1111/mec.15253.

Sverrisdóttir, E., E.H.R. Sundmark, H.Ø. Johnsen, H.G. Kirk, T. Asp, et al. 2018. The Value of

Expanding the Training Population to Improve Genomic Selection Models in Tetraploid

Potato. Frontiers in Plant Science 9: 1118. doi: 10.3389/fpls.2018.01118.

Tayeh, N., A. Klein, M.-C. Le Paslier, F. Jacquin, H. Houtin, et al. 2015. Genomic Prediction in

Pea: Effect of Marker Density and Training Population Size and Composition on

Prediction Accuracy. Frontiers in Plant Science 6: 941. doi: 10.3389/fpls.2015.00941.

Torello Marinoni, D., N. Valentini, E. Portis, A. Acquadro, C. Beltramo, et al. 2018.

Construction of a high-density genetic linkage map and QTL analysis for hazelnut

breeding. Acta Hortic. (1226): 25–30. doi: 10.17660/ActaHortic.2018.1226.3.

USDA Economic Research Service. 2025. Tree nuts bearing acreage, yield, gross returns per

acre, production, grower price, and supply and availability.

VanRaden, P.M. 2008. Efficient Methods to Compute Genomic Predictions. Journal of Dairy

Science 91(11): 4414–4423. doi: 10.3168/jds.2007-0980.

Waldmann, P. 2019. On the Use of the Pearson Correlation Coefficient for Model Evaluation in

Genome-Wide Prediction. Front. Genet. 10. doi: 10.3389/fgene.2019.00899.

Werner, C.R., R.C. Gaynor, G. Gorjanc, J.M. Hickey, T. Kox, et al. 2020. How Population

Structure Impacts Genomic Selection Accuracy in Cross-Validation: Implications for

Practical Breeding. Front. Plant Sci. 11. doi: 10.3389/fpls.2020.592977.

Wu, R., C.-X. Ma, I. Painter, and Z.-B. Zeng. 2002. Simultaneous Maximum Likelihood

689   Estimation of Linkage and Linkage Phases in Outcrossing Species. Theoretical

690   Population Biology 61(3): 349–363. doi: 10.1006/tpbi.2002.1577.

691 Würschum, T. 2012. Mapping QTL for agronomic traits in breeding populations. Theor Appl

692   Genet 125(2): 201–210. doi: 10.1007/s00122-012-1887-6.

693 Zhang, A., H. Wang, Y. Beyene, K. Semagn, Y. Liu, et al. 2017. Effect of Trait Heritability,

694   Training Population Size and Marker Density on Genomic Prediction Accuracy

695   Estimation in 22 bi-parental Tropical Maize Populations. Frontiers in Plant Science 8:

696   1916. doi: 10.3389/fpls.2017.01916.

697 Zhao, Y., M. Gowda, W. Liu, T. Würschum, H.P. Maurer, et al. 2013. Choice of shrinkage

698   parameter and prediction of genomic breeding values in elite maize breeding populations.

699   Plant Breeding 132(1): 99–106. doi: 10.1111/pbr.12008.

700 Zhou, Y., M.I. Vales, A. Wang, and Z. Zhang. 2017. Systematic bias of correlation coefficient

701   may explain negative accuracy of genomic prediction. Briefings in Bioinformatics 18(5):

702   744–753. doi: 10.1093/bib/bbw064.

FIGURES

Figure 1. Digital image acquisition and processing pipeline. Staged images of subsamples of 30

inshell nuts (**A**) and kernels (**B**) were acquired using a tethered DSLR camera. QR codes were

scanned, with identifying information used for file management. Blue boxes containing

individual nuts or kernels were then cropped, with the dimensions of each box being used as a

scale for converting pixel to spatial resolution. RGB indexing was used to create a binary mask

of each nut or kernel, with bounding boxes fit to each mask to obtain phenotypic measures.

Figure 2. Histograms for six traits measured using digital imagery. Blue distributions

correspond to the Wisconsin diversity population of *C. americana*, while red distributions

correspond to the interspecific biparental populations in Minnesota. Length and height

measurements for inshell (**A**, **B**) and kernel (**C**, **D**) illustrate the larger size of nuts from the

interspecific biparental populations in Minnesota, the paternal parents of which are commercial

C. avellana cultivars. Slightly greater variance is also observed in the kernel measurements from

the interspecific biparental populations. In contrast, higher percent kernel (measured

volumetrically) was observed in the *C. americana* diversity population in Wisconsin (**E**), while

the two populations had nearly identical distributions for percent kernel when measured by

weight (**F**).

Figure 3. PCA biplots of kernel and nut morphological characteristics, specifically inshell

length, inshell height, kernel length, kernel height, percentage of kernel by volume, and

percentage of kernel by weight. Parallel vectors indicate a positive correlation between the two

traits, while vectors oriented 180 degrees from each other indicate negative correlation, and

vectors at right angles reflect independence between the relevant traits. In all analyses, kernel

height and length, and inshell height and length, were tightly positively correlated with each

other, while volumetric and gravimetric measures of percent kernel were more loosely positively

correlated. **A:** In the *C. americana* diversity population in Wisconsin, inshell and kernel traits

were also somewhat positively correlated, and uncorrelated with percent kernel metrics. **B:** In

the interspecific biparental populations in Minnesota, the positive correlation between inshell and

kernel measures was substantially reduced, to such a degree that percent kernel metrics

approached a negative correlation with inshell traits. **C:** When both populations were assessed

together, correlations appeared much more similar to the *C. americana* population in Wisconsin

(likely due to its larger size), with a slight weakening of the positive correlation between the two

measure of percent kernel.

Figure 4. LOD profiles of the composite interval mapping results in the Eric\Jeff interspecific

biparental F_1 population for six morphological nut traits. **A**: inshell length; **B:** inshell height; **C:**

kernel length; **D:** kernel height; **E:** percent kernel (measured volumetrically); **F:** percent kernel

(measure gravimetrically).

Figure 5. Assessment of the impact of marker density on pairwise estimates relatedness within

both the *C. americana* diversity panel, and the interspecific biparental populations. First, the **G**

matrix was calculated with the complete marker set, and then for each number of markers along

the *x*-axis, a **G** matrix was calculated again using StageWise. These matrices were then

subtracted to give a scalar difference, representing the deviation in total estimates of relatedness

at varying levels of marker density. **A:** results for the *C. americana* diversity panel in Wisconsin;

B: results for the interspecific biparental populations in Minnesota.

Figure 6. The impact of training population design on prediction accuracy in the *C. americana*

diversity panel in Wisconsin. **A:** The impact of decreasing the relatedness of the training

population to the predicted sample. Declining prediction accuracy is observed as the most

related individuals are removed from the training population, until an asymptotically minimum

prediction accuracy is reached once the twenty most-closely related individuals have been

removed. **B:** The impact of increasing population size on prediction accuracy, when these twenty

most-20 individuals have been removed. Population size is increased from 100 to 275

individuals, with marginal increases in prediction accuracy observed for four traits (inshell and

kernel height, inshell and kernel length), while prediction accuracy for both measures of percent

kernel were unaffected.

TABLES

Table 1. Results of composite interval mapping results in the Eric\Jeff interspecific biparental F_1

population for six morphological nut traits. For each QTL which exceeded the permutation-test

determined LOD threshold, the position is reported in cM along the respective linkage group. In

addition, additive effects of both alleles, and dominance effects are reported. R^2 values for the

individual QTLs, as well as the complete model are also given.

Inshell height

LG	Pos (cM)	α_p	α_q	δ_{pq}	R^2 (%)
1	17.346	-0.473	-0.294	0.457	19.239
2	6.234	0.016	-0.027	-0.429	13.592
8	9.623	0.068	0.490	-0.182	9.884
11	0.012	-0.093	0.435	-0.223	10.809
All	-	-	-	-	48.513

Inshell length

LG	Pos (cM)	α_p	α_q	δ_{pq}	R^2 (%)
1	17.062	-0.400	-0.254	0.386	19.517
2	3.564	-0.047	0.027	-0.573	19.066
6	38.255	-0.359	0.090	-0.272	7.574
8	32.854	-0.038	0.351	-0.300	9.723
All	-	-	-	-	55.472

Kernel height

LG	Pos (cM)	α_p	α_q	δ_{pq}	R^2 (%)
1	51.295	0.101	-0.262	-0.201	7.976
2	12.013	0.063	-0.186	-0.246	9.129
8	34.421	-0.167	0.397	-0.196	7.310
All	-	-	-	-	19.809

Kernel length

LG	Pos (cM)	α_p	α_q	δ_{pq}	R^2 (%)
4	9.422	0.152	-0.075	0.151	8.926
9	29.042	0.243	0.515	-0.440	8.076
10	4.213	-0.081	-0.080	0.290	7.279
11	18.915	-0.111	-0.084	-0.242	7.237
All	-	-	-	-	37.344

Percent kernel (gravimetric)

LG	Pos (cM)	α_p	α_q	δ_{pq}	R^2 (%)
6	27.213	0.021	-0.006	0.012	20.323
8	52.938	0.006	0.043	-0.017	10.622
9	35.709	0.002	-0.004	0.022	8.259
All	-	-	-	-	37.143

Percent kernel (volumetric)

LG	Pos (cM)	α_p	α_q	δ_{pq}	R^2 (%)
1	17.005	0.003	0.009	-0.0181	17.082
3	71.099	-0.002	-0.016	0.0007	7.434
6	32.960	-0.023	-0.025	0.0002	14.203
9	7.961	0.006	0.011	0.0134	12.035
All	-	-	-	-	45.142

770 **Table 2.** Heritabilities of six morphological nut traits in the *C. americana* diversity population in

771 Wisconsin and the interspecific biparental families in Minnesota.

Trait	Wisconsin	Minnesota
Inshell height	0.585	0.718
Inshell length	0.510	0.654
Kernel height	0.678	0.301
Kernel length	0.527	0.347
Percent kernel (grav.)	0.615	0.553
Percent kernel (vol.)	0.513	0.542

Trait	Wisconsin – StageWise-r^2	Wisconsin – Pearson-r^2	Minnesota – StageWise-r^2	Minnesota – Pearson-r^2
Height (in-shell)	0.55	0.90	0.88	0.87
Length (in-shell)	0.51	0.85	0.84	0.75
Height (kernel)	0.62	0.93	0.44	0.51
Length (kernel)	0.53	0.83	0.48	0.42
% kernel (grav.)	0.61	0.92	0.74	0.75
% kern (vol.)	0.53	0.87	0.71	0.73

775 **Table 3.** Impact of relatedness on prediction accuracy in the interspecific biparental populations

776 in Minnesota. For each of the three F_1 families, leave-one-out validation was performed using

777 the entirety of each of the three families, with average Pearson-r^2 values being calculated. The

778 prediction population which generated the highest prediction accuracy is highlighted.

Sample population	Prediction Population	Prediction accuracy
Gibs\York	Eric\Jeff	-0.128
Gibs\York	**Gibs\OSU**	**0.281**
Gibs\York	Gibs\York	-0.0545
Eric\Jeff	Eric\Jeff	-0.723
Eric\Jeff	**Gibs\OSU**	**0.404**
Eric\Jeff	Gibs\York	-0.0219
Gibs\OSU	**Eric\Jeff**	**0.245**
Gibs\OSU	Gibs\OSU	-0.638
Gibs\OSU	Gibs\York	0.220

A *C. americana* diversity panel

Inshell Height · Inshell Length · Kernel Height · Kernel Length · Perc Kern Grav · Perc Kernel Vol

B Interspecific biparental populations