The human neuroblastoma SH-SY5Y cell line as a model to assess β-amyloid neurotoxicity: A systematic review and meta-analysis

Nathalia R. Pinheiro[a,i][*][†]; Clarissa F. D. Carneiro[b,i][*]; Giulia S. Cancelliero[a,i][*]; Giovanna O. Nogueira[a]; Glaucia M. Almeida[a]; Nathalia Fernandes[c,d,e,i]; Ana Paula Wasilewska-Sampaio[f]; Samantha M. Martins[g]; Antonio Felix[h,i].; Olavo B. Amaral[b,i#]; Adriano Sebollela[.a,i#]

[a]Department of Biochemistry and Immunology, Ribeirao Preto Medical School, University of Sao Paulo, Ribeirao Preto, SP, Brazil.
[b]Institute of Medical Biochemistry Leopoldo de Meis, Federal University of Rio de Janeiro, Brazil.
[c]Institute of Biological Sciences, Federal University of Amazonas, Manaus, Amazonas, Brazil.
[d]Mountain Research Center, Campus of Santa Apolónia, Polytechnic Institute of Bragança, Portugal.
[e]Laboratory for Sustainability and Technology in Mountain Regions, Campus of Santa Apolónia, Polytechnic Institute of Bragança, Portugal.
[f]Brazilian Biosciences National Laboratory (LNBio), Brazilian Centre for Research in Energy and Materials (CNPEM), SP, Brazil.
[g]Laboratório de Produtos Bioativos, Federal University of Rio de Janeiro, Campus Macaé, Macaé, RJ, Brazil.
[h]Department of Medicine, University of Pernambuco, Garanhuns, PE, Brazil.
[i]Brazilian Reproducibility Initiative in preclinical Systematic review and meta-Analysis (BRISA).

These authors contributed equally to this work

 # Corresponding authors:
Adriano Sebollela
sebollela@fmrp.usp.br

Olavo B. Amaral
olavo@bioqmed.ufrj.br

† present affiliation:
MUSC, USA

Abstract

The SH-SY5Y human neuroblastoma cell line is widely used as an *in vitro* model of β-amyloid (Aβ) neurotoxicity in Alzheimer's disease (AD). However, the lack of standardized protocols for assessing Aβ toxicity - including differentiation strategies for SH-SY5Y cells - limits the comparability of results across studies. To address these issues, we conducted a systematic review and meta-analysis to evaluate how methodological factors influence Aβ-induced toxicity in SH-SY5Y cells. We included 359 eligible studies encompassing 1,192 MTT-based comparisons of cell viability between Aβ-treated and control SH-SY5Y cells. A three-level meta-analysis estimated mean cell viability after Aβ exposure at 63% of control levels (95% CI [61.6; 64.3]), with very high heterogeneity (I^2=99.6%). Meta-regression identified significant associations between increased toxicity and higher Aβ concentrations, longer exposure durations, and the use of peptide preparations described as fibrils. Conversely, differentiation protocols, duration, and cell density did not significantly influence toxicity outcomes. Reporting quality was often poor, with frequent omissions regarding cell line origin, authentication, contamination testing, Aβ preparation details and nature of the experimental unit. Overall, our findings show robust Aβ toxicity in SH-SY5Y cells, primarily driven by dose, exposure time, and Aβ aggregation state, but not cell differentiation status. Our conclusions highlight the critical need for better reporting of Aβ exposure parameters to enhance reproducibility and translational potential in AD research.

Keywords: SH-SY5Y, β-amyloid, Alzheimer's disease, differentiation, neurotoxicity, cell line, MTT assay.

Background

Cognitive impairment in Alzheimer's disease (AD) is widely attributed to synaptic dysfunction induced by aggregates of the β-amyloid peptide (Aβ) (1–3). Enzymatic cleavage of the amyloid precursor protein (APP) on the extracellular surface of neurons by BACE-1 and γ-secretase generates different isoforms of the Aβ peptide, with the most abundant isoforms having 40 (Aβ40) or 42 amino acid residues (Aβ42) (4). These peptides can assemble into structurally diverse aggregates with varying degrees of solubility and toxicity (5–9). Despite substantial efforts to characterize the neurotoxic properties of different Aβ species, progress has been hindered by inconsistencies across experimental models, Aβ concentrations, exposure durations, and aggregation protocols (6,10,11). This lack of standardization contributes to the limited generalizability of preclinical findings and may partly explain the high failure rates of Aβ-targeting therapies in clinical trials (3,12,13).

The SH-SY5Y cell line is a widely used model in neuroscience research, including studies on AD (11,14–16). Generated through multiple rounds of subcloning and selection from the parental SK-N-SH human neuroblastoma cell line, SH-SY5Y cells can be employed either in their undifferentiated state or following differentiation into a mature neuron-like phenotype, with all-trans retinoic acid (ATRA) being the preferential agent for this purpose (17,18). ATRA acts through the inhibition of cellular proliferation accompanied by increased expression of β-tubulin III, a cytoskeletal protein typical of mature neurons (14,19) and favours the emergence of a cholinergic phenotype (18). When associated with brain-derived neurotrophic factor (BDNF), ATRA further enhances neuronal characteristics, including the formation of synaptic structures and axonal vesicle transport (14,20). Alternative differentiation protocols have also been explored, including

the use of specialized media (e.g., N2) or the addition of alternative growth factors (e.g., FGF) (21).

Several studies, including those using the SH-SY5Y cell line (20), have proposed that toxicity mechanisms elicited by Aβ aggregates involve direct interactions with synaptic structures (22,23). This raises concerns about the suitability of undifferentiated SH-SY5Y cells, which do not present extended neurites or functional synapses, as a model for studying Aβ-induced toxicity. Moreover, SH-SY5Y cells differentiated using distinct culture media exhibit differential toxicity responses to Aβ42 aggregates exposure (24), suggesting that the differentiation mechanistic route and final phenotype distribution of the cell culture impact toxicity outcomes. Such differences may explain the wide range of Aβ concentrations used in studies using SH-SY5Y as a model – including some that are much higher than those found *in vivo* (25,26). The combination of variability in differentiation protocols, Aβ aggregation preparation, and Aβ concentrations is likely to affect cellular responses and contribute to poor commensurability across studies (16,24).

Motivated by the widespread use of SH-SY5Y cells as a model to investigate Aβ toxicity, we conducted a systematic review and meta-analysis to determine the main factors influencing Aβ toxicity in this cell line. Specifically, we examined parameters related to Aβ exposure - including aggregation state, concentration, and duration - as well as cell line variables such as differentiation protocol and cell density. In addition, we carried out a comprehensive assessment of methodological aspects and reporting quality in the included articles. Our results point out windows of opportunity for improving reporting standards and may offer practical guidance for optimizing the use of SH-SY5Y cells in future studies of Aβ neurotoxicity and AD pathology.

Methods

Search Strategy

The study protocol was preregistered at the Open Science Framework (**10.17605/OSF.IO/EV4TZ**), and deviations from the preregistered protocol are described in https://osf.io/jc5bz. Initially, we created a list of 14 sentinel articles to guide the development of search strategies (Supplementary Table 1). Our final search string comprised two components, one with keywords related to the *in vitro* model and the other to AD or the β-amyloid peptide (Supplementary Table 2). We conducted this search among titles and abstracts of articles indexed in PubMed and EMBASE on October 18, 2020, filtering for those written in English. The total number of articles retrieved from each database is reported in Supplementary Table 2.

Title and abstract screening

Reviewers analyzed articles and removed duplicates using Rayyan (27). Inclusion criteria were (i) articles written in English that (ii) presented original results, (iii) described experiments with administration of the β-amyloid peptide of synthetic origin or purified from a recombinant source, (iv) described experiments with SH-SY5Y cells as a model, and (v) were not retracted. The title and abstract screening step was carried out following the same criteria in the screening pilots, with each article evaluated by a single reviewer and 10% of articles undergoing double screening to calculate agreement levels. The agreement between reviewer pairs was measured by absolute percentage and Cohen's kappa.

Full-text screening

During full-text screening, each article included was assessed by a reviewer distinct from the one who had performed the initial title and abstract screening. Inclusion

criteria were the same as in the previous step, with the addition of the following: (i) use of SH-SY5Y cells exposed to the Aβ peptide in its isoform(s) 1-37 to 1-43; (ii) use of a control group of SH-SY5Y cells, subject to the same culture conditions but not exposed to Aβ; (iii) use of toxicity outcomes measuring cell viability, cell death or oxidative stress markers; (iv) presentation of sample size, mean and standard deviation (SD) or standard error of the mean (SEM) for each group. If any of the required statistics were unclear or missing, the corresponding authors were contacted to request the necessary information. At this stage, the figure containing experiments that met the inclusion criteria was recorded using Airtable (https://airtable.com/).

Data Extraction

Figure data were extracted by a reviewer distinct from those who included the article in the screening stages. Variables were collected at the level of article, figure/table, results, and protocol and encompassed article metadata, experimental design, results, and protocol variables such as control and treatment group outcomes and details on Aβ, cell culture and treatment conditions. The complete set of extracted variables is described in Supplementary Table 3. Gsys (GSYS 2.4.7) was used to extract values from the graphs and Airtable (https://www.airtable.com/) was used to record data.

Data Processing

Data sheets with data extracted by different reviewers were combined using R (4.4.3) (28). Extracted data for the description of experimental units was reviewed and categorized. Information on the number of plated cells per well and plate size was used to calculate cell density as a new variable. The description of medium supplements was reviewed, which led to the creation of two dichotomous variables to indicate the use of antibiotics and glutamine. After this manual cleaning step, data was imported back into R to conduct all subsequent cleaning, analysis, and visualizations. Comparisons with no

data on either group's means or sample size, as well as comparisons with missing variation for the treated group, were excluded from the analyses. When variability was reported as SEM, the SD was calculated as SEM×√sample size. If the meaning of error bars was unclear, they were assumed to represent SEM. If error bars were presented for the treated group only, we assumed that paired normalization was performed (with each treated unit divided by a particular control value).

Data Synthesis

All analysis scripts and datasets are available at OSF ([10.17605/OSF.IO/EV4TZ](10.17605/OSF.IO/EV4TZ)). Packages used are listed in Supplementary Table 4 (29–33). All analyses were conducted with both log-transformed ratio of means (ROM/ROMC) and standardized mean difference (SMD) as effect size measures. When the control group had no SD or SEM reported, we considered it as zero and used measure = "ROMC" or "SMC" in the escalc function in metafor. Measure = "ROM" or "SMD" was used when SD or SEM was reported for the control group. For all ratio of means analyses, we transformed the resulting effect estimates to a linear scale (i.e., % of control) when displaying results by taking the natural exponential of the meta-analytical results.

Random-effects meta-analyses (two-level models) were calculated using the restricted maximum likelihood estimator for τ^2. Based on these results, publication bias was assessed by trim-and-fill analyses using both R0 and L0 estimators, and by Egger's regression using the inverse of SEM as the predictor variable. Three-level meta-analysis models were calculated using random effects at both the article and experiment levels.

Three-level univariate meta-regressions were performed using differentiation duration (no differentiation = 0 days), Aβ aggregation state, Aβ concentration, Aβ exposure duration and cell density. For meta-regressions using Aβ concentration as a moderator, we excluded comparisons in which Aβ concentrations were higher than 100

μM (18 comparisons), as high concentrations of this peptide in solution are prone to fast aggregation into insoluble species (34). Three-level multivariate meta-regressions used the maximum likelihood estimator of τ^2, and all combinations of variables from the selected list were tested in multivariable models. The best models were ranked by corrected Akaike Information Criteria (AICc). Additionally, we calculated the R^2 for the model and performed a Q test of moderators for each variable (including all dummy variables for each categorical moderator) to obtain p-values for individual variables.

Results

Selection of experiments

The study selection process is illustrated in the PRISMA flowchart (Figure 1). A total of 9,391 articles were screened, resulting in 1,364 articles included for full-text screening. Mean agreement between reviewers in the sample of articles undergoing double screening was 86.6%, with a Cohen's kappa of 0.67 (see https://osf.io/bg4qt). From the 1,364 articles analyzed in the full-text screening step, 553 were included for data extraction.

After the second screening, an initial analysis of the toxicity readout methods (Supplementary Table 5) revealed metabolic viability-based assays to be the most common readout, with 1,386 comparisons (54% of total). MTT was the most widely used reagent in these assays (>80% of comparisons; Supplementary Table 6). Based on the widespread use of these assays, we opted to extract data only from these comparisons for feasibility reasons. During data extraction, 215 additional comparisons in the selected articles were identified as eligible by reviewers, which led to a total of 444 articles and 1,601 comparisons. Of these, 226 comparisons were excluded due to missing information on means or variability, and 183 were excluded due to lack of information on sample size, leaving 1,192 comparisons from 359 articles for analysis.

Figure 1. PRISMA flowchart of the systematic review. The flowchart summarizes the screening rounds and data extraction, which resulted in 359 articles, including 1,192 comparisons between control and treatment groups, selected for data analysis.

Sample description

Included articles were published between 1998 and 2021 (Supplementary Figure 1) and presented between 1 and 52 comparisons each, with a median of 2 and a mean of

3.3 comparisons per article (Supplementary Figure 2). In most of the selected studies (69%), the assay was used to test whether a given intervention could reverse the toxicity elicited by Aβ. Cell origin was reported in only 55% of articles. Reported sources were mostly cell banks (51% of articles), predominantly from the American Type Culture Collection (ATCC) (30%) or the European Collection of Authenticated Cell Cultures (ECACC) (8%) (Supplementary Table 7). The most commonly used culture medium was a combination of DMEM and F12 (36%), followed by DMEM (31%). In 8.5% of comparisons, culture medium was not reported. Most experiments included antibiotics in the culture medium (72%), and approximately 40% of the experiments included glutamine supplementation. In most experiments, the cell culture medium was supplemented with 10% fetal bovine/calf serum (FBS/FCS), but a wide range of serum types and concentrations was observed (Supplementary Table 8).

Table 1 summarizes variables related to the Aβ exposure protocol. All experimental comparisons but one described a single exposure to Aβ. Aβ1-42 was the most used isoform (81% of comparisons), followed by Aβ1-40 (18%), which is expected given the implication of these two isoforms in AD (35); (36); (8). In most experiments, neither the source of the peptide nor the species' identity was reported (52% and 80%, respectively). The oligomeric (soluble) form was the most prevalent aggregation status reported (38%), followed by fibrils (9%). Almost half of the experiments did not provide a clear description of the state of aggregation (45%) and likely used mixtures of different aggregates (soluble and insoluble). A clear description of the control (unexposed) condition (e.g., medium only, vehicle, etc.) used was missing in 44% of comparisons. Exposure duration and concentration of Aβ were reported in 99% of cases. A wide range of values was observed, with a mean (± S.D.) exposure duration of 33.2 h (± 16.9) and a mean Aβ concentration of 41.5 μM (± 875.4).

Table 1. Aβ exposure protocols.

Treatment variable	Category	Comparisons	
		N	%
Aβ isoform	1-42	963	80.8
	1-40	217	18.2
	1-43	10	0.8
	1-38	2	0.2
Aβ origin	Synthetic	524	44.0
	Recombinant	52	4.4
	Not reported	616	51.7
Aβ species identity	Human	232	19.5
	Rat	5	0.4
	Not reported	955	80.1
Aβ aggregation	Oligomers	462	38.8
	Fibrils	103	8.6
	Monomers	92	7.7
	Not reported	535	44.9
Control description	Vehicle	344	28.9
	Medium only	286	24.0
	Other	41	3.4
	Not reported	521	43.7
Treatment variable	**Estimate**	**Data**	
Exposure duration (hours)	Reported	99.1%	
	Mean (SD)	33.2 (16.8)	
	Median (min, max)	24 (0, 144)	
Concentration (μM)	Reported	98.6%	
	Mean (SD)	41.5 (875.4)	
	Median (min, max)	$10 \ (10^{-5}, 3\times10^{4})$	

The percentage of each Aβ treatment protocol variable is related to the 1192 included comparisons. The number (N), mean, standard deviation (SD), median, minimum (min), and maximum (max) values of exposure duration and concentration variables are displayed.

Variables related to the differentiation protocol are summarized in Table 2. SHSY-5Y cells underwent differentiation in 270 comparisons (23%). Culture medium

composition used for differentiation was detailed less often than media used for standard cell culturing. As for routine culturing, DMEM and DMEM+F12 were the most frequently reported medium options. The medium was supplemented with FBS/FCS in 47% of comparisons, with 5% or less being the most common concentration range (32%). Regarding the differentiation method, ATRA was employed in 51.5% of comparisons, while ATRA plus (ATRA associated with another agent, such as BDNF, N2, or FGF) appeared in 30%. In almost 20% of comparisons with differentiated cells, the duration of differentiation was not reported, and a similar fraction did not describe the concentration of ATRA. Among comparisons describing these variables, mean differentiation duration was 6.6 (± 3.2) days, and mean ATRA concentration was 9.2 μM (± 3.4).

Table 2. Differentiation protocol variables.

Protocol variable	Category	Comparisons	
		N	**%**
Method	ATRA	139	51.5
	ATRA plus	80	29.6
	Other	25	9.3
	Not reported	26	9.6
Serum type	FBS/FCS	127	47
	No serum	15	5.6
	Not reported	128	47.4
Serum concentration	5% or less	86	31.9
	10%	41	15.2
	No serum	15	5.6
	Not reported	128	47.4
Medium	DMEM_F12	111	41.1
	DMEM	38	14.1
	Other	25	9.3
	Not reported	96	35.6
Antibiotics	Yes	66	24.4
	Not reported	204	75.6
Glutamine	Yes	38	14.1
	No	21	7.8
	Not reported	211	78.1
Protocol variable	**Estimate**	**Data**	
Differentiation duration (days)	Reported	82.2%	
	Mean (SD)	6.6 (3.2)	
	Median (min, max)	6 (1, 14)	
ATRA concentration (µM)	Reported	80.7%	
	Mean (SD)	9.2 (3.4)	
	Median (min, max)	10 (0, 20)	

Numbers and percentages of comparisons for each category are presented. Percentages are relative to the 270 comparisons in which cells underwent differentiation. The number (N), mean, standard deviation (SD), median, and range of differentiation duration (days), and concentration of ATRA (µM) are also shown.

Reporting quality

Article-level variables related to reporting are presented in Table 3. None of the selected studies reported a sample size calculation, and protocol registration was reported in only 10 studies (2.8%). A conflict of interest statement was present in 51% of studies and was more frequent in recent articles (Supplementary Figure 3). In a significant proportion of articles (44.6%), the origin of the cells was not reported. Cell line authentication and testing for mycoplasma contamination were reported in only 2 and 3 articles, respectively, and protocols for these tests were not provided in any of them.

Table 3. Article-level reporting and cell line quality control.

Variable	Category	N	%
Reporting	Sample size calculation	0	0
	Conflict of interest statement	183	51
	Pre-registration	10	2.8
Cell source	Cell banks	182	50.7
	Donation	17	4.7
	Unclear/Not reported	160	44.6
Cell line authentication	Yes	2	0.6
	No/Not reported	357	99.4
Cell line mycoplasma testing	Yes	3	0.8
	No/Not reported	356	99.2

Numbers (N) of articles and percentages (in relation to the 359 included articles) for each category are presented.

Moreover, the meaning of sample sizes was not always clear. Although a generic description such as "independent experiments" or "replicates" was usually provided (ca. 64% of comparisons combined; Table 4), there is no guarantee that authors agree on the meaning of these terms. For instance, one may refer to "independent experiments" as

either repeating the assay using a single culture batch, independent culture batches, or different passages of the same batch.

Table 4. Experimental unit descriptions.

N definition	N	%
Independent experiments	542	45.5
Replicates	217	18.2
Experiments	108	9.1
Wells	24	2
Assays	19	1.6
Independent determinations	13	1.1
Samples	12	1
Determinations	8	0.7
Independent experimental measurements	6	0.5
Biological replicates	5	0.4
Others	9	0.8
Not informed	229	19.2

"Others" included cell cultures, independent repetitions, independent replicates, independent runs, independent sets of studies, and observations.

Main meta-analyses

We initially ran a random-effect meta-analysis including all 1,192 comparisons based on standardized mean differences as the effect size measure, as planned in our preregistered protocol. This model yielded an effect size (Hedges' g) of -5.47 (95% CI [-5.7, -5.2], p <0.0001) (Figure 2A) and high heterogeneity (I^2 = 93.77%, Q-test p value < 0.0001), indicating that the retrieved differences in effect size would be very unlikely to occur by sampling error alone.

The use of standardized effect sizes, however, presented two problems. First, in a significant number of studies (51.6%), no error bars for the control group were presented (likely because each treated unit had its value normalized by a particular control unit). In these cases, standardized effect sizes cannot be accurately estimated, as pooling the error bars for both groups will not provide the true pooled standard deviation. Second, the uncertainties over the meaning of experimental units (Table 4) raised questions about whether the degree of independence between experimental units was commensurable between studies. This concern was further strengthened by the presence of studies reporting extremely small standard deviations between experimental units, leading to very large standardized effect sizes - potentially due to units not being truly independent from each other in each group (as in the case of technical replicates). Beyond that concern, the mere fact that error bars were very small led to unreliable estimates of SD or SEM from graphs (and thus to unreliable standardized effect sizes).

Because of this, we chose to use the ratio of means as our main effect size measure, as it (a) provides results that are commensurable between normalized and non-normalized data and (b) is not distorted by underestimation of variability (although studies with very small standard errors will still have greater weight in meta-analyses). Using this approach, the random-effects meta-analysis yielded a log ratio of means ([95% CI]) of -0.46 [-0.48, -0.44], p<0.0001, which corresponds to mean cell viability in the Aβ-treated group being 63% [61.6%, 64.3%] of that observed in controls (Figure 2B). Heterogeneity was still very high (I^2=99.6%; Q-test p-value <0.0001). All our subsequent analyses are based on this measure, but alternative analyses using standardized effect sizes are presented as supplementary material.

Figure 2. Orchard plots of two-level random-effects meta-analyses. Standardized effect sizes (A) or relative effect sizes expressed as ratios of means (B) results of two-level random-effects meta-analyses using all 1,192 comparisons included in this study. Each comparison is represented by a grey circle, with its position on the x-axis indicating the effect size (i.e., standardized difference between groups in A, % viability in the treated group relative to control in B) and its size indicating the inverse of standard errors. The green dashed line indicates the absence of effect (i.e., zero difference or 100% viability).

Publication Bias Analysis

Publication bias was assessed by funnel plot asymmetry (Figure 3) and Egger's regression test (p=0.1587). Trim-and-fill analysis obtained with the R0 method showed strong evidence of publication bias (Supplementary Table 9), with 86 missing studies and an adjusted meta-analysis effect size of -0.40 [-0.42, -0.37], while the L0 method found no missing studies and thus yielded no difference from the main meta-analysis. The results observed in Figure 3 show that experiments in which viability in the treated group is above 100% are rarely reported. This is arguably expected, as in most experiments (~70%), Aβ exposure was used with the purpose of testing for reversal of its toxic effect with another intervention. In these cases, it is likely that the absence of toxicity is interpreted as a methodological failure that precludes evaluation of the treatment and thus ends up unreported.

Figure 3. Funnel plot for all 1,192 comparisons included in this study. Each comparison is represented by a grey circle, with its position on the x-axis indicating the cell viability in the Aβ-treated group (% of control) and on the y-axis indicating the study precision (1/SEM). The overall effect is indicated by the dashed green line.

Three-level meta-analyses

Given the nested structure of the data, with a mean of 3.3 comparisons per article, we conducted a three-level meta-analysis, using the experiment and study of origin as independent sources of variability nested within each other. We found the meta-analytical estimate to be similar to that of the two-level analysis, with a log ratio of means ([95% CI]) of -0.50 [-0.53, -0.47], p<0.0001, which corresponds to a mean cell viability in the Aβ-treated group of 61% [59%, 63%], p<0.0001. The variance ([95% CI]) was 0.09 [0.08, 0.1] with I^2=66.3% between comparisons and 0.04 [0.03, 0.06] with I^2=33.4% between articles, indicating that the heterogeneity between experiments within an article was

larger than that between articles. This may reflect the use of different experimental conditions (e.g., Aβ concentrations, exposure duration, etc.) within individual articles, including some that may be expected to have no effect and are used as negative controls.

Meta-regressions

The primary goal of our meta-analysis was not to obtain an effect estimate for the overall sample of articles, but rather to explore how different variables impact this effect. Before data collection, we selected the following variables to analyze in meta-regressions: differentiation method, differentiation duration, Aβ state of aggregation, Aβ concentration, Aβ exposure duration, cell density, and outcome measure; the latter was excluded as we only extracted data from cell viability assays. We also removed the differentiation method due to the limited number of comparisons in which a differentiation protocol was carried out. Analyses are based on all results for which the variable was reported and are shown in Table 5 and Figure 4 (with meta-regressions based on SMD available as Supplementary Table 10 and Supplementary Figure 4). Toxicity of Aβ was associated with higher concentrations, longer exposure, and Aβ preparations reported as fibrils or with no aggregation status reported (as compared to Aβ monomers). Differentiation duration showed a trend of association with lower toxicity, which was more significant when analysis was carried out using SMDs. Cell density, on the other hand, had no significant impact on effect sizes.

Table 5. Three-level meta-regression models for protocol variables.

Moderators and categories	Sample size	Effect size [95% C.I.]	Cell Viability	R^2	p-value
Duration of differentiation	1144	-0.51 [-0.54, -0.48]	60% [58.3%, 61.9%]	2.7%	0.0595
(day)		+0.01 [-0.0004, +0.02]	+1% [0%, +2%]		
State of aggregation (reference=monomers)	1192 (92)	-0.51 [-0.55, -0.47]	60% [57.7%, 62.5%]	2.2%	0.0004
Fibrils	103	-0.11 [-0.2, -0.02]	-10.4% [-18.1%, -2%]		0.0215
Oligomers	462	+0.05 [-0.01, +0.11]	+5.1% [-1, +11.6%]		0.123
Unclear	535	-0.13 [-0.24, -0.02]	-12.2% [-21.3%, -2%]		0.016
Aβ concentration	1123	-0.35 [-0.39, -0.32]	70.5% [67.7%, 72.6%]	15.7%	<0.0001
(μM)		-0.013 [-0.015, -0.011]	-1.3% [-1.5%, -1.1%]		
Aβ exposure duration	1180	-0.35 [-0.41, -0.30]	70.5% [66.4%, 74.1%]	2.7%	< 0.0001
(day)		-0.11 [-0.14, -0.07]	-10.4% [-13.1%, -6.8%]		
Cell density	690	-0.50 [-0.54, -0.46]	60.7% [58.3%, 63.1%]	0%	0.798
10^3 cells/cm^2		-4.2x10^{-12} [-3.6x10^{-12}, 2.8x10^{-11}]	0% [0%, 0%]		

Models have separate nested levels for articles and individual comparisons. Grey lines contain sample sizes, intercept effect sizes as ratio of means and cell viability, R^2 for the moderator and Q-test p-values for the moderator. White lines contain betas indicating the additional contribution of each unit/category to the effect size (as well as sample sizes and p-values for individual categories). For categorical variables, reference groups are described in the first column, with the sample size for these groups indicated in parentheses in the second column. I^2 and intercept p-values for all moderators were ~99.6% and <0.0001, respectively. Moderator R^2 values are calculated as the difference between total variances in the model with no moderators and in the tested model, divided by the total variance in the model with no moderators. Aβ concentrations above 100 μm were excluded from the concentration meta-regression.

Figure 4. Univariate meta-regressions. Each experiment is represented by an open circle, with size corresponding to the inverse of variance (i.e., larger circles are more precise). Aβ concentrations above 100 μm were excluded from the concentration meta-regression.

While univariate meta-regressions are informative, they can be confounded by covariation of the included variables. To account for this possibility, we also ran multivariate meta-regressions including the same variables used in the univariate analysis. Only comparisons with complete data for all variables were eligible for inclusion in the multivariate meta-regression. As a result, 571 comparisons were excluded, and 621 were retained for analysis. We then tested all 64 possible models, accounting for the combination of all variables. Table 6 shows the best model selected by the corrected Akaike Information Criteria (AICc), which includes Aβ aggregation status, Aβ concentration, and Aβ exposure duration and matches what was found by univariate meta-regressions. A similar multivariate meta-regression based on SMD was carried out, with results available in Supplementary Table 11.

Table 6. Three-level multivariate meta-regression model for testing selected protocol variables.

Model and categories	Sample size	Effect size [95% C.I.]	Cell viability	p-value
Aβ aggregation + Concentration + Exposure duration	621	-0.12 [-0.25, +0.02]	88.7% [77.9%, 102%]	<0.0001
Fibrils		-0.21 [-0.34, -0.07]	-18.9% [-28.8%, -6.8%]	0.004
Oligomers		-0.06 [-0.18, +0.07]	-5.8% [-16.5%, +7.3%]	0.369
Unclear		-0.14 [-0.27, -0.02]	-13.1% [-23.7%, -2%]	0.026
Aβ Concentration		-1.2×10^{-2} [-1.5×10^{-2}, -1×10^{-2}]	-1.2% [-1.5%, -1%]	<0.0001
Exposure duration		-0.1 [-0.14, -0.05]	-9.5% [-13.1%, -4.9%]	<0.0001

Models have separate nested levels for articles and individual comparisons. The covariates tested were differentiation duration, Aβ state of aggregation, Aβ concentration, Aβ exposure duration, and cell density. Out of the 64 tested models, only the best fit, as selected by AICc, is shown (Akaike weight and AICc of 0.28 and 324.1, respectively). In the grey line are the sample size, intercept effect sizes (as absolute mean differences and cell viability) and p-value for the full range of moderators. White lines present betas indicating the additional contribution of each unit/category to the effect size and cell viability (as well as sample sizes and p-values for individual variables). I^2, R^2 and intercept p-value for the model were 99.4%, 19.3% and 0.087, respectively.

Discussion

The human neuroblastoma cell line SH-SY5Y has been extensively employed as a cell model to investigate the toxicity of Aβ, as indicated by the large number of published articles retrieved from our search. To the best of our knowledge, this is the first systematic review to synthesize published data on Aβ-induced toxicity in SH-SY5Y cells. Our findings include both a scoping review of methodological and reporting practices at the article level and a meta-analysis and meta-regression to identify factors that impact the observed toxicity outcomes. The evidence compiled in this review demonstrates that Aβ exposure reduces the viability of SH-SY5Y cells, as assessed by the MTT reduction assay, suggesting impairment of mitochondrial metabolic activity and, by extension, neuronal viability.

Approximately 70% of the studies included in our review employed SH-SY5Y cells to evaluate interventions aimed at counteracting Aβ-induced toxicity. This predominant use of a neuroprotection-focused framework may partly explain the lower-than-expected number of studies reporting preserved or increased cell viability following Aβ exposure, as observed in the funnel plot. Such outcomes may have been perceived as methodologically incongruent with the study objectives (as demonstrating neurotoxicity as a prerequisite for testing protective effects) and therefore deemed unpublishable or uninformative. This aligns with prior findings on publication bias in biomedical sciences (37).

Poor reporting quality significantly hindered the interpretation and synthesis of results across studies. Notably, the absence of standardized definitions for the experimental unit in *in vitro* research limited our ability to interpret reported sample sizes accurately. A wide range of ambiguous descriptions was used, often failing to clearly distinguish between technical and biological replicates. This lack of clarity may partly

account for the unusually large standardized effect sizes observed in some comparisons, as technical replicates typically exhibit lower variability than biological replicates, potentially leading to inflated effect sizes when the former are incorrectly treated as independent observations. Additional factors that may contribute to these extreme effect sizes include inaccuracies in the extraction of means and error values from figures - particularly when error bars were small - as well as selective reporting practices (i.e., exclusion of outliers), and eventually data fabrication.

Among the 359 articles from which data were extracted, 44.6% did not report the origin of the cell line used, and 4.7% stated that it was acquired by donation. Along with the absence of cell line authentication, this raises concerns about the reliability of the cell lines used. This is particularly relevant for cell lines like SH-SY5Y, which, being tumor-derived, are especially prone to phenotypic and genetic drift during prolonged culture (38). Furthermore, the risk of cross-contamination between cell lines remains a well-documented issue (38,39), and, without routine authentication procedures, such contamination may go undetected, especially when morphological differences are subtle or absent. These concerns, when considered alongside others, such as poor definition of experimental units, highlight the need for rigorous assessment of internal validity of *in vitro* studies (40). The development and implementation of standardized tools to evaluate risk of bias in cell culture research is therefore critical for advancing evidence synthesis in this field, and ongoing initiatives are beginning to address this gap (41–43).

In terms of reporting quality, a significant proportion of experiments failed to specify the source of Aβ or the species origin of its amino acid sequence. Although it is reasonable to assume the peptide used is usually synthetic and based on the human sequence, clearer reporting of these variables is recommended, as recombinantly produced peptides or sequences from rats were used in some studies. Likewise, almost

half of the experiments did not provide a clear description of the aggregation state of Aβ and likely used heterogeneous mixtures of soluble and insoluble aggregates. This lack of proper description is concerning, given the differential toxicity of soluble and insoluble Aβ aggregates (6,8,44). The clinical implication of this differential role of distinct Aβ aggregates is underscored by findings reporting that amyloid plaque burden does not consistently correlate with cognitive decline in Alzheimer's disease (45,46). It is therefore strongly recommended to carefully select and characterize Aβ aggregation status when using SH-SY5Y cells to evaluate novel therapeutic compounds for Aβ toxicity, with the ultimate goal of developing effective treatments for Alzheimer's disease.

The mean Aβ concentration used across the studies included in this review was 41.5 μM, a high value compared to the estimated concentrations found *in vivo*. For instance, the levels of soluble Aβ oligomers found in brain tissue from AD patients have been reported to be in the low picogram per gram range (47,48), and neurotoxicity has been induced *in vitro* by human brain-derived Aβ oligomers at concentrations as low as 50 nM (49). Interestingly, the median Aβ concentration in our review, 10 μM, was significantly lower than the mean, indicating a right-skewed distribution driven by a subset of studies using concentrations in the millimolar range. This discrepancy points to a potential lack of awareness regarding the low aqueous solubility of Aβ - particularly the 42-residue isoform - and highlights a broader issue of insufficient standardization in *in vitro* neurotoxicity models. Despite these caveats, our meta-analysis revealed a robust and statistically significant reduction in cell viability following Aβ exposure, consistent across both standard random-effects models and three-level models accounting for clustering at the study level.

Of potential relevance to future studies, our meta-regressions identified a significant association between both Aβ concentration and exposure duration with cell

viability, with higher concentrations and longer exposure predictably correlating with lower cell viability. These findings align with the established dose- and time-dependent nature of Aβ-induced neurotoxicity. In contrast, differentiation status showed only a nonsignificant trend towards decreased toxicity with longer differentiation periods. This result was somewhat unexpected, given that Aβ toxicity has been shown to involve direct interactions with synaptic structures (22,23,50), which are less abundant in undifferentiated SH-SY5Y cells than in differentiated ones (18,51). The observed trend may therefore be spurious or influenced by methodological heterogeneity. In line with this notion, individual studies have reported increased susceptibility to Aβ toxicity in differentiated SH-SY5Y cells relative to undifferentiated ones (24). The absence of a robust effect in our meta-analysis may reflect the influence of uncontrolled confounding variables or the possibility that, at supraphysiological Aβ concentrations, toxicity mechanisms may become less selective and less dependent on neuronal maturity.

We also examined the impact of Aβ aggregation state on cellular toxicity. Interestingly, compared to monomeric Aβ, which has been reported to be non-toxic and may even play physiological roles at low concentrations (52,53), both fibrillar Aβ and preparations of undefined aggregation status were associated with significantly higher toxicity. In contrast, Aβ preparations classified as oligomeric were associated with slightly lower toxicity, although this difference was not statistically significant. This finding was also unexpected, given the robust body of evidence pointing to soluble oligomers as the most neurotoxic Aβ species (2,46,54). As with differentiation status, we cannot rule out the possibility that poor reporting, inadequate biochemical characterization and classification of Aβ preparations, or other unidentified confounders may have biased this assessment.

Finally, a growing number of studies have reported that highly toxic Aβ assemblies exist along the continuum between soluble oligomers and insoluble fibrils. These species are distinct from typical protofibrils - an on-pathway to fibrils species - and are commonly found in close interaction with mature fibrils (55,56). It seems reasonable to consider that preparations categorized as fibrils, as well as those that were not characterized ("unclear"), may contain and release such toxic assemblies, in contrast to typical fibril-free oligomeric preparations. Therefore, the observed higher Aβ toxicity retrieved from studies using fibrils or unclear aggregation status could be instigated by intermediate species that are absent in well-characterized Aβ soluble oligomeric preparations. Future studies should prioritize rigorous biochemical characterization of Aβ aggregation states in order to clarify the contributions of specific assemblies to neurotoxicity in SH-SY5Y cell models.

Limitations

A key limitation of this study stems from the absence of a standardized terminology to describe the experimental unit in the included studies. Despite our best efforts, it is possible that some technical replicates were misclassified as biological replicates. As outlined by Lazic et al. (2018), such misclassification can inflate the sample sizes and evidential strength of individual studies, potentially biasing the overall effect estimates (57). While we sought to mitigate this risk by avoiding the use of standardized mean differences, studies with artificially precise data are still likely to weigh more in a meta-analysis. Additionally, our approach to variable categorization in the meta-regression, particularly regarding Aβ species, may have introduced misclassification bias. Beyond instances where the aggregation state was explicitly unreported, it is possible that other preparations were incorrectly categorized due to inconsistencies or limitations in how aggregation states were described. This lack of precise characterization may have

affected the accuracy of subgroup comparisons and should be addressed in future primary studies through improved reporting and biochemical validation.

Another limitation of this study relates to the time frame of data collection. Due to feasibility constraints and the extended duration of data extraction and analysis, our literature search included only articles published up to October 2020. A follow-up search using the original search terms, limited to articles published from October 2020 onward, retrieved 3,550 articles in PubMed and 2,061 in Embase (see Supplementary Figure 5). Based on the inclusion rate observed in our review, we estimate that approximately 124 of these articles would have met our inclusion criteria, representing a potential 34% increase in the number of included comparisons. While we acknowledge this as a significant limitation, the large number of studies already included in our analysis and the robustness of the overall effect size suggest that the inclusion of more recent data would be unlikely to substantially alter our main conclusions.

Recommendations for future studies

Improved reporting of methodological details and clearer definitions of experimental units are essential for enhancing the transparency and reproducibility of *in vitro* research. To address these issues, we recommend the adoption of standardized, unambiguous terminologies for the classification of replicates and experimental design. Lazic et al. (2018) advocate for replacing ambiguous terms such as 'technical' and 'biological' replicates with more precise descriptors - namely, biological, experimental, and observational units - which help clarify the hierarchical structure of the experimental setup (57). In cell line studies, the use of standardized language specifying the variable components between experimental units (e.g., passage number, culture plate, or experimental day) would further aid in reproducibility and comparability across studies. Additionally, routine implementation of quality control measures - such as cell line

authentication and mycoplasma testing - is strongly encouraged, as these practices enhance the reliability and translational relevance of findings derived from cell-based models. For systematic reviews in particular, the development and dissemination of tools and guidelines for assessing risk of bias in *in vitro* studies would represent a valuable step toward improving the rigor and interpretability of evidence synthesis in this field.

Conclusions

Our meta-analysis confirms that Aβ exposure significantly reduces SH-SY5Y cell viability, as measured by the metabolic-based cell viability assays. The magnitude of this effect is primarily influenced by Aβ concentration, exposure duration, and aggregation state. In contrast, other factors such as differentiation status and cell density did not show consistent effects across studies. Our analysis also identified widespread methodological shortcomings, including inadequate reporting practices, lack of cell line authentication, and insufficient characterization of Aβ preparations. These limitations reduce the reliability and reproducibility of individual studies and may affect the interpretability of compiled findings. Overall, our findings support the continued use of SH-SY5Y cells as a relevant *in vitro* model for investigating Aβ-induced neurotoxicity. However, they also highlight the urgent need for improved methodological rigor, standardized reporting, and quality control measures to enhance the model's reliability and translational value in AD research.

List of abbreviations

AD – Alzheimer's Disease

AICc – Corrected Akaike Information Criteria

APP – Amyloid Precursor Protein

ATRA – All Trans Retinoic Acid

ATCC – American Type Culture Collection

Aβ – Beta-amyloid peptide

BDNF – Brain-Derived Neurotrophic Factor

CI – Confidence Interval

DMEM – Dulbecco's Modified Eagle Medium

ECACC – European Collection of Authenticated Cell Cultures

FBS – Fetal Bovine Serum

FCS – Fetal Calf Serum

FGF – Fibroblast growth factor

F12 – Nutrient Mixture F12

MTT – 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide

N – Number

N2 – N-2 Supplement

ROM – Ratio of Means

SEM – Standard Error of the Mean

SD – Standard Deviation

SMD – Standardized Mean Difference

Declarations

Acknowledgements

The authors acknowledge the contributions of the BRISA Consortium members to the planning, conduct, and interpretation of the results of this study. The authors would also like to thank Wolfgang Viechtbauer for input on the analysis.

Funding

NRP, GSC, GMA and GON received pre-doctoral fellowships from FAPESP (São Paulo Research Foundation – grants 18/10721-0, 22/12904-0 and 21/12263-2), CAPES (Coordination for the Improvement of Higher Education Personnel), or CNPq (The Brazilian National Council for Scientific and Technological Development). OBA is funded by FAPERJ (E-26/200.824/2021 and E-26/204.061/2024), CNPq (310813/2021-2) and the Serrapilheira Institute. AS is supported by grants from FAPESP (21/10925-8) and CNPq (productivity fellowship).

Availability of data and materials

All data generated and analyzed during this study are included in the manuscript, supporting information or available at indicated databases.

Authors contributions

Conceptualization, methodology, project administration, resources, supervision, validation, and visualization of the study were carried out by NRP, CFDC, GSC, OBA, and AS. These authors were also responsible for the original draft preparation as well as the critical review and editing of the manuscript. Funding acquisition was secured by OBA and AS. Statistical analyses were performed mainly by CFDC, with contributions from NRP, GSC, and OBA. All authors contributed to data curation and investigation and

reviewed the manuscript, collectively assuming responsibility for the decision to submit it for publication.

Competing interests

All authors declare that they have no competing interests.

References

1.   Hardy J, Selkoe DJ. The Amyloid Hypothesis of Alzheimer's Disease: Progress and Problems on the Road to Therapeutics. Science (1979) [Internet]. 2002;297(5580):353–6. Available from: https://www.science.org/doi/10.1126/science.1072994

2.   Cline EN, Bicca MA, Viola KL, Klein WL. The Amyloid-β Oligomer Hypothesis: Beginning of the Third Decade. Perry G, Avila J, Moreira PI, Sorensen AA, Tabaton M, editors. Journal of Alzheimer's Disease [Internet]. 2018;64(s1):S567–610. Available from: https://www.medra.org/servlet/aliasResolver?alias=iospress&doi=10.3233/JAD-179941

3.   Sehar U, Rawat P, Reddy AP, Kopel J, Reddy PH. Amyloid Beta in Aging and Alzheimer's Disease. Int J Mol Sci [Internet]. 2022;23(21):12924. Available from: https://www.mdpi.com/1422-0067/23/21/12924

4.   Querfurth HW, LaFerla FM. Alzheimer's Disease. New England Journal of Medicine [Internet]. 2010;362(4):329–44. Available from: http://www.nejm.org/doi/abs/10.1056/NEJMra0909142

5.   Velasco PT, Heffern MC, Sebollela A, Popova IA, Lacor PN, Lee KB, et al. Synapse-Binding Subpopulations of Aβ Oligomers Sensitive to Peptide Assembly Blockers and scFv Antibodies. ACS Chem Neurosci [Internet]. 2012;3(11):972–81. Available from: https://pubs.acs.org/doi/10.1021/cn300122k

6.   Benilova I, Karran E, De Strooper B. The toxic Aβ oligomer and Alzheimer's disease: an emperor in need of clothes. Nat Neurosci [Internet]. 2012;15(3):349–57. Available from: https://www.nature.com/articles/nn.3028

7.   Sebollela A, Mustata GM, Luo K, Velasco PT, Viola KL, Cline EN, et al. Elucidating Molecular Mass and Shape of a Neurotoxic Aβ Oligomer. ACS Chem Neurosci [Internet]. 2014;5(12):1238–45. Available from: https://pubs.acs.org/doi/10.1021/cn500156r

8.   De S, Wirthensohn DC, Flagmeier P, Hughes C, Aprile FA, Ruggeri FS, et al. Different soluble aggregates of Aβ42 can give rise to cellular toxicity through different mechanisms. Nat Commun [Internet]. 2019;10(1):1541. Available from: https://www.nature.com/articles/s41467-019-09477-3

9.   Michno W, Blennow K, Zetterberg H, Brinkmalm G. Refining the amyloid β peptide and oligomer fingerprint ambiguities in Alzheimer's disease: Mass spectrometric molecular characterization in brain, cerebrospinal fluid, blood, and plasma. J Neurochem. 2021 Oct 27;159(2):234–57.

10.   Jan A, Hartley DM, Lashuel HA. Preparation and characterization of toxic Aβ aggregates for structural and functional studies in Alzheimer's disease research. Nat Protoc [Internet]. 2010;5(6):1186–209. Available from: https://www.nature.com/articles/nprot.2010.72

11. Fontana IC, Zimmer AR, Rocha AS, Gosmann G, Souza DO, Lourenco M V, et al. Amyloid-β oligomers in cellular models of Alzheimer's disease. J Neurochem [Internet]. 2020;155(4):348–69. Available from: https://onlinelibrary.wiley.com/doi/10.1111/jnc.15030

12. Cummings JL, Morstorf T, Zhong K. Alzheimer's disease drug-development pipeline: few candidates, frequent failures. Alzheimers Res Ther [Internet]. 2014;6(4):37. Available from: https://alzres.biomedcentral.com/articles/10.1186/alzrt269

13. Mehta D, Jackson R, Paul G, Shi J, Sabbagh M. Why do trials for Alzheimer's disease drugs keep failing? A discontinued drug perspective for 2010-2015. Expert Opin Investig Drugs. 2017 Jun 3;26(6):735–9.

14. Agholme L, Lindström T, Kågedal K, Marcusson J, Hallbeck M. An In Vitro Model for Neuroscience: Differentiation of SH-SY5Y Cells into Cells with Morphological and Biochemical Characteristics of Mature Neurons. Journal of Alzheimer's Disease [Internet]. 2010;20(4):1069–82. Available from: https://www.medra.org/servlet/aliasResolver?alias=iospress&doi=10.3233/JAD-2010-091363

15. Forster JI, Köglsberger S, Trefois C, Boyd O, Baumuratov AS, Buck L, et al. Characterization of Differentiated SH-SY5Y as Neuronal Screening Model Reveals Increased Oxidative Vulnerability. SLAS Discovery [Internet]. 2016;21(5):496–509. Available from: https://linkinghub.elsevier.com/retrieve/pii/S247255522207099X

16. De Medeiros LM, De Bastiani MA, Rico EP, Schonhofen P, Pfaffenseller B, Wollenhaupt-Aguiar B, et al. Cholinergic Differentiation of Human Neuroblastoma SH-SY5Y Cell Line and Its Potential Use as an In vitro Model for Alzheimer's Disease Studies. Mol Neurobiol [Internet]. 2019;56(11):7355–67. Available from: http://link.springer.com/10.1007/s12035-019-1605-3

17. Biedler JL, Roffler-Tarlov S, Schachner M, Freedman LS. Multiple Neurotransmitter Synthesis by Human Neuroblastoma Cell Lines and Clones. Cancer Res [Internet]. 1978;38. Available from: files/42087/Biedler et al. - Multiple Neurotransmitter Synthesis by Human Neuroblastoma Cell Lines and Clones.pdf

18. Kovalevich J, Langford D. Considerations for the Use of SH-SY5Y Neuroblastoma Cells in Neurobiology. In: Amini S, White MK, editors. Neuronal Cell Culture [Internet]. Totowa, NJ: Humana Press; 2013. p. 9–21. Available from: https://link.springer.com/10.1007/978-1-62703-640-5_2

19. Encinas M, Iglesias M, Liu Y, Wang H, Muhaisen A, Ceña V, et al. Sequential Treatment of SH-SY5Y Cells with Retinoic Acid and Brain-Derived Neurotrophic Factor Gives Rise to Fully Differentiated, Neurotrophic Factor-Dependent, Human Neuron-Like Cells. J Neurochem [Internet]. 2000;75(3):991–1003. Available from: https://onlinelibrary.wiley.com/doi/10.1046/j.1471-4159.2000.0750991.x

20. Krishtal J, Bragina O, Metsla K, Palumaa P, Tõugu V. In situ fibrillizing amyloid-beta 1-42 induces neurite degeneration and apoptosis of differentiated SH-SY5Y cells. Ginsberg SD, editor. PLoS One [Internet]. 2017;12(10):e0186636. Available from: https://dx.plos.org/10.1371/journal.pone.0186636

21. Pedersen AK, Pfeiffer A, Karemore G, Akimov V, Bekker-Jensen DB, Blagoev B, et al. Proteomic investigation of Cbl and Cbl-b in neuroblastoma cell differentiation highlights roles for SHP-2 and CDK16. iScience [Internet]. 2021;24(4):102321. Available from: https://linkinghub.elsevier.com/retrieve/pii/S2589004221002893

22. Lacor PN, Buniel MC, Furlow PW, Sanz Clemente A, Velasco PT, Wood M, et al. Aβ Oligomer-Induced Aberrations in Synapse Composition, Shape, and Density Provide a Molecular Basis for Loss of Connectivity in Alzheimer's Disease. The Journal of Neuroscience [Internet]. 2007;27(4):796–807. Available from: https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.3501-06.2007

23. Alfonso S, Kessels HW, Banos CC, Chan TR, Lin ET, Kumaravel G, et al. Synapto-depressive effects of amyloid beta require <scp>PICK</scp> 1. European Journal of Neuroscience. 2014 Apr 8;39(7):1225–33.

24. Krishtal J, Metsla K, Bragina O, Tõugu V, Palumaa P. Toxicity of Amyloid-β Peptides Varies Depending on Differentiation Route of SH-SY5Y Cells. Journal of Alzheimer's Disease [Internet]. 2019;71(3):879–87. Available from: https://www.medra.org/servlet/aliasResolver?alias=iospress&doi=10.3233/JAD-190705

25. Seubert P, Vigo-Pelfrey C, Esch F, Lee M, Dovey H, Davis D, et al. Isolation and quantification of soluble Alzheimer's β-peptide from biological fluids. Nature [Internet]. 1992;359(6393):325–7. Available from: https://www.nature.com/articles/359325a0

26. Hu X, Crick SL, Bu G, Frieden C, Pappu R V, Lee JM. Amyloid seeds formed by cellular uptake, concentration, and aggregation of the amyloid-beta peptide. Proceedings of the National Academy of Sciences [Internet]. 2009;106(48):20324–9. Available from: https://pnas.org/doi/full/10.1073/pnas.0911281106

27. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. Syst Rev [Internet]. 2016;5(1):210. Available from: http://systematicreviewsjournal.biomedcentral.com/articles/10.1186/s13643-016-0384-4

28. Team RC. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2024. Available from: https://www.R-project.org/

29. Viechtbauer W. Conducting Meta-Analyses in *R* with the **metafor** Package. J Stat Softw [Internet]. 2010;36(3). Available from: http://www.jstatsoft.org/v36/i03/

30. Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome to the Tidyverse. J Open Source Softw [Internet]. 2019;4(43):1686. Available from: https://joss.theoj.org/papers/10.21105/joss.01686

31. Kossmeier M, Tran US, Voracek M. Power-Enhanced Funnel Plots for Meta-Analysis: The Sunset Funnel Plot. Z Psychol [Internet]. 2020;228(1):43–9. Available from: https://econtent.hogrefe.com/doi/10.1027/2151-2604/a000392

32. Calcagno V. glmulti: Model Selection and Multimodel Inference Made Easy [Internet]. 2020. Available from: https://CRAN.R-project.org/package=glmulti

33. Wickham H, Bryan J. R Packages: Organize, Test, Document, and Share Your Code [Internet]. O'Reilly Media; 2023. Available from: https://books.google.com.br/books?id=kTHFEAAAQBAJ

34. Burdick D, Soreghan B, Kwon M, Kosmoski J, Knauer M, Henschen A, et al. Assembly and aggregation properties of synthetic Alzheimer's A4/beta amyloid peptide analogs. Journal of Biological Chemistry [Internet]. 1992;267(1):546–54. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0021925818485298

35. Glenner GG, Wong CW. Alzheimer's disease: Initial report of the purification and characterization of a novel cerebrovascular amyloid protein. Biochem Biophys Res Commun [Internet]. 1984;120(3):885–90. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0006291X84801904

36. Walsh DM, Lomakin A, Benedek GB, Condron MM, Teplow DB. Amyloid β-Protein Fibrillogenesis. Journal of Biological Chemistry [Internet]. 1997;272(35):22364–72. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0021925819657030

37. Fanelli D. Negative results are disappearing from most disciplines and countries. Scientometrics [Internet]. 2012;90(3):891–904. Available from: http://link.springer.com/10.1007/s11192-011-0494-7

38. Fusenig NE, Capes-Davis A, Bianchini F, Sundell S, Lichter P. The need for a worldwide consensus for cell line authentication: Experience implementing a mandatory requirement at the International Journal of Cancer. PLoS Biol. 2017 Apr 17;15(4):e2001438.

39. Capes-Davis A, Theodosopoulos G, Atkin I, Drexler HG, Kohara A, MacLeod RAF, et al. Check your cultures! A list of cross-contaminated or misidentified cell lines. Int J Cancer [Internet]. 2010;127(1):1–8. Available from: https://onlinelibrary.wiley.com/doi/10.1002/ijc.25242

40. Whaley P, Blain RB, Draper D, Rooney AA, Walker VR, Wattam S, et al. Identifying assessment criteria for *in vitro* studies: a method and item bank. Toxicological Sciences. 2024 Oct 1;201(2):240–53.

41. Tran L, Tam DNH, Elshafay A, Dang T, Hirayama K, Huy NT. Quality assessment tools used in systematic reviews of in vitro studies: A systematic review. BMC Med Res Methodol. 2021 Dec 8;21(1):101.

42. Network the DIA, Barthélemy NR, Li Y, Joseph-Mathurin N, Gordon BA, Hassenstab J, et al. A soluble phosphorylated tau signature links tau, amyloid and the evolution of stages of dominantly inherited Alzheimer's disease. Nat Med [Internet]. 2020;26(3):398–407. Available from: https://www.nature.com/articles/s41591-020-0781-z

43. Mathisen GH, Svendsen C, Vist GE, Husøy T, Ames HM, Bearth A, et al. Identification of concepts of importance for the assessment of internal validity of *in vitro* toxicology studies using a modified Delphi technique. Evidence-Based Toxicology [Internet]. 2025;3(1):2551013. Available from: https://www.tandfonline.com/doi/full/10.1080/2833373X.2025.2551013

44. Figueiredo CP, Clarke JR, Ledo JH, Ribeiro FC, Costa C V, Melo HM, et al. Memantine Rescues Transient Cognitive Impairment Caused by High-Molecular-Weight A Oligomers But Not the Persistent Impairment Induced by Low-Molecular-Weight Oligomers. Journal of Neuroscience [Internet]. 2013;33(23):9626–34. Available from: https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.0482-13.2013

45. Price JL, Morris JC. Tangles and plaques in nondemented aging and ?preclinical? Alzheimer's disease. Ann Neurol [Internet]. 1999;45(3):358–68. Available from: https://onlinelibrary.wiley.com/doi/10.1002/1531-8249(199903)45:3<358::AID-ANA12>3.0.CO

46. Selkoe DJ, Hardy J. The amyloid hypothesis of Alzheimer's disease at 25 years. EMBO Mol Med [Internet]. 2016;8(6):595–608. Available from: https://www.embopress.org/doi/10.15252/emmm.201606210

47. Mc Donald JM, Savva GM, Brayne C, Welzel AT, Forster G, Shankar GM, et al. The presence of sodium dodecyl sulphate-stable Aβ dimers is strongly associated with Alzheimer-type dementia. Brain [Internet]. 2010;133(5):1328–41. Available from: https://academic.oup.com/brain/article-lookup/doi/10.1093/brain/awq065

48. Yang T, Hong S, O'Malley T, Sperling RA, Walsh DM, Selkoe DJ. New ELISAs with high specificity for soluble oligomers of amyloid β-protein detect natural Aβ oligomers in human brain but not CSF. Alzheimer's & Dementia [Internet]. 2013;9(2):99–112. Available from: https://alz-journals.onlinelibrary.wiley.com/doi/10.1016/j.jalz.2012.11.005

49. Gong Y, Chang L, Viola KL, Lacor PN, Lambert MP, Finch CE, et al. Alzheimer's disease-affected brain: Presence of oligomeric Aβ ligands (ADDLs) suggests a molecular basis for reversible memory loss. Proceedings of the National Academy of Sciences [Internet]. 2003;100(18):10417–22. Available from: https://pnas.org/doi/full/10.1073/pnas.1834302100

50. Marttinen M, Takalo M, Natunen T, Wittrahm R, Gabbouj S, Kemppainen S, et al. Molecular Mechanisms of Synaptotoxicity and Neuroinflammation in Alzheimer's Disease. Front Neurosci [Internet]. 2018;12:963. Available from: https://www.frontiersin.org/article/10.3389/fnins.2018.00963/full

51. Lopes GS, Lico DTP, Silva-Rocha R, De Oliveira RR, Sebollela A, Paçó-Larson ML, et al. A phylogenetically conserved hnRNP type A/B protein from squid brain. Neurosci Lett [Internet]. 2019;696:219–24. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0304394019300023

52. Giuffrida ML, Caraci F, Pignataro B, Cataldo S, De Bona P, Bruno V, et al. β-Amyloid Monomers Are Neuroprotective. The Journal of Neuroscience [Internet]. 2009;29(34):10582–7. Available from: https://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.1736-09.2009

53. Jeong H, Shin H, Hong S, Kim Y. Physiological Roles of Monomeric Amyloid-β and Implications for Alzheimer's Disease Therapeutics. Exp Neurobiol [Internet]. 2022;31(2):65–88. Available from: https://www.en-journal.org/journal/view.html?doi=10.5607/en22004

54. Tolar M, Hey J, Power A, Abushakra S. Neurotoxic Soluble Amyloid Oligomers Drive Alzheimer's Pathogenesis and Represent a Clinically Validated Target for Slowing Disease Progression. Int J Mol Sci [Internet]. 2021;22(12):6355. Available from: https://www.mdpi.com/1422-0067/22/12/6355

55. Lasagna-Reeves CA, Glabe CG, Kayed R. Amyloid-β Annular Protofibrils Evade Fibrillar Fate in Alzheimer Disease Brain. Journal of Biological Chemistry [Internet]. 2011;286(25):22122–30. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0021925819489199

56. Bigi A, Cascella R, Chiti F, Cecchi C. Amyloid fibrils act as a reservoir of soluble oligomers, the main culprits in protein deposition diseases. BioEssays [Internet]. 2022;44(11):2200086. Available from: https://onlinelibrary.wiley.com/doi/10.1002/bies.202200086

57. Lazic SE, Clarke-Williams CJ, Munafò MR. What exactly is 'N' in cell culture and animal experiments? PLoS Biol [Internet]. 2018;16(4):e2005282. Available from: https://dx.plos.org/10.1371/journal.pbio.2005282