Conspecific attraction in songbirds during spring migration

André Desrochers ⓞ
Sciences du bois et de la forêt
Université Laval
Québec, Qc, G1V 0A6
andre.desrochers@sbf.ulaval.ca

Laetitia Desbordes

Observatoire d'oiseaux de Tadoussac
Les Bergeronnes, Qc, G0T 1G0
ladesbordes@hotmail.fr

November 5, 2025

ABSTRACT

Conspecific attraction has often been hypothesized as a mechanism that could facilitate migration and territory acquisition. We studied conspecific attraction by songbirds during spring migration between 2019 and 2025 at the Dunes of Tadoussac, Quebec, Canada. This location is known to produce spectacular migration events with thousands of passerine birds flying at close range from the observers. We recorded sequences of individual birds moving just over the ground in a ~20 m wide corridor along the edge of the St. Lawrence estuary. We also conducted hourly counts to detect conspecific clusters at a coarser temporal scale. Birds were much more likely to be following conspecifics at close range than other species, as evidenced by randomization tests, after accounting for species composition and abundance during each specific migration event. Conspecific sequences were mostly of two individuals, but much larger sequences occurred, up to 103 consecutive individuals of the same species. We found no evidence of leading species, i.e. species that were more likely to lead, rather than follow, another species. Hourly species counts were often much more variable than expected from a random (Poisson) process, at least in the case of Tennessee and Magnolia Warblers, providing evidence for conspecific attraction at a coarser temporal scale. This study provides the first detailed evidence of conspecific attraction with passerine birds during migration.

Keywords migration • conspecific attraction • Tadoussac • Quebec • randomization test • warblers • bird count • Poisson distribution

1 Introduction

Bird migrations often assemble birds of different species in stopover sites and along migration corridors (Gauthreaux 1972; Cohen and Satterfield 2020). This phenomenon, termed *co-migration*, fosters interspecific interactions, leading to nonrandom species assemblages (Van Doren et al. 2025; DeSimone et al. 2024). Those interspecific interactions may be agonistic (Recher and Recher 1969; Novcic 2018), occasionally lethal (Ydenberg et al. 2004; Furey et al. 2018), and potentially detrimental to populations (Ydenberg, Tavera, and Lank 2022). However, co-migration can also confer benefits, as interactions may yield useful information, whether inadvertently or intentionally (Danchin et al. 2004; Simons 2004). Research has long demonstrated predator avoidance advantages in flocking birds during daytime (Morse 1970; Pulliam 1973; Lima and Dill 1990). More recently, research found evidence of acoustic communication and interpreted it as a potential benefit during night migration, especially in challenging conditions such as poor

visibility or obstacles (Gayk and Mennill 2023; DeSimone et al. 2024; Hüppop and Hilgerloh 2012). In daylight, following conspecifics or heterospecifics en route to breeding grounds may similarly share navigational cues (Couzin et al. 2005). For example, groups of soaring hawks ('kettles') may provide social information to multiple species, in this case, about the location of thermal currents that reduce energy expenditure (Leshem and YomTov 1996; Kerlinger and Gauthreaux 1985).

Given migration's inherent risks (Sillett and Holmes 2002), birds likely leverage available cues — especially from conspecifics — to enhance navigation, speed, and stopover site selection. Yet, the social dimensions of migratory strategies remain underexplored (Schofield et al. 2018; DeSimone et al. 2024). While heterospecific information exchange occurs (Seppänen et al. 2007), it is unclear whether conspecifics provide more useful information than heterospecifics, particularly during active flight. If conspecifics offer distinct advantages, we expect aggregations of the latter within co-migrating flocks. We know conspecific aggregation exists within migrating mixed-species flocks of waterfowl (Eddleman, Patterson, and Knopf 1985; White and James 1978), waders (Erwin 1983; Folmer, Olff, and Piersma 2010) and parrots (Ferdinand et al. 2023). But the presence of conspecific aggregations remains poorly documented in mixed-species flocks of passerines, within or outside migration. Gauthreaux (1972) observed aggregations of conspecifics in 17 species during co-migration in the southeastern United States, but lack of detail prevents a distinction between deliberate and fortuitous clustering. More recently, Gayk et al. (2023) found that temporally adjacent nocturnal migrants in northern Ontario make similar nocturnal flight calls, suggesting conspecific aggregations.

Social interactions between individuals are notoriously difficult to study in the case of co-migrating passerines, because of their tendency to move in loose flocks, mostly at night. However, certain locations along migration routes funnel movements, thus facilitating the observation of possible species aggregations. One such location is the northern coast of the St. Lawrence river estuary, Quebec, Canada (Ibarzabal, Côté, and Drolet 2009). In this location, we tested whether co-migrating passerines formed conspecific aggregations during spring migration events, and whether certain species played a leading role during uninterrupted flights. We looked for conspecific attraction at a fine temporal scale (seconds) and a coarser temporal scale (hours).

2 Methods

2.1 Study Area

We carried out this study at the dunes of Tadoussac, Québec, Canada (48.157°N, 69.665°W). This site is on the northern shore of the St. Lawrence river estuary, and is a patchwork of open sandy areas interspersed with deciduous (*Betula sp.*, *Populus sp.*) and coniferous (*Pinus sylvestris*, *Picea glauca*) stands, approximately 50 m above sea level. The dunes of Tadoussac are known as a major hawk migration corridor (Ibarzabal 1999; Brisson-Curadeau, Elliott, and Côté 2020), as well as their spring migration events during which well over 10,000 passerines can sometimes be observed daily, especially near the end of the season (Figure 1). Most of the time, birds fly overhead 10 to 100 m above ground, sometimes they fly over the water, and sometimes birds fly just above ground in a very narrow corridor in the steep slope that borders the estuary. In the spring, flights are generally part of a 'redetermined' migration because most birds fly along the coast in a southwest direction, i.e. contrary to their expected direction (Gauthreaux 1978).

2.2 Field methods

We conducted all observations from the top of the slope bordering the estuary. The observation site was situated just above a part of the slope devoid of vegetation, thus facilitating the observation of individual birds, mostly at distances from 10 to 20 m from the observers (Figure 2).

Redetermined migration events (thereafter, 'migration events' or 'events') are frequent on the site (Tadoussac Bird Observatory, unpubl. data). Birds during those events often fly ~ 50 m or higher above ground, in a > 100 m wide corridor. In other cases, they fly hundreds of meters offshore. In the latter cases, most of the individuals are unidentifiable to species. However, when strong winds from north or northwest occur, moving birds tend to fly just above ground in the steep slope facing the estuary, along a ~ 20 m wide path. This 'river' of birds enables the measurement of sequences of individuals identified to species.

We recorded sequences of birds on 12 migration events, each in a separate date, in late May or early June 2019, 2020, 2022, 2024, and 2025. Observations were all performed between 05:00 and 18:05 (EDT) and lasted from 0.8 to 4.6 hours. Each time, two experienced ornithologists recorded sequences of individual birds as they flew from left (NE) to right (SW). One of the ornithologists (coauthor LD) identified each bird, either by naked eye or with binoculars, and named the species or genus out loud. Most species examined here exhibit conspicuous field marks which facilitate migration. Despite this, there was insufficient time in most cases to reliably determine the sex of the bird, and in 12 percent of cases we were unable to identify to species. The second observer (coauthor AD) recorded each bird identification on a smartphone with a CyberTracker (Steventon et al. 2011; Liebenberg et al. 2017) custom-made

Figure 1: Redetermined migration events at the dunes of Tadoussac, Québec, Canada. Shaded areas are local spring migration seasons for most passerines (15 April - 5 June).

(a) Bird observation area. (b) American Redstart in flight.

Figure 2: Scenes from the study area.

application showing species acronyms on the screen, requiring just one tap to log species or taxon, with an exact timestamp. This method enabled data collection at rates exceeding 1 bird per second. Despite this, observers were sometimes overwhelmed by large numbers of bird, leading to sequences of unidentified birds.

In addition to the measurement of bird sequences, we performed hourly counts of all birds identifiable to species, either flying low or overhead. We retained only counts from the species observed also during our species sequence sessions (above), but retained counts on all dates, irrespective of migration events.

2.3 Statistical analysis

We performed all data preparation and analysis with R/RStudio (R Studio Team 2024; R Core Team 2023). We collapsed all recorded birds into a table with each line representing one individual and its exact passage time. We excluded non-passerine birds, leaving 87783 birds for analysis (Table 1).

For each of the migration events, we counted the number of cases where an individual bird was followed by a conspecific. We generated an empirical random distribution of the number of conspecific sequences for each event by reshuffling data 9,999 times. This procedure thus accounts for unequal species counts and missing data. We consid-

Table 1: Species and numbers of individuals analysed in redetermined migration events, dunes of Tadoussac, Québec.

Scientific	Common Name	Code	Number
Cardellina canadensis	Canada Warbler	CAWA	14
Cardellina pusilla	Wilson's Warbler	WIWA	50
Catharus fuscescens	Veery	VEER	6
Catharus guttatus	Hermit Thrush	HETH	4
Catharus minimus	Gray-cheeked Thrush	GCTH	3
Catharus ustulatus	Swainson's Thrush	SWTH	134
Empidonax alnorum	Alder Flycatcher	ALFL	14
Empidonax flaviventris	Yellow-bellied Flycatcher	YBFL	30
Empidonax minimus	Least Flycatcher	LEFL	37
Geothlypis philadelphia	Mourning Warbler	MOWA	6
Leiothlypis peregrina	Tennessee Warbler	TEWA	482
Leiothlypis ruficapilla	Nashville Warbler	NAWA	29
Mniotilta varia	Black-and-white Warbler	BAWW	13
Parkesia noveboracensis	Northern Waterthrush	NOWA	11
Seiurus aurocapilla	Ovenbird	OVEN	6
Setophaga americana	Northern Parula	NOPA	80
Setophaga caerulescens	Black-throated Blue Warbler	BTBW	42
Setophaga castanea	Bay-breasted Warbler	BBWA	615
Setophaga coronata	Yellow-rumped Warbler	YRWA	3807
Setophaga fusca	Blackburnian Warbler	BLBW	103
Setophaga magnolia	Magnolia Warbler	MAWA	710
Setophaga palmarum	Palm Warbler	PAWA	4
Setophaga pensylvanica	Chestnut-sided Warbler	CSWA	19
Setophaga ruticilla	American Redstart	AMRE	159
Setophaga striata	Blackpoll Warbler	BLPW	44
Setophaga tigrina	Cape May Warbler	CMWA	899
Setophaga virens	Black-throated Green Warbler	BTNW	102
Vireo olivaceus	Red-eyed Vireo	REVI	15
Vireo philadelphicus	Philadelphia Vireo	PHVI	47
Vireo solitarius	Blue-headed Vireo	BHVI	5

ered observed counts of conspecific sequences as significantly clustered if their quantile was > 0.995, thus accounting for multiple testing.

To determine if species counts per hour are more clustered than expected under a random (Poisson) distribution, while accounting for varying total counts across hours, we used a Monte Carlo method using Variance-to-Mean Ratios (VMR). We retained all survey dates with at least 5 hourly counts, and one species or mode with at least 10 individuals. For each hour across the days with sufficient data (n = 84), we calculated the observed VMR of counts for each species. We built a null empirical distribution by generating Poisson-distributed counts, accounting for hourly variation of total counts (λ) for all species, repeating this 999 times to compute simulated VMRs. We calculated the quantile of the observed VMR in the corresponding null distribution, i.e., the proportion of simulated VMRs that were greater than the observed VMR. We interpreted quantiles > 0.95 as evidence for clustering.

3 Results

3.1 Sequences of birds

On average, we observed 939 birds per session (range: [113, 3565]). During observation sessions, an average of 6.2 birds/min flowed, but it was highly variable, with long pauses interrupted with intense flows, in some cases exceeding 50 birds/min (Figure 3).

In each migration event, conspecific sequences were significantly more frequent (P < 0.005) than expected if the birds, in the same frequency distribution of species, were mixed independently (Figure 4). Conspecific sequences were often

Figure 3: Flow of birds during migration redetermination events, Tadoussac, Québec.

more than 2 birds (max 103 conspecifics in a single run), and the number of 2-bird conspecific sequences (694) was significantly smaller than that expected from a Poisson (random) process (971; $\chi^2 = 22.9$, $P < 0.001$).

Given the large number of species, interspecific sequences were very variable and most of them occurred a small number of times. However, species sequences that occurred 30 times or more were not asymmetric; that is, there was no evidence that species exerted net leader or follower roles (Table 2).

3.2 Hourly clustering

Twenty of the 24 species exhibited significant clustering of hourly counts, at least on some days (Table 3), but with strong day-to-day variation. The tendency for clustering by hours was consistent only for Tennessee Warbler and Magnolia Warbler (Table 3).

Figure 4: Numbers of conspecific sequences observed vs. expected by chance, on all observation dates. Grey histograms are empirical density functions of conspecific sequences based on the null hypothesis (species independent), and red arrows show actual counts of conspecific sequences. After each date, numbers of species and individuals are shown.

Table 2: Sequences of birds of different species. Only cases with 30+ sequences are shown. with P-values calculated with Fisher's exact test (two-tailed, null hypothesis of equal frequencies).

		Species 1...		
Species 1	Species 2	Leading	Following	P
American Redstart	Cape May Warbler	20	23	0.83
American Redstart	Magnolia Warbler	30	24	0.70
American Redstart	Yellow-rumped Warbler	33	24	0.45
Bay-breasted Warbler	Cape May Warbler	68	76	0.72
Bay-breasted Warbler	Magnolia Warbler	67	61	0.80
Bay-breasted Warbler	Tennessee Warbler	45	54	0.57
Bay-breasted Warbler	Yellow-rumped Warbler	165	164	1.00
Black-throated Blue Warbler	Yellow-rumped Warbler	19	16	0.81
Black-throated Green Warbler	Yellow-rumped Warbler	25	18	0.52
Blackburnian Warbler	Yellow-rumped Warbler	27	26	1.00
Cape May Warbler	Magnolia Warbler	74	70	0.91
Cape May Warbler	Tennessee Warbler	45	38	0.64
Cape May Warbler	Yellow-rumped Warbler	278	313	0.32
Magnolia Warbler	Tennessee Warbler	32	25	0.57
Magnolia Warbler	Yellow-rumped Warbler	176	176	1.00
Northern Parula	Yellow-rumped Warbler	20	19	1.00
Swainson's Thrush	Yellow-rumped Warbler	26	19	0.53
Tennessee Warbler	Yellow-rumped Warbler	116	112	0.93

Table 3: Conspecific clustering by hour, sorted by decreasing order. Days are numbers of dates with at least 10 individuals of the species observed. Clustering is measured as Variance-to-Mean Ratios (VMRs) with their 95 % limits. A VMR of 1.0 would be expected if hourly counts were random (Poisson distributed). Days clustered are numbers of days where a significant clustering occurred. A paired-sampled, one-tailed Wilcoxon test was applied in cases with at least 10 days, to test if species exhibited consistent hourly clustering.

Species	Days	Variance to Mean Ratio	Days clustered	Wilcoxon P
Bay-breasted Warbler	19	66.8 [0, 147]	11	0.07
Tennessee Warbler	41	62.2 [0, 135]	16	< 0.001
Yellow-rumped Warbler	75	32.1 [12, 52]	14	0.29
Cape May Warbler	26	23.1 [9.6, 37]	9	0.22
Magnolia Warbler	22	19.3 [0, 39]	9	0.03
Blackburnian Warbler	10	19.2 [0, 50]	5	0.22
American Redstart	8	15.8 [0, 37]	6	
Northern Parula	5	11.9 [0, 28]	1	
Least Flycatcher	3	8.4 [0, 29]	1	
Blackpoll Warbler	3	8.2 [0, 33]	0	
Black-throated Green Warbler	11	8.2 [1.5, 15]	3	0.21
Swainson's Thrush	9	7.9 [4.4, 11]	6	
Wilson's Warbler	4	7.8 [0, 19]	2	
Red-eyed Vireo	1	6.7	1	
Yellow-bellied Flycatcher	1	6.2	1	
Philadelphia Vireo	3	5.7 [0, 14]	2	
Nashville Warbler	8	4.8 [1, 9]	2	
Chestnut-sided Warbler	4	4.6 [0, 9]	2	
Black-throated Blue Warbler	5	4.5 [0, 10]	1	
Orange-crowned Warbler	1	4.3	1	
Mourning Warbler	1	3.4	1	
Northern Waterthrush	2	2.5 [0, 14]	0	
Black-and-white Warbler	2	2.5 [1.4, 4]	0	
Canada Warbler	3	2.4 [1.2, 4]	0	

4 Discussion

Given that most migratory songbirds exhibit species-specific peak migration dates, conspecific clustering at coarse temporal scales (e.g., weeks) is expected. But to our knowledge, this is the first detailed evidence of conspecific attraction at finer temporal scales, hours and seconds, during migration. On given days, overall migration intensity usually varied markedly among hours, but we found that single-species numbers often varied even more, providing evidence for conspecific attraction at the temporal resolution of hours. At a finer, behavioral, resolution of seconds, conspecific attraction was evident, and occurred on each opportunity we had to observed precise sequences of birds actively flying.

Most of the literature on conspecific attraction has focused on the breeding season, following the pioneering study by Stamps (1988). While poorly documented, conspecific attraction during migration has often been inferred from casual or indirect observations. Gauthreaux speculated that conspecifics possibly gather in flocks at dawn before completion of their solo flight across the Gulf of Mexico (Gauthreaux 1972). Balcomb also believed that most passerine birds flew singly at night (Balcomb 1977). An early study in Louisiana suggested the occurrence of conspecific attraction during landfall at dawn (Gauthreaux 1972). Hamilton (1962) and Ball (1952) also came to the conclusion of conspecific attraction triggered or maintained by calls. A more recent study (Gayk and Mennill 2023) documented spatial aggregation of similar nocturnal flight calls in Ontario, again suggesting that passerines of the same species flock together during nocturnal migration.

As for the hourly clustering, its sporadic occurrence in most species could reflect opportunistic responses to temporary cues like food, weather, or predation risk. The consistent clustering in Tennessee and Magnolia and possibly, Bay-breasted Warblers might indicate stronger social drivers.

Our evidence for conspecific attraction at the scales of seconds and hours is novel, and before before attempting to place it in a conceptual frame, replication will be needed. For now, we can only speculate on the mechanisms leading to those patterns. One plausible explanation would be the facilitation of information sharing among individuals with similar ecological needs (Couzin et al. 2005; Leshem and YomTov 1996; Kerlinger and Gauthreaux 1985). For instance, collective decision-making in conspecific flocks could enhance the detection of favorable winds or food patches, as suggested by studies on collective navigation in migratory birds (Simons 2004). In contrast, heterospecific flocks might introduce conflicting cues due to differing migratory strategies or destinations, reducing the reliability of group-level decisions.

From an evolutionary perspective, conspecific attraction during migration may also reflect selection pressures related to territory acquisition and more generally, breeding preparation. Migratory songbirds, particularly males, often arrive at breeding grounds earlier than females - a phenomenon known as protandry (Morbey and Ydenberg 2001) — and establishing conspecific networks en route could facilitate territory defense or mate attraction upon arrival. Flying with conspecifics might allow individuals to assess competitors or form alliances.

The preference for conspecific over heterospecific associations contrasts with some evidence of mixed-species flocks during migration, particularly at stopover sites (Goodale et al. 2010). These heterospecific groupings are often attributed to predator avoidance or resource competition, benefits that may be more pronounced during stationary phases than active flight. Our findings, focused on in-flight behavior, suggest that the costs of coordinating with heterospecifics—such as mismatched flight speeds or navigational goals—may outweigh these advantages during transit.

Several limitations in our study should be addressed in future research. First, we were unable to determine the sex of most birds, including those with sexually dimorphic plumages thus precluding inference about birds flying in pairs, or comparisons of conspecific attraction within sex. Second, we can only speculate whether conspecific attraction was driven by navigation, communication, or social mechanisms. Experiments manipulating flock composition or simulating conspecific cues could clarify these drivers.

We were unable to follow individual birds for more than a few seconds, thus we cannot determine how ephemeral are species aggregations. It is possible that some individuals fly with their mate, which could explain the tendency of conspecifics to be together. However, two observations point against this explanation. First, there were not more sequences of two conspecifics than expected by chance - in fact there were significantly fewer. Furthermore, as noted earlier, males usually migrate sooner in the season than females in many passerine species (Morbey and Ydenberg 2001).

In conclusion, our finding that songbirds preferentially fly with conspecifics during migration underscores the importance of species-specific sociality in shaping migratory behavior. This conspecific attraction likely enhances navigational efficiency, reinforces communication, and prepares individuals for breeding, offering a selective advantage in a challenging life-history stage. These insights contribute to our understanding of social dynamics in migratory birds

and open avenues for exploring how such behaviors influence population connectivity and resilience. As migratory songbirds face increasing threats from habitat loss and climate change, unraveling these social strategies may also inform conservation efforts, ensuring that critical group behaviors are preserved across their annual cycle.

5 Acknowledgments

We thank birders at the Observatoire d'oiseaux de Tadoussac, who not only provided help by finding birds, but also entertainment from lively discussions in those frequent moments when no birds are to be found. Thanks especially to Alexandre Terrigeol, Pascal Côté, and Jean-François Therrien, for comments on earlier versions of this manuscript. Funding for this study was provided by Université Laval department funds, and Explos-Nature.

References

Balcomb, Richard. 1977. "The Grouping of Nocturnal Passerine Migrants." *The Auk* 94 (3): 479–88.

Ball, Stanley C. 1952. *Fall Bird Migration on the Gaspé Peninsula*. Yale University Bulletin.

Brisson-Curadeau, Émile, Kyle H. Elliott, and Pascal Côté. 2020. "Factors Influencing Fall Departure Phenology in Migratory Birds That Bred in Northeastern North America." *The Auk* 137 (1): ukz064.

Cohen, Emily B., and Dara A. Satterfield. 2020. "'Chancing on a Spectacle:' Cooccurring Animal Migrations and Interspecific Interactions." *Ecography* 43 (11): 1657–71.

Couzin, Iain D., Jens Krause, Nigel R. Franks, and Simon A. Levin. 2005. "Effective Leadership and Decision-Making in Animal Groups on the Move." *Nature* 433 (7025): 513–16.

Danchin, Étienne, Luc-Alain Giraldeau, Thomas J. Valone, and Richard H. Wagner. 2004. "Public Information: From Nosy Neighbors to Cultural Evolution." *Science* 305 (5683): 487.

DeSimone, Joely G., Lucaske W. DeGroote, Stuart A. MacKenzie, Jennifer C. Owen, Andrea J. Patterson, and Emily B. Cohen. 2024. "Persistent Species Relationships Characterize Migrating Bird Communities Across Stopover Sites and Seasons." *Proceedings of the National Academy of Sciences* 121 (34): e2322063121.

Eddleman, William R., Craig T. Patterson, and Fritz L. Knopf. 1985. "Interspecific Relationships Between American Coots and Waterfowl During Fall Migration." *The Wilson Bulletin* 97 (4): 463–72.

Erwin, R. Michael. 1983. "Feeding Habitats of Nesting Wading Birds: Spatial Use and Social Influences." *The Auk* 100 (4): 960–70.

Ferdinand, Vanessa, Elle Pattenden, Donald J. Brightsmith, and Elizabeth A. Hobson. 2023. "Inferring the Decision Rules That Drive Co-Foraging Affiliations in Wild Mixed-Species Parrot Groups." *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* 378 (1878): 20220101.

Folmer, Eelke O, Han Olff, and Theunis Piersma. 2010. "How Well Do Food Distributions Predict Spatial Distributions of Shorebirds with Different Degrees of Selforganization?" *Journal of Animal Ecology* 79 (4): 747–56.

Furey, Nathan B., Jonathan B. Armstrong, David A. Beauchamp, and Scott G. Hinch. 2018. "Migratory Coupling Between Predators and Prey." *Nature Ecology & Evolution* 2 (12): 1846–53.

Gauthreaux, Jr., Sidney A. 1972. "Behavioral Responses of Migrating Birds to Daylight and Darkness: A Radar and Direct Visual Study." *The Wilson Bulletin*, 136–48.

———. 1978. "Importance of the Daytime Flights of Nocturnal Migrants: Redetermined Migration Following Displacement." In *Animal Migration, Navigation, and Homing: Symposium Held at the University of Tübingen, August 17–20, 1977*, 219–27. Springer.

Gayk, Zach G., and Daniel J. Mennill. 2023. "Acoustic Similarity of Flight Calls Corresponds with the Composition and Structure of Mixed-Species Flocks of Migrating Birds: Evidence from a Three-Dimensional Microphone Array." *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 378 (1878): 20220114.

Goodale, Eben, Guy Beauchamp, Robert D. Magrath, James C. Nieh, and Graeme D. Ruxton. 2010. "Interspecific Information Transfer Influences Animal Community Structure." *Trends in Ecology & Evolution* 25 (6): 354–61.

Hamilton, William J. 1962. "Evidence Concerning the Function of Nocturnal Call Notes of Migratory Birds." *The Condor* 64 (5): 390–401.

Hüppop, Ommo, and Gudrun Hilgerloh. 2012. "Flight Call Rates of Migrating Thrushes: Effects of Wind Conditions, Humidity and Time of Day at an Illuminated Offshore Platform." *Journal of Avian Biology* 43 (1): 85–90.

Ibarzabal, Jacques. 1999. "Tadoussac: Un Site de Migration Des Oiseaux de Proie." *Le Naturaliste Canadien* 123 (1): 11–18.

Ibarzabal, Jacques, Pascal Côté, and Bruno Drolet. 2009. "Tadoussac: Sentinelle Ornithologique de La Forêt Boréale Au Parc National Du Saguenay." *Le Naturaliste Canadien* 133 (3): 27–31.

Kerlinger, Paul, and Jr. Gauthreaux Sidney A. 1985. "Seasonal Timing, Geographic Distribution, and Flight Behavior of Broad-Winged Hawks During Spring Migration in South Texas: A Radar and Visual Study." *The Auk* 102 (4): 735–43.

Leshem, Yossi, and Yoram YomTov. 1996. "The Use of Thermals by Soaring Migrants." *Ibis* 138 (4): 667–74.

Liebenberg, Louis, Justin Steventon, Nate Brahman, Karel Benadie, James Minye, and Horekhwe Karoha Langwane. 2017. "Smartphone Icon User Interface Design for Non-Literate Trackers and Its Implications for an Inclusive Citizen Science." *Biological Conservation* 208: 155–62.

Lima, Steven L., and Larry M. Dill. 1990. "Behavioral Decisions Made Under the Risk of Predation: A Review and Prospectus." *Canadian Journal of Zoology* 68 (4): 619–40.

Morbey, Yolanda E., and Ronald C. Ydenberg. 2001. "Protandrous Arrival Timing to Breeding Areas: A Review." *Ecology Letters* 4 (6): 663–73.

Morse, Douglass H. 1970. "Ecological Aspects of Some Mixed-Species Foraging Flocks of Birds." *Ecological Monographs* 40 (1): 119–68.

Novcic, Ivana. 2018. "Aggression in Flocks of Foraging Shorebirds During Spring Stopover at Delaware Bay, USA." *Waterbirds* 41 (1): 73–79.

Pulliam, H. Ronald. 1973. "On the Advantages of Flocking." *Journal of Theoretical Biology* 38: 419–22.

R Core Team. 2023. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

R Studio Team. 2024. *RStudio: Integrated Development for r. RStudio, Inc., Boston, MA URL Http://Www.rstudio.com/*.

Recher, Harry F, and Judy A Recher. 1969. "Some Aspects of the Ecology of Migrant Shorebirds. II. Aggression." *The Wilson Bulletin*, 140–54.

Schofield, Lynn N., Jill L. Deppe, Theodore J. Zenzal, Michae lP. Ward, Robert H. Diehl, Rachel T. Bolus, and Frank R. Moore. 2018. "Using Automated Radio Telemetry to Quantify Activity Patterns of Songbirds During Stopover." *The Auk: Ornithological Advances* 135 (4): 949–63.

Seppänen, Janne-Tuomas T., Jukka T. Forsman, Mikko Mönkkönen, and Robert L. Thomson. 2007. "Social Information Use Is a Process Across Time, Space, and Ecology, Reaching Heterospecifics." *Ecology* 88 (7): 1622–33.

Sillett, T. Scott, and Richard T. Holmes. 2002. "Variation in Survivorship of a Migratory Songbird Throughout Its Annual Cycle." *Journal of Animal Ecology* 71 (2): 296–308.

Simons, Andrew M. 2004. "Many Wrongs: The Advantage of Group Navigation." *Trends in Ecology & Evolution* 19 (9): 453–55.

Stamps, J. A. 1988. "Conspecific Attraction and Aggregation in Territorial Species." *American Naturalist* 131: 329–47.

Steventon, Justin, Louis Liebenberg, M. Derbecker, V. Bapat, and D. G. Miles. 2011. *CyberTracker. Available from: Www.cybertracker.org*.

Van Doren, Benjamin M., Joely G. DeSimone, Josh A. Firth, Friederike Hillemann, Zach Gayk, Emily Cohen, and Andrew Farnsworth. 2025. "Social Associations Across Species During Nocturnal Bird Migration." *Current Biology* 35 (4): 898–904.e4.

White, Donald H., and Douglas James. 1978. "Differential Use of Fresh Water Environments by Wintering Waterfowl of Coastal Texas." *The Wilson Bulletin* 90 (1): 99–111.

Ydenberg, Ronald C., Robert W. Butler, David B. Lank, Barry D. Smith, and John Ireland. 2004. "Western Sandpipers Have Altered Migration Tactics as Peregrine Falcon Populations Have Recovered." *Proceedings of the Royal Society B: Biological Sciences* 271: 1263–69.

Ydenberg, Ronald C., Eveling A. Tavera, and David B. Lank. 2022. "Danger, Risk and Antipredator Behavior in the Life History of Longdistance Migratory Sandpipers." *Journal of Avian Biology* 2022 (6): e03002.