1   **Habitat and complex life cycles promote morphological diversity in salamander limb bones**

2   Jonathan M. Huie[1,2], R. Alexander Pyron[1], Sandy M. Kawano[1,3]

3   [1] Department of Biological Sciences, The George Washington University, Washington, DC, USA
4   [2] Department of Ecology and Evolution, University of California Irvine, Irvine, CA, USA
5   [3] Department of Biology, University of Virginia, Charlottesville, VA, USA
6

7   Corresponding Author: Jonathan M. Huie
8   Corresponding Author's Email: jonathanmhuie@gmail.com
9

10   Running Title: Habitat and life cycle shape salamander limb bones

ABSTRACT

Salamander evolution featured multiple transitions between water and land that likely

promoted distinct adaptations in limb bones for buoyancy control versus increased load-bearing

capacities, respectively. Many extant species spend their entire lives either in water or on land,

while, others undergo water-land transitions within their lifetime. However, exposure to both

environments may impose competing demands that restrict adaptive evolution for a particular

habitat. Using a 3D morphological dataset of 133 species spanning the phylogenetic and

ecological breadth of salamanders, we find that the external and internal morphology of limb

bones are evolutionarily decoupled, which increases the evolvability of limb bones in response to

diverse mechanical demands. Terrestrial salamanders have stiffer bones with greater resistance to

fracture, while aquatic species have denser bones that are hypothesized to aid in buoyancy

regulation. We uncover a functional trade-off between stiffness and density that promotes stiff

yet lightweight bones in terrestrial lineages. Released from terrestrial constraints, aquatic

paedomorphs have disparate external morphologies, whereas terrestrial direct developers

consistently share a rod-like bone shape. Aquatic and terrestrial multiphasic taxa show less

morphological divergence than monophasic species living in comparable habitats but are not

constrained by their complex life cycle. Multiphasic species have distinct external limb bone

shapes that have evolved as fast or faster than monophasic species. Taken together, we propose

that the trade-offs imposed by different habitats and complex life cycles increase limb bone

diversity by promoting alternate evolutionary pathways.

Keywords: adaptation, evolution, habitat, locomotion, trade-off, water-to-land transition

INTRODUCTION

Habitat transitions expose animals to new physical demands that often promote adaptive changes to the musculoskeletal system (Easterling et al. 2022; Koehl 2023). However, many animals have complex life cycles, which involve ecological shifts over ontogeny that expose them to conflicting selective pressures (Hanken 1992; Laudet 2011; McMenamin and Parichy 2013; Truman 2019). While metamorphosis potentially decouples selective pressures across life stages (Ebenman 1992; Hanken 1992; Moran 1994), complex life cycles may limit overall adaptive responses if the different life stages occupy habitats with dramatically different demands. For example, water-land transitions involve dramatic ecological shifts with profound consequences for the anatomy and function of limbs (Bejder and Hall 2002; Pierce et al. 2020; Dickson et al. 2021; Molnar et al. 2021a). Compared to terrestrial habitats, limbs used underwater experience lower mechanical loads due to positive buoyancy and the use of locomotor modes that minimize limb-substrate interactions, such as axial-based swimming (Azizi and Horton 2004; Young and Blob 2015; Gamel et al. 2024). In contrast, terrestrial locomotion requires the limbs to support body weight against gravity and generate larger propulsive forces than aquatic locomotion that can exceed body mass by orders of magnitude (Biewener and Taylor 1986; Biewener and Patek 2018). Consequently, the move onto land is associated with the evolution of stronger limb bones to avoid fracture (Dickson et al. 2021). However, we currently lack a clear understanding of whether complex life cycles, that expose individuals to water and land, may constrain biomechanical adaptations during evolutionary transitions.

Salamander (Amphibia: Urodela) evolution is characterized by repeated water-land and life cycle transitions, making them an advantageous system to investigate the factors that affect morphological evolution in the limbs (Baken and Adams 2019; Bonett et al. 2022). Many salamanders have a two- or three-part life cycle (multiphasic), that begins with aquatic larvae that subsequently metamorphose into adults with varying degrees of terrestriality (Bonett et al. 2022). However, even terrestrial multiphasic species return to the water for reproduction. Other lineages have monophasic life cycles and either forego metamorphosis (paedomorphism) or undergo metamorphosis in ovum (direct development), which are associated with obligate aquatic or terrestrial habits, respectively (Bonett et al. 2022). Free from contrasting environmental demands during ontogeny, monophasic salamanders may have limb bones more finely tuned to their respective environments and exhibit greater ecomorphological divergence than multiphasic species.

The external morphology of limb bones is often associated with habitat use, but the internal morphology may reflect mechanical adaptations that cannot be gleaned from external morphology alone (Kilbourne and Hutchinson 2019; Keeffe and Blackburn 2022; Rickman et al. 2023; Vera et al. 2020, 2025). Externally, long and gracile bones facilitate speed and maneuverability while short robust bones are associated with strength and increased force production (Polly and Hall 2007). Internally, compact or solid ('dense') limb bones with thickened cortical walls are commonly found in aquatic tetrapods; presumably for buoyancy

regulation (Wall 1983; Canoville and Laurin 2010; Houssaye 2013). While denser limb bones
are more fortified, they are also heavier, which can reduce mobility on land. Alternatively,
terrestrial species might have evolved bones that are less dense and therefore more lightweight,
but with cross-sectional morphologies that still increase stiffness (Currey and Alexander 1985).
The second moment of area, a proxy for 'stiffness' measured from cross-sectional images,
reflects a bone's ability to resist applied forces, such as those experienced during locomotion
(Lieberman et al. 2004; Huie et al. 2022). Stiffer morphologies typically entail hollower bones
and wider cross-sections that distribute material further from the centroid, with the caveat that
they must be thick enough to avoid buckling and fracturing. Contrasting demands between
lightweight versus dense bones likely impose a trade-off between stiffness and density that is
more pronounced in terrestrial lineages compared to aquatic species due to the larger effects of
gravity on land. A question that then arises is how lineages respond to these opposing demands
while also transitioning between conflicting life stages.
A potential strategy for mitigating the consequences of functional trade-offs is to
decouple locomotor structures. For instance, weaker evolutionary integration between external
and internal limb bone morphologies would allow them to evolve independently when exposed
to different selective pressures (Rickman et al. 2023). Similarly, decoupling of the forelimb and
hindlimb bones would enable them to take on more diverse functional roles (Gatesy and Dial
1996; Vera et al. 2025). During most limb-based locomotion in salamanders, the hindlimbs serve
as the primary propulsors and the forelimbs mainly act as brakes (Kawano et al. 2015; Dickson
et al. 2021; Kawano and Blob 2022). The external forces applied to the limbs of the aquatic newt
Pleurodeles waltl during terrestrial locomotion are lower and more divergent between their
forelimbs and hindlimbs compared to the terrestrial salamander *Ambystoma tigrinum* (Kawano
and Blob 2022). Additionally, the demands of terrestrial locomotion are thought to constrain
limb length and maintain stronger covariance between the limbs of terrestrial salamanders
compared to aquatic lineages (Ledbetter and Bonett 2019). Thus, hindlimb bones may be
substantially stiffer than forelimb bones in more aquatic lineages. Yet, terrestrial salamanders
interact with their environment in diverse ways (i.e., climbing, digging, jumping) (Blankers et al.
2012; Baken and Adams 2019) that may promote greater functional decoupling between their
limbs to accommodate different loading regimes. Variation in the developmental timing of
forelimbs and hindlimbs may also influence patterns of limb integration. Many paedomorphs and
multiphasic species develop their limbs disjointly, while direct developers hatch with all four
limbs already developed (Bonett and Ledbetter 2022), suggesting that direct developers may
exhibit stronger levels of covariance between limbs than other life cycle strategies.
We examined the effects of habitat preference, life cycle strategy, and functional trade-
offs on the external and internal morphology of limb bones using micro-computed tomography
(micro-CT) scans of 133 species spanning the phylogenetic and ecological breadth of
salamanders. We used 3D geometric morphometrics to compare external shape, cross-sectional
geometries to compare density and second moment of area ('stiffness'), and phylogenetic
comparative methods to quantify the tempo and mode of evolutionary patterns in the stylopods

(i.e., humerus and femur). We hypothesized that terrestrial salamanders have stiff yet lightweight
limb bones, represented by a functional trade-off (i.e., strong negative correlation) between
stiffness and density. Without the constraints of terrestrial locomotion, aquatic species may
exhibit greater morphological disparity and a functional trade-off that is weaker or absent. We
also hypothesized that multiphasic species would exhibit less morphological divergence than
their monophasic counterparts due to exposure to ontogenetic habitat transitions in the former.
Finally, we hypothesized that the mechanical requirements of terrestrial locomotion promoted
stronger integration between forelimbs and hindlimbs. Our work enhances our understanding of
how biomechanical factors and contrasting ontogenetic demands influence the evolutionary
changes of locomotor structures across water-land transitions.
124
METHODS
Morphological and taxon sampling
We examined the humeri and femora of 133 salamander species, including
representatives from all 10 families and 65 of the 69 (94%) extant genera (AmphibiaWeb 2025).
Each species was represented by a single specimen due to the limited availability of public CT
scans. All specimens were considered adults or large juveniles based on their body size. Most
scans were obtained from MorphoSource.org (Boyer et al. 2016; Blackburn et al. 2024), but 23
additional scans were generated for this study. Scanning was conducted at the Friday Harbor
Labs Karel F. Liem Bio-Imaging Center, using a Bruker Skyscan 1173, or at the University of
Michigan Museum of Zoology (UMMZ) Micro(μ)CT Scanning Laboratory, using a Nikon XT H
225ST. One humerus and one femur per specimen were selected for morphological analysis,
except for sirens (Sirenidae) that have secondarily lost their hindlimbs. To increase data quality,
all sampled limb bones were required to be at least 100 slices long. We used 3D Slicer (version
5.6.1) with the SlicerMorph toolkit (version 23d21fd) to visualize, isolate, and generate models
of each limb bone (Kikinis et al. 2014; Rolfe et al. 2021). Bones selected from the left side of the
body were mirrored so they appeared to originate from the right side.
141
Ancestral reconstruction of habitat preference and life cycle strategy
Salamanders were classified based on their habitat preferences (i.e., aquatic, semi-
aquatic, or terrestrial) and life cycle strategy (i.e., paedomorphic, multiphasic, direct developer).
Most classifications were curated from previous studies or repositories (Fabre et al. 2020;
Louppe et al. 2025; AmphibiaWeb 2025), and modified as needed to more accurately reflect the
natural history of each taxon. We based habitat preferences on where adults spend most of their
time, with the semi-aquatic group being represented by species that live at - and regularly move
across - the interface of aquatic and terrestrial environments. Newts with a multi-year terrestrial
'eft' stage but an aquatic adult stage or long aquatic breeding season were also considered semi-
aquatic. While some multiphasic species exhibit facultative paedomorphism, we primarily
sampled metamorphosed adults. In these cases, habitat and life cycle classifications were coded
based on the characteristics of the sampled individuals (Table S1). Viviparous salamanders were

not included in this study due to small sample sizes. A caveat with our classification approach is
that salamanders are unlikely to match these discrete categories perfectly, especially semi-
aquatic taxa, but they allow us to test general hypotheses surrounding habitat use and life cycle
strategy.
158   We reconstructed the joint evolutionary history of salamander habitat and life cycle
strategy using the most comprehensive time-calibrated salamander phylogeny to date (Stewart
and Wiens 2025). We pruned the tree to include only species in our taxonomic sampling. Species
sampled for morphology but absent from the phylogeny (*Bolitoglossa diminuta*, *Hypselotriton*
wolterstorffi, and *Plethodon grobmani*) were added by replacing the tips of taxa in the same
species complex (*Bolitoglossa aureogularis*, *Hypselotriton yuannensis*, and *Plethodon savannah*,
respectively). We used the CorHMM R package v2.8 (Beaulieu et al. 2022) to reconstruct the
evolution of habitat preference and life cycle strategy with hidden Markov models (Boyko and
Beaulieu 2021). We fit three transition rate models (equal rate, symmetric, and all-rates different)
with and without a hidden state parameter, and all with and without dual transitions for a total of
12 models. We generated 1,000 stochastic character mappings using parameters from the best
fitting model based on weighted AIC values. All analyses were performed in R version 4.0.2 (R
Core Team 2023).
171
172   *External limb bone shape*
173   We used 3D geometric morphometrics to compare the external shapes of the humeri and
femora. Salamander limb bones have few identifiable landmarks; therefore, we used a
pseudolandmark approach. We generated 193 pseudolandmarks on a representative humerus
(*Aneides hardii*) and 190 on a representative femur (*Plethodon elongatus*) using the
PseudoLMGenerator module from the SlicerMorph toolkit in 3D Slicer (Rolfe et al. 2021)
(Figure S1). The pseudolandmarks were transferred from the representative templates to the
remaining humeral and femoral models with the ALPACA module (Porto et al. 2021). Landmark
files were imported into R, where we performed sensitivity analyses to assess whether our results
were robust to the number of landmarks using the "LaSEC" (Landmark Sampling Evaluation
Curve) function in the *LaMBDA* package v0.1.1 (Watanabe 2018). Then we performed
generalized Procrustes Superimpositions on the humerus and femur data sets using the "gpagen"
function in the *geomorph* package v4.0.8 (Baken et al. 2021). To visualize the major axes of
shape variation we conducted principal component analysis (PCA) with a covariance matrix
using the "gm.prcomp" function in *geomorph*.
187
188   *Internal limb bone morphology*
189   We measured two cross-sectional traits that are proxies for bone stiffness and density.
Specifically, we calculated the maximal second moment of area irrespective of any anatomical
axes and bone compactness, respectively, using the *SegmentGeometry* module for 3D Slicer
(Huie et al. 2022). We calculated the cross-sectional traits across the middle 10% of the limb
bones' length, where mechanical loads are predicted to be greatest (Biewener and Taylor 1986),

and calculated the average values for each bone. To isolate the effects of cross-sectional shape on
bone stiffness, we normalized the second moment of area of the bones with the second moment
of area of solid circles with the same cross-sectional area (Huie et al. 2022). These normalized
second moment of area values (referred to as "stiffness") represent how well a bone's shape can
resist bending relative to a solid rod. Bone compactness (referred to as "density") measures the
ratio between the cross-sectional area of the cortical bone and the total cross-sectional area (area
of the cortical bone and vacuities within the section). Total cross-sectional areas were calculated
from solid versions of our limb models generated with the *SurfaceWrapSolidify* module in 3D
Slicer (Weidert et al. 2020).
203
Statistical analyses
We performed phylogenetic ANOVAs to test the effects of body size, habitat preference,
life cycle strategy, and their interactions on limb bone morphology using the "procD.pgls"
geomorph function. Due to the absence of some habitat and life cycle combinations *in situ*, we
tested for an effect of ecotype (combined habitat and life cycle categories) on morphology.
Separate ANOVAs were performed for each bone and trait using a similar formula: trait ~
log(size) + ecotype + log(size) * ecotype. We used snout-vent length (SVL) as a proxy for size
because locomotor forces are expected to scale with body size. For the phylogenetic ANOVAs
involving the cross-sectional traits, we fit an Ornstein–Uhlenbeck (OU) error matrix because we
determined that an OU model of evolution was a better fit for the data than a Brownian Motion
(BM) model with the "fitContinuous" function in the *geiger* package v2.0.11 (Pennell et al.
2014). Where ecotype had a significant effect on shape ($p \leq 0.05$) or there was an interaction
between size and ecotype, we performed pairwise post-hoc tests to compare the mean shape
differences and allometric slopes between ecotypes, respectively. Humeral analyses were
performed with and without sirens but, because the results were similar (Table S3-4), we report
the results that included the sirens.
We also performed phylogenetic paired t-tests to compare differences in mean stiffness
and density between the humerus and femur. Separate analyses were conducted to compare the
humerus and femur of each ecotype using the "phyl.pairedttest" function in the *phytools* R
package v2.4 (Revell 2024).
To characterize the effect of habitat preference and life cycle on morphological diversity,
we estimated the disparity of the bone shapes and cross-sectional traits with the
"morphol.disparity" function in *geomorph*. We performed a separate analysis on each trait and
used ecotype as the discrete variable. For humeral traits, we performed the analyses with and
without the sirens.
229
230 *Correlated evolution of limb bone traits*
231   A functional trade-off between stiffness and density would be represented by a negative
232 correlation between the traits. To compare the evolutionary correlation between stiffness and
233 density across the different ecotypes, we employed the *ratematrix* R package v1.2.4 (Caetano

and Harmon 2017) to implement a Bayesian approach for estimating phylogenetic variance-covariance matrices (Caetano and Harmon 2019). We used the 'Q' transition-rate matrix from the ancestral state reconstruction to generate a distribution of 1,000 stochastic character maps for the evolutionary history of the five ecotypes to account for uncertainty. Then we ran multiple Markov chain Monte Carlo (MCMC) in parallel for five million generations each with 25% burn-in and discarded non-convergent chains until five convergent chains remained. We used the default parameters for all other priors. We ran separate analyses for the humerus (with and without sirens) and femur data sets.

We also investigated the evolutionary integration between forelimb and hindlimb traits across ecotypes. The Bayesian *ratematrix* approach was used to compare the correlations between the humeri and femora regarding stiffness and density. To assess the evolutionary integration of the high-dimensional geometric morphometric datasets, we used the "phylo.integration" function in *geomorph* to perform a two-block partial least-squares analysis to quantify shape covariance between the external shapes of the humerus and femur. Separate analyses were run to compare the limbs within each ecotype. Sirens were omitted from all tests that quantified the integration between the forelimbs and hindlimbs.

Evolution of limb bone shape

We tested whether habitat preference and life cycle strategy influenced the evolution of cross-sectional shapes with an evolutionary model fitting approach. We fit each trait with a set of 26 evolutionary models using the recently described *hOUwie* framework implemented in the *OUwie* R package v2.13 (Boyko and Beaulieu 2021). Parameterized model structures were used to test for both correlated (CD; character dependent) and uncorrelated (CID; character independent) evolution between the cross-sectional traits, habitat preference, and life cycle strategy. We first fit a BM model (BM1) with a single evolutionary rate parameter (σ^2), and a single-rate and single adaptive optimum (θ) OU model (OU1). We also fit more complex models with varied rate parameters (BMV, OUV), trait optima (OUM, OUMV), or rates and trait optima (OUMV) per selective regime. We fit three versions of the BMV, OUV, OUM, and OUMV models with modified model structures to reflect three regime classifications based on habitat preference, life cycle strategy, and the combined ecotypes. Because the evolution of cross-sectional shapes may be affected by factors unaccounted for in our analyses, we also fit CID versions of the BMV, OUV, OUM, and OUMV models each with an additional rate category to account for hidden states. Each model was fitted with 100 stochastic maps per iteration, and with adaptive sampling enabled for the CID models. Model-averaged evolutionary rate and trait optima were estimated for each ecotype using AICc scores to weight the models.

There are currently no reliable methods to model the effects of a discrete character, much less the joint effect of two characters, on high-dimensional data in an OU framework (Adams and Collyer 2019). Therefore, we used a multivariate Brownian Motion framework implemented in *geomorph* to estimate differences in evolutionary rates for external limb bone shape. We used

273   the five ecotype categories as the discrete variable and estimated humeral rates with and without
274   sirens.
275
RESULTS
Evolutionary history of habitat and life cycle
Evolutionary changes in habitat and life cycle were best described by symmetric
transition rates and simultaneous changes (Figure 1). The ancestor of modern salamanders had a
high probability of being either an aquatic paedomorph (47%) or terrestrial multiphasic species
(48%), and a small chance of being semi-aquatic and multiphasic (5%). We recovered multiple
transitions and reversals between aquatic, semi-aquatic, and terrestrial habitats for multiphasic
species. We also estimated up to ten independent transitions to aquatic paedomorphism from a
multiphasic ancestor and nine reversals. Consistent with previous studies, we found a single
transition to direct development and two or three independent reacquisitions of an aquatic larval
stage (Chippindale et al. 2004; Bonett et al. 2014).
287
Disparity in external morphologies
Salamanders displayed a broad range of humeral and femoral shapes with similar axes of
diversification (Figure 2). The primary axes of separate morphospaces for humeri and femora
(principal component 1, PC 1) both depicted the contrast between gracile and robust bones,
consistent with modifications for mobility versus strength, respectively (Polly and Hall 2007;
Young et al. 2014). PC 2 of the humeral and femoral morphospaces captured variation in the size
and presence of the ossified humeral and femoral heads. PC 3 reflected variation in the curvature
of the bone shaft (diaphysis), possibly reflecting differences in the predictability of loading
patterns (Bertram and Biewener 1988).
Terrestrial salamanders displayed lower morphological disparities in humerus and femur
shapes compared to aquatic lineages (Table S2). Aquatic paedomorphs had the most diverse
humeri and femora that were 2.8x (2.3x without sirens) and 1.9x more diverse than those of the
terrestrial direct developers, respectively (Table S2). Multiphasic species exhibited intermediate
disparities that were 0.96x-1.8x more diverse than direct developers. In both morphospaces,
paedomorphs spanned the range of values along PC 1 and PC 3 and uniquely occupied a region
of PC 2 associated with absent or highly reduced ossified humeral heads. Terrestrial direct
developers were generally restricted to regions of the morphospaces associated with more rod-
like and gracile bones. Multiphasic plethodontids also had more rod-like bones, but most
multiphasic species from other families had more robust bones. All multiphasic species had
larger epiphyses than aquatic paedomorphs.
Phylogenetic ANOVAs indicated that body size, ecotype, and their interactions explained
significant variation in humerus and femur shape ($p \leq 0.002$) (Table S3). Pairwise tests supported
findings that paedomorphs had humeri and femora morphologies that were distinct from
multiphasic species ($p \leq 0.015$); no other comparisons were statistically significant (Table S4).
Paedomorphs also exhibited distinct relationships between bone shape and body size compared

to all ecotypes except for aquatic biphasic species (p ≤ 0.017) (Table S4). Larger aquatic
paedomorphs had more robust bones than smaller species, whereas other ecotypes maintained
relatively similar shapes at all body sizes (Figure S2).
316
317   *Disparity in internal morphologies*
318   Terrestrial salamanders had stiffer humeral and femoral cross-sectional shapes, while
aquatic salamanders had denser and more solid bones than terrestrial species (Figure 3). In
contrast to patterns of external shape disparity, terrestrial salamanders exhibited more variation
in their cross-sectional shapes than aquatic species (Table S2). Direct developers had cross-
sectional shapes that were 2.9-8.4x (3.2-12.9x without sirens) more diverse than those of the
paedomorphs (Table S2). Again, the multiphasic species were generally intermediate disparities
that were 0.7-6.0x that of paedomorphs (1.2-9.2x without sirens). Within the direct developers,
temperate genera (e.g., *Aneides*, *Hydromantes*, and *Plethodon*) generally had stiffer bones than
Neotropical lineages (e.g., *Bolitoglossa*, *Chiropterotriton*, *Pseudoeurycea*). Among the semi-
aquatic species, some newts (Salamandridae) and Asiatic salamanders (Hynobiidae) had the
stiffest and least dense cross-sectional shapes of all species examined.
329   Phylogenetic ANOVAs indicated that variation in stiffness and density is associated with
differences between ecotypes (Table S3). Ecotype had a strong effect on most traits (p ≤ 0.003),
except for femoral stiffness (p = 0.053). Pairwise comparisons revealed that terrestrial direct
developers had significantly stiffer humeri than other ecotypes (p ≤ 0.021), except for semi-
aquatic species, which also had stiffer humeri than the aquatic ecotypes (p ≤ 0.035) (Table S4).
Direct developers and semi-aquatic species also had significantly stiffer femora than aquatic
paedomorphs (p ≤ 0.048) (Table S4). Meanwhile, all ecotypes had significantly denser humeri
than direct developers (p ≤ 0.011), but only the aquatic paedomorphs and semi-aquatic species
had significantly denser femora than direct developers (p ≤ 0.010) (Table S4).
338   Humeral stiffness and density were also influenced by the interaction between ecotype
and body size (p ≤ 0.040) (Table S3). Direct developers and semi-aquatic species exhibited
steeper positive relationships between humeral stiffness and body size than aquatic paedomorphs
(p ≤ 0.049), indicating that the former ecotypes have proportionally stiffer bones at larger body
sizes (Figure S2). Direct developers also exhibited a steeper negative relationship between
humeral density and body size than paedomorphs, terrestrial biphasic species, and semi-aquatic
species (p ≤ 0.045) (Figure S2).
345   Despite taking on a larger propulsatory role during limb-based locomotion, the femora
346   were not significantly stiffer than the humeri for any ecotype (p ≥ 0.230) (Table 5). Nevertheless,
347   a few species did exhibit considerably stiffer femora than humeri, such as *Pleurodeles waltl*.
348   There were also no differences in density (p ≥ 0.251), except for aquatic multiphasic species that
349   had denser femora than humeri (t = 3.287, p = 0.009) (Table 5).
350
Functional trade-off between stiffness and density

Stiffness and density exhibited negative evolutionary correlations, providing evidence
for a functional trade-off (Figure 4A, B). Terrestrial direct developers displayed stronger median
correlation values and therefore stronger trade-off between stiffness and density for both limbs
(humerus/femur r = -0.880/-0.929) (Figure 4A, B). Meanwhile, aquatic paedomorphs had
comparatively weaker and/or positive correlations (humerus/femur r = -0.506/0.041; when
excluding siren humeri r = -0.340). However, for the humerus, aquatic (r = -0.233) and terrestrial
(r = - 0.341) multiphasic species had relatively weak correlations but semi-aquatic species
displayed a strong correlation (r = -0.830). For the femur, multiphasic species exhibited
intermediate correlations compared to monophasic species; terrestrial (r = -0.723) and semi-
aquatic (r = -0.629) multiphasic species had stronger correlations than aquatic multiphasic
lineages (r = -0.192).
363

Evolutionary integration between limbs
Habitat preference and life cycle strategy had different effects on the evolutionary
integration of external and internal morphologies between limbs. All ecotypes exhibited
relatively strong signals of integration between the external shapes of their humeri and femora,
based on phylogenetic two-block partial least-squares tests. Paedomorphs had the strongest
integration (r-pls = 0.973, p = 0.006), followed by direct developers (r-pls = 0.929, p = 0.001),
terrestrial (r-pls = 0.945, p = 0.001) and aquatic (r-pls = 0.927, p = 0.021) multiphasic species
and then semi-aquatic species (r-pls = 0.882, p = 0.001). In contrast to external shape, levels of
integration between the cross-sectional traits of the humeri and femora were strongest in direct-
developers (stiffness/density r = 0.635/0.790) and semi-aquatic species (stiffness/density r =
0.654/0.750) (Figure 4C, D). Aquatic multiphasic species (r = 0.155) had stronger integration
between humeral and femoral stiffness than terrestrial multiphasic species (r = -0.029), but the
inverse was true for density (r = 0.608 and 0.563, respectively). Meanwhile, paedomorphs
displayed the weakest correlations (stiffness/density r = -0.253/0.035).
378

Evolutionary rates and models
Estimates of net evolutionary rates showed that the external shapes of the paedomorphic
humeri and femora evolved 5.6x (6.3x without sirens) and 5.3x faster than those of terrestrial
direct developers, respectively (Table S6). The limb bones of all multiphasic salamanders
evolved at intermediate rates, 1.6-3.4x faster than those of the direct developers. We tested 26
models of evolution for all cross-sectional traits, and the best-fitting were generally character
dependent Ornstein–Uhlenbeck models with different adaptive optima for each habitat type
(OUM) or different optima and evolutionary rates (OUMV) (Tables S7–12). Model-averaged
estimates of evolutionary rates indicated that the traits of the direct developers evolved 1.2–8.1x
(1.2–7.9x without sirens) faster than the paedomorphs. Multiphasic species evolved at
comparable rates, 0.6-9.8x faster than paedomorphs (0.8–8.3x without sirens) (Table S13).
Model-averaged trait optima generally resembled the empirically measured values, with more

terrestrial species associated with stiffer bones and aquatic species with denser bones (Table
S13).
393
DISCUSSION
Evolutionary transitions to new environments often promote functional changes to
locomotor structures, but adaptations may be mediated by variation in life cycle strategies. We
leveraged the rich ecomorphological diversity in salamanders to investigate the role of complex
life cycles in shaping the morphological evolution of limb bones across water-land transitions.
We found that aquatic and terrestrial salamanders have evolved contrasting limb bone
morphologies and that multiphasic species display distinct evolutionary patterns and
morphologies compared to monophasic species. Thus, we propose that complex life cycles
increase, rather than constrain, diversity by promoting alternate morphological strategies capable
of addressing the different demands of aquatic and terrestrial habitat use. Furthermore, external
and internal regions of the limb bones have responded differently to habitat and life cycle
transitions, suggesting that evolutionary decoupling provides more viable pathways for
addressing functional and life history trade-offs.
407
Morphological differences between ecotypes
We found evidence that terrestriality promotes stiff yet lightweight bones. Many
terrestrial salamanders have limb bones with ossified epiphyses and stiffer cross-sectional
morphologies that indicate they can resist greater gravitational and locomotor loads compared to
aquatic lineages (Carter et al. 1998; Lieberman et al. 2004; Huie et al. 2022). Conversely, aquatic
salamanders have repeatedly evolved denser limb bones, suggesting that aquatic taxa are
exposed to comparable selective pressures, perhaps functioning as ballasts and increasing
substrate contact during station holding or underwater walking (Canoville and Laurin 2010;
Sanchez et al. 2010; Houssaye 2013; Fabbri et al. 2022). We also uncovered a functional trade-
off between density and stiffness, which is more pronounced among terrestrial species. When
evolving stiffer cross-sectional shapes (e.g. cross-sections with thinner walls or wider diameters),
less dense limb bones also evolve in terrestrial salamanders (Figure 4A, B), presumably as a
means for weight-reduction (Toyama et al. 2023). Furthermore, terrestrial direct developers have
limb bones that are slender and rod-like. While large robust bones are stronger and intuitively
better for dealing with the demands of terrestriality, they are also heavier and therefore more
difficult and energetically expensive to move on land. These findings support that terrestriality
constrains the axes of skeletal diversification in terrestrial tetrapods (Ledbetter and Bonett 2019).
Despite exhibiting narrower axes of variation, terrestrial salamanders exhibit considerable
variation in cross-sectional morphology. Variation in locomotor gaits and behavior result in
different magnitudes and orientations of mechanical loads being applied to the limbs that likely
affect internal morphology. For instance, some scansorial salamanders (*Aneides* and
Hydromantes) had some of the stiffest humeri and femora in our dataset, which could be
adaptations for counteracting higher bone stresses and supporting a larger propulsatory role of

the forelimbs during climbing (Wang et al. 2015; Munteanu et al. 2023). Moreover, salamander femora are not substantially stiffer than the humeri despite the former playing a larger role in propulsion on land, consistent with findings for the terrestrial *Ambystoma tigrinum* (Kawano et al. 2015). It is likely that non-locomotor behaviors also expose the humeri to unique mechanical demands that promote stiffer bones, such as sexually dimorphic behaviors. Males of salamandrids and hynobiids have thicker forelimbs for grabbing females during courtship (amplexus) or fighting other males (Park et al. 1996; Reinhard et al. 2015). Semi-fossorial species (*Ambystoma* spp.) also use their forelimbs for non-locomotor behaviors but did not have exceptionally stiff cross-sectional shapes expected of forelimb burrowers (Semlitsch 1983). However, other factors influence bone stiffness besides morphology, including material properties (Erickson et al. 2002; Lieberman et al. 2004). As a result, material properties may complement or mediate structural changes in response to different environmental demands.

Additionally, differences in body size also may help explain the variation in cross-sectional morphology. Because heavier animals experience disproportionately larger forces on their bones than lighter animals, the selective advantages of stiff yet lightweight bones would be greater in larger species (Currey and Alexander 1985). We found some evidence that larger animals have proportionally stiffer bones, indicated by the steeper slope between stiffness and body size among terrestrial salamanders (Figure S2). Conversely, most salamanders are relatively small animals that produce correspondingly small locomotor forces (Kawano et al. 2015; Kawano and Blob 2022). Thus, fortifying limb bones with cortical bone and increasing density may have minimal effects on mobility, especially direct developers that already have fairly gracile limb bones. That might explain why many miniature species examined herein (e.g., *Thorius macdougalli*, *Desmognathus wrighti*, *Nototriton* spp., *Parvimolge townsendi*) have relatively solid cross-sections (density values > 0.95) (Uzzell 1961; Hanken 1982). Although, *Thorius* exhibits hyper-ossification of their limb bones that coincides with sexual maturation rather than a particular body size (Hanken 1982), suggesting that not all changes in bone shape reflect mechanical adaptations for terrestrial locomotion.

Complex life cycles promote diversity

Life cycle complexities do not influence all aspects of the salamander skeleton equally. Our results on the external shape of limb bones are consistent with previous findings that the crania and feeding apparatus of multiphasic salamanders are less morphologically diverse than those of paedomorphs, but more diverse than those of direct developers (Fabre et al. 2020; Louppe et al. 2025). Those authors proposed that complete metamorphosis, and not just the presence of a complex life cycle, constrains cranial diversity in multiphasic and direct developing species. This appears to be plausible for external limb bone shape but not for cross-sectional shape, where direct developers and multiphasic exhibit greater disparity and higher rates of morphological evolution than paedomorphs. Different from skeletal evolution, multiphasic salamanders have more constrained body shapes than monophasic species attributed to the constraints associated with conflicting environmental demands throughout ontogeny

(Bonett and Blair 2017). Our findings provide additional support that complex life cycles may
limit morphological diversification in multiphasic species, but the magnitude varies across traits.
Aquatic paedomorphs and terrestrial direct developers exhibit greater ecomorphological
divergence between their limb bone shapes than multiphasic species in those same environments.
This likely reflects the narrower range of selective pressures associated with consistent habitat
use over ontogeny, whereas multiphasic species are exposed to contrasting physical demands
associated with living in water and on land across life stages (Wilson 2005; Bonett and Blair
2017). It also suggests that limb development is not entirely decoupled by metamorphosis,
supported by observations of limited anatomical changes between life stages (Ashley-Ross
1992). Therefore, multiphasic species have evolved limb bones influenced by both aquatic and
terrestrial demands. Interspecific variation in the timing and duration of life history
characteristics that affect time spent in water or on land (i.e., the aquatic larval stage, terrestrial
eft stage, or aquatic breeding season) likely influence the relative impacts of the environment and
primary agents of selection acting on bone morphology (Weaver et al. 2020; Bonett and
Ledbetter 2022; Bonett et al. 2022). For example, many newts have exceptionally stiff bones,
possibly reflecting their terrestrial juvenile phases that can last seven years and are absent in
other semi-aquatic taxa (Forester and Lykens 1991).
The relative differences in shape, allometry, and evolutionary rates between multiphasic
species living in different environments do not always parallel the patterns observed among
monophasic species. Many multiphasic species have robust external limb bone shapes not
observed in monophasic species and cross-sectional traits that rival direct developers with the
most extreme stiffness and density values. These results indicate that multiphasic species are not
intermediate versions of monophasic species but instead have their own distinct evolutionary
trajectories. Similarly, semi-aquatic lineages are not simply intermediates between aquatic and
terrestrial species. These findings suggest that the repeated losses of multiphasic life cycles are
unlikely to represent mechanisms for escaping the constraints imposed by complex life cycles.
Instead, despite potential life cycle constraints, multiphasic species have highly evolvable limb
bones that may have played a role in the reacquisition of an aquatic larval stage and semi-aquatic
habitat preferences (Chippindale et al. 2004; Bonett et al. 2014).
500
Decoupling of locomotor structures
Weak evolutionary covariance between the external and internal limb bone morphologies
are reflected in the independent correlations with habitat and life cycles. We posit that their
decoupling has expanded the locomotor capabilities of direct developers and plethodontid
salamanders more broadly and, in part, enabled their ecological diversification. Specifically, we
propose that the long and gracile plethodontid limb bones increase mobility and locomotor speed
(Young et al. 2014), while the internal morphologies can reflect diverse mechanical demands.
That is evidenced by the fact that plethodontids occupy diverse microhabitats (i.e., burrows,
trees, rocks, caves, streams) (Blankers et al. 2012; Baken and Adams 2019) and employ
terrestrial locomotor modes rarely observed outside of plethodontids (e.g., jumping and

climbing) (Brown and Deban 2020; Aretz et al. 2022; Huie et al. 2025). In contrast, aquatic
paedomorphs have conserved cross-sectional shapes but diverse external shapes, perhaps as the
result of relaxed selection on walking kinematics, body shape, and body size in aquatic
environments (Ashley-Ross and Bechtel 2004; Azizi and Horton 2004; Bonett and Blair 2017).
The only shared external feature of paedomorph limb bones is the reduction or absence of
ossified epiphyses, which are formed by cartilage in aquatic salamanders (Molnar 2021b).
Ecological transitions have also influenced the evolutionary correlation between limbs.
All ecotypes exhibit relatively strong integration between the external and internal shapes of
their limbs (r-pls = 0.88-0.98), possibly reflecting their relatively conserved form of limb-based
locomotion (Pierce et al. 2020). Surprisingly, paedomorphs have the strongest integration of
external shapes but the weakest covariance between their cross-sectional traits. The latter is
aligned with the reduced integration between forelimb and hindlimb lengths among aquatic
salamanders and the loss of hindlimbs in aquatic sirens (Ledbetter and Bonett 2019), perhaps
reflecting more decoupled mechanical demands. In contrast, direct developers exhibit strong
patterns of integration between all humeral and femoral traits, which may reflect the shared load-
bearing demands imposed onto the forelimbs and hindlimbs associated with living on land
(Kawano et al. 2015; Kawano and Blob 2022).
Multiphasic species also exhibit considerable variation in the evolutionary integration of
their forelimbs and hindlimbs. Semi-aquatic species exhibit some of the weakest integration
between their external shapes but strongest integration between cross-sectional traits. Because
terrestrial multiphasic species do not exhibit these patterns, they are likely dissociated from the
demands of terrestrial locomotion and instead reflect unexamined factors. For example, variation
in the developmental timing of the forelimbs and hindlimbs may influence patterns of integration
in adults. Pond-dwelling larvae exhibit a pronounced lag time between forelimb and hindlimb
development, while steam-dwelling larvae (and direct developers) experience less of a lag likely
due to an increased need for station holding and underwater walking in streams compared to
ponds (Shubin and Wake 2003). Investigations into whether larval development influences levels
of integration and if that varies across life stages could yield insights into the factors that
generate limb diversity.
540
Future directions
In this study we found that interactions between habitat and complex life cycle shape the
evolutionary morphology of salamander limb bones. Functional trade-offs between stiffness and
density have seemingly promoted stiff yet lightweight bones on land and restricted how these
traits covary. Yet, because terrestrial salamanders are relatively small animals, additional work is
needed to empirically test whether the forces acting on the limb bones of terrestrial species place
demands on the musculoskeletal system strong enough to constrain morphological evolution in
the limbs. Walking alone is unlikely to cause bone breakage due to the high safety factors
('margins against failure') found in salamanders (Kawano et al. 2013), but other terrestrial

activities like running, jumping, or combat may promote the stiffer shapes that we observed (Biewener and Patek 2018).

Additionally, we suggest that complex life cycles promote distinct limb bone morphologies to manage the contrasting aquatic and terrestrial demands experienced within a single lifetime. Future work should examine limb bones over ontogeny to assess the magnitude of change that occurs in morphology, integration, and mechanical properties across metamorphosis and habitat transitions. Doing so will directly address whether metamorphosis can decouple limb bone development between life stages and if complex life cycles necessitate alternative morphologies.

Finally, examining specific clades in greater detail will reveal more nuanced factors affecting limb evolution. For instance, it is unclear to what extent sexual dimorphism and courtship behaviors in salamandrids influence the properties of their limb bones (Reinhard et al. 2015). We also recognize that the semi-aquatic ecotype as defined in this study may be overly broad because it includes species that spend most of their lives at the water's edge as well as newts with three life stages and multiple transitions between aquatic and terrestrial stages. We suspect that the different semi-aquatic strategies have distinct consequences on limb bone evolution, and encourage future studies to test this hypothesis at a finer scale than what was focused on in this study. In sum, further investigation into how the biomechanical properties of salamander limb bones relate to life cycle strategy, locomotion, and other functional applications would provide a deeper understanding of the selective pressures governing musculoskeletal evolution.

ACKNOWLEDGEMENTS

We thank P. Millers for providing access to Burke Museum specimens amidst the COVID-19 pandemic, D. Grossnickle for retrieving the specimens, and A. Summers and the Karel F. Liem Bio-Imaging facility for free access to a micro-CT scanner. We also thank H. Martens, R. Nagesan, and G. Schneider for providing CT scans of UMMZ specimens. We thank D. Caetano for helping with the *ratematrix* analyses. We also thank two anonymous reviewers for their helpful feedback on earlier drafts of this manuscript.

FUNDING

This work was supported by a US National Science Foundation Graduate Research Fellowship [DGE-1746914], a Washington Biologists' Field Club Award, and Wilbur V. Harlan Research Fellowship from The George Washington University to JMH; US National Science Foundation grants DBI-0905765, DEB-1441719, and DEB-1655737 to RAP; and lab start-up and University Facilitating Funds from The George Washington University and a Research Publication Grant from the American Association of University Women [award number 015943] to SMK.

AUTHOR CONTRIBUTIONS

J.M.H. and S.M.K. designed research; J.M.H. performed research; J.M.H. and R.A.P. analyzed
data; and J.M.H., R.A.P., and S.M.K. wrote the paper.

DATA AVAILABILITY
The morphological data and R script files used for this study can be found on GitHub
(https://github.com/jmhuie/Salamander_Limb_Bone_Evo).

REFERENCES
Adams, D. C., and M. L. Collyer. 2019. Phylogenetic Comparative Methods and the Evolution of
        Multivariate Phenotypes. Annu. Rev. Ecol. Evol. Syst. 50:405–425.
AmphibiaWeb. 2025. <https://amphibiaweb.org> University of California, Berkeley, CA, USA.
        Accessed 10 Oct 2025.
Aretz, J. M., C. E. Brown, and S. M. Deban. 2022. Vertical locomotion in the arboreal
        salamander Aneides vagrans. J. Zool. 316:72–79.
Ashley-Ross, M. A. 1992. The comparative myology of the thigh and crus in the salamanders
        Ambystoma tigrinum and Dicamptodon tenebrosus. J. Morphol. 211:147–163.
Ashley-Ross, M. A., and B. F. Bechtel. 2004. Kinematics of the transition between aquatic and
        terrestrial locomotion in the newt Taricha torosa. J. Exp. Biol. 207:461–474.
Azizi, E., and J. M. Horton. 2004. Patterns of axial and appendicular movements during aquatic
        walking in the salamander *Siren lacertina*. Zoology 107:111–120.
Baken, E. K., and D. C. Adams. 2019. Macroevolution of arboreality in salamanders. Ecol. Evol.
        9:7005–7016.
Baken, E. K., M. L. Collyer, A. Kaliontzopoulou, and D. C. Adams. 2021. geomorph v4.0 and
        gmShiny: Enhanced analytics and a new graphical interface for a comprehensive
        morphometric experience. Methods Ecol. Evol. 12:2355–2363.
Beaulieu, J., B. O'Meara, J. Oliver, and J. Boyko. 2022. corHMM: Hidden Markov Models of
        Character Evolution.
Bejder, L., and B. K. Hall. 2002. Limbs in whales and limblessness in other vertebrates:
        mechanisms of evolutionary and developmental transformation and loss. Evol. Dev.
        4:445–458.
Bertram, J. E. A., and A. A. Biewener. 1988. Bone curvature: Sacrificing strength for load
        predictability? J. Theor. Biol. 131:75–92.
Biewener, A. A., and C. R. Taylor. 1986. Bone Strain: A Determinant of Gait and Speed? J. Exp.
        Biol. 123:383–400.
Biewener, A., and S. Patek. 2018. Animal Locomotion. Oxford University Press.
Blackburn, D. C., D. M. Boyer, J. A. Gray, J. Winchester, J. M. Bates, S. L. Baumgart, E.
        Braker, D. Coldren, K. W. Conway, A. D. Rabosky, N. de la Sancha, C. B. Dillman, J. L.
        Dunnum, C. M. Early, B. W. Frable, M. W. Gage, J. Hanken, J. A. Maisano, B. D.
        Marks, K. P. Maslenikov, J. E. McCormack, R. S. Nagesan, G. G. Pandelis, H. L.
        Prestridge, D. L. Rabosky, Z. S. Randall, M. B. Robbins, L. A. Scheinberg, C. L.

Spencer, A. P. Summers, L. Tapanila, C. W. Thompson, L. Tornabene, G. J. Watkins-Colwell, L. J. Welton, the oVert Project Team, and E. L. Stanley. 2024. Increasing the impact of vertebrate scientific collections through 3D imaging: The openVertebrate (oVert) Thematic Collections Network. BioScience 74:169–186.

Blankers, T., D. C. Adams, and J. J. Wiens. 2012. Ecological radiation with limited morphological diversification in salamanders. J. Evol. Biol. 25:634–646.

Bonett, R. M., and A. L. Blair. 2017. Evidence for complex life cycle constraints on salamander body form diversification. Proc. Natl. Acad. Sci. 114:9936–9941. Proceedings of the National Academy of Sciences.

Bonett, R. M., and N. M. Ledbetter. 2022. Paedomorphic salamanders are larval in form and patterns of limb emergence inform life cycle evolution. Dev. Dyn. 251:934–941.

Bonett, R. M., N. M. Ledbetter, A. J. Hess, M. A. Herrboldt, and M. Denoël. 2022. Repeated ecological and life cycle transitions make salamanders an ideal model for evolution and development. Dev. Dyn. 251:957–972.

Bonett, R. M., M. A. Steffen, S. M. Lambert, J. J. Wiens, and P. T. Chippindale. 2014. Evolution of paedomorphosis in plethodontid salamanders: ecological correlates and re-evolution of metamorphosis. Evolution 68:466–482.

Boyer, D. M., G. F. Gunnell, S. Kaufman, and T. M. McGeary. 2016. MORPHOSOURCE: ARCHIVING AND SHARING 3-D DIGITAL SPECIMEN DATA. Paleontol. Soc. Pap. 22:157–181. Cambridge University Press.

Boyko, J. D., and J. M. Beaulieu. 2021. Generalized hidden Markov models for phylogenetic comparative datasets. Methods Ecol. Evol. 12:468–478.

Brown, C. E., and S. M. Deban. 2020. Jumping in arboreal salamanders: A possible tradeoff between takeoff velocity and in-air posture. Zoology 138:125724.

Caetano, D. S., and L. J. Harmon. 2019. Estimating Correlated Rates of Trait Evolution with Uncertainty. Syst. Biol. 68:412–429.

Caetano, D. S., and L. J. Harmon. 2017. ratematrix: An R package for studying evolutionary integration among several traits on phylogenetic trees. Methods Ecol. Evol. 8:1920–1927.

Canoville, A., and M. Laurin. 2010. Evolution of humeral microanatomy and lifestyle in amniotes, and some comments on palaeobiological inferences. Biol. J. Linn. Soc. 100:384–406.

Canoville, A., and M. Laurin. 2009. Microanatomical diversity of the humerus and lifestyle in lissamphibians. Acta Zool. 90:110–122.

Carter, D. R., B. Mikić, and K. Padian. 1998. Epigenetic mechanical factors in the evolution of long bone epiphyses. Zool. J. Linn. Soc. 123:163–178.

Chippindale, P. T., R. M. Bonett, A. S. Baldwin, and J. J. Wiens. 2004. Phylogenetic evidence for a major reversal of life-history evolution in plethodontid salamanders. Evolution 58:2809–2822.

Currey, J. D., and R. M. Alexander. 1985. The thickness of the walls of tubular bones. J. Zool. 206:453–468. Wiley Online Library.

Dickson, B. V., J. A. Clack, T. R. Smithson, and S. E. Pierce. 2021. Functional adaptive landscapes predict terrestrial capacity at the origin of limbs. Nature 589:242–245. Nature Publishing Group.

Easterling, C. M., M. A. Kolmann, and M. K. O'Donnell. 2022. The Lesser-Known Transitions: Organismal Form and Function across Abiotic Gradients. Integr. Comp. Biol. 62:829–839.

Ebenman, B. 1992. Evolution in Organisms that Change Their Niches during the Life Cycle. Am. Nat. 139:990–1021. [The University of Chicago Press, The American Society of Naturalists].

Erickson, G. M., J. Catanese, and T. M. Keaveny. 2002. Evolution of the biomechanical material properties of the femur. Anat. Rec. 268:115–124.

Fabbri, M., G. Navalón, R. B. J. Benson, D. Pol, J. O'Connor, B.-A. S. Bhullar, G. M. Erickson, M. A. Norell, A. Orkney, M. C. Lamanna, S. Zouhri, J. Becker, A. Emke, C. Dal Sasso, G. Bindellini, S. Maganuco, M. Auditore, and N. Ibrahim. 2022. Subaqueous foraging among carnivorous dinosaurs. Nature 603:852–857. Nature Publishing Group.

Fabre, A.-C., C. Bardua, M. Bon, J. Clavel, R. N. Felice, J. W. Streicher, J. Bonnel, E. L. Stanley, D. C. Blackburn, and A. Goswami. 2020. Metamorphosis shapes cranial diversity and rate of evolution in salamanders. Nat. Ecol. Evol. 4:1129–1140. Nature Publishing Group.

Forester, D. C., and D. V. Lykens. 1991. Age Structure in a Population of Red-Spotted Newts from the Allegheny Plateau of Maryland. J. Herpetol. 25:373–376. Society for the Study of Amphibians and Reptiles.

Gamel, K. M., S. Pinti, and H. C. Astley. 2024. Ground Reaction Forces And Energy Exchange During Underwater Walking. Integr. Org. Biol. obae013.

Gatesy, S. M., and K. P. Dial. 1996. Locomotor Modules and the Evolution of Avian Flight. Evolution 50:331–340.

Hanken, J. 1982. Appendicular skeletal morphology in minute salamanders, genus *Thorius* (amphibia: Plethodontidae): Growth regulation, adult size determination, and natural variation. J. Morphol. 174:57–77.

Hanken, J. 1992. Life history and morphological evolution. J. Evol. Biol. 5:549–557.

Houssaye, A. 2013. Bone histology of aquatic reptiles: what does it tell us about secondary adaptation to an aquatic life? Biol. J. Linn. Soc. 108:3–21.

Huie, J. M., A. P. Summers, and S. M. Kawano. 2022. SegmentGeometry: A Tool for Measuring Second Moment of Area in 3D Slicer. Integr. Org. Biol. 4:obac009.

Huie, J. M., G. Park, and S. M. Kawano. 2025. Salamander ecomorphology reveals a unique suite of climbing adaptations. Proc. R. Soc. B. 292:20251295.

Kawano, S. M., and R. W. Blob. 2022. Terrestrial force production by the limbs of a semi-aquatic salamander provides insight into the evolution of terrestrial locomotor mechanics. J. Exp. Biol. 225:jeb242795.

Kawano, S. M., D. R. Economy, M. S. Kennedy, D. Dean, and R. W. Blob. 2015. Comparative limb bone loading in the humerus and femur of the tiger salamander *Ambystoma tigrinum* : testing the 'mixed-chain' hypothesis for skeletal safety factors. J. Exp. Biol. jeb.125799.

Keeffe, R., and D. C. Blackburn. 2022. Diversity and function of the fused anuran radioulna. J. Anat. 241:1026–1038.

Kikinis, R., S. D. Pieper, and K. G. Vosburgh. 2014. 3D Slicer: A Platform for Subject-Specific Image Analysis, Visualization, and Clinical Support. Pp. 277–289 *in* F. A. Jolesz, ed. Intraoperative Imaging and Image-Guided Therapy. Springer, New York, NY.

Kilbourne, B. M., and J. R. Hutchinson. 2019. Morphological diversification of biomechanical traits: mustelid locomotor specializations and the macroevolution of long bone cross-sectional morphology. BMC Evol. Biol. 19:37.

Koehl, M. A. R. 2023. Of corpses, ghosts and mirages: biomechanical consequences of morphology depend on the environment. J. Exp. Biol. 226:jeb245442.

Laudet, V. 2011. The Origins and Evolution of Vertebrate Metamorphosis. Curr. Biol. 21:R726–R737.

Laurin, M., M. Girondot, and M.-M. Loth. 2004. The evolution of long bone microstructure and lifestyle in lissamphibians. Paleobiology 30:589–613.

Ledbetter, N. M., and R. M. Bonett. 2019. Terrestriality constrains salamander limb diversification: Implications for the evolution of pentadactyly. J. Evol. Biol. 32:642–652.

Lieberman, D. E., J. D. Polk, and B. Demes. 2004. Predicting long bone loading from cross-sectional geometry. Am. J. Phys. Anthropol. 123:156–171.

Louppe, V., K. Le Verger, J. Clavel, G. S. Ferreira, C. Kyriakouli, E. L. Stanley, E. C. Watt, C. Bardua, C. Pérez-Ben, N. B. Fröbisch, V. Bothe, R. Boistel, G. Daghfous, A. Goswami, and A.-C. Fabre. 2025. Complete metamorphosis promotes morphological and functional diversity in Caudata. Funct. Ecol. 39:1732–1747.

McMenamin, S. K., and D. M. Parichy. 2013. Chapter Five - Metamorphosis in Teleosts. Pp. 127–165 *in* Y.-B. Shi, ed. Current Topics in Developmental Biology. Academic Press.

Molnar, J. L., J. R. Hutchinson, R. Diogo, J. A. Clack, and S. E. Pierce. 2021a. Evolution of forelimb musculoskeletal function across the fish-to-tetrapod transition. Sci. Adv. 7:eabd7457. American Association for the Advancement of Science.

Molnar, J. L. 2021b. Variation in Articular Cartilage Thickness Among Extant Salamanders and Implications for Limb Function in Stem Tetrapods. Front. Ecol. Evol. 9.

Moran, N. A. 1994. Adaptation and Constraint in the Complex Life Cycles of Animals. Annu. Rev. Ecol. Syst. 25:573–600. Annual Reviews.

Munteanu, V. D., K. M. Diamond, and R. W. Blob. 2023. Limb bone strains during climbing in green iguanas (Iguana iguana): testing biomechanical release as a mechanism promoting morphological transitions in arboreal vertebrates. J. Exp. Biol. 226:jeb245175.

Park, S.-R., D.-S. Park, and S. Y. Yang. 1996. Courtship, fighting behaviors and sexual
dimorphism of the salamander, Hynobius leechii. Korean J. Zool. 39:437–446.
ZOOLOGICAL SOCIETY OF KOREA.
Pennell, M. W., J. M. Eastman, G. J. Slater, J. W. Brown, J. C. Uyeda, R. G. FitzJohn, M. E.
Alfaro, and L. J. Harmon. 2014. geiger v2.0: an expanded suite of methods for fitting
macroevolutionary models to phylogenetic trees. Bioinformatics 30:2216–2218.
Pierce, S. E., L. P. Lamas, L. Pelligand, N. Schilling, and J. R. Hutchinson. 2020. Patterns of
Limb and Epaxial Muscle Activity During Walking in the Fire Salamander, *Salamandra*
salamandra. Integr. Org. Biol. 2:obaa015.
Polly, P. D., and B. Hall. 2007. Limbs in mammalian evolution. Fins Limbs Evol. Dev.
Transform. 15:245–268. University of Chicago Press Chicago.
Porto, A., S. Rolfe, and A. M. Maga. 2021. ALPACA: A fast and accurate computer vision
approach for automated landmarking of three-dimensional biological structures. Methods
Ecol. Evol. 12:2129–2144.
R Core Team. 2023. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria.
Reinhard, S., S. Renner, and A. Kupfer. 2015. Sexual dimorphism and age of Mediterranean
salamanders. Zoology 118:19–26.
Revell, L. J. 2024. phytools 2.0: an updated R ecosystem for phylogenetic comparative methods
(and other things). PeerJ 12:e16505. PeerJ Inc.
Rickman, J., A. E. Burtner, T. J. Linden, S. E. Santana, and C. J. Law. 2023. Size And
Locomotor Ecology Have Differing Effects on the External and Internal Morphologies of
Squirrel (Rodentia: Sciuridae) Limb Bones. Integr. Org. Biol. 5:obad017.
Rolfe, S., S. Pieper, A. Porto, K. Diamond, J. Winchester, S. Shan, H. Kirveslahti, D. Boyer, A.
Summers, and A. M. Maga. 2021. SlicerMorph: An open and extensible platform to
retrieve, visualize and analyse 3D morphology. Methods Ecol. Evol. 12:1816–1825.
Semlitsch, R. D. 1983. Burrowing ability and behavior of salamanders of the genus *Ambystoma*.
Can. J. Zool. 61:616–620.
Shubin, N. H., and D. B. Wake. 2003. Morphological variation, development, and evolution of
the limb skeleton of salamanders. Amphib. Biol. 5:1782–1808. Surrey Beatty and Sons
Chipping Norton, Australia.
Stewart, A. A., and J. J. Wiens. 2025. A time-calibrated salamander phylogeny including 765
species and 503 genes. Mol. Phylogenet. Evol. 204:108272.
Toyama, K. S., A. Tinius, and D. L. Mahler. 2023. Evidence supporting an evolutionary trade-off
between material properties and architectural design in Anolis lizard long bones.
Evolution qpad208.
Truman, J. W. 2019. The Evolution of Insect Metamorphosis. Curr. Biol. 29:R1252–R1268.
Elsevier.

Uzzell, T. M. 1961. Calcified Hyoid and Mesopodial Elements of Plethodontid Salamanders.
Copeia 1961:78–86. [American Society of Ichthyologists and Herpetologists (ASIH),
Allen Press].
Vera, M. C., J. L. Ferretti, V. Abdala, and G. R. Cointry. 2020. Biomechanical properties of
anuran long bones: correlations with locomotor modes and habitat use. Journal of
Anatomy 236:1112–1125.
Vera, M. C., J. Fratani, G. R. Cointry, and V. Abdala. n.d. Biomechanical Properties of Long
Bones and Degrees of Morphological Integration Between the Fore and Hindlimbs in
Anuran Species With Different Habitat Uses. Journal of Experimental Zoology Part B:
Molecular and Developmental Evolution n/a.
Wall, W. P. 1983. The Correlation between High Limb-Bone Density and Aquatic Habits in
Recent Mammals. J. Paleontol. 57:197–207. Paleontological Society.
Wang, Z., Z. Dai, W. Li, A. Ji, and W. Wang. 2015. How do the substrate reaction forces acting
on a gecko's limbs respond to inclines? Sci. Nat. 102:7.
Watanabe, A. 2018. How many landmarks are enough to characterize shape and size variation?
PLOS ONE 13:e0198341.
Weaver, S., D. B. Shepard, and K. H. Kozak. 2020. Developmental life history is associated with
variation in rates of climatic niche evolution in a salamander adaptive radiation*.
Evolution 74:1804–1814.
Weidert, S., S. Andress, C. Linhart, E. M. Suero, A. Greiner, W. Böcker, C. Kammerlander, and
C. A. Becker. 2020. 3D printing method for next-day acetabular fracture surgery using a
surface filtering pipeline: feasibility and 1-year clinical results. Int. J. Comput. Assist.
Radiol. Surg. 15:565–575.
Wilson, R. S. 2005. Consequences of Metamorphosis for the Locomotor Performance and
Thermal Physiology of the Newt *Triturus cristatus*. Physiol. Biochem. Zool. 78:967–975.
Young, J. W., R. Danczak, G. A. Russo, and C. D. Fellmann. 2014. Limb bone morphology,
bone strength, and cursoriality in lagomorphs. J. Anat. 225:403–418.
Young, V. K. H., and R. W. Blob. 2015. Limb bone loading in swimming turtles: changes in
loading facilitate transitions from tubular to flipper-shaped limbs during aquatic
invasions. Biol. Lett. 11:20150110.

816
817 **Figure 1.** Phylogeny of 133 salamander species sampled for morphological analysis, showing
818 the joint ancestral state reconstruction of habitat and life cycle. Insets depict representative
819 humeri and femora and their cross-sectional shapes at the midshaft (not to scale).

Humerus

Femur

Aquatic/Paedomorphic   Aquatic/Multiphasic   Semi-aquatic/Multiphasic   Terrestrial/Multiphasic   Terrestrial/Direct Developer

Figure 2. Phylomorphospaces illustrating the shape diversity of salamander humeri (left) and femora (right). Separate principal component analyses were used to identify the primary axes of variation for each limb bone. Lateral and dorsal perspectives of bones from representative taxa depict the extremes of each phylomorphospace: a) *Nyctanolis pernix*, b) *Andrias japonicus*, c) *Siren intermedia*, d) *Salamandrina terdigitata*, e) *Pseudobranchus striatus*, f) *Bolitoglossa porrasorum*, g) *Chiropterotriton magnipes*, h) *Ambystoma maculatum*, i) *Amphiuma tridactylum*, j) *Bradytriton silus*, k) *Hypselotriton wolterstorffi*, l) *Ensatina eschscholtzii*.

Figure 3. Variation in the stiffness and density of salamander humeri (left) and femora (right)
across habitats and life cycle strategies. Boxplots show the median, upper and lower quartiles,
interquartile range and outliers as determined by the 1.5 interquartile rule. Illustrations along the
y-axes depict graphical representations of the changes in cross-sectional morphology. Some
outliers are labeled: a) *Paradactylodon persicus*, b) *Aneides lugubris*, c) *Pleurodeles waltl*, d)
Tylototriton tagaliensis, e) *Ambystoma mabeei*, f) *Hydromantes platycephalus*, g)
Onychodactylus japonicus, and g) *Amphiuma tridactylum*.

Figure 4. Evolutionary relationships between cross-sectional traits within and across limbs. Scatter plot points were slightly jittered to enhance readability. Insets depict density plots of the correlation (r) values estimated by Bayesian *ratematrix* analyses. Humerus plot (A) depicts the results from the analysis that included sirens. The strong negative correlations between stiffness and density within bones indicate a functional trade-off between traits (A, B). Meanwhile, strong positive correlations between the humeral and femoral traits indicate integration between the limbs (C, D).