1

Combinatorial Treatment of Glioblastoma with Temozolomide (TMZ) Plus 5-Ethynyl-2'-deoxyuridine (EdU)

10 Humeyra Kaanoglu[1,*], Yasemin K. Akyel[1,*], Adebimpe Adefolaju[2], Alain Valdivia[2], Dominique
11 Higgins[3], Caroline Mosely[4], Corey J. Haswell[4], William C. Zamboni[4], Shawn D. Hingtgen[2], Andrew
12 B. Satterlee[2, 5], and Aziz Sancar[1, 6]

13
14

15 [1]Department of Biochemistry and Biophysics, University of North Carolina School of Medicine,
16 Chapel Hill, North Carolina
17
18 [2]Division of Pharmacoengineering and Molecular Pharmaceutics, Eshelman School of Pharmacy
19 University of North Carolina, Chapel Hill, North Carolina
20
21 [3]Department of Neurosurgery, University of North Carolina, Chapel Hill, North Carolina
22
23 [4]UNC Advanced Translational Pharmacology and Analytical Chemistry Lab, UNC Eshelman
24 School of Pharmacy, University of North Carolina at Chapel Hill, North Carolina
25
26 [5]Eshelman Innovation, University of North Carolina, Chapel Hill, North Carolina
27
[6]**Correspondence:** aziz_sancar@med.unc.edu

*Equal contribution

Conflict of Interest Disclosure Statement: Aziz Sancar, Humeyra Kaanoglu and Yasemin Akyel
filed a patent application with title "METHODS OF TREATING CANCER OF CENTRAL
NERVOUS SYSTEM COMPRISING 5-ETHYNYL-2'-DEOXYURIDINE AND TEMOZOLOMIDE"
and U.S. Provisional Application No. 6/893,131 on October 3, 2025.

ABSTRACT

Glioblastoma (GBM) is the most aggressive malignant primary brain tumor in adults, with incidence peaking in later life. It is commonly treated with surgery followed by administration of ionizing radiation and the DNA alkylating agent temozolomide (TMZ). Even though this regimen confers some progression-free survival, there is essentially no cure with the median survival with the standard of care being about 12 months. Currently, several alternative approaches are being developed to improve upon this outcome. We have already shown EdU alone effectively treats GBM, and we now study efficacy of TMZ+EdU combination therapy. TMZ+EdU significantly improves antitumor efficacy compared to either single-agent therapy against GBM cell lines *in vitro*, against three different orthotopic GBM xenograft models, and against passage-zero patient GBM tumor tissues engrafted within an organotypic brain slice culture (OBSC)-based platform. Together, these data suggest that EdU could be effective alongside standard of care TMZ in patients with GBM.

STATEMENT OF SIGNIFICANCE (50-WORD)

By simultaneously engaging distinct DNA repair pathways, combination of TMZ and EdU produced unprecedented survival benefits and synergistic effects in preclinical GBM models, offering a promising new avenue in the treatment of GBM.

INTRODUCTION

Despite the application of multidisciplinary treatment approaches combining surgical, radiotherapeutic, and pharmacologic interventions, glioblastoma (GBM) remains one of the most treatment-resistant human cancers, with progression and recurrence as defining features (1–4). At the molecular level, GBM arises through a constellation of genetic alterations, including loss of heterozygosity on chromosome 10q, EGFR amplification, p16INK4a deletion, PTEN mutation, and, at later stages, TP53 and IDH1 mutations (5,6). Currently, the standard of care for GBM consists of surgery followed by ionizing radiation and temozolomide (TMZ) drug therapy (7–9)., During TMZ treatment, eventually MGMT (methylguanine methyl transferase) promoter hypermethylation causes silencing of MGMT which plays an essential role in TMZ-initiated death of malignant tumor cells. However, after an initial positive response to multimodal therapy, ~80% of GBM patients experience recurrence at or near the original tumor site (10–13), often accompanied by new genetic alterations acquired under treatment pressure (14,15). At this stage, therapeutic options are limited, and even with standard of care, median survival remains around 12 months (4,13,16). Recently, a number of new drugs/intervention methods have been developed to improve the management of GBM (17). Among these, 5-ethynyl-2-deoxyuridine (EdU), conceptually, appears to be unique in terms of being immune to resistance development due to its high similarity to thymidine (see **Fig. 1** for the chemical structures of TMZ and EdU).

EdU is commonly used to study DNA replication via "click chemistry" (18–21). During our attempt to employ EdU to analyze nucleotide excision repair patch size in human cell lines, we discovered that EdU incorporated into the human genome is recognized as "damage" and removed from the genome by the classic human dual incision mechanism in the form of 26-27 nt-long oligomers, leading to a futile cycle of incorporation into DNA and its removal (22). This realization, combined with previous reports on toxicity of EdU in tissue culture as well as its established property of crossing the blood brain barrier led to the notion that EdU might be effective against brain cancer (23,24). In our previous study, we showed that EdU is effective against GBM cell lines *in vitro*, orthotopic *in vivo* mouse models of GBM established with intracranial engrafts of mouse glioma and human GBM cells, and living, passage-zero patient GBM tissues tested *ex vivo* on an organotypic brain slice culture (OBSC)-based platform (25). Here, building on these proof-of-concept studies, we sought to investigate the therapeutic potential of combining TMZ with EdU for GBM, reasoning that their distinct mechanisms of action and the conceptual improbability of

resistance to EdU—given its close structural similarity to thymidine—might reduce recurrence
rates and even enable durable remission in a subset of patients. To this end, in this study, we
tested the synergistic effect of TMZ+EdU in three different GBM cell lines *in vitro*, in *in vivo*
orthotopic mouse models of GBM established with these three GBM cell lines, and in living,
passage-zero GBM patient tissues seeded on OBSCs. Our results show the combination of TMZ
and EdU as a new and promising therapeutic avenue, yielding synergistic effects in GBM and
having the potential to ultimately improve the poor prognosis faced by patients with this disease.
97
RESULTS
Targeted localization and pharmacokinetic features of EdU *in vivo*
In our previous study, we showed the targeted localization of EdU into the human GBM cell lines
seeded on OBSCs (25). In this study, we started out by testing the targeted localization of EdU
into tumor cells *in vivo*. To this end, we established orthotopic GBM tumors in female athymic
nude mice by inoculating FLuc expressing U87 human GBM cells into their brain parenchyma.
Since U87 tumor cells form large and well-demarcated tumors upon intracranial inoculation, we
chose this *in vivo* orthotopic mouse model of GBM to test the targeted localization of EdU into
tumor cells in the brain (26,27). Upon confirming tumor establishment via bioluminescent imaging
(BLI), we randomized these mice into four different groups to be injected with PBS Control, 200
mg/kg EdU, 5 mg/kg TMZ, or 5 mg/kg TMZ + 200 mg/kg EdU. These injections were administered
for 3 consecutive days, and the brains of mice were harvested the following day to perform
hematoxylin and eosin (H&E), nuclear (DAPI) and EdU staining. In **Figure 2**, the presence of
tumor mass is seen with H&E staining. Once the tumor cells were confirmed with H&E, EdU
staining via click chemistry along with nuclear staining (DAPI) was performed on the slices from
the same area of the brains. As negative controls, there was no EdU signal detected in either
tumor or non-tumor cells in the brains of mice administered with PBS or 5 mg/kg TMZ, pointing
out the specificity of EdU staining via click chemistry (**Fig 2A, 2B, 2E and 2F**). On the other hand,
the EdU signal (green) was highly localized to the tumor cells in mice treated with both 200 mg/kg
EdU (**Fig. 2C**) and 5 mg/kg TMZ + 200 mg/kg EdU (**Fig. 2D**). The quantification of EdU-positive
cells in tumor versus non-tumor cells revealed ~83 times more EdU-positive cells in 200 mg/kg
EdU-treated group and ~70 times more EdU positive cells in 5 mg/kg TMZ and 200 mg/kg EdU-
treated group (**Fig. 2G, ****p<0.0001 for both comparisons**). The residual EdU signal observed
in **Fig. 2C-2D** was localized mostly to the subventricular zone (SVZ) of the brain, which is known
to harbor some proliferative cells (28). These results confirmed the highly targeted localization of
EdU into tumor cells *in vivo*. To our knowledge, EdU's distribution in blood plasma and brain has
not been characterized before. To characterize this biodistribution and clearance of EdU *in vivo*,
we next performed a pharmacokinetic study with healthy female athymic nude mice. We
administered a single dose of 200 mg/kg EdU, collected blood plasma and brain samples from
mice at six different time points ranging from 15 mins to 6 hrs, and obtained the EdU concentration
over time via LC-MS/MS (**Fig. 3A**). In plasma, the maximum concentration of EdU was reached
at 15 minutes at ~73 μg/ml, with a half-life of ~10 minutes (**Fig. 3B**). On the other hand, in brain,
the maximum concentration of EdU was reached at 15 minutes at ~4.8 μg/ml, with a half-life of
~30 minutes (**Fig. 3C**). The mean $AUC_{0-\infty}$ values for plasma and brain were 42.14 and 4.92,
respectively, corresponding to a brain-to-plasma (B/P) AUC ratio of approximately 0.12 for EdU.
Together, these results supported our hypothesis that EdU gets specifically localized into the
tumor cells in the brain *in vivo* and identified the pharmacokinetic parameters of EdU in blood
plasma and brain tissue for the first time.
136
Testing TMZ+EdU combination therapy *in vitro* in human GBM cells and *in vivo* in
orthotopic mouse models of GBM
It has been found that combinations of various mutations in oncogenes and tumor suppressor
genes give rise to GBM (5). Human GBM cell lines derived from patients are widely used in testing

drug candidates both *in vitro* and *in vivo*. To recapitulate this heterogeneous genetic profile, in our
study, we used 3 human GBM cell lines with different sets of mutations (27,29–34) in testing the
combination of TMZ and EdU: U87, GBM8, and LN229. All of these cell lines are engineered to
express firefly luciferase (FLuc) and thus bioluminescence signal measured upon incubation with
luciferin is used as a measure of viability. We tested these cell lines with the standard of care drug
TMZ and EdU, and the TMZ+EdU combination. Testing was done by cell survival assay *in vitro*
and by monitoring survival and tumor growth *in vivo* in female athymic nude mice bearing
orthotopic xenografts of these 3 cell lines.
149
Testing TMZ+EdU combination therapy in vitro in U87 cell line and in mice bearing
orthotopic xenografts of U87
We first tested for a potential synergistic effect of TMZ and EdU combination on U87 cells *in vitro*
before proceeding to *in vivo* with this cell line. We have reported the IC50 value of EdU for this
cell line as 40 μM (25). Cell survival assay with the combination of EdU at this concentration and
various doses of TMZ showed that these two drugs are synergistic, especially at the higher doses
of TMZ (**Fig. S1A**). This *in vitro* synergism between TMZ and EdU encouraged us to test this
combination *in vivo* in the mice bearing orthotopic xenografts of U87 cells.
To establish the *in vivo* orthotopic mouse model with the U87 cell line, we inoculated U87 cells
intracranially into the brain parenchyma of female athymic nude mice and confirmed
establishment of tumors via BLI. Informed by our results on the pharmacokinetic features of EdU,
we decided to administer EdU at 200 mg/kg every Monday-Friday at a once daily frequency and
TMZ at 5 mg/kg every Monday, Wednesday, and Friday once every other day for a total of 6
weeks. For the combination group, mice received the TMZ injections in the morning and EdU
injections in the afternoon with the indicated regimens for 6 weeks. Upon randomization of mice
into PBS control and three treatment groups on Day 7 (**Fig. S1B**), we monitored the tumor growth
of these mice via BLI. Neither 200 mg/kg EdU nor 5 mg/kg TMZ treatment showed a successful
suppression of tumor growth while the treatment with the combination of 200 mg/kg EdU and 5
mg/kg TMZ suppressed the tumor growth and decreased the tumor signal below the baseline
level (**Fig. 4A**). This suppression lasted as long as the injections continued. However, once the
injections were completed by Day 46, the tumors recurred in 7-10 days, and the mice started to
reach humane end point after ~15 days following the discontinuation of the treatment. **Figure 4B**
shows the BLI images of representative mice throughout the study (see **Fig. S1B-G** for extended
BLI images). While treatment with single-agent 200 mg/kg and 5 mg/kg TMZ extended the survival
by ~30% and ~50%, respectively, compared to the PBS control group (p<0.0008 and p<0.0001,
respectively), combination of 5 mg/kg TMZ and 200 mg/kg EdU extended the survival by more
than 130% compared to PBS control group (p<0.0001). Additionally, combination group extended
the survival by ~60% and ~80% compared to the single-agent TMZ and EdU groups, respectively
(**Fig. 3C,** p<0.0001 for both comparisons). At the study endpoint of this study, 1 out of 8 mice in
the combination group showed no detectable tumor BLI signal, and the brain of this mouse did
not show detectable tumor cells at or around the tumor cell inoculation site (**Fig. S1G-H**).
Moreover, at the study endpoint, ~%12 of mice (1 out of 8 mice) in 5 mg/kg TMZ + 200 mg/kg
EdU group was alive. Thus, we concluded that TMZ and EdU combination showed a synergistic
effect *in vivo* in suppressing the tumor growth and increasing the survival in mice bearing
orthotopic xenografts of U87 cells.
185
Testing the synergistic effect of TMZ and EdU combination in vitro in the GBM8 cell line
and in mice bearing orthotopic xenografts of GBM8
Upon obtaining these results with the U87 *in vivo* model, we moved on to test the potential
synergistic effect of the combination of TMZ and EdU on GBM8 cell line *in vitro*. Previously, the
IC50 value of EdU for this cell line was identified as ~50 nM (25). Cell survival assay combining

this IC50 value of EdU with various doses of TMZ showed synergistic to additive effect at the high
doses of TMZ (**Fig. S2A**). Although the synergism observed in GBM8 was not as high as that in
U87 in cell survival assay, it was encouraging enough to proceed with GBM8 cell line to establish
an *in vivo* orthotopic mouse model of GBM.

To this end, we started out by inoculating GBM8 cells intracranially into the brain parenchyma of
female athymic nude mice. After confirmation of tumor establishment via BLI, mice were
randomized into PBS control and treatment groups on Day 9 (**Fig. S2B**). Considering the inherent
sensitivity of GBM8 cell line to TMZ, we chose two different TMZ doses in this model; the first
being low dose (1 mg/kg) and the second being high dose (5 mg/kg). For the dose of EdU
treatment, we used 200 mg/kg again as in U87 *in vivo* model, considering the fast clearance of
the molecule. The combination groups featured combination of 1 mg/kg TMZ and 200 mg/kg and
the combination of 5 mg/kg TMZ and 200 mg/kg, separately. As in the U87 *in vivo* model, 6 weeks
of injections were administered with the M-F and MWF treatment regimens for EdU and TMZ,
respectively. Unlike the U87 *in vivo* model, following the end of 6 weeks of injections, we wanted
to investigate tumor recurrence and response to maintenance therapy in all treatment groups.
Thus, we set a recurrence criteria per mouse where treatment was resumed when the tumor's
BLI signal increased to ≥10× its baseline (initial) value and a discontinuation criteria per mouse
where reinjections were discontinued once the tumor's BLI signal returned to and stabilized at the
baseline level. The tumor growth was assessed via BLI in two phases: first phase during the
administration of 6 weeks of injections (Day 7 to Day 50) and the second phase during the
monitoring of tumor recurrence and response to as-needed maintenance therapy (Day 50 to study
endpoint, Day 170).

Assessment of tumor growth and recurrence during the first phase of treatment and the second phase of maintenance therapy in mice bearing orthotopic xenografts of GBM8

Assessment of tumor growth and recurrence during the first phase of treatment and the
second phase of maintenance therapy in mice bearing orthotopic xenografts of GBM8
During the first phase, we observed suppression of tumor growth in all mice except the ones that
received PBS and 1 mg/kg TMZ (**Fig. 5A**, **5C**, **S2B**, **S2C**, **S3A-F**). During the second phase, when
the initial treatment regimen had been discontinued, we started to observe recurrence in some
mice in the 5 mg/kg TMZ, 200 mg/kg EdU, and 1 mg/kg TMZ + 200 mg/kg EdU groups and
initiated maintenance therapy on recurrent mice on Days 59 and 63 (**Fig. 5B**, **S3B**, **S3C**, **S3F**).
Recurrence occurred in ~60% (4 out of 7) of mice in the 5 mg/kg TMZ group, 100% (6 out of 6)
of mice in the 200 mg/kg EdU group, and ~15% (1 out of 7) of mice in the 1 mg/kg TMZ + 200
mg/kg EdU group. In contrast, no recurrence was observed in any mice in the 5 mg/kg TMZ + 200
mg/kg EdU group (**Fig. 5B**, **S3D**). According to the tumors' BLI signals, none of the recurred mice
in the 5 mg/kg TMZ group responded to maintenance therapy while 50% of the recurred mice in
the 200 mg/kg EdU group responded to maintenance therapy and showed remission (**Fig. 5B**,
S3B, **S3C**). One case of recurrence observed in the 1 mg/kg TMZ + 200 mg/kg EdU group also
showed suppression of tumor growth and eventually remission upon receiving maintenance
therapy (**Fig. 5B, 5C** and **S3F**; see **Fig. S2B-G** for extended BLI images). From these results, we
concluded that maintenance therapy with single-agent EdU and combination of TMZ and EdU
helped with suppressing the tumors while maintenance therapy with single-agent TMZ did not
provide any benefit in suppressing the tumors.
232
Monitoring survival in mice bearing orthotopic xenografts of GBM8

Monitoring survival in mice bearing orthotopic xenografts of GBM8
In addition to assessing tumor growth and recurrence, we also monitored survival throughout the
study in the *in vivo* GBM8 model. At the study endpoint, single-agent treatments with 1 mg/kg
TMZ, 5 mg/kg TMZ, and 200 mg/kg EdU extended the survival significantly by ~10%, ~200%, and
~260%, respectively, compared to the PBS control group (p=0.0037, p=0.0005 and p=0.0011,
respectively, **Fig. 5D**). Moreover, 50% of mice (3 out of 6 mice) in 200 mg/kg EdU group and
~30% of mice (2 out of 7 mice) in 5 mg/kg TMZ group were alive by the study endpoint.

Importantly, all of the mice in 1 mg/kg TMZ + 200 mg/kg EdU and 5 mg/kg TMZ + 200 mg/kg
combination groups were alive and tumor-free by the study endpoint, conferring a complete
survival at the study endpoint in this stringent patient-derived model and highlighting the strong
therapeutic potential of TMZ+EdU combination (**Fig. 5D**). We concluded from these results that
the combination of TMZ and EdU conferred a synergistic effect in terms of both suppressing the
tumor growth and extending the survival in mice bearing orthotopic xenografts of GBM8 cells.
Combined with the results of the U87 *in vivo* model, these results supported our hypothesis that
combining TMZ and EdU could yield a greater therapeutic effect than either of these drugs confers
alone.
249
Testing TMZ+EdU combination therapy in vitro in LN229 cell line and in mice bearing
orthotopic xenografts of LN229
Upon observing these synergistic results with the GBM8 *in vivo* model, we wished to test this
synergistic effect in a third orthotopic *in vivo* model using LN229 cell line. Before establishing the
in vivo orthotopic model with this cell line, we performed cell survival assay to test the synergistic
effect of TMZ+EdU *in vitro* by combining the IC50 value of EdU for LN229 (~20 μM, as identified
previously (25)) with various doses of TMZ. As in the other two GBM cell lines, the TMZ+EdU
combination was more efficient than either drug alone at higher concentrations of TMZ (**Fig S4A**).
These synergistic results led us to test the synergism of TMZ+EdU combination in the *in vivo*
orthotopic model with LN229 cell line.
Accordingly, we created the *in vivo* orthotopic model by inoculating LN229 cells intracranially into
the brain parenchyma of female athymic nude mice. Upon confirming the establishment of tumors
via BLI, we stratified the mice into PBS control and five treatment groups on Day 4 (**Fig. S4B**).
We chose the same treatment groups and treatment regimen with 6 weeks of injections as in the
GBM8 *in vivo* model. By the third week of injections, mice in PBS control group started to reach
humane end point while the tumors' BLI signals in all the other groups, except 200 mg/kg EdU
group, decreased below the baseline level, indicating suppression of tumor growth (**Fig. 6A**, **6B**,
S4C). On the other hand, 200 mg/kg EdU treatment was only able to stabilize the tumors' BLI
signals at ~3 times the baseline level signal. By the completion of 6 weeks of injections on Day
46, all the mice in the PBS control group reached humane endpoint and the tumors' BLI signals
in 200 mg/kg EdU groups started to increase. However, the tumors' BLI signals in TMZ single-
agent and TMZ+EdU combination groups decreased further and stabilized ~0.01 times the
baseline levels (**Fig. 6A**). Around a month after the completion of injections by Day 70, mice
treated with 200 mg/kg EdU started to reach humane endpoint while tumors' BLI signals in
combination or single-agent TMZ groups remained lower than the baseline level signal (**Fig. 6A**,
S4D-F). Additionally, around Day 80 in 25% (2 out of 8 mice) of mice in 1 m/kg TMZ group and
on Day 85 in ~12% (1 out of 8 mice) of mice in 5 mg/kg TMZ group, we started to observe some
recurrence (**Fig. 6A**, **6B, S4G**). However, this recurrence was not observed in either of the
combination groups. For survival, treatment with 200 mg/kg EdU extended the median survival
significantly by ~110% compared to the PBS control group (p=0.0002, **Fig. 6C**). At conclusion of
the study, none of the mice in 5 mg/kg TMZ + 200 mg/kg EdU group reached the humane
endpoint, resulting in a striking and significant increase in median survival compared to both PBS
control and single-agent EdU treatment groups (p<0.0001 and p=0.0008, respectively, **Fig.6C**).
Similarly, 1 mg/kg TMZ + 200 mg/kg EdU group extended the survival significantly compared to
both PBS control and single-agent EdU group (p<0.0001 and p=0.0018, respectively, **Fig.6C**). In
1 mg/kg TMZ + 200 mg/kg EdU group, only one mouse reached humane endpoint due to orbital
infection, although its tumor had way lower BLI signal compared to the baseline level,
corresponding to ~88% survival at the study endpoint. However, different from our results in U87
and GBM8 *in vivo* models, in the LN229 *in vivo* model, we observed that combination of TMZ and
EdU performed as well as the single-agent TMZ groups, leading to an additive effect in

suppressing the tumor growth and prolonging the median survival (**Fig. 6C**). Collectively, the
results from our *in vivo* orthotopic mouse models of GBM established with U87, GBM8 and LN229
cells provided preclinical evidence that TMZ+EdU combination could elicit a synergistic to partly
additive effect in GBM, supporting the progression of this combination to early-phase clinical
testing.

Toxicity profiling of mice treated with EdU, TMZ and their combination in *in vivo* orthotopic mouse model of U87

To assess treatment-associated toxicity, we collected peripheral organs (intestine, kidney, spleen,
and liver) from mice bearing orthotopic xenografts of U87 at the humane endpoint. Organs were
fixed, processed, paraffin-embedded, sectioned, and stained with hematoxylin and eosin (H&E).
Slides were histopathologically evaluated and scored for organ-specific signs of toxicity (**Fig. S5**).
In the small intestine of U87 tumor-bearing mice, minimal to mild crypt epithelial
necrosis/apoptosis was observed in 8 of 10 animals treated with 200 mg/kg EdU. Regenerative
activity, evidenced by increased mitotic figures, was also present in some animals. In mice treated
with 5 mg/kg TMZ + 200 mg/kg EdU, no overt crypt necrosis/apoptosis was seen, but increased
mitotic figures were detected in 2 of 5 animals, suggesting earlier crypt injury followed by
regeneration. These mice survived ~30 days longer than EdU-only animals, supporting the
interpretation that the intestinal epithelium had recovered from prior EdU-related injury. No crypt
changes were observed in mice treated with 5 mg/kg TMZ or PBS. The mild severity and evident
reversibility of the crypt epithelial injury indicate a non-adverse effect. In the kidney, minimal
multifocal cortical tubular epithelial degeneration/regeneration was noted in 2 of 10 EdU-treated
animals and 1 of 5 animals receiving the TMZ+EdU combination, but not in TMZ-only or PBS-
treated mice. As this lesion is a common background finding and may occur with general
debilitation, its low incidence and severity suggest a non-adverse, possibly incidental effect. In
the spleen, increased extramedullary hematopoiesis (EMH) was observed in 7 of 10 mice treated
with EdU and 4 of 5 mice treated with TMZ+EdU, but not in TMZ-only animals and only minimally
in 1 of 7 PBS controls. This finding is considered EdU-related and likely reflects bone marrow
suppression. Slight decreases in splenic lymphocyte cellularity were also seen in EdU- and
combination-treated mice; however, similar changes occurred in PBS controls, suggesting a
stress-related rather than treatment-specific effect. In the liver, no hepatocellular injury was
detected. Minor group differences in hepatocellular vacuolation were consistent with expected
diurnal variation in glycogen content (hepatocytes from animals euthanized earlier in the day
typically exhibit greater glycogen-associated vacuolation than those euthanized later) (35–38).
Overall, these observations demonstrate that EdU, alone or in combination with TMZ, induced
only mild and reversible histopathological changes in peripheral organs. This conclusion is also
supported by the recovery of body weight after treatment in mice bearing U87, GBM8, and LN229
tumors (**Fig. S6A-C**).

In addition to these toxicity profiling of the peripheral organs, we also wanted to test the effect of
the treatment with EdU, TMZ and their combination on complete blood counts. To this end, we
collected blood samples from mice bearing LN229 tumors on three different days: two of them
during the 6 weeks of injections (Day 16 and Day 30) and one after the completion of 6 weeks of
injections (Day 58). According to the results of complete blood counts, there were only minor and
reversible differences between the counts of White Blood Cells (WBCs), Lymphocytes (LYMPH),
Neutrophils (NEUT), Basophils (BASO), Eosinophils (EO), and Red Blood Cells (RBC) values in
PBS control group on Day 16 and in all the other groups. Additionally, for all the counts of these
different cells, the values in all groups were mainly at or above the normal lower limit values by
Day 58 (**Fig. S7A-F**). These results, together with the results on the histopathological and body
weight changes, further confirmed that EdU, alone or in combination with TMZ, induced mild,
transient, and reversible toxicity.

Testing TMZ+EdU combination therapy against living, passage-zero GBM patient tumor tissues *ex vivo*

Upon collecting these promising *in vivo* results, we proceeded to test the efficacy of TMZ+EdU on living, passage-zero GBM patient tissues seeded atop organotypic brain slice cultures (25,39,40). In our previous investigation of EdU as a potential therapeutic for GBM, we observed differential response across all patient tumors tested with monotherapy EdU or TMZ (25). In these new experiments, we observed a similar trend. We exposed four IDH-wild-type WHO-classified uncultured patient GBM tissues to single-agent EdU or TMZ as well as a combination of both compounds. 3 out of 4 of these tissue samples (PTS125, PTS152, and PTS153) were from patients who clinically showed rapid tumor progression after receiving TMZ.

During treatment, TMZ and/or EdU were added under the transwell membranes supporting each tumor-bearing OBSC and allowed to diffuse through the OBSCs and into the tumor cells. All tumors were treated with EdU and/or TMZ at doses of 0, 100, 250, and 1000µM. Combination therapy of [TMZ + EdU] was given at doses of [0, 0], [100, 100], [250, 250] and [500 µM, 500 µM]. Among the tissues tested, we observed several doses where EdU alone or in combination with TMZ performed better than TMZ alone (**Fig. 7A-D**). Using the Zero Interaction Potency (ZIP) Synergy score (41,42) that measures the difference between observed effects of a combinatorial regimen and expected effects if individual drugs are not interactive, we found that the effect of the combinatorial therapy ranged from additive ($0 < ZIP < 10$ in PT158, PT125, PT152) to synergistic ($ZIP > 10$ in PT153) (**Fig. 7E**).

GBM tumors have significant intra- and inter-tumoral heterogeneity; therapies that are effective for some patients may not be effective for others, and vice versa. We therefore tested EdU against two passage-zero patient GBM tissues in combination with a range of other FDA-approved chemotherapies. We chose carboplatin and lomustine because of their cell cycle-independent anti-tumor activity to complement EdU's hypothesized activity during the S-phase (43,44), and also because lomustine is frequently used as a control arm in trials and sometimes considered the standard of care after GBM progression (43,45). We chose etoposide because its mechanism of action through the stabilization of the topoisomerase II-DNA cleavage complex is very different from EdU's and may elicit different downstream compensatory mechanisms. It has also been well-tolerated in heavily pretreated patients with refractory gliomas (46,47). We also chose the poly (ADP-ribose) polymerase-1 inhibitor (PARPi) Olaparib because it is under investigation as a radio and chemosensitizer (48,49). As such, it might potentiate EdU's ability to target glioma stem cells (GSCs) which have upregulated PARP that promotes base excision repair and the repair of single-strand breaks in treated cancer cells (50).

To assess these exploratory combinatorial regimens, we first used doses previously determined to be close to the ED50 values of each single agent. We then combined these drugs with 1000 µM EdU. Across all drugs tested in combination, we observed more killing in the combination groups than when the drug was used alone (**Fig. 7F**, **7G**). Notably, at the same doses, each patient tissue responded differently to each drug. For instance, in PTS152, there was a 2.9-fold increase in tumor killing when EdU was added to etoposide, but just a 1.2 fold increase in PTS153, (**Fig. 7F**, **7G**), further highlighting the need for more personalized and patient-centric models that capture patient-to-patient heterogeneity.

DISCUSSION

As the most common primary malignancy of the central nervous system, GBM carries an extremely poor prognosis, with only ~7% of patients surviving five years after diagnosis (51–53). Therefore, there is an urgent need to develop new treatment modalities for GBM. Current efforts focus on investigating the therapeutic potential of integrating different drugs into combinatorial

388 treatments with TMZ, the first-line chemotherapeutic for GBM. However, despite trials and
389 preclinical studies with agents such as lomustine (54), bevacizumab (55), and PARP inhibitors
390 (56,57), none of these combinations have received FDA approval to date.
391
392 In this work, we demonstrated in (i) GBM cell lines *in vitro*, (ii) orthotopic mouse models of GBM
393 *in vivo*, and (iii) living, uncultured GBM patient tissues on OBSC that EdU shows a synergistic
394 effect in GBM when combined with TMZ. Moreover, for the first time, we characterized the
395 pharmacokinetic features of EdU and confirmed the targeted localization of EdU into GBM tumor
396 cells *in vivo* in both single-agent EdU-treated and combination-treated mice.
397
398 Our results on pharmacokinetic characteristics of EdU *in vivo* highlighted the fast clearance of
399 EdU with limited penetrance to the blood brain barrier and suggested that there is room for
400 improvement to modulate the bioavailability of EdU via formulation. Formulating EdU would
401 increase the bioavailability and in turn the penetrance of EdU to blood brain barrier, potentiating
402 the therapeutic effect we observed in combination with TMZ.
403
In our U87 *in vivo* model, we observed only a partial increase in survival with single-agent TMZ
and EdU while neither was able to suppress the tumor growth. However, combination of TMZ and
EdU showed a truly synergistic effect in this model by both suppressing the tumor growth and
extending the survival drastically. On the other hand, in our GBM8 *in vivo* model, single-agent
EdU and TMZ (5 mg/kg) treatments conferred considerable survival benefit compared to PBS
control group. Nevertheless, we also observed recurrence in these single-agent treatment groups
with partial to no response to maintenance therapy. For instance, we observed recurrence and no
response to maintenance therapy in ~60% of the mice treated with 5 mg/kg TMZ, reflecting the
acquired resistance following an initial positive response to TMZ treatment. Notably, this TMZ
resistance was overcome in mice treated with 5 mg/kg TMZ + 200 mg/kg EdU since no recurrence
was observed in these mice by the study endpoint. These results point out the power of TMZ+EdU
combination in targeting the TMZ-resistant tumors which is a prevalent occurrence in the clinic.
416
Surprisingly, in our third *in vivo* model with LN229, we observed that single-agent TMZ treatments
initially performed as well as the combination treatment groups, indicating an additive effect in this
model. However, both single-agent TMZ groups exhibited recurrence following the completion of
six weeks of injections. Combination groups in this LN229 model, on the other hand, were still
able to avoid this recurrence, similar to the results in GBM8 model while the survival benefit these
combination treatments conferred was not as drastic as in GBM8 model. The difference among
these different *in vivo* models of GBM could be attributed to the heterogeneous nature of this
disease and the differences among the genetic alterations harbored by these GBM cell lines.
Additionally, further *in vivo* studies in these and other models featuring various doses of EdU in
combination with TMZ could identify lower doses of EdU that could still show synergistic effect.
427
Furthermore, on our OBSC platform, TMZ+EdU acted in an additive or synergistic manner against
all four passage-zero patient tumor tissues. Notably, these findings demonstrate that the positive
interactions between EdU and TMZ extend beyond *in vitro* and animal models to actual
heterogeneous human tumors. The more muted responses measured here reflect what one may
expect to observe in the clinical setting, where tumors with heterogeneity in proliferation rate and
mutational drivers are typically not responsive to TMZ alone and respond incrementally to
combination therapies. The heterogeneity of patient-to-patient response also highlights the need
for more personalized and patient-centric models in drug development. Our OBSC platform may
therefore serve as a functional correlative biomarker to identify patients most likely to benefit from
TMZ+EdU or other combinatorial regimens. Expanding OBSC testing of EdU to a larger,

molecularly diverse cohort could refine the estimated prevalence of synergy, enable biomarker
discovery, and further advance functional precision medicine efforts in GBM.
440
New therapeutic avenues to pursue: EdU's potential in combination with various other
chemotherapeutics
Our results on combining EdU with various other chemotherapeutics against GBM patient tumor
tissues revealed the transformative role EdU could play in alternative treatment regimens for
GBM. By targeting complementary mechanisms of action in these combinations, we found
patient-specific sensitivities to different combination therapies. Future studies further testing the
mechanism and therapeutic potential of these combinatorial regimens will shed light on the clinical
utility of each in GBM. Introducing such personalized combination regimens into clinical GBM
treatment could provide much-needed alternatives for patients with GBM.
450
MATERIALS & METHODS
Cell Lines
U87 (purchased from ATCC in 2012, catalog #: HTB-14, RRID: CVCL_0022) and LN229
(purchased from ATCC in 2012, catalog #: CRL-2611, RRID: CVCL_0393) cells were cultured in
1X DMEM supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin in
T75 flasks. Patient-derived GBM8 cells (a kind gift from gift from H. Wakimoto, Massachusetts
General Hospital) (29,30) were cultured in neurobasal-A medium (Gibco, catalog #: 12349015)
supplemented with 1 mg heparin (STEMCELL Technologies, catalog #: 07980), 10 µg FGF
(Shenandoah Biotechnology, catalog #: 100-146), 10 mL B27 supplement (Life Technologies Inc.,
catalog #: A3582801), 7.5 mL L-glutamine (Life Technologies Inc., catalog #: 10888022), 10 µg
EGF (Shenandoah Biotechnology, catalog #: 100-26-100ug), 2.5 mL N2 supplement (Life
Technologies Inc., catalog #: 17502048), and 2.5 mL Antibiotic-Antimycotic (100X) (Thermo
Fisher Scientific, catalog #: 15240062) in T75 flasks.
464
Lentiviral vectors
The following lentiviral vector (LV) was used in this study: mCherry protein fused to firefly
luciferase (LV–mCh-FL). This LV also encodes a puromycin resistance gene. U87, GBM8 and
LN229 cell lines were transduced with this mCherry-FLuc Lentivirus. Following transduction,
mCherry expression was confirmed and cells expressing mCherry were selected using puromycin
treatment.
471
In vitro **survival assay via BLI**
The cells were counted using a Countess Cell Counter (Thermo Fisher Scientific, catalog #:
AMQAX1000) and seeded in triplicate with a 2500 cells/well density in black bottom 96 well plate.
Following 24 h incubation, the cells were treated with increasing doses of single agent TMZ
(Sigma-Aldrich, catalog #: T2577-100MG), single agent EdU (Sigma-Aldrich, catalog #: 900584-
500MG). For combination, IC50 value of the cell line for EdU was combined with increasing doses
of TMZ. At the end of 72 h treatment, the cell viability was measured via BLI using an AMI optical
imaging system. The data was analyzed using Aura software, Microsoft Excel, and GraphPad
Prism to plot dose-response curves ('Non-linear Regression (curve fit)' option used in GraphPad
Prism).
482
Assessing synergism between TMZ and EdU *in vitro*
The cell viability data obtained from 3 biological replicates of *in vitro* survival assay via BLI of U87,
LN229 and GBM8 cell lines for each single dose of single agent EdU, TMZ or their combination
were fed in CompuSyn software as Fa (fraction affected). The CompuSyn software then
generated "Logarithmic Combination Index Plot" for drug combinations for each cell line where
log of Combination Index (CI) is plotted against the Fa (58–60). In this plot, the log(CI)<0,

log(CI)=0 and log(CI)>0 show synergism, additive effect and antagonism between EdU and TMZ,
respectively.
491
Establishment of *in vivo* orthotopic xenograft mouse models of human GBM
All animal experiments were conducted in accordance with institutional guidelines and approved
by the Institutional Animal Care and Use Committee (IACUC) at the University of North Carolina
at Chapel Hill (UNC-CH) under UNC IACUC protocol 25-016. Female athymic nude mice (in-
house bred) were housed under sterile conditions in an AAALAC-accredited animal facility at
UNC-CH.

U87, LN229, and GBM8 glioma cells expressing luciferase were cultured and prepared for
intracranial (i.c.) inoculation into the brains of athymic female nude mice. U87 and LN229 cells
were detached using 0.05% Trypsin-EDTA (1X; Gibco, catalog #25300-054) and incubated for 5
minutes. GBM8 cells were detached using 1X Accutase (STEMCELL Technologies, catalog #:
07920) and incubated for 5 minutes, with declumping by pipetting at 2.5 minutes. After
detachment, cells were counted using a Countess II Cell Counter, and suspensions were
prepared in PBS at final concentrations of 3.3×10^4 cells/μL (U87), 5×10^4 cells/μL (GBM8), and
8.3×10^4 cells/μL (LN229). For intracranial inoculation, mice were anesthetized with isoflurane
(2.0–2.5% for induction; 2.5–3.0% for maintenance) and placed in a stereotactic apparatus
following confirmation of adequate anesthesia via a negative toe-pinch reflex. After surgical site
preparation, a midline scalp incision was made (<10 mm) and a ~1 mm burr hole was drilled to
facilitate needle insertion for intracranial injection. Each mouse was injected with a 3 μL volume
of the corresponding cell suspension of U87, GBM8 or LN229 cells using the following coordinates
relative to bregma: mediolateral (ML) = 2.5 mm, anteroposterior (AP) = 0.5 mm, dorsoventral (DV)
= 2 mm. The total cells inoculated were 100,000 cells per mouse for U87 (n=40), 250,000 cells
per mouse for LN229 (n=50), and 150,000 cells per mouse for GBM8 (n=42). The wounds were
closed with 3M Vetbond™ tissue adhesive. Mice were monitored during and post-surgery and
received meloxicam once daily for 48 hours postoperatively as established in the approved animal
protocol.

Monitoring tumor growth via BLI in *in vivo* orthotopic xenograft mouse models of human GBM
Four to nine days post-implantation of U87, GBM8 or LN229 cells into the brains of mice, mice
were injected with IVISbrite D-Luciferin Potassium Salt Bioluminescent Substrate (1g, XenoLight)
(Revvity, catalog #: 122799-5) at 150 mg/kg, anesthetized with isoflurane and imaged via
bioluminescence imaging (BLI) with IVIS Spectrum optical imaging system. The BLI signals
recorded were analyzed using the LivingImage software and mice were stratified into treatment
groups based on tumor burden as assessed by BLI. The monitoring of tumor growth in all of these
in vivo models continued throughout the study durations at a frequency of 2-3 times a week.
526
Administration of EdU, Temozolomide (TMZ), and their combination, monitoring of tumor growth and evaluation of therapeutic response in *in vivo* orthotopic xenograft mouse models of human GBM
For the U87 and LN229 models, mice received a total of 6 weeks of injections. The treatment
regimen for EdU was 200 mg/kg EdU once every weekday (M-F) and for TMZ either 1 mg/kg or
5 mg/kg TMZ every other weekday (MWF). For the combination group, the mice received TMZ
injections in the morning and EdU injections in the afternoon with the same regimens. The tumor
growth was monitored via bioluminescence imaging (BLI) of the mice using IVIS Spectrum device
through intraperitoneally injecting luciferin. The mice started to receive treatment with EdU, TMZ
and their combination 6 days and 4 days post tumor cell inoculation for U87 model and LN229
model, respectively. For the GBM8 model, the mice started to receive 6 weeks of injections with

the abovementioned regimens for EdU, TMZ and their combination 7 days post tumor cell
inoculation and the tumor growth was monitored similarly. Different from the other models, the
injections restarted once a mouse's tumor recurred in GBM8 model. The recurrence criterion was
recording a BLI signal that is 10 times higher than the initial signal recorded 7 days post tumor
cell inoculation. The injections continued until a consistently low BLI signal (4 BLI readings
showing either the same initial BLI signal or lower). The injections were discontinued once this
stopping criterion was met. Tumor growth was continuously monitored by BLI throughout the
duration of the studies in all models. In the U87 and LN229 models, mice underwent a 6-week
treatment regimen. EdU was administered at 200 mg/kg once daily on weekdays (Monday–
Friday), while TMZ was given at either 1 mg/kg or 5 mg/kg every other weekday (Monday,
Wednesday, Friday). For combination treatment, mice received TMZ injections in the morning and
EdU in the afternoon, following the same individual dosing schedules. Tumor growth was
monitored via bioluminescence imaging (BLI) using the IVIS Spectrum system, following
intraperitoneal injection of luciferin. Treatment began 6 days post–tumor cell inoculation for the
U87 model and 4 days post-inoculation for the LN229 model. In the GBM8 model, treatment also
began 7 days after tumor cell inoculation and followed the same 6-week regimen. Tumor
progression was monitored via BLI, as in the other models. Unique to GBM8, treatment resumed
upon tumor recurrence, defined as a BLI signal ≥10x the baseline signal recorded on day 7 post-
inoculation. Injections continued until the BLI signal returned to and remained at baseline or lower
for four consecutive readings, at which point treatment was discontinued. Tumor growth was
continuously monitored by BLI throughout the duration of the studies in all models, along with
survival tracking. Mice were also observed daily or every other day for signs of neurological
symptoms, ≥20% body weight loss, or decline in body condition score. Animals exhibiting any of
these signs were euthanized. Survival analysis was performed using the log-rank (Mantel–Cox)
test in GraphPad Prism.
563
Blood collection from mice bearing orthotopic xenografts of LN229 cells for complete
blood counts
Blood samples were obtained via a small venous nick on the mouse tail using a 22-gauge needle.
This procedure was performed on mice bearing orthotopic xenografts of LN229 cells and receiving
different treatment regimens and complete blood counts were assessed on days 16, 30, and 58
following tumor implantations. Blood collection volumes adhered strictly to the approved animal
use protocol. A maximum of three sampling sessions were conducted at bi-weekly intervals, with
each individual collection limited to approximately 30-50 µL. Complete blood counts (CBCs) were
performed on whole blood using an IDEXX ProCyte Dx™ Hematology Analyzer. Parameters
measured included indices of leukocyte, erythrocyte, and platelet lineages.
574
Processing, embedding, sectioning, staining and imaging of tissues from peripheral
organs of mice bearing orthotopic xenografts of U87 and GBM8 Cells
Formalin-fixed tissues were processed in a Leica Pegasus tissue processer, embedded in paraffin
(Leica Paraplast) using a HistoCore Arcadia H. Tissues were then sectioned on a microtome
(Leica, catalog #: RM2255) at 5 microns onto positively charged slides (Fisher Scientific, catalog
#: 12-550-15). For brains, the tissues were trimmed and sectioned around the site of tumor cell
inoculation in the right hemisphere at coordinates of 0.2 mm anterior-posterior (AP), 2.0 mm
mediolateral (ML), and 3.5 mm dorsoventral (DV).
H&E staining and imaging
Prior to H&E staining, slides were air dried overnight and baked for 30 minutes at 60 degrees
Celsius. H&E stains were performed using the Autostainer XL from Leica Biosystems. The slides
were deparaffinized in graded ethanol, then stained with Hematoxylin (Epredia, catalog #: 7211)
for 2 mins and Eosin -Y (Epredia, catalog #: 71311) for 1 min with respective washes in between.

Clarifier 2 (catalog #: 7402) and Bluing (catalog #: 7111) solutions from Epredia were used to
differentiate the reaction. After staining, slides were then dehydrated in graded ethanol, ending in
xylene, and coverslipped with Surgipath Micromount (Leica, catalog #: 3801731). Stained slides
were digitally imaged in the Aperio AT2 (Leica Microsystems) using 20x objective.

EdU and DAPI staining and imaging

The slides with brain tissues at the site of tumor inoculation were deparaffinized in graded ethanol.
They were then washed once with 1X Dulbecco's PBS (Gibco, catalog #: 14190144) for 10
minutes and once with 0.1% Triton X-100 (Sigma-Aldrich, catalog #: T9284-500ML) in PBS for 10
minutes in Coplin jars at room temperature. The slides were then permeabilized, blocked,
washed, and stained for EdU with Click-iT Plus Alexa Fluor picolyl azide toolkit (Thermo Fisher
Scientific, catalog #: C10641) and for DAPI with 1 µg/ mL Hoechst 33258, pentahydrate (bis-
benzimide) (Thermo Fisher Scientific, catalog #: H3569) as previously described (25). Upon
completion of these stainings, the slides were mounted with ProLong Gold Antifade Mountant
(Thermo Fisher Scientific, catalog #: P36934) and cured overnight.

The next day, slides were digitalized using the Aperio ScanScope FL (Aperio Technologies Inc).
The digital images were captured in each channel by 20x objective (0.468 µm/pixel resolution)
using line-scan camera technology (U.S. Patent 6,711,283). The adjacent 1 mm stripes captured
across the entire slide were aligned into a contiguous digital image by an image composer. Images
were archived in PSC's eSlide Manger database (Leica Biosystems).

OBSC generation

All animal work related to OBSC generation was approved by the Institutional Animal Care and
Use Committee at the University of North Carolina at Chapel Hill under protocol 22-171. P8
Sprague-Dawley rat pups were used for all OBSC preparation. Animals were housed with one
mother and no more than ten pups. All animals were euthanized prior to weaning. OBSCs were
generated as previously described. In brief, 300um coronal sections of rat brains were generated
with a VT1000S vibratome and suspended on a cell culture insert immersed in 1ml of brain slice
media consisting of "Neurobasal-A medium supplemented with 10% heat-inactivated pig serum,
5% heat-inactivated rat serum, 1 mM L-glutamine, 10 mM KCl, 10 mM HEPES, 1 mM sodium
pyruvate and 100 U/mL penicillin-streptomycin." (39,40) The slices were then incubated at 37°C,
5% CO2 and 95% humidity.

Patient tissue identification, procurement, and processing

All brain tumor specimens were collected at University of North Carolina Hospitals. A total of four
patients with primary or recurrent IDH-wild-type WHO-classified GBM diagnosis were included in
the present study. Appropriate written informed consent by parent/guardian and assent as
applicable was obtained under protocols where ethical approval was given by the Institutional
Review Board (IRB) in the Office of Human Research Ethics at the University of North Carolina
at Chapel Hill. All patients consented to non-interventional IRB Protocol #23-0834 (LCCC2212;
ClinicalTrials.gov NCT05978557), which began on 27/07/2023; estimated end date 01/03/2028.
The laboratory study team had no access to identifying patient information; de-identified or coded
diagnostic data describing tumor type and mutational status was provided by a Study Coordinator
or Honest Broker.

Upon standard of care surgery, leftover tumor tissues not necessary for diagnosis were
deidentified and transferred to the laboratory on wet ice. The tissue samples were immediately

washed with PBS and red blood cells (RBC) were lysed with an RBC lysis buffer. The tissue
samples were then mechanically dissociated with scalpels and cryopreserved at 150 mg of tissue
per ml of CryoStor CS10.
636

Patient tissue processing and drug killing

Originally cryopreserved patient tumor tissue samples were processed for drug dosing according
to established protocols at the Screening Live Cancer Explants Core Facility at UNC (25,39). In
brief, tissue samples were retrieved from liquid nitrogen storage, rapidly thawed, washed with
PBS and passed through a 100um strainer to approximate a single-cell suspension. The tissue
was then exposed to a LV–mCh-FL lentivirus to express both fluorescent mCherry and
bioluminescent firefly luciferase reporters. On Day 0, 1 mg of tissue was seeded atop each
hemisphere of brain slices suspended in a cell culture insert. On D1, corresponding doses of EdU,
TMZ, carboplation (Sigma Aldrich, catalog #: C2538-100MG), lomustine (Selleck Chemicals,
catalog #: S1840), Olaparib (Selleck Chemicals, catalog #: S1060), or etoposide (Sigma Aldrich,
E1383-100MG) diluted in 1 ml of brain slice media were added underneath the inserts. The treated
seeded tumors were then incubated for 3 days and bioluminescence readouts were obtained from
an AMI optical imaging system.

Synergy score calculation

652 All doses of drugs tested were exported into Synergy finder https://synergyfinder.org/ and synergy
653 scores were automatically calculated. Synergy scores less than -10, between -10 and +10, and
654 greater than +10 indicate antagonistic, additive, or synergistic interactions respectively.

Pharmacokinetic Study

Pharmacokinetic (PK) analyses were conducted in healthy 12-week-old female Nu/Nu mice. Each
animal received a single intraperitoneal (IP) dose of EdU (200 mg/kg). Blood and brain samples
were collected at 15 min, 30 min, 1 h, 2 h, and 4 h after the dose (n = 5 per time point) and at 6 h
(n = 3) under isoflurane anesthesia. Approximately 0.8–1.0 mL of blood was obtained via cardiac
puncture into K2EDTA tubes, immediately placed on ice, and centrifuged (5000 rpm, 5 min, 4 °C)
to obtain plasma. Brains were snap-frozen in liquid nitrogen. All samples were stored at −80 °C
until analysis.

LC–MS/MS Quantification of EdU in Plasma and Brain

For sample preparation, brains were weighed and diluted 1:2 with phosphate-buffered saline
(PBS), then homogenized using zirconium oxide beads in 2 mL tubes at 5000 RPM on a Precellys
24 homogenizer (VWR Bead Mill Homogenizer), applied twice for 20 seconds each.

Plasma (50 µL) or brain homogenate (diluted 1:1 with control plasma) was precipitated with
methanol containing D3-methotrexate (100 ng/mL) as the internal standard. After vortexing and
centrifugation (10,000 RPM, 10 min, 4 °C), 50 µL of the supernatant was diluted with 100 µL of
water for injection.

Liquid chromatography was performed on an Agilent 1260 Infinity II system equipped with a
Waters XSelect HSS T3 column (50 × 2.1 mm, 3.5 µm) at 30 °C. The mobile phases were (A)
water with 0.1% formic acid and (B) methanol with 0.1% formic acid. The gradient was 10% B (0–
1.0 min) to 50% B (3.0 min), returning to 10% B for a total run time of 7.0 min, at a flow rate of 0.2
mL/min. Injection volume was 4 µL.

Mass spectrometric detection used an Agilent 6470 triple quadrupole with AJS-ESI in positive-ion
MRM mode. The following transitions were monitored: m/z 253→136.9 (plasma) and 253→117
(tissue) for EdU, and m/z 458.2→311.2 for the internal standard. Quantification was performed

using peak-area ratios relative to matrix-matched calibration standards, following method
validation.

Pharmacokinetic analysis

PK parameters were determined by non-compartmental analysis as described previously (61).
Maximum plasma concentration (C_{max}) and time to maximum concentration (t_{max}) were derived
directly from concentration-time profiles. The area under the concentration–time curve from 0 to
24 hours (AUC_{0-24h}) was calculated using the linear trapezoidal rule. The terminal elimination rate
constant (kel) was obtained from the log-linear portion of the plasma concentration–time curve,
and the elimination half-life ($t_{1/2}$) was calculated as $ln2/k_{el.}$ The area under the curve extrapolated
to infinity ($AUC_{0-\infty}$) was determined as AUC_{0-24h} + ($C_{24h/kel}$). Apparent clearance (CL/F) was
calculated as $Dose/AUC_{total}$, and apparent volume of distribution (Vd/F) as $(CL/F)/k_{el}$.

Acknowledgements:

We wish to thank UNC Pathology Services Core (PSC) and **Nicholas Pankow** in the UNC PSC
for expert technical assistance with Histopathology and Digital Pathology. The UNC PSC is
supported in part by an NCI Center Core Support Grant (P30CA016086). We also thank **Dr. Rani
Suzanne Sellers** for expert assistance with toxicity scoring of peripheral organs and
histopathologic review of brain and peripheral organs. All experiments using passage-zero patient
tumor tissues was completed within the Screening Live Cancer Explants Core Facility at UNC.
This research is supported by the National Center for Advancing Translational Sciences through
NC TraCS (K12TR004416), and National Institute of Health (NIH) Grants ES033414 and
GM118102. H.K. is supported by Fulbright PhD Grant. We would like to also extend our thanks
to UNC Biomedical Research Imaging Center, UNC Advanced Translational Pharmacology and
Analytical Chemistry (ATPAC) Core, UNC Animal Clinical Chemistry Core, and UNC SLiCE Core.

REFERENCES

1. Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJB, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. N Engl J Med. 2005 Mar 10;352(10):987–96.

2. Rong L, Li N, Zhang Z. Emerging therapies for glioblastoma: current state and future directions. J Exp Clin Cancer Res. 2022 Apr 15;41(1):142.

3. Singh S, Dey D, Barik D, Mohapatra I, Kim S, Sharma M, et al. Glioblastoma at the crossroads: current understanding and future therapeutic horizons. Signal Transduct Target Ther. 2025 Jul 9;10(1):213.

4. Sipos D, Raposa BL, Freihat O, Simon M, Mekis N, Cornacchione P, et al. Glioblastoma: Clinical Presentation, Multidisciplinary Management, and Long-Term Outcomes. Cancers (Basel). 2025 Jan 5;17(1).

5. Haque A, Banik NL, Ray SK. Molecular alterations in glioblastoma: potential targets for immunotherapy. Prog Mol Biol Transl Sci. 2011;98:187–234.

6. Radaelli E, Ceruti R, Patton V, Russo M, Degrassi A, Croci V, et al. Immunohistopathological and neuroimaging characterization of murine orthotopic xenograft models of glioblastoma multiforme recapitulating the most salient features of human disease. Histol Histopathol. 2009 Jul;24(7):879–91.

7. Davis ME. Glioblastoma: overview of disease and treatment. Clin J Oncol Nurs. 2016 Oct 1;20(5 Suppl):S2-8.

8. Hanif F, Muzaffar K, Perveen K, Malhi SM, Simjee SU. Glioblastoma Multiforme: A Review of its Epidemiology and Pathogenesis through Clinical Presentation and Treatment. Asian Pac J Cancer Prev. 2017 Jan 1;18(1):3–9.

9. Goenka A, Tiek D, Song X, Huang T, Hu B, Cheng S-Y. The many facets of therapy resistance and tumor recurrence in glioblastoma. Cells. 2021 Feb 24;10(3).

10. Silver DJ, Lathia JD. Therapeutic injury and tumor regrowth: tumor resection and radiation establish the recurrent glioblastoma microenvironment. EBioMedicine. 2018 May;31:13–4.

11. Rapp M, Baernreuther J, Turowski B, Steiger H-J, Sabel M, Kamp MA. Recurrence pattern analysis of primary glioblastoma. World Neurosurg. 2017 Jul;103:733–40.

12. Jiang H, Yu K, Li M, Cui Y, Ren X, Yang C, et al. Classification of progression patterns in glioblastoma: analysis of predictive factors and clinical implications. Front Oncol. 2020 Nov 3;10:590648.

13. Kawauchi D, Ohno M, Honda-Kitahara M, Miyakita Y, Takahashi M, Yanagisawa S, et al. Clinical characteristics and prognosis of Glioblastoma patients with infratentorial recurrence. BMC Neurol. 2023 Jan 7;23(1):9.

755   14.   Yoon N, Kim H-S, Lee JW, Lee E-J, Maeng L-S, Yoon WS. Targeted genomic sequencing
756   reveals different evolutionary patterns between locally and distally recurrent glioblastomas.
757   Cancer Genomics Proteomics. 2020;17(6):803–12.

758   15.   Wang L, Jung J, Babikir H, Shamardani K, Jain S, Feng X, et al. A single-cell atlas of
759   glioblastoma evolution under therapy reveals cell-intrinsic and cell-extrinsic therapeutic
760   targets. Nat Cancer. 2022 Dec 20;3(12):1534–52.

761   16.   Chehade G, Lawson TM, Lelotte J, Daoud L, Di Perri D, Whenham N, et al. Long-term
762   survival in patients with IDH-wildtype glioblastoma: clinical and molecular characteristics.
763   Acta Neurochir (Wien). 2023 Apr;165(4):1075–85.

764   17.   Lindsey-Boltz LA, Sancar A. New biochemical approaches for treatment of glioblastoma.
765   J Biol Chem. 2025 Sep 20;110748.

766   18.   Fantoni NZ, El-Sagheer AH, Brown T. A Hitchhiker's Guide to Click-Chemistry with Nucleic
767   Acids. Chem Rev. 2021 Jun 23;121(12):7122–54.

768   19.   Salic A, Mitchison TJ. A chemical method for fast and sensitive detection of DNA synthesis
769   in vivo. Proc Natl Acad Sci USA. 2008 Feb 19;105(7):2415–20.

770   20.   Zeng C, Pan F, Jones LA, Lim MM, Griffin EA, Sheline YI, et al. Evaluation of 5-ethynyl-
771   2'-deoxyuridine staining as a sensitive and reliable method for studying cell proliferation in
772   the adult nervous system. Brain Res. 2010 Mar 10;1319:21–32.

773   21.   Maltsev DI, Mellanson KA, Belousov VV, Enikolopov GN, Podgorny OV. The bioavailability
774   time of commonly used thymidine analogues after intraperitoneal delivery in mice: labeling
775   kinetics in vivo and clearance from blood serum. Histochem Cell Biol. 2022
776   Feb;157(2):239–50.

777   22.   Wang L, Cao X, Yang Y, Kose C, Kawara H, Lindsey-Boltz LA, et al. Nucleotide excision
778   repair removes thymidine analog 5-ethynyl-2'-deoxyuridine from the mammalian genome.
779   Proc Natl Acad Sci USA. 2022 Aug 30;119(35):e2210176119.

780   23.   Ross HH, Rahman M, Levkoff LH, Millette S, Martin-Carreras T, Dunbar EM, et al.
781   Ethynyldeoxyuridine (EdU) suppresses in vitro population expansion and in vivo tumor
782   progression of human glioblastoma cells. J Neurooncol. 2011 Dec;105(3):485–98.

783   24.   Ligasová A, Strunin D, Friedecký D, Adam T, Koberna K. A fatal combination: a
784   thymidylate synthase inhibitor with DNA damaging activity. PLoS ONE. 2015 Feb
785   11;10(2):e0117459.

786   25.   Kaanoglu H, Adefolaju A, Fraley C, Shobande M, Dasari R, Mann B, et al. Repurposing
787   the DNA Labeling Agent EdU for Therapy against Heterogeneous Patient Glioblastoma.
788   Mol Cancer Ther. 2025 Aug 1;24(8):1213–25.

789   26.   Han J-H, Yoon JS, Chang D-Y, Cho KG, Lim J, Kim S-S, et al. CXCR4-STAT3 Axis Plays
790   a Role in Tumor Cell Infiltration in an Orthotopic Mouse Glioblastoma Model. Mol Cells.
791   2020 Jun 30;43(6):539–50.

792   27.   Schulz JA, Rodgers LT, Kryscio RJ, Hartz AMS, Bauer B. Characterization and
793   comparison of human glioblastoma models. BMC Cancer. 2022 Aug 3;22(1):844.

28.   Faiz M, Acarin L, Castellano B, Gonzalez B. Proliferation dynamics of germinative zone
cells in the intact and excitotoxically lesioned postnatal rat brain. BMC Neurosci. 2005 Apr
12;6:26.

29.   Sasportas LS, Kasmieh R, Wakimoto H, Hingtgen S, van de Water JAJM, Mohapatra G,
et al. Assessment of therapeutic efficacy and fate of engineered human mesenchymal stem
cells for cancer therapy. Proc Natl Acad Sci USA. 2009 Mar 24;106(12):4822–7.

30.   Wakimoto H, Kesari S, Farrell CJ, Curry WT, Zaupa C, Aghi M, et al. Human glioblastoma-
derived cancer stem cells: establishment of invasive glioma models and treatment with
oncolytic herpes simplex virus vectors. Cancer Res. 2009 Apr 15;69(8):3472–81.

31.   Sullivan JP, Nahed BV, Madden MW, Oliveira SM, Springer S, Bhere D, et al. Brain tumor
cells in circulation are enriched for mesenchymal gene expression. Cancer Discov. 2014
Nov;4(11):1299–309.

32.   Sesen J, Dahan P, Scotland SJ, Saland E, Dang V-T, Lemarié A, et al. Metformin inhibits
growth of human glioblastoma cells and enhances therapeutic response. PLoS ONE. 2015
Apr 13;10(4):e0123721.

33.   DepMap Cell Line Summary [Internet]. [cited 2025 Oct 14]. Available from:
https://depmap.org/portal/cell_line/ACH-000075?utm_source=chatgpt.com&tab=overview

34.   Ishii N, Maier D, Merlo A, Tada M, Sawamura Y, Diserens A-C, et al. Frequent Co-
Alterations ofTP53, p16/CDKN2A, p14 ,PTEN Tumor Suppressor Genes in Human Glioma
Cell Lines. Brain Pathol. 1999 Jul;9(3):469–79.

35.   Liver, Hepatocyte - Glycogen Accumulation and Depletion - Nonneoplastic Lesion Atlas
[Internet].   [cited   2025   Oct   14].   Available   from:
https://ntp.niehs.nih.gov/atlas/nnl/hepatobiliary-system/liver/Hepatocyte-
GlycogenAccumulationandDepletion?utm_source=chatgpt.com

36.   Fuller RW, Diller ER. Diurnal variation of liver glycogen and plasma free fatty acids in rats
fed ad libitum or single daily meal. Metab Clin Exp. 1970 Mar;19(3):226–9.

37.   Rothacker DL, Kanerva RL, Wyder WE, Alden CL, Maurer JK. Effects of variation of
necropsy time and fasting on liver weights and liver components in rats. Toxicol Pathol.
1988;16(1):22–6.

38.   Wells MY, Weisbrode SE, Maurer JK, Capen CC, Bruce RD. Variable hepatocellular
vacuolization associated with glycogen in rabbits. Toxicol Pathol. 1988;16(3):360–5.

39.   Mann B, Zhang X, Bell N, Adefolaju A, Thang M, Dasari R, et al. A living ex vivo platform
for functional, personalized brain cancer diagnosis. Cell Rep Med. 2023 Jun
20;4(6):101042.

40.   Satterlee AB, Dunn DE, Lo DC, Khagi S, Hingtgen S. Tumoricidal stem cell therapy
enables killing in novel hybrid models of heterogeneous glioblastoma. Neuro Oncol. 2019
Dec 17;21(12):1552–64.

41.   Yadav B, Wennerberg K, Aittokallio T, Tang J. Searching for Drug Synergy in Complex
Dose-Response Landscapes Using an Interaction Potency Model. Comput Struct
Biotechnol J. 2015 Sep 25;13:504–13.

42.   Ianevski A, He L, Aittokallio T, Tang J. SynergyFinder: a web application for analyzing
835   drug combination dose-response matrix data. Bioinformatics. 2017 Aug 1;33(15):2413–5.

836   43.   Weller M, Le Rhun E. How did lomustine become standard of care in recurrent
837   glioblastoma? Cancer Treat Rev. 2020 Jul;87:102029.

838   44.   Bristol-Myers Squibb Company. Paraplatin(carboplatin) [package insert] [Internet]. 2010
839   [cited   2025   Oct   16].   Available   from:
840   https://www.accessdata.fda.gov/drugsatfda_docs/label/2010/020452s005lbl.pdf

841   45.   Zeyen T, Treiber A, Schneider M, Potthoff A-L, Schaub C, Roever L, et al. Second-line
842   temozolomide   in   first   recurrent   MGMT-methylated   glioblastoma   after
843   lomustine/temozolomide: Efficacy and safety. Neurooncol Adv. 2025 Apr 28;7(1):vdaf084.

844   46.   Reyhanoglu G, Tadi P. Etoposide. StatPearls. Treasure Island (FL): StatPearls Publishing;
845   2025.

846   47.   van der Meulen M, Chahal M, Mason WP. The value of etoposide for recurrent glioma.
847   Can J Neurol Sci. 2024 Jul;51(4):509–12.

848   48.   Dhanavath N, Bisht P, Jamadade MS, Murti K, Wal P, Kumar N. Olaparib: A
849   chemosensitizer for the treatment of glioblastoma. Mini Rev Med Chem. 2025;25(5):374–
850   85.

851   49.   Hanna C, Kurian KM, Williams K, Watts C, Jackson A, Carruthers R, et al.
852   Pharmacokinetics, safety, and tolerability of olaparib and temozolomide for recurrent
853   glioblastoma: results of the phase I OPARATIC trial. Neuro Oncol. 2020 Dec
854   18;22(12):1840–50.

855   50.   Lesueur P, Lequesne J, Grellard J-M, Dugué A, Coquan E, Brachet P-E, et al. Phase I/IIa
856   study of concomitant radiotherapy with olaparib and temozolomide in unresectable or
857   partially resectable glioblastoma: OLA-TMZ-RTE-01 trial protocol. BMC Cancer. 2019 Mar
858   4;19(1):198.

859   51.   Kaina B. Temozolomide, procarbazine and nitrosoureas in the therapy of malignant
860   gliomas: update of mechanisms, drug resistance and therapeutic implications. J Clin Med.
861   2023 Nov 30;12(23).

862   52.   Slika H, Karimov Z, Alimonti P, Abou-Mrad T, De Fazio E, Alomari S, et al. Preclinical
863   models and technologies in glioblastoma research: evolution, current state, and future
864   avenues. Int J Mol Sci. 2023 Nov 14;24(22).

865   53.   Price M, Ballard C, Benedetti J, Neff C, Cioffi G, Waite KA, et al. CBTRUS Statistical
866   Report: Primary Brain and Other Central Nervous System Tumors Diagnosed in the United
867   States in 2017-2021. Neuro Oncol. 2024 Oct 6;26(Supplement_6):vi1–85.

868   54.   Herrlinger U, Tzaridis T, Mack F, Steinbach JP, Schlegel U, Sabel M, et al. Lomustine-
869   temozolomide combination therapy versus standard temozolomide therapy in patients with
870   newly diagnosed glioblastoma with methylated MGMT promoter (CeTeG/NOA-09): a
871   randomised, open-label, phase 3 trial. Lancet. 2019 Feb 16;393(10172):678–88.

55. Chinot OL, Wick W, Mason W, Henriksson R, Saran F, Nishikawa R, et al. Bevacizumab
plus radiotherapy-temozolomide for newly diagnosed glioblastoma. N Engl J Med. 2014
Feb 20;370(8):709–22.

56. Jones AB, Tuy K, Hawkins CC, Quinn CH, Saad J, Gary SE, et al. Temozolomide and the
PARP Inhibitor Niraparib Enhance Expression of Natural Killer Group 2D Ligand ULBP1
and Gamma-Delta T Cell Cytotoxicity in Glioblastoma. Cancers (Basel). 2024 Aug
15;16(16).

57. Yuan AL, Ricks CB, Bohm AK, Lun X, Maxwell L, Safdar S, et al. ABT-888 restores
sensitivity in temozolomide resistant glioma cells and xenografts. PLoS ONE. 2018 Aug
28;13(8):e0202860.

58. Chou T-C, Talalay P. Quantitative analysis of dose-effect relationships: the combined
effects of multiple drugs or enzyme inhibitors. Adv Enzyme Regul. 1984 Jan;22:27–55.

59. Chou: ComboSyn Inc; Paramus,(NJ): 2005 - Google Scholar [Internet]. [cited 2025 Oct
12].   Available   from:
https://scholar.google.com/scholar_lookup?title=ComboSyn%20Inc%20Paramus%20NJ&
author=TC%20Chou&author=N%20Martin&publication_year=2005&

60. Chou T-C. Drug combination studies and their synergy quantification using the Chou-
Talalay method. Cancer Res. 2010 Jan 15;70(2):440–6.

61. Akyel YK, Seyhan NO, Gül Ş, Çelik M, Taşkın AC, Selby CP, et al. The impact of circadian
rhythm disruption on oxaliplatin tolerability and pharmacokinetics in Cry1-/-Cry2-/- mice
under constant darkness. Arch Toxicol. 2025 Apr;99(4):1417–29.

 FIGURE LEGENDS

Figure 1: Chemical structures of temozolomide **(A)** and 5-ethynyl-2'-deoxyuridine (EdU) **(B)**.

Figure 2: Targeted localization of EdU into GBM tumor cells *in vivo* in single-agent EdU-
and TMZ+EdU combination-treated mice. Hematoxylin and eosin (H&E), nuclear (DAPI), EdU
staining, and merged DAPI and EdU staining pictures in the brains of PBS- **(A)**, 5 mg/kg TMZ-
(B), 200 mg/kg EdU- **(C)**, and 5 mg/kg TMZ + 200 mg/kg EdU-treated **(D)** mice bearing orthotopic
xenografts of U87. **(E)** Quantification of EdU-positive cells in non-tumor cells in the brains of PBS-
(n=4), 200 mg/kg EdU- (n=4), 5 mg/kg TMZ- (n=4), and 5 mg/kg TMZ + 200 mg/kg EdU-treated
(n=4) mice bearing orthotopic xenografts of U87 (One-way ANOVA, ****$p<0.0001$). **(F)**
Quantification of EdU-positive cells in tumor cells in the brains of PBS- (n=4), 200 mg/kg EdU-
(n=4), 5 mg/kg TMZ- (n=4), and 5 mg/kg TMZ + 200 mg/kg EdU-treated (n=4) mice bearing
orthotopic xenografts of U87 (One-way ANOVA, ****$p<0.0001$). **(G)** Quantification of EdU-positive
cells in tumor versus non-tumor cells in the brains of 200 mg/kg EdU- (n=4) and 5 mg/kg TMZ +
200 mg/kg EdU-treated (n=4) mice bearing orthotopic xenografts of U87 (Unpaired, two-tailed t-
test, ****$p<0.0001$).

Figure 3: Pharmacokinetic features of EdU in blood plasma and brain. (A) Overview of the
sample collection and analysis to identify pharmacokinetic parameters of EdU in blood plasma
and brain of female athymic nude mice. **(B)** EdU concentration versus time curve in blood plasma,
and Area Under Curve (AUC), maximum concentration (C_{max}), time to reach maximum
concentration (t_{max}), half-life ($t_{1/2}$), apparent clearance (CL/F) and volume of distribution (V_d)
parameters for EdU in blood plasma. **(C)** EdU concentration versus time curve in brain, and Area
Under Curve (AUC), maximum concentration (C_{max}), time to reach maximum concentration (t_{max}),
and half-life ($t_{1/2}$) parameters for EdU in brain.

Figure 4: Testing the efficacy of TMZ+EdU in *in vivo* orthotopic U87 mouse model of GBM.
(A) Bioluminescence imaging (BLI) signal fold changes of the tumors of mice bearing orthotopic
xenografts of U87 and treated with either PBS (n=8), 200 mg/kg EdU (n=8), 5 mg/kg TMZ (n=8),
or 5 mg/kg TMZ + 200 mg/kg EdU (n=8). Arrows under the x axis denote the days on which six
weeks of EdU and TMZ injections were administered. **(B)** Representative BLI images of mice
treated with either PBS, 200 mg/kg EdU, 5 mg/kg TMZ, or 5 mg/kg TMZ + 200 mg/kg EdU on
Days 7, 23, 31, 38, and 58. **(C)** Kaplan Meier survival plot of mice bearing orthotopic xenografts
of U87 and treated with either PBS, 200 mg/kg EdU, 5 mg/kg TMZ, or 5 mg/kg TMZ + 200 mg/kg
EdU (log-rank test).

Figure 5: Testing the efficacy of TMZ+EdU in *in vivo* orthotopic GBM8 mouse model of
GBM. (A) Bioluminescence imaging (BLI) signal fold changes of the tumors of mice bearing
orthotopic xenografts of GBM8 and treated with either PBS (n=5), 200 mg/kg EdU (n=6), 5 mg/kg
TMZ (n=7), 5 mg/kg TMZ + 200 mg/kg EdU (n=7), 1 mg/kg TMZ (n=6), or 1 mg/kg TMZ + 200
mg/kg EdU (n=7) during the first phase of treatment (day 0 to Day 50). Arrows under the x axis
denote the days on which six weeks of EdU and TMZ injections were administered. **(B)**
Bioluminescence imaging (BLI) signal fold change of the tumors of mice bearing orthotopic
xenografts of GBM8 and treated with either PBS (n=5), 200 mg/kg EdU (n=6), 5 mg/kg TMZ (n=7),
5 mg/kg TMZ + 200 mg/kg EdU (n=7), 1 mg/kg TMZ (n=6), or 1 mg/kg TMZ + 200 mg/kg EdU
(n=7) during the second phase of maintenance therapy (day 50 to Day 170). Arrows on the graph
denote the day of the initiation of maintenance therapy in corresponding color-coded group. **(C)**
Representative BLI images of mice bearing orthotopic xenografts of GBM8 and treated with either
PBS, 200 mg/kg EdU, 5 mg/kg TMZ, 5 mg/kg TMZ + 200 mg/kg EdU, 1 mg/kg TMZ, or 1mg/kg
TMZ + 200 mg/kg EdU on Days 9, 42, 133, and 168. **(D)** Kaplan Meier survival plot of mice bearing
orthotopic xenografts of GBM8 and treated with either PBS, 200 mg/kg EdU, 5 mg/kg TMZ, 5
mg/kg TMZ + 200 mg/kg EdU, 1 mg/kg TMZ, or 1mg/kg TMZ + 200 mg/kg EdU (log-rank test).

Figure 6: Testing the efficacy of TMZ+EdU in *in vivo* orthotopic LN229 mouse model of GBM. (A) Bioluminescence imaging (BLI) signal fold changes of the tumors of mice bearing orthotopic xenografts of LN229 and treated with either PBS (n=7), 200 mg/kg EdU (n=7), 5 mg/kg TMZ (n=8), 5 mg/kg TMZ + 200 mg/kg EdU (n=8), 1 mg/kg TMZ (n=7), or 1 mg/kg TMZ + 200 mg/kg EdU (n=8). Arrows under the x axis denote the days on which six weeks of EdU and TMZ injections were administered. **(B)** Representative BLI images of mice treated with either PBS, 200 mg/kg EdU, 5 mg/kg TMZ, 5 mg/kg TMZ + 200 mg/kg EdU, 1 mg/kg TMZ, or 1 mg/kg TMZ + 200 mg/kg EdU on Days 4, 24, 70, and 98. **(C)** Kaplan Meier survival plot of mice bearing orthotopic xenografts of LN229 and treated with either PBS, 200 mg/kg EdU, 5 mg/kg TMZ, 5 mg/kg TMZ + 200 mg/kg EdU, 1 mg/kg TMZ, or 1 mg/kg TMZ + 200 mg/kg EdU (log-rank test).

Figure 7: Testing the efficacy of TMZ+EdU in living, uncultured GBM patient tissues seeded on OBSC. Dose-response curves and grouped dose-response bar graphs of GBM patient tissues PT153 **(A)**, PT158 **(B)**, PT125 **(C)**, and PT152 **(D)** treated with single-agent EdU, TMZ or TMZ+EdU (Two-way ANOVA, ****p<0.0001, ***p<0.001). **(E)** Zero Interaction Potency (ZIP) Synergy Score values and ZIP p values for TMZ+EdU combination tested on four GBM patient tissues. High-throughput combo screening of PT153 **(F)** and PT153 **(G)** with 1000 µM EdU in combination with carboplatin (50 µM), etoposide (10 µM), lomustine (250 µM), Olaparib (10 µM), and TMZ (2000 µM).

Figure S1: Testing the synergism of TMZ+EdU *in vitro* in U87 cell line, extended BLI images and H&E staining on the brains of mice bearing orthotopic xenografts of U87 cells. (A) Cell survival assay via BLI and CompuSyn assessment of synergy with combination index (CI) values on U87 cell line with the combination of 40 µM EdU (IC50 for U87) and various doses of TMZ (log(CI)<0, log(CI)=0 and log(CI)>0 show synergism, additive effect and antagonism between EdU and TMZ, respectively). Extended BLI images of mice bearing orthotopic xenografts of U87 cells on Days 7 **(B)**, 21 **(C)**, 31 **(D)**, 38 **(E)**, 48 **(F)**, and 58 **(G)**. **(H, I)** H&E staining showing no detectable tumor on the brains of 2 surviving mice from the 5 mg/kg TMZ + 200 mg/kg EdU treatment group.

Figure S2: Testing the synergism of TMZ+EdU *in vitro* in GBM8 cell line, extended BLI images of mice bearing orthotopic xenografts of GBM8 Cells. (A) Cell survival assay via BLI and CompuSyn assessment of synergy with combination index (CI) values on GBM8 cell line with the combination of 50 nM EdU (IC50 for GBM8) and various doses of TMZ (log(CI)<0, log(CI)=0 and log(CI)>0 show synergism, additive effect and antagonism between EdU and TMZ, respectively). Extended BLI images of mice bearing orthotopic xenografts of GBM8 cells on Days 9 **(B)**, 42 **(C)**, 94 **(D)**, 133 **(E)**, 150 **(F)**, and 168 **(G)**.

Figure S3: Spaghetti plots of bioluminescence imaging (BLI) signal fold changes of the tumors of mice bearing orthotopic xenografts of GBM8 and treated with either PBS (n=5) **(A)**, 200 mg/kg EdU (n=6) **(B)**, 5 mg/kg TMZ (n=7) **(C)**, 5 mg/kg TMZ + 200 mg/kg EdU (n=7) **(D)**, 1 mg/kg TMZ (n=6) **(E)**, or 1 mg/kg TMZ + 200 mg/kg EdU (n=7) **(F)**.

Figure S4: Testing the synergism of TMZ+EdU *in vitro* in LN229 cell line and extended BLI images of mice bearing orthotopic xenografts of LN229 Cells. (A) Cell survival assay via BLI and CompuSyn assessment of synergy with combination index (CI) values on LLN229 cell line with the combination of 20 µM EdU (IC50 for LN229) and various doses of TMZ (log(CI)<0, log(CI)=0 and log(CI)>0 show synergism, additive effect and antagonism between EdU and TMZ, respectively). Extended BLI images of mice bearing orthotopic xenografts of GBM8 cells on Days 4 **(B)**, 24 **(C)**, 64 **(D)**, 70 **(E)**, 73 **(F)**, and 98 **(G)**.

Figure S5: Toxicity profiling of the mice bearing orthotopic xenografts of U87 cells. Summary of toxicity findings on the peripheral organs (small intestine, kidneys, spleen, and liver) of mice bearing orthotopic xenografts of U87 cells and treated with either PBS, 200 mg/kg EdU, 5 mg/kg TMZ, or 5 mg/kg TMZ + 200 mg/kg EdU. Abbreviations: 200 mg/kg EdU abbreviated as

"EdU", 5 mg/kg TMZ abbreviated as "TMZ", 5 mg/kg TMZ + 200 mg/kg EdU abbreviated as
"TMZ+EdU". Grading: 0=no finding; 1=minimal finding; 2=mild finding; 3=moderate finding;
4=marked finding; 5=severe finding.

Figure S6: Mean body weight and mean body weight change of the mice bearing orthotopic
xenografts of U87 (A), GBM8 (B), or LN229 (C) cells.

Figure S7: Complete blood counts for mice bearing orthotopic xenografts of LN229 cells
treated with either PBS, 200 mg/kg EdU, 5 mg/kg TMZ, 5 mg/kg TMZ + 200 mg/kg EdU, 1
mg/kg TMZ, or 1 mg/kg TMZ + 200 mg/kg EdU on Days 16, 30 and 58. Complete counts of
white blood cells (WBCs) **(A)**, lymphocytes (LYMPH) **(B)**, neutrophils **(C)**, basophils (BASO) **(D)**,
eosinophils (EO) **(E)**, and red blood cells (RBC) **(F)** for mice bearing orthotopic xenografts of
LN229 cells treated with either PBS, 200 mg/kg EdU, 5 mg/kg TMZ, 5 mg/kg TMZ + 200 mg/kg
EdU, 1 mg/kg TMZ, or 1 mg/kg TMZ + 200 mg/kg EdU on Days 16, 30 and 58. Lower limit of 95%
CI of mean blood counts for female BALB/c nude mice (n=136): White blood cells: 2.97 K/µL,
lymphocytes: 1.25 K/µL, neutrophils: 1.02 K/µL, basophils: 0.00 K/µL, eosinophils: 0.01 K/µL,
and red blood cells: 8.87 M/µL. One-way ANOVA, *$p < 0.05$. Abbreviations: 200 mg/kg EdU
abbreviated as "EdU", 5 mg/kg TMZ abbreviated as "TMZ (H)", 5 mg/kg TMZ + 200 mg/kg EdU
abbreviated as "TMZ (H) + EdU", 1 mg/kg TMZ abbreviated as "TMZ (L)", and 1 mg/kg TMZ +
200 mg/kg EdU abbreviated as "TMZ (L) + EdU".

Figure 1

A

Temozolomide (TMZ)

B

5-ethynyl-2'-deoxyuridine (EdU)

Figure 2

Figure 3

A

B

Plasma Concentration-Time Curve of EdU (200 mg/kg) in Healthy Female Nu/Nu Mice

Plasma Pharmacokinetic Parameters of EdU in Healthy Nu/Nu/ Mice	
AUC_{0-4h} (μ.h/ml)	42.14 ± 2.54
$AUC_{0-\infty}$ (μ.h/ml)	42.14 ± 2.54
C_{max} (μg/ml)	73.25 ± 2.28
t_{max} (h)	0.25
$t_{1/2}$ (h)	0.19 ± 0.01
CL/F (ml/h)	125.12 ± 7.14
V_d (ml)	34.56 ± 0.3

C

Brain Concentration-Time Curve of EdU (200 mg/kg) in Healthy Female Nu/Nu Mice

Brain Pharmacokinetic Parameters of EdU in Healthy Nu/Nu/ Mice	
AUC_{0-4h} (μ.h/ml)	4.89 ± 0.18
$AUC_{0-\infty}$ (μ.h/ml)	4.92 ± 0.2
C_{max} (μg/ml)	4.78 ± 0.45
t_{max} (h)	0.25
$t_{1/2}$ (h)	0.45 ± 0.07

Figure 4

A

BLI Signal Fold Changes of U87-bearing Mice

B

C

Figure 5

Figure 6

Figure 7

E

Patient ID	Drug1	Drug2	ZIP Synergy Score	ZIP Synergy p value	Status of Interaction Between Drugs
153	EdU	TMZ	14.3	5.06E-02	Synergistic
158	EdU	TMZ	3.0	4.64E-01	Additive
125	EdU	TMZ	2.7	7.07E-01	Additive
152	EdU	TMZ	2.2	7.75E-01	Additive

F PTS153 High ThroughPut EdU Combo Screening

G PTS152 High ThroughPut EdU Combo Screening