TMEM55B controls lipolysis via lysosomal pH and ER–lysosome calcium signaling

Nagi Mukae[1], Honoka Maki[1] & Michiko Shirane[1]*

[1]Department of Molecular Biology, Graduate School of Pharmaceutical Sciences, Nagoya City University, Nagoya, Aichi, Japan.

*Corresponding: M.S. (shiram@phar.nagoya-cu.ac.jp)

Keywords:
TMEM55B, PDZD8, Lysosomes, Endoplasmic reticulum, Calcium, Membrane contact sites

ABSTRACT

A cholesterol transfer protein PDZD8 promotes lipolysis by facilitating endosomal maturation through membrane contact sites (MCSs) between endoplasmic reticulum (ER) and endolysosomes (LE/Ly). Endosomal maturation are suggested to be associated with endosomal acidification and calcium dynamics. However, the detailed mechanism by which the ER protein PDZD8 promotes endosomal maturation remained unclear. Here, we identified TMEM55B as a PDZD8-binding protein that localizes to lysosomes and is involved in cholesterol metabolism. TMEM55B functions as a regulator of the v-ATPase complex. Suppression of TMEM55B expression reduced lysosomal acidification, leading to impaired lipid droplet degradation, as well as decreased lysosomal calcium release and uptake, which in turn diminished calcium-induced calcium release (CICR) at ER–LE/Ly MCSs. These findings indicate that the lysosomal protein TMEM55B regulates lysosomal acidification and calcium dynamics through its interaction with the ER protein PDZD8, thereby promoting endosomal maturation and lipid metabolism.

Introduction

ER- LE/Ly MCSs are not formed by random contacts between the ER and endosomes but are mediated by specific membrane tethering complexes. ORP1L and VAP were first identified as tethering factors of ER–LE/Ly MCSs, and subsequent studies revealed that many sterol transfer proteins—including ORP, STARD, and Lam families, as well as NPC1, Protrudin, and PDZD8—also function at these sites[1-3]. The tethering of ER–LE/Ly MCSs is strengthened during endosomal maturation, a process involving both endosomal acidification and Ca^{2+} dynamics. Cholesterol and Ca^{2+} therefore serve as key messengers mediating cross-talk between these two organelles at ER–LE/Ly MCSs[4].

Cells acquire cholesterol through two main pathways: de novo synthesis and endocytosis of lipoproteins (LPs)[5-7]. In the brain, however, neurons lack the ability to synthesize cholesterol and instead rely on supply from glial cells. Cholesterol produced in glia is esterified to cholesteryl esters (CEs), packed into HDL-like lipoproteins, transported to neurons, internalized via LDLR-mediated endocytosis, and hydrolyzed to free cholesterol by lysosomal acidic lipases[7,8]. This cholesterol is then trafficked to the ER and redistributed to other organelles for cellular functions[4,9].

CEs stored in lipid droplets (LDs) are degraded through lipophagy, a form of microautophagy in which LDs fuse with lysosomes, enabling lipids to be hydrolyzed by lysosomal acidic lipases[10-13]. Endocytosed LP particles are degraded in a similar manner.

Thus, endosomal maturation, which depends on lysosomal acidification, is essential for
metabolizing CEs from both LDs and LPs into cholesterol.

PDZD8 is an ER membrane protein originally identified as a Protrudin-binding partner,
with both proteins acting as tethers at ER–LE/Ly MCSs[14-20]. PDZD8 belongs to the
TULIP super family and mediates cholesterol transport through its SMP domain[16,21,22].
Because Protrudin interacts with VAP via its FFAT motif, the PDZD8–Protrudin–VAP
complex is thought to function in lipid transport[3]. PDZD8 thereby contributes to
cholesterol metabolism by promoting lipolysis through endosomal maturation in
neurons[23]. Consistently, PDZD8-deficient mice exhibit CE accumulation in the brain due
to impaired lipolysis, along with memory and emotional dysfunction[24,25]. CE
accumulation is also observed in the brains of Alzheimer's disease (AD) patients and AD
model mice[26-28]. In particular, neurons carrying the APOE4 allele—the strongest genetic
risk factor for AD—show CE accumulation with lipofuscin formation due to impaired
lysosomal function[29,30].

Compared with cholesterol metabolism, research on lysosomal calcium dynamics is
relatively recent, and many questions remain unresolved. As endosomes undergo luminal
acidification, their Ca^{2+} concentration increases, and lysosomes eventually become a
major intracellular Ca^{2+} store alongside the ER and mitochondria. Ca^{2+} uptake into the
endosomal lumen promotes intraluminal vesicle fusion, thereby facilitating endosomal
maturation[31-34]. Conversely, calcium-induced calcium release (CICR)—in which a small
amount of Ca^{2+} efflux from lysosomes trigger large scale release from the ER IP_3
receptors and ryanodine receptors—has emerged as an important mechanism of transient
Ca^{2+} signaling[35-39]. Endosomal acidification activates lysosomal Ca^{2+} channels, such as
two-pore channel-2 (TPC2) and TRPML1, which promote Ca^{2+} release from lysosomes.
Although Ca^{2+} dynamics at ER–LE/Ly MCSs have been implicated in diverse cellular
processes[34,40], their detailed mechanisms remain poorly understood.

TMEM55B has been reported to perform multiple functions. It was first identified as a
phosphatase that hydrolyzes the 4-position phosphate of $PtdIns(4,5)P_2$ to generate
$PtdIns(5)P$[41,42]. TMEM55B is a lysosomal membrane protein that interacts with the v-
ATPase and Ragulator complexes, thereby contributing to v-ATPase activation and
subsequent mTORC1 signaling at the lysosomal membrane[43]. Consistent with its
importance, TMEM55B knockout mice are embryonic lethal[43]. Because v-ATPase is a
proton pump essential for endolysosomal acidification, its dysfunction leads to various
lysosomal storage diseases[44-46]. TMEM55B also cooperates with TFEB and JIP4 to
promote dynein-dependent retrograde lysosomal transport[47] and participate in lysosomal

oxidative stress responses[48]. In addition, TMEM55B regulates the expression and activity

of LDLR thereby modulating lipoprotein uptake and cholesterol metabolism[49,50].

In this study, we identified TMEM55B as a PDZD8-interacting lysosomal protein that

regulates cholesterol metabolism via lysosomal acidification and Ca^{2+} dynamics at ER–

LE/Ly MCSs.

Results

PDZD8 interacts with the lysosomal protein TMEM55B.

In neuronal cells, the lipid transfer protein PDZD8 promotes cholesterol metabolism via

endosomal maturation; however, the mechanism by which this ER protein regulates

lysosomal activity remains unclear. To identify lysosomal proteins interacting with

PDZD8, we established a Neuro2A cell line stably expressing 3×FLAG-tagged PDZD8.

The PDZD8 complex was isolated by anti-FLAG immunoprecipitation, separated by

SDS-PAGE, digested with trypsin, and analyzed by LC-MS/MS. In parallel, the same

analysis was performed using cells transfected with empty vector to exclude nonspecific

binding. Proteins reproducibly detected across three independent experiments were

extracted, and semiquantitative analysis based on identification frequency (IF)[51]

identified 196 proteins with IF > 0.01. Among those, 13 endosome-localized proteins

were selected using GO Term and UniProt subcellular localization annotations (Fig. 1A).

This group included the lysosomal protein TMEM55B and its binding partners, v-ATPase

subunit Atp6v0d1 and Lamtor1 (Fig. 1A)[43]. Notably, Lamtor1 and TMEM55B are

lysosome-localized proteins associated with cholesterol metabolism, suggesting a

potential role in lipolysis-mediated cholesterol processing. Since TMEM55B regulates

the v-ATPase complex, we focused on TMEM55B as a possible upstream regulator of

PDZD8-mediated lysosomal function.

TMEM55B is not essential for ER–LE/Ly tethering.

Because PDZD8 functions as a tether at ER–LE/Ly MCSs, we examined whether its

binding partner TMEM55B also contributes to this process. TMEM55B expression was

suppressed in HeLa cells using siRNA (siTMEM55B), resulting in a marked reduction of

both RNA and protein levels compared with control cells (siControl) (Fig. 1B, C). To

assess ER–LE/Ly contacts, we employed a split-GFP system in which GFP fluorescence

is reconstituted only when ER and lysosomal membranes are closely apposed[16,52]. Cells

were co-transfected with ERj1-GFP(1–10) (ER-localized fragment) and LAMP1-

GFP(11) (lysosomal fragment), together with mCherry to normalize expression levels.

The GFP/mCherry fluorescence ratio per cell, reflecting ER–LE/Ly MCS formation, was

quantified (Fig. 1D). No significant difference in GFP/mCherry ratio was observed
between siTMEM55B and siControl cells (Fig. 1E). These results demonstrate that,
unlike PDZD8, TMEM55B is not required for tethering of ER–LE/Ly MCSs.
127
TMEM55B regulates lysosomal acidification and activity.
TMEM55B interacts with the v-ATPase complex and has been implicated in its regulation.
We therefore hypothesized that TMEM55B contributes to lysosomal acidification and
activity via v-ATPase. To test this, HeLa cells transfected with siControl or siTMEM55B
were stained with LysoTracker (Fig. 2A) or LysoSensor (Fig. 2C), both of which increase
in fluorescence intensity as lysosomal pH decreases. In both assays, siTMEM55B cells
showed significantly reduced fluorescence compared with controls, indicating impaired
lysosomal acidification (Fig. 2B, D). To assess the functional consequences in neurons,
we suppressed TMEM55B expression in PC12 cells (Fig. 2E) and evaluated lipolysis
using EGFP-PLIN2 as a lipid droplet (LD) marker and LysoTracker Red as a lysosomal
marker (Fig. 2F). TMEM55B knockdown cells exhibited enlarged LDs (Fig. 2G) with
increased LD fluorescence intensity per cell (Fig. 2H), as well as reduced LysoTracker
fluorescence (Fig. 2I), consistent with defective lysosomal acidification and impaired
lipolysis. Furthermore, LD–lysosome fusion events were significantly decreased in
siTMEM55B cells compared with controls (Fig. 2J). Together, these results demonstrate
that TMEM55B promotes lysosomal acidification and activity, thereby facilitating LD
degradation and lipolysis.
145
TMEM55B promotes Ca^{2+} release from the ER.
Intracellular Ca^{2+} elevation promotes lysosome fusion with other organelles[44], suggesting
that lipolysis mediated by the PDZD8–TMEM55B complex could involve Ca^{2+} signaling
at ER–LE/Ly MCSs. Since the ER is the major intracellular Ca^{2+} store, we investigated
whether TMEM55B regulates Ca^{2+} release or uptake from the ER. HeLa cells were
transfected with the ER-targeted Ca^{2+} indicator G-CEPIA1er[53], which binds ER luminal
Ca^{2+} and emits GFP fluorescence, and colocalization with the ER marker KDEL was
confirmed by confocal microscopy (Fig. 3A, B). To assess ER Ca^{2+} dynamics, siControl
or siTMEM55B cells were stimulated with ATP to induce transient ER Ca^{2+} release,
followed by Ca^{2+} reuptake to the ER. In siTMEM55B cells, ATP-induced Ca^{2+} release as
well as reuptake were markedly reduced compared with controls (Fig. 3C, D).
Quantitative analysis revealed that siTMEM55B cells exhibited significantly decreased
rates of ER Ca^{2+} release (Fig. 3E), slower subsequent Ca^{2+} reuptake (Fig. 3F), and delayed

time to peak Ca²⁺ release (Fig. 3G). These findings indicate that TMEM55B facilitates

ATP-stimulated transient Ca²⁺ release from the ER.

TMEM55B promotes Ca²⁺ release from lysosomes.

We next investigated whether TMEM55B regulates Ca²⁺ release from lysosomes and how

this affects ER Ca²⁺ dynamics. Lysosomal Ca²⁺ release and uptake were monitored using

Ly-GG, which binds cytosolic Ca²⁺ at the lysosomal membrane and emits GFP

fluorescence[54] (Fig. 4A). Co-expression of Ly-GG with the lysosomal marker LAMP1

confirmed colocalization (Fig. 4B). ATP stimulation of siControl HeLa cells induced

transient Ca²⁺ release from lysosomes. In siTMEM55B cells, the magnitude (Fig. 4C) and

rate (Fig. 4D) of lysosomal Ca²⁺ release were significantly reduced compared to controls.

The subsequent Ca²⁺ reuptake rate was also decreased (Fig. 4E), indicating impaired

lysosomal Ca²⁺ dynamics. As a control, we examined mitochondrial Ca²⁺ using

CEPIA3mt[53], which binds mitochondrial matrix Ca²⁺, together with the mitochondrial

marker Su9-RFP[55] (Fig. 4F), confirmed colocalization (Fig. 4G). ATP stimulation

induced transient mitochondrial Ca²⁺ increases as well as decrease, which were not

significantly different between siControl and siTMEM55B cells (Fig. 4H–J). These

results indicate that TMEM55B specifically regulates lysosomal Ca²⁺ dynamics without

affecting mitochondrial Ca²⁺.

Discussion

Lysosomes serve as central hubs for cellular metabolism, coordinating degradation,

nutrient recycling, and signaling. Proper lysosomal acidification and calcium homeostasis

are essential for lipid metabolism and neuronal function, and their dysregulation has been

increasingly linked to neurodegenerative diseases. Despite this, the molecular

mechanisms connecting lysosomal function to lipid regulation remain incompletely

understood.

In this study, we identify TMEM55B as a key lysosomal regulator that interacts with the

ER protein PDZD8 to control lysosomal acidification and calcium dynamics. TMEM55B

promotes endosomal maturation and lipid droplet degradation, revealing a novel

mechanism linking lysosomal activity to lipid metabolism. These findings provide new

insights into cellular lipid homeostasis and suggest potential targets for disorders

associated with lysosomal dysfunction and cholesterol metabolism.

The lipid transfer protein PDZD8 mediates tethering of ER–LE/Ly membrane contact

sites (MCSs) and facilitates cholesterol metabolism through lipolysis via endosomal

maturation. However, the mechanism by which PDZD8 promotes endosome maturation

has remained unclear. Through complex analysis in neuronal cells, we identified Lamtor1
and TMEM55B as lysosome-localized, cholesterol-associated proteins among LE/Ly
candidates. Given that TMEM55B regulates the v-ATPase complex, we focused on its
intracellular functions.

We found that TMEM55B contributes to lysosomal acidification and activation. Loss of
TMEM55B impaired lysosomal Ca^{2+} release and uptake, as well as ATP-stimulated Ca^{2+}
release from the ER, while mitochondrial Ca^{2+} dynamics remained unaffected. In our
system, lysosomal Ca^{2+} release peaked within 180 s after ATP stimulation (Fig. 4C),
whereas ER Ca^{2+} release peaked later (Fig. 3C), consistent with calcium-induced calcium
release (CICR) from the ER triggered by lysosomal Ca^{2+} release. These results indicate
that TMEM55B promotes endosomal maturation via lysosomal acidification and
activation, and facilitates CICR at ER–LE/Ly MCSs, thereby supporting lipolysis (Fig.
5).

The PDZD8–TMEM55B complex thus plays a central role in ER–LE/Ly MCS functions,
including endosomal maturation, Ca^{2+} dynamics, and cholesterol metabolism.
Nevertheless, the effects of TMEM55B deficiency on lysosomal fusion and lipid droplet
degradation were partial, failing to recapitulate the strong phenotype observed with
PDZD8 deficiency. This suggests that additional, as-yet-unidentified mechanisms
contribute to PDZD8-mediated lipolysis.

In summary, TMEM55B enhances lysosomal acidification via v-ATPase regulation,
thereby increasing lysosomal enzyme activity and promoting lipolysis. Lysosomal
acidification also activates Ca^{2+} channels, facilitating CICR at ER–LE/Ly MCSs and
supporting cholesterol metabolism through lysosomal fusion. Direct evidence linking
TMEM55B-mediated CICR to enhanced lipolysis, however, remains lacking, and further
studies are needed to clarify this mechanism.

220
221

Methods
Plasmids
EGFP and mCherry vectors were from Clontech (Mountain View, CA, US). Human
cDNAs encoding PLIN2 (GenBank accession number NM_007408.4), LAMP1
(GenBank accession number BC025335), and KDEL with the signal sequence of
calreticulin (GenBank accession number NM_004343) were amplified from HeLa cells
by the polymerase chain reaction (PCR) with PrimeSTAR HS DNA polymerase (Takara,
Shiga, Japan). pCMV-G-CEPIA1er (Addgene plasmid #58215) and pCMV-CEPIA3mt
(Addgene plasmid #58219) were obtained from Addgene (Watertown, MA, US). Ly-GG

was kindly provided by Dr. CW Taylor (University of Cambridge, UK). Vectors for split-
GFP analysis, ERj1–GFP(1–10) and LAMP1–GFP(11), were kindly provided by Dr.
Tamura (Yamagata University, Japan). The pSu9 cDNA was kindly provided by Dr.
Mihara (Kyushu University, Japan). The mRFP cDNA was kindly provided by Dr. R.Y.
Tsien (University of California, San Diego, US).
236

237 **Antibodies**
238 Antibodies to TMEM55B were from Proteintech (Rosemont, IL, USA); antibodies to
239 HSP90 were from BD Biosciences (Franklin Lakes, NJ, USA).
240

Cell culture, transfection, and retrovirus infection
HeLa cells (CCL-2, purchased from ATCC, US) were cultured under a humidified
atmosphere of 5% CO_2 at 37°C in Dulbecco's modified Eagle's medium (Invitrogen Life
Technologies) supplemented with 10% fetal bovine serum (Invitrogen Life Technologies).
The culture medium for Plat-E cells was also supplemented with blasticidin (10 μg/ml).
Cells were transfected with the use of XtremeGene9 reagents (Roche Diagnostics,
Mannheim, Germany) according to the manufacturer's protocol. For retroviral infection,
Plat-E cells were transiently transfected with pMX-puro–based vectors and then cultured
for 48 h. The retroviruses in the resulting culture supernatants were used to infect
Neuro2A cells, and the cells were then subjected to selection with puromycin (1 μg/ml).
251

Identification and categorization of PDZD8-associated proteins
Neuro2a cells infected with a retrovirus encoding FLAG–tagged mouse PDZD8 or with
the corresponding empty virus were solubilized in a lysis buffer (40 mM HEPES-NaOH
[pH 7.5], 150 mM NaCl, 10% glycerol, 0.5% Triton X-100, 1 mM Na_3VO_4, 25 mM NaF,
aprotinin [10 μg/ml], leupeptin [10 μg/ml], 1 mM phenylmethylsulfonyl fluoride) to
prepare whole-cell extracts. Lysates were cleared by centrifugation at 20,400 × g for 15
min at 4 °C. The clarified supernatants were subjected to affinity purification with anti-
FLAG (M2)–agarose affinity gel (Sigma-Aldrich), and then proteins eluted with FLAG
peptide (Sigma-Aldrich) was separated by SDS-PAGE and stained with silver. The
stained gels were sliced into pieces, and the abundant proteins therein were subjected to
in-gel digestion with trypsin. The resulting peptides were dried, dissolved in a mixture of
0.1% trifluoroacetic acid and 2% acetonitrile, and then applied to a nanoflow LC system
(Advanced UHPLC; Michrom BioResources, Auburn, CA, USA) equipped with an L-
column (C18, 0.15 by 50 mm, particle size of 3 μm; CERI, Tokyo, Japan). LC–MS/MS
analysis was performed with an Orbitrap Velos Pro instrument (Thermo Fisher Scientific).

All MS/MS spectra were compared with protein sequences in the International Protein

Index (IPI, European Bioinformatics Institute) mouse version 3.44 with the use of the

MASCOT algorithm. Assigned high-scoring peptide sequences were considered for

correct identification. Identified peptides from 3 independent experiments were integrated

and regrouped by IPI accession number. Proteins detected in ≥2 of 3 independent

experiments from FLAG-PDZD8–expressing cells, but absent from control cells, were

considered PDZD8-associated. Semi-quantitative abundance was estimated from

identification frequency (IF; number of identified peptides per protein normalized to the

number of theoretically detectable tryptic peptides). Functional categorization of

PDZD8-associated proteins was categorized with use of the UniProtKB database

(https://www.uniprot.org) based on Gene Ontology (GO) annotations. The proteins listed

in figure 1a shows endosome-related (GO:0005768), as well as columns for lysosome

(GO:0005764) or cholesterol-related as well.

RNA interference, reverse transcription, and real-time PCR analysis

Stealth siRNAs targeted to human TMEM55B (siTMEM55B#1, 5'-

ACGUGGAAGGCAAGAUGCAUCAGCA-3'; siTMEM55B#2, 5'-

CAAGAUGCAUCAGCAUGUAGUCAAA-3'), rat TMEM55B (siTMEM55B#1, 5'-

UGCUGAUGCAUCUUGCCUUCCACGU-3'; siTMEM55B#2, 5'-

CAAGAUGCAUCAGCAUGUAGUCAAA-3'), and non-targeting control were obtained

from Invitrogen Life Technologies. (Invitrogen Life Technologies, Carlsbad, CA, USA).

Human siRNAs were introduced into HeLa cells with the use of Lipofectamine

RNAiMAX (Invitrogen Life Technologies), and rat siRNAs were introduced into PC12

cells with the use of a Cell Line Nucleofector Kit V (VCA-1003) and Nucleofector

instrument (Lonza, Basel, Switzerland) according to the manufacturer's protocol. The

cells were then cultured for 48 h (for HeLa) or 72 h (for PC12). Total RNA was isolated

from cells with the use of an RNeasy Mini Kit (Qiagen, Hilden, Germany) according to

the manufacturer's protocol and was subjected to reverse transcription (RT) with the use

of a QuantiTect Rev Transcription Kit (Qiagen). The resulting cDNA was subjected to

real-time PCR analysis with Power SYBR Green PCR Master Mix in an ABI-Prism 7000

sequence detection system (Applied Biosystems, Foster City, CA). The amounts of each

target mRNA were calculated and normalized by that of GAPDH mRNA. Primers for

PCR (forward and reverse, respectively) were as follows: 5'-

GTCAGTGGTGGACCTGACCTG-3' and 5'-AAAGTGGTCGTTGAGGGCAAT-3' for

human GAPDH; 5'-CGAGTCTGCCAATCTCTCATC-3' and 5'-

AGTGCGGTCTGTGAACTCTGT-3' for human TMEM55B; 5'-

AAGGCTGAGAATGGGAAGCTG-3' and 5'-GGAGATGATGACCCTTTTGGC-3' for
rat GAPDH; 5'-CTAGCCCGGATAGTGGGAGTG-3' and 5'-
TTCCTGGAGGTGCATTCTTGA-3' for rat TMEM55B.
306

Fluorescence imaging of live cells

LysoTracker Red DND-99 (0.2 µM) or LysoSensor Green DND-189 (0.2 µM) (Thermo
Fisher Scientific, Rockford, IL, USA) were added to cells for 1 h at 37 °C according to
the manufacturer's protocol. HeLa and PC12 cells expressing plasmid-encoded
fluorescent proteins for 24 h and 96 h respectively, or metabolically labeled with
fluorescent probes were observed with an LSM800 confocal microscope (Zeiss,
Oberkochen, Germany), and the images were processed for calculation of fluorescence
intensity with ZEN imaging software (Zeiss) as well as Fiji (ImageJ, https://fiji.sc).
315

Ca^{2+} imaging

Cell media of HeLa cells expressing organelle-targeted Ca^{2+} indicators were pre-
incubated with HEPES-buffered saline (HBS; 137 mM NaCl, 5.9 mM KCl, 2.2 mM CaCl$_2$,
1.2 mM MgCl$_2$, 10 mM HEPES, 14 mM glucose, pH 7.4) for 1 h at 37 °C and then
stimulated with ATP (200 µM) (Wako, Japan). The images were captured by Operetta
high-content screening system (PerkinElmer, Waltham, MA, USA) using a ×20 objective
(λ_ex 460–490 nm; λ_em 500–550 nm) at the time point of ATP addition (t = 0 s) and
continued time points (t = X s) at a rate of one frame per 20 sec for total 1240 sec (G-
CEPIA) or 180 sec (Ly-GG and CEPIA3mt). The concentration of Ca^{2+} for each time
points was calculated by $\Delta F/F_0 = (F_x - F_0)/F_0$. Images were analyzed by Harmony
software (PerkinElmer).
327

Immunoblot analysis

HeLa cells were lysed in Triton lysis buffer containing 40 mM HEPES-NaOH (pH 7.6),
150 mM NaCl, 0.5% Triton X-100, 10 mM MgCl$_2$, 1 mM Na$_3$VO$_4$, 25 mM NaF, 1 mM
EDTA, 1 mM PMSF, leupeptin (10 µg/mL), and 10% glycerol. After 10 min on ice,
lysates were centrifuged at 15,000 × g for 10 min at 4 °C, and protein concentrations were
determined with the Pierce BCA Protein Assay (Thermo Fisher Scientific). Extracts were
resolved by SDS–PAGE, transferred to Immobilon-P membranes (Merck Millipore,
Darmstadt, Germany), and probed with primary antibodies. Signals were developed with
SuperSignal reagents (Thermo Fisher Scientific) and imaged on an LAS-4000 (GE
Healthcare, Waukesha, WI, USA).
338

Statistical analysis

Statistical analyses were performed with BellCurve for Excel software (Social Survey Research Information, Tokyo, Japan). Normality of data was assessed with the Shapiro–Wilk test. If the normality assumption was not met, the Kruskal–Wallis test with the Steel's multiple comparisons test were applied. If data were normally distributed, One-way repeated-measures analysis of variance (ANOVA) with the Dunnett's multiple comparisons test were performed. Quantitative data are presented as mean ± SE. Statistical significance of a P value is indicated as $0.01 \leq p < 0.05$ (*); $0.001 \leq p < 0.01$ (**); $p < 0.001$ (***); $p \geq 0.05$ (n.s.). Sample size (n) are in the figure legends.

References

Luo, J., Jiang, L. Y., Yang, H. & Song, B. L. Intracellular Cholesterol Transport by Sterol Transfer Proteins at Membrane Contact Sites. *Trends Biochem Sci* **44**, 273-292, doi:10.1016/j.tibs.2018.10.001 (2019).

Wong, L. H., Gatta, A. T. & Levine, T. P. Lipid transfer proteins: the lipid commute via shuttles, bridges and tubes. *Nat Rev Mol Cell Biol* **20**, 85-101, doi:10.1038/s41580-018-0071-5 (2019).

Shirane, M. Lipid Transfer-Dependent Endosome Maturation Mediated by Protrudin and PDZD8 in Neurons. *Front Cell Dev Biol* **8**, 615600, doi:10.3389/fcell.2020.615600 (2020).

van der Kant, R. & Neefjes, J. Small regulators, major consequences - Ca(2)(+) and cholesterol at the endosome-ER interface. *J Cell Sci* **127**, 929-938, doi:10.1242/jcs.137539 (2014).

Ikonen, E. Cellular cholesterol trafficking and compartmentalization. *Nat Rev Mol Cell Biol* **9**, 125-138, doi:10.1038/nrm2336 (2008).

Goldstein, J. L. & Brown, M. S. A century of cholesterol and coronaries: from plaques to genes to statins. *Cell* **161**, 161-172, doi:10.1016/j.cell.2015.01.036 (2015).

Luo, J., Yang, H. & Song, B. L. Mechanisms and regulation of cholesterol homeostasis. *Nat Rev Mol Cell Biol* **21**, 225-245, doi:10.1038/s41580-019-0190-7 (2020).

Mehta, N. *et al.* The evolving role of cholesteryl ester transfer protein inhibition beyond cardiovascular disease. *Pharmacol Res* **197**, 106972, doi:10.1016/j.phrs.2023.106972 (2023).

9   Martin, M. G., Pfrieger, F. & Dotti, C. G. Cholesterol in brain disease:
sometimes determinant and frequently implicated. *EMBO Rep* **15**, 1036-1052,
doi:10.15252/embr.201439225 (2014).

10   Zechner, R., Madeo, F. & Kratky, D. Cytosolic lipolysis and lipophagy: two
sides of the same coin. *Nat Rev Mol Cell Biol* **18**, 671-684,
doi:10.1038/nrm.2017.76 (2017).

11   Schuck, S. Microautophagy - distinct molecular mechanisms handle cargoes of
many sizes. *J Cell Sci* **133**, doi:10.1242/jcs.246322 (2020).

12   Haidar, M., Loix, M., Bogie, J. F. J. & Hendriks, J. J. A. Lipophagy: a new
player in CNS disorders. *Trends Endocrinol Metab* **32**, 941-951,
doi:10.1016/j.tem.2021.08.010 (2021).

13   Lan, Z. Q., Ge, Z. Y., Lv, S. K., Zhao, B. & Li, C. X. The regulatory role of
lipophagy in central nervous system diseases. *Cell Death Discov* **9**, 229,
doi:10.1038/s41420-023-01504-z (2023).

14   Shirane, M. & Nakayama, K. I. Protrudin induces neurite formation by
directional membrane trafficking. *Science* **314**, 818-821,
doi:10.1126/science.1134027 (2006).

15   Guillen-Samander, A., Bian, X. & De Camilli, P. PDZD8 mediates a Rab7-
dependent interaction of the ER with late endosomes and lysosomes. *Proc Natl*
Acad Sci U S A **116**, 22619-22623, doi:10.1073/pnas.1913509116 (2019).

16   Shirane, M. *et al.* Protrudin and PDZD8 contribute to neuronal integrity by
promoting lipid extraction required for endosome maturation. *Nat Commun* **11**,
4576, doi:10.1038/s41467-020-18413-9 (2020).

17   Elbaz-Alon, Y. *et al.* PDZD8 interacts with Protrudin and Rab7 at ER-late
endosome membrane contact sites associated with mitochondria. *Nat Commun*
11, 3645, doi:10.1038/s41467-020-17451-7 (2020).

18   Khan, H., Chen, L., Tan, L. & Im, Y. J. Structural basis of human PDZD8-Rab7
interaction for the ER-late endosome tethering. *Sci Rep* **11**, 18859,
doi:10.1038/s41598-021-98419-5 (2021).

19   Gao, Y., Xiong, J., Chu, Q. Z. & Ji, W. K. PDZD8-mediated lipid transfer at
contacts between the ER and late endosomes/lysosomes is required for neurite
outgrowth. *J Cell Sci* **135**, doi:10.1242/jcs.255026 (2022).

20   Thakur, R. S. & O'Connor-Giles, K. M. PDZD8 promotes autophagy at ER-
lysosome membrane contact sites to regulate activity-dependent synaptic
growth. *Cell Rep* **44**, 115483, doi:10.1016/j.celrep.2025.115483 (2025).

21   Alva, V. & Lupas, A. N. The TULIP superfamily of eukaryotic lipid-binding
proteins as a mediator of lipid sensing and transport. *Biochim Biophys Acta*
1861, 913-923, doi:10.1016/j.bbalip.2016.01.016 (2016).
22   Wong, L. H. & Levine, T. P. Tubular lipid binding proteins (TULIPs) growing
everywhere. *Biochim Biophys Acta Mol Cell Res* **1864**, 1439-1449,
doi:10.1016/j.bbamcr.2017.05.019 (2017).
23   Morita, K. *et al.* PDZD8-deficient mice accumulate cholesteryl esters in the
brain as a result of impaired lipophagy. *iScience* **25**, 105612,
doi:10.1016/j.isci.2022.105612 (2022).
24   Al-Amri, A. H. *et al.* PDZD8 Disruption Causes Cognitive Impairment in
Humans, Mice, and Fruit Flies. *Biol Psychiatry* **92**, 323-334,
doi:10.1016/j.biopsych.2021.12.017 (2022).
25   Kurihara, Y. *et al.* PDZD8-deficient mice manifest behavioral abnormalities
related to emotion, cognition, and adaptation due to dyslipidemia in the brain.
Mol Brain **16**, 11, doi:10.1186/s13041-023-01002-4 (2023).
26   Chan, R. B. *et al.* Comparative lipidomic analysis of mouse and human brain
with Alzheimer disease. *J Biol Chem* **287**, 2678-2688,
doi:10.1074/jbc.M111.274142 (2012).
27   van der Kant, R. *et al.* Cholesterol Metabolism Is a Druggable Axis that
Independently Regulates Tau and Amyloid-beta in iPSC-Derived Alzheimer's
Disease Neurons. *Cell Stem Cell* **24**, 363-375 e369,
doi:10.1016/j.stem.2018.12.013 (2019).
28   Nugent, A. A. *et al.* TREM2 Regulates Microglial Cholesterol Metabolism upon
Chronic Phagocytic Challenge. *Neuron* **105**, 837-854 e839,
doi:10.1016/j.neuron.2019.12.007 (2020).
29   Blanchard, J. W. *et al.* APOE4 impairs myelination via cholesterol dysregulation
in oligodendrocytes. *Nature* **611**, 769-779, doi:10.1038/s41586-022-05439-w
(2022).
30   Guo, J. L. *et al.* Decreased lipidated ApoE-receptor interactions confer
protection against pathogenicity of ApoE and its lipid cargoes in lysosomes. *Cell*
188, 187-206 e126, doi:10.1016/j.cell.2024.10.027 (2025).
31   Mayran, N., Parton, R. G. & Gruenberg, J. Annexin II regulates multivesicular
endosome biogenesis in the degradation pathway of animal cells. *EMBO J* **22**,
3242-3253, doi:10.1093/emboj/cdg321 (2003).

32   Lloyd-Evans, E. *et al.* Niemann-Pick disease type C1 is a sphingosine storage
disease that causes deregulation of lysosomal calcium. *Nat Med* **14**, 1247-1255,
doi:10.1038/nm.1876 (2008).

33   Bissig, C. *et al.* Viral infection controlled by a calcium-dependent lipid-binding
module in ALIX. *Dev Cell* **25**, 364-373, doi:10.1016/j.devcel.2013.04.003
(2013).

34   Bretou, M. *et al.* Accumulation of APP C-terminal fragments causes
endolysosomal dysfunction through the dysregulation of late endosome to
lysosome-ER contact sites. *Dev Cell* **59**, 1571-1592 e1579,
doi:10.1016/j.devcel.2024.03.030 (2024).

35   Churchill, G. C. *et al.* NAADP mobilizes Ca(2+) from reserve granules,
lysosome-related organelles, in sea urchin eggs. *Cell* **111**, 703-708,
doi:10.1016/s0092-8674(02)01082-6 (2002).

36   Calcraft, P. J. *et al.* NAADP mobilizes calcium from acidic organelles through
two-pore channels. *Nature* **459**, 596-600, doi:10.1038/nature08030 (2009).

37   Galione, A. A primer of NAADP-mediated Ca(2+) signalling: From sea urchin
eggs to mammalian cells. *Cell Calcium* **58**, 27-47,
doi:10.1016/j.ceca.2014.09.010 (2015).

38   Kilpatrick, B. S. *et al.* An Endosomal NAADP-Sensitive Two-Pore Ca(2+)
Channel Regulates ER-Endosome Membrane Contact Sites to Control Growth
Factor Signaling. *Cell Rep* **18**, 1636-1645, doi:10.1016/j.celrep.2017.01.052
(2017).

39   Yuan, Y. *et al.* Two-pore channel-2 and inositol trisphosphate receptors
coordinate Ca(2+) signals between lysosomes and the endoplasmic reticulum.
Cell Rep **43**, 113628, doi:10.1016/j.celrep.2023.113628 (2024).

40   Zheng, Q. *et al.* Calcium transients on the ER surface trigger liquid-liquid phase
separation of FIP200 to specify autophagosome initiation sites. *Cell* **185**, 4082-
4098 e4022, doi:10.1016/j.cell.2022.09.001 (2022).

41   Ungewickell, A. *et al.* The identification and characterization of two
phosphatidylinositol-4,5-bisphosphate 4-phosphatases. *Proc Natl Acad Sci U S*
A **102**, 18854-18859, doi:10.1073/pnas.0509740102 (2005).

42   Zou, J., Marjanovic, J., Kisseleva, M. V., Wilson, M. & Majerus, P. W. Type I
phosphatidylinositol-4,5-bisphosphate 4-phosphatase regulates stress-induced
apoptosis. *Proc Natl Acad Sci U S A* **104**, 16834-16839,
doi:10.1073/pnas.0708189104 (2007).

43   Hashimoto, Y., Shirane, M. & Nakayama, K. I. TMEM55B contributes to
lysosomal homeostasis and amino acid-induced mTORC1 activation. *Genes*
Cells **23**, 418-434, doi:10.1111/gtc.12583 (2018).
44   Xu, H. & Ren, D. Lysosomal physiology. *Annu Rev Physiol* **77**, 57-80,
doi:10.1146/annurev-physiol-021014-071649 (2015).
45   Banerjee, S. & Kane, P. M. Regulation of V-ATPase Activity and Organelle pH
by Phosphatidylinositol Phosphate Lipids. *Front Cell Dev Biol* **8**, 510,
doi:10.3389/fcell.2020.00510 (2020).
46   Ballabio, A. & Bonifacino, J. S. Lysosomes as dynamic regulators of cell and
organismal homeostasis. *Nat Rev Mol Cell Biol* **21**, 101-118,
doi:10.1038/s41580-019-0185-4 (2020).
47   Willett, R. *et al.* TFEB regulates lysosomal positioning by modulating
TMEM55B expression and JIP4 recruitment to lysosomes. *Nat Commun* **8**,
1580, doi:10.1038/s41467-017-01871-z (2017).
48   Jeong, E., Willett, R., Rissone, A., La Spina, M. & Puertollano, R. TMEM55B
links autophagy flux, lysosomal repair, and TFE3 activation in response to
oxidative stress. *Nat Commun* **15**, 93, doi:10.1038/s41467-023-44316-6 (2024).
49   Medina, M. W. *et al.* Transmembrane protein 55B is a novel regulator of cellular
cholesterol metabolism. *Arterioscler Thromb Vasc Biol* **34**, 1917-1923,
doi:10.1161/ATVBAHA.113.302806 (2014).
50   Qin, Y. *et al.* Phosphatidylinositol-(4,5)-Bisphosphate Regulates Plasma
Cholesterol Through LDL (Low-Density Lipoprotein) Receptor Lysosomal
Degradation. *Arterioscler Thromb Vasc Biol* **40**, 1311-1324,
doi:10.1161/ATVBAHA.120.314033 (2020).
51   Matsumoto, M. *et al.* Large-scale proteomic analysis of tyrosine-
phosphorylation induced by T-cell receptor or B-cell receptor activation reveals
new signaling pathways. *Proteomics* **9**, 3549-3563,
doi:10.1002/pmic.200900011 (2009).
52   Kakimoto, Y. *et al.* Visualizing multiple inter-organelle contact sites using the
organelle-targeted split-GFP system. *Sci Rep* **8**, 6175, doi:10.1038/s41598-018-
24466-0 (2018).
53   Suzuki, J. *et al.* Imaging intraorganellar Ca2+ at subcellular resolution using
CEPIA. *Nat Commun* **5**, 4153, doi:10.1038/ncomms5153 (2014).
54   Atakpa, P., Thillaiappan, N. B., Mataragka, S., Prole, D. L. & Taylor, C. W.
IP(3) Receptors Preferentially Associate with ER-Lysosome Contact Sites and

Selectively Deliver Ca(2+) to Lysosomes. *Cell Rep* **25**, 3180-3193 e3187,
doi:10.1016/j.celrep.2018.11.064 (2018).
55 Otera, H. *et al.* A novel insertion pathway of mitochondrial outer membrane
proteins with multiple transmembrane segments. *J Cell Biol* **179**, 1355-1363,
doi:10.1083/jcb.200702143 (2007).
517

Acknowledgements and Funding Information
We thank Y. Tamura for split-GFP cDNAs; E. Taylor for Ly-GG cDNA; K. Mihara for
Su9 cDNA; K. I. Nakayama and T. Ohnishi for proteome analysis; K. Sato and K. Kato
for Ca^{2+} imaging analysis; and the Research Equipment Sharing Center at Nagoya City
University for assistance. This work was supported in part by KAKENHI grants from
Japan Society for the Promotion of Science (JSPS) to M.S. (24K02122).
524

525 **Author contributions statement**
526 M.S. designed experiments, supervised the study, analyzed the data, and wrote the
527 manuscript. N.M. and H.M. performed experiments.
528

Additional information
Data availability
The MS data have been deposited in the JPOST Repository Database as follows:
URL: https://repository.jpostdb.org
JPID and PXID are as follows;
Mock-FLAG complex in Neuro2A; (1) JPST004039, PXD067776; (2) JPST004040,
PXD067777; (3) JPST004041, PXD067779.
PDZD8-FLAG complex in Neuro2A; (1) JPST004042, PXD067780; (2) JPST004043,
PXD067782; (3) JPST004044, PXD067783.
538

539 **Supplementary datasets**
540 Raw data is provided within the supplementary information files.
541

542 **Competing interests**
543 The authors declare no competing interests.
544

545 **Correspondence**
546 Correspondence and requests for materials should be addressed to M.S.
547 (shiram@phar.nagoya-cu.ac.jp)

548
549

Figure Legends

Fig. 1. Identification of PDZD8-associated proteins by proteomic analysis. (**A**) Endosome-related proteins identified as PDZD8-associated proteins. Protein abundance was estimated semiquantitatively by identification frequency (IF). Asterisks indicate proteins previously reported as TMEM55B-associated. Lysosome (Ly)-localized and cholesterol (Chol)-related proteins are also indicated. Proteins were categorized based on Gene Ontology (GO) terms and UniProtKB annotations. (**B, C**) HeLa cells were transfected with siControl, siTMEM55B#1, or siTMEM55B#2. Total RNA was analyzed by RT-qPCR for TMEM55B mRNA (B), and protein levels were examined by immunoblotting with anti-TMEM55B and anti-HSP90 antibodies (C). Data are shown as mean ± SE (n = 9 for RNA analysis). ***$p < 0.001$ (one-way ANOVA with Dunnett's multiple comparisons). (**D**) HeLa cells were transfected with siControl, siTMEM55B#1, or siTMEM55B#2 together with split-GFP vectors ERj1–GFP(1–10) and LAMP1–GFP(11), along with mCherry as a transfection control. Scale bar, 10 μm. (**E**) Quantification of split-GFP fluorescence normalized to mCherry intensity as in (D). n = 38, 36, and 37 cells for siControl, siTMEM55B#1, and siTMEM55B#2, respectively. n.s., not significant (Kruskal–Wallis test with Steel's multiple comparisons).

Fig. 2. TMEM55B depletion impairs lysosomal acidification and lipolysis. (**A, C**) HeLa cells transfected with siControl, siTMEM55B#1, or siTMEM55B#2 were stained with LysoTracker Red (A) or LysoSensor Green (C) and analyzed by confocal microscopy. Boxed areas (top panels) are shown at higher magnification (bottom panels). Scale bars, 10 μm (top and middle) and 5 μm (bottom). (**B, D**) Quantification of LysoTracker Red (B) or LysoSensor Green (D) mean fluorescence intensity per lysosome in (A) (n = 746, 497, and 452 lysosomes for siControl, siTMEM55B#1, and siTMEM55B#2, respectively) or (C) (n = 703, 847, and 794 lysosomes, respectively). (**E**) RT-qPCR analysis of TMEM55B mRNA in PC12 cells transfected with siControl, siTMEM55B#1, or siTMEM55B#2 (mean ± SE, n = 6). (**F**) PC12 cells were transfected with siControl, siTMEM55B#1, or siTMEM55B#2 together with EGFP-PLIN2, and stained with LysoTracker Red. (**G**) Quantification of lipid droplet (LD) size (μm²) of EGFP-PLIN2 puncta in (F) (n = 627, 450, and 317 LDs for siControl, siTMEM55B#1, and siTMEM55B#2, respectively). (**H**) Quantification of mean EGFP-PLIN2 fluorescence intensity per cell in (F) (n = 52, 41, and 35 cells for siControl, siTMEM55B#1, and siTMEM55B#2, respectively). (**I**) Quantification of mean LysoTracker Red fluorescence intensity per cell in (F) (n = 52, 41, and 35 cells, respectively). (**J**) Pearson's correlation coefficient between EGFP-PLIN2 and LysoTracker Red per cell in (F) (n = 52, 41, and

35 cells, respectively). ***p < 0.001, **p < 0.01 (Kruskal–Wallis test with Steel's
multiple comparisons test in B, D, and G–I); **p < 0.01, *p < 0.05 (one-way ANOVA
with Dunnett's multiple comparisons in E and J).

Fig. 3. Depletion of TMEM55B impairs Ca^{2+} release from the ER. (**A**) Schematic
representation of G-CEPIA1er, a Ca^{2+} indicator with the G-GECO1 domain facing the
ER lumen. (**B**) Confocal images of HeLa cells co-expressing G-CEPIA1er (green) and
the ER marker mCherry-KDEL (red). Scale bar, 10 μm. (**C**) Time lapse analysis of G-
CEPIA1er fluorescence ($\Delta F/F_0$) in cells transfected with siControl (gray, n = 30 fields),
siTMEM55B#1 (red, n = 30), or siTMEM55B#2 (pink, n = 30). ATP (200 μM) was added
at t = 0 s; Data are mean ± SE. (**D**) Quantification of peak $\Delta F/F_0$ values in (C).
(**E**) Quantification of the slope of G-CEPIA1er $\Delta F/F_0$ during 0–120 s after ATP addition
($\Delta F/F_0 \cdot s^{-1}$). (**F**) Quantification of time (s) to peak after ATP addition for G-CEPIA1er
$\Delta F/F_0$. (**G**) Quantification of the slope of G-CEPIA1er $\Delta F/F_0$ during 900 s after the peak
($\Delta F/F_0 \cdot s^{-1}$). Data are mean ± SE; ***p < 0.001, *p < 0.05 (Kruskal–Wallis test with
Steel's multiple comparisons in D, F, and G); ***p < 0.001 (one-way ANOVA with
Dunnett's multiple comparisons in E).

Fig. 4. Depletion of TMEM55B impairs lysosomal Ca^{2+} dynamics. (**A**) Schematic
representation of Ly-GG, a Ca^{2+} indicator with the G-GECO1 domain facing the cytosolic
side of the lysosomal membrane. (**B**) Confocal images of HeLa cells co-expressing Ly-
GG (green) and the lysosomal marker mCherry–LAMP1 (red). Scale bar, 10 μm. (**C**)
Time lapse analysis of Ly-GG fluorescence ($\Delta F/F_0$) in cells transfected with siControl
(gray, n = 29 fields), siTMEM55B#1 (red, n = 29), or siTMEM55B#2 (pink, n = 26). ATP
(200 μM) was added at t = 0 s; Data are mean ± SE and normalized by subtracting values
from ATP-free cells, in which fluorescence declined due to photobleaching. (**D, E**)
Quantification of Ly-GG slopes during 0–60 s (D) and 60–160 s (E) ($\Delta F/F_0 \cdot s^{-1}$). (**F**)
Schematic representation of CEPIA3mt, a Ca^{2+} indicator with the GCaMP2 domain
facing the mitochondrial matrix. (**G**) Confocal images of HeLa cells co-expressing
CEPIA3mt (green) and the mitochondrial marker Su9–RFP (red). Scale bar, 10 μm. (**H**)
Time lapse analysis of CEPIA3mt fluorescence ($\Delta F/F_0$) in cells transfected with control
siRNA (gray, n = 25 fields), siTMEM55B#1 (red, n = 26), or siTMEM55B#2 (pink, n =
24). ATP (200 μM) was added at t = 0 s; Data are mean ± SE. (**I, J**) Quantification of
CEPIA3mt slopes during 0–60 s (I) and 60–160 s (J) ($\Delta F/F_0 \cdot s^{-1}$). Data are mean ± SE;
***p < 0.001; (Kruskal–Wallis test with Steel's multiple comparisons in D and E); n.s.,
not significant (one-way ANOVA with Dunnett's multiple comparisons in I and J).

Fig. 5. Model of the mechanism where TMEM55B promotes lipolysis via lysosomal acidification and calcium regulation at ER-Ly MCSs. ER protein PDZD8 is associated with lysosome protein TMEM55B that promotes lysosomal acidification through regulation of v-ATPase. Lysosomal acidification then promotes activity of lysosomal acidic lipases that promote lipolysis and subsequently degradation of CEs to cholesterol, as well as calcium channels that promote CICR. CICR at ER-Ly MCSs may also contribute to lipolysis by fusion of LDs or LPs with lysosomes. ER, endoplasmic reticulum; MCS, membrane contact site; v-ATPase, vacuolar-ATPase; TPC2, two-pore channel 2; TRPML1, transient receptor potential mucolipin 1; RyR, ryanodine receptor; IP$_3$R, inositol 1,4,5-trisphosphate receptor; CICR, Ca^{2+}-induced Ca^{2+} release; LD, lipid droplet; LP, lipoprotein; CE, cholesteryl ester; Chol, cholesterol.

A

Gene symbol	Protein name	IF	Ly	Chol
Zfyve27	ISOFORM 1 OF ZINC FINGER FYVE DOMAIN-CONTAINING PROTEIN 27.	1.095		
Stmn2	STATHMIN-2.	0.500		
Vac14	PROTEIN VAC14 HOMOLOG.	0.324		
*Atp6v0d1	VACUOLAR PROTON PUMP SUBUNIT D 1.	0.200	✓	
Sh3gl1	ENDOPHILIN-A2.	0.125		
Gm364	SIMILAR TO TRANSMEMBRANE 9 SUPERFAMILY MEMBER 2.	0.111		
*Lamtor1	UPF0404 PROTEIN C11ORF59 HOMOLOG.	0.111	✓	✓
Tfrc	TRANSFERRIN RECEPTOR PROTEIN 1.	0.100		
*Tmem55b	TRANSMEMBRANE PROTEIN 55B.	0.083	✓	✓
Fig4	SAC DOMAIN-CONTAINING PROTEIN 3.	0.048		
Sun2	PROEIN UNC-84 HOMOLOG B.	0.032		
Vps35	VACUOLAR PROTEIN SORTING-ASSOCIATED PROTEIN 35.	0.026	✓	
Kif13a	KINESIN-LIKE PROTEIN KIF13A.	0.014		

B

C

D

E

Scale: 10 μm

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5