TMEM55B controls lipolysis via lysosomal pH and ER–lysosome calcium signaling

Nagi Mukae[1], Honoka Maki[1] & Michiko Shirane[1]*

[1]Department of Molecular Biology, Graduate School of Pharmaceutical Sciences, Nagoya City University, Nagoya, Aichi, Japan.

*Corresponding: M.S. (shiram@phar.nagoya-cu.ac.jp)

Keywords:
TMEM55B, PDZD8, Lysosomes, Endoplasmic reticulum, Calcium, Membrane contact sites

ABSTRACT

Endolysosomal maturation is accompanied by luminal acidification and increased Ca^{2+} concentration. Mature lysosomes function as major intracellular Ca^{2+} storage organelles, alongside the endoplasmic reticulum (ER) and mitochondria. Recent studies have highlighted a role for lysosomal Ca^{2+} in calcium-induced calcium release (CICR), where local lysosomal Ca^{2+} efflux triggers subsequent ER Ca^{2+} release at ER–lysosome membrane contact sites (MCSs). However, the molecular mechanisms underlying this process remain incompletely understood. In this study, we identified the lysosome-localized protein TMEM55B in a PDZD8-associated protein complex, a factor previously implicated in lipid transfer and membrane tethering at ER–late endosome/lysosome (LE/Ly) contacts. Loss of TMEM55B disrupted lysosomal acidification and impaired lipid droplet degradation. TMEM55B knockdown also reduced lysosomal Ca^{2+} release and uptake, accompanied by attenuated CICR at ER–LE/Ly MCSs. Together, these findings suggest that TMEM55B contributes to the regulation of lysosomal pH and Ca^{2+} dynamics and may influence Ca^{2+} signaling between lysosomes and the ER.

Introduction

ER- LE/Ly MCSs are not formed by random contacts between the ER and endosomes but are mediated by specific membrane tethering complexes. ORP1L and VAP were first identified as tethering factors of ER–LE/Ly MCSs, and subsequent studies revealed that many sterol transfer proteins—including ORP, STARD, and Lam families, as well as NPC1, Protrudin, and PDZD8—also function at these sites[1-3]. The tethering of ER–LE/Ly MCSs is strengthened during endosomal maturation, a process involving both endosomal acidification and Ca^{2+} dynamics. Cholesterol and Ca^{2+} therefore serve as key messengers mediating cross-talk between these two organelles at ER–LE/Ly MCSs[4].

Cells acquire cholesterol through two main pathways: de novo synthesis and endocytosis of lipoproteins (LPs)[5-7]. In the brain, however, neurons lack the ability to synthesize cholesterol and instead rely on supply from glial cells. Cholesterol produced in glia is esterified to cholesteryl esters (CEs), packed into HDL-like lipoproteins, transported to neurons, internalized via LDLR-mediated endocytosis, and hydrolyzed to free cholesterol by lysosomal acidic lipases[7,8]. This cholesterol is then trafficked to the ER and redistributed to other organelles for cellular functions[4,9].

CEs stored in lipid droplets (LDs) are degraded through lipophagy, a form of microautophagy in which LDs fuse with lysosomes, enabling lipids to be hydrolyzed by

lysosomal acidic lipases[10-13]. Endocytosed LP particles are degraded in a similar manner.

Thus, endosomal maturation, which depends on lysosomal acidification, is essential for

metabolizing CEs from both LDs and LPs into cholesterol.

PDZD8 is an ER membrane protein originally identified as a Protrudin-binding partner,

with both proteins acting as tethers at ER–LE/Ly MCSs[14-21]. PDZD8 belongs to the

TULIP super family and mediates cholesterol transport through its SMP domain[16,22,23].

Because Protrudin interacts with VAP via its FFAT motif, the PDZD8–Protrudin–VAP

complex is thought to function in lipid transport[3]. PDZD8 thereby contributes to

cholesterol metabolism by promoting lipolysis through endosomal maturation in

neurons[24]. Consistently, PDZD8-deficient mice exhibit CE accumulation in the brain due

to impaired lipolysis, along with memory and emotional dysfunction[25,26]. CE

accumulation is also observed in the brains of Alzheimer's disease (AD) patients and AD

model mice[27-29]. In particular, neurons carrying the APOE4 allele—the strongest genetic

risk factor for AD—show CE accumulation with lipofuscin formation due to impaired

lysosomal function[30,31].

Compared with cholesterol metabolism, research on lysosomal calcium dynamics is

relatively recent, and many questions remain unresolved. As endosomes undergo luminal

acidification, their Ca^{2+} concentration increases, and lysosomes eventually become a

major intracellular Ca^{2+} store alongside the ER and mitochondria. Ca^{2+} uptake into the

endosomal lumen promotes intraluminal vesicle fusion, thereby facilitating endosomal

maturation[32-35]. Conversely, calcium-induced calcium release (CICR)—in which a small

amount of Ca^{2+} efflux from lysosomes trigger large scale release from the ER IP_3

receptors and ryanodine receptors—has emerged as an important mechanism of transient

Ca^{2+} signaling[36-40]. Endosomal acidification activates lysosomal Ca^{2+} channels, such as

two-pore channel-2 (TPC2) and TRPML1, which promote Ca^{2+} release from lysosomes.

Although Ca^{2+} dynamics at ER–LE/Ly MCSs have been implicated in diverse cellular

processes[35,41], their detailed mechanisms remain poorly understood.

TMEM55B has been reported to perform multiple functions. It was first identified as a

phosphatase that hydrolyzes the 4-position phosphate of $PtdIns(4,5)P_2$ to generate

$PtdIns(5)P$[42,43]. TMEM55B is a lysosomal membrane protein that interacts with the v-

ATPase and Ragulator complexes, thereby contributing to v-ATPase activation and

subsequent mTORC1 signaling at the lysosomal membrane[44]. Consistent with its

importance, TMEM55B knockout mice are embryonic lethal[44]. Because v-ATPase is a

proton pump essential for endolysosomal acidification, its dysfunction leads to various

lysosomal storage diseases[45-47]. TMEM55B also cooperates with TFEB and JIP4 to

promote dynein-dependent retrograde lysosomal transport[48] and participate in lysosomal

oxidative stress responses[49]. In addition, TMEM55B regulates the expression and activity
of LDLR thereby modulating lipoprotein uptake and cholesterol metabolism[50,51].
In this study, we identified lysosomal protein TMEM55B in a PDZD8 complex that may
regulate cholesterol metabolism via lysosomal acidification and Ca^{2+} dynamics at ER–
LE/Ly MCSs.
94
Results
Identification of lysosomal protein TMEM55B in the PDZD8 complex.
In neuronal cells, the lipid transfer protein PDZD8 promotes cholesterol metabolism via
endosomal maturation; however, the mechanism by which this ER protein regulates
lysosomal activity remains unclear. To identify lysosomal proteins interacting with
PDZD8, we established a Neuro2A cell line stably expressing 3×FLAG-tagged PDZD8.
The PDZD8 complex was isolated by anti-FLAG immunoprecipitation, separated by
SDS-PAGE, digested with trypsin, and analyzed by LC-MS/MS. In parallel, the same
analysis was performed using cells transfected with empty vector to exclude nonspecific
binding. Proteins reproducibly detected across three independent experiments were
extracted, and semiquantitative analysis based on identification frequency (IF)[52]
identified 196 proteins with IF > 0.01. Among those, 13 endosome-localized proteins
were selected using GO Term and UniProt subcellular localization annotations (Fig. 1A).
This group included the lysosomal protein TMEM55B and its binding partners, v-ATPase
subunit Atp6v0d1 and Lamtor1 (Fig. 1A)[44]. Notably, Lamtor1 and TMEM55B are
lysosome-localized proteins associated with cholesterol metabolism, suggesting a
potential role in lipolysis-mediated cholesterol processing. Since TMEM55B regulates
the v-ATPase complex, we focused on TMEM55B as a possible upstream regulator of
PDZD8-mediated lysosomal function.
114
TMEM55B is not essential for ER–LE/Ly tethering.
Because PDZD8 functions as a tether at ER–LE/Ly MCSs, we examined whether its
binding partner TMEM55B also contributes to this process. TMEM55B expression was
suppressed in HeLa cells using siRNA (siTMEM55B), resulting in a marked reduction of
both RNA and protein levels compared with control cells (siControl) (Fig. 1B, C). To
assess ER–LE/Ly contacts, we employed a split-GFP system in which GFP fluorescence
is reconstituted only when ER and lysosomal membranes are closely apposed[16,53]. Cells
were co-transfected with ERj1-GFP(1–10) (ER-localized fragment) and LAMP1-
GFP(11) (lysosomal fragment), together with mCherry to normalize expression levels.
The GFP/mCherry fluorescence ratio per cell, reflecting ER–LE/Ly MCS formation, was

quantified (Fig. 1D). No significant difference in GFP/mCherry ratio was observed
between siTMEM55B and siControl cells (Fig. 1E). Compared with prior reports,
including our own showing reduced ER–LE/Ly tethering upon Protrudin or PDZD8
knockdown, TMEM55B knockdown caused only a minor effect, suggesting it is unlikely
a major contributor to tethering of ER–LE/Ly MCSs.
130

TMEM55B regulates lysosomal acidification and activity.
TMEM55B interacts with the v-ATPase complex and has been implicated in its regulation.
We therefore hypothesized that TMEM55B contributes to lysosomal acidification and
activity via v-ATPase. To test this, HeLa cells transfected with siControl or siTMEM55B
were stained with LysoTracker (Fig. 2A) or LysoSensor (Fig. 2C), both of which increase
in fluorescence intensity as lysosomal pH decreases. In both assays, siTMEM55B cells
showed significantly reduced fluorescence compared with controls, indicating impaired
lysosomal acidification (Fig. 2B, D). To assess the functional consequences in neurons,
we suppressed TMEM55B expression in PC12 cells (Fig. 2E) and evaluated lipolysis
using EGFP-PLIN2 as a lipid droplet (LD) marker and LysoTracker Red as a lysosomal
marker (Fig. 2F). TMEM55B knockdown cells exhibited enlarged LDs (Fig. 2G) with
increased LD fluorescence intensity per cell (Fig. 2H), as well as reduced LysoTracker
fluorescence (Fig. 2I), consistent with defective lysosomal acidification and impaired
lipolysis. Furthermore, LD–lysosome fusion events were significantly decreased in
siTMEM55B cells compared with controls (Fig. 2J). Together, these results demonstrate
that TMEM55B may promote lysosomal acidification and activity, thereby facilitating
LD degradation and lipolysis. However, since the signals from the Lysotracker and
Lysosensor alone cannot reliably quantify lysosomal pH, a colorimetric probe that
directly measures lysosomal pH would be useful.
150

TMEM55B promotes Ca^{2+} release from the ER.
Intracellular Ca^{2+} elevation promotes lysosome fusion with other organelles[45], suggesting
that lipolysis mediated by the PDZD8–TMEM55B complex could involve Ca^{2+} signaling
at ER–LE/Ly MCSs. Since the ER is the major intracellular Ca^{2+} store, we investigated
whether TMEM55B regulates Ca^{2+} release or uptake from the ER. HeLa cells were
transfected with the ER-targeted Ca^{2+} indicator G-CEPIA1er[54], which binds ER luminal
Ca^{2+} and emits GFP fluorescence, and colocalization with the ER marker KDEL was
confirmed by confocal microscopy (Fig. 3A, B). To assess ER Ca^{2+} dynamics, siControl
or siTMEM55B cells were stimulated with ATP to induce transient ER Ca^{2+} release,
followed by Ca^{2+} reuptake to the ER. In siTMEM55B cells, ATP-induced Ca^{2+} release as

well as reuptake were markedly reduced compared with controls (Fig. 3C, D).
Quantitative analysis revealed that siTMEM55B cells exhibited significantly decreased
rates of ER Ca^{2+} release (Fig. 3E), slower subsequent Ca^{2+} reuptake (Fig. 3F), and delayed
time to peak Ca^{2+} release (Fig. 3G). These findings indicate that TMEM55B may
facilitates ATP-stimulated transient Ca^{2+} release from the ER.
166
TMEM55B promotes Ca^{2+} release from lysosomes.
We next investigated whether TMEM55B regulates Ca^{2+} release from lysosomes and how
this affects ER Ca^{2+} dynamics. Lysosomal Ca^{2+} release and uptake were monitored using
Ly-GG, which binds cytosolic Ca^{2+} at the lysosomal membrane and emits GFP
fluorescence[55] (Fig. 4A). Co-expression of Ly-GG with the lysosomal marker LAMP1
confirmed colocalization (Fig. 4B). ATP stimulation of siControl HeLa cells induced
transient Ca^{2+} release from lysosomes. In siTMEM55B cells, the magnitude (Fig. 4C) and
rate (Fig. 4D) of lysosomal Ca^{2+} release were significantly reduced compared to controls.
The subsequent Ca^{2+} reuptake rate was also decreased (Fig. 4E), indicating impaired
lysosomal Ca^{2+} dynamics. As a control, we examined mitochondrial Ca^{2+} using
CEPIA3mt[54], which binds mitochondrial matrix Ca^{2+}, together with the mitochondrial
marker Su9-RFP[56] (Fig. 4F), confirmed colocalization (Fig. 4G). ATP stimulation
induced transient mitochondrial Ca^{2+} increases as well as decrease, which were not
significantly different between siControl and siTMEM55B cells (Fig. 4H–J). These
results indicate that TMEM55B may specifically regulates lysosomal Ca^{2+} dynamics
without affecting mitochondrial Ca^{2+}.
183
Discussion
Lysosomes serve as central hubs for cellular metabolism, coordinating degradation,
nutrient recycling, and signaling. Proper lysosomal acidification and calcium homeostasis
are essential for lipid metabolism and neuronal function, and their dysregulation has been
increasingly linked to neurodegenerative diseases. Despite this, the molecular
mechanisms connecting lysosomal function to lipid regulation remain incompletely
understood.
In this study, we identify TMEM55B as a key lysosomal regulator that interacts with the
ER protein PDZD8 to control lysosomal acidification and calcium dynamics. Through
these functions, TMEM55B in PDZD8 complex may contribute to endosomal maturation
and lipid droplet degradation, revealing a novel mechanism linking lysosomal activity to
lipid metabolism. These findings provide new insights into cellular lipid homeostasis and

suggest potential targets for disorders associated with lysosomal dysfunction and
cholesterol metabolism.

However, immunoprecipitation-immunoblotting analysis of endogenous proteins
yielded signals below the detection threshold, and that of overexpressed exogenous
proteins resulted in non-specific interactions, preventing us from demonstrating direct
binding between TMEM55B and PDZD8. Therefore, while both proteins reside within
the same complex, the interaction may be indirect rather than direct, and TMEM55B-
dependent regulation of lysosomal acidification and calcium dynamics may occur
independently of PDZD8.

Because Ly-GG can detect Ca^{2+} originating from the ER in proximity to lysosomes,
reduced ER-derived local Ca^{2+} release may also contribute to the decreased Ly-GG signal
following TMEM55B knockdown.

The lipid transfer protein PDZD8 mediates tethering of ER–LE/Ly membrane contact
sites (MCSs) and facilitates cholesterol metabolism through lipolysis via endosomal
maturation. However, the mechanism by which PDZD8 promotes endosome maturation
has remained unclear. Through complex analysis in neuronal cells, we identified Lamtor1
and TMEM55B as lysosome-localized, cholesterol-associated proteins among LE/Ly
candidates. Given that TMEM55B regulates the v-ATPase complex, we focused on its
intracellular functions.

We found that TMEM55B contributes to lysosomal acidification and activation. Loss of
TMEM55B impaired lysosomal Ca^{2+} release and uptake, as well as ATP-stimulated Ca^{2+}
release from the ER, while mitochondrial Ca^{2+} dynamics remained unaffected. In our
system, lysosomal Ca^{2+} release peaked within 180 s after ATP stimulation (Fig. 4C),
whereas ER Ca^{2+} release peaked later (Fig. 3C), consistent with calcium-induced calcium
release (CICR) from the ER triggered by lysosomal Ca^{2+} release. These results indicate
that TMEM55B promotes endosomal maturation via lysosomal acidification and
activation, and facilitates CICR at ER–LE/Ly MCSs, thereby supporting lipolysis (Fig.
5).

The PDZD8–TMEM55B complex thus plays a central role in ER–LE/Ly MCS functions,
including endosomal maturation, Ca^{2+} dynamics, and cholesterol metabolism.
Nevertheless, the effects of TMEM55B deficiency on lysosomal fusion and lipid droplet
degradation were partial, failing to recapitulate the strong phenotype observed with
PDZD8 deficiency. This suggests that additional, as-yet-unidentified mechanisms
contribute to PDZD8-mediated lipolysis.

In summary, TMEM55B enhances lysosomal acidification via v-ATPase regulation,
thereby increasing lysosomal enzyme activity and promoting lipolysis. Lysosomal

acidification also activates Ca²⁺ channels, facilitating CICR at ER–LE/Ly MCSs and
supporting cholesterol metabolism through lysosomal fusion. Direct evidence linking
TMEM55B-mediated CICR to enhanced lipolysis, however, remains lacking, and further
studies are needed to clarify this mechanism.
236
Limitation of this study
We were unable to demonstrate direct binding between endogenous or exogenous PDZD8
and TMEM55B proteins. Attempts to express TMEM55B at endogenous levels for rescue
experiments were also unsuccessful. Additionally, assessing V-ATPase assembly
following TMEM55B knockdown proved technically difficult and was beyond the scope
of this work. Although TMEM55B deficiency caused partial defects in lysosomal fusion
and lipid droplet degradation, the phenotype was less pronounced than that observed with
PDZD8 loss. These findings imply that additional, unidentified mechanisms may
contribute to PDZD8-dependent lipolysis, which remain unresolved in this study.
246
Methods
Plasmids
EGFP and mCherry vectors were from Clontech (Mountain View, CA, US). Human
cDNAs encoding PLIN2 (GenBank accession number NM_007408.4), LAMP1
(GenBank accession number BC025335), and KDEL with the signal sequence of
calreticulin (GenBank accession number NM_004343) were amplified from HeLa cells
by the polymerase chain reaction (PCR) with PrimeSTAR HS DNA polymerase (Takara,
Shiga, Japan). pCMV-G-CEPIA1er (Addgene plasmid #58215) and pCMV-CEPIA3mt
(Addgene plasmid #58219) were obtained from Addgene (Watertown, MA, US). Ly-GG
was kindly provided by Dr. CW Taylor (University of Cambridge, UK). Vectors for split-
GFP analysis, ERj1–GFP(1–10) and LAMP1–GFP(11), were kindly provided by Dr.
Tamura (Yamagata University, Japan). The pSu9 cDNA was kindly provided by Dr.
Mihara (Kyushu University, Japan). The mRFP cDNA was kindly provided by Dr. R.Y.
Tsien (University of California, San Diego, US).
261
262   **Antibodies**
263   Antibodies to TMEM55B were from Proteintech (Rosemont, IL, USA); antibodies to
264   HSP90 were from BD Biosciences (Franklin Lakes, NJ, USA).
265
Cell culture, transfection, and retrovirus infection

HeLa cells (CCL-2, purchased from ATCC, US) were cultured under a humidified
atmosphere of 5% CO_2 at 37°C in Dulbecco's modified Eagle's medium (Invitrogen Life
Technologies) supplemented with 10% fetal bovine serum (Invitrogen Life Technologies).
The culture medium for Plat-E cells was also supplemented with blasticidin (10 μg/ml).
Cells were transfected with the use of XtremeGene9 reagents (Roche Diagnostics,
Mannheim, Germany) according to the manufacturer's protocol. For retroviral infection,
Plat-E cells were transiently transfected with pMX-puro–based vectors and then cultured
for 48 h. The retroviruses in the resulting culture supernatants were used to infect
Neuro2A cells, and the cells were then subjected to selection with puromycin (1 μg/ml).
276

Identification and categorization of PDZD8-associated proteins
Neuro2a cells infected with a retrovirus encoding FLAG–tagged mouse PDZD8 or with
the corresponding empty virus were solubilized in a lysis buffer (40 mM HEPES-NaOH
[pH 7.5], 150 mM NaCl, 10% glycerol, 0.5% Triton X-100, 1 mM Na_3VO_4, 25 mM NaF,
aprotinin [10 μg/ml], leupeptin [10 μg/ml], 1 mM phenylmethylsulfonyl fluoride) to
prepare whole-cell extracts. Lysates were cleared by centrifugation at 20,400 × g for 15
min at 4 °C. The clarified supernatants were subjected to affinity purification with anti-
FLAG (M2)–agarose affinity gel (Sigma-Aldrich), and then proteins eluted with FLAG
peptide (Sigma-Aldrich) was separated by SDS-PAGE and stained with silver. The
stained gels were sliced into pieces, and the abundant proteins therein were subjected to
in-gel digestion with trypsin. The resulting peptides were dried, dissolved in a mixture of
0.1% trifluoroacetic acid and 2% acetonitrile, and then applied to a nanoflow LC system
(Advanced UHPLC; Michrom BioResources, Auburn, CA, USA) equipped with an L-
column (C18, 0.15 by 50 mm, particle size of 3 μm; CERI, Tokyo, Japan). LC–MS/MS
analysis was performed with an Orbitrap Velos Pro instrument (Thermo Fisher Scientific).
All MS/MS spectra were compared with protein sequences in the International Protein
Index (IPI, European Bioinformatics Institute) mouse version 3.44 with the use of the
MASCOT algorithm. Assigned high-scoring peptide sequences were considered for
correct identification. Identified peptides from 3 independent experiments were integrated
and regrouped by IPI accession number. Proteins detected in ≥2 of 3 independent
experiments from FLAG-PDZD8–expressing cells, but absent from control cells, were
considered PDZD8-associated. Semi-quantitative abundance was estimated from
identification frequency (IF; number of identified peptides per protein normalized to the
number of theoretically detectable tryptic peptides). Functional categorization of
PDZD8-associated proteins was categorized with use of the UniProtKB database
(https://www.uniprot.org) based on Gene Ontology (GO) annotations. The proteins listed

in figure 1a shows endosome-related (GO:0005768), as well as columns for lysosome
(GO:0005764) or cholesterol-related as well.

RNA interference, reverse transcription, and real-time PCR analysis

Stealth siRNAs targeted to human TMEM55B (siTMEM55B#1, 5'-
ACGUGGAAGGCAAGAUGCAUCAGCA-3'; siTMEM55B#2, 5'-
CAAGAUGCAUCAGCAUGUAGUCAAA-3'), rat TMEM55B (siTMEM55B#1, 5'-
UGCUGAUGCAUCUUGCCUUCCACGU-3'; siTMEM55B#2, 5'-
CAAGAUGCAUCAGCAUGUAGUCAAA-3'), and non-targeting control were obtained
from Invitrogen Life Technologies. (Invitrogen Life Technologies, Carlsbad, CA, USA).
Human siRNAs were introduced into HeLa cells with the use of Lipofectamine
RNAiMAX (Invitrogen Life Technologies), and rat siRNAs were introduced into PC12
cells with the use of a Cell Line Nucleofector Kit V (VCA-1003) and Nucleofector
instrument (Lonza, Basel, Switzerland) according to the manufacturer's protocol. The
cells were then cultured for 48 h (for HeLa) or 72 h (for PC12). Total RNA was isolated
from cells with the use of an RNeasy Mini Kit (Qiagen, Hilden, Germany) according to
the manufacturer's protocol and was subjected to reverse transcription (RT) with the use
of a QuantiTect Rev Transcription Kit (Qiagen). The resulting cDNA was subjected to
real-time PCR analysis with Power SYBR Green PCR Master Mix in an ABI-Prism 7000
sequence detection system (Applied Biosystems, Foster City, CA). The amounts of each
target mRNA were calculated and normalized by that of GAPDH mRNA. Primers for
PCR (forward and reverse, respectively) were as follows: 5'-
GTCAGTGGTGGACCTGACCTG-3' and 5'-AAAGTGGTCGTTGAGGGCAAT-3' for
human GAPDH; 5'-CGAGTCTGCCAATCTCTCATC-3' and 5'-
AGTGCGGTCTGTGAACTCTGT-3' for human TMEM55B; 5'-
AAGGCTGAGAATGGGAAGCTG-3' and 5'-GGAGATGATGACCCTTTTGGC-3' for
rat GAPDH; 5'-CTAGCCCGGATAGTGGGAGTG-3' and 5'-
TTCCTGGAGGTGCATTCTTGA-3' for rat TMEM55B.

Fluorescence imaging of live cells

LysoTracker Red DND-99 (0.2 µM) or LysoSensor Green DND-189 (0.2 µM) (Thermo
Fisher Scientific, Rockford, IL, USA) were added to cells for 1 h at 37 °C according to
the manufacturer's protocol. HeLa and PC12 cells expressing plasmid-encoded
fluorescent proteins for 24 h and 96 h respectively, or metabolically labeled with
fluorescent probes were observed with an LSM800 confocal microscope (Zeiss,

Oberkochen, Germany), and the images were processed for calculation of fluorescence

intensity with ZEN imaging software (Zeiss) as well as Fiji (ImageJ, https://fiji.sc).

Ca^{2+} imaging

Cell media of HeLa cells expressing organelle-targeted Ca^{2+} indicators were pre-

incubated with HEPES-buffered saline (HBS; 137 mM NaCl, 5.9 mM KCl, 2.2 mM CaCl$_2$,

1.2 mM MgCl$_2$, 10 mM HEPES, 14 mM glucose, pH 7.4) for 1 h at 37 °C and then

stimulated with ATP (200 µM) (Wako, Japan). The images were captured by Operetta

high-content screening system (PerkinElmer, Waltham, MA, USA) using a ×20 objective

(λ_ex 460–490 nm; λ_em 500–550 nm) at the time point of ATP addition (t = 0 s) and

continued time points (t = X s) at a rate of one frame per 20 sec for total 1240 sec (G-

CEPIA) or 180 sec (Ly-GG and CEPIA3mt). The concentration of Ca^{2+} for each time

points was calculated by $\Delta F/F_0 = (F_x - F_0)/F_0$. Images were analyzed by Harmony

software (PerkinElmer).

Immunoblot analysis

HeLa cells were lysed in Triton lysis buffer containing 40 mM HEPES-NaOH (pH 7.6),

150 mM NaCl, 0.5% Triton X-100, 10 mM MgCl$_2$, 1 mM Na$_3$VO$_4$, 25 mM NaF, 1 mM

EDTA, 1 mM PMSF, leupeptin (10 µg/mL), and 10% glycerol. After 10 min on ice,

lysates were centrifuged at 15,000 × g for 10 min at 4 °C, and protein concentrations were

determined with the Pierce BCA Protein Assay (Thermo Fisher Scientific). Extracts were

resolved by SDS–PAGE, transferred to Immobilon-P membranes (Merck Millipore,

Darmstadt, Germany), and probed with primary antibodies. Signals were developed with

SuperSignal reagents (Thermo Fisher Scientific) and imaged on an LAS-4000 (GE

Healthcare, Waukesha, WI, USA).

Statistical analysis

Statistical analyses were performed with BellCurve for Excel software (Social Survey

Research Information, Tokyo, Japan). Normality of data was assessed with the Shapiro–

Wilk test. If the normality assumption was not met, the Kruskal–Wallis test with the

Steel's multiple comparisons test were applied. If data were normally distributed, One-

way repeated-measures analysis of variance (ANOVA) with the Dunnett's multiple

comparisons test were performed. Quantitative data are presented as mean ± SE.

Statistical significance of a P value is indicated as $0.01 \leq p < 0.05$ (*); $0.001 \leq p < 0.01$

(**); $p < 0.001$ (***); $p \geq 0.05$ (n.s.). Sample size (n) are in the figure legends.

References

1 Luo, J., Jiang, L. Y., Yang, H. & Song, B. L. Intracellular Cholesterol Transport
by Sterol Transfer Proteins at Membrane Contact Sites. *Trends Biochem Sci* **44**,
273-292, doi:10.1016/j.tibs.2018.10.001 (2019).

2 Wong, L. H., Gatta, A. T. & Levine, T. P. Lipid transfer proteins: the lipid
commute via shuttles, bridges and tubes. *Nat Rev Mol Cell Biol* **20**, 85-101,
doi:10.1038/s41580-018-0071-5 (2019).

3 Shirane, M. Lipid Transfer-Dependent Endosome Maturation Mediated by
Protrudin and PDZD8 in Neurons. *Front Cell Dev Biol* **8**, 615600,
doi:10.3389/fcell.2020.615600 (2020).

4 van der Kant, R. & Neefjes, J. Small regulators, major consequences - Ca(2)(+)
and cholesterol at the endosome-ER interface. *J Cell Sci* **127**, 929-938,
doi:10.1242/jcs.137539 (2014).

5 Ikonen, E. Cellular cholesterol trafficking and compartmentalization. *Nat Rev*
Mol Cell Biol **9**, 125-138, doi:10.1038/nrm2336 (2008).

6 Goldstein, J. L. & Brown, M. S. A century of cholesterol and coronaries: from
plaques to genes to statins. *Cell* **161**, 161-172, doi:10.1016/j.cell.2015.01.036
(2015).

7 Luo, J., Yang, H. & Song, B. L. Mechanisms and regulation of cholesterol
homeostasis. *Nat Rev Mol Cell Biol* **21**, 225-245, doi:10.1038/s41580-019-0190-
7 (2020).

8 Mehta, N. *et al.* The evolving role of cholesteryl ester transfer protein inhibition
beyond cardiovascular disease. *Pharmacol Res* **197**, 106972,
doi:10.1016/j.phrs.2023.106972 (2023).

9 Martin, M. G., Pfrieger, F. & Dotti, C. G. Cholesterol in brain disease:
sometimes determinant and frequently implicated. *EMBO Rep* **15**, 1036-1052,
doi:10.15252/embr.201439225 (2014).

10 Zechner, R., Madeo, F. & Kratky, D. Cytosolic lipolysis and lipophagy: two
sides of the same coin. *Nat Rev Mol Cell Biol* **18**, 671-684,
doi:10.1038/nrm.2017.76 (2017).

11 Schuck, S. Microautophagy - distinct molecular mechanisms handle cargoes of
many sizes. *J Cell Sci* **133**, doi:10.1242/jcs.246322 (2020).

12 Haidar, M., Loix, M., Bogie, J. F. J. & Hendriks, J. J. A. Lipophagy: a new
player in CNS disorders. *Trends Endocrinol Metab* **32**, 941-951,
doi:10.1016/j.tem.2021.08.010 (2021).

13 Lan, Z. Q., Ge, Z. Y., Lv, S. K., Zhao, B. & Li, C. X. The regulatory role of
lipophagy in central nervous system diseases. *Cell Death Discov* **9**, 229,
doi:10.1038/s41420-023-01504-z (2023).

14 Shirane, M. & Nakayama, K. I. Protrudin induces neurite formation by
directional membrane trafficking. *Science* **314**, 818-821,
doi:10.1126/science.1134027 (2006).

15 Guillen-Samander, A., Bian, X. & De Camilli, P. PDZD8 mediates a Rab7-
dependent interaction of the ER with late endosomes and lysosomes. *Proc Natl*
Acad Sci U S A **116**, 22619-22623, doi:10.1073/pnas.1913509116 (2019).

16 Shirane, M. *et al.* Protrudin and PDZD8 contribute to neuronal integrity by
promoting lipid extraction required for endosome maturation. *Nat Commun* **11**,
4576, doi:10.1038/s41467-020-18413-9 (2020).

17 Elbaz-Alon, Y. *et al.* PDZD8 interacts with Protrudin and Rab7 at ER-late
endosome membrane contact sites associated with mitochondria. *Nat Commun*
11, 3645, doi:10.1038/s41467-020-17451-7 (2020).

18 Khan, H., Chen, L., Tan, L. & Im, Y. J. Structural basis of human PDZD8-Rab7
interaction for the ER-late endosome tethering. *Sci Rep* **11**, 18859,
doi:10.1038/s41598-021-98419-5 (2021).

19 Gao, Y., Xiong, J., Chu, Q. Z. & Ji, W. K. PDZD8-mediated lipid transfer at
contacts between the ER and late endosomes/lysosomes is required for neurite
outgrowth. *J Cell Sci* **135**, doi:10.1242/jcs.255026 (2022).

20 Thakur, R. S. & O'Connor-Giles, K. M. PDZD8 promotes autophagy at ER-
lysosome membrane contact sites to regulate activity-dependent synaptic
growth. *Cell Rep* **44**, 115483, doi:10.1016/j.celrep.2025.115483 (2025).

21 Yang, H. *et al.* LYVAC/PDZD8 is a lysosomal vacuolator. *Science* **389**,
eadz0972, doi:10.1126/science.adz0972 (2025).

22 Alva, V. & Lupas, A. N. The TULIP superfamily of eukaryotic lipid-binding
proteins as a mediator of lipid sensing and transport. *Biochim Biophys Acta*
1861, 913-923, doi:10.1016/j.bbalip.2016.01.016 (2016).

23 Wong, L. H. & Levine, T. P. Tubular lipid binding proteins (TULIPs) growing
everywhere. *Biochim Biophys Acta Mol Cell Res* **1864**, 1439-1449,
doi:10.1016/j.bbamcr.2017.05.019 (2017).

24 Morita, K. *et al.* PDZD8-deficient mice accumulate cholesteryl esters in the
brain as a result of impaired lipophagy. *iScience* **25**, 105612,
doi:10.1016/j.isci.2022.105612 (2022).

25   Al-Amri, A. H. *et al.* PDZD8 Disruption Causes Cognitive Impairment in
445   Humans, Mice, and Fruit Flies. *Biol Psychiatry* **92**, 323-334,
446   doi:10.1016/j.biopsych.2021.12.017 (2022).

447   26   Kurihara, Y. *et al.* PDZD8-deficient mice manifest behavioral abnormalities
448   related to emotion, cognition, and adaptation due to dyslipidemia in the brain.
449   *Mol Brain* **16**, 11, doi:10.1186/s13041-023-01002-4 (2023).

450   27   Chan, R. B. *et al.* Comparative lipidomic analysis of mouse and human brain
451   with Alzheimer disease. *J Biol Chem* **287**, 2678-2688,
452   doi:10.1074/jbc.M111.274142 (2012).

453   28   van der Kant, R. *et al.* Cholesterol Metabolism Is a Druggable Axis that
454   Independently Regulates Tau and Amyloid-beta in iPSC-Derived Alzheimer's
455   Disease Neurons. *Cell Stem Cell* **24**, 363-375 e369,
456   doi:10.1016/j.stem.2018.12.013 (2019).

457   29   Nugent, A. A. *et al.* TREM2 Regulates Microglial Cholesterol Metabolism upon
458   Chronic Phagocytic Challenge. *Neuron* **105**, 837-854 e839,
459   doi:10.1016/j.neuron.2019.12.007 (2020).

460   30   Blanchard, J. W. *et al.* APOE4 impairs myelination via cholesterol dysregulation
461   in oligodendrocytes. *Nature* **611**, 769-779, doi:10.1038/s41586-022-05439-w
462   (2022).

463   31   Guo, J. L. *et al.* Decreased lipidated ApoE-receptor interactions confer
464   protection against pathogenicity of ApoE and its lipid cargoes in lysosomes. *Cell*
465   **188**, 187-206 e126, doi:10.1016/j.cell.2024.10.027 (2025).

466   32   Mayran, N., Parton, R. G. & Gruenberg, J. Annexin II regulates multivesicular
467   endosome biogenesis in the degradation pathway of animal cells. *EMBO J* **22**,
468   3242-3253, doi:10.1093/emboj/cdg321 (2003).

469   33   Lloyd-Evans, E. *et al.* Niemann-Pick disease type C1 is a sphingosine storage
470   disease that causes deregulation of lysosomal calcium. *Nat Med* **14**, 1247-1255,
471   doi:10.1038/nm.1876 (2008).

472   34   Bissig, C. *et al.* Viral infection controlled by a calcium-dependent lipid-binding
473   module in ALIX. *Dev Cell* **25**, 364-373, doi:10.1016/j.devcel.2013.04.003
474   (2013).

475   35   Bretou, M. *et al.* Accumulation of APP C-terminal fragments causes
476   endolysosomal dysfunction through the dysregulation of late endosome to
477   lysosome-ER contact sites. *Dev Cell* **59**, 1571-1592 e1579,
478   doi:10.1016/j.devcel.2024.03.030 (2024).

479   36   Churchill, G. C. *et al.* NAADP mobilizes Ca(2+) from reserve granules,
480   lysosome-related organelles, in sea urchin eggs. *Cell* **111**, 703-708,
481   doi:10.1016/s0092-8674(02)01082-6 (2002).

482   37   Calcraft, P. J. *et al.* NAADP mobilizes calcium from acidic organelles through
483   two-pore channels. *Nature* **459**, 596-600, doi:10.1038/nature08030 (2009).

484   38   Galione, A. A primer of NAADP-mediated Ca(2+) signalling: From sea urchin
485   eggs to mammalian cells. *Cell Calcium* **58**, 27-47,
486   doi:10.1016/j.ceca.2014.09.010 (2015).

487   39   Kilpatrick, B. S. *et al.* An Endosomal NAADP-Sensitive Two-Pore Ca(2+)
488   Channel Regulates ER-Endosome Membrane Contact Sites to Control Growth
489   Factor Signaling. *Cell Rep* **18**, 1636-1645, doi:10.1016/j.celrep.2017.01.052
490   (2017).

491   40   Yuan, Y. *et al.* Two-pore channel-2 and inositol trisphosphate receptors
492   coordinate Ca(2+) signals between lysosomes and the endoplasmic reticulum.
493   *Cell Rep* **43**, 113628, doi:10.1016/j.celrep.2023.113628 (2024).

494   41   Zheng, Q. *et al.* Calcium transients on the ER surface trigger liquid-liquid phase
495   separation of FIP200 to specify autophagosome initiation sites. *Cell* **185**, 4082-
496   4098 e4022, doi:10.1016/j.cell.2022.09.001 (2022).

497   42   Ungewickell, A. *et al.* The identification and characterization of two
498   phosphatidylinositol-4,5-bisphosphate 4-phosphatases. *Proc Natl Acad Sci U S*
499   *A* **102**, 18854-18859, doi:10.1073/pnas.0509740102 (2005).

500   43   Zou, J., Marjanovic, J., Kisseleva, M. V., Wilson, M. & Majerus, P. W. Type I
501   phosphatidylinositol-4,5-bisphosphate 4-phosphatase regulates stress-induced
502   apoptosis. *Proc Natl Acad Sci U S A* **104**, 16834-16839,
503   doi:10.1073/pnas.0708189104 (2007).

504   44   Hashimoto, Y., Shirane, M. & Nakayama, K. I. TMEM55B contributes to
505   lysosomal homeostasis and amino acid-induced mTORC1 activation. *Genes*
506   *Cells* **23**, 418-434, doi:10.1111/gtc.12583 (2018).

507   45   Xu, H. & Ren, D. Lysosomal physiology. *Annu Rev Physiol* **77**, 57-80,
508   doi:10.1146/annurev-physiol-021014-071649 (2015).

509   46   Banerjee, S. & Kane, P. M. Regulation of V-ATPase Activity and Organelle pH
510   by Phosphatidylinositol Phosphate Lipids. *Front Cell Dev Biol* **8**, 510,
511   doi:10.3389/fcell.2020.00510 (2020).

512   47   Ballabio, A. & Bonifacino, J. S. Lysosomes as dynamic regulators of cell and
513   organismal homeostasis. *Nat Rev Mol Cell Biol* **21**, 101-118,
514   doi:10.1038/s41580-019-0185-4 (2020).

48   Willett, R. *et al.* TFEB regulates lysosomal positioning by modulating
TMEM55B expression and JIP4 recruitment to lysosomes. *Nat Commun* **8**,
1580, doi:10.1038/s41467-017-01871-z (2017).
49   Jeong, E., Willett, R., Rissone, A., La Spina, M. & Puertollano, R. TMEM55B
links autophagy flux, lysosomal repair, and TFE3 activation in response to
oxidative stress. *Nat Commun* **15**, 93, doi:10.1038/s41467-023-44316-6 (2024).
50   Medina, M. W. *et al.* Transmembrane protein 55B is a novel regulator of cellular
cholesterol metabolism. *Arterioscler Thromb Vasc Biol* **34**, 1917-1923,
doi:10.1161/ATVBAHA.113.302806 (2014).
51   Qin, Y. *et al.* Phosphatidylinositol-(4,5)-Bisphosphate Regulates Plasma
Cholesterol Through LDL (Low-Density Lipoprotein) Receptor Lysosomal
Degradation. *Arterioscler Thromb Vasc Biol* **40**, 1311-1324,
doi:10.1161/ATVBAHA.120.314033 (2020).
52   Matsumoto, M. *et al.* Large-scale proteomic analysis of tyrosine-
phosphorylation induced by T-cell receptor or B-cell receptor activation reveals
new signaling pathways. *Proteomics* **9**, 3549-3563,
doi:10.1002/pmic.200900011 (2009).
53   Kakimoto, Y. *et al.* Visualizing multiple inter-organelle contact sites using the
organelle-targeted split-GFP system. *Sci Rep* **8**, 6175, doi:10.1038/s41598-018-
24466-0 (2018).
54   Suzuki, J. *et al.* Imaging intraorganellar Ca2+ at subcellular resolution using
CEPIA. *Nat Commun* **5**, 4153, doi:10.1038/ncomms5153 (2014).
55   Atakpa, P., Thillaiappan, N. B., Mataragka, S., Prole, D. L. & Taylor, C. W.
IP(3) Receptors Preferentially Associate with ER-Lysosome Contact Sites and
Selectively Deliver Ca(2+) to Lysosomes. *Cell Rep* **25**, 3180-3193 e3187,
doi:10.1016/j.celrep.2018.11.064 (2018).
56   Otera, H. *et al.* A novel insertion pathway of mitochondrial outer membrane
proteins with multiple transmembrane segments. *J Cell Biol* **179**, 1355-1363,
doi:10.1083/jcb.200702143 (2007).
544

Acknowledgements and Funding Information

We thank Y. Tamura for split-GFP cDNAs; E. Taylor for Ly-GG cDNA; K. Mihara for
Su9 cDNA; K. I. Nakayama and T. Ohnishi for proteome analysis; K. Sato and K. Kato
for Ca^{2+} imaging analysis; and the Research Equipment Sharing Center at Nagoya City
University for assistance. This work was supported in part by KAKENHI grants from
Japan Society for the Promotion of Science (JSPS) to M.S. (24K02122).

Author contributions statement

M.S. designed experiments, supervised the study, analyzed the data, and wrote the manuscript. N.M. and H.M. performed experiments.

Additional information

Data availability

The MS data have been deposited in the JPOST Repository Database as follows:

URL: https://repository.jpostdb.org

JPID and PXID are as follows;

Mock-FLAG complex in Neuro2A; (1) JPST004039, PXD067776; (2) JPST004040, PXD067777; (3) JPST004041, PXD067779.

PDZD8-FLAG complex in Neuro2A; (1) JPST004042, PXD067780; (2) JPST004043, PXD067782; (3) JPST004044, PXD067783.

Supplementary datasets

Raw data are provided within the supplementary information file.

Competing interests

The authors declare no competing interests.

Correspondence

Correspondence and requests for materials should be addressed to M.S. (shiram@phar.nagoya-cu.ac.jp)

Figure Legends

Fig. 1. Identification of PDZD8-associated proteins by proteomic analysis. (**A**) Endosome-related proteins identified as PDZD8-associated proteins. Protein abundance was estimated semiquantitatively by identification frequency (IF). Asterisks indicate proteins previously reported as TMEM55B-associated. Lysosome (Ly)-localized and cholesterol (Chol)-related proteins are also indicated. Proteins were categorized based on Gene Ontology (GO) terms and UniProtKB annotations. (**B, C**) HeLa cells were transfected with siControl, siTMEM55B#1, or siTMEM55B#2. Total RNA was analyzed by RT-qPCR for TMEM55B mRNA (B), and protein levels were examined by immunoblotting with anti-TMEM55B and anti-HSP90 antibodies (C). Data are shown as mean ± SE (n = 9 for RNA analysis). ***p < 0.001 (one-way ANOVA with Dunnett's multiple comparisons). (**D**) HeLa cells were transfected with siControl, siTMEM55B#1, or siTMEM55B#2 together with split-GFP vectors ERj1–GFP(1–10) and LAMP1–GFP(11), along with mCherry as a transfection control. Scale bar, 10 μm. (**E**) Quantification of split-GFP fluorescence normalized to mCherry intensity as in (D). n = 38, 36, and 37 cells for siControl, siTMEM55B#1, and siTMEM55B#2, respectively. n.s., not significant (Kruskal–Wallis test with Steel's multiple comparisons).

Fig. 2. TMEM55B depletion impairs lysosomal acidification and lipolysis. (**A, C**) HeLa cells transfected with siControl, siTMEM55B#1, or siTMEM55B#2 were stained with LysoTracker Red (A) or LysoSensor Green (C) and analyzed by confocal microscopy. Boxed areas (top panels) are shown at higher magnification (bottom panels). Scale bars, 10 μm (top and middle) and 5 μm (bottom). (**B, D**) Quantification of LysoTracker Red (B) or LysoSensor Green (D) mean fluorescence intensity per lysosome in (A) (n = 746, 497, and 452 lysosomes for siControl, siTMEM55B#1, and siTMEM55B#2, respectively) or (C) (n = 703, 847, and 794 lysosomes, respectively). (**E**) RT-qPCR analysis of TMEM55B mRNA in PC12 cells transfected with siControl, siTMEM55B#1, or siTMEM55B#2 (mean ± SE, n = 6). (**F**) PC12 cells were transfected with siControl, siTMEM55B#1, or siTMEM55B#2 together with EGFP-PLIN2, and stained with LysoTracker Red. (**G**) Quantification of lipid droplet (LD) size (μm²) of EGFP-PLIN2 puncta in (F) (n = 627, 450, and 317 LDs for siControl, siTMEM55B#1, and siTMEM55B#2, respectively). (**H**) Quantification of mean EGFP-PLIN2 fluorescence intensity per cell in (F) (n = 52, 41, and 35 cells for siControl, siTMEM55B#1, and siTMEM55B#2, respectively). (**I**) Quantification of mean LysoTracker Red fluorescence intensity per cell in (F) (n = 52, 41, and 35 cells, respectively). (**J**) Pearson's correlation coefficient between EGFP-PLIN2 and LysoTracker Red per cell in (F) (n = 52, 41, and

35 cells, respectively). ***$p < 0.001$, **$p < 0.01$ (Kruskal–Wallis test with Steel's

multiple comparisons test in B, D, and G–I); **$p < 0.01$, *$p < 0.05$ (one-way ANOVA

with Dunnett's multiple comparisons in E and J).

Fig. 3. Depletion of TMEM55B impairs Ca^{2+} release from the ER. (**A**) Schematic

representation of G-CEPIA1er, a Ca^{2+} indicator with the G-GECO1 domain facing the

ER lumen. (**B**) Confocal images of HeLa cells co-expressing G-CEPIA1er (green) and

the ER marker mCherry-KDEL (red). Scale bar, 10 μm. (**C**) Time lapse analysis of G-

CEPIA1er fluorescence ($\Delta F/F_0$) in cells transfected with siControl (gray, n = 30 fields),

siTMEM55B#1 (red, n = 30), or siTMEM55B#2 (pink, n = 30). ATP (200 μM) was added

at t = 0 s; Data are mean ± SE. (**D**) Quantification of peak $\Delta F/F_0$ values in (C).

(**E**) Quantification of the slope of G-CEPIA1er $\Delta F/F_0$ during 0–120 s after ATP addition

($\Delta F/F_0 \cdot s^{-1}$). (**F**) Quantification of time (s) to peak after ATP addition for G-CEPIA1er

$\Delta F/F_0$. (**G**) Quantification of the slope of G-CEPIA1er $\Delta F/F_0$ during 900 s after the peak

($\Delta F/F_0 \cdot s^{-1}$). Data are mean ± SE; ***$p < 0.001$, *$p < 0.05$ (Kruskal–Wallis test with

Steel's multiple comparisons in D, F, and G); ***$p < 0.001$ (one-way ANOVA with

Dunnett's multiple comparisons in E).

Fig. 4. Depletion of TMEM55B impairs lysosomal Ca^{2+} dynamics. (**A**) Schematic

representation of Ly-GG, a Ca^{2+} indicator with the G-GECO1 domain facing the cytosolic

side of the lysosomal membrane. (**B**) Confocal images of HeLa cells co-expressing Ly-

GG (green) and the lysosomal marker mCherry–LAMP1 (red). Scale bar, 10 μm. (**C**)

Time lapse analysis of Ly-GG fluorescence ($\Delta F/F_0$) in cells transfected with siControl

(gray, n = 29 fields), siTMEM55B#1 (red, n = 29), or siTMEM55B#2 (pink, n = 26). ATP

(200 μM) was added at t = 0 s; Data are mean ± SE and normalized by subtracting values

from ATP-free cells, in which fluorescence declined due to photobleaching. (**D, E**)

Quantification of Ly-GG slopes during 0–60 s (D) and 60–160 s (E) ($\Delta F/F_0 \cdot s^{-1}$). (**F**)

Schematic representation of CEPIA3mt, a Ca^{2+} indicator with the GCaMP2 domain

facing the mitochondrial matrix. (**G**) Confocal images of HeLa cells co-expressing

CEPIA3mt (green) and the mitochondrial marker Su9–RFP (red). Scale bar, 10 μm. (**H**)

Time lapse analysis of CEPIA3mt fluorescence ($\Delta F/F_0$) in cells transfected with control

siRNA (gray, n = 25 fields), siTMEM55B#1 (red, n = 26), or siTMEM55B#2 (pink, n =

24). ATP (200 μM) was added at t = 0 s; Data are mean ± SE. (**I, J**) Quantification of

CEPIA3mt slopes during 0–60 s (I) and 60–160 s (J) ($\Delta F/F_0 \cdot s^{-1}$). Data are mean ± SE;

***$p < 0.001$; (Kruskal–Wallis test with Steel's multiple comparisons in D and E); n.s.,

not significant (one-way ANOVA with Dunnett's multiple comparisons in I and J).

Fig. 5. Model of the mechanism where TMEM55B promotes lipolysis via lysosomal

acidification and calcium regulation at ER-Ly MCSs. ER protein PDZD8 is associated

with lysosome protein TMEM55B that promotes lysosomal acidification through

regulation of v-ATPase. Lysosomal acidification then promotes activity of lysosomal

acidic lipases that promote lipolysis and subsequently degradation of CEs to cholesterol,

as well as calcium channels that promote CICR. CICR at ER-Ly MCSs may also

contribute to lipolysis by fusion of LDs or LPs with lysosomes. ER, endoplasmic

reticulum; MCS, membrane contact site; v-ATPase, vacuolar-ATPase; TPC2, two-pore

channel 2; TRPML1, transient receptor potential mucolipin 1; RyR, ryanodine receptor;

IP$_3$R, inositol 1,4,5-trisphosphate receptor; CICR, Ca^{2+}-induced Ca^{2+} release; LD, lipid

droplet; LP, lipoprotein; CE, cholesteryl ester; Chol, cholesterol.

A

Gene symbol	Protein name	IF	Ly	Chol
Zfyve27	ISOFORM 1 OF ZINC FINGER FYVE DOMAIN-CONTAINING PROTEIN 27.	1.095		
Stmn2	STATHMIN-2.	0.500		
Vac14	PROTEIN VAC14 HOMOLOG.	0.324		
*Atp6v0d1	VACUOLAR PROTON PUMP SUBUNIT D 1.	0.200	✓	
Sh3gl1	ENDOPHILIN-A2.	0.125		
Gm364	SIMILAR TO TRANSMEMBRANE 9 SUPERFAMILY MEMBER 2.	0.111		
*Lamtor1	UPF0404 PROTEIN C11ORF59 HOMOLOG.	0.111	✓	✓
Tfrc	TRANSFERRIN RECEPTOR PROTEIN 1.	0.100		
*Tmem55b	TRANSMEMBRANE PROTEIN 55B.	0.083	✓	✓
Fig4	SAC DOMAIN-CONTAINING PROTEIN 3.	0.048		
Sun2	PROEIN UNC-84 HOMOLOG B.	0.032		
Vps35	VACUOLAR PROTEIN SORTING-ASSOCIATED PROTEIN 35.	0.026	✓	
Kif13a	KINESIN-LIKE PROTEIN KIF13A.	0.014		

B

C

D

Scale: 10 μm

E

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5