Trends of the Passerines avifauna are linked climatic variability and habitat management in a protected artificial wetland in southeast Iberia.

Ignacio García Peiró [1,2]

[1] **Academic affiliation**: Departamento de Ecología & Hidrología. Facultad de Biología. Universidad de Murcia. ES - Spain. Email: ignacio.peiro@yahoo.es

[2] **Corresponding author**: SEO/Birdlife-ALICANTE. C/ El Salvador, 17-4D. 03203 Elche (Alicante). ES-Spain. Email: Ignacio.peiro@yahoo.es

Abstract

The Mediterranean region is facing significant impacts on populations of biota, particularly birds, due to climate variability and habitat alteration. Consequently, the ecological diversity of many ecosystems has diminished, and the abundance of bird species has declined. This study employs a novel strategy in a seasonal freshwater artificial wetland in south-east Iberia establishing species-specific index trends for common passerine birds by comparing them with climate and habitat fragmentation data. A climatic overview indicates minimal trends towards warmer and wetter conditions in terms of temperature and rainfall over the 15-year study period. This pattern persisted over a 30-year period, resulting in the formation of a Mediterranean climatic sub regional scenario with warm, dry conditions that may influence the passerine assemblage in an analogous manner to that observed in other Mediterranean assemblages. One notable feature is that this assemblage in the 1990s was characterized by low natural diversity and specialization when it was a highly disturbed wetland. However, it became natural by the mid-2000s, with diversity values like those observed in other central Mediterranean wetlands. Trends used to evaluate species-specific patterns revealed that 98% of species had an unknown trend status during the study period, with a trend slope ranging from 0 to 15%, which was not substantially different than zero, probably due to the unequal standardization methodology and the short time studied. Further research using *Generalized Linear Mixed Models* (GLMM) revealed that precipitation and habitat alteration had the most detrimental effect on species. Of the observed species, 60% were affected by both factors, the majority of which were Palearctic (20%). The reduction of wintering areas and shifting migration routes may account for these effects. Further research is required on longer temporal data to clarify the evolution of the avifauna in this wetland.

Keywords: El Hondo Natural Park, habitat fragmentation, local climate, passerines, SE Iberia.

1.Introduction

The habitat and space provided by the Mediterranean are very favorable for some of the North African faunal traits that are exclusive to the southern Mediterranean, especially in the south-part of Iberia (Cheylan 1991, Martínez-Freiría et al. 2010, Husemann et al. 2014). In the class Aves, passerines represent a real hotspot of biodiversity (Covas & Blondel 1998). One of its most interesting features is the scarcity of Mediterranean bird species, with European and Palearctic elements making up the majority of the region's composition (Voous 1960, Blondel 1987). They are important elements of study of the evolutionary processes (Pons et al 2016).

Freshwater areas, particularly wetlands (marshes, lagoons, swamps), are declining in number and surface area due to significant water deficits caused by climate change since the turn of the century (Perennou et al. 2012; Lefebvre et al. 2019). The loss of Mediterranean wetlands has reduced the biological diversity of passerines (Paracuellos 1993, Peiró 2018), many of which are extremely important European-Turkestanian or Turkestanian-Mediterranean faunistic elements (Kirwan et al. 2010, Rustamov 2015, Peiró & Esteve 2002). These elements are vital for biodiversity conservation, ecological study, and understanding evolutionary processes since they play an important role in the early stages of plant successions (see Schulze-Hagen and Leisler 2011, Abramova 2019) and as secondary habitat (Bozó et al 2020).

The preceding assumptions suggest that the loss of specialists may enhance the extinction risk of ecosystems through biotic homogenization, with generalist winners replacing losers' specialists (McKinney & Lockwood 1999; Le Viol et al. 2012). This is primarily due to anthropogenic causes (Smart et al. 2006, De Victor et al. 2008a), while climate can play an influence (Davey et al. 2011).

The shift towards patchy wetlands (Paracuellos 2006 & 2008) has facilitated the proliferation of noteworthy Palearctic specialist songbirds across Europe, which congregate in reed-bed

masses. This phenomenon is exemplified by the Bearded Reedling *Panurus biarmicus* (Erard 1966, Kumberloeve 1968, Mead & Pearson 1974, Loison & Godin 1975, Olsson 1975, Antoniazza & Levéque 1977, Sluys 1982) is listed as Vulnerable in Spain (Vera 2021). In the study area, their small populations are situated at the southernmost extent of the species' distribution in Western Europe (Gosler & Mogyorósi, 1997) and may have dispersed to southern Iberian counterparts from the south-east of France, where this species was equally common and highly specialized since 1830, and where temperatures increased from 1960 (Galewski & Devictor, 2016). Currently, the species is almost extinct in the study area (Belenguer et al., 2016).

Inside this scenario, the analysis of bird trends using different monitoring data has been highly important to detect the impact of climate change on songbird trends, and applied in Western Europa (Reif et al. 2008, van Turnhout et al. 2008) and Eastern Europe (Zalakevičius 1998, Bozó et al. 2023, Tamás et al. 2023), and the results have shown to be highly variable (Jiguet et al. 2010), but few studies yet don't make use the TRIM methodology despite Bird Monitoring Schemes were set long ago (Vorisek & Marchant 2003, Vorirsek et al. 2008). However, it appears that little trend research has been carried out in wetlands (Žalakevičius & Švažas 2005, Van Turnhout et al. 2010) and many few using long-term ringing data and showing an overall decrease in the European avifauna (Petras & Vrezec 2022).

Based on the aforementioned arguments, the current study aims to: 1) provide a comprehensive overview of the climatology and zoogeography of the study area that has not been yet adequately covered (Peiró & Esteve 2001, Peiró 2006) and 2) address the question of how much abiotic (local climate) and biotic (anthropogenic) factors can influence the population trends of a Mediterranean wetland passerine assemblage subject to significant artificial management of water cycles and vegetation, as well as how these factors affected the homogenization of the assemblage and provide additional insights for the years to come.

2. Material and Methods

2.1. The study region, area and site.

With an estimated surface area of 260,000 Ha, the extreme southeastern Iberian span the easternmost portion of Iberia's Baetic southern plateau (Armada et al. 2018). Almería (37°58'N - 2°45'W), Murcia (37°58'N - 01°33'W), and Alicante (38°34' N - 00°48'W) are its three main provinces. The region has a semiarid Valencian climate (Costa 2019) and a Thermo-Mediterranean Iberian climate (Rivas-Martínez 1983, Rivas-Martinez & Armaiz 1984). The weather is seasonal, especially in the spring and autumn when it's rainier.

The study area "*El Hondo Natural Park*" (abbreviated EH, 38º'16'N - 00º 41'W) is in the middle-south of the province of Alicante. It is a 2.400-hectare inland freshwater swamp reservoir that was completed around the turn of the 20th century (Medina 1985) and designated as a regional natural park in 1988 (GVA 1988). Additionally, it was referred to as "El Hondo Swamp" in 1989 as RAMSAR site number 14 (Potenciano & Andueza 2006) and renamed as site number 455 (Bernués & Mateache 2020) because it maintains large populations of endangered waterbirds (Medina & Robledano 1995). A rich community of medium-to-small reed-bird species, some of which are currently threatened, and developed inside the eutrophic vegetation between man-made structures (Medina & García 1984, Peiró 1999).

The 800-meter line between two ponds is sited in the core area of the park and forms a succession of halophytic vegetation that reaches the park's boundaries (Peiró & Esteve 2001). The line included of saltmarshes of the species *Salicornia*, *Suaeda*, and *Arthrocnemum*, as well as common reeds *Phragmites australis* at various growth stages (Navarro 1985, Peiró 2006).

2.2. The Climatic data treatment.

I used annual mean temperatures and rainfall as indexes of abiotic factors or climatic
variability. Forecast variables were collected from the "*Estación Experimental Agraria of*
Elche (EEA-IVIA)", a meteorological station located 11 km northeast of the park at 38º14'N
1º25'W and 62 m above sea level. The data was gathered over a 15-year period (1991-2005,
period D1) in EEA notepads and online from IVIA http://riegos.ivia.es/datos-meteorologicos.
Long-time datasets for a 30-year period (1991-2020, period D2) were used in comparison
with the first dataset. The climatic data consisted of 1) annual and monthly average
temperatures (°C) and 2) overall annual and monthly precipitation (mm). To see the overall
summary of climate, additional variables were used: 1) mean annual temperature in spring
(March-May), summer (June-September), autumn (July-October), and winter (November-
February); and 2) overall annual winter (November-February).

The research area's bird seasonality is achieved, however certain climatic data, particularly
the annual temperature did not reach normality (Wilk's Shapiro test: W = 0.815, P = 0.006).
However, because rainfall varies so much, parametric tests based on the alternative
hypothesis of outlier existence were employed (Grubbs Test, Grubbs 1969). Standard
deviations (SD) were shown with the means. Two-tailed significance values were set at P <
0.05.

2.3. The Monitoring management

Strong management practices were implemented at the research site, mostly using human and
animal grazing and harvesting (wildfires, reed cutting, and weed-clearing base) (Peiró 2006).
By doing these steps, we were able to raise the reed's height and gradually remove its

structure during the study period. The yearly average height of reeds was selected as a
measure of habitat change brought about by management. From 1991 to 2005, I measured the
height of the reed beds on both sides of the study path in order to collect field data for
assessing these changes. One to four months following each management intervention, I
randomly picked one to twenty reed stems and used a meter rule to measure the height (in
cm) from the base to the seeds. Management actions were conducted almost every year
during the study period and consisted mainly in harvesting, grazing and wildfires of the
reedbed path.

2.4. The Bird's data recording and monitoring

First, throughout annual surveys (1991-2005, N = 15, period D1), the author implemented a
Flexible Constant Effort Site mist-netting program in the Park's core region with varied
ringing visits, which typically lasted four hours after sunrise. Only comparisons of
biodiversity were conducted using the incomplete datasets from 2006 to 2012 (N = 7, period
D3). According to Peterson et al. (1977) and Svensson (1992), I metal-ringed every bird
(Ministery of Environment-MADRID, SPAIN), measured each bird, and then released them
all to the same location. Nearly every season from D1 was covered by the ringing and
handling in D1 and D3 (1) (N = 15 years). I attempted to evaluate potential assemblage
changes over long periods of time using some standardized ringing data from period D3.

I mostly used period 1 for data monitoring. By catching them, ringed birds were measured
and standardized. Standardization procedures consisted by dividing the number of individuals
by trapping effort [(individuals/(m2·hour)) ·1000]. This was done differently for each of the
following annual presences at the site: breeding (March-October, B), wintering (November-
February; W), sedentary (January-December, S), spring migrant (March-May, SM), autumn
migrant (July-October, AM), and occasional (O) (Table 1). Sum spring-autumn trapping
effort (SO) was used to consider birds with both migratory statuses. Wintering birds with
spring and autumn migratory populations were taken into consideration with an overall
November-February trapping effort, while birds mostly in breeding status but with spring and
autumn migrant populations were taken into consideration with an overall March-October
trapping effort. The annual and seasonal abundances and trapping effort by species are

presented in Table 1. According to Poulin et al. (2000), the normalized breeding trapping

effort used to measure the assemblage of reed bed species was set at 1.300 m2/hour, which

satisfies the requirements at least during the study site's breeding seasons.

I performed a checklist of passerines from EH that were initially categorized into faunistic

elements by Voous (1960) and listed, classified, and given English names by Rouco et al.

(2022). The plotting of this general template was the same in Tables 1-4. I calculated the

individual biodiversity distributions (Shannon's Index) between periods (D1 and D3) to see

whether or not the bird assemblage was similarly distributed between periods (Van Turnhout

et al 2007). This allowed me to determine whether the bird assemblage may have altered

between then. Since bird diversity between assemblages did not meet normality (Shapiro

Wilk's tests: D1 = 0.585; P < 0.001; D3 = 0.008; p < 0.001), median diversities were

evaluated using the non-parametric Mann-Wittney-Wilcoxon rank sum tests (MWW, Sokal &

Rolhf 1981). Because the relative number of specialists and generalist species is likely to

affect community functions (Julliard et al. 2006), I attempt to provide a suitable index of

specialisation of each species within the assemblage using the species-specific specialisation

index (SSI) derived from Morelli (Appendix 1 in Morelli et al. 2020). According to DeVictor

et al. (2008b), this index is based on how frequently each species occurs in each type of

habitat or land use. As a result, each bird species' specialization index (SSI) may have values

ranging from 0 to 1, which correspond to low and high levels of specialization, respectively.

The specific weighted SSI of the assemblage in each time was calculated by multiplying the

SSI index for each species by the number of individuals. The abundance of each species, as

indicated by the number of individuals, is taken into consideration when computing the

ensemble of community specialization in this index. In order to determine whether the two

periods were equally distributed on time and because the SSI between assemblages in the

period D1 and D3 did not meet normality (SSI-1 = 0.585; P < 0.001; SSI-3 = 0.008; p <

0.001), the median weighted SSI for D1 and D2 were compared using the Mann Wittney

Wilcoxon tests for two independent samples.

2.5. The Population trends and climatic and management effects

For analyzing species-specific trends, I employed Trends & Indexes for Monitoring TRIM

software (Pannekoek & Van-Strien 2001), which was developed from R-Project v. 44.2

software (R-Core Team 2020) by the "rtrim" package (Bogaart et al. 2024). I implemented

"rtrim" using Model 2 based in Poisson regression method (Van-Strien et al. 2004,
Pannekoek & Van-Strien 2001) which accounts for only one slope in the overall model (Ter
Braak et al. 1994). To indexing the bird abundances data and run the TRIM they were
transformed into integers. Null zero assignments were one-added and missing values were
transformed into not analyzed NA values and afterwards imputed (Ter Braak et al. 1994). See
Table 2 for complete explanations. Using the "glm2" package (Marschner 2011), I employed
Generalized Linear Mixed Models (GLMM; McCullagh & Nelder 1983) to find some
relationships between climate and habitat transformation. The response variable was the
abundance of each bird, and the predictive variables were annual mean temperatures, annual
total precipitation, and reed height. For more details on the analysis, refer to Table 3.

3. Results

3.1. The overall climatic sketch and trends of weather variables.

The average temperature during the coldest month (January, period D2) was 19.4 ± 1.54,
while the average rainfall during the warmest month (July) was 1.4 ± 2.99 mm. The total
annual precipitation averaged at 241.8 ± 88.14 mm, with an average annual temperature of
$19.4° \pm 0.96$ recorded for the period of 1991-2020. Over the period D1 annual temperatures in
EH decreased (Temperature = - 0.029·Year + 76.399; $r^2 = 0.018$, $F_{1,13} = 0.2372$, $P = 0.634$)
and rainfall increased (Precipitation = 1.920·Year – 3610.731, $r^2 = 0.016$, $F_{1,13} = 0.219$, $P =$
0.648) these trends were similar in D2 (Temperature: Pearson's $r = - 0.077$, $P = 0.685$;
Rainfall: $r = 0.286$; $P = 0.126$). Large scale temperatures were influenced spring temperatures
since they got higher determination coefficient on the linear relationships ($r^2 = 0.723$, $P <$
0.001). Likewise, rainfall in current autumn was the greatest driver of overall precipitation (r^2
$= 0.521$, $P < 0.001$). Punctual storminess may lead extreme rainfalls as occur in other central
Mediterranean areas (Diodato & Bellocchi 2010) and determinates the highest variability of
rainfall (CV = 35%) compared to temperatures (CV = 5%) since outliers on the large-scale
precipitations in 2019 (469 mm; Grubbs Tests, G = 2.509, U = 0.776, P = 0.124) could be
responsible for the positive correlations of rainfall with year so this scenario represents an
apparent dry and medium- warm cycle in the study area during the last century (1991-2020)
which could extend to the southern subregion of Alicante and subsequently, due to climate

change, rainfalls became shorter but more intense in an episodic manner (Diodato et al 2011)

so in climatic terms, conformed a posterior cycle (1921-1924) in a wetter and warmer period.

This is coincident with the great local climate variability in Mediterranean areas depicted by

Xoplaki (2002).

3.2. The Zoogeography and trends in the avifauna.

Seventy percent of the assemblage's species (N = 38) were of Palearctic, European, and

European-Turkestanian origin. A reminiscence was of Turkestanian-Mediterranean origin. An

annual trapping effort (r = 0.800, P < 0.001) that grew weakly on time (r = 0.073, P = 0.795)

had a considerable influence on the yearly total number of species, which declined weakly on

time (r = - 0.092, P = 0.734). The sole species with European status, the Common Chaffinch

Fringilla coelebs, saw a significant rise (42%) over the 1991–2005 study period, while the

Eurasian Penduline tit *Remiz pendulinus* saw a reduction of 1.4%.

During the study period, the remaining assemblage had an ambiguous trend status with a

positive trend slope ranging from 0 to 15%, primarily non-significantly. Even though the

overall diversity was 13% greater in the following years (1991-2005: 1.875 nats, 2006-2012:

2.163 nats), the median diversity across periods was statistically significant (D1: 0.010 nats,

D3: 0.029 nats; Wilcoxon-Mann-Wittney Test; W = 351.5, P = 0.012). Additionally, the

community became increasingly less specialized over time as a result of the overall SSI index

declining by 82% (D1: 1070.3, D3: 193.2) as a result of 32% of species (N = 12) being lost

and three species being added: two Mediterranean, the Woodchat shrike *Lanius senator* and

the Spectacled warbler *Sylvia conspicillata* and one Palearctic, the Western yellow wagtail *Motacilla flava* (Table 3).

3.3. The relationships with climate and habitat transformation.

Simple (linear) models that primarily used one variable to explain annual abundance did not significantly fitted most species (Table 4). 78% of the species in the collection were influenced by climate factors, particularly temperature and precipitation, which mostly determined the species' interannual abundance. Most of these species were Palearctic (16%) and Euro-Turquestanian (21%). 29% of species were impacted by habitat change, primarily in the Palearctic (13%).

4. Discussion

Southeastern Iberia experiences a Mediterranean climate characterized by hot, dry summers and mild, wet winters. The primary meteorological factors influencing this climate include: 1.- Temperature variability: rising temperatures have been observed in the Mediterranean region, significantly affecting local ecosystems and species distributions. For instance, the abundance of migratory birds like Savi's warbler *Locustella luscinioides* has shown a positive correlation with intra-annual temperatures, indicating that temperature fluctuations can impact breeding and migratory patterns (Peiró 2024), 2.- Precipitation patterns: precipitation in SE Iberia is characterized by seasonal variability, with significant impacts on local flora and fauna. The region has experienced decreasing precipitation trends, which can lead to drought conditions, particularly in summer. This has been linked to the adaptation and migration of various species, as well as changes in vegetation dynamics (Carrasco et al. 2018, Alambiaga et al. 2021) and 3.- Climatic oscillations: the climate dynamics in SE Iberia are also influenced by larger climatic patterns such as the North Atlantic Oscillation (NAO) and the El Niño Southern Oscillation (ENSO). These oscillations can lead to variations in precipitation and temperature, affecting the overall climate stability in the region. These factors collectively shape the unique climatic conditions of southeastern Iberia, influencing biodiversity, species distribution, and ecological interactions.

The findings indicate that the majority of the species in EH are generalists, with only a limited number of reed-marsh specialists (*Acrocephalus*, *Cettia*, *Locustella*, Cramp 1992, Kennerley & Pearson 2010) making up 25% of the bird assemblage. The arrival of four species of passerines, not specialists, was the primary cause of the diversity increaseThe Zoogeography of EH is primarily Palearctic, European and European-Turkestan, owing the wintering of Palearctic wintering heavyweights of Common Chiffchaff *Phylloscopus collybita* (PC) and the summer European-Turkestan Eurasian Reed Warbler *Acrocephalus scirpaceus* (AS). These two (AS and PC) are quantitatively the bulk species, increasing in abundance at rates of 5–14% each year, respectively. However, they are adversely affected by habitat fragmentation and annual temperature increases. There may be a decline in the PC population but a rise in AS in the future because to rising temperatures and habitat resiliency.

In the 1960s, Turkestan-Mediterranean elements, such as scrub warblers like the Sardinian Warbler *Sylvia melanocephala* (SM) and two reed-marsh specialists, the Cetti's Warbler *Cettia cetti* (CC) and the Moustached warbler *Acrocephalus melanopogon* (AM), spread towards central Europe from southern Europe and southwest Asia. More recently, in the case of AM, they likely came from wintering birds of populations of central and eastern European origin (Bozó et al. 2023), and there may have been an over-dispersal of breeding populations from the eastern Iberian regions to the south (Ceresa et al. 2015).

Diversity in 1991-2005 in EH falls well below the values detected in other Mediterranean wetlands (Malavasi et al. 2009), indicating that the study area is human-disturbed and is highly below the range of diversity of natural or semi-natural central Mediterranean wetlands (2.8-2.5 nats, Malavasi et al. 2009). The huge increase in diversity from 2006 to 2012 is indicative of transformation towards a natural wetland.

Denser reeds and less rainfall appear to be more conducive to the population dynamics of the earlier specialists, whose abundances were adversely affected by habitat management and rainfall, respectively. We could anticipate a notable rise in the population of Turkestan-Mediterranean elements (AM & CC) in this ecosystem in the event of future warming, less precipitation, and the restoration of the natural habitat in EH. Since reedbed alterations (clearing) have a detrimental impact on SM, boosting habitat densification would have a significant favourable impact on their abundances.

The Reed bunting, another reed marsh specialist, is only found in EH as an overwintering
bird and has not been breeding there since the 1970-80s (Medina & Garcia 1984, Medina
1985). It is likely that the earliest breeders here were of the Iberian subspecies *witherby*,
which is found in the widely recognised breeding nucleus in the central (Alambiaga et al.
2021) and eastern marshlands of Iberia (Carrasco et al. 2018).Simple (linear) models that
explained the annual abundances on climate and habitat alteration were mostly not significant
and this is in accordance with the findings of Jiguet et al (2010) which hypothesizes that
trends in European abundances do not meet lineal relationships. According to Leviol et al
(20102), European bird communities are dominated by native generalist species, indicating a
significant homogenization process in different regions. The authors propose that this rapid
and non-random change in community composition is due to anthropogenic activities, but the
effects of climate change are also very important (Davey et al 2011).

Conclussions

Future research during longer cycles is needed to ascertain the possible changes in the
species' composition due to climate warming and management use. To determine the
potential changes in the species' composition brought on by climate change and management
use, more research is required.

335
336
337　**References**

338
Abramova, I. V. 2023. Dynamics of the Abundance of Bird Species during Succession of Oak
Forests in Southwestern Belarus. Biology Bulletin 50: 1864-1874.

Alambiaga, I., Carrasco, M., Ruiz, C., Mesquita-Joanes, F.& Monrós, J. S. 2021. Population
trends and habitat selection of threatened marsh passerines in a protected Mediterranean
wetland. Avian Conserv. Ecol. 16: 23.

Antoniazza, M & Levéque R. 1977. La Mésange à moustaches (Panurus biarmicus), une
nouvelle espéce nicheuse de l'avifaune suisse. Nos Oiseaux. 34: 93-110.

Armada, X. L., Grau-Mira, I., Haselgrove, C., Rebay-Salisbury, K. & Wells, P. 2018.The
Iberian Peninsula. The Oxford Handbook of the European Iron Age, pp 1-43.

Belenguer, R., López Iborra, G. M., Dies, J. I. & Castany i Àlvaro, J. 2016. Dramatic decline

of the bearded reedling, Panurus biarmicus, in Spanish Mediterranean wetlands. Animal
Biodiversity and Conservation. 39:17–27.

Bernues, M & Mateache, P. Ficha Informativa RAMSAR EL Hondo sitio número 455, 2020,
https://rsis.ramsar.org/ris/455

Blondel, J. 1987. History and development of bird faunas in the Mediterranean
region. *Travaux et Documents des Laboratoires de Géologie de Lyon* 99 : 231-238.

Bogaart P, Loo, V. M & Pannekoek J. 2024. rtrim: Trends and Indices for Monitoring Data.
R-package version 2.3.0.

Bonham, P. F. & Robertson, J. C. M. 1975. The spread of Cetti's Warbler in north-west
Europe. British Birds. 68: 393-408.

Bozó, L. 2020.The role of reedbeds in secondary habitats during the migration and breeding
of reed warblers. Ornis Hungarica. 28:76-91.

Bozó, L., Németh, Á., & Csörgő, T. 2023. Long-term population changes of the Moustached
Warbler in a Central Hungarian wetland habitat. Ornis Hungarica 31: 89-98.

Carrasco, L., Vera, P., Belda, E. J., & Monrós, J. S. 2018. Combining remote sensing and
field mapping methods to study the vegetation dynamics within a coastal wetland and
determine the habitat effects of a threatened bird species (Emberiza schoeniclus witherbyi).
Journal of Coastal Research 34: 42-49.

Ceresa, F., Belda, E. J., Kvist, L., Rguibi Idrissi, H., & Monrós, J. S. 2015. Does
fragmentation of wetlands affect gene flow in sympatric Acrocephalus warblers with different
migration strategies? Journal of Avian Biology 46: 577-588.

Cheylan, G. 1991. Patterns of Pleistocene turnover, current distribution and speciation among
Mediterranean mammals. Biogeography of Mediterranean invasions pp 227-262.

Costa, M., 2019. Pisos bioclimáticos y series de vegetación en el área valenciana. Cuadernos
de Geografía de la Universidad de València.

Covas, R., & Blondel, J. 1998. Biogeography and history of the Mediterranean bird
fauna. Ibis140: 395-407.

Cramp, S Ed. 1992. The Birds of the Western Palearctic. Vol VI. Oxford University Press.
Oxford.

Davey, C. M., Chamberlain, D. E., Newson, S. E., Noble, D. G. & Johnston, A. 2011 Rise of
the Generalists: evidence for climate driven homogenization in avian communities. Glob.
Ecol. Biogeogr. 21: 568–578.

Devictor, V., Julliard, R., Clavel, J., Jiguet, F., Lee, A. & Couvet, D. 2008. Functional biotic

homogenization of bird communities in disturbed landscapes. Glob. Ecol. Biogeogr. 17: 252–
261.

Devictor, V., Julliard, R., & Jiguet, F. 2008b. Distribution of specialist and generalist species
along spatial gradients of habitat disturbance and fragmentation. Oikos 117: 507–514.

Devictor, V., Clavel, J., Julliard, R., Lavergne, S., Mouillot, D., Thuiller, W & Mouquet, N.
2010. Defining and measuring ecological specialization. Journal of Applied Ecology 47: 15-
25.

Diodato, N., & Bellocchi, G. 2010. Storminess and environmental changes in the
Mediterranean Central Area. Earth Interactions, 14(5), 1-16.

Diodato, N., Bellocchi, G., Romano, N., & Chirico, G. B. 2011. How the aggressiveness of
rainfalls in the Mediterranean lands is enhanced by climate change. Climatic Change 108:
591-599.

Erard Ch. 1966. Invasion de Mèsanges à moustaches (Panurus biarmicus L.). Alauda 34 :
240-242.

Forsman, J. T., & Mönkkönen, M. 2003. The role of climate in limiting European resident
bird populations. *Journal of Biogeography 30*: 55-70.

Galewski, T., & Devictor, V. 2016. When common birds became rare: historical records shed
light on long-term responses of bird communities to global change in the largest wetland of
France. PLoS One 11: e0165542.

Gosler A & Mogyorósi D. 1997. Bearded Tit. In Hagenmeijer EJ. and Blair MJ. (eds). The
EBCC Atlas of European Breeding Birds: Their Distribution and Abundance. TAD Poyser,
London. pp 628–629.

Grubbs, F. E. 1969. Procedures for detecting outlying observations in
samples. *Technometrics*, *11*: 1-21.

GVA. 1988. Decreto 187/1988, de 12 de diciembre, del Consell de la Generalitat Valenciana,
de declaración de Paraje Natural de la Comunidad Valenciana de El Hondo (DOGV 991 -
25.01.1989).

Husemann, M., Schmitt, T., Zachos, F. E., Ulrich, W. & Habel, J. C. 2014. Palaearctic
biogeography revisited: evidence for the existence of a North African refugium for Western
Palaearctic biota. *Journal of Biogeography* 41: 81-94.

Jiguet, F., Gregory, R. D., Devictor, V., Green, R. E., Vorisek, P., Van Strien, A., & Couvet, D.
2010. Population trends of European common birds are predicted by characteristics of their
climatic niche. *Global change biology* 16*: 497-505.

Jiguet, F., Devictor, V., Ottvall, R., Van Turnhout, C., Van der Jeugd, H., & Lindström, Å. 2010. Bird population trends are linearly affected by climate change along species thermal ranges. Proceedings of the Royal Society B: Biological Sciences 277: 3601-3608.

Julliard, R., Clavel, J., Devictor, V., Jiguet, F., & Couvet, D. 2006 Spatial segregation of specialists and generalists in bird communities. *Ecology letters* 9: 1237-1244.

Kennerley, P., & Pearson, D. 2010. Reed and bush warblers. A&C Black.

Kirwan, G., Demirci, B., Welch, H., Boyla, K., Özen, M., Castell, P., & Marlow, T. 2010. *The birds of Turkey*. Bloomsbury Publishing.

Kumberloeve H. 1968. Sur la situation subsepecifique des Mesànges à moustaches (Panurus biarmicus) en Asia Minuere et les alentours. Aves 6 : 61.

Le Viol, I., Jiguet, F., Brotons, L., Herrando, S., Lindström, Å., Pearce-Higgins, J. W, & Devictor, V. 2012. More and more generalists: two decades of changes in the European avifauna. *Biology letters* 8: 780-782.

Lefebvre, G., Redmond, L., Germain, C., Palazzi, E., Terzago, S., Willm, L., & Poulin, B. 2019. Predicting the vulnerability of seasonally flooded wetlands to climate change across the Mediterranean Basin. Science of the Total Environment, 692, 546-555.

Leisler, B., & Catchpole, C. K. 1992. The evolution of polygamy in European reed warblers of the genus Acrocephalus: a comparative approach. *Ethology ecology & evolution 4* :225-243.

Loison M. & Godin J. 1975. Presence estivale de la Mesànge à moustaches (Panurus biarmicus) en Hainaut occidental. Aves 12: 31-32.

López, G & Monrós, J. 2004. Bigotudo, *Panurus biarmicus*. In Madroño, A., González, C & Atienza J.C (Eds): Libro Rojo de las Aves de España. Dirección General para la Biodiversidad-SEO/Bidlife. Madrid, pp 341-344.

MacKenzie R., Warren. M. 2015. Topic B4: Wetlands and Ecosystem Services. PPT. In: SWAMP toolbox: Theme B, Section B4, 2015; pp. 53.

Malavasi, R., Battisti, C., & Carpaneto, G. M. 2009. Seasonal bird assemblages in a Mediterranean patchy wetland: corroborating the intermediate disturbance hypothesis. *Polish Journal of Ecology* 57: 171-179.

Marschner, I. C. 2011. glm2: Fitting generalized linear models with convergence problems. The R Journal 3: 12-15

Martínez-Freiría, F., Lizana, M., do Amaral, J. P., & Brito, J. C. 2010. Spatial and temporal segregation allows coexistence in a hybrid zone among two Mediterranean vipers (Vipera aspis and V. latastei). Amphibia-Reptilia, 31: 195-212.

McCullagh, P. & Nelder, J. A.1983. Generalized Linear Models, Chapman & Hall.

McKinney, M. L. & Lockwood, J. L. 1999. Biotic homogenization: a few winners replacing
many losers in the next mass extinction. Trends Ecol. Evol. 14: 450–453.

Mead CJ. and Pearson DJ. 1974. Bearded Reedling populations in England and Holland. Bird
Study 21 : 211-214.

Medina, J. D. N., & García, J. N. 1984. La avifauna de los embalses de «el hondo»
(Alicante). Mediterránea: Serie de Estudios Biológicos 7: 117-137.

Medina, J.D.N. 1985 *Estudio Ornitológico de "El Hondo"*. Caja de Ahorros de Meditarraneo.
Alicante.

Medina, J. D. N & Robledano, F. 1995. La Cerceta Pardilla *Marmaronetta angustirostris* en
España. ICONA-MAPA Colección Técnica. Instituto Nacional para la Conservación de la
Naturaleza, Madrid, Spain.

Møller, A. P., Rubolini, D., & Lehikoinen, E. 2008. Populations of migratory bird species that
did not show a phenological response to climate change are declining. Proceedings of the
National Academy of Sciences 105: 16195-16200.

Morelli, F., Benedetti, Y., & Callaghan, C. T. 2020. Ecological specialization and population
trends in European breeding birds. Global Ecology and Conservation 22: e00996.

Morelli, F., Benedetti, Y., Møller, A. P., & Fuller, R. A. 2019. Measuring avian specialization.
Ecology and Evolution 9: 8378-8386.

Olsson V. 1975. Bearded Reedling populations in Scandinavia. Bird Study 22:116-117.

Pannekoek, J & Van-Strien, A. 2001. TRIM 3 manual (trends & indices for monitoring data).
http://www. ebcc. info/trim. Html.

Paracuellos, M. 1993. Fenología anual de la ornitofauna en las Salinas de Guardias Viejas
(Almería). Calidad ornítica. Alytes 6: 317-333.

Paracuellos, M. 1994. Dinámica anual de la comunidad de Paseriformes en un saladar litoral
del sudeste ibérico. Doñana, Acta Vertebrata 21: 119-130.

Paracuellos, M. 1996. Dinámica anual de la comunidad de Paseriformes en carrizales costeros
del sudeste ibérico. Doñana, Acta Vertebrata 23: 33-44.

Paracuellos, M. 2006. Relationships of songbird occupation with habitat configuration and
bird abundance in patchy reed beds. Ardea 94: 87-98

Paracuellos, M. 2008. Effects of long-term habitat fragmentation on a wetland bird
community. *Revue d'Ecologie, Terre et Vie* 63 : 227-238.

Peiro, I. G. 2024. Local climate variability, phenology and morphological traits of the long-
distance migrants Savis warbler Locustella luscinioides and Sedge warbler Acrocephalus
schoenobaenus in reedbeds of a man-made wetland in SE Iberia. *bioRxiv*, 2024-05.

Peiró, I. G & Esteve, M.A. 2002. Ecología de los paseriformes del carrizal del Parque Natural
del Hondo. Inst.Est. Juan Gil Albert. Alicante.

Peiró, I. G. 2018. The biodiversity of small passerine birds at wetlands: species loss and
effects of climate change. Biodiversity International Journal 2: 297-301.

Peiró, I.G. 1999. Passeriformes del Parque Natural del Hondo (España)-Passeriformes of the
El Hondo Natural Park (Spain). Sehumed 10:7-8.

Peiró, I.G. 2006. Estudios ornitológicos aplicados a la gestión del hábitat de los paseriformes
del carrizal del Parque Natural de El Fondo (Alicante). PhD Thesis. University of Murcia.

Perennou, C., Beltrame, C., Guelmami, A., Tomàs Vives, P, & Caessteker, P. 2012. Existing
areas and past changes of wetland extent in the Mediterranean region: an overview. *Ecologia*
mediterranea 38: 53-66.

Peterson, R., Mountfort, G. & Hollom, P. 1977. Guía de campo de las aves de España y de
Europa. Ed. *Omega*. Barcelona.

Petras, T& Vrezec, A. 2022. Long-Term Ringing Data on Migrating Passerines Reveal
Overall Avian Decline in Europe. Diversity 14: 905.

Pons, J. M., Thibault, J. C., Aymí, R., Grussu, M., Muntaner, J., Olioso, G., & Fuchs, J. 2016.
The role of western Mediterranean islands in the evolutionary diversification of the spotted
flycatcher Muscicapa striata, a long□distance migratory passerine species. *Journal of Avian*
Biology 47: 386-398.

Potenciano, A.B & Andueza, I.L. 2006. *Ficha Informativa de humedales de RAMSAR (F.I.R).*
El Hondo.

Poulin, B., Lefebvre, G. & Pilard, P. 2000. Quantifying the breeding assemblage of reedbed
passerines with mist-net and point-count surveys. Journal of Field Ornithology 71: 443-454.

R Core-Team 2020. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Reif, J., Voříšek, P., Šťastný, K., Koschová, M., & Bejček, V. 2008. The impact of climate
change on long□term population trends of birds in a central European country. Animal
Conservation 11: 412-421.

Rivas-Martínez, S. 1983. Pisos bioclimáticos de España. *Lazaroa* 5: 33-43.

Rivas-Martinez, S., & Armaiz, C. 1984. Bioclimatologia y vegetacion en la Peninsula Ibérica. Bulletin de La Société Botanique de France. Actualités Botaniques 131: 110–120.

Robinson, R. A., Freeman, S. N., Balmer, D. E., & Grantham, M. J. 2007. Cetti's Warbler Cettia cetti: analysis of an expanding population. *Bird Study* 54: 230-235.

Rouco, M., Copete, J. L., de Juana, E., Gil-Velasco, M., Lorenzo, J. A., Martín, M., & Santos, D. M. 2019. Lista de las aves de España. *SEOBirdlife. Madrid.*

Rustamov, E. A. 2015. An annotated checklist of the birds of Turkmenistan. *Sandgrouse*, *37*(1), 28-56.

Schulze-Hagen, K., & Leisler, B. 2011. Reed warblers: diversity in a uniform bird family. *Zeist Netherlands: KNNV Uitgeverij.*

Sluys K.1982. Bearded Tits (Panurus biarmicus) in the Netherlands and England: comments on hibridization theory. J. Orn. 123:175-182.

Smart, S. M.,Thompson, K., Marrs, R. H., Le Duc, M. G., Maskell, L. C., & Firbank, L. G. 2006 Biotic homogenization and changes in species diversity across human-modified ecosystems. Proceedings of the Royal Society B: Biological Sciences, 273: 2659-2665.

Sokal, R. R & Rolhf, F.J. 1981 *Biometry*. Freeman & Co. San Francisco.

Svensson, L. 1992 Identification guide to European passerines. Lars Svensson, Stockholm.

Tamás E.A., Kalocsa B., Németh A., Marinov M. 2023. Survey of the Moustached warbler (Acrocephalus melanopogon) populations and migration routes in Europe, role of the Black Sea population, pp. 96-104. In Gastescu, P., Bretcan, P. (edit., 2023), Water resources and wetlands, 6th International Hybrid Conference Water resources and wetlands, 13-17 September 2023, Tulcea (Romania).

Ten Brink P, Badura T, Farmer A, Russi D. 2012. The Economics of Ecosystem and Biodiversity for Water and Wetlands: A Briefing Note. Institute for European Environmental Policy, London.

Ter Braak, C.J.F., Van Strien, A.J. Meijer, R & Verstrael,T.J. 1994. Analysis of monitoring data with many missing values: which method? In: E.J.M. Hagemeijer & T.J. Verstrael (eds.), 1994. Bird Numbers 1992 Distribution, monitoring and ecological aspects. Proceedings of the 12th International Conference of IBCC and EOAC, Noordwijkerhout, The Netherlands. Statistics Netherlands, Voorburg/Heerlen & SOVON, Beek-Ubbergen, pp. 663-673.

Van Turnhout, C. V., Willems, F., Plate, C., Strien, A. V., Teunissen, W., Dijk, A. V., & Foppen, R. P. B. 2008. Monitoring common and scarce breeding birds in the Netherlands: applying a post-hoc stratification and weighting procedure to obtain less biased population trends. Revista Catalana d'Ornitologia, 24: 15-29.

Van Strien, A., Pannekoek, J. Hagemeijer, W & Verstrael, T 2004. A log-linear Poisson
regression method to analyze bird monitoring data. In: Anselin, A. (ed.) Bird Numbers 1995,
Proceedings of the International Conference and 13th Meeting of the European Bird Census
Council, Pärnu, Estonia. Bird Census News 13: 33-39

Van Turnhout, C. A., Foppen, R. P., Leuven, R. S., Siepel, H., & Esselink, H. 2007. Scale-
dependent homogenization: changes in breeding bird diversity in the Netherlands over a 25-
year period. Biological Conservation, 134: 505-516.

Van Turnhout, C. A. M., Hagemeijer, E. J. M., & Foppen, R. P. B. 2010 Long-term population
developments in typical marshland birds in The Netherlands. Ardea, 98: 283-299.

Vera, P. 2021. Bigotudo, *Panurus biarmicus*. In López-Jiménez, N. (Ed.): Libro Rojo de las
Aves de España, pp. 125-136. SEO/BirdLife. Madrid, pp 576-580.

Voous, K. H. 1960. *Atlas of European birds*. Thomas Nelson & Sons Limited.

Voous, K. H. 1963. The concept of faunal elements or faunal types. Proc. Int. Orn, Congress,
13: 1104- 1108.

Vorisek, P. & Marchant J.H. 2003. Review of largescale generic population monitoring
schemes in Europe. Bird Census News 16: 14–38.

Vorisek, P., Gregory, R. D., Van Strien, A. J., & Meyling, A. G. 2008. Population trends of 48
common terrestrial bird species in Europe: results from the Pan-European Common Bird
Monitoring Scheme. Revista Catalana d'Ornitologia 24: 4-14.

Wickham H, François R, Henry L, Müller K, Vaughan, D. 2023. dplyr: A Grammar of Data
Manipulation. R package version 1.1.4. https://CRAN.R-project.org/package=dplyr.

Xoplaki, E. 2002. Climate variability over the Mediterranean. PhD Thesis. University of
Bern.

Žalakevičius, M & Švažas, S. 2005. Global climate change and its impact on wetlands and
waterbird populations. Acta Zoologica Lituanica, 15: 211-217.

Zalakevičius, M. 1998. Global climate change impact on breeding bird species and their
populations in Lithuania: status, population estimate and distribution trends. Acta Zoologica
Lituanica, 8: 3-16.

Table 1. List of species ringed in EH in 1991-2005. Standardized abundances of each species in mist nets [individuals/m^{2}net* hour) *1000] are plotted. Abundances were calculated differently according to the annual occurrence of the trapping effort at the site. Breeding (March-October, B), wintering (November-February; W), sedentary (January-December, S), spring migrant (March-May, SM), autumn migrant (July-October, AM), occasional (O). Birds with both migratory phenology were considered as an adding spring-autumn trapping effort. Birds with a mainly breeding status but with spring and autumn migrant populations, marked as (*), were considered with an overall trapping effort March-October and wintering birds with spring and autumn migrant populations, marked as (*) were considered with a summed trapping effort November-February and June-September. Summed total abundances between 1991 and 2005 are also plotted for each species. Abundances were transformed into indices for TRIM analyses. Birds were classified and English named according to Rouco et al (2022). Faunal elements follow Voous (1960).

Common name	Scientific name	Faunal element	Annual status	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Barn Swallow	Hirundo rustica	Holarctic	Breeding	0,227	0,087	0,000	0,063	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,213
Cetti's warbler	Cettia cetti	Turkestanian-Mediterranean	Wintering	20,741	9,003	3,055	9,396	4,770	12,364	16,892	7,927	77,413	29,037	0,000	0,000	7,368	0,000	11,615
Southern Grey Shrike	Lanius meridionalis	Holarctic	Occasional	0,000	0,024	0,000	0,000	0,000	0,000	0,101	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Willow Warbler	Phylloscopus trochilus	Palaearctic	Migrant	0,000	0,033	0,143	0,142	0,137	0,000	0,287	0,573	0,186	0,000	0,000	0,000	0,000	0,000	0,498
Common Chiffchaff	Phylloscopus collybita	Palaearctic	Wintering	0,454	4,637	5,185	3,833	9,142	14,013	18,070	82,289	14,742	11,973	0,890	1,088	4,586	0,000	11,700
Great Reed Warbler	Acrocephalus arundinaceus	European-Turkestanian	Breeding*	1,818	0,671	1,246	1,068	0,104	0,643	1,794	2,939	1,421	0,000	0,000	4,549	3,583	1,372	5,743
Moustached Warbler	Acrocephalus melanopogon	Turkestanian-Mediterranean	Wintering	5,926	7,325	1,222	4,027	1,101	4,755	2,413	4,844	18,215	13,754	0,000	0,000	7,368	0,000	12,168
Aquatic warbler	Acrocephalus paludicola	European	Migrant	0,000	0,033	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Sedge Warbler	Acrocephalus schoenobaenus	European-Turkestanian	Migrant	0,000	0,133	0,071	0,000	0,000	0,159	0,000	0,573	0,558	0,273	0,000	0,106	0,000	0,000	0,000
Common Reed Warbler	Acrocephalus scirpaceus	European-Turkestanian	Breeding *	6,363	8,136	7,677	8,042	12,779	11,699	18,583	28,654	21,846	11,464	0,890	23,438	17,125	10,837	41,906
Common Grasshopper Warbler	Locustella naevia	European-Turkestanian	Migrant	0,000	0,033	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Savi's Warbler	Locustella luscinioides	European-Turkestanian	Migrant	0,000	0,066	0,167	0,000	0,000	0,143	0,000	0,000	0,000	0,000	0,000	0,106	0,095	0,000	0,249
Zitting Cisticola	Cisticola juncidis	Indian-African	Breeding	0,000	0,321	0,034	0,000	0,104	0,129	0,000	2,204	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Great Tit	Parus major	Palaearctic	Occasional	0,000	0,049	0,047	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Eurasian Penduline Tit	Remiz pendulinus	Palaearctic	Wintering	0,000	1,679	0,687	0,746	1,101	0,634	0,000	0,000	0,000	2,292	0,000	0,000	0,460	0,000	0,000
Eurasian Blackcap	Sylvia atricapilla	European	Wintering	0,000	1,679	0,764	2,088	1,101	0,000	0,965	0,000	13,661	6,113	0,000	0,000	0,000	0,000	1,106
Garden Warbler	Sylvia borin	European	Migrant	0,000	0,000	0,000	0,071	0,000	0,000	0,143	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Dartford Warbler	Sylvia undata	Mediterranean	Wintering	0,000	0,305	0,076	0,000	0,367	0,317	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Subalpine Warbler	Sylvia cantillans	Mediterranean	Migrant	0,000	0,033	0,024	0,000	0,137	0,159	0,000	0,573	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Sardinian Warbler	Sylvia melanocephala	Turkestanian-Mediterranean	Sedentary	0,000	0,318	0,351	0,177	0,388	0,000	0,101	0,200	0,000	0,000	0,000	0,000	0,000	0,000	1,536
Bearded Reedling	Panurus biarmicus	Palaearctic	Sedentary	0,422	0,832	0,537	1,061	0,775	0,183	0,304	0,000	0,171	0,373	0,000	4,945	2,170	1,502	4,455
Melodious Warbler	Hippolais polyglotta	European	Occasional	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,200	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Eurasian Wren	Troglodytes troglodytes	Holarctic	Occasional	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,186	0,000	0,000	0,000	0,000	0,000
Common blackbird	Turdus merula	Palaearctic	Occasional	0,000	0,024	0,023	0,044	0,000	0,000	0,000	0,000	0,342	0,186	0,000	0,000	0,000	0,000	0,000
Song Trush	Turdus philomelos	European	Wintering	0,000	0,000	0,000	0,015	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Spotted Flycatcher	Muscicapa striata	European-Turkestanian	Occasional	0,000	0,024	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
European Robin	Erithacus rubecula	European	Wintering	0,000	0,305	0,305	0,746	1,101	3,804	1,448	1,321	4,554	5,349	0,000	0,000	0,000	0,000	0,553
Bluethroat	Luscinia svecica	Palaearctic	Migrant	0,292	1,063	0,690	0,638	0,137	0,477	0,143	2,294	0,744	2,461	0,000	1,586	2,469	0,000	4,481
Common Nightingale	Luscinia megarhynchos	European	Occasional	0,000	0,033	0,024	0,000	0,000	0,159	0,287	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,249
European Stonechat	Saxicola rubicola	Palaearctic	Sedentary	0,000	0,294	0,164	0,044	0,078	0,000	0,000	0,601	0,000	0,000	0,000	0,000	0,000	0,000	0,000
House Sparrow	Passer domesticus	Palaearctic	Sedentary	0,000	0,000	0,048	0,071	0,137	0,000	0,287	0,573	0,372	0,000	0,000	0,000	0,000	0,000	0,000
Eurasian Tree Sparrow	Passer montanus	European	Occasional	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,573	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Water Pipit	Anthus pratensis	Palaearctic	Wintering	0,000	0,000	0,305	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	1,106
Common Chaffinch	Fringilla coelebs	European	Wintering	0,000	0,000	0,076	0,597	1,101	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
European Greenfinch	Chloris chloris	European-Turkestanian	Occasional	0,000	0,066	0,048	0,000	0,137	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
European Goldfinch	Carduelis carduelis	European-Turkestanian	Occasional	0,000	0,000	0,048	0,000	0,274	0,000	0,143	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
European Serin	Serinus serinus	Mediterranean	Occasional	0,000	0,100	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000	0,000
Common Reed Bunting	Emberiza schoeniclus	Palaearctic	Wintering	0,000	0,763	0,000	0,597	2,935	0,000	0,000	0,000	0,000	0,764	0,000	0,000	0,460	0,000	4,425

Sampling Effort

				1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Breeding	March - October			4401	34292	29700	15917	9626	7778	7803	2722	5630	3926	10112	10112	13956	7290	4701
Wintering	November-February			338	6553	13095	6705	2726	3154	2072	2271	220	1309	0	0	2172	0	1808
Sedentary	January-December			4738	40846	42795	22622	12902	10933	9875	4993	5850	5369	382	10112	16128	7990	6509
Occasional	January-December			4738	40846	42795	22622	12902	10933	9875	4993	5850	5369	382	10112	16128	7990	6509
Migrant	March-May/ July-October			3422	30092	42000	14117	7313	6292	6969	1744	5374	3657	9456	9456	10530	5670	4017

Table 2. Species trends in EH for D1 period in EH and in Europa (1980-2022) (2) based on implemented TRIM software (Pannekoek & Van-Strien 2001). The overall trend based on multiplicative slopes jointly with Standard Errors (SE) are depicted. If the slope value is 1, there is no trend. If >1, there is a positive trend, if <1, the trend is negative. The percentage of change from imputed data is estimated from the slope. For instance, 1.08 means an 8% increase per year, 0.93 means a 7% decline per year. Data overdispersion was tested by the Wald test (values > 1 and significant and indicates that there is overdispersion and they fit a Poisson Regression equation. Trend status of each species is as the pointed out by Vorisek et al (2008). Best models are less sparse and those with greater Akaike's value (AIC). Data from (2) was obtained online in https://pecbms.info/trends-and-indicators/species-trends/.

Common name	Scientific name	Faunal element	EH Trend (1991-2005)						European Trend (1980-2022)	
			Slope (SE)	Wald	P	EH Change	AIC	EH Status	Slope (SE)	European Change
Barn Swallow	Hirundo rustica	Holarctic	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,994 (0,002)	-0,560
Cetti's wabler	Cettia cetti	Turkestanian-Mediterranean	1,057 (1,217)	0,570	0,570	5,7	137,700	Uncertain	0,021 (0,006)	-0,979
Southern Grey Shrike	Lanius meridionalis	Holarctic	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	Not Available	
Willow Warbler	Phylloscopus trochilus	Palaearctic	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,986 (0,004)	-1,410
Common Chiffchaff	Phylloscopus collybita	Palaearctic	1,146 (0,399)	0,970	0,702	14,6	959,430	Uncertain	1,015 (0,001)	1,460
Great Reed Warbler	Acrocephalus arundinaceus	European-Turkestanian	1,088 (0,049)	5,590	0,018	8,8	-21,230	Uncertain	1,003 (0,067)	0,290
Moustached Warbler	Acrocephalus melanopogon	Turkestanian-Mediterranean	1,097 (0,050)	6,120	0,014	9,7	0,740	Uncertain	Not Available	
Aquatic warbler	Acrocephalus paludicola	European	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	Not Available	
Sedge Warbler	Acrocephalus schoenobaenus	European-Turkestanian	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	1,000 (0,004)	0,000
Common Reed Warbler	Acrocephalus scirpaceus	European-Turkestanian	1,052 (0,110)	5,750	0,017	5,2	39,760	Uncertain	0,997 (0,001)	-0,300
Common Grashopper Warbler	Locustella naevia	European-Turkestanian	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,979 (0,007)	-2,210
Savi's Warbler	Locustella luscinoides	European-Turkestanian	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	Not Available	
Zitting Cisticola	Cisticola juncidis	Indian-African	1,000 (0,064)	0,000	1,000	0,0	-23,660	Uncertain	1,003 (0,002)	0,310
Great Tit	Parus major	Palaearctic	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	1,005 (0,001)	0,470
Eurasian Penduline Tit	Remiz pendulinus	Palaearctic	0,986 (0,062)	0,070	0,791	-1,4	-22,920	Uncertain	Not Available	
Eurasian Blackcap	Sylvia atricapilla	European	1,089 (0,092)	2,410	0,121	8,9	-1,550	Uncertain	1,028 (0,001)	2,760
Garden Warbler	Sylvia borin	European	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,991 (0,001)	-0,890
Dartford Warbler	Sylvia undata	Mediterranean	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	Not Available	
Subalpine Warbler	Sylvia cantillans	Mediterranean	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	1,047 (0,015)	4,700
Sardinian Warbler	Sylvia melanocephala	Turkestanian-Mediterranean	1,018 (0,056)	0,690	0,405	1,8	-23,930	Uncertain	1,009 (0,008)	0,860
Bearded Reedling	Panurus biarmicus	Palaearctic	1,099 (0,051)	6,330	0,012	9,9	-20,680	Uncertain	Not Available	
Melodious Warbler	Hippolais polyglotta	European	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,995 (0,002)	-0,490
Eurasian Wren	Troglodytes troglodytes	Holarctic	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	1,013 (0,001)	1,300
Common blackbird	Turdus merula	Palaearctic	1,000 (0,059)	0,190	0,666	0,0	-24,260	Uncertain	1,008 (0,001)	0,750
Song Trush	Turdus philomelos	European	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	1,006 (0,001)	0,550
Spotted Flycatcher	Muscicapa striata	European-Turkestanian	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,984 (0,002)	-1,600
European Robin	Erithacus rubecula	European	1,055 (0,072)	1,190	0,277	5,5	-17,290	Uncertain	1,008 (0,001)	0,790
Bluethroat	Luscinia svecica	Palaearctic	1,095 (0,051)	5,200	0,022	9,5	-22,310	Uncertain	Not Available	
Common Nightingale	Luscinia megarhynchos	European	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,989 (0,002)	-1,040
European Stonechat	Saxicola rubicola	Palaearctic	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,979 (0,005)	-2,140
House Sparrow	Passer domesticus	Palaearctic	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,984 (0,002)	-1,630
Eurasian Tree Sparrow	Passer montanus	European	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,981 (0,004)	-1,900
Water Pipit	Anthus pratensis	Palaearctic	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,981 (0,003)	-1,930
Common Chaffinch	Fringilla coelebs	European	1,000 (0,059)	237,880	0,000	42,0	-23,990	Strong increase	0,999 (0,000)	-0,080
European Greenfinch	Chloris chloris	European-Turkestanian	1,063 (0,173)	0,350	0,712	6,3	148,050	Uncertain	0,993 (0,002)	-0,660
European Goldfinch	Carduelis carduelis	European-Turkestanian	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	1,016 (0,003)	1,620
European Serin	Serinus serinus	Mediterranean	1,000 (0,059)	0,000	1,000	0,0	-26,000	Uncertain	0,973 (0,004)	**-2,670**
Common Reed Bunting	Emberiza schoeniclus	Palaearctic	1,063 (0,058)	2,670	0,102	6,3	-20,320	Uncertain	0,991 (0,001)	-0,890

Table 3. General diversity classification based on Shannon's Diversity Index for D1 and D2 periods. Species Specific Index (SSI-1 and SSI-2, see methods) are depicted only for D1 period.

604
605
606

Common name	Scientific name	Faunal element	Individuals		Diversity		Species-Specific Index	Weighted Species-Specific Index	
			1991-2005	2006-2012	1991-2005	2006-2012	1991-2005	1991-2005	2006-2012
Barn Swallow	*Hirundo rustica*	*Holarctic*	6	2	0,008	0,013	0,739	4,434	1,478
Cetti's wabler	*Cettia cetti*	*Turkestanian-Mediterranean*	376	23	0,198	0,093	0,159	59,784	3,657
Southern Grey Shrike	*Lanius meridionalis*	*Holarctic*	2		0,003		0,159	0,318	0,000
Willow Warbler	*Phylloscopus trochilus*	*Palaearctic*	16	5	0,019	0,029	0,068	1,088	0,340
Common Chiffchaff	*Phylloscopus collybita*	*Palaearctic*	1207	148	0,346	0,296	0,384	463,488	56,832
Great Reed Warbler	*Acrocephalus arundinaceus*	*European-Turkestanian*	254	94	0,155	0,235	0,159	40,386	14,946
Moustached Warbler	*Acrocephalus melanopogon*	*Turkestanian-Mediterranean*	221	93	0,141	0,233	0,225	49,725	20,925
Aquatic warbler	*Acrocephalus paludicola*	*European*	1		0,002		0,434	0,434	
Sedge Warbler	*Acrocephalus schoenobaenus*	*European-Turkestanian*	14	5	0,017	0,029	0,324	4,536	1,620
Common Reed Warbler	*Acrocephalus scirpaceus*	*European-Turkestanian*	2029	359	0,364	0,367	0,173	351,017	62,107
Common Grasshopper Warbler	*Locustella naevia*	*European-Turkestanian*	1	1	0,002	0,008	0,281	0,281	0,281
Savi's Warbler	*Locustella luscinoides*	*European-Turkestanian*	13	3	0,016	0,019	0,180	2,340	0,540
Zitting Cisticola	*Cisticola juncidis*	*Indian-African*	20	9	0,023	0,046	0,455	9,100	4,095
Great Tit	*Parus major*	*Palaearctic*	4	1	0,006	0,008	0,101	0,404	0,101
Eurasian Penduline Tit	*Remiz pendulinus*	*Palaearctic*	34	2	0,035	0,013	0,137	4,658	0,274
Eurasian Blackcap	*Sylvia atricapilla*	*European*	53	12	0,049	0,057	0,225	11,925	2,700
Garden Warbler	*Sylvia borin*	*European*	2		0,003		0,193	0,386	
Dartford Warbler	*Sylvia undata*	*Mediterranean*	5		0,007		0,346	1,730	
Subalpine Warbler	*Sylvia cantillans*	*Mediterranean*	5	1	0,007	0,008	0,259	1,295	0,259
Sardinian Warbler	*Sylvia melanocephala*	*Turkestanian-Mediterranean*	49	34	0,046	0,123	0,340	16,660	11,560
Bearded Reedling	*Panurus biarmicus*	*Palaearctic*	227	50	0,144	0,160	0,094	21,338	4,700
Melodious Warbler	*Hippolais polyglotta*	*European*	1		0,002		0,225	0,225	
Eurasian Wren	*Troglodytes troglodytes*	*Holarctic*	1		0,002		0,303	0,303	
Common blackbird	*Turdus merula*	*Palaearctic*	6		0,008		0,213	1,278	
Song Trush	*Turdus philomelos*	*European*	1	8	0,002	0,042	0,254	0,254	2,032
Spotted Flycatcher	*Muscicapa striata*	*European-Turkestanian*	1		0,002		0,275	0,275	
European Robin	*Erithacus rubecula*	*European*	41	19	0,040	0,081	0,000		0,000
Bluethroat	*Luscinia svecica*	*Palaearctic*	152	35	0,109	0,126			
Common Nightingale	*Luscinia megarhynchos*	*European*	6	2	0,008	0,013	0,173	1,038	0,346
European Stonechat	*Saxicola rubicola*	*Palaearctic*	24	3	0,026	0,019	0,412	9,888	1,236
House Sparrow	*Passer domesticus*	*Palaearctic*	9	2	0,012	0,013	0,159	1,431	0,318
Eurasian Tree Sparrow	*Passer montanus*	*European*	1	2	0,002	0,013	0,324	0,324	0,648
Water Pipit	*Anthus pratensis*	*Palaearctic*	6		0,008		0,520	3,120	
Common Chaffinch	*Fringilla coelebs*	*European*	9		0,012		0,048	0,432	
European Greenfinch	*Chloris chloris*	*European-Turkestanian*	5		0,007		0,399	1,995	
European Goldfinch	*Carduelis carduelis*	*European-Turkestanian*	5	1	0,007	0,008	0,254	1,270	0,254
European Serin	*Serinus serinus*	*Mediterranean*	3		0,005		0,455	1,365	0,000
Common Reed Bunting	*Emberiza schoeniclus*	*Palaearctic*	30	21	0,032	0,087	0,050	1,500	1,050
Woodchat Shrike	*Lanius senator*	*Mediterranean*		1		0,008	0,290		0,290
Western Yellow Wagtail	*Motacilla flava*	*Palaearctic*		1		0,008	0,193		0,193
Spectacled Warbler	*Sylvia conspicilliata*	*Mediterranean*		1		0,008	0,434		0,434

Table 4. *General Linearized Mixed Models* (GLMM) implemented by package "*glm2*" (Marschner 2011) in the R-Project software (R-Core Team 2921) on the specific abundances as fixed dependent variables with climatic and management variables as response-proxy independent variables. Only the main interactions among independent variables were considered. Slopes of the Linear Regression Models are depicted. Minor AIC's values were used to select the best model. ANOVA's F- value with degrees of freedom and determination coefficient (r2) of the selected model and p-values of the regression. Significant (P < 0.05) in bold. C = habitat; T = Temperature; P = Precipitation

Common name	Faunal element	Scientific name	GLM MODEL								
			Slope 1	Slope 2	Intercept	AIC score	Wighted AIC	df	F	r2	p
Barn Swallow	Holarctic	Hirundo rustica	0,1443·T		-0,23943	-29,416	0,242	1,13	0,147	0,031	0,530
Cetti's wabler	Turkestanian-Mediterranean	Cettia cetti	-0,0173·C		23,7632	102,800	0,425	1,9	0,058	0,009	0,777
Southern Grey Shrike	Holarctic	Lanius meridionalis	-0,0371·T		0,7478	-40,915	0,468	1,13	5,936	0,314	**0,030**
Willow Warbler	Palaearctic	Phylloscopus trochilus	-0,0008·P		0,319308	-4,309	0,395	1,13	1,366	0,095	0,264
Common Chiffchaff	Palaearctic	Phylloscopus collybita	-9,994·T		255,994	149,983	0,794	1,1	0,111	0,011	0,746
Great Reed Warbler	European-Turkestanian	Acrocephalus arundinaceus	-0,006·C		3,7382	50,152	0,395	1,1	2,253	0,184	0,164
Moustached Warbler	Turkestanian-Mediterranean	Acrocephalus melanopogon	-0,328·P		13,6693	76,836	1,000	1,1	1,036	0,162	0,194
Aquatic warbler	European	Acrocephalus paludicola	0,001·T		-0,0111095	-98,388	0,067	1,13	0,090	0,007	0,769
Sedge Warbler	European-Turkestanian	Acrocephalus schoenobaenus	-0,082·T	-0,002·P	2,074	-4,530	0,452	2,12	2,470	0,292	0,124
Common Reed Warbler	European-Turkestanian	Acrocephalus scirpaceus	-0,034·C		28,21971	93,481	0,431	1,1	2,218	0,182	0,167
Common Grashopper Warbler	European-Turkestanian	Locustella naevia	0,001·T		0,0110195	-98,388	0,426	1,13	0,090	0,007	0,769
Savi's Warbler	European-Turkestanian	Locustella luscinoides	0,000·P		0,090071	-28,630	0,426	1,13	0,429	0,032	0,524
Zitting Cisticola	Indian-African	Cisticola juncidis	-0,143·T	-0,002·P	3,413976	31,274	1,000	2,12	0,450	0,070	0,648
Great Tit	Palaearctic	Parus major	0,004·T		-0,074	-68,282	0,459	1,13	0,484	0,036	0,498
Eurasian Penduline Tit	Palaearctic	Remiz pendulinus	-0,002·C		1,332564	29,334	0,457	1,9	1,178	0,116	0,306
Eurasian Blackcap	European	Sylvia atricapilla	-0,028·P	-0,004·C	9,351298	65,849	0,324	2,8	1,375	0,256	0,307
Garden Warbler	European	Sylvia borin	0,000·P	-	-0,018795	-50,646	0,484	1,13	0,933	0,067	0,352
Dartford Warbler	Mediterranean	Sylvia undata	-0,027·T		0,61217	-7,128	0,283	1,1	0,396	0,038	0,543
Subalpine Warbler	Mediterranean	Sylvia cantillans	-0,037·T		7857	-10,526	0,362	1,13	0,781	0,056	0,393
Sardinian Warbler	Turkestanian-Mediterranean	Sylvia melanocephala	-0,001·C		0,634724	17,247	0,309	1,1	1,375	0,121	0,268
Bearded Reedling	Palaearctic	Panurus biarmicus	-0,007·C		3,3159	-47,904	0,385	1,1	3,441	0,256	0,093
Melodious Warbler	European	Hippolais polyglotta	-0,000·P		0,037816	-41,698	0,411	1,13	0,207	0,022	0,601
Eurasian Wren	Holarctic	Troglodytes troglodytes	-0,000·P		0,0666	-44,758	0,519	1,13	1,706	0,116	0,214
Common blackbird	Palaearctic	Turdus merula	-0,001·P		0,20642	-27,777	0,640	1,13	5,191	0,285	**0,040**
Song Trush	European	Turdus philomelos	0,028·T		0,53586	-44,605	0,428	1,1	9,915	0,498	**0,010**
Spotted Flycatcher	European-Turkestanian	Muscicapa striata	0,000·T		0,00735	-110,552	0,426	1,13	0,090	0,007	0,769
European Robin	European	Erithacus rubecula	-0,004·C		2,88132	49,497	1,000	1,9	0,443	0,050	0,511
Bluethroat	Palaearctic	Luscinia svecica	-0,008·C		3,70832	34,612	0,500	1,1	13,608	0,578	**0,004**
Common Nightingale	European	Luscinia megarhynchos	-0,037·T		0,7487	-40,920	0,486	1,13	5,936	0,314	**0,030**
European Stonechat	Palaearctic	Saxicola rubicola	-0,000·P		0,18037	-0,689	0,433	1,13	0,491	0,364	0,496
House Sparrow	Palaearctic	Passer domesticus	0,002·C		0,02016	-14,088	0,080	1,1	0,406	0,039	0,835
Eurasian Tree Sparrow	European	Passer montanus	-0,027·T		0,5321	-45,617	0,518	1,13	0,486	0,036	0,060
Water Pipit	Palaearctic	Anthus pratensis	0,018·T		-0,3164	-18,755	0,260	1,1	0,472	0,045	0,508
Common Chaffinch	European	Fringilla coelebs	0,004·P	0,001·C	0,331342	4,991	0,537	2,8	6,519	0,620	**0,021**
European Greenfinch	European-Turkestanian	Chloris chloris	-0,000·P		-0,0413	-64,309	0,540	1,13	1,848	0,124	0,109
European Goldfinch	European-Turkestanian	Carduelis carduelis	-0,000·P		0,05	-44,341	0,379	1,13	0,498	0,037	0,493
European Serin	Mediterranean	Serinus serinus	0,002·T		-0,025712	-72,969	0,426	1,13	0,090	0,007	0,765
Common Reed Bunting	Palaearctic	Emberiza schoeniclus	-0,000·C		1,97828	43,656	0,292	1,9	0,808	0,082	0,329