Paenitracins, a novel family of bacitracin-type nonribosomal peptide antibiotics produced

by plant-associated *Paenibacillus* species

Nataliia V. Machushynets [a, *], Somayah S. Elsayed [a], Chao Du [a], Vladyslav Lysenko [b], Mercedes

de la Cruz [c], Pilar Sanchez [c], Olga Genilloud [c], Nathaniel I. Martin [b], Mark R. Liles [d], and Gilles P.

van Wezel [a, e, *]

[a] Molecular Biotechnology Group, Institute of Biology, Leiden University, Sylviusweg 72, 2333 BE,

Leiden, The Netherlands

[b] Biological Chemistry Group, Institute of Biology, Leiden University, Sylviusweg 72, 2333 BE,

Leiden, The Netherlands

[c] Fundación MEDINA, Health Sciences Technology Park, Avda Conocimiento 34, 18016,

Granada, Spain.

[d] Department of Biological Sciences, Auburn University, 120 W Samford Av, AL 36849, Auburn,

AL, USA

[e] Department of Microbial Ecology, Netherlands Institute of Ecology, Droevendaalsesteeg 10,

6708 PB, Wageningen, The Netherlands

*Corresponding authors:

Nataliia V. Machushynets, email: n.v.machushynets@biology.leidenuniv.nl; Gilles P. van Wezel,

email: g.wezel@biology.leidenuniv.nl.

ABSTRACT

The growing threat of antimicrobial resistance necessitates the discovery of novel antibiotics with activity against drug-resistant pathogens. Members of the genus *Paenibacillus* are a rich source of nonribosomal peptides (NRPs), including well-known antibiotics such as polymyxins, paenibacterin and tridecaptins. Here we use a targeted Mass-QL-based mass spectrometry approach to identify the NRPs produced by a collection of 227 taxonomically diverse plant-associated *Paenibacillus* strains, providing detailed insights into their NRP-producing potential. Using MassQL to zoom in specifically on NRPs containing basic amino acids, we discovered a novel family of bacitracins, which we designated paenitracins. The paenitracins are the first bacitracin-type peptides reported in *Paenibacillus*, and are distinguished from canonical bacitracins by three previously unseen amino acid substitutions. The paenitracins exhibit potent activity against Gram-positive pathogens, including vancomycin-resistant *Enterococcus faecium* E155. Our work provides a novel metabolomics- and genomics-guided workflow for the discovery of bioactive NRPs as a strategy to prioritize natural product chemical space and accelerate antibiotic discovery.

IMPORTANCE

Members of the genus *Paenibacillus* play an important role in soil ecology, producing a range of important nonribosomal peptides (NRPs) that protect their eukaryotic host. A collection of plant-associated *Paenibacillus* spp. was analyzed for their phylogenetic and metabolic diversity. We developed a novel discovery pipeline that combines feature-based molecular networking with MassQL queries to systematically prioritize bioactive NRPs

containing basic amino acids. Thus we provide a comprehensive genus-wide inventory of

NRPs produced by *Paenibacillus* spp. We thereby identified the paenitracins, a new

subfamily of bacitracins active against multidrug-resistant Gram-positive pathogens. Our

pipeline enables the discovery of novel peptidic natural products to accelerate the

prioritization of chemical space for antibiotics.

Keywords:

Paenibacillus; antimicrobial resistance; natural product biosynthesis; nonribosomal

peptides; biosynthetic gene cluster; metabolic networking.

INTRODUCTION

Antimicrobial resistance (AMR) continues to challenge global health, and the discovery of new antibiotics is therefore a prerequisite to be able to keep fighting infectious diseases caused by multidrug-resistant pathogens (1). Bacterial natural products (NPs) represent a rich reservoir of antibacterial compounds, many of which have been successfully translated into clinical applications (2). A major challenge is to prioritize the huge extant genomic space for natural products (3).

Members of the bacterial phylum Bacillota (formerly Firmicutes) are well known for their potential to produce a wide range of antimicrobials, in particular nonribosomally synthesized peptides (NRPs) (4-6). Numerous bioactive NRPs produced by *Bacillus* and *Paenibacillus* have been identified, including octapeptins, paenibacterins, polypeptins, tridecaptins, etc (5-7). NRPs such as polymyxins and bacitracins that are derived from these microorganisms, have found application in clinical settings (8, 9). A large fraction of the NRPs contain positively charged amino acids and possess potent antimicrobial activities against Gram-positive and Gram-negative pathogens (10, 11). The mode of action polymyxins, involves binding to the lipid A component of lipopolysaccharides (LPSs) in the outer membrane of Gram-negative bacteria. Other NRPs, like bacitracin produced by *Bacillus subtilis* and *Bacillus licheniformis*, act as inhibitors of cell wall biosynthesis. (9) Specifically, in the presence of a divalent metal ion, most commonly Zn^{2+}, bacitracin binds to and sequesters the membrane-associated phospholipid undecaprenyl pyrophosphate ($C_{55}PP$) (12, 13). Bacitracin A is the major bioactive congener with the highest clinical relevance, and it is mainly used topically to treat infections caused by Gram-positive bacteria (14). The bacitracin family is composed of structurally similar macrocyclic dodecapeptides containing a unique N-terminal aminothiazoline moiety (15).

Previously reported bacitracin A-F congeners vary at positions 1, 5, and 8 of the peptide

sequence, with substitutions among Ile, Leu, and Val (16).

One of the challenges in NP discovery is the rapid identification of previously

reported compounds, termed structural dereplication (17). Liquid chromatography coupled

with high-resolution mass spectrometry (LC-MS) is widely used at the early stages of

dereplication studies. Molecular formulas of the detected molecules are determined

through the exact mass using the Seven Golden Rules, Sirius 2, and MSFINDER

software; and subsequently queried in various NP databases like DNP, UNPD,

ChemSpider, and REAXYS; to annotate the molecular structures (18).

In order to further identify compounds in LC-MS-based metabolite profiling

workflows, the introduction of the Global Natural Products Social Molecular Networking

(GNPS MN) has appeared as a powerful tool to process tandem mass spectrometry (MS)

data and perform molecular networking (MN) (19). The MN concept is based on organizing

and visualizing tandem MS data through a spectral similarity map, revealing the presence

of homologous MS^2 fragmentations. As structurally related compounds share similar

fragmentation spectra, their nodes tend to gather together and create clusters of analogs

(20). The known compounds are annotated by matching their MS^2 spectra against an

available spectral library. Moreover, combination of MN with annotation methods, such as

MS2LDA (21), NAP (22), or DEREPLICATOR (23) accelerates the identification of

unannotated ions (24). Mass Spec Query Language (MassQL) has been developed based

on advancements in molecular networking to explore underutilized MS/MS data further

(25). MassQL captures the unique characteristics of MS data, including isotopic patterns,

diagnostic fragmentation, and neutral loss; and establishes a comprehensive MS

terminology for searching MS patterns across datasets. Additionally, NP prioritization

strategies have been applied based on the combination of molecular networking with

different sources of information (e.g., genomic, taxonomic, geographic, bioactivity, and/or

spectral data) (26-29). Despite the ability of current tools to dereplicate NPs upstream of

the isolation process, effective strategies for the early detection of potentially bioactive

NPs within crude extracts are still lacking.

In this study, we provide a full phylogenomic analysis of 227 *Paenibacillus* spp. and

their potential to produce NRPs. We introduce an early-stage prioritization method that

combines feature-based MN (FBMN) and MassQL search to target basic amino acid-

containing NRPs with potential antimicrobial properties that are produced by a collection

of plant-associated *Paenibacillus* spp. Basic amino acid signatures were detected and

mapped onto a massive multi-informative molecular network. This led to the discovery of

new bacitracin congeners, which we designated paenitracins, with potent bioactivity

against Gram-positive pathogens, including vancomycin-resistant *E. faecium* E155.

RESULTS AND DISCUSSION

Taxonomic and chemical diversity of a collection of *Paenibacillus* strains

Considering the great pharmacological and biotechnological potential of the metabolites produced by *Paenibacillus*, we here analyzed 227 *Paenibacillus* isolates from the Auburn University (USA) plant-associated microbial strain collection. These *Paenibacillus* species had been isolated from the root surface of field-grown corn (*Z. mays*) and soybean (*G. max*) grown in Dunbar (Nebraska), Carrol (Iowa), Whitewater (Wisconsin), Sparta (Illinois), Troy (Ohio) and Tallassee (Alabama) (Table S1). Taxonomic analysis of the strains was conducted by comparing the 16S rRNA genes from the isolates, and a maximum-likelihood tree was constructed (Figure 1).

In search of novel NRPs, we investigated the chemical diversity of the specialized metabolites produced by *Paenibacillus* isolates. For this purpose, all 227 isolates of the collection were cultured individually on TSA media in 96 deep-well plates for 72 h and extracted with isopropyl alcohol (IPA) supplemented with 0.1% (v/v) formic acid (FA). Subsequently, crude extracts were tested against Gram-positive (*Staphylococcus aureus* ATCC 29213) and Gram-negative (*Escherichia coli* ATCC 25922) indicator strains to assess the antibacterial activity of the strains. Nine crude extracts exhibited significant growth inhibition of *S. aureus*, while 22 crude extracts showed bioactivity against *E. coli* (Figure 1, Table S1). These extracts were then subjected to LC-MS/MS analysis to identify the natural products accountable for the observed bioactivities. LC-MS/MS data were processed with MzMine 2 (30) and exported for GNPS FBMN. Prior to uploading the data to GNPS, media-derived mass features were eliminated. A mass exclusion threshold of 300 *m/z* was implemented to minimize the occurrence of mass features originating from small metabolites that are not relevant to our study. FBMN analysis resulted in generating

molecular network consisting of 3096 parent ions (nodes) connected through 3692 edges

(Figure 2). Characteristics of the natural products, such as annotation, *m/z* value and

species-specific molecule production were visualized using Cytoscape 3.9.1 (31).

Mass spectrometry-based molecular networking allows clustering of molecules with

similar MS/MS fragmentation patterns, which stem from the similarity in their structures

(19). The clusters allow visual exploration of the metabolites produced by the strains,

enabling rapid dereplication of known compounds and visualization of unknown

compounds (20). Known metabolites were dereplicated by matching the experimentally

derived MS/MS fragmentation patterns with the annotated and curated MS/MS spectra

within the GNPS database. Since the spectral library search did not work well for NRPs

from *Paenibacillus*, an in-house database was created of the MS data for known

lipopeptides from *Paenibacillus* and used to dereplicate known metabolites. This identified

lipopeptides belonging to the families of fusaricidins, polymyxins and tridecaptins (Table

S3, Figure 2, Figures S1-S3). These dereplicated mass features were used to propagate

the annotation to other connected mass features. Despite the identification of a substantial

number of putative congeners of known lipopeptides in the network, the majority of the

produced metabolites remained unidentified.

Targeting cationic lipopeptides using MassQL-assisted fragment search

Several of the identified lipopeptides are well-known antibiotics with potent activity,

typically attributed to their net positive charge (Table S3), which enhances electrostatic

interactions with the negatively charged components on the surface of bacterial cells, such

as the cell membrane or lipopolysaccharides (LPSs) (32). Therefore, we sought to detect

specifically those lipopeptides with potential bioactive properties that contained one or

multiple positively charged amino acid residues. To accomplish this, a feature-based

spectral similarity network was paired with a MassQL (33) search query to target the

MS/MS spectra that contained the most common positively charged amino acids. The

fragment search of b ions of 2,4-diaminobuturic acid (Dab) (m/z 101.0709), ornithine (Orn)

(m/z 115.0871), lysine (Lys) (m/z 129.1027), arginine (Arg) (m/z 157.1089) and the

immonium ion of histidine (His) (m/z 110.0718) was conducted with the accuracy of m/z

0.005 and at minimum intensity of 5% (Figure 3).

MassQL search of Dab-containing parent ions highlighted previously dereplicated

polymyxin and tridecaptin spectral families (Figure 4). The identification of different

polymyxin (34, 35) and tridecaptin (36) congeners within numerous *Paenibacillus* extracts

underscored the chemical diversity of our strain collection. Despite the structural

heterogeneity of polymyxins and tridecaptins, a shared characteristic is the occurrence of

multiple Dab amino acid residues. Furthermore, MassQL revealed a spectral family of

compounds that contained two distinct amino acid residues, namely ornithine and lysine

(Figure 5A). This cluster represented triply-charged ions with a mass range of 1500–1650

Da. Interestingly, the compounds from this cluster were produced exclusively by two

isolates from the entire strain collection, namely *Paenibacillus* sp. JJ-602 and

Paenibacillus sp. JJ-1115. These compounds could not be dereplicated by spectral library

and in-house database search. Extensive literature search and manual MS/MS

dereplication resulted in the annotation of these compounds as paenibacterin A

congeners (Figure 5B) (10).

Multiple positively charged lysine and ornithine residues contribute to the overall

charge of paenibacterins and enhance interactions with negatively charged LPS from the

outer membrane of Gram-negative bacteria (37).

Discovery of a novel family of bacitracins

In addition to the annotated lipopeptide classes, we identified multiple unknown molecular

families that contained compounds with basic amino acids such as lysine, ornithine, and

histidine (Figure 3). One molecular family of Lys-containing compounds stood out, as the

production was specific to only a limited number of isolates (Figure S4). The Lys-

containing molecular family could be distinguished into three sub-families (Figure 6A).

Sub-family 1 and 2 represent doubly charged ions with a mass range of 1080 to 1150 Da.

These compounds were similar to recently discovered lipopeptide paenilipoheptins A and

B produced by *P. polymyxa* E681 (38, 39). The third sub-family comprised doubly charged

molecular ions with significantly larger masses compared to paenilipoheptins (Figure 6A).

The edge connecting a node in sub-family 3 (*m/z* 712.8836) with the node due to

paenilipoheptin D (*m/z* 530.8307) has a cosine score of 0.6, indicating low similarity. A

mirror plot comparing the MS/MS spectra of the two mass features showed no matching

dipeptide fragment ions, and the only shared ions were due to a Lys residue and

fragments of Trp (Figure S6).

MS/MS spectra of all the mass features within this sub-family exhibited an ion with

an *m/z* 199.0897. However, these mass features could not be matched to known microbial

natural products. Analysis of the MS/MS spectra of the parent compound with *m/z*

712.8836 revealed that a fragment ion with an *m/z* 199.0897 has a molecular formula of

$C_9H_{15}N_2OS$, which corresponds to a thiazoline-containing substructure commonly found

in the peptide antibiotic bacitracin, so far exclusively identified from the genus *Bacillus*.

(40) Based on the fact that the b_2, b_3, b_4, and b_5 ions were shared between the parent

mass with *m/z* 712.8836 and bacitracin A (41), we concluded that their pentapeptide side

chains are identical. The differences observed in b_6–b_{11} ions indicated that the structural

distinctions between the two metabolites lie in the heptapeptide ring. We did not observe

fragment ions corresponding to Orn, Phe or His, which are found in the structure of

bacitracin A, while we could ascertain the presence of an ion corresponding to a Trp

residue (*m/z* 187.0864). Moreover, the dipeptide fragment ions Trp-Leu (*m/z* 300.171),

Trp-Thr (*m/z* 288.1352), and Leu-Leu (*m/z* 227.1764) were detected. By tracing the series

of both *b* and *y* ions, we elucidated the amino acid sequence of the compound with *m/z*

712.8836   as   (Ile-Cys)–Leu/Ile–Glu–Leu/Ile–Lys–Leu/Ile–Leu/Ile–Trp–Thr–Asp–Asn

(Figure S7). The MS/MS studies used did not allow us to discriminate between Leu and

Ile in positions 3, 5, 7 and 8 of the peptide amino acid sequence. Notably, this sequence

diverges from any known bacitracin analogues, indicating the discovery of a novel branch

of the important bacitracin-family of NRPs.

To purify the compound for more detailed studies, *Paenibacillus* sp. JJ-778 was

cultured in a 10 L MHB media and subjected to extraction using Diaion® HP-20 beads,

followed by elution with 100% IPA supplemented with 0.1% (v/v) formic acid. Multiple

rounds of chromatographic purification resulted in the isolation of compounds (**1**) and (**2**)

(Figures S8, S9). Both compounds had the same molecular formula of $C_{67}H_{106}N_{15}O_{17}S$,

based on their exact mass of *m/z* 1424.7584 for [M+H]$^+$, highly similar to the calculated

mass of 1424.7611. 1D and 2D NMR analysis of **1** confirmed the amino acid sequence

(Figure 7), and together with the molecular formula dictated by the accurate mass, the

structure of **1** was determined as a new congener of bacitracin (Table S4, Figures S10-

S16). NMR analysis identified Leu residues in positions 3 and 7, and Ile in positions 5 and

8. The ^{1}H NMR spectra of **1** and **2** were almost identical, suggesting that they are

diastereomers (Figure S17). Accordingly, the new compounds were designated as

paenitracin A (**1**) and B (**2**).

To verify the identity and stereochemistry of specific residues, compounds 1 and 2

were analyzed using Marfey's method (Table S5, S6) (42). This analysis confirmed the

presence of D-Glu, L-Lys, L-Thr, and L-Ile as single isomers. Additionally, it was found that

both L and D isomers of Leu were present in the molecules, and it was also observed that

the Asn and Asp residues exhibit different stereochemistry. However, the stereochemistry

of the thiazoline moiety and Trp could not be determined due to their instability under the

conditions used for Marfey's analysis. The 1H NMR spectra of molecules **1** and **2** were

nearly identical, with the exception of the region corresponding to the thiazoline moiety,

indicating that these molecules are diastereomers (Figure S17) with stereochemical

ambiguity arising from the thiazoline portion. Previous studies on bacitracin have

demonstrated that racemization at the Cys position within the thiazoline moiety completely

abolishes activity, whereas racemization at the Ile position only slightly impacts activity,

showing a preference for L-stereochemistry (43). Consequently, we propose that the

variation between molecules **1** and **2** is due to the racemization of the terminal Ile residue,

with compound **1** exhibiting D-stereochemistry and compound **2** showing L-

stereochemistry, as the latter was found to be marginally more potent.

Bacitracin, originally isolated from *Bacillus licheniformis* and *B. subtilis*, targets

primarily Gram-positive bacteria such as *Staphylococcus*, *Streptococcus*, *Enterococcus*

and *Clostridium difficile* (44). Bacitracin and its congeners are synthesized by a

nonribosomal peptide synthetase (NRPS) complex encoded by the bacitracin synthetase

operon, which consists of *bacA*, *bacB*, and *bacC* genes. To explore the biosynthesis of

the novel paenitracin, the genome of *Paenibacillus* sp. JJ-778 was sequenced and

assembled using Falcon (version 1.8.1) (45). Analysis using antiSMASH 7.1.0 (46) readily

identified a 12-modular gene cluster with 77% similarity to the bacitracin BGC of *B.*

licheniformis ATCC 10716. In silico analysis of the A domain substrate specificity of BacA,

BacB, and BacC predicted variations in amino acid composition at positions 7, 9, and 10

compared to the well-known bacitracin A (Figure 6B).

All known bacitracin congeners isolated and identified so far exhibit similar amino

acid compositions, with the Ile residues occasionally replaced by Val (Figure 6B). In

contrast, paenitracin contains Leu, Trp and Thr instead of Orn, Phe and His at the amino

acid positions 7, 9 and 10, respectively. This makes paenitracin a structurally distant

relative of the bacitracins.

Mode of action and bioactivity of paenitracins

Bacitracins specifically target the unique bacterial phospholipid undecaprenyl

pyrophosphate (C_{55}-PP). C_{55}PP is the membrane-associated phospholipid that anchors

lipid II and shuttles it between the cell's cytoplasm where it is synthesized, and the

membrane exterior, where it is incorporated into the growing cell wall (47). To see if

paenitracin indeed has the same mode of action as bacitracin, the C_{55}PP dependence of

paenitracin activity was tested using antibacterial activity antagonization assays in the

presence or absence of zinc (48). For this, paenitracin was mixed with C_{10}-PP, a more

soluble analogue of C_{55}-PP, (15) and incubated with cell suspension of *S. aureus* ATCC

29213. Antagonization assays showed that the activity of paenitracin B was antagonized

by C_{10}PP exclusively in the presence of Zn^{2+} (Table S7). This indicates that paenitracin

binds to C_{55}-PP of Gram-positive bacteria in a zinc-dependent manner and thus has the

same mode of actions as bacitracins.

To obtain insights into the antibacterial spectrum for paenitracin A and B, we

performed minimum inhibitory concentration (MIC) assays against a panel of

representative pathogenic strains, namely, *E. coli* ATCC 25922, *B. subtilis* 168, *S. aureus*

ATCC 29213, *S. aureus* USA300, *E. faecium* E155 and *E. faecium* E980. The media used

in the MIC assay was supplemented with 0.3 mM $ZnSO_4$ to provide the divalent metal ion

required for bacitracin's activity (43). The activity of the paenitracin diastereomers was

compared to that of commercially sourced bacitracin A, which served as a positive control.

Paenitracin B exhibited activity against Gram-positive bacteria, with MICs ranging from 1

to 8 µg/mL, comparable to commercial bacitracin (Figure 6C). *B. subtilis* 168, previously

reported to be resistant to bacitracin, also exhibited reduced susceptibility to paenitracins

A and B, with MIC values of 32 µg/mL. Importantly, paenitracin B exhibited activity against

vancomycin-resistant *E. faecium* E155 with MICs of 2 µg/mL, which is lower compared to

the MIC of the commercially available bacitracin A of natural origin. Considering the

divergent structure of paenitracin as compared to the previously reported congeners, the

mode of action should be investigated further.

In conclusion, MS-based indexing of NRPs from 227 plant-associated

Paenibacillus isolates revealed extensive chemical diversity. Molecular networking

combined with MassQL identified several classes of lipopeptides, whereby we specifically

focused on NRPs containing basic residues. This revealed the presence of distant

relatives of bacitracin, which we designated paenitracins. Paenitracin A and B showed

potent bioactivity against Gram-positive ESKAPE pathogens, including vancomycin-

resistant *E. faecium* E155. Unlike previously described bacitracin variants, paenitracins

show substantial structural divergence and represent the first example of bacitracin-family

NRPs in *Paenibacillus*. These findings highlight the untapped chemical space within this

genus and demonstrate the power of combining metabolomic indexing with targeted

mining strategies to accelerate the discovery of structurally unique antibiotics.

MATERIALS AND METHODS

General experimental procedures

NMR spectra were recorded on a Bruker Ascend 850 MHz NMR spectrometer (Bruker BioSpin GmbH). Data were analyzed using MestReNova 14 software (Mestrelab Research, Santiago de Compostela, Spain). High-performance liquid chromatography (HPLC) purification was performed on a Waters preparative HPLC system composed of a 1525 pump, a 2707 autosampler, a 2998 PDA detector, and a Water fraction collector III. All solvents and chemicals were of HPLC or LC-MS grade, depending on the experiment.

Isolation and identification of bacterial strains

Paenibacillus strains were obtained from the Auburn University Plant-Associated Microbial strain collection and had previously been isolated from the root surface of field-grown corn (*Zea mays*) and soybean (*Glycine max*) plants grown in Dunbar (Nebraska), Carrol (Iowa), Whitewater (Wisconsin), Sparta (Illinois), Troy (Ohio) and Tallassee (Alabama) (Table S1).

The phylogenetic tree was constructed using 16S rRNA gene sequences obtained through PCR amplification with universal bacterial primers 27F and 1492R (49). Each PCR amplicon was purified and used for Sanger sequencing. Subsequently, 16S rRNA gene sequences were assembled into consensus sequences and aligned with reference sequences. These include 220 sequences from our study, 41 sequences from type strains of *Paenibacillus* spp. downloaded from GenBank (Table S2) and sequence of *Bacillus subtilis* IAM 12118 (GenBank accession NR_112116.2). The alignment is done using ClustalOmega (v 1.2.2) (50). Aligned sequences are trimmed to the length of 1,452 bp to

remove extending regions from a few sequences and also to preserve as many variable

regions as possible. The trimmed alignment is re-aligned and a neighbor-joining

consensus tree is generated using Geneious (v 2024.0.3) tree function, with Tamura-Nei

genetic distance model and *B. subtilis* as outgroup. The generated tree is then uploaded

to ITOL for visualization (51).

Growth of *Paenibacillus* spp. and natural product extraction

Our method for cultivation and extraction of specialized metabolites was adapted from

previously described procedures (52). Briefly, frozen stocks of 227 *Paenibacillus* spp. from

the Auburn University strain collection were inoculated into 1600 µL of Tryptic Soy Agar

(TSA, Bacto Soybean-Casein Digest Medium, 30 g/L) in 2.0 mL 96 deep well plates

(Thermo Scientific, Nunc 2.0 mL DeepWell Plate). After 72 h, cultures were used to

inoculate 200 µL Tryptic Soy Broth (TSB, Bacto Soybean-Casein Digest Medium, 30 g/L)

with a pin replicator in 96 well plates. These cultures were incubated overnight at 30 °C

and used as an inoculum for the second 2.0 mL 96-deep well plate containing 600 µL

TSA. Plates were sealed with 96 Well-Cap Mats (Thermo Scientific, Nunc 96 Well-Cap

Mats), and incubated at 30 °C for 72 h. The cultures were extracted with 300 µL 100%

isopropanol acidified with 0.1% formic acid. The plates were resealed with the same 96

Well-Cap Mats, sonicated for 10 min, and extracted for an additional 50 min. 200 µL of

these crude extracts were transferred into a pre-washed 96 well plate (Agilent

Technologies, 96 well plates, 0.5 mL, polypropylene) and lyophilized to dryness. The

extraction protocol was repeated once more for a total extract volume of 400 µL. Dried

samples were redissolved in 160 µL of methanol:water (1:1 v/v).

Genome sequencing, assembly and annotation

*Paenibacillu*s sp. JJ-778 was grown in TSB at 30 °C and 220 rpm for 24 h. DNA was extracted as described (53). DNA quality was verified by agarose gel electrophoresis. PacBio sequencing and assembly was performed by Novogene (UK). Generally, libraries were prepared using SMRTbell template prep kit (PacBio, USA) according to manufacturer instructions. Sequencing was then performed using PacBio Sequel platform in continuous long reads mode. Assembly was done using Falcon (version 1.8.1) (45). The complete genome sequence of *Paenibacillus* sp. JJ-778 has been deposited in GenBank under the accession number CP194855. BGCs in these genomes were annotated using AntiSMASH (version 7.1.0) (46).

Data-dependent LC-ESI-HRMS/MS

LC-MS/MS acquisition was performed using Shimadzu Nexera X2 ultra-high-performance liquid chromatography (UPLC) system, with an attached photodiode array detector (PDA), coupled to Shimadzu 9030 QTOF mass spectrometer, equipped with a standard electrospray ionization (ESI) source unit. A total of 2 μL was injected into a Waters Acquity HSS C_{18} column (1.8 μm, 100 Å, 2.1 × 100 mm) and data acquisition was performed as previously described (54). Briefly, the gradient used was 5% B for 1 min, 5–85% B for 9 min, 85–100% B for 1 min, and 100% B for 4 min. The column was re-equilibrated to 5% B for 3 min before the next run was started. All the samples were analyzed in positive polarity, using data-dependent acquisition mode. In this regard, full scan MS spectra (*m/z* 100–1700, scan rate 10 Hz, ID enabled) were followed by two data-dependent MS/MS spectra (*m/z* 100–1700, scan rate 10 Hz, ID disabled) for the two most intense ions per

scan. The ions were fragmented using collision-induced dissociation (CID) with fixed

collision energy (CE 20 eV) and excluded for 1 s before being re-selected for

fragmentation. The parameters used for the ESI source were: interface voltage 4 kV,

interface temperature 300 °C, nebulizing gas flow 3 L/min, and drying gas flow 10 L/min.

MZmine 2 parameters

Prior to statistical analysis, mzXML files were imported into Mzmine 2.53 (30) and

processed as previously described (54). Briefly, mass ion peaks were detected for

MS^1 and MS^2 at a noise level of 2.0E2 and 0.0E0, respectively (positive polarity, mass

detector: centroid), and their chromatograms were built using ADAP chromatogram

builder. The detected peaks were smoothed, and the chromatograms were deconvoluted

(algorithm: local minimum search; chromatographic threshold: 90; search minimum in RT

range: 0.05; minimum relative height: 1%; minimum ratio of peak top/edge: 2; peak

duration: 0.03–3.00 min). The detected peaks were deisotoped (monotonic shape;

maximum charge: 3; representative isotope: most intense). Peak lists from different

samples were aligned (weight for RT = weight for m/z = 50; compare the isotopic pattern

with a minimum score of 50%). Missing peaks detected in at least one of the samples

were filled with the gap-filling algorithm (RT tolerance: 0.1 min, m/z tolerance was set to

0.002 m/z or 15 ppm). Duplicate peaks were filtered. Only the features with MS/MS data

were exported to a GNPS-FBMN.

In addition, all features that originate from the culture medium were removed by

retaining only features with an average peak intensity of at least 100 times greater in the

bacterial extracts than in the culture medium extracts. Features that are present in the

blanks with an intensity higher than 5000 were also removed. In the end, features with

m/z less than 300 were filtered out.

Molecular network analysis and MassQL search

The resulting feature quantification table (CSV file) and MS/MS spectrum files (in mgf

format) were uploaded to the GNPS webserver (http://gnps.ucsd.edu) (19). The LC-

MS/MS data for the 227 *Paenibacillus* isolates were analyzed using the Feature-Based

Molecular Networking tool (FBMN) (20). Briefly, the precursor ion mass tolerance was set

to 0.005 Da and the MS/MS fragment ion tolerance to 0.05 Da. A molecular network was

then created where edges were filtered to have a cosine score above 0.5 and more than

3 matched peaks. The molecular networks were visualized using Cytoscape software

version 3.9.1 (31) and displayed using an unweighted force-directed layout. The data are

publicly accessible in the MassIVE repository (MSV000094386) and the networking

results   and   parameters   can   be   found   at

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=2f1e0ae1728142249ac4e841d5a72e

f4 . The MassQL tool version 31.4 (33) was employed to search for specific MS/MS

fragments that correspond to the *b* or immonium ions of positively charged amino acids.

The fragment search of *b* ions of 2,4-diaminobuturic acid (*m/z* 101.0709), ornithine (*m/z*

115.0871), lysine (*m/z* 129.1027), and arginine (*m/z* 157.1089); together with the

immonium ion of histidine (*m/z* 110.0718), was conducted with an accuracy of *m/z* 0.005

and minimum intensity of 5%.

Up-scale fermentation, extraction, and isolation

Paenibacillus sp. JJ-778 was grown at 30 °C on tryptic soy agar (TSA) for 72 h and three colonies were inoculated into TSB and incubated at 30 °C overnight. This inoculum (1%) was used to inoculate thirteen 2 L Erlenmeyer flasks containing 0.75 L of sterile MHB and fermented at 30 °C while shaking at 200 rpm for 72 h. To extract the specialized metabolites produced by *Paenibacillus* sp. JJ-778 supernatant was collected by centrifugation and subjected to extraction using Diaion® HP-20 (Resindion, Mitsubishi), followed by elution with 100% IPA supplemented with 0.1% (v/v) formic acid. The crude extract was adsorbed onto silica gel (pore size 60 Å, 70–230 mesh, Sigma Aldrich), and loaded on a silica column, which was eluted using varying gradients of EtOAc/MeOH. The fractions eluted with EtOAc-MeOH (1:3) supplemented with 0.1% (v/v) formic acid were combined, reconstituted in 50% ACN and injected into the preparative SunFire column (19 × 150 mm), which was eluted with H_2O:ACN gradient of 15–60% in 30 min, at a flow rate of 15 mL/min, to yield three fractions. Fraction 3 was further separated on the semi-preparative SunFire column (10 × 250 mm) at 3 mL/min using a H_2O–ACN gradient of 30–50% in 30 min to yield paenitracin A (**1**, 0.3 mg) and paenitracin B (**2**, 0.2 mg).

Marfey`s analysis

The stereochemistry of chiral centers present at α carbons were assigned by applying derivatization methods coupled with chromatographic analysis. The advanced Marfey's method using L-FDAA (1-fluoro-2-4-dinitrophenyl-5-L-alanine amide) established the absolute configurations of amino acids (55). The general method for Marfey`s analysis was conducted as described) (42). Briefly, a sample of peptide (30 µg) in 6M HCl (100 µL)

was heated to 100 °C in a sealed vial for 8−12 h using heating block, after which the

hydrolysate was concentrated to dryness at 40 °C under a stream of dry N_2. The

hydrolysate was then treated with 1 M $NaHCO_3$ (20 µL) and L-FDAA (1% solution in

acetone, 40 µL) at 40 °C for 1 h, after which the reaction was neutralized with 1 M HCl (20

µL). An aliquot of the analyte was diluted 50 times with H_2O/ACN (1:1) and injected (2 µL)

into an HRMS instrument following the standard protocol of the analysis. The analyte

amino acid content was assessed by comparison to authentic standards. The authentic

standards were prepared via a similar procedure.

Antibacterial activity assays and MIC determination

For *E. coli* ATCC 25922 and *S. aureus* ATCC 29213, single colonies were inoculated into

20 mL of MHII broth medium in 250 mL Erlenmeyer flask and incubated overnight at 37

°C with shaking at 220 rpm and then diluted to obtain an assay inoculum of 5 x 10^5

CFU/mL. Aztreonam and vancomycin were used as reference positive controls for each

pathogen. Absorbance at OD_{612} nm was measured at T_0 (zero time) using an EnVision™

Microplate Reader (PerkinElmer), and assay plates were incubated at 37 °C. After that,

absorbance at OD_{612} nm was measured at T_f (final time) using the same microplate reader

equipment. The percentage of growth inhibition was calculated using the following

normalization:

469
$$\%\text{Inhibition} = 100 \times \left\{ 1 - \frac{([T_{f\text{Sample}} - T_{0\text{Sample}}] - [T_{f\text{Blank}} - T_{0\text{Blank}}])}{([T_{f\text{Growth}} - T_{0\text{Growth}}] - [T_{f\text{Blank}} - T_{0\text{Blank}}])} \right\}$$

$T_{0\text{Sample}}$, the absorbance of the strain growth in the presence of sample measured at zero

time;

$T_{fSample}$, the absorbance of the strain growth in the presence of sample measured at final

time;

$T_{0Growth}$, the absorbance of the strain growth in the absence of sample measured at zero

time;

$T_{fGrowth}$, the absorbance of the strain growth in the absence of sample measured at the

final time;

T_{0Blank}, the absorbance of the broth medium (blank) measured at zero time;

T_{fBlank}, the absorbance of the broth medium (blank) measured at the final time.

The Genedata Screener software (Genedata, Inc., Basel, Switzerland) was used to

process and analyze the data. The activity of the samples was expressed as a percentage

of growth inhibition, where (-100%) represented the total growth inhibition of the target

microorganism and 0% represented the total growth of the target microorganism in the

assay. The Z' factor predicts the robustness of an assay by considering the mean and

standard deviation of both positive and negative controls (56). The robust Z' factor (RZ'

factor) is based on the Z' factor, but standard deviations and means are replaced by the

robust standard deviations and medians, respectively. In all experiments performed in this

work, the RZ' factor obtained was between 0.9173 and 0.9684.

For MIC tests, blood agar plates were inoculated with glycerol stocks of *E. coli*

ATCC 25922, *B. subtilis* 168, *S. aureus* ATCC 29213, *S. aureus* USA300 (clinical isolate

from Texas Children's Hospital, USA), *E. faecium* E155 and *E. faecium* E980 and

incubated overnight at 37 °C. Following incubation, 3 mL of tryptic soy broth (TSB) was

inoculated with an individual colony and incubated at 37 °C with shaking at 220 rpm until

the cultures reached the exponential phase. The bacterial suspensions were then diluted

100-fold in cation-adjusted Mueller-Hinton broth (MHB) or Luria-Bertani (LB) broth

supplemented with 0.3 M $ZnSO_4$ to reach a bacterial cell density of 10^6 CFU/mL. In

parallel, test compounds were 2-fold serially diluted with cation-adjusted MHB or LB

supplemented with 0.3 M $ZnSO_4$ in polypropylene 96-well plates to achieve a final volume

of 50 µL per well. An equal volume of bacterial suspension (10^6 CFU/mL) was added to

the wells. The well plates were sealed with an adhesive membrane and incubated for 20

hours at 37 °C. The reported MIC values are based on three technical triplicates and are

defined as the lowest concentration of the compound that prevented the visible growth of

bacteria.

In vitro antagonization of paenitracin antibacterial activity

Antibacterial activity antagonization assay was performed as previously described (48).

Briefly, geranyl pyrophosphate ammonium salt ($C_{10}PP$) (1 mg/mL in methanol [aqueous

10 mM NH_4OH (7:3), Sigma Aldrich, Taufkirchen, Germany] was used as an antagonist

of paenitracin A in either the presence or absence of 0.3 mM $ZnSO_4$. Paenitracin A was

mixed with $C_{10}PP$ (five-fold molar excess relative to paenitracin) or added to empty wells

(absence of antagonist) giving in all cases 50 µL of total volume in each well with

paenitracin A at 4× MIC in the presence and absence of antagonist as well as in the

presence and absence of 0.3 mM $ZnSO_4$, in triplicate. Subsequently, 50 µL of *S. aureus*

ATCC 29213 with OD_{600} of 0.005 was added to the test compounds in the microtiter plate

to achieve a final antibiotic concentration of 2× MIC. The samples were incubated at 37

°C for 24 h with constant shaking and inspected for visible bacterial growth.

Acknowledgments

We thank D. van der Horst for technical assistance with LC-MS analysis, to Jesús Martín

for assistance in compound annotation and to Marnix Medema for stimulating discussions.

The work was supported by the NACTAR program of The Netherlands Organization for

Scientific Research (NWO), Grant 16440.

Competing Interests Declaration

The authors declare no conflicts of interests.

REFERENCES

1.   Jesudason T. 2024. WHO publishes updated list of bacterial priority pathogens. *The Lancet Microbe* 5.

2.   Newman DJ, Cragg GM. 2020. Natural products as sources of new drugs over the nearly four decades from 01/1981 to 09/2019. *J Nat Prod* 83:770-803.

3.   Medema MH, van Wezel GP. 2025. New solutions for antibiotic discovery: Prioritizing microbial biosynthetic space using ecology and machine learning. *PLOS Biology* 23:e3003058.

4.   Aleti G, Sessitsch A, Brader G. 2015. Genome mining: prediction of lipopeptides and polyketides from *Bacillus* and related Firmicutes. *Comput Struct Biotechnol J* 13:192-203.

5.   Cochrane SA, Vederas JC. 2016. Lipopeptides from *Bacillus* and *Paenibacillus* spp.: A gold mine of antibiotic candidates. *Med Res Rev* 36:4-31.

6.   Olishevska S, Nickzad A, Déziel E. 2019. *Bacillus* and *Paenibacillus* secreted polyketides and peptides involved in controlling human and plant pathogens. *Appl Microbiol Biotechnol* 103:1189-1215.

7.   Pilz M, Cavelius P, Qoura F, Awad D, Brück T. 2023. Lipopeptides development in cosmetics and pharmaceutical applications: A comprehensive review. *Biotechnol Adv* 67:108210.

8.   Falagas ME, Kasiakou SK. 2005. Colistin: the revival of polymyxins for the

management of multidrug-resistant Gram-negative bacterial infections. *Clin Infect*

Dis 40:1333-41.

9.   Johnson BA, Anker H, Meleney FL. 1945. Bacitracin: a new antibiotic produced by

a member of the *B. subtilis gr*oup. *Science* 102:376-377.

10.   Li YX, Zhong Z, Zhang WP, Qian PY. 2018. Discovery of cationic nonribosomal

peptides as Gram-negative antibiotics through global genome mining. *Nat*

Commun 9:3273.

11.   Cardoso P, Glossop H, Meikle TG, Aburto Medina A, Conn CE, Sarojini V, Valery

C. 2021. Molecular engineering of antimicrobial peptides: microbial targets, peptide

motifs and translation opportunities. *Biophys Rev* 13:35-69.

12.   Stone KF, Strominger JL. 1971. Mechanism of action of bacitracin: complexation

with metal ion and C_{55}-isoprenyl pyrophosphate. *Proc Natl Acad Sci U S A* 68.

13.   Siewert G, Strominger JL. 1967. Bacitracin: an inhibitor of the dephosphorylation

of lipid pyrophosphate, an intermediate in the biosynthesis of the peptidoglycan of

bacterial cell walls. *Proc Natl Acad Sci U S A* 57.

14.   Katz BE, Fisher AA. 1987. Bacitracin: a unique topical antibiotic sensitizer. *Journal*

of the American Academy of Dermatology 17:1016-1024.

15.   Economou NJ, Cocklin S, Loll PJ. 2013. High-resolution crystal structure reveals

molecular details of target recognition by bacitracin. *Proc Natl Acad Sci USA* 110.

16.   Ikai Y, Oka H, Hayakawa J, Matsumoto M, Saito M, Harada K, Mayumi Y, Suzuki

M. 1995. Total structures and antimicrobial activity of bacitracin minor components.

J Antibiot (Tokyo) 48:233-42.

17.   Hubert J, Nuzillard JM, Renault JH. 2017. Dereplication strategies in natural

product research: How many tools and methodologies behind the same concept?

Phytochem Rev 16:55-95.

18.   Gaudêncio SP, Bayram E, Lukić Bilela L, Cueto M, Díaz-Marrero AR,

Haznedaroglu BZ, Jimenez C, Mandalakis M, Pereira F, Reyes F, Tasdemir D.

2023. Advanced methods for natural products discovery: bioactivity screening,

dereplication, metabolomics profiling, genomic sequencing, databases and

informatic tools, and structure elucidation. *Mar Drugs* 21:308.

19.   Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, Nguyen DD,

Watrous J, Kapono CA, Luzzatto-Knaan T, Porto C, Bouslimani A, Melnik AV,

Meehan MJ, Liu WT, Crusemann M, Boudreau PD, Esquenazi E, Sandoval-

Calderon M, Kersten RD, Pace LA, Quinn RA, Duncan KR, Hsu CC, Floros DJ,

Gavilan RG, Kleigrewe K, Northen T, Dutton RJ, Parrot D, Carlson EE, Aigle B,

Michelsen CF, Jelsbak L, Sohlenkamp C, Pevzner P, Edlund A, McLean J, Piel J,

Murphy BT, Gerwick L, Liaw CC, Yang YL, Humpf HU, Maansson M, Keyzers RA,

Sims AC, Johnson AR, Sidebottom AM, Sedio BE, et al. 2016. Sharing and

community curation of mass spectrometry data with Global Natural Products Social

Molecular Networking. *Nat Biotechnol* 34:828-837.

20.   Nothias LF, Petras D, Schmid R, Dührkop K, Rainer J, Sarvepalli A, Protsyuk I,

Ernst M, Tsugawa H, Fleischauer M, Aicheler F, Aksenov AA, Alka O, Allard PM,

Barsch A, Cachet X, Caraballo-Rodriguez AM, Da Silva RR, Dang T, Garg N,

Gauglitz JM, Gurevich A, Isaac G, Jarmusch AK, Kameník Z, Kang KB, Kessler N,

Koester I, Korf A, Le Gouellec A, Ludwig M, Martin H C, McCall L-I, McSayles J,

Meyer SW, Mohimani H, Morsy M, Moyne O, Neumann S, Neuweger H, Nguyen

NH, Nothias-Esposito M, Paolini J, Phelan VV, Pluskal T, Quinn RA, Rogers S,

Shrestha B, Tripathi A, van der Hooft JJJ, et al. 2020. Feature-based molecular

networking in the GNPS analysis environment. *Nat Microbiol* 17:905-908.

21.   van der Hooft JJJ, Wandy J, Young F, Padmanabhan S, Gerasimidis K, Burgess

KEV, Barrett MP, Rogers S. 2017. Unsupervised discovery and comparison of

structural families across multiple samples in untargeted metabolomics. *Anal Chem*

89:7569-7577.

22.   da Silva RR, Wang M, Nothias LF, van der Hooft JJJ, Caraballo-Rodríguez AM,

Fox E, Balunas MJ, Klassen JL, Lopes NP, Dorrestein PC. 2018. Propagating

annotations of molecular networks using in silico fragmentation. *PLOS Comput Biol*

14:e1006089.

23.   Mohimani H, Gurevich A, Mikheenko A, Garg N, Nothias LF, Ninomiya A, Takada

K, Dorrestein PC, Pevzner PA. 2017. Dereplication of peptidic natural products

through database search of mass spectra. *Nat Chem Biol* 13:30-37.

24. de Jonge NF, Mildau K, Meijer D, Louwen JJR, Bueschl C, Huber F, van der Hooft

JJJ. 2022. Good practices and recommendations for using and benchmarking

computational metabolomics metabolite annotation tools. *Metabolomics* 18:103.

25. Damiani T, Jarmusch AK, Aron AT, Petras D, Phelan VV, Zhao HN, Bittremieux W,

Acharya DD, Ahmed MMA, Bauermeister A, Bertin MJ, Boudreau PD, Borges RM,

Bowen BP, Brown CJ, Chagas FO, Clevenger KD, Correia MSP, Crandall WJ,

Crüsemann M, Fahy E, Fiehn O, Garg N, Gerwick WH, Gilbert JR, Globisch D,

Gomes PWP, Heuckeroth S, James CA, Jarmusch SA, Kakhkhorov SA, Kang KB,

Kessler N, Kersten RD, Kim H, Kirk RD, Kohlbacher O, Kontou EE, Liu K, Lizama-

Chamu I, Luu GT, Luzzatto Knaan T, Mannochio-Russo H, Marty MT, Matsuzawa

Y, McAvoy AC, McCall L-I, Mohamed OG, Nahor O, Neuweger H, et al. 2025. A

universal language for finding mass spectrometry data patterns. *Nature Methods*

22:1247-1254.

26. Olivon F, Allard PM, Koval A, Righi D, Genta-Jouve G, Neyts J, Apel C,

Pannecouque C, Nothias LF, Cachet X, Marcourt L, Roussi F, Katanaev VL,

Touboul D, Wolfender JL, Litaudon M. 2017. Bioactive natural products

prioritization using massive multi-informational molecular networks. *ACS Chem*

Biol 12:2644-2651.

27. Nothias LF, Nothias-Esposito M, da Silva R, Wang M, Protsyuk I, Zhang Z,

Sarvepalli A, Leyssen P, Touboul D, Costa J, Paolini J, Alexandrov T, Litaudon M,

Dorrestein PC. 2018. Bioactivity-based molecular networking for the discovery of

630   drug leads in natural product bioassay-guided fractionation. *J Nat Prod* 81:758-

631   767.

632 28. Olivon F, Retailleau P, Desrat S, Touboul D, Roussi F, Apel C, Litaudon M. 2020.

633   Isolation of picrotoxanes from austrobuxus carunculatus using taxonomy-based

634   molecular networking. *J Nat Prod* 83:3069-3079.

635 29. Crüsemann M, O'Neill EC, Larson CB, Melnik AV, Floros DJ, da Silva RR, Jensen

636   PR, Dorrestein PC, Moore BS. 2017. Prioritizing natural product diversity in a

637   collection of 146 bacterial strains based on growth and extraction protocols. *J Nat*

638   *Prod* 80:588-597.

639 30. Pluskal T, Castillo S, Villar-Briones A, Oresic M. 2010. MZmine 2: modular

640   framework for processing, visualizing, and analyzing mass spectrometry-based

641   molecular profile data. *BMC Bioinformatics* 11:395.

642 31. Shannon P, Markiel A, Ozier O, Baliga N, Wang J, Ramage D, Amin N,

643   Schwikowski B, Ideker T. 2003. Cytoscape: a software environment for integrated

644   models of biomolecular interaction networks. *Genome Res* 13:2498-2504.

645 32. Straus SK, Hancock REW. 2006. Mode of action of the new antibiotic for Gram-

646   positive pathogens daptomycin: Comparison with cationic antimicrobial peptides

647   and lipopeptides. *Biochim Biophys Acta* 1758:1215-1223.

648 33. Jarmusch A, Aron A, Petras D, Phelan V, Bittremieux W, Acharya D, Ahmed M,

649   Bauermeister A, Bertin M, Boudreau P, Borges R, Bowen B, Brown C, Chagas F,

650   Clevenger K, Correia M, Crandall W, Crüsemann M, Damiani T, Fiehn O, Garg N,

Gerwick W, Gilbert J, Globisch D, Gomes PW, Heuckeroth S, James A, Jarmusch

S, Kakhkhorov S, Kang KB, Kersten R, Kim H, Kirk R, Kohlbacher O, Kontou E, Liu

K, Lizama-Chamu I, Luu G, Knaan TL, Marty M, McAvoy A, McCall L-I, Mohamed

O, Nahor O, Niedermeyer THJ, Northen T, Overdahl K, Pluskal T, Rainer J, Reher

R, et al. 2022. A universal language for finding mass spectrometry data patterns

doi:10.1101/2022.08.06.503000. bioRxiv.

34. Vogler K, Studer RO. 1966. The chemistry of the polymyxin antibiotics. *Experientia*

22:345-354.

35. Jian Li, Roger L Nation, John D Turnidge, Robert W Milne, Kingsley Coulthard,

Craig R Rayner, Paterson DL. 2006. Colistin: the re-emerging antibiotic for

multidrug-resistant Gram-negative bacterial infections. *Lancet Infect Dis* 6:589-

601.

36. Bann SJ, Ballantine RD, Cochrane SA. 2021. The tridecaptins: non-ribosomal

peptides that selectively target Gram-negative bacteria. *RSC Med Chem* 12:538-

551.

37. Knolhoff AM, Zheng J, McFarland MA, Luo Y, Callahan JH, Brown EW, Croley TR.

2015. Identification and structural characterization of naturally-occurring broad-

spectrum cyclic antibiotics isolated from *Paenibacillus*. *J Am Soc Mass Spectrom*

26:1768-1779.

38.   Machushynets NV, Lysenko V, Du C, Slingerland CJ, Elsayed SS, Liles MR, Martin
671   NI, van Wezel GP. 2025. Exploring the chemical space of *Paenibacillus* NRPs and
672   discovery of paenilipoheptin B. *Organic Letters* 27:2821-2825.

673   39.   Lysenko V, Machushynets NV, van Dam JL, Sterk FAC, Speer A, Ram AFJ,
674   Slingerland CJ, Van Wezel GP, Martin NI. 2025. Total synthesis, structure
675   elucidation, and bioactivity evaluation of the cyclic lipopeptide natural product
676   paenilipoheptin A. *Organic Letters* 27:2826-2831.

677   40.   Qin F, Zhang H, Liu H, Zhang X. 2020. High performance liquid chromatography-
678   quadrupole/Time of Flight–tandem mass spectrometry for the characterization of
679   components in bacitracin. *Chromatographia* 83.

680   41.   Suleiman SA, Song F, Su M, Hang T, Song M. 2017. Analysis of bacitracin and its
681   related substances by liquid chromatography tandem mass spectrometry. *J Pharm
682   Anal* 7:48-55.

683   42.   Lysenko V, Son S, Theriault ME, Slingerland CJ, Hauk G, Cleenewerk L, Speer A,
684   Berger JM, Lewis K, Martin NI. 2024. Total synthesis and structural reassignment
685   of the antitubercular natural product evybactin. *Chem Eur J*
686   doi:10.1002/chem.202403767:e202403767.

687   43.   Buijs N, Vlaming HC, van Haren MJ, Martin NI. 2022. Synthetic studies with
688   bacitracin A and preparation of analogues containing alternative zinc binding
689   groups. *ChemBioChem* 23:e202200547.

690   44.   André B. 2005. Antimicrobial agents: antibacterials and antifungals. ASM Press,
691   Washington, D.C.

692   45.   Chin CAO, Peluso P, Sedlazeck FA-O, Nattestad M, Concepcion GT, Clum A,
693   Dunn CA-O, O'Malley R, Figueroa-Balderas R, Morales-Cruz A, Cramer GA-O,
694   Delledonne M, Luo C, Ecker JR, Cantu DA-O, Rank DR, Schatz MC. 2016. Phased
695   diploid genome assembly with single-molecule real-time sequencing. *Nat Methods*
696   13:1050–1054.

697   46.   Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, Fetter A, Terlouw
698   BR, Metcalf WW, Helfrich EJN, van Wezel GP, Medema MH, Weber T. 2023.
699   antiSMASH 7.0: new and improved predictions for detection, regulation, chemical
700   structures and visualisation. *Nucleic Acids Res* 51:W46-W50.

701   47.   Storm DR. 1974. Mechanism of bacitracin action: a specific lipid-peptide
702   interaction. *Annals of the New York Academy of Sciences* 235:387-398.

703   48.   Buijs NP, Vlaming HC, Kotsogianni I, Arts M, Willemse J, Duan Y, Alexander FM,
704   Cochrane SA, Schneider T, Martin NI. 2024. A classic antibiotic reimagined:
705   Rationally designed bacitracin variants exhibit potent activity against vancomycin-
706   resistant pathogens. *Proceedings of the National Academy of Sciences*
707   121:e2315310121.

708   49.   Lane D. 1991. 16S/23S rRNA sequencing. Wiley.

709   50.   Sievers F, Higgins DG. 2018. Clustal Omega for making accurate alignments of
710   many protein sequences. *Protein Sci* 27:135-145.

51.   Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for

phylogenetic tree display and annotation. *Nucleic Acids Res* 49:W293-W296.

52.   Nguyen DD, Melnik AV, Koyama N, Lu X, Schorn M, Fang J, Aguinaldo K,

Lincecum TL, Ghequire MGK, Carrion VJ, Cheng TL, Duggan BM, Malone JG,

Mauchline TH, Sanchez LM, Kilpatrick AM, Raaijmakers JM, De Mot R, Moore BS,

Medema MH, Dorrestein PC. 2016. Indexing the *Pseudomonas* specialized

metabolome enabled the discovery of poaeamide B and the bananamides. *Nat*

Microbiol 2:16197.

53.   Kieser T, Bibb, M. J., Buttner, M. J., Chater, K. F. & Hopwood, D. A. 2000. Practical

Streptomyces genetics. John Innes Foundation.

54.   van Bergeijk DA, Elsayed SS, Du C, Santiago IN, Roseboom AM, Zhang L, Carrión

VJ, Spaink HP, van Wezel GP. 2022. The ubiquitous catechol moiety elicits

siderophore and angucycline production in *Streptomyces*. *Commun Chem* 5.

55.   Marfey P. 1984. Determination of D-amino acids. II. Use of a bifunctional reagent,

1, 5-difluoro-2, 4-dinitrobenzene. *Carlsberg Res Commun* 49:591-596.

56.   Zhang J-H, Chung TDY, Oldenburg KR. 1999. A simple statistical parameter for

use in evaluation and validation of high throughput screening assays. *J Biomol*

Screen 4:67-73.

FIGURES

Figure 1. Phylogenetic tree based on 16S rRNA sequences of plant-associated

***Paenibacillus* isolates and closely related type strains.** The tree was obtained by

neighbor-joining analysis. For an overview of the strains see Tables S1 and S2. Type

strains are indicated with green squares. The tree was rooted using *Bacillus subtilis* IAM

12118 as an outgroup. Antibacterial activity of the strains against *E. coli* ATCC 25922 and

S. aureus ATCC 29213 is indicated with purple and red squares, respectively. The tree

was generated using Geneious v2024.0.3.

Figure 2. Molecular network of mass features detected in the bacterial extracts from the *Paenibacillus* isolates. Each node represents a parent ion detected in one or more strains. Different node colors represent distinct molecular families. The red nodes highlight the specific annotated compound in each family, with the corresponding chemical structure displayed next to the network.

Figure 3. MassQL-annotated molecular network of the ions detected in the bacterial

extracts from *Paenibacillus* isolates. Colors of the nodes, as indicated in the figure,

highlight parent ions that contain specific basic amino acid residues (Dab, Orn, His, Lys

or combinations thereof) in the structures.

Figure 4. Dab-containing lipopeptide families highlighted by MassQL search.

Molecular families of polymyxins (A) and tridecaptins (B) consist of multiple structural

analogues, reflecting high chemical diversity. Pie charts were mapped to the nodes to

represent the relative precursor ion intensities within each extract. Note that both

polymyxins and tridecaptins contain multiple Dab residues.

Figure 5. MassQL-based detection of paenibacterins. A. MassQL-annotated

molecular family of paenibacterins. **B.** Chemical structure of paenibacterin A.

A. Paenitracin

B. *B. licheniformis* ATCC 10716

Ile Cys Leu D-Glu Ile Lys D-Orn Ile D-Phe His D-Asp Asn

Paenibacillus sp. JJ-778

bacA bacB bacC

Ile Cys Leu D-Glu Ile Lys D-X Ile D-Tyr Thr D-Asp Asn

Known bacitracin congeners

	1	2	3	4	5	6	7	8	9	10	11	12	
	I	C	L	E	I	K	O	I	F	H	D	N	Bacitracin A₁
	V	C	L	E	I	K	O	I	F	H	D	N	Bacitracin B₁
	V	C	L	E	I	K	O	V	F	H	D	N	Bacitracin C₁
	I	C	L	E	I	K	L	I	W	T	D	N	**Paenitracin**

C.

	PaenA	PaenB	BacA	MIC (µg/mL)
E. faecium E980				1
E. faecium E155				
S. aureus ATCC 29213				
S. aureus USA 300 (MRSA)				
B. subtilis 168				
E. coli ATCC 25922				32

Figure 6. Molecular family of Lys-containing NRPs highlighted by MassQL search.

A. Exploring the mass features from sub-family 3 led to the discovery of new bacitracin analog, designated as paenitracin. **B.** The BGC and its predicted amino acid substrates of new paenitracin from the genome of *Paenibacillus* sp. JJ-778 was compared to the reference BGC from *B. licheniformis* ATCC 10716. The amino acid sequence of paenitracin was compared to those of known bacitracins from *B. licheniformis*. Colored amino acids are unique to paenitracin. **C.** MIC data of paenitracin A and B and bacitracin against a panel of human pathogenic bacteria.

781 **Figure 7. Correlations obtained by COSY, HMBC, and NOESY measurements in**

782 **NMR of paenitracin A (1).**