mRNA delivery of genetically encoded mosaic-8 pan-sarbecovirus RBD vaccines

Alexander A. Cohen[1], Jennifer R. Keeffe[1], Lusineh Manasyan[1], Indeever Madireddy[1], Ange-Célia I. Priso Fils[1], Kim-Marie A. Dam[2], Haley E. Stober[2], Rory A. Hills[,3,4,7], Woohyun J. Moon[5], Paulo J.C. Lin[5], Mark R. Howarth[3], Magnus A.G. Hoffmann[2,6,8*], Pamela J. Bjorkman[1,8*]

Affiliations
[1]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA 91125, USA

[2]Gladstone Institutes, San Francisco, CA 94158, USA

[3]Department of Pharmacology, University of Cambridge, Tennis Court Road, Cambridge, CB2 1PD, UK

[4]Department of Biochemistry, University of Oxford, Oxford, OX1 3QU, UK

[5]Acuitas Therapeutics, Vancouver, BC, V6T 1Z3, Canada

[6]Department of Medicine, University of California, San Francisco, CA 94143, USA

[7]Present address: Department of Pathology, Stanford University, School of Medicine, Palo Alto, CA 94305, USA

[8]Lead contact

*Correspondence: magnus.hoffmann@gladstone.ucsf.edu, bjorkman@caltech.edu

Keywords
Deep mutational scanning; epitope mapping; mRNA and protein nanoparticle vaccines; pandemic preparedness; pan-sarbecovirus vaccine; systems serology; SySPEM (systems serology-polyclonal epitope mapping)

Summary
Global health remains threatened by spillovers of zoonotic SARS-like betacoronaviruses
(sarbecoviruses) that could be mitigated by a pan-sarbecovirus vaccine[1]. We described
elicitation of potently neutralizing and cross-reactive anti-sarbecovirus antibodies by
mosaic-8 nanoparticles (NPs) displaying eight different sarbecovirus spike receptor-binding
domains (RBDs) as 60 copies of eight individual RBDs[2-6] (mosaic-8 RBD-NPs) or 30 copies
of two "quartets," each presenting four tandemly-arranged RBDs[7] (dual quartet RBD-NPs).
To facilitate manufacture of a broadly protective mosaic-8 vaccine, we generated membrane-
bound RBD quartets that can be genetically encoded and delivered via mRNA: dual quartet
RBD-mRNA and dual quartet RBD-EABR-mRNA, which utilizes ESCRT- and ALIX-
binding region (EABR) technology that promotes immunogen presentation on cell surfaces
and circulating enveloped virus-like particles (eVLPs)[8]. Immunization with mRNA
immunogens elicited equivalent or improved binding breadths, neutralization potencies, T
cell responses, and targeting of conserved RBD epitopes across sarbecoviruses,
demonstrating successful conversion of protein-based mosaic-8 RBD vaccines to mRNA
formats. Systems serology[9] showed that the mRNA vaccines elicited balanced IgG subclass
responses with increased Fcγ receptor-binding IgGs, consistent with potentially superior Fc
effector functions. A new technique, Systems Serology-Polyclonal Epitope Mapping
(SySPEM), revealed distinct IgG-subclass-specific epitope targeting signatures across
mRNA and protein-based vaccine modalities. These results demonstrate successful
conversion of mosaic-8 RBD-NPs to mRNA or EABR-mRNA vaccines that provide easy
manufacturing and enhanced protection from future pandemic sarbecovirus outbreaks.

Introduction

Three coronaviruses have spilled over from animal hosts to cause human epidemics or pandemics in the past 25 years: SARS-CoV (hereafter SARS-1) in the early 2000s, MERS-CoV in 2012, and SARS-CoV-2 (hereafter SARS-2) in 2019[10]. Two of these coronaviruses, SARS-1 and SARS-2, are members of the SARS-like betacoronavirus (sarbecovirus) subgenus. Although SARS-1 is not currently circulating, SARS-2 continues to infect humans, driving emergence of new variants that necessitate regular updates of current COVID-19 vaccines[11]. Since we cannot predict how SARS-2 will continue to mutate or which zoonotic sarbecovirus will next infect humans, we need a pan-sarbecovirus vaccine that does not need updating to provide broad protection against new SARS-2 variants and future sarbecovirus spillovers[1].

Spike trimers of coronaviruses function in host cell entry after one or more of the receptor-binding domains (RBDs) adopt an "up" position to permit interactions with a host cell receptor[12]. RBD targeting has been suggested for COVID-19 vaccine development as RBDs are the main target of neutralizing antibodies (Abs)[13]. We previously described a vaccine approach designed to elicit broad Ab responses to diverse sarbecoviruses in which RBDs from multiple sarbecoviruses were covalently linked using the SpyCatcher-SpyTag system[2-5] to a SpyCatcher-mi3 protein nanoparticle[14] (NP)[2-5]. In this approach, mosaic-8 RBD-NPs displayed eight different sarbecovirus spike RBD antigens attached randomly to the 60 positions of the mi3 NP (Fig. 1a,b). We hypothesized that B cells bearing cross-reactive receptors (B cell receptors; BCRs) recognizing features in common between adjacent non-identical RBDs would be preferentially activated compared to B cells with BCRs recognizing immunodominant strain-specific epitopes[3] (Fig. 1a). To investigate this prediction, we used deep mutational scanning (DMS)[15] to map RBD epitopes recognized by Abs from animals immunized with mosaic-8 RBD-NPs or homotypic RBD-NPs (i.e., only presenting SARS-2 RBD), which revealed targeting of more conserved RBD regions by mosaic-8 RBD-NP–elicited Abs compared with targeting of more variable RBD regions by homotypic RBD-NP–elicited Abs[3-7]. This finding rationalized results from challenge experiments in which mosaic-8 RBD-NPs protected from both a matched viral challenge (i.e., from a virus represented by an RBD on the NP) and a mismatched challenge (i.e., from a virus not represented by an RBD on the NP), whereas homotypic RBD-NPs protected only from a matched challenge[3]. In addition to initial experiments in SARS-2–naïve animals, we also found broader Ab responses elicited by mosaic-8 RBD-NPs in animals that were pre-vaccinated with COVID-19 vaccines[4]. Taken together, these results suggest that a mosaic-8 RBD-based vaccine given as a COVID-19 booster could provide increased protection from SARS-2 variants and prevent future sarbecovirus spillover(s) from causing an epidemic or pandemic.

Manufacturing of a mosaic-8 RBD-NP vaccine is a challenging process that requires production of nine protein components: eight different RBDs and a NP to which the RBDs are coupled. Making an mRNA-based version of a mosaic-8 vaccine would also pose challenges due to the requirement of synthesizing eight individual modified mRNAs. The RBD-encoding mRNAs could either be packaged into individual lipid nanoparticles (LNPs) or co-formulated into a single LNP. In the first case, some cells would not take up all eight mRNA LNPs, therefore presenting fewer instances of different RBDs adjacent to each other, potentially reducing the hypothesized avidity effects to activate B cells with maximally cross-reactive BCRs (Fig. 1a). In the second case, co-formulation of eight mRNAs encoding each RBD into a single LNP would present technical and regulatory challenges. A potential solution relevant to converting a mosaic-8 vaccine into an mRNA format involves the reported modification of the mosaic-8 RBD-NP protein-based immunogen into an mi3 NP presenting two RBD "quartets" (each containing four RBDs arranged in tandem), which was shown to elicit equally broad and potent Abs as the original mosaic-8 RBD-

NP that presents monomeric RBDs[7] (Fig. 1c; dual quartet RBD-NP). Dual quartet RBD-NPs require manufacture of only three components (two RBD quartets and a NP), and by synthesizing mRNAs encoding membrane-bound RBD quartets, could be more easily converted to an mRNA-LNP format than mosaic-8 RBD-NPs. In addition to simpler manufacturing, mRNA-LNP vaccines offer an advantage over protein-based vaccines in that viral antigens undergo translation in the cytoplasm, thereby providing a source of viral peptides for presentation by MHC class I molecules to activate CD8+ cytotoxic T cells[16].

Figure 1. Protein- and mRNA-based pan-sarbecovirus immunogens.
(a) Avidity hypothesis. Left: Membrane-bound strain-specific BCRs (pale yellow) using avidity to recognize a strain-specific epitope (pale yellow triangle) on antigens attached to a homotypic NP. Middle: Strain-specific BCRs binding with only a single Fab (i.e., not using avidity) to a strain-specific epitope (triangle) on an antigen attached to a mosaic NP. Right: Cross-reactive BCRs using avidity to recognize a common epitope (blue circle) presented on different antigens attached to a mosaic NP, but not to strain-specific epitopes (triangles). Only a fraction of the 60 attached RBDs are shown for clarity. (b) Sarbecovirus phylogenetic tree (made using a Jukes-Cantor generic distance model with Geneious Prime® 2023.1.2) calculated from amino acid sequences of RBDs aligned using Clustal Omega[17]. Viruses with RBDs included in RBD quartets and mosaic-8 RBD-NPs are indicated with an asterisk. Scale bar = phylogenetic distance of 0.08 nucleotide substitutions per site. (c) Schematics of protein-based and mRNA-based pan-sarbecovirus immunogens (not to scale; showing only a fraction of antigens for clarity). (d) Schematics of mRNA constructs encoding RBD dual quartet immunogens used in this study.

Conversion of dual RBD quartet immunogens to an mRNA format (Fig. 1c; dual quartet RBD-mRNA) also offers the possibility of using the ESCRT- and ALIX-binding region (EABR) mRNA vaccine platform, which results in presentation of immunogens on cell surfaces and on circulating enveloped virus-like particles (eVLPs), thereby combining attributes of both mRNA and protein-NP vaccines[8] (Fig. 1c; dual quartet RBD-EABR-mRNA). eVLP formation is achieved by appending a short EABR sequence to the cytoplasmic domain of the mRNA-encoded immunogen, which recruits cellular ESCRT proteins to induce eVLP budding from the plasma membrane[8]. We previously showed that two immunizations with mRNA-LNP encoding spike-EABR elicited potent CD8+ T-cell responses and superior neutralizing antibody responses against original and variant SARS-2 compared to conventional spike-encoding mRNA-LNP and purified spike-EABR eVLPs, improving neutralizing titers >10-fold against Omicron-based variants.

Here, we show that the advantages of protein-based mosaic-8 RBD-NP vaccines can be transferred to an mRNA-LNP vaccine format. We compared immune responses in mice immunized with dual quartet RBD-mRNA versions of mosaic-8 immunogens (dual quartet RBD-mRNA and dual quartet RBD-EABR-mRNA) with responses to two versions of protein-based mosaic-8 NPs (dual quartet RBD-NP[7] and EABR-based dual quartet RBD-eVLPs) or to homotypic SARS-2 RBD-eVLPs. Immunization with mRNA immunogens elicited equivalent or improved binding breadths, neutralization potencies, T cell responses, and targeting of conserved RBD epitopes across sarbecoviruses compared to protein-based mosaic-8 immunogens, demonstrating successful conversion of the mosaic-8 RBD vaccine to mRNA formats that enable streamlined, scalable manufacturing of a pan-sarbecovirus vaccine.

Results

mRNA-encoded RBD quartets are presented on cell surfaces and eVLPs
mRNA-LNP can direct cell surface expression of a variety of gene products, including viral antigens[18]. We investigated whether protein-based soluble dual quartets could be modified to induce cell surface expression by creating constructs that encode an RBD quartet followed by a linker, transmembrane anchoring sequence, and cytoplasmic tail (Fig. 1d; RBD quartets 4a and 4b). The eight RBDs in the 4a and 4b quartet sequences correspond to the RBDs used in previous studies investigating dual quartet RBD-NPs constructed from soluble quartets[7] and to RBDs in a mosaic-8 RBD-NP constructed from soluble monomeric RBDs[2] (Fig. 1b). We used flow cytometry

to verify presentation of membrane-bound RBD quartets on the surface of cells transfected with
mRNA encoding RBD quartet 4a or RBD quartet 4b by staining with C118, a human mAb that
recognizes a conserved RBD epitope present on all eight RBDs[19,20]. Flow cytometry results
showed high and equivalent levels of RBD quartets 4a and 4b on cell surfaces when each was
expressed separately (Extended Data Fig. 1a).

We next investigated the use of EABR technology, which modifies membrane proteins delivered
by mRNA-LNP to induce self-assembly and budding of eVLPs, resulting in immunogen
presentation on cell surfaces and circulating eVLPs[8], for delivering quartet RBD sequences. eVLP
assembly is induced by appending a short amino acid sequence to the cytoplasmic domain of a
membrane protein, which recruits proteins from the host ESCRT pathway that drives budding for
many enveloped viruses[21]. Addition of an EABR sequence has been shown to produce eVLPs for
a variety of membrane proteins including SARS-2 spike[8], HIV-1 Env[8], influenza hemagglutinin[22],
RSV fusion protein[8] (class I viral fusion protein trimers), Andes virus glycoprotein (a tetramer)[23],
CCR5 (a G protein-coupled receptor with multiple transmembrane regions)[8], epidermal growth
factor receptor[23], and MHC class I proteins[24]. Compared to non-EABR RBD quartets, cells
expressing EABR-modified RBD quartets showed lower cell surface expression by flow cytometry
using the pan-sarbecovirus RBD-specific C118 Ab[19,20] (Extended Data Fig. 1a). Lower cell surface
expression for an EABR-containing membrane protein, which was also found when comparing
SARS-2 spike trimer expression plus and minus an EABR sequence[8], is presumably due to
incorporation of a subset of cell surface proteins into released eVLPs that are not retained on the
cell surface.

To evaluate whether the RBD quartet 4a-EABR and 4b-EABR constructs induced secretion of
RBD quartet-displaying eVLPs, we subjected supernatants of quartet 4a- or 4b–transfected cells
to an established eVLP purification procedure[8]. ELISAs were consistent with efficient eVLP
formation for EABR mRNAs, showing high quartet 4a or 4b protein levels in the purified eVLP
samples for EABR, but not non-EABR, quartets (Extended Data Fig. 1b). We also used Abs
specific for quartet 4a (anti-Rs4081 RBD)[25] and 4b (anti-WIV1 RBD)[26] to examine co-transfected
cells by flow cytometry, demonstrating that RBD quartets 4a and 4b were co-expressed at the
surface of individual cells (Extended Data Fig. 1c). To confirm that both quartets were
incorporated into the same eVLP, we used a sandwich ELISA to probe eVLPs resulting from an
RBD quartet 4a-EABR and 4b-EABR co-transfection. eVLPs were purified from co-transfected
cells, captured with anti-WIV1 RBD Ab (quartet 4b), and detected with anti-Rs4081 RBD Ab
(quartet 4a), verifying the presence of both quartets on dual quartet eVLPs (Extended Data Fig.
1d). As expected, no binding was detected for samples prepared in the same way from cells co-
transfected with non-EABR versions of the quartets or cells only transfected with quartet 4a-EABR
or 4b-EABR (Extended Data Fig. 1d). Taken together, these data show that mRNA-encoded RBD
quartets were co-expressed on the surface of transfected cells and that the two EABR-modified
quartets are co-displayed on eVLPs.

Dual quartet RBD mRNA immunogens elicit cross-reactive Abs
We next compared Ab responses in mice immunized with mRNA-encoded dual quartets to
immunizations with protein-based dual quartet RBD-NPs or with purified dual quartet RBD-
eVLPs. For mRNA immunizations, mRNA encoding RBD quartets 4a and 4b were co-formulated
into lipid nanoparticles (LNP) to generate dual quartet immunogens (dual quartet RBD-mRNA
and dual quartet RBD-EABR-mRNA), which were each injected intramuscularly at RNA doses of
1.5 µg or 0.5 µg (Fig. 2a, Extended Data Fig. 2a). We also included cohorts of mice that were
immunized with 5 µg of dual quartet RBD-NPs (calculated based on RBD content) in the presence

of adjuvant (Fig. 2a) and cohorts immunized with adjuvanted purified RBD-eVLPs (dual quartet
RBD-eVLP or SARS-2 RBD-eVLP; 5 µg RBD content per dose), the latter representing a
homotypic immunogen control presenting only one type of RBD (Fig. 1c).

Because binding and neutralizing Ab responses both correlate with protection in humans and
animals vaccinated with COVID-19 mRNA vaccines[27-29], we analyzed immunized mouse sera by
ELISA to evaluate Ab binding responses and used pseudovirus assays[30] to evaluate neutralization
(Fig. 2b,c; Extended Data Fig. 2b). When comparing mRNA-based to protein-based immunogens,
we used a 1.5 µg mRNA dose and a 5 µg (RBD content) dose for the protein-based immunogens
(Fig. 2b,c), which represent standard doses used for these vaccine modalities in mice[2,8,31]. We also
compared the mRNA and mRNA-EABR immunogens using a 0.5 µg mRNA dose to investigate
potential dose-sparing effects of the EABR technology (Extended Data Fig. 2b). To evaluate
potential sarbecovirus strain-specific differences in Ab binding and neutralization properties, we
displayed data to show responses to individual sarbecovirus antigens (Fig. 2b,c; left panels) and
also as the overall geometric mean (geomean) response elicited by the different cohorts across
evaluated antigens, a form that allows statistical evaluation of differences across different cohorts
(Fig. 2b,c; right panels).

The highest mean binding titers for sera from the five cohorts across a panel of matched and
mismatched sarbecovirus RBDs were found for dual quartet RBD-mRNA and dual quartet RBD-
EABR-mRNA (Fig. 2b, left). Thus, genetically encoded RBD quartets delivered by a 1.5 µg dose
of mRNA-LNPs were significantly more effective at eliciting cross-reactive RBD-binding Abs
across a panel of sarbecovirus RBDs than 5 µg of the dual quartet RBD-NP protein-based
immunogen (Fig. 2b, right; p<0.0001 for comparisons of dual quartet RBD-mRNA and dual
quartet RBD-EABR-mRNA). In addition, both dual quartet RBD-mRNA and dual quartet RBD-
EABR-mRNA exhibited significantly higher geomean binding titers than the protein-based dual
quartet RBD-eVLPs and homotypic SARS-2 RBD-eVLPs (Fig. 2b, right; p<0.0001 for all
comparisons of mRNA-LNP immunogens and protein-based immunogens). SARS-2 RBD-eVLPs
exhibited significantly lower binding titers across the RBD panel than the dual quartet RBD-NPs
and purified dual quartet RBD-eVLPs (Fig. 2b, right; p<0.0001 for all comparisons except
comparing with dual quartet RBD-eVLP, where p=0.0057), as expected based on results for
homotypic RBD-NPs[2,3].

We also compared neutralizing titers elicited by the five cohorts across a panel of matched and
mismatched sarbecovirus strains (Fig. 2c, left). When comparing neutralization titers by strain
between the five immunization cohorts, dual quartet RBD-mRNA, dual quartet RBD-EABR-
mRNA, and dual quartet RBD-NP exhibited the highest titers across the pseudovirus panel except
for neutralization of SARS-2 D614G, where homotypic SARS-2 RBD-eVLPs showed induction
of slightly higher titers (Fig. 2c, left). Geomean neutralization titers for the five cohorts (Fig. 2c,
right) were more similar to each other than the mean binding titers (Fig. 2b, right) but dual quartet
RBD-EABR-mRNA induced significantly higher mean neutralizing titers than dual quartet RBD-
NP (p=0.0152), dual quartet RBD-eVLP (p=0.0152), and SARS-2 RBD-eVLP (p=0.0470).

To investigate the effects of lower doses of the mRNA immunogens, we measured binding and
neutralization titers elicited by 0.5 µg doses of dual quartet RBD-mRNA and dual quartet RBD-
EABR-mRNA immunizations (Extended Data Fig. 2a,b). Binding titers against the RBDs tested
were not statistically different for the two mRNA immunogens, except for binding to the BM48-
31 RBD, in which case the dual quartet RBD-EABR-mRNA immunization induced significantly
higher binding than the dual quartet RBD-mRNA immunization (Extended Data Fig. 2b, left;

p=0.0089). Neutralization titers were significantly different for three viruses: SARS-2 D614G,
where titers were higher for dual quartet RBD-mRNA (p=0.00013), and for SARS-1 (p=0.00024)
and WIV1 (p<0.0001), where titers were higher for dual quartet RBD-EABR-mRNA (Extended
Data Fig. 2b, right panel). These results are consistent with the more potent neutralization of non-
SARS-2 viruses induced by dual quartet RBD-EABR-mRNA compared with dual quartet RBD-
mRNA at the 1.5 µg dose (Fig. 2c).
234
Despite eliciting high binding titers against both SARS-2 variants and non-SARS-2
sarbecoviruses, neutralization titers for mosaic-8 immunogens (dual quartet RBD mRNAs, dual
quartet RBD-NP, and dual quartet RBD-eVLP) were lower against SARS-2 variants than against
non-SARS-2 sarbecoviruses (Fig. 2c, Extended Data Fig. 2b). Importantly, however, binding Abs
can trigger immune responses to clear infected cells through Fc-mediated effector functions[32]
and/or complement activation[33]. Indeed, human mAbs that neutralized early SARS-2 strains but
lost neutralizing activity against SARS-2 Omicron variants retained Fc effector functions[32], and
non-neutralizing human anti-RBD mAbs protected from a SARS-2 challenge in animal models[34].
Thus, even in the absence of strong neutralizing titers against SARS-2 variants, the ability of
mosaic-8 immunogens to elicit Abs that exhibit broad binding across sarbecovirus RBDs is a
desirable trait for a protective pan-sarbecovirus vaccine.
246
247
248

Figure 2. Dual quartet RBD mRNA immunogens elicit cross-reactive Abs.
ELISA and neutralization results for terminal bleed serum samples (see also Extended Data Fig. 2). (a) Immunization regimen. Left: Mice were primed at day 0, boosted at day 28, and samples were collected from a terminal bleed at day 77 or 78. Right: Colors used to identify immunization cohorts and symbols indicating a sarbecovirus antigen or pseudovirus that is matched (filled in data points; gray shading around name) or mismatched (unfilled data points; black outline around name). Sarbecovirus strain names are colored in panels b and c according to clade. (b,c) Results for RBD-binding ELISAs (panel b) and pseudovirus neutralization assays (panel c) for serum samples from day 77 or 78. Dashed horizontal lines indicate detection limits for assays. Left: Geomeans of RBD-binding ELISA half-maximal effective dilution (ED_{50}) values (panel b) or neutralization half-maximal inhibitory dilution (ID_{50}) values (panel c) across a panel of viral antigens or pseudoviruses for sera from animals in each cohort (geomeans are plotted as symbols with geometric standard deviations indicated by error bars). Mean ED_{50} or ID_{50} values across antigens are connected by colored lines corresponding to the immunization cohort. Right: Box and whisker plots of geomean responses (ELISA ED_{50}s in panel b; neutralization ID_{50}s in panel c) with individual data points representing geomean responses across n=10 sera to a single antigen or

266 pseudovirus. Geomean titers were compared pairwise between immunization cohorts by Tukey's
267 multiple comparison test with the Geisser-Greenhouse correction (as calculated by GraphPad
268 Prism). Significant differences between cohorts are indicated by asterisks: $p<0.05 = *$, $p<0.01 =$
269 $**$, $p<0.001 = ***$, $p<0.0001 = ****$.
270
271

mRNA-encoded dual quartets elicit potent T cell responses

273 T cell responses to the various immunogens were evaluated on day 77/78 in splenocytes from
274 immunized mice using enzyme-linked immunosorbent spot (ELISpot) assays[35]. Secretion of IFN-
275 γ (a cytokine produced by CD4+ T helper 1 (T_H1) cells and cytotoxic CD8+ T cells) and IL-4 (a
276 cytokine produced by CD4+ T helper 2 (T_H2) and T follicular helper cells[36]) was measured after
277 splenocytes were stimulated with a pool of SARS-2 spike peptides.
278
The dual quartet RBD-mRNA and dual quartet RBD-EABR-mRNA cohorts exhibited potent IFN-
γ responses, consistent with activation of RBD-specific cytotoxic CD8+ T cells, whereas
low/undetectable IFN-γ responses were found for the protein-based dual quartet RBD-NP, dual
quartet RBD-eVLP, and SARS-2 RBD-eVLP cohorts, results consistent with previous findings for
mice immunized with purified spike-EABR eVLPs[8] and the fact that mRNA-LNP immunogens,
but not protein immunogens, promote intracellular expression of antigens that serve as sources for
peptides loaded onto MHC class I proteins[37]. We found strong IL-4 responses for both the mRNA
and protein-based immunogen cohorts (Fig. 3).
287
288 Thus, mRNA-based mosaic-8 dual RBD quartet immunizations induce superior (INF-γ) or
289 equivalent (IL-4) release of cytokines compared with protein-based mosaic-8 immunogens.
290

291
Figure 3. mRNA-encoded dual quartets elicit potent T cell responses. ELISpot assay data for
SARS-2 RBD-specific IFN-γ (left) and IL-4 (right) responses of splenocytes from BALB/c mice
that were immunized with the indicated immunogens (immunization regimen in Fig. 2a). Results
are shown as spots per $3x10^5$ cells for individual mice (colored circles) presented as the median
(bars) and standard deviation (horizontal lines). Cohorts were compared by Tukey's multiple
comparison test calculated by GraphPad Prism. Significant differences between cohorts linked by
horizontal lines are indicated by asterisks: $p<0.05 = *$, $p<0.0001 = ****$.
299

mRNA-encoded and protein-based dual quartets elicit Abs targeting similar epitopes

301 To investigate which RBD epitopes were targeted by Abs elicited by the different vaccine
302 modalities, we used DMS to evaluate the effects of all possible amino acid changes in an RBD on
303 binding to polyclonal serum Abs[15,38-43]. To interpret results, we classified RBD residues by

epitopes that were defined based on structural properties and conservation/variability across sarbecoviruses: class 1 and class 2 RBD epitopes exhibit more sequence variability across sarbecoviruses and SARS-2 variants, and class 4, class 1/4, and portions of class 3 and class 5 epitopes are more conserved[20,44-46] (Fig. 4a). In some DMS analyses of polyclonal serum Abs, we observed weak escape profiles that lacked clear features, which we defined as a "polyclass," i.e., a more balanced, response[4]. We interpreted polyclass responses as resulting from sera containing multiple classes of anti-RBD Abs based on control experiments in which we evaluated equimolar mixtures of anti-RBD monoclonal Abs that targeted multiple epitopes[4]. For the current experiments, we analyzed serum using yeast display libraries encoding RBDs derived from SARS-2 WA1 (matched) and SARS-1 (mismatched) spike proteins.

DMS results showed that the epitopes targeted in both RBD libraries by the mRNA-encoded dual quartet RBD immunogens were similar overall to each other and to epitopes targeted by the protein-based dual quartet RBD-NP immunogen (Fig. 4b,c; Table 1, Extended Data Figs. 3, 4). In particular, the dual quartet immunogens, like mosaic-8 RBD-NPs[3,4], did not primarily elicit Abs against the immunodominant and variable class 1 and class 2 RBD epitopes, as observed for Abs elicited by homotypic SARS-2 RBD NPs[3,4] and spike-based mRNA vaccines[47]. Instead, elicited Abs primarily targeted class 4 and class 5 RBD epitopes, except for the Abs raised by immunization with dual quartet RBD-NPs, where class 5 was the predominant targeted epitope.

Although dual quartet RBD immunizations resulted in a mixed class 4/class 5 RBD response when evaluated against the SARS-2 WA1 and SARS-1 RBD libraries (Fig. 4b,c; Table 1, Extended Data Figs. 3, 4), there were notable differences. For example, both RNA-based immunogens elicited similar class 4 and class 5 RBD responses, but dual quartet RBD-mRNA sera showed more of a class 1/class 2 RBD response than dual quartet RBD-EABR-mRNA sera against the SARS-2 WA1 RBD library (Fig. 4b; Table 1), perhaps explaining why dual quartet RBD-mRNA elicited higher neutralization titers against the matched SARS-2 D614G pseudovirus (Fig. 2c, Extended Data Fig. 2b). Against the SARS-1 RBD library, each of the dual quartet immunogens induced strong class 4 and class 5 responses. In addition, weak escape by Abs against the variable class 1 and class 2 RBD epitopes was seen for dual quartet RBD-mRNA, dual quartet RBD-EABR-mRNA, and dual quartet RBD-NP (Table 1). Finally, one mouse in the dual quartet RBD-EABR-mRNA group and two in the dual quartet RBD-NP group elicited polyclass responses, consistent with the ability of these dual quartet immunogens to induce Abs against multiple RBD epitopes.

We conclude that converting dual quartet RBD immunogens into an mRNA format retains the ability of mosaic-8 vaccines to elicit Abs mainly against more conserved RBD regions, as previously observed in DMS experiments comparing protein NPs presenting dual RBD quartets or eight monomeric RBDs[7].

342
343
Figure 4. Dual quartet RBD-mRNA immunogens elicit Ab responses against conserved class
4 and class 5 RBD epitopes.
(a) Sequence conservation of 16 sarbecovirus RBDs calculated using ConSurf[48] shown on a
surface representation of SARS-2 RBD (PDB 7BZ5). Class 1, 2, 3, 4, 1/4, and 5 anti-RBD Ab
epitopes[20,44-46] are outlined in dots in different colors using information from representative
structures of Abs bound to SARS-2 spike or RBD (C102: PDB 7K8M; C002: PDB 7K8T, S309:
PDB 7JX3; CR3022: PDB 7LOP; C118: PDB 7RKV; WRAIR-2063: PDB 8EOO). (b,c) Left:
Line plots for DMS results using a SARS-2 WA1 RBD library (b) and SARS-1 library (c) from
the indicated number of mice immunized with the immunogens listed above each line plot. X-axis:
RBD residue number. Y-axis: sum of the Ab escape of all mutations at a site (larger numbers =
more Ab escape). Each line represents one antiserum with thick lines showing the average across
the n = 5 or 6 sera in each group. Lines are colored according to RBD epitopes in panel a. Right:
Average site-total Ab escape calculated for results from n = 5 or 6 serum samples for the indicated
RBD yeast display libraries. Mice were immunized with the immunogens listed, and results were

Immunogen	Animal	WA1 DMS Profile						SARS-1 DMS Profile				
		class 1	class 2	class 3	class 4	class 5		class 1	class 2	class 3	class 4	class 5
Dual Quartet RBD-mRNA	9076	None	None	None	Moderate	None		None	Weak	None	Strong	Moderate
Dual Quartet RBD-mRNA	9077	None	Weak	None	None	Weak		None	Weak	None	None	Strong
Dual Quartet RBD-mRNA	9078	Weak	Moderate	None	None	Weak		None	None	None	Strong	Strong
Dual Quartet RBD-mRNA	9079	Weak	None	None	Weak	None		—	—	—	—	—
Dual Quartet RBD-mRNA	9080	Weak	None	None	None	Strong		None	Weak	None	Weak	Strong
Dual Quartet RBD-mRNA	9081	None	None	None	None	Moderate		Weak	Weak	None	Strong	Strong
Dual Quartet RBD-EABR-mRNA	9091	None	None	None	Weak	Strong		None	Weak	Weak	Strong	Strong
Dual Quartet RBD-EABR-mRNA	9092	Polyclass	Polyclass	Polyclass	Polyclass	Polyclass		None	Weak	Weak	Strong	Strong
Dual Quartet RBD-EABR-mRNA	9093	None	None	None	Moderate	None		None	Weak	None	Strong	Strong
Dual Quartet RBD-EABR-mRNA	9094	None	None	None	None	None		None	Weak	None	Strong	Strong
Dual Quartet RBD-EABR-mRNA	9095	None	None	None	None	None		Weak	None	None	None	Moderate
Dual Quartet RBD-EABR-mRNA	9096	None	None	None	None	Moderate		None	Weak	None	Moderate	Strong
Dual Quartet RBD-NP	9043	None	None	None	None	Moderate		None	Weak	None	Strong	Strong
Dual Quartet RBD-NP	9044	None	None	None	None	None		None	Weak	None	Strong	None
Dual Quartet RBD-NP	9045	Polyclass	Polyclass	Polyclass	Polyclass	Polyclass		—	—	—	—	—
Dual Quartet RBD-NP	9049	None	None	None	None	Moderate		None	None	None	None	Strong
Dual Quartet RBD-NP	9050	None	None	None	None	Moderate		None	None	None	Strong	None
Dual Quartet RBD-NP	9051	Polyclass	Polyclass	Polyclass	Polyclass	Polyclass		Weak	None	None	Moderate	Moderate

Escape fraction

Polyclass	all classes <0.5	None	<0.5	Weak	0.5-1	Moderate	>1-2	Strong		>2

Table 1. Summary of DMS data showing elicitation of strong Ab responses against class 4 and class 5 RBD epitopes by dual quartet RBD-mRNA immunogens.
Results for individual mice are shown with ID numbers indicated. DMS was conducted using a SARS-2 WA1 RBD library (left) or a SARS-1 RBD library (right). DMS profiles were classified as polyclass, weak, moderate, or strong as indicated by the maximum escape fraction for any RBD position in each RBD epitope class. Colors and different designations for epitopes were assigned as indicated at the bottom (white entries for animals 9079 and 9045 indicate that no data were available). See also Fig. 4 and Extended Data Figs. 3, 4.

mRNA-encoded dual quartet RBD immunogens elicit balanced IgG2a/IgG1 responses
IgG subclasses induce distinct Fc effector functions, including opsonization, phagocytosis, and Ab-dependent cell-mediated cytotoxicity, through differential binding of their Fc domains to Fc gamma receptors (FcγRs)[49]; e.g., mouse IgG1, IgG2a, and IgG2b bind the activating receptor FcγR3 and the inhibitory receptor FcγR2b, IgG2a and IgG2b bind the activating receptor FcγR4, and mouse IgG3 does not bind detectably to any of these FcγRs[50]. Of the activating receptors, FcγR3 is more widely expressed than FcγR4, with FcγR4 mainly expressed on monocytes/macrophages and neutrophils and FcγR3 found on those cells plus dendritic cells, natural killer cells, basophils, mast cells, and eosinophils[50]. In immunized mice, IgG2a is associated with T_H1-type immune responses, effective activation of complement, and protective interactions with FcγRs, which are critical functions for clearing viruses[51]. IgG1, although usually efficient in neutralization of viruses, is associated with T_H2-type responses, which are thought to be less effective than T_H1 responses in anti-viral immune defense[52].

We used systems serology[9] to evaluate the IgG distributions elicited by mRNA versus protein
immunogens by comparing binding of IgG1, IgG2a, IgG2b, IgG3, FcγR2b-, FcγR3-, or FcγR4-
binding IgGs, and total IgG (defined as IgGs isolated using a pan-IgG Ab) elicited by the five
immunogens to a panel of spike trimers and RBDs derived from SARS-2 variants and other
sarbecoviruses. Results are presented to show the elicited responses from individual mice to each
antigen (Fig. 5) and geomean responses across the antigens for each immunogen (Extended Data
Fig. 5).

Both dual quartet RBD-mRNA immunogens elicited high levels of IgG1 and IgG2a against a broad
panel of sarbecovirus spikes and RBDs, with somewhat higher levels of IgG2a than IgG1 (Fig. 5,
Extended Data Fig. 5), demonstrating a desirable mixture of IgGs for an anti-viral vaccine of
induced T_H1 and T_H2 responses. In addition, the mRNA-based immunogens each elicited
significantly higher IgG2a and FcγR4-binding IgG responses than the protein-based immunogens
(p<0.0001) (Extended Data Fig. 5). The protein-based dual quartet RBD-NP and dual quartet
RBD-eVLP immunogens elicited higher levels of IgG1 than IgG2a, consistent with their
administration with Addavax adjuvant, which enhances T_H2-biased immune responses[53]. Dual
quartet RBD-eVLPs also elicited significantly higher IgG2a responses than dual quartet RBD-NPs
(p<0.0001). Interestingly, dual quartet RBD-EABR-mRNA elicited significantly higher IgG2b
(p<0.0001), as well as FcγR3- (p=0.0013) and FcγR4- (p=0.0021) binding IgGs, than dual quartet
RBD-mRNA. Finally, consistent with earlier demonstrations of the more limited sarbecovirus
recognition properties of Abs raised in response to homotypic RBD-NPs[2-4,25], the SARS-2 RBD-
eVLP immunogen elicited IgGs mainly against SARS-2 and closely related RBDs (Fig. 5).
412

413   Taken together, these results demonstrate that the mRNA versions of dual quartet immunogens
414 elicit balanced IgG responses that recognize a broad range of sarbecovirus antigens, arguing for
415 their likely efficacy to protect against diverse sarbecoviruses.

417
418
Figure 5. mRNA-encoded dual quartet RBD immunogens elicit balanced IgG subclass and
potent FcγR-binding responses.
MFI = median fluorescent intensity. Binding of IgG1, IgG2a, IgG2b, IgG3, FcγR2b-binding IgGs,
FcγR3-binding IgGs, FcγR4-binding IgGs, and total IgG to the indicated spikes, RBDs, or non-

sarbecovirus control proteins. Antigen names (y-axes) are colored according to spike or RBD
clades. Matched antigens are indicated with gray shading around the name, and mismatched
antigens are indicated with a black outline around the name. Each column represents binding data
from an individual mouse, and immunization cohorts are separated by a vertical black line. A
soluble influenza hemagglutinin trimer (CA-09 sHA) was used as a control to evaluate non-
specific binding. See also Extended Data Fig. 5.
429
Systems Serology-Polyclonal Epitope Mapping (SySPEM)
To further explore potential differences in RBD recognition properties by different IgG subclasses
within polyclonal Ab samples, we combined systems serology with epitope mapping in a new
methodology we call Systems Serology-Polyclonal Epitope Mapping (SySPEM) (Extended Data
Fig. 6), by analogy to Electron Microscopy-based Polyclonal Epitope Mapping[54] (EMPEM). For
the epitope mapping aspect of SySPEM, we created epitope knock-out (KO) mutants in the SARS-
2 WA1 RBD by introducing one or more potential N-linked glycosylation sites (PNGSs) to direct
addition of N-glycan(s) in RBD epitopes that are recognized by either class 1 and class 2, class 2
only, class 3, class 1/4, class 4, or class 5 anti-RBD mAbs (Extended Data Fig. 7a). Addition of
N-glycan(s) at the introduced PNGS(s) was verified by SDS-PAGE of purified RBD KO proteins,
and correct folding and epitope blocking were verified by ELISA using a panel of control mAbs
(Extended Data Fig. 7b,c). We measured binding of total polyclonal IgG and IgG subsets to
unmodified (wildtype; wt) WA1 RBD and to each of RBD KO mutants, defining a SySPEM score
for each IgG sample/epitope pair as [1.0 – (RBD KO binding / RBD wt binding) x 100]; thus,
higher SySPEM scores indicate increased binding by a particular IgG class to that epitope
(Extended Data Fig. 6). We present SySPEM results in three formats to visualize epitope
recognition across immunogens and antibody subclasses: (*i*) Median SySPEM scores from
individual mice divided into distinct RBD epitopes recognized by different IgG classes with
statistical comparisons between immunogen cohorts (Fig. 6). This comparison highlights
differences in epitope recognition across immunogen cohorts. (*ii*) Median SySPEM scores for each
IgG class within a given immunogen cohort with statistical comparisons between targeted epitopes
(Extended Data Fig. 8). This shows the epitope distribution for a given immunogen/IgG class pair,
allowing for epitope profile comparisons between different immunogens, analogous to a DMS
profile (e.g., Fig. 4). (*iii*) Uniform Manifold Approximation and Projection (UMAP)[55] format to
reduce the dimensions of the data set (Fig. 7). These figures provide hierarchical views of the
SySPEM data: statistical comparisons across immunogens (Fig. 6), detailed within-cohort analyses
(Extended Data Fig. 8), and global similarity mapping of antibody recognition patterns (Fig. 7).
457
458 SySPEM scores validated the epitope mapping approach using RBD KO mutants by showing
459 significantly higher scores for total IgGs elicited by homotypic SARS-2 RBD-eVLPs than by other
460 immunogens against variable epitopes (class 1/class 2, class 2) and little to no targeting of more
461 conserved epitopes (class 4, class 1/4, class 5) (Fig. 6), as previously observed by DMS for
462 homotypic SARS-2 RBD-NPs[3].
463
For IgG1, the mRNA- and protein-based dual quartet immunogens showed the highest class 5
SySPEM scores, with only dual quartet RBD-mRNA exhibiting a score that was not significantly
higher than that of homotypic SARS-2 RBD-eVLPs (Fig. 6). In addition, dual quartet RBD-eVLPs
and dual quartet RBD-EABR-mRNA exhibited class 4 and class 1/4 IgG1 SySPEM scores that
were significantly higher than respective scores for SARS-2 RBD-eVLPs. Similarly, for IgG2a,
the dual quartet immunogens all elicited significantly higher class 5 responses than homotypic
SARS-2 RBD-eVLP, with both dual quartet RBD-EABR-mRNA and dual quartet RBD-eVLP
featuring the highest class 5 SySPEM scores. Dual quartet RBD-EABR-mRNA, dual quartet RBD-

NP, and dual quartet RBD-eVLP also elicited higher class 4 SySPEM scores than SARS-2 RBD-
eVLP. Interestingly, although not significant, dual quartet RBD-mRNA elicited a higher class
1/class 2 SySPEM score than dual quartet RBD-EABR-mRNA, and the opposite trend was
observed for class 5. These differences might explain the more potent SARS-2 D614G
neutralization elicited by dual quartet RBD-mRNA compared to dual quartet RBD-EABR-mRNA
(Fig. 2c), since class 1 and class 2 anti-RBD Abs are potent against SARS-2 WA1[44]. However, the
combination of higher class 5 and lower class 1/class 2 responses elicited by the dual quartet RBD-
EABR-mRNA immunogen represents a more favorable epitope targeting profile for broad pan-
sarbecovirus vaccine efforts. For IgG2b and IgG3, SySPEM scores were more similar across the
different RBD epitope categories, although dual quartet RBD-eVLP showed significantly higher
class 4 and class 5 SySPEM scores than SARS-2 RBD-eVLP.
483
For total IgG, dual quartet RBD-EABR-mRNA and dual quartet RBD-NP showed higher class 5
SySPEM scores than SARS-2 RBD-eVLP, with dual quartet RBD-eVLP showing the highest
scores, and dual quartet RBD-mRNA showing class 5 scores that were not significantly higher
than those exhibited by SARS-2 RBD-eVLP (Fig. 6). Dual quartet RBD-EABR-mRNA exhibited
significantly higher class 4 and class 5 SySPEM scores for total IgG than dual quartet RBD-
mRNA, likely driven by the stronger class 4 and class 5 targeting by IgG2a for this comparison
(Fig. 6). Class 4 responses were higher for dual quartet RBD-EABR-mRNA, dual quartet RBD-
NP, and dual quartet RBD-eVLP all exhibiting significantly higher class 4 SySPEM scores than
SARS-2 RBD-eVLP, with dual quartet RBD-eVLP showing the highest class 4, 1/4, and 5 scores.
493
FcγR2b- and FcγR3-binding IgG SySPEM scores were similar across the RBD epitopes, with dual
quartet RBD-EABR-mRNA, dual quartet RBD-NP, and dual quartet RBD-eVLP all exhibiting
significantly higher class 5 SySPEM scores than SARS-2 RBD-eVLP (Fig. 6). Dual quartet RBD-
eVLP and, to a lesser extent, dual quartet RBD-EABR-mRNA exhibited higher class 4 SySPEM
scores than SARS-2 RBD-eVLPs. For FcγR4-binding IgGs, dual quartet RBD-EABR-mRNA,
dual quartet RBD-NP, and dual quartet RBD-eVLP showed significantly higher class 4 and class
5 SySPEM scores than SARS-2 RBD-eVLP.
501
To identify all epitopes recognized by sera elicited by a given immunogen, we displayed the
SySPEM results in a format in which SySPEM scores for recognition of individual RBD epitopes
by different IgG classes are shown (Extended Data Fig. 8). These results highlight increased
targeting of class 5 RBD epitopes by IgG1, IgG2a, IgG2b, and IgG3 subclasses elicited by the four
dual quartet immunogens, but not by homotypic SARS-2 RBD-eVLPs. Interestingly, displaying
the data in this format showed that in some cases for a given immunogen, different IgG subclasses
can exhibit distinct epitope profiles, e.g., IgG1 and IgG2a from dual quartet EABR-mRNA cohort
sera, where IgG1 showed higher class 1 and class 2 targeting than IgG2a, for which only class 5
targeting was high.
511
In order to visualize global patterns of epitope recognition between the different immunogens, we
made UMAP plots[55] to project epitope profiles into two- or three-dimensional plots (Fig. 7). For
2D UMAP plots (Fig. 7a-e), clusters of points indicate samples that behaved similarly across all
epitope knockouts, with ellipses indicating the spread of responses from each immunogen, and
centroid markers (indicated by an X) indicating the average epitope mapping profile for each
immunogen. When comparing IgG1 (Fig. 7a), IgG2a (Fig. 7b), IgG1, IgG2a, IgG2b, and IgG3
(Fig. 7c), and total IgG (Fig. 7d), epitope profiles for different immunogens showed similar trends
with SARS-2 RBD-eVLP clustering farthest from the four dual quartet immunogens, and the dual

quartet immunogens showing more overlap with each other. Interestingly, dual quartet RBD-
mRNA mapped closer to SARS-2 RBD-eVLP in all cases (Fig. 7a-e), as is especially apparent for
IgG2a (Fig. 7b), total IgG (Fig. 7d), and the FcγRs (Fig. 7e). By contrast, dual quartet RBD-EABR-
mRNA clustered further away from SARS-2 RBD-eVLP and closer to dual quartet RBD-eVLP.
Dual quartet RBD-NP mapped similarly to dual quartet RBD-EABR-mRNA. To further assess
differences in epitope recognition between different immunogens, a three-dimensional UMAP plot
was generated using all data (Fig. 7f). Trends in the 3D map were consistent with the 2D UMAPs
in that responses to the dual quartet immunogens were distinct from responses to SARS-2 RBD-
eVLP and responses to dual quartet RBD-EABR-mRNA tended to map between dual quartet
RBD-mRNA and the protein-based dual quartets, consistent with the EABR vaccine modality that
combines traditional mRNA- and protein-based vaccinations[8]. In addition, the finding that
SySPEM scores for dual quartet RBD-mRNA overlapped more with scores for SARS-2 RBD-
eVLP than scores for the other dual quartet immunogens is consistent with more class 1/class 2
RBD epitope targeting by dual quartet RBD-mRNA and SARS-2 RBD-eVLP (Table 1).
534
Overall, the SySPEM studies confirmed targeting of more conserved RBD epitopes by the mRNA-
based dual quartet immunogens, particularly the EABR-modified version, indicating successful
conversion of the desirable properties of protein-based mosaic-8 immunogens. With detailed
analyses, however, we found distinct epitope profiles for IgG subclass and FcγR-binding IgGs:
i.e., the same antigens presented via different modalities (mRNA plus and minus EABR sequences
and protein antigens on NPs or eVLPs) showed divergent SySPEM epitope profiles. In addition,
SySPEM profiling of binding data can rationalize differences in neutralization profiles when
comparing two similar immunogens presented differently (e.g., dual quartet RBD-EABR-mRNA
versus dual quartet RBD-mRNA).
544
545

546
547
Figure 6. SySPEM score comparisons show differences in epitope recognition across
immunogen cohorts.
SySPEM scores from individual mice were determined for RBD epitopes (columns) recognized
by different IgG classes (rows, IgG subclass plus different FcγR-binding IgGs) with statistical
comparisons between immunogen cohorts (colors). A SySPEM value of 0 indicates that none of
the IgGs in that sample were affected by the glycan addition and therefore the sample did not
contain IgGs that recognize that epitope, and a SySPEM value of 100 indicates that all IgGs in that

sample recognized that epitope (Extended Data Fig. 6). A SySPEM value between 0 and 100
indicates the proportion of IgGs in a sample that recognized the epitope that was blocked by glycan
addition in the RBD KO mutant. Box and whisker plots of SySPEM scores with individual data
points representing one mouse are shown. Significant differences between cohorts were calculated
using Tukey's HSD posthoc test and linked by vertical lines indicated by asterisks: $p<0.05 = *$,
$p<0.01 = **$, $p<0.001 = ***$, $p<0.0001 = ****$. See also Extended Data Fig. 8 and Fig. 7.
561

562
563
Figure 7. SySPEM UMAP analyses.
Uniform Manifold Approximation and Projection (UMAP)[55] was used to project multi-
dimensional SySPEM scores into two (panels a-e) or three (panel f) dimensions for (a) IgG1 (b)
IgG2a, (c) IgG1, IgG2a, IgG2b, and IgG3, (d) Total IgG, (e) FcγR-binding IgGs, and (f) Total IgG,
FcγR-binding IgGs, and IgG subclasses. Samples with similar antibody–epitope recognition
profiles cluster together, with colored ellipses indicating the variance within each immunogen
group and centroid markers (large Xs) showing the group mean. (f) Three-dimensional UMAP
projection for all IgG subclasses and FcγR-binding IgGs with centroid markers (large Xs). See also
Fig. 6, Extended Data Figs. 6, 8.
573
Discussion

mRNA-based COVID-19 vaccines have been critical for preventing severe disease and death after
SARS-2 infection[56], thereby helping to keep the worldwide pandemic in check. A recent
modification of Moderna's mRNA-1273 vaccine involved changing the encoded antigen from a
SARS-2 spike trimer to a membrane-bound spike N-terminal domain and RBD[57]. The new
vaccine, mRNA-1283, exhibited equivalent or superior induced immune responses compared to
mRNA-1273 in mice[57] and humans[58]. These results, together with the immunodominance of RBDs
in SARS-2 infections and vaccinations[13], support further development of RBD-based COVID-19
and broader CoV vaccines.

We previously described protein-based RBD-NP immunogens that show promise as a pan-
sarbecovirus vaccine[2-5,7,25,26]. However, manufacturing a protein-based mosaic-8 NP vaccine
presents challenges. For example, mosaic-8 RBD-NP requires expression and purification of nine
protein components (eight RBDs and a NP) to generate a conjugated RBD-NP, which then requires
additional purification and characterization. mRNA vaccines offer the possibility of simpler
manufacturing[59] but developing an mRNA vaccine that delivers eight individual RBD constructs
would also be challenging, especially since mRNA delivery might not achieve co-display of all
eight RBDs on cell surfaces and/or eVLPs.

Here, we report an efficient approach to develop an mRNA-encoded version of a broadly cross-
reactive mosaic-8 RBD vaccine[2-5,7]. The new approach, inspired by protein-based dual quartet
RBD-NPs[7], involves genetically encoding sarbecovirus RBDs arranged as two membrane-bound
quartets, each comprising four tandemly arranged RBD ectodomains. RBD quartets offer the
opportunity for optimal Ab induction by displaying a greater number of RBDs per transmembrane
region compared with presentation of individual RBDs. The mRNA immunogens described here
display dual RBD quartets at cell surfaces (dual quartet RBD-mRNA) or at both the surfaces of
cells and secreted eVLPs (dual quartet RBD-EABR-mRNA) when an EABR cytoplasmic domain
sequence is included[8]. Our analyses showed that dual quartet mRNA immunogens elicited
equivalent or superior Ab responses, improved Fc effector functions, and increased T cell
responses compared with protein-based dual quartet RBD-NPs. In addition, the dual quartet
mRNA immunogens induced balanced T_H1/T_H2 responses, thereby stimulating both cellular and
humoral immune responses. Thus, dual quartet RBD-mRNA immunogens retain the ability of
conventional spike mRNA-LNP vaccines to activate cytotoxic CD8+ T cells and helper CD4+ T
cells that can stimulate Ab-producing B cells[60]. Finally, and of critical importance for a potential
pan-sarbecovirus vaccine, DMS experiments showed that mRNA-based dual RBD quartet
immunogens induced Abs against conserved RBD epitopes, as required for a pan-sarbecovirus
vaccine and as observed for protein-based mosaic-8 RBD-NP immunogens[2-5,7,25,26].

For characterizing dual quartet immunogens, we describe a new systems serology technique,
SySPEM, that quantifies the degree of binding to individual epitopes on an antigen across IgG
subclasses and FcγR-binding IgGs, therefore allowing for comparisons of epitope profiles elicited
by different immunogens that are not possible using traditional systems serology[8]. SySPEM allows
epitope mapping in a more indirect, but higher throughput, way than, e.g., electron microscopy-
based EMPEM[61-64] but can deliver some of the same information while also delineating epitopes
recognized by different Ab classes. In contrast to epitope mapping competition ELISAs[65],
multiplexed SySPEM analyses use minimal sample volumes, do not rely on a panel of mAbs of
known epitopes, and provide more precise epitope definition based on the location of added N-
glycans. In the present study, SySPEM was used to show that the same antigen (dual quartet RBDs)
delivered through mRNA- or protein-based vaccine modalities elicited distinct IgG class-specific

epitope profiles, suggesting that the way an antigen is presented can impact its epitope recognition
by the immune system. Elucidating the mechanism by which this occurs warrants further study.
625
Here, SySPEM showed that, although overall binding and neutralizing Ab titers were similar, the
dual quartet RBD-EABR-mRNA immunogen elicited stronger targeting of conserved class 4 and
class 5 epitopes and weaker targeting of variable class 1/class 2 epitopes than the dual quartet
RBD-mRNA immunogen, suggesting that the EABR approach could improve the effectiveness of
an mRNA-encoded dual quartet RBD pan-sarbecovirus vaccine. For example, immunogen
presentation on circulating eVLPs would promote wider antigen distribution to potentially engage
larger populations of B cells[8,66]. Indeed, circulating RBD-eVLPs resulting from a dual quartet
RBD-EABR-mRNA immunogen offer similar advantages to a recently described mRNA-
delivered RBD protein nanoparticle[67] but also encode membrane-bound antigens at cell surfaces
that could trigger additional immune responses. eVLP budding in the EABR approach might
facilitate continuous replenishment of the cell surface with antigen, potentially increasing overall
levels of antigen being displayed. The small size of eVLPs compared with a cell also facilitates
efficient internalization by antigen-presenting cells[61-64], which could result in enhanced antigenic
peptide presentation by class II MHC and activation of T follicular helper cells to promote B cell
activation and germinal center formation[66]. Improvements for mRNA vaccines by including
EABR appear to be general, as the EABR approach was shown to enhance vaccine-induced
humoral responses against SARS-2[2-5,7], RSV[67], and influenza[22]. We also showed that a bivalent
Wuhan-Hu-1/BA.5 spike-EABR-mRNA booster elicited more balanced targeting of multiple RBD
epitopes in pre-vaccinated mice than conventional monovalent and bivalent spike-mRNA
boosters, which primarily targeted variable RBD epitopes[66]. Taken together, our findings suggest
that EABR-mRNA delivery of mosaic and multivalent immunogens improves targeting of
conserved epitopes to enhance Ab breadth.
648
A pan-sarbecovirus vaccine candidate evaluated in mice would ideally elicit broad-binding IgG2a
and IgG2b subclasses that promote robust Fc effector functions[49,50] to target conserved epitopes
and elicit IgGs that bind to activating Fc receptors (FcγR3 and FcγR4). Using SySPEM, we show
that dual RBD quartet immunogens, including the two mRNA-encoded versions, fulfill these
criteria. There are many future potential applications for this new technology for design and
evaluation of vaccines against a variety of viruses and other pathogens. For example, studies to
establish correlates of protection of vaccines could include evaluations of SySPEM parameters to
determine the relative protective effects of epitope targeting by different classes of IgGs, thus
establishing desirable parameters for future vaccine efforts. In addition, SySPEM can be used to
determine whether some epitopes elicit IgGs with different potential Fc effector functions, again
to inform immunogen design for eliciting optimal immune responses against a particular pathogen.
660
661 In summary, our data support advancement of dual quartet RBD-mRNA immunogens into clinical
662 development as a next-generation booster vaccine to provide protection from emerging SARS-2
663 variants and to reduce the chances of a future zoonotic sarbecovirus spillover from causing a new
664 epidemic or pandemic.
665
666

667 **Methods**
668

Membrane-bound RBD quartet constructs
Sequences of the extracellular domains of RBD quartets 4a and 4b were based on previously
reported SpyTag-Quartet (Genbank PP13603) and SpyTag-Alternate Quartet (GenBank
PP136032, Addgene plasmid ID 214728) sequences[7]. mRNA constructs encoded a mouse Ig heavy
chain signal peptide, codon-optimized RBD quartets 4a and 4b in which individual RBDs were
separated by nine-residue Gly/Ser/Thr linkers followed by a five-residue Gly/Ser linker, the
SARS-1 spike transmembrane domain, and a truncated spike cytoplasmic tail[68] (GenBank
AAP13441.1, residues [1218-1235]). For RBD quartet 4a-EABR and 4b-EABR constructs, an
endocytosis-preventing motif (EPM)[8], four-residue Gly/Ser linker, and the EABR[8] sequence were
appended to the C-terminus. For in vitro assays, these constructs were cloned into the IVTpro T7
plasmid (Takara 6144), containing the immunogen open reading frame flanked by 5′ and 3′
untranslated regions (UTRs) and a genetically encoded poly(A) tail.
681

mRNA synthesis
For in vitro assays, IVTpro T7 plasmids encoding codon-optimized RBD quartet 4a, RBD quartet
4b, RBD quartet 4a-EABR, and RBD quartet 4b-EABR constructs were linearized by BspQI
restriction digestion (NEB, R0712) and subsequently purified using the Zymo DNA Clean &
Concentrator-25 kit (Zymo, D4034). mRNA was generated by in vitro transcription using the
IVTpro™ T7 mRNA Synthesis Kit (Takara, 6144) following manufacturer's instructions, with the
addition of the CleanCap Reagent AG (3' OMe) (TriLink, N-7413) and complete substitution of
uridine with N1-Methylpseudouridine-5'-Triphosphate (TriLink, N-108; TriLink) to reduce
immunogenicity[8]. mRNA was then purified by lithium chloride precipitation and an ethanol wash
following standard protocols[69]. Purified mRNA was resuspended in nuclease-free water and stored
at –80 °C.
693

For mRNAs used for in vivo immunization studies, constructs were synthesized by RNAcore
(https://www.houstonmethodist.org/research-cores/rnacore/) using proprietary manufacturing
protocols including incorporation of the CleanCap Reagent AG (3' OMe) (TriLink, N-7413) and
complete substitution of uridine with N1-Methylpseudouridine-5'-Triphosphate (TriLink, N-108).
mRNAs were purified by oligo-dT affinity purification. To remove double-stranded RNA
contaminants, mRNAs were further purified using a cellulose-based method as described[70]. In
brief, cellulose fibers (0.2 g/mL) were equilibrated in chromatography buffer (10mM HEPES pH
7.2, 0.1 mM EDTA, 125 mM NaCl, 16% ethanol) and added to microcentrifuge spin columns.
mRNA diluted in chromatography buffer was added and shaken vigorously at room temperature
for 30 minutes. The flow-through containing single-stranded mRNA was collected and purified
with a sodium acetate precipitation and an ethanol wash. The final purified mRNA was filtered
through a 0.2 µm filter before storage at –80 °C[8,70].
mRNA transfections and flow cytometry
HEK293T cells were seeded at 1×10^6 cells per well in 6-well plates and incubated for 20 h before
transfection with 2 µg of mRNA encoding RBD quartet 4a, RBD quartet 4b, and/or their EABR-
fused counterparts (4a-EABR, 4b-EABR) using Lipofectamine™ MessengerMax™ (Thermo
Fisher Scientific). Forty-eight hours post-transfection, supernatants were collected for eVLP
purification by ultracentrifugation on a 20 % sucrose cushion as described[8]. Cells were gently
detached, resuspended in PBS+ (PBS + 2 % FBS), and 100 µL aliquots were taken for flow
cytometry.

For the experiment shown in Extended Data Fig. 1a, cells were stained with the cross-reactive anti-
RBD mAb C118[19,20] (2.5 μg/mL) for 30 min in the dark at room temperature, washed, and
subsequently stained with Alexa Fluor® 647-conjugated anti-human IgG secondary antibody
(Invitrogen, A21445; 1:4,000) for 15 min. After a final wash, cells were resuspended in PBS+ and
analyzed on an Attune NxT Flow Cytometer (Thermo Fisher).
719
For the experiment shown in Extended Data Fig. 1c, cells were either singly transfected or co-
transfected with combinations of RBD quartet 4a and 4b constructs and stained simultaneously
with 2.5 μg/mL of Alexa Fluor® 647-conjugated anti-Rs4081 RBD[25] and Alexa Fluor® 488-
conjugated M8a-7 IgG (anti-WIV1 RBD)[26] mAbs for 30 min at room temperature in the dark.
After washing and resuspension, samples were analyzed on an Attune NxT Flow Cytometer.
Results were plotted using FlowJo 10.5.3 software.
726

Production of dual quartet and SARS-2 RBD-eVLPs

Dual quartet RBD-eVLP and SARS-2 RBD-eVLP samples were generated as described[8] by
transfecting Expi293F cells (Gibco) cultured in Expi293F expression media (Gibco) on an orbital
shaker at 37°C and 8% CO_2. To generate dual quartet RBD-eVLPs, DNA plasmids encoding RBD
quartet 4a-EABR and RBD quartet 4b-EABR constructs were mixed at a ratio of 1:1. 72 hours
post-transfection, cells were centrifuged at 400 x g for 10 min, supernatants were passed through
a 0.45 μm syringe filter and concentrated using Amicon Ultra-15 centrifugal filters with 100 kDa
molecular weight cut-off (Millipore). eVLPs were purified by ultracentrifugation at 50,000 rpm
(135,000 x g) for 2 hours at 4°C using a TLA100.3 rotor and an Optima™ TLX ultracentrifuge
(Beckman Coulter) on a 20% w/v sucrose cushion. After removal of supernatants by aspiration,
pellets were re-suspended in 200 μL sterile PBS at 4°C overnight. To remove residual debris,
samples were centrifuged at 10,000 x g for 10 min, and supernatants were collected. eVLPs were
further purified by SEC using a Superose 6 Increase 10/300 column (Cytiva) equilibrated with
PBS. Peak fractions corresponding to eVLPs were combined and concentrated to 250-500 μL in
Amicon Ultra-4 centrifugal filters with 100 kDa molecular weight cut-off. Samples were aliquoted
and stored at -20°C.
743

The amount of SARS-2 RBD on SARS-2 RBD-eVLPs and the amount of dual quartet RBD on
dual quartet RBD-eVLPs were determined by quantitative Western blot analysis. Various dilutions
of SEC-purified eVLP samples and known amounts of either soluble SARS-2 RBD protein (Sino
Biological, 40591-V08H-B-20) or purified soluble SpyTagged dual quartet RBDs (SpyTagged
RBD quartets 4a and 4b were mixed at a ratio of 1:1) were separated by SDS-PAGE and transferred
to nitrocellulose membranes (0.45 μm) (Thermo Fisher Scientific, LC2001). Rabbit anti-SARS-2
S1 polyclonal IgG (Thermo Fisher Scientific, PA5-116916; 1:1,000) and HRP-conjugated goat
anti-rabbit IgG (Abcam, ab98467; 1:10,000) were used. Protein bands were visualized using ECL
Prime Western Blotting Detection Reagent (Cytiva, RPN2232). Band intensities of the respective
standards and eVLP sample dilutions were measured using ImageJ to determine RBD amounts.
754

LNP encapsulation of mRNAs

Purified dual quartet RBD-encoding mRNAs (RBD quartet 4a + RBD quartet 4b or RBD quartet
4a-EABR + RBD quartet 4b-EABR; 1:1 ratio) for in vivo studies were formulated in LNPs as
previously described[71]. 1,2-distearoyl-*sn*-glycero-3-phosphocholine, cholesterol, a PEG lipid, and
an ionizable cationic lipid dissolved in ethanol were rapidly mixed with an aqueous acidic solution
containing mRNA using an in-line mixer. The ionizable lipid and LNP composition are described
in the international patent application WO2017075531(2017). The post in-line solution was
dialyzed with PBS to remove the ethanol and displace the acidic solution. LNP was measured for

size (60-65 nm) and polydispersity (PDI < 0.075) by dynamic light scattering (Malvern Nano ZS
Zetasizer). Encapsulation efficiencies were >97% as measured by the Quant-iT Ribogreen Assay
(Invitrogen).
766
Protein expression
RBDs, RBD quartets, and soluble sarbecovirus spike-6P[72] (SARS-2 WA1 and JN.1) or spike-2P
(XBB.1.5, SARS-1, Rf1, Rs4091, Yun11, BM48-31, BtKY72) trimers were expressed as
described[3,6,7]. Briefly, AviTag and/or His-tagged proteins were purified from transiently-
transfected Expi293F cells (Gibco) by nickel affinity chromatography (HisTrap HP, Cytiva) and
SEC (Superose 6 Increase 10/300, Cytiva)[3,44,73]. Fractions corresponding to proteins of interest
were pooled, concentrated, and stored at 4ºC. Biotinylated proteins were generated as described[74]
by co-transfection of AviTag/His-tagged spike and RBD constructs with a plasmid encoding an
endoplasmic reticulum-directed BirA enzyme (kind gift from Michael Anaya, Caltech). The anti-
Rs4081, anti-RmYN02, and anti-WIV1 RBD mAb IgGs from Extended Data Fig. 1 and the mAb
IgGs and human ACE2-Fc protein shown in Extended Data Fig. 7c were expressed, purified, and
classified as recognizing the RBD epitopes as described[20,25,26,44].
779
Preparation of RBD-NPs
For making dual quartet RBD-NPs, SpyCatcher003-mi3–C-Tag (Addgene 159995) subunits were
expressed in BL21 (DE3) *E. coli* (Agilent) as described[75]. SpyCatcher003-mi3–C-Tag bacterial
cell pellets were resuspended in 20 mL of 20 mM Tris-HCl pH 8.5, 300 mM NaCl, supplemented
with 0.1 mg/mL lysozyme, 1 mg/mL cOmplete mini EDTA-free protease inhibitor (Roche), and
1.0 mM PMSF. After incubation at 4 °C for 45 min with end-over-end mixing, an Ultrasonic
Processor equipped with a microtip (Cole-Parmer) was used to perform sonication on ice (four
times for 60 s, 50% duty-cycle), and cell debris were cleared by centrifugation (35,000 x g for
45 min at 4 °C). SpyCatcher003-mi3–C-Tag was purified by ammonium sulfate precipitation (170
mg per mL of lysate) and resuspension, followed by SEC using a HiPrep Sephacryl S-400 HR 16-
60 column (GE Healthcare) equilibrated with PBS as described[75]. Dual quartet RBD-NPs were
generated by incubating purified SpyCatcher003-mi3–C-Tag with a 1:1 molar ratio of purified
SpyTagged RBD quartets at 4 ºC in PBS.
793
Immunization of mice
Mouse procedures were approved by the Labcorp Institutional Animal Care and Use Committee.
Six- to 7-week-old female BALB/c mice from Charles River Laboratories (RRID:
IMSR_JAX:000664) were housed at Labcorp Drug Development, Denver, PA for immunizations.
All animals were healthy upon receipt and monitored during a 7-day acclimation period. Mice
were assigned randomly to experimental groups of 10 animals. Mouse cages were kept in a
climate-controlled room (68 - 79 ºC) at 50 ± 20% relative humidity and provided with Rodent Diet
#5001 (Purina Lab Diet) ad libitum.
802
Mice were immunized by intramuscular injection on days 0 and 28 with 1.5 or 0.5 µg dual quartet
RBD-mRNA, 1.5 or 0.5 µg dual quartet RBD-EABR-mRNA, 5 µg dual quartet RBD-NP, 5 µg
dual quartet RBD-eVLP, or 5 µg SARS-2 RBD-eVLP (based on RBD content for protein-based
immunogens). Prior to immunizations, dual quartet RBD-NP, dual quartet RBD-eVLP, and SARS-
2 RBD-eVLP immunogens were mixed with Addavax adjuvant (InvivoGen; 50% v/v). Serum
samples for ELISAs and neutralization assays were obtained on indicated days by retro-orbital
bleeding.
810
ELISAs

For eVLP sandwich ELISAs, 96-well high-binding plates (Corning, #9018) were coated overnight
at 4 °C with capture monoclonal antibody (5 µg/mL in 0.1 M NaHCO₃, pH 9.8). Plates were
blocked with 3% BSA and 0.1% Tween-20 in TBS (TBS-T) for 30 min at room temperature,
followed by addition of serially diluted purified eVLP samples for 2 h. After washing, biotinylated
detection antibodies (5 µg/mL in blocking buffer) were added for 2 h at room temperature. Plates
were washed three times with TBS-T, incubated with streptavidin-HRP (Abcam, 7403; 1:20,000)
for 30 min, washed again, and developed with 1-Step™ Ultra TMB-ELISA substrate solution
(ThermoFisher). Reactions were stopped with 1 N HCl and absorbance was measured at 450 nm.

For the ELISA in Extended Data Fig. 1b, plates were coated with either anti-Rs4081[25] or anti-
RmYN02[25] mAbs and detected with biotinylated C118[19,20]. For the ELISA in Extended Data Fig.
1d, plates were coated with anti-WIV1[26] and detected with biotinylated anti-Rs4081[25].

For serum ELISAs, purified His-tagged RBD (2.5 µg/mL in 0.1 M sodium bicarbonate buffer pH
9.8) was coated onto Nunc® MaxiSorp™ 384-wellplates (Sigma) and incubated overnight at 4˚C.
After blocking with 3% bovine serum albumin (BSA), 0.1% Tween 20 in Tris-buffered saline
(TBS-T) for 1 hour at room temperature, blocking solution was removed by aspiration. For serum
ELISAs shown in Fig. 2 and Extended Data Fig. 2, mouse serum (1:100 initial dilution) was
serially diluted by 3.1-fold in TBS-T/3% BSA and then added to plates for 3 hours at room
temperature, followed by washing with TBS-T. Plates were then incubated for 1 hour with a
1:100,000 dilution of HRP-conjugated anti-IgG secondary (goat anti-mouse IgG; Abcam; RRID:
AB_955439), and plates were washed with TBS-T. For mAb ELISAs shown in Extended Data
Fig. 7, purified mAbs were serially diluted by 3.1-fold in TBS-T/3% BSA and then added to plates
for 3 hours at room temperature, followed by washing with TBS-T. Plates were then incubated for
1 hour with a 1:100,000 dilution of HRP-conjugated anti-IgG secondary (goat anti-human IgG;
SouthernBiotech 2014-05), and plates were washed with TBS-T. SuperSignal™ ELISA Femto
Substrate (ThermoFisher) was added as per manufacturer's instructions, and luminescence was
read at 425 nm. Data were collected in duplicate and midpoint titers (ED$_{50}$ values) were obtained
using Graphpad Prism 10.5.0 assuming a four-parameter dose-response curve fit.

Pseudovirus neutralization assays
Lentivirus-based pseudoviruses were produced in HEK293T cells (RRID:CVCL_0063) cultured
in Dulbecco's modified Eagle's medium (DMEM, Gibco) supplemented with 10% heat-
inactivated fetal bovine serum (FBS, Bio-Techne), 1% Penicillin/Streptomycin (Gibco), and 1%
L-Glutamine (Gibco) at 37 °C and 5% CO₂. HEK293T-hACE2 cells (RRID:CVCL_A7UK)[43] and
high-hACE2 HEK-293T (kindly provided by Kenneth Matreyek, Case Western Reserve
University) target cells for neutralization assays were cultured in DMEM (Gibco) supplemented
with 10% heat-inactivated FBS (Bio-Techne), 5 mg/mL gentamicin (Sigma-Aldrich), and 5
mg/mL blasticidin (Gibco) at 37 °C and 5% CO₂.
848
Lentiviral-based viruses were prepared as described[19,76] using genes encoding spike sequences
lacking C-terminal residues in the cytoplasmic tail: deletions of 21 residues (SARS-2 variants) or
19 residues (SARS-1, Khosta-2-SARS-1 chimera, BtKY72-SARS-1 chimera). BtKY72
(containing K493Y/T498W substitutions) and Khosta-2 pseudoviruses were made with chimeric
spikes in which the RBD from SARS-1 (residues 323-501) was substituted with the RBD from
BtKY72 K493Y/T498W (residues 327-503) or Khosta-2 (residues 324-500) as described.[77]. Cells
were co-transfected with HIV-1-based lentiviral plasmids, a luciferase reporter gene, and a
coronavirus spike construct, resulting in lentivirus-based pseudovirions expressing a sarbecovirus

spike protein. Supernatants were harvested 48-72 hours post-transfection, filtered, and stored at -
80 °C. Pseudovirus infectivity was determined by titration using HEK293T-hACE2 cells.
859
Prior to neutralization assays, serum from immunized mice was heat-inactivated at 56 ºC for 10
minutes. Heat-inactivated serum was three-fold serially diluted and incubated with pseudovirus
for 1 hour at 37˚C, then the serum/virus mixture was added to HEK293T-hACE2 target cells or
high-hACE2 HEK-293T cell line expressing hACE2 encoded with a consensus Kozak sequence
(for SHC014 assays; kindly provided by Kenneth Matreyek, Case Western Reserve University)
and incubated for 48 hours at 37ºC. After removing media, cells were lysed with Britelite Plus
reagent (Revvity Health Sciences), and luciferase activity was measured as relative luminesce units
(RLUs). Relative RLUs were normalized to RLUs from cells infected with pseudotyped virus in
the absence of antiserum. Half-maximal inhibitory dilutions (ID_{50} values) were derived in
AntibodyDatabase[78] using 4-parameter nonlinear regression.
870
ELISpot assays
Mice were euthanized on day 77 or 78, and spleens were collected and processed as described[8].
Briefly, spleens were homogenized using a gentleMACS Octo Dissociator (Miltenyi Biotec). Cells
were passed through a 70 µm tissue screen, centrifuged at 1,500 rpm for 10 min, and resuspended
in CTL-Test™ media (ImmunoSpot) containing 1% GlutaMAX™ supplement (Gibco). ELISpot
assays were performed as described[8]. A SARS-CoV-2 spike PepMix™ pool of 315 peptides (15-
mers with 11 amino acid overlap; JPT Peptide Technologies) was added to mouse IFNγ/IL4
double-color ELISpot plates (ImmunoSpot) at a concentration of 2 µg/mL, and 300,000
splenocytes were added per well. Plates were incubated at 37ºC for 24 hours. Biotinylated
detection reagents, streptavidin-alkaline phosphatase (AP), and substrate reagents were added
according to the manufacturer's guidelines. To stop the reactions, plates were gently rinsed with
water three times. Plates were air-dried for two hours in a running laminar flow hood. Numbers of
spots were quantified using a CTL ImmunoSpot S6 Universal-V Analyzer (Immunospot).
884
DMS Epitope Mapping
DMS experiments to map epitopes recognized by serum Abs were performed in biological
duplicates using independent mutant RBD yeast libraries (WA1[79] and SARS-1[80] generously
provided by Tyler Starr, University of Utah) as described[3,41]. Sera that had been heat inactivated
for 30 min at 56 °C were incubated twice with 50 OD units of AWY101 yeast transformed with
an empty vector in order to remove non-specific yeast-binding Abs. Expression of RBDs in
libraries was induced in galactose-containing synthetic defined medium with Casamino acids
(6.7g/L Yeast Nitrogen Base, 5.0 g/L Casamino acids, 1.065 g/L MES acid, and 2% w/v galactose
plus 0.1% w/v dextrose). After 18 hours of induction, cells were washed twice and incubated with
serum under conditions of gentle agitation for 1 hour at room temperature. Cells were labeled for
1 hour with a secondary Ab (1:200 Alexa Fluor-647-conjugated goat anti-mouse-IgG Fc-gamma,
Jackson ImmunoResearch 115-605-008, RRID:AB_2338904) after washing twice.
897
Stained yeast cells were examined by fluorescence-activated cell sorting (FACS) using a Sony
SH800 cell sorter. Cells were gated to capture RBD mutants that showed reduced Ab binding
compared to a control. Cells were collected for each sample until ~5 x 10^6 RBD+ cells were
processed, which corresponds to ~5 x 10^5- 1 x 10^6 RBD+ Ab-escaped cells. These cells were grown
overnight in synthetic defined media (6.7 g/L Yeast Nitrogen Base, 5.0 g/L Casamino acids, 1.065
g/L MES acid, and 2% w/v dextrose, 100 U/mL penicillin, 100 µg/mL streptomycin) to expand
cells prior to plasmid extraction. DNA extraction and Illumina sequencing were done as described[7]
(raw sequencing data will be available on NCBI SRA). Escape fractions were computed as

described[7,41] using Swift DMS[7] (available upon request). Escape scores were calculated using a
filter to remove variants with mutations that escaped binding because of poor expression, >1 amino
acid mutation, or low sequencing counts[7,81].

Escape map visualizations (static line plots, logo plots, and structural depictions) were created
using Swift DMS[7]. Line heights show the escape score for a particular amino acid substitution, as
described[7]. In some visualizations, sites were categorized based on RBD epitope region[20,44-46]:
class 1 (pink; RBD residues 403, 405, 406, 417, 420, 421, 453, 455-460, 473-478, 486, 487, 489,
503, 504); class 2 (purple; residues 472, 479, 483-485, 490-495), class 3 (blue; residues 341, 345,
346, 437-450, 496, 498-501,), class 4 (orange; residues 365-390, 408), class 5 (green; residues
352-357, 396, 462-468). In structural depictions of DMS results, an RBD surface (PDB 6M0J)
was colored by the site-wise escape metric at each site, with red scaled at an escape fraction of 2.0.
Residues exhibiting the highest escape fractions were labeled with their residue number, which
was colored according to epitope class. Logo plot residues are colored according to RBD epitopes
within different classes as indicated on the legend.

We stratified DMS escape fraction values into four groups. Escape fractions for each RBD
substitution range from 0 (no cells with this substitution were sorted into the escape bin) to 1 (all
cells with this substitution were sorted into the serum Ab escape bin)[40]. The sum of escape fractions
for all substitutions at a specific site is represented by the total escape peak[40]. DMS profiles with
total escape peaks <0.5 at all sites were classified as polyclass responses. DMS profiles with total
escape peaks of 0.5 to 1, >1 to 2, or >2 at one or more sites were classified as weak, moderate, or
strong escape profiles, respectively, corresponding to their RBD epitope (Table 1).

Systems serology: Ab subclass and FcγR binding profiling
Serum samples from immunized mice were analyzed in systems serology[9] experiments as
described[6] using modifications of a Luminex assay to quantify the levels of antigen-specific Ab
subclasses and FcγR binding profiles, which allows for simultaneous detection and quantification
of multiple analytes (i.e., different antigens) in a single sample[82]. In brief, avidin (Sigma-Aldrich
Catalog #: A9275-25MG) was coupled by carbodiimide-NHS ester chemistry to magnetic
Luminex microspheres (LuminexCorp) using Sulfo-NHS (ThermoFisher™ Catalog number
24510) and 1-Ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC)
(ThermoFisher™ Catalog number 22980) according to the manufacturer's instructions.
Microspheres were blocked for 30 min with 1% BSA in PBS pH 7.4 (assay buffer) and then washed
twice with PBS. Microspheres were incubated with biotinylated antigen (sarbecovirus spike trimer,
RBD, or control antigen) at room temperature for 2 hours in assay buffer followed by blocking
with 10 µM biotin (Millipore Sigma, Catalog B4501-100MG). Antigen-coupled microspheres
were combined then incubated with heat-inactivated serum samples for 1 hour at room temperature
in 96-well plates (Corning) at 1:200 or 1:1000 dilutions. Unbound IgGs were removed by washing
twice with 200 µL of assay buffer. Total IgG was detected using a PE-conjugated secondary Ab
recognizing mouse IgG (Southern Biotech 1030-09S). For evaluating antigen binding by different
IgG subclasses, secondary Abs against mouse IgG subclasses (Southern Biotech 1070-09S, 1080-
09S, 1090-09S, 1100-09S; PE-coupled anti-IgG1, IgG2a, IgG2b, IgG3) were added at a 1:1000
dilution in assay buffer and incubated for 1 hour with continuous shaking at room temperature.
Excess primary and secondary Abs were removed by washing twice with 200 µL of assay buffer.
Beads were resuspended in 200 µL of assay buffer, run in duplicate, and flowed in single file past
two lasers on a Luminex™ FLEXMAP 3D™ Instrument System. In a Luminex assay, each bead
region is impregnated with a different ratio of fluorescent dyes such that bead region identity, and

thereby antigen identity, is interrogated by the first laser. A second laser interrogates the signal of
secondary Ab binding. Median fluorescence intensity (MFI) was calculated for all samples.
956
For evaluating FcγR-binding IgGs, soluble ectodomains of 6xHis-tagged FcγR2b, FcγR3, and
FcγR4 were prepared as described[6] and biotinylated by co-expression with BirA enzyme in
Expi293T cells[74]. Biotinylated FcγRs were purified on a HisTrap column (VWR) according to the
manufacturer's instructions followed by SEC and then bound to PE-streptavidin (eBioscience).
Labeled FcγRs were diluted 1:200 in assay buffer and incubated with serum-coated microspheres
for 1 hour at room temperature with continuous shaking. Unbound primary and PE-labeled FcγR
were removed as described for anti-IgG subclass secondary Abs. Beads were resuspended and run
on a Luminex™ FLEXMAP 3D™ Instrument System to derive MFI values as described above.
965
966 Log$_{10}$-transformed heatmaps of antigen-specific IgG responses were generated using Python
967 (v3.9.16) with the Pandas (v2.0.3), NumPy (v1.23.5), Matplotlib (v3.7.1), and Seaborn (v0.12.2)
968 packages. A soluble influenza hemagglutinin trimer (listed as CA-09 HA in Fig. 5; expressed and
969 purified as described[83]) was used as a control to evaluate non-specific binding.
970
SySPEM
One or more potential N-linked glycosylation site sequons (Asn-x-Ser/Thr)[84] were introduced at
solvent-exposed residue(s) within the class 1 and class 2, class 2, class 3, class 4, class 1/4, or class
5 RBD epitopes[20,44-46] of the WA1 RBD (Extended Data Fig. 7a). Addition of N-glycan(s) was
confirmed by SDS-PAGE as a shift in electrophoretic mobility to a higher apparent molecular
weight (Extended Data Fig. 7b). Proper folding of RBD KO mutants and expected effects of added
N-glycans were verified by assaying the binding of characterized mAbs or a human ACE2-Fc
construct[20] by ELISA (Extended Data Fig. 7c). Binding of serum subsets to RBD wt and RBD KO
proteins was measured using a Luminex multiplexed bead-based immunoassay as described above.
The SySPEM score for each IgG sample/epitope pair was calculated as [1.0 – (RBD KO binding
/ RBD wt binding) x 100]. SySPEM scores range from limits of 100 (none of the IgGs in the
sample bound to the RBD KO; i.e., all of the IgGs recognize the epitope that was targeted by the
glycan addition) to 0 (all of the IgGs in the sample bound to the RBD KO; i.e., none of the IgGs
recognize the epitope that was targeted by the glycan addition) (Extended Data Fig. 6). Python
(v3.9.16) with the Pandas (v2.0.3), NumPy (v1.23.5), Matplotlib (v3.7.1), and Seaborn (v0.12.2)
packages were used for box and whisker plots of SySPEM data. For UMAP visualizations in Fig.
7, SySPEM scores of each serum sample were organized into a feature matrix where each column
contains an epitope class SySPEM score, and each row corresponds to an individual sample.
Features were standardized using the standard scalar function on sckit-learn 1.4.2. UMAP was
then performed with umap-learn (0.5.9.post2) on the standardized matrix. UMAP parameters: the
nearest neighbors (n-neighbors) parameter was set to 12 and minimum distance (min-dist) was set
to 0.25 to preserve the best local and global structures on the UMAP plot. Seaborn (v0.12.2) was
used to plot resulting UMAPs.
994
Quantification and statistical analyses
We used pairwise comparisons, a method to evaluate sets of mean binding titers against individual
viral strains for different immunization cohorts, as described previously[4] to determine if results
from different immunized cohorts were significantly different from each other. For ELISAs and
neutralizations (Fig. 2), we used analysis of variance (ANOVA) followed by Tukey's multiple
comparison post hoc tests with the Geisser-Greenhouse correction, with pairing by viral strain, of
ED$_{50}$s/ID$_{50}$s (converted to log$_{10}$ scale) calculated using GraphPad Prism 10.5.0 to identify

statistically significant titer differences between immunized groups for ELISAs. For comparisons of neutralizing titers by strain (Extended Data Fig. 2), statistically significant titer differences between immunized groups for each given strain were determined using unpaired t test calculated using GraphPad Prism 10.5.0. For ELISpot assays, statistically significant differences between immunized cohorts were calculated using ordinary one-way ANOVA followed by Tukey's multiple comparison test using GraphPad Prism.

For statistical analysis of systems serology results, responses were aggregated by computing the geometric mean (geomean) across replicate samples for each immunogen-antigen combination. Log_{10}-transformed geomeans were compared across immunogen groups using Tukey's multiple comparison post hoc tests with the Geisser-Greenhouse correction, with pairing by viral strain calculated using GraphPad Prism 10.5.0. Box-and-whisker plots were generated for each IgG subclass or FcγR-binding IgG, displaying individual antigen-level geomeans per immunogen. Python (v3.9.16) with statsmodels (v0.14.4) packages were used for SySPEM Tukey's HSD posthoc statistical analyses. Statistical significance was annotated using brackets and asterisk notation ($p < 0.05$ *, $p < 0.01$ **, $p < 0.001$ ***, $p < 0.0001$ ****).

Materials, data, and code availability
DMS raw sequencing data will be available on the NCBI SRA under BioProject PRJNA1067836, BioSample when the NCBI SRA is back online. SySPEM scores for each IgG sample/epitope pair will be made publicly available upon publication. Materials are available upon request to corresponding authors with a signed material transfer agreement, and other information required to analyze the data in this paper is available from the lead contacts upon request. Code used for data processing and visualization of DMS, systems serology, and SySPEM results is available upon request. The work is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0) license, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. To view a copy of this license, visit https://creativecommons.org/licenses/by/4.0/. This license does not apply to figures/photos/artwork or other content included in the article that is credited to a third party; obtain authorization from the rights holder before using such material.

Acknowledgements
We thank Annie Rorick, Han Gao, and Priyanthi Gnanapragasam for experimental assistance, Jesse Bloom (Fred Hutchinson Cancer Research Center) and Tyler Starr (University of Utah) for RBD libraries, Ryan McNamara and Galit Alter (Ragon Institute) for help with Systems Serology experiments, Jost Vielmetter, Luisa Segovia, Alyssa Player, Annie Lam, and the Caltech Beckman Institute Protein Expression Center for protein production, Igor Antoshechkin and the Caltech Millard and Muriel Jacobs Genetics and Genomics Laboratory for Illumina sequencing, Michael Anaya for help with Luminex experiments, Dom Aylard, Katie Lippert, and the Gladstone Flow Cytometry Core for help with flow cytometry, and Anthony West and Ying Tam (Acuitas) for critical reading of the manuscript. These studies were funded by Wellcome Leap (P.J.B.), the National Institutes of Health P01-AI165075 (P.J.B.), DP5OD033362 (M.A.G.H.), and 1K99AI185267 (A.A.C.), the Medical Research Council (MRC grant no. MR/Y011910/1 to M.R.H.), the Merkin Institute for Translational Research (Caltech) (P.J.B.), and the James B. Pendleton Charitable Trust that supports the Gladstone Flow Cytometry Core. This manuscript is the result of funding in whole or in part by the National Institutes of Health (NIH). It is subject to the NIH Public Access Policy. Through acceptance of this federal funding, NIH has been given a right to make this manuscript publicly available in PubMed Central upon the Official Date of Publication, as defined by NIH.

Author Contributions

Conceptualization: A.A.C., J.R.K., R.A.H., M.R.H., M.A.G.H., P.J.B.; Methodology: A.A.C., M.A.G.H.; Software: A.A.C.; Investigation: A.A.C., J.R.K., LM, IM, A.-C.P.F., K.-M.A.D., HES, M.A.G.H.; Resources: R.A.H., W.J.M., P.J.C.L.; Writing – original draft: A.A.C., J.R.K., K.-M.A.D., M.A.G.H., P.J.B.; Writing – review and editing: A.A.C., J.R.K., K.-M.A.D., R.A.H., M.R.H., M.A.G.H., P.J.B.; Visualization: A.A.C., J.R.K., K.-M.A.D., M.A.G.H., P.J.B.; Supervision: M.R.H., M.A.G.H., P.J.B.; Funding: M.A.G.H., P.J.B..

P.J.B. and A.A.C. are inventors on a US patent application (17/523,813) filed by the California Institute of Technology that covers mosaic RBD-NPs. M.A.G.H. and P.J.B. are inventors on a patent application (US17/835,751; international: PCT/US2022/032702) filed by the California Institute of Technology that covers the EABR technology. M.R.H. is an inventor on patents on spontaneous amide bond formation (EP2534484 and UK Intellectual Property Office 1706430.4) and a SpyBiotech co-founder and shareholder. P.J.B., M.A.G.H., A.A.C., J.R.K., K-M.A.D., R.A.H., and M.R.H. are co-inventors on a provisional US patent application filed by the California Institute of Technology that covers dual quartet mRNA immunogens. P.J.B. is a scientific advisor for Vaccine Company, Inc. M.A.G.H. is a scientific consultant for Vaccine Company, Inc. W.J.M. and P.J.C.L. are employees of Acuitas Therapeutics, a company developing LNP delivery technology; P.J.C.L. holds equity in Acuitas Therapeutics.

1081 Literature cited
1082

1   Whittaker, C., Barnsley, G., Mesa, D. O., Cox, V., Laydon, D. J., Tan, C. W., Zhu, F.,
Johnson, R., Doohan, P., Whittles, L. K., Nedjati-Gilani, G., Winskill, P., Hogan, A. B.,
Deol, A., Mukandavire, C., Hauck, K., Lye, D. C. B., Wang, L. F., Watson, O. J. & Ghani,
A. C. Quantifying the impact of a broadly protective sarbecovirus vaccine in a future
SARS-X pandemic. *Nat Commun* **16**, 8495 (2025).

2   Cohen, A. A., Gnanapragasam, P. N. P., Lee, Y. E., Hoffman, P. R., Ou, S., Kakutani, L.
M., Keeffe, J. R., Wu, H. J., Howarth, M., West, A. P., Barnes, C. O., Nussenzweig, M. C.
& Bjorkman, P. J. Mosaic nanoparticles elicit cross-reactive immune responses to zoonotic
coronaviruses in mice. *Science* **371**, 735–741 (2021).

3   Cohen, A. A., van Doremalen, N., Greaney, A. J., Andersen, H., Sharma, A., Starr, T. N.,
Keeffe, J. R., Fan, C., Schulz, J. E., Gnanapragasam, P. N. P., Kakutani, L. M., West, A.
P., Jr., Saturday, G., Lee, Y. E., Gao, H., Jette, C. A., Lewis, M. G., Tan, T. K., Townsend,
A. R., Bloom, J. D., Munster, V. J. & Bjorkman, P. J. Mosaic RBD nanoparticles protect
against challenge by diverse sarbecoviruses in animal models. *Science* **377**, eabq0839
(2022).

4   Cohen, A. A., Keeffe, J. R., Schiepers, A., Dross, S. E., Greaney, A. J., Rorick, A. V., Gao,
H., Gnanapragasam, P. N. P., Fan, C., West, A. P., Jr., Ramsingh, A. I., Erasmus, J. H.,
Pata, J. D., Muramatsu, H., Pardi, N., Lin, P. J. C., Baxter, S., Cruz, R., Quintanar-Audelo,
M., Robb, E., Serrano-Amatriain, C., Magneschi, L., Fotheringham, I. G., Fuller, D. H.,
Victora, G. D. & Bjorkman, P. J. Mosaic sarbecovirus nanoparticles elicit cross-reactive
responses in pre-vaccinated animals. *Cell* **187**, 5554–5571 e5519 (2024).

5   Wang, E., Cohen, A. A., Caldera, L. F., Keeffe, J. R., Rorick, A. V., Adia, Y. M.,
Gnanapragasam, P. N. P., Bjorkman, P. J. & Chakraborty, A. K. Designed mosaic
nanoparticles enhance cross-reactive immune responses in mice. *Cell* **188**, 1036–1050
e1011 (2025).

6   Cohen, A. A., Keeffe, J. R., Rorick, A. V., Rho, S., Priso Fils, A.-C., Manasyan, L., Gao,
H., Gnanapragasam, P. N. P., Funke, H. H., Randolph, T. W., Garcea, R. L. & Bjorkman,
P. J. Broad anti-sarbecovirus responses elicited by a single administration of mosaic-8
RBD-nanoparticle vaccine prepared using atomic layer deposition. *iScience* (2025).

7   Hills, R. A., Tan, T. K., Cohen, A. A., Keeffe, J. R., Keeble, A. H., Gnanapragasam, P. N.
P., Storm, K. N., Rorick, A. V., West, A. P., Jr., Hill, M. L., Liu, S., Gilbert-Jaramillo, J.,
Afzal, M., Napier, A., Admans, G., James, W. S., Bjorkman, P. J., Townsend, A. R. &
Howarth, M. R. Proactive vaccination using multiviral Quartet Nanocages to elicit broad
anti-coronavirus responses. *Nat Nanotechnol* **19**, 1216–1223 (2024).

8   Hoffmann, M. A. G., Yang, Z., Huey-Tubman, K. E., Cohen, A. A., Gnanapragasam, P. N.
P., Nakatomi, L. M., Storm, K. N., Moon, W. J., Lin, P. J. C., West, A. P., Jr. & Bjorkman,
P. J. ESCRT recruitment to SARS-CoV-2 spike induces virus-like particles that improve
mRNA vaccines. *Cell* **186**, 2380–2391 e2389 (2023).

9   Ackerman, M. E., Barouch, D. H. & Alter, G. Systems serology for evaluation of HIV
vaccine trials. *Immunol Rev* **275**, 262–270 (2017).

10   Andersen, K. G., Rambaut, A., Lipkin, W. I., Holmes, E. C. & Garry, R. F. The proximal
origin of SARS-CoV-2. *Nature Medicine* **26**, 450–452 (2020).

11   Roederer, A. L., Cao, Y., St Denis, K., Sheehan, M. L., Li, C. J., Lam, E. C., Gregory, D.
J., Poznansky, M. C., Iafrate, A. J., Canaday, D. H., Gravenstein, S., Garcia-Beltran, W. F.
& Balazs, A. B. Ongoing evolution of SARS-CoV-2 drives escape from mRNA vaccine-
induced humoral immunity. *Cell Rep Med* **5**, 101850 (2024).

12 Hills, F. R., Geoghegan, J. L. & Bostina, M. Architects of infection: A structural overview
of SARS-related coronavirus spike glycoproteins. *Virology* **604**, 110383 (2025).
13 Kleanthous, H., Silverman, J. M., Makar, K. W., Yoon, I. K., Jackson, N. & Vaughn, D.
W. Scientific rationale for developing potent RBD-based vaccines targeting COVID-19.
NPJ Vaccines **6**, 128 (2021).
14 Bruun, T. U. J., Andersson, A. C., Draper, S. J. & Howarth, M. Engineering a Rugged
Nanoscaffold To Enhance Plug-and-Display Vaccination. *ACS Nano* **12**, 8855–8866
(2018).
15 Starr, T. N., Greaney, A. J., Hilton, S. K., Ellis, D., Crawford, K. H. D., Dingens, A. S.,
Navarro, M. J., Bowen, J. E., Tortorici, M. A., Walls, A. C., King, N. P., Veesler, D. &
Bloom, J. D. Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain
Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295–1310.e1220 (2020).
16 Gote, V., Bolla, P. K., Kommineni, N., Butreddy, A., Nukala, P. K., Palakurthi, S. S. &
Khan, W. A Comprehensive Review of mRNA Vaccines. *Int J Mol Sci* **24** (2023).
17 Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R.,
McWilliam, H., Remmert, M., Soding, J., Thompson, J. D. & Higgins, D. G. Fast, scalable
generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol
Syst Biol* **7**, 539 (2011).
18 Hou, X., Zaks, T., Langer, R. & Dong, Y. Lipid nanoparticles for mRNA delivery. *Nat Rev
Mater*, 1–17 (2021).
19 Robbiani, D. F., Gaebler, C., Muecksch, F., Lorenzi, J. C. C., Wang, Z., Cho, A., Agudelo,
M., Barnes, C. O., Gazumyan, A., Finkin, S., Hagglof, T., Oliveira, T. Y., Viant, C.,
Hurley, A., Hoffmann, H. H., Millard, K. G., Kost, R. G., Cipolla, M., Gordon, K.,
Bianchini, F., Chen, S. T., Ramos, V., Patel, R., Dizon, J., Shimeliovich, I., Mendoza, P.,
Hartweger, H., Nogueira, L., Pack, M., Horowitz, J., Schmidt, F., Weisblum, Y.,
Michailidis, E., Ashbrook, A. W., Waltari, E., Pak, J. E., Huey-Tubman, K. E., Koranda,
N., Hoffman, P. R., West, A. P., Jr., Rice, C. M., Hatziioannou, T., Bjorkman, P. J.,
Bieniasz, P. D., Caskey, M. & Nussenzweig, M. C. Convergent antibody responses to
SARS-CoV-2 in convalescent individuals. *Nature* **584**, 437–442 (2020).
20 Jette, C. A., Cohen, A. A., Gnanapragasam, P. N. P., Muecksch, F., Lee, Y. E., Huey-
Tubman, K. E., Schmidt, F., Hatziioannou, T., Bieniasz, P. D., Nussenzweig, M. C., West,
A. P., Keeffe, J. R., Bjorkman, P. J. & Barnes, C. O. Broad cross-reactivity across
sarbecoviruses exhibited by a subset of COVID-19 donor-derived neutralizing antibodies.
Cell reports **36**, 109760 (2021).
21 Votteler, J. & Sundquist, W. I. Virus budding and the ESCRT pathway. *Cell Host Microbe*
14, 232–241 (2013).
22 Zhang, Y., Wang, C., Zheng, Y., Chen, F., Feng, Y., Fang, L., Wang, Z., Fu, Z. F., Zhou,
M. & Zhao, L. A chimeric Ad5-Envp-VLP vaccine platform confers broad-spectrum 1
immunity against emerging and re-emerging pathogens *bioRxiv* (2025).
23 Gonzalez-Magaldi, M., Gullapalli, A., Papoulas, O., Liu, C., Leung, A. Y., Guo, L., Brilot,
A. F., Marcotte, E. M., Ke, Z. & Leahy, D. J. Structure and organization of full-length
epidermal growth factor receptor in extracellular vesicles by cryo-electron tomography.
Proc Natl Acad Sci U S A **122**, e2424678122 (2025).
24 Olson, B. A., Huey-Tubman, K. E., Mao, Z., Hoffmann, M. A. G., Murray, R. M. & Mayo,
S. L. Recruiting ESCRT to single-chain heterotrimer peptide-MHCI releases antigen-
presenting vesicles that stimulate T cells selectively. (2025).
25 Fan, C., Keeffe, J. R., Malecek, K. E., Cohen, A. A., West, A. P., Jr., Baharani, V. A.,
Rorick, A. V., Gao, H., Gnanapragasam, P. N. P., Rho, S., Alvarez, J., Segovia, L. N.,
Hatziioannou, T., Bieniasz, P. D. & Bjorkman, P. J. Cross-reactive sarbecovirus antibodies

induced by mosaic RBD nanoparticles. *Proc Natl Acad Sci U S A* **122**, e2501637122
(2025).
26　Fan, C., Cohen, A. A., Park, M., Hung, A. F., Keeffe, J. R., Gnanapragasam, P. N. P., Lee,
Y. E., Gao, H., Kakutani, L. M., Wu, Z., Kleanthous, H., Malecek, K. E., Williams, J. C.
& Bjorkman, P. J. Neutralizing monoclonal antibodies elicited by mosaic RBD
nanoparticles bind conserved sarbecovirus epitopes. *Immunity* **55**, 2419–2435 e2410
(2022).
27　Goldblatt, D., Alter, G., Crotty, S. & Plotkin, S. A. Correlates of protection against SARS-
CoV-2 infection and COVID-19 disease. *Immunol Rev* **310**, 6–26 (2022).
28　Corbett, K. S., Nason, M. C., Flach, B., Gagne, M., O'Connell, S., Johnston, T. S., Shah,
S. N., Edara, V. V., Floyd, K., Lai, L., McDanal, C., Francica, J. R., Flynn, B., Wu, K.,
Choi, A., Koch, M., Abiona, O. M., Werner, A. P., Moliva, J. I., Andrew, S. F., Donaldson,
M. M., Fintzi, J., Flebbe, D. R., Lamb, E., Noe, A. T., Nurmukhambetova, S. T., Provost,
S. J., Cook, A., Dodson, A., Faudree, A., Greenhouse, J., Kar, S., Pessaint, L., Porto, M.,
Steingrebe, K., Valentin, D., Zouantcha, S., Bock, K. W., Minai, M., Nagata, B. M., van
de Wetering, R., Boyoglu-Barnum, S., Leung, K., Shi, W., Yang, E. S., Zhang, Y., Todd,
J. M., Wang, L., Alvarado, G. S., Andersen, H., Foulds, K. E., Edwards, D. K., Mascola,
J. R., Moore, I. N., Lewis, M. G., Carfi, A., Montefiori, D., Suthar, M. S., McDermott, A.,
Roederer, M., Sullivan, N. J., Douek, D. C., Graham, B. S. & Seder, R. A. Immune
correlates of protection by mRNA-1273 vaccine against SARS-CoV-2 in nonhuman
primates. *Science* **373**, eabj0299 (2021).
29　Gilbert, P. B., Montefiori, D. C., McDermott, A. B., Fong, Y., Benkeser, D., Deng, W.,
Zhou, H., Houchens, C. R., Martins, K., Jayashankar, L., Castellino, F., Flach, B., Lin, B.
C., O'Connell, S., McDanal, C., Eaton, A., Sarzotti-Kelsoe, M., Lu, Y., Yu, C., Borate, B.,
van der Laan, L. W. P., Hejazi, N. S., Huynh, C., Miller, J., El Sahly, H. M., Baden, L. R.,
Baron, M., De La Cruz, L., Gay, C., Kalams, S., Kelley, C. F., Andrasik, M. P., Kublin, J.
G., Corey, L., Neuzil, K. M., Carpp, L. N., Pajon, R., Follmann, D., Donis, R. O., Koup,
R. A., Immune Assays Team section, s., Moderna, I. T. s. s., Coronavirus Vaccine
Prevention Network /Coronavirus Efficacy Team section, s. & United States Government
/Co, V. P. N. B. T. s. s. Immune correlates analysis of the mRNA-1273 COVID-19 vaccine
efficacy clinical trial. *Science* **375**, 43–50 (2022).
30　Schmidt, F., Weisblum, Y., Muecksch, F., Hoffmann, H.-H., Michailidis, E., Lorenzi, J. C.
C., Mendoza, P., Rutkowska, M., Bednarski, E., Gaebler, C., Agudelo, M., Cho, A., Wang,
Z., Gazumyan, A., Cipolla, M., Caskey, M., Robbiani, D. F., Nussenzweig, M. C., Rice,
C. M., Hatziioannou, T. & Bieniasz, P. D. Measuring SARS-CoV-2 neutralizing antibody
activity using pseudotyped and chimeric viruses. *Journal of Experimental Medicine* **217**
(2020).
31　Corbett, K. S., Edwards, D. K., Leist, S. R., Abiona, O. M., Boyoglu-Barnum, S., Gillespie,
R. A., Himansu, S., Schafer, A., Ziwawo, C. T., DiPiazza, A. T., Dinnon, K. H., Elbashir,
S. M., Shaw, C. A., Woods, A., Fritch, E. J., Martinez, D. R., Bock, K. W., Minai, M.,
Nagata, B. M., Hutchinson, G. B., Wu, K., Henry, C., Bahl, K., Garcia-Dominguez, D.,
Ma, L., Renzi, I., Kong, W. P., Schmidt, S. D., Wang, L., Zhang, Y., Phung, E., Chang, L.
A., Loomis, R. J., Altaras, N. E., Narayanan, E., Metkar, M., Presnyak, V., Liu, C., Louder,
M. K., Shi, W., Leung, K., Yang, E. S., West, A., Gully, K. L., Stevens, L. J., Wang, N.,
Wrapp, D., Doria-Rose, N. A., Stewart-Jones, G., Bennett, H., Alvarado, G. S., Nason, M.
C., Ruckwardt, T. J., McLellan, J. S., Denison, M. R., Chappell, J. D., Moore, I. N.,
Morabito, K. M., Mascola, J. R., Baric, R. S., Carfi, A. & Graham, B. S. SARS-CoV-2
mRNA vaccine design enabled by prototype pathogen preparedness. *Nature* **586**, 567–571
(2020).

32 Paciello, I., Maccari, G., Pantano, E., Andreano, E. & Rappuoli, R. High-resolution map
of the Fc functions mediated by COVID-19-neutralizing antibodies. *Proc Natl Acad Sci U*
S A **121**, e2314730121 (2024).
33 Abu-Humaidan, A. H. A., Ahmad, F. M., Awajan, D., Jarrar, R. F. & Alaridah, N. Anti-
SARS-Cov-2 S-RBD IgG Formed after BNT162b2 Vaccination Can Bind C1q and
Activate Complement. *J Immunol Res* **2022**, 7263740 (2022).
34 Clark, J. J., Hoxie, I., Adelsberg, D. C., Sapse, I. A., Andreata-Santos, R., Yong, J. S.,
Amanat, F., Tcheou, J., Raskin, A., Singh, G., Gonzalez-Dominguez, I., Edgar, J. E.,
Bournazos, S., Sun, W., Carreno, J. M., Simon, V., Ellebedy, A. H., Bajic, G. & Krammer,
F. Protective effect and molecular mechanisms of human non-neutralizing cross-reactive
spike antibodies elicited by SARS-CoV-2 mRNA vaccination. *Cell reports* **43**, 114922
(2024).
35 Ranieri, E., Popescu, I. & Gigante, M. CTL ELISPOT assay. *Methods Mol Biol* **1186**, 75–
86 (2014).
36 Leehan, K. M. & Koelsch, K. A. T Cell ELISPOT: For the Identification of Specific
Cytokine-Secreting T Cells. *Methods Mol Biol* **1312**, 427–434 (2015).
37 Rock, K. L., Reits, E. & Neefjes, J. Present Yourself! By MHC Class I and MHC Class II
Molecules. *Trends Immunol* **37**, 724–737 (2016).
38 Greaney, A. J., Loes, A. N., Crawford, K. H. D., Starr, T. N., Malone, K. D., Chu, H. Y. &
Bloom, J. D. Comprehensive mapping of mutations in the SARS-CoV-2 receptor-binding
domain that affect recognition by polyclonal human plasma antibodies. *Cell Host Microbe*
29, 463–476 e466 (2021).
39 Greaney, A. J., Loes, A. N., Gentles, L. E., Crawford, K. H. D., Starr, T. N., Malone, K.
D., Chu, H. Y. & Bloom, J. D. Antibodies elicited by mRNA-1273 vaccination bind more
broadly to the receptor binding domain than do those from SARS-CoV-2 infection. *Science*
translational medicine **13** (2021).
40 Greaney, A. J., Starr, T. N., Barnes, C. O., Weisblum, Y., Schmidt, F., Caskey, M., Gaebler,
C., Cho, A., Agudelo, M., Finkin, S., Wang, Z., Poston, D., Muecksch, F., Hatziioannou,
T., Bieniasz, P. D., Robbiani, D. F., Nussenzweig, M. C., Bjorkman, P. J. & Bloom, J. D.
Mapping mutations to the SARS-CoV-2 RBD that escape binding by different classes of
antibodies. *Nat Commun* **12**, 4196 (2021).
41 Greaney, A. J., Starr, T. N., Gilchuk, P., Zost, S. J., Binshtein, E., Loes, A. N., Hilton, S.
K., Huddleston, J., Eguia, R., Crawford, K. H. D., Dingens, A. S., Nargi, R. S., Sutton, R.
E., Suryadevara, N., Rothlauf, P. W., Liu, Z., Whelan, S. P. J., Carnahan, R. H., Crowe, J.
E., Jr. & Bloom, J. D. Complete Mapping of Mutations to the SARS-CoV-2 Spike
Receptor-Binding Domain that Escape Antibody Recognition. *Cell Host Microbe* **29**, 44–
57 e49 (2021).
42 Starr, T. N., Czudnochowski, N., Liu, Z., Zatta, F., Park, Y. J., Addetia, A., Pinto, D.,
Beltramello, M., Hernandez, P., Greaney, A. J., Marzi, R., Glass, W. G., Zhang, I.,
Dingens, A. S., Bowen, J. E., Tortorici, M. A., Walls, A. C., Wojcechowskyj, J. A., De
Marco, A., Rosen, L. E., Zhou, J., Montiel-Ruiz, M., Kaiser, H., Dillen, J. R., Tucker, H.,
Bassi, J., Silacci-Fregni, C., Housley, M. P., di Iulio, J., Lombardo, G., Agostini, M.,
Sprugasci, N., Culap, K., Jaconi, S., Meury, M., Dellota, E., Jr., Abdelnabi, R., Foo, S. C.,
Cameroni, E., Stumpf, S., Croll, T. I., Nix, J. C., Havenar-Daughton, C., Piccoli, L.,
Benigni, F., Neyts, J., Telenti, A., Lempp, F. A., Pizzuto, M. S., Chodera, J. D., Hebner,
C. M., Virgin, H. W., Whelan, S. P. J., Veesler, D., Corti, D., Bloom, J. D. & Snell, G.
SARS-CoV-2 RBD antibodies that maximize breadth and resistance to escape. *Nature* **597**,
97–102 (2021).

43 Starr, T. N., Greaney, A. J., Addetia, A., Hannon, W. W., Choudhary, M. C., Dingens, A.
S., Li, J. Z. & Bloom, J. D. Prospective mapping of viral mutations that escape antibodies
used to treat COVID-19. *Science* **371**, 850–854 (2021).
44 Barnes, C. O., Jette, C. A., Abernathy, M. E., Dam, K.-M. A., Esswein, S. R., Gristick, H.
B., Malyutin, A. G., Sharaf, N. G., Huey-Tubman, K. E., Lee, Y. E., Robbiani, D. F.,
Nussenzweig, M. C., West, A. P. & Bjorkman, P. J. SARS-CoV-2 neutralizing antibody
structures inform therapeutic strategies. *Nature* **588**, 682–687 (2020).
45 Jensen, J. L., Sankhala, R. S., Dussupt, V., Bai, H., Hajduczki, A., Lal, K. G., Chang, W.
C., Martinez, E. J., Peterson, C. E., Golub, E. S., Rees, P. A., Mendez-Rivera, L., Zemil,
M., Kavusak, E., Mayer, S. V., Wieczorek, L., Kannan, S., Doranz, B. J., Davidson, E.,
Yang, E. S., Zhang, Y., Chen, M., Choe, M., Wang, L., Gromowski, G. D., Koup, R. A.,
Michael, N. L., Polonis, V. R., Rolland, M., Modjarrad, K., Krebs, S. J. & Joyce, M. G.
Targeting the Spike Receptor Binding Domain Class V Cryptic Epitope by an Antibody
with Pan-Sarbecovirus Activity. *J Virol* **97**, e0159622 (2023).
46 Cui, L., Li, T., Lan, M., Zhou, M., Xue, W., Zhang, S., Wang, H., Hong, M., Zhang, Y.,
Yuan, L., Sun, H., Ye, J., Zheng, Q., Guan, Y., Gu, Y., Xia, N. & Li, S. A cryptic site in
class 5 epitope of SARS-CoV-2 RBD maintains highly conservation across natural isolates.
iScience **27**, 110208 (2024).
47 Bangaru, S., Jackson, A. M., Copps, J., Fernandez-Quintero, M. L., Torres, J. L., Richey,
S. T., Nogal, B., Sewall, L. M., Torrents de la Pena, A., Rehman, A., Guebre-Xabier, M.,
Girard, B., Das, R., Corbett-Helaire, K. S., Seder, R. A., Graham, B. S., Edwards, D. K.,
Patel, N., Smith, G. & Ward, A. B. Structural serology of polyclonal antibody responses to
mRNA-1273 and NVX-CoV2373 COVID-19 vaccines. *Cell reports* **44**, 115986 (2025).
48 Landau, M., Mayrose, I., Rosenberg, Y., Glaser, F., Martz, E., Pupko, T. & Ben-Tal, N.
ConSurf 2005: the projection of evolutionary conservation scores of residues on protein
structures. *Nucleic Acids Res* **33**, W299–302 (2005).
49 Crescioli, S., Jatiani, S. & Moise, L. With great power, comes great responsibility: the
importance of broadly measuring Fc-mediated effector function early in the antibody
development process. *MAbs* **17**, 2453515 (2025).
50 Bruhns, P. & Jonsson, F. Mouse and human FcR effector functions. *Immunol Rev* **268**, 25–
51 (2015).
51 Hwang, H. S., Lee, Y. T., Kim, K. H., Seo, H. S., Yang, K. S., Cho, H. & Kang, S. M.
Roles of the Fc Receptor gamma-Chain in Inducing Protective Immune Responses after
Heterologous Vaccination against Respiratory Syncytial Virus Infection. *Vaccines (Basel)*
9 (2021).
52 Aleebrahim-Dehkordi, E., Molavi, B., Mokhtari, M., Deravi, N., Fathi, M., Fazel, T.,
Mohebalizadeh, M., Koochaki, P., Shobeiri, P. & Hasanpour-Dehkordi, A. T helper type
(Th1/Th2) responses to SARS-CoV-2 and influenza A (H1N1) virus: From cytokines
produced to immune responses. *Transplant Immunology* **70** (2022).
53 Han, Q., Gao, X., Chu, Z., Wang, X., Eisa Addoma Adam, F., Zhang, S., Jia, Y., Qiu, X.,
Wang, X. & Yang, Z. Truncated chicken MDA5 enhances the immune response to
inactivated NDV vaccine. *Vet Immunol Immunopathol* **208**, 44–52 (2019).
54 Turner, H. L., Jackson, A. M., Richey, S. T., Sewall, L. M., Antanasijevic, A., Hangartner,
L. & Ward, A. B. Protocol for analyzing antibody responses to glycoprotein antigens using
electron-microscopy-based polyclonal epitope mapping. *STAR Protoc* **4**, 102476 (2023).
55 McInnes, L., Healy, J. & Melville, J. *UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction, v3*, <https://arxiv.org/abs/1802.03426v3> (2020).
56 Wong, B. K. & Mabbott, N. A. Systematic review and meta-analysis of COVID-19 mRNA
vaccine effectiveness against hospitalizations in adults. *Immunother Adv* **4**, ltae011 (2024).

57 Stewart-Jones, G. B. E., Elbashir, S. M., Wu, K., Lee, D., Renzi, I., Ying, B., Koch, M.,
Sein, C. E., Choi, A., Whitener, B., Garcia-Dominguez, D., Henry, C., Woods, A., Ma, L.,
Montes Berrueta, D., Avena, L. E., Quinones, J., Falcone, S., Hsiao, C. J., Scheaffer, S.
M., Thackray, L. B., White, P., Diamond, M. S., Edwards, D. K. & Carfi, A. Domain-based
mRNA vaccines encoding spike protein N-terminal and receptor binding domains confer
protection against SARS-CoV-2. *Science translational medicine* **15**, eadf4100 (2023).
58 Chalkias, S., Dennis, P., Petersen, D., Radhakrishnan, K., Vaughan, L., Handforth, R.,
Rossi, A., Wahid, R., Edwards, D. K., Feng, J., Deng, W., Zhou, H., De Windt, E.,
Urdaneta, V., Paila, Y., Girard, B., Faust, S. N., Walsh, S. R., Cosgrove, C. A., Miller, J.
& Das, R. Efficacy, immunogenicity, and safety of a next-generation mRNA-1283
COVID-19 vaccine compared with the mRNA-1273 vaccine (NextCOVE): results from a
phase 3, randomised, observer-blind, active-controlled trial. *The Lancet Infectious*
Diseases (2025).
59 Pardi, N., Hogan, M. J., Porter, F. W. & Weissman, D. mRNA vaccines - a new era in
vaccinology. *Nat Rev Drug Discov* **17**, 261–279 (2018).
60 Mahrokhian, S. H., Tostanoski, L. H., Vidal, S. J. & Barouch, D. H. COVID-19 vaccines:
Immune correlates and clinical outcomes. *Hum Vaccin Immunother* **20**, 2324549 (2024).
61 Foged, C., Brodin, B., Frokjaer, S. & Sundblad, A. Particle size and surface charge affect
particle uptake by human dendritic cells in an in vitro model. *Int J Pharm* **298**, 315–322
(2005).
62 Joshi, V. B., Geary, S. M. & Salem, A. K. Biodegradable particles as vaccine delivery
systems: size matters. *AAPS J* **15**, 85–94 (2013).
63 Zepeda-Cervantes, J., Ramirez-Jarquin, J. O. & Vaca, L. Interaction Between Virus-Like
Particles (VLPs) and Pattern Recognition Receptors (PRRs) From Dendritic Cells (DCs):
Toward Better Engineering of VLPs. *Front Immunol* **11**, 1100 (2020).
64 Kim, C., Kim, J. D. & Seo, S. U. Nanoparticle and virus-like particle vaccine approaches
against SARS-CoV-2. *J Microbiol* **60**, 335–346 (2022).
65 Morris, G. E. in *The Protein Protocols Handbook Springer Protocols Handbooks* Ch.
Chapter 96, 595–600 (1996).
66 Fan, C., Cohen, A. A., Dam, K. A., Rorick, A. V., Priso Fils, A. I., Yang, Z.,
Gnanapragasam, P. N. P., Segovia, L. N., Huey-Tubman, K. E., Moon, W. J., Lin, P. J. C.,
Bjorkman, P. J. & Hoffmann, M. A. G. Bivalent mRNA booster encoding virus-like
particles elicits potent polyclass RBD antibodies in pre-vaccinated mice. *bioRxiv* (2025).
67 Chai, P., Shi, Y., Li, X., Yang, M., Liu, X., Liu, M., Yu, J., Yin, X., Li, D., Li, K., Kong,
X., Zhang, Q., Wang, H., Sun, X., Li, J., Li, L., Li, D., Pang, L., Lu, X. & Duan, Z.
Improved mRNA-based RSV vaccine with PreF forming enveloped virus-like particles.
NPJ Vaccines **10**, 152 (2025).
68 McBride, C. E., Li, J. & Machamer, C. E. The cytoplasmic tail of the severe acute
respiratory syndrome coronavirus spike protein contains a novel endoplasmic reticulum
retrieval signal that binds COPI and promotes interaction with membrane protein. *J Virol*
81, 2418–2428 (2007).
69 Walker, S. E. & Lorsch, J. RNA purification--precipitation methods. *Methods Enzymol*
530, 337–343 (2013).
70 Baiersdorfer, M., Boros, G., Muramatsu, H., Mahiny, A., Vlatkovic, I., Sahin, U. & Kariko,
K. A Facile Method for the Removal of dsRNA Contaminant from In Vitro-Transcribed
mRNA. *Mol Ther Nucleic Acids* **15**, 26–35 (2019).
71 Pardi, N., Tuyishime, S., Muramatsu, H., Kariko, K., Mui, B. L., Tam, Y. K., Madden, T.
D., Hope, M. J. & Weissman, D. Expression kinetics of nucleoside-modified mRNA

delivered in lipid nanoparticles to mice by various routes. *J Control Release* **217**, 345–351
(2015).

72 Hsieh, C. L., Goldsmith, J. A., Schaub, J. M., DiVenere, A. M., Kuo, H. C., Javanmardi,
K., Le, K. C., Wrapp, D., Lee, A. G., Liu, Y., Chou, C. W., Byrne, P. O., Hjorth, C. K.,
Johnson, N. V., Ludes-Meyers, J., Nguyen, A. W., Park, J., Wang, N., Amengor, D.,
Lavinder, J. J., Ippolito, G. C., Maynard, J. A., Finkelstein, I. J. & McLellan, J. S. Structure-
based design of prefusion-stabilized SARS-CoV-2 spikes. *Science* **369**, 1501–1505 (2020).

73 Wang, Z., Muecksch, F., Cho, A., Gaebler, C., Hoffmann, H. H., Ramos, V., Zong, S.,
Cipolla, M., Johnson, B., Schmidt, F., DaSilva, J., Bednarski, E., Ben Tanfous, T., Raspe,
R., Yao, K., Lee, Y. E., Chen, T., Turroja, M., Milard, K. G., Dizon, J., Kaczynska, A.,
Gazumyan, A., Oliveira, T. Y., Rice, C. M., Caskey, M., Bieniasz, P. D., Hatziioannou, T.,
Barnes, C. O. & Nussenzweig, M. C. Analysis of memory B cells identifies conserved
neutralizing epitopes on the N-terminal domain of variant SARS-Cov-2 spike proteins.
Immunity **55**, 998–1012 e1018 (2022).

74 Tykvart, J., Sacha, P., Barinka, C., Knedlik, T., Starkova, J., Lubkowski, J. & Konvalinka,
J. Efficient and versatile one-step affinity purification of in vivo biotinylated proteins:
expression, characterization and structure analysis of recombinant human glutamate
carboxypeptidase II. *Protein Expr Purif* **82**, 106–115 (2012).

75 Rahikainen, R., Rijal, P., Tan, T. K., Wu, H. J., Andersson, A. C., Barrett, J. R., Bowden,
T. A., Draper, S. J., Townsend, A. R. & Howarth, M. Overcoming Symmetry Mismatch in
Vaccine Nanoassembly through Spontaneous Amidation. *Angew Chem Int Ed Engl* **60**,
321–330 (2021).

76 Crawford, K. H. D., Eguia, R., Dingens, A. S., Loes, A. N., Malone, K. D., Wolf, C. R.,
Chu, H. Y., Tortorici, M. A., Veesler, D., Murphy, M., Pettie, D., King, N. P., Balazs, A.
B. & Bloom, J. D. Protocol and Reagents for Pseudotyping Lentiviral Particles with SARS-
CoV-2 Spike Protein for Neutralization Assays. *Viruses* **12** (2020).

77 Seifert, S. N., Bai, S., Fawcett, S., Norton, E. B., Zwezdaryk, K. J., Robinson, J., Gunn, B.
& Letko, M. An ACE2-dependent Sarbecovirus in Russian bats is resistant to SARS-CoV-
2 vaccines. *PLoS Pathog* **18**, e1010828 (2022).

78 West, A. P., Jr., Scharf, L., Horwitz, J., Klein, F., Nussenzweig, M. C. & Bjorkman, P. J.
Computational analysis of anti-HIV-1 antibody neutralization panel data to identify
potential functional epitope residues. *Proc Natl Acad Sci U S A* **110**, 10598–10603 (2013).

79 Starr, T. N., Greaney, A. J., Hannon, W. W., Loes, A. N., Hauser, K., Dillen, J. R., Ferri,
E., Farrell, A. G., Dadonaite, B., McCallum, M., Matreyek, K. A., Corti, D., Veesler, D.,
Snell, G. & Bloom, J. D. Shifting mutational constraints in the SARS-CoV-2 receptor-
binding domain during viral evolution. *Science* **377**, 420–424 (2022).

80 Lee, J., Zepeda, S. K., Park, Y. J., Taylor, A. L., Quispe, J., Stewart, C., Leaf, E. M.,
Treichel, C., Corti, D., King, N. P., Starr, T. N. & Veesler, D. Broad receptor tropism and
immunogenicity of a clade 3 sarbecovirus. *Cell Host Microbe* **31**, 1961–1973 e1911
(2023).

81 Greaney, A. J., Starr, T. N., Eguia, R. T., Loes, A. N., Khan, K., Karim, F., Cele, S., Bowen,
J. E., Logue, J. K., Corti, D., Veesler, D., Chu, H. Y., Sigal, A. & Bloom, J. D. A SARS-
CoV-2 variant elicits an antibody response with a shifted immunodominance hierarchy.
PLoS Pathog **18**, e1010248 (2022).

82 Brown, E. P., Dowell, K. G., Boesch, A. W., Normandin, E., Mahan, A. E., Chu, T.,
Barouch, D. H., Bailey-Kellogg, C., Alter, G. & Ackerman, M. E. Multiplexed Fc array for
evaluation of antigen-specific antibody effector profiles. *J Immunol Methods* **443**, 33–44
(2017).

83   Cohen, A. A., Yang, Z., Gnanapragasam, P. N. P., Ou, S., Dam, K. A., Wang, H. &
Bjorkman, P. J. Construction, characterization, and immunization of nanoparticles that
display a diverse array of influenza HA trimers. *PLoS One* **16**, e0247963 (2021).
84   Lehle, L. & Bause, E. Primary structural requirements for N- and O-glycosylation of yeast
mannoproteins. *Biochimica et Biophysica Acta (BBA) - General Subjects* **799**, 246–251
(1984).
1426
1427

1428   Supplementary Information for

1429

1430

1431

<div align="center">Genetic encoding of mosaic-8 pan-sarbecovirus RBD vaccine for mRNA delivery</div>

1432   Alexander A. Cohen[1], Jennifer R. Keeffe[1], Lusineh Manasyan[1], Indeever Madireddy[1],

1433   Ange-Célia I. Priso Fils[1], Kim-Marie A. Dam[2], Haley E. Stober[2], Rory A. Hills[,3,4,7], Woohyun J.

1434   Moon[5], Paulo J.C. Lin[5], Mark R. Howarth[3], Magnus A.G. Hoffmann[2,6,8*], Pamela J. Bjorkman[1,8*]

1435

1436
Extended Data Figure 1. mRNA-encoded dual quartets are presented on cells and eVLPs.
(a) Flow cytometry using C118, a human pan-sarbecovirus RBD-specific mAb[19,20], demonstrating
that transfections of mRNAs encoding EABR and non-EABR versions of RBD quartet 4a or
quartet 4b resulted in expression on the surface of transfected cells. Untransfected cells are shown
as controls (unstained or stained with C118). (b) Left: ELISA schematic. RBD quartet levels were
measured by capturing purified eVLPs with an anti-Rs4081 RBD mAb[25] (recognizes RBD quartet
4a but not quartet 4b) or an anti-RmYN02 RBD mAb[25] (recognizes RBD quartet 4b but not quartet
4a) and detecting with the pan-sarbecovirus RBD-specific C118 mAb[19,20]. Right: ELISA showing
that purified single quartet eVLPs purified from supernatants of cells transfected with mRNA
encoding RBD quartet 4a-EABR or RBD quartet 4b-EABR displayed RBD quartets on their
surface. The mean absorbance of two replicates is shown as a function of eVLP dilution with error
bars representing standard deviations. (c) Flow cytometry demonstrating that RBD quartets 4a and
4b were co-expressed on the surfaces of individual cells. RBD quartet 4a was detected using an
anti-Rs4081 RBD mAb[25] and RBD quartet 4b was detected using anti-WIV1 RBD mAb[26].
Untransfected cells stained with the anti-Rs4081 and anti-WIV1 RBD mAbs are shown as controls.
The percentage of cells in each quadrant is shown. (d) Left: ELISA schematic. Purified eVLPs
were evaluated in a sandwich ELISA in which an anti-WIV1 RBD mAb[26] (recognizes RBD quartet
4b but not quartet 4a) was used as a capture Ab and an anti-Rs4081 RBD mAb[25] (recognizes

quartet 4a but not quartet 4b) was used for detection. Right: ELISA showing detection of both
RBD quartets 4a and 4b on the surfaces of eVLPs purified from supernatants of cells co-transfected
with mRNAs encoding RBD quartet 4a-EABR and 4b-EABR constructs. The mean absorbance of
two replicates is shown as a function of eVLP dilution with error bars representing standard
deviations. Binding was not detected for purified supernatant samples from untransfected cells
(control) or from supernatants from cells transfected with RBD quartet 4a, RBD quartet 4a-EABR,
RBD quartet 4b, RBD quartet 4b-EABR, or RBD quartet 4a plus RBD quartet 4b (all data were
plotted but some data points near 0.0 A_{450} are obscured by others).
1463

1464

1465

1466

Extended Data Figure 2. Dual quartet mRNAs elicited cross-reactive Abs.

Data are shown for ELISA and neutralization analyses for terminal bleed serum samples (see also
Fig. 2). (a) Immunization regimen. Left: Mice were primed at day 0, boosted at day 28, and samples
were collected from a terminal bleed at day 77 or 78. Right: Colors used to identify immunizations
and symbols indicating a matched (filled in square data points; gray shading around name) or
mismatched (unfilled square data points; black outline around name) sarbecovirus antigen.
Sarbecovirus strain names are colored in panel b according to clade. (b) RBD-binding ELISA (left)
and pseudovirus neutralization (right) results for serum samples from day 77 or 78 after the prime
immunization. Immunogens are shown compared with the cohorts immunized with 0.5 μg of an
mRNA-based immunogen. Dashed horizontal lines indicate the limits of detection for each assay.
Left: Geomeans of ED_{50} values for animals in each cohort (symbols with geometric standard
deviations indicated by error bars) are connected by colored lines. Mean titers against RBDs from
indicated sarbecoviruses were compared pairwise across immunization cohorts by Tukey's
multiple comparison test with the Geisser-Greenhouse correction (as calculated by GraphPad
Prism). Right: Neutralization potencies for serum samples from day 77 or 78 after immunization
presented as half-maximal inhibitory dilutions (ID_{50} values) of sera against pseudoviruses from
the indicated coronavirus strains. Significant differences between the two cohorts are indicated by
asterisks: $p<0.05$ = *, $p<0.01$ = **, $p<0.001$ = ***, $p<0.0001$ = ****.

1485
1486
Extended Data Figure 3. DMS line and logo plots (WA1 RBD library) for individual mice.
DMS line plots (left) and logo plots (right) for results from individual mice (identified by 4-digit
numbers) immunized with the indicated immunogens. X-axes of line and logo plots show RBD
residue numbers, and y-axes of line plots show the sum of the Ab escape for all substitutions at an
RBD residue (larger numbers indicate increased Ab escape). Sites with the strongest Ab escape
are shown in logo plots; tall letters represent the most frequent substitutions at a site. Logo plot
residues are colored according to RBD epitopes within different classes[20,44-46] as indicated on the
legend.

1495
1496
Extended Data Figure 4. DMS line and logo plots (SARS-1 RBD library) for individual mice.
DMS line plots (left) and logo plots (right) for results from individual mice (identified by 4-digit
numbers) immunized with the indicated immunogens. X-axes of line and logo plots show RBD
residue numbers, and y-axes of line plots show the sum of the Ab escape for all substitutions at an
RBD residue (larger numbers indicate increased Ab escape). Sites with the strongest Ab escape
are shown in logo plots; tall letters represent the most frequent substitutions at a site. Logo plot
residues are colored according to RBD epitopes within different classes[20,44-46] as indicated on the
legend.

1506
Extended Data Figure 5. mRNA-encoded dual quartet RBD immunogens elicit balanced IgG subclass and potent FcγR-binding responses.

MFI = Median fluorescent intensity. For IgG1, IgG2a, IgG2b, IgG3, FcγR2b-binding IgGs, FcγR3-binding IgGs, FcγR4-binding IgGs, and total IgG, geomean MFI values of the individual responses shown in panel a for each cohort binding to different spikes or RBDs are represented as points in a box and whisker plot and compared pairwise across immunization cohorts by Tukey's multiple comparison test calculated by GraphPad Prism. Significant differences between cohorts linked by vertical lines in panels b and c are indicated by asterisks: $p<0.05$ = *, $p<0.01$ = **, $p<0.001$ = ***, $p<0.0001$ = ****.

See also Fig. 5.

1518

1519   **Extended Data Figure 6. Schematic of SySPEM approach.**

Extended Data Figure 7. Characterization of RBD N-glycan mutants used for SySPEM analysis.

(a) Top: Locations of the added N-glycan(s) in RBD KO mutants, as shown by highlighting of one or more residues that were changed to Asn in the introduced PNGS(s). Bottom: Sequence conservation of 16 sarbecovirus RBDs calculated using ConSurf[48] shown on a surface representation of SARS-2 RBD (PDB 7BZ5). Class 1, 2, 3, 4, 1/4, and 5 anti-RBD Ab epitopes[20,44-46] are outlined in dots in different colors using information from representative structures of Abs bound to SARS-2 spike or RBD (C102: PDB 7K8M; C002: PDB 7K8T, S309: PDB 7JX3; CR3022: PDB 7LOP; C118: PDB 7RKV; WRAIR-2063: PDB 8EOO). (b) SDS-PAGE analysis of purified wt RBD and RBD KO mutants. Molecular weight marker positions are shown on the left with the molecular weight indicated in kilodaltons. KO mutants are listed with the RBD epitope class affected by the N-glycan addition(s) and residue number(s) of Asn residue(s) to which N-linked glycan(s) were added. (c) Results of ELISA showing ratio of binding of characterized mAbs or human ACE2-Fc[20] to wt RBD versus the RBD KO mutants listed on the left. ELISA EC_{50} values for binding of each reagent to wt RBD and the six RBD KO mutants were derived using , and the

1537   fold reduction in binding to each RBD KO mutant was calculated as EC_{50} RBD KO / EC_{50} RBD

1538   wt. Classifications of RBD epitopes recognized by the characterized mAbs and ACE-2 Fc are

1539   taken from refs[20,25,26,44-46].

1540

1541

Extended Data Figure 8. SySPEM score comparisons show distinct epitope profiles across immunogen cohorts.

SySPEM scores from individual mice in each immunogen cohort (columns) were determined for each IgG class (rows, IgG subclass plus different FcγR-binding IgGs) with statistical comparisons

between targeted epitopes (colors). A SySPEM value of 0 indicates that none of the IgGs in that sample were affected by the glycan addition and therefore the sample did not contain IgGs that recognize that epitope, and a SySPEM value of 100 indicates that all IgGs in that sample recognized that epitope (Extended Data Fig. 6). A SySPEM value between 0 and 100 indicates the proportion of IgGs in a sample that recognized the epitope that was blocked by glycan addition in the RBD KO mutant. Box and whisker plots of SySPEM scores with individual data points representing one mouse are shown. Significant differences between cohorts were calculated using Tukey's HSD posthoc test and linked by vertical lines indicated by asterisks: $p<0.05$ = *, $p<0.01$ = **, $p<0.001$ = ***, $p<0.0001$ = ****. See also Figs. 6, 7.