1

Anthropogenic nitrogen deposition decouples relationships with decomposing microbes, altering

SOM molecular composition, but not molecular complexity or diversity

Brooke E. Propson[1,2], William A. Argiroff[3], Grace A. Cagle[1], Rima A. Upchurch[3], Donald R.

Zak[3,4], A. Stuart Grandy[5], and Zachary B. Freedman[1*]

[1]Department of Soil and Environmental Sciences

University of Wisconsin, Madison, WI, USA 53706

[2]University of Michigan Biological Station

9133 Biological Rd, Pellston, MI, USA 49769

[3]School for Environment and Sustainability

[4]Ecology and Evolutionary Biology

University of Michigan, Ann Arbor, MI, USA 48109

[5]Department of Natural Resources and the Environment

University of New Hampshire, Durham, NH, USA 03824

*Corresponding author:

Zachary B. Freedman, zfreedman@wisc.edu (ORCID: 0000-0001-9160-7470)

Abstract

Soil organic matter (SOM) consists of different biochemical constituents that exist across

a spectrum of chemical complexity from simple to complex, and the relative abundance of these

potential substrates for microbial metabolism can have consequences for the fate of soil C, and

consequently the pace of climate change. However, current understanding of the mechanistic

controls on SOM biochemistry and decay, and how these may be affected by climate change, is

still incomplete. As such, the primary objectives of this study are to assess (1) the trajectory of

organic matter decomposition and stabilization from undecayed plant root litter to mineral SOM

and (2) how these processes are altered by anthropogenic nitrogen (N) deposition by using high-

resolution SOM biochemical and microbiome datasets and a long-term field experiment in the

Upper Great Lakes region of North America. N deposition did not significantly alter the

biochemical composition of SOM in the undecomposed and 1-year decomposed samples but did

alter mineral SOM biochemistry. This shift was attributed to changes in the relative abundance of

specific biochemical compounds rather than changes in more broadly defined compound classes,

such as lipids, for example. A reduced association between SOM biochemistry and microbial

decomposer communities under N deposition was also observed. Taken together, atmospheric N

deposition altered the composition of mineral SOM and decoupled the relationship between

microbial composition and SOM biochemistry, likely contributing to the previously observed

reduction in the decomposition of SOM while not significantly altering SOM's overall

biochemistry.

Keywords: SOM, biochemistry, bacteria, fungi, py-GCMS, decomposition

1. Introduction

Soil organic matter (SOM) constitutes the largest terrestrial organic carbon reservoir, and changes in SOM decomposition dynamics under fluctuating environmental conditions could intensify or mitigate impacts of climate change. Inputs from plants, through the deposition of above- and below-ground plant residues and the release of exudates from plant roots represents the largest organic matter input to soils[1,2]. Litter chemical composition, which is often estimated in terms of its nitrogen (N) or lignin contents[3,4], is regarded as one of the primary controls over decomposition rates and SOM accumulation and the fate of these inputs is governed by soil microbial activity[5–9]. In recent decades, there has been a shift from traditional theory of SOM chemical recalcitrance, where biochemically recalcitrant lignified material was thought to constitute the majority of SOM because of its theorized resistance to microbial decomposition[10,11], towards a continuum model of SOM complexity that proposes microbial-derived products as the primary constituents of SOM and the potential importance of molecular diversity in enabling SOM persistence[12–15].

Biochemical compounds that comprise litter exist across a spectrum of chemical complexity from simple to complex substrates and the relative abundance of these potential substrates for microbial metabolism can have consequences for the fate of C stored as mineral SOM[7,14,16]. When plant litter undergoes microbially mediated breakdown during decomposition, the resulting products will differ depending on the initial biochemical quality of the plant litter, as well as the composition of the microbial community decomposing that material[7,17]. The potential importance of biochemical complexity and diversity for SOM persistence lies in the different microbial requirements for metabolizing different biochemical compounds, both for compounds that are structurally dissimilar, as well as ones that are largely similar but may exhibit minor

differences, such as differences in the position of functional groups (for example, *ortho-* versus

para- benzoic acid)[15,18]. This rise in the acknowledgement and potential importance of a higher

resolution understanding of SOM is a deviation from the standard practice in which compounds

are typically grouped together as classes (e.g., lipids, etc.) that are assumed to behave similarly

through the decomposition process. This results in individual compounds being grouped into

classes that are inherently less refined. However, with the conceptual move away from stable SOM

largely consisting of recalcitrant material towards a spectrum of which biochemical complexity

and diversity enable SOM persistence, research has shown that certain SOM biochemical

constituents are resistant to decomposition despite being in a compound class that was traditionally

viewed as "labile" (ex: long-chain lipids)[19]. As such, there is increasing emphasis on the potential

importance of SOM molecular complexity and diversity in driving SOM persistence[12,15,20–22].

Critically, there is a need to determine how this proposed relationship between SOM molecular

diversity and persistence responds to changing climatic conditions.

To better understand how anthropogenic N deposition, a pervasive driver of global

environmental change, affects the cycling and storage of SOM, we conducted a 37-year field

experiment to simulate future rates of anthropogenic N deposition. After 20 years of experimental

N deposition inputs, we found that anthropogenic N deposition slowed the rate and extent of

organic matter decomposition, resulting in the accumulation of lignified material in mineral soil

derived primarily from fine root litter[23–25]. This change in SOM biochemistry in response to N

deposition was associated with decreases in lignolytic fungi and positively associated with

saprotrophic bacteria[23,26–28]. However, lignin, and its identifiable derivatives, often comprises

$\leq 10\%$ mineral SOM[19,29–32], meaning a substantial majority of SOM biochemistry has been

considered unaltered by N deposition. A higher resolution analysis of SOM chemistry and the soil

microbiome through decomposition and N deposition remains a critical gap in our knowledge on

the persistence of SOM that built up under anthropogenic N deposition.

By using high resolution SOM biochemical and soil microbiome datasets, in the context of

a long-term field study, the primary objective of this study was to assess the effect of experimental

N deposition on the trajectory of organic matter decomposition and stabilization from undecayed

plant root litter to mineral SOM. We assessed the effect of experimental N deposition on SOM

molecular abundance, diversity, and complexity during microbial decay by establishing a

decomposition gradient from undecomposed fine roots to mineral SOM. Here, we define SOM

molecular diversity analogous to species richness and evenness, where molecular richness refers

to the number of different biochemical compounds in SOM and molecular evenness describes their

relative abundance. We define SOM molecular complexity as how structurally (dis)similar

biochemical compounds that constitute SOM are to one another[33]. More molecularly complex

SOM would thus have fewer structural similarities between compounds. We hypothesized that

experimental N deposition will elicit changes in organic matter biochemistry at the individual

compound-level that are not detected when assessed at the less refined compound class-level, and

these compound substitutions may have functional implications for storage of SOM in the system.

We further hypothesized that changes in SOM biochemistry will be associated with changes in

microbial community composition that can explain the observed biochemical changes.

2. Methods

2.1. Description of study sites

The Michigan Nitrogen Deposition Gradient Study was established in 1987 to examine the

effects of atmospheric N deposition on ecosystem biogeochemical processes. Spanning a 500-km

distance, the study area in Michigan, USA, consists of four replicate sugar maple (*Acer saccharum*)

dominated northern hardwood forest stands on sandy spodosol soil (Typic Haplorthods of the

Kalkaska series). The stands contain a thick organic horizon that includes a mat of fine roots

(Oe/Oa) at its boundary with the A horizon. The stands are floristically and edaphically matched

and encompass the full latitudinal range of northern hardwood forests in eastern North America

and the Upper Great Lakes Region[34], allowing us to generalize our findings across this widespread

ecologically and economically important ecosystem (Table S1). At each site ($n = 4$), there are six

30-m by 30-m plots, three of which received experimental N additions (30 kg $NO_3{}^-$N ha^{-1} yr^{-1}) in

the form of $NaNO_3$ pellets applied annually in six equal applications throughout the growing

season beginning in 1994 and three of which receive ambient atmospheric N deposition (Figure

1).

2.2. Field-based experiment, sampling, and sample processing

This field-based experiment was designed to produce samples that constitute a gradient of

fine root decomposition, spanning undecomposed fine roots, 1-year decomposed fine roots, and

more highly decomposed mineral soil organic matter. To do this, samples were collected in

September and October 2013 by collecting soil cores to a depth of 10cm (5-cm-diameter), which

included both Oe/Oa and A horizons[23,35]. Cores were transported on ice to the University of

Michigan and stored at -20°C. Cores were thawed and passed through a 2mm sieve, and first-

through third-order fine roots were collected and pooled at the plot level[36,37]. Roots were rinsed to

remove mineral soil and dried at 60°C for 24 hours. The outcome of this processing yielded the

"undecomposed" and "mineral SOM" sample fractions (Figure 1). For this purpose, roots from

fresh Oe/Oa horizon samples are considered undecomposed in that relative to the other sample

types in this experiment, they have undergone decomposition to a lesser degree as indicated by a

light coloration and lack of observable decay. Organic material in the mineral soil represents

organic matter that has undergone the greatest degree of decomposition and exists as a relatively

stable form of SOM compared to organic horizon material[38].

To obtain the intermediate, "decomposed" samples, a subset of the undecomposed fine roots

from each plot were sterilized and placed in mesh litter bags, deployed in the field, and collected

after 12 months. Prior to field incubation, fine roots were sterilized to remove soil microbes using

ethylene oxide[39], a process that minimally alters organic matter biochemistry[40], as compared to

other methods such as autoclaving[41]. Sterilized fine roots (~2g dry weight) were placed into

15x15cm mesh litter bags constructed with 300 µm polyester mesh on top and 20µm polyester

mesh on bottom to allow for microfauna and fungal hyphae to enter the bags[35,42]. Litter bags were

deployed in their respective plots to their original location in the soil profile at the boundary of the

Oe/Oa and A horizons ($n = 3$ litter bags/plot at 6 plots/site; $n = 72$ litter bags total). Litter bags

were placed in the field in June 2014 and collected in June 2015, immediately storing each bag on

ice. The contents of each litter bag were homogenized by hand and composited at the plot level.

The decomposed fine root samples were subsampled for biochemical analyses, dried at 60°C for

24 hours, and the remaining decomposed fine root material was stored at -80°C for microbial

community analyses. Supporting justification and additional details on sample collection and

processing can be found in Argiroff et al. (2019)[23].

2.3. Organic matter biochemistry

Subsets of each of the three sample types (undecomposed fine roots, decomposed fine roots,

and mineral soil; herein referred to as three different stages of decomposition) were ground with a

ball mill and assessed for organic matter biochemistry using pyrolysis-GC/MS[43]. Samples were

pyrolyzed at 600°C in quartz tubes for 20 seconds using a DS Pyroprobe 5150 pyrolyzer and

analyzed using a ThermoTrace GC Ultra gas chromatograph (Thermo Fisher Scientific, Walthan,

MA, USA) and ITQ 900 mass spectrometer (Thermo Fisher Scientific)[23,44] at the University of

New Hampshire. Compound peaks were identified and assigned using the Automated Mass

Spectral Deconvolution and Identification Systems (AMDIS v2.65) software, the National

Institute of Standards and Technology (NIST) compounds library, and published literature[13,43].

Organic matter compounds are expressed as the % relative abundance of total sample peak area

and classified at the individual compound-level and grouped by compound source origin (e.g.,

lignin-derivatives, phenolics, aromatics, proteins, N-bearing, polysaccharides, and lipids).

Critically, individual compounds observed in pyrolysis-GC/MS analysis are pyrolysis products of

a source compound and are classified according to the nature of their source compound.

Identifiable compounds with potentially multiple or unknown origins were classified as

unspecified[13]. In total, this effort yielded 260 identifiable compounds, each assigned to one of the

8 compound source classes described above (Figure 1).

2.4. Fungal and bacterial community composition

Fungal and bacterial community composition was determined from the remaining subset of

decomposed root litter and characterized using ribosomal DNA sequence abundances. Supporting

justification and additional details on DNA isolation, PCR amplification, amplicon sequencing,

and sequence quality control can be found in Argiroff et al. (2019)[23]. Briefly, total genomic DNA

from three replicate subsamples taken from each litter bag was isolated using the DNeasy Plant

Mini Kit (Qiagen, Germantown, MD) and checked for quality with a NanoDrop 8000

Spectrophotometer (Thermo Scientific) and gel electrophoresis. Replicate extractions from each

litter bag were pooled and stored at -80°C prior to PCR amplification. PCR amplification was

performed using the LROR and LR3 primers[45] that target the D1-D2 region of the 28S rRNA gene.

The V1-V3 regions of the bacterial 16S rRNA gene were targeted using the primers 27f and 519r[46].

PCR products were purified using the MinElute PCR Purification Kit (Qiagen) and the quality of

the purified PCR products was assessed as described above. Sequencing was performed at the

University of Michigan DNA Sequencing Core on 16 SMRT chips with a PacBio RS II system

(Pacific Biosciences, Menlo Park, CA). Mean amplicon lengths were 688 bp for fungal 28S and

525 bp for bacterial 16S rRNA genes. Sequences were processed using mothur v1.40.5[47]. Fungal

sequences were aligned against a 28S reference alignment from the RDP LSU training set[48] and

bacterial sequences were aligned against the 16S SILVA v132 reference alignment[49]. Sequences

were clustered into operational taxonomic units (OTUs) for fungal and bacterial sequences at 99%

and 97% sequence similarity, respectively. The most abundant sequence for each OTU was used

as the representative for that OTU, and taxonomic assignments were made using the RDP classifier

with the LSU training set v11 for fungi[50] and the SILVA v132 reference alignment with the naïve

Bayesian classifier[51] in mothur for bacteria.

2.5. Statistical Analyses

To assess the effect of experimental N deposition on the trajectory of organic matter

biochemistry during decay, we assessed compositional change elicited by the treatment in a three-

step approach at two increasing levels of biochemical resolution. At the least resolved level, herein

referred to as the compound class-level, we used two-way analysis of variance (ANOVA) to test

the effect of experimental N deposition, site, and their interaction on log_2-transformed compound

class ($n = 8$) abundances for each of the three decomposition stages. At the higher resolution level

of analysis, herein referred to as the individual compound-level, we performed PERMANOVA on

Euclidean distance matrices of log_2-transformed relative abundances of individual compounds at

the three decomposition stages testing for the effect of decomposition stage, treatment, site, and

their interactions on organic matter biochemistry. In the case of a significant PERMANOVA

outcome, we performed PERMDISP[52] to determine whether this effect was driven by differences

in composition, heterogenous variance (i.e., dispersion), or both. The third step in our approach,

at the individual compound-level, we additionally performed differential abundance analysis

(Wald test)[53,54] to test for the effect of N deposition on individual compound abundances in

decomposition stages with a significant PERMANOVA treatment effect.

We measured organic matter molecular diversity of each sample ($n = 24$ per decomposition

stage; 72 total) by calculating molecular richness and evenness using the vegan package (v2.6.4)[55]

in R and performed two-way ANOVA to test the effect of decomposition stage, N deposition, and

their interaction on each molecular diversity metric. When applicable, pairwise comparisons were

made using Tukey's honest significant difference test (HSD) to control for multiple comparisons[56].

Molecular complexity values were calculated using the Bertz/Hendrickson/Ihlenfeldt formula[57,58]

computed by Cactvs 3.4.8.18 (PubChem release 2021.10.14)[59] and are estimates of how

complicated a compound's molecular structure is in terms of both the elements contained and

structural symmetry. We created molecular networks to represent changes in molecular complexity

between compounds present in the ambient and experimental N deposition treatment through

decomposition using the chemodiv package (v0.3.0)[33,60] in R and performed ANOVA to test the

effect of N deposition on molecular complexity.

To assess how experimental N deposition altered microbial community composition and

relationships with organic matter biochemistry, we performed Mantel tests on $\log_2$-transformed

Euclidean distances of organic matter biochemistry and Hellinger-transformed Bray-Curtis

distances of bacterial, fungal, and saprotroph communities (as defined using FUNGuild in R)[61].

To determine the taxa that most differed in relative abundance across the decomposition gradient,

differential abundance analysis was performed on the bacterial and fungal communities using

DESeq2 with *P*-value adjustment for multiple comparisons using the false discovery rate

method[62,63].

Due to the broad geographic expanse of our experiment and inherent heterogeneity of the

soil environment, we accepted statistical significance at $\alpha = 0.1$. All data processing, statistical

analyses, and visualizations were performed and produced in R (v4.2.2)[64] and RStudio (version

2022.12.0+353)[65], relying extensively on R packages such as vegan (v2.6.4)[55], easyanova

(v11.0)[66], and the collection of packages comprising the tidyverse (v1.3.1)[67].

3. Results

3.1 Compound class biochemistry

At the less-refined compound class-level, experimental N deposition significantly

increased the relative abundance of lignin-derived compounds in the mineral SOM (+53%, *P* =

0.09; Figure S1; Table S2) across all sites. However, the experimental N deposition did not affect

the abundance of any other compound class at any stage of decomposition (Table S3 and S4).

3.2 Individual compound biochemistry

When assessing the effect of experimental N deposition on the higher resolution, individual

compound relative abundances across the three decomposition stages, the composition of

individual compounds differed dependent on decomposition stage, site, and the N deposition

treatment (decomposition stage × site × treatment; F = 1.29, *P* = 0.05; Figure 2; Table S5). Pairwise

PERMANOVA at each decomposition stage revealed a significant effect of experimental N

deposition on mineral SOM biochemistry (F = 4.26, *P* = 0.004; Figure 2), and no effect on the

organic matter biochemistry at the undecomposed (F = 0.18, *P* = 0.93) or 1-yr decomposed (F =

0.20, *P* = 0.92) stages of decomposition (Table S5). Permutation tests of homogeneity of

multivariate dispersion on the organic matter biochemistry at each of the three decomposition

stages were nonsignificant ($P > 0.1$; Table S6), confirming that PERMANOVA detected

differences in mineral SOM biochemical composition at the individual compound-level between

the ambient and experimental N deposition treatment, rather than detecting changes in variability

among samples (i.e., multivariate dispersion). Differential abundance tests of individual compound

abundances in mineral SOM revealed that experimental N deposition led to the differential

abundance of 19 out of 216 total compounds detected in the mineral SOM (Figure 3). Of these

differentially abundant compounds, 7 (3.2% of the total number of compounds) were significantly

more abundant in the N deposition treatment and 12 compounds (5.6% of total) were significantly

more abundant in the ambient treatment (all $P \leq 0.1$; Table S7). Compounds more abundant under

experimental N deposition compared to ambient conditions included: phenol, 3-methyl- (aromatic

combustion product), anthracene (aromatic), 4-guaiacylpropane (lignin), benzeneacetic acid, 4-

hydroxy-3-methoxy- (lignin), benzene 1-methoxy-4-methyl- (lignin), n-decane (lipid), and 3-

pyridinol (N-bearing). Compounds more abundant in the ambient treatment included: benzene 1-

ethenyl-4-methoxy- (lignin), propene (lipid), isocrotonic acid (lipid), 3-methylpyridazine (N-

bearing), ethanone, 1-(1-methyl-1H-pyrrol-2-yl)- (N-bearing), 2,5-furandione, 3-methyl- (N-

bearing), 2(5H)-furanone, 5-methyl- (polysaccharide), 1,2-cyclohexanedione (polysaccharide),

acetic anhydride (polysaccharide), benzenepropanenitrile (protein), fluoranthene (unspecified),

and (E)-1,3-butadien-1-ol (unspecified).

3.3 Biochemical diversity and complexity

Experimental N deposition did not affect molecular species richness (F = 0, $P = 0.97$) or

evenness (F = 0.39, $P = 0.53$) at any of the three decomposition stages (Figure 4; Table S8). Given

the observed treatment differences in biochemical composition in the mineral SOM only, we

assessed the effect of experimental N deposition on molecular complexity at only the mineral SOM

decomposition stage. To determine how experimental N deposition affects molecular complexity

(i.e., how structurally similar compounds that constitute SOM are to one another), compound co-

occurrence and structural dissimilarity matrices were generated for mineral SOM biochemistry in

both treatments. Under ambient conditions in the mineral SOM, molecular network analysis

revealed 197 nodes, representing compounds, and 2450 edges, representing the structural

similarities present between those compounds (Figure 5). Molecular network analysis of the

mineral SOM biochemistry in experimental N deposition revealed it did not significantly alter the

number of compounds present (n_{nodes} = 187) nor the structural similarities between compounds

(n_{edges} = 2502; Figure 5). Molecular complexity values calculated using the

Bertz/Hendrickson/Ihlenfeldt formula affirmed that the N deposition treatment did not

significantly alter individual compound molecular complexity in the mineral SOM (F = 0, P =

0.997; Figure S2).

Decomposition stage had a significant effect on molecular richness (F = 9.49, P<0.001)

and evenness (F = 12.22, P<0.001; Figure 4; Table S8). HSD-adjusted pairwise comparisons

revealed significantly higher molecular richness at the 1-yr decomposed stage compared to the

undecomposed (post hoc P = 0.001) and mineral SOM (post hoc P<0.001; Figure 4; Table S9).

Molecular evenness was significantly higher in the 1-yr decomposed (post hoc P = 0.002) and

mineral SOM (post hoc P<0.001) stages of decomposition as compared to the undecomposed root

litter (Figure 4; Table S9). There was no difference in molecular evenness between the 1-yr

decomposed and mineral SOM decomposition stages (post hoc P = 0.39; Figure 4; Table S9). In

the undecomposed root litter, molecular network analysis revealed 209 nodes and 2095 edges,

while the 1-yr decomposed root litter network had 225 nodes and 3372 edges, and the mineral

SOM network had 216 nodes and 3022 edges (Figure S3). Molecular complexity values calculated

using the Bertz/Hendrickson/Ihlenfeldt formula affirmed that decomposition stage did not

significantly alter individual compound molecular complexity at any point (F = 0.01, P = 0.99,

Figure S4).

3.4 Microbial community associations with SOM biochemistry

Fungal and bacterial communities were differentially associated with organic matter

biochemistry at the three decomposition stages in the ambient and N deposition treatments (Figure

6). In the ambient treatment, bacterial community composition was significantly correlated with

organic matter biochemistry in the undecomposed root litter (R = 0.29, P = 0.07) and weakly

associated in the mineral SOM (R = 0.26, P = 0.12) but exhibited no relationship with the organic

matter biochemistry in the 1-yr decomposed root litter (Table S10). Under experimental N

deposition, bacterial community composition and organic matter biochemistry were not correlated

at any of the three decomposition stages. In the ambient treatment, fungal community composition

was also not correlated with the 1-yr decomposed root litter (Table S10) and was significantly

correlated with the organic matter biochemistry in the mineral SOM (R = 0.20, P = 0.09). Under

experimental N deposition, fungal community composition was significantly correlated with the

1-yr decomposed root litter biochemistry (R = 0.23, P = 0.04) and marginally associated in the

mineral SOM (R = 0.17, P = 0.11) but exhibited no relationship with the organic matter

biochemistry in the undecomposed root litter (Table S10; Figure 6). In the ambient treatment,

saprotroph community composition was marginally associated with organic matter biochemistry

in the undecomposed root litter (R = 0.18, P = 0.15) and significantly correlated with the mineral

SOM (R = 0.30, P = 0.06) but exhibited no relationship with the organic matter biochemistry in

the 1-yr decomposed root litter (Table S10; Figure 6). Under experimental N deposition,

saprotroph community composition and organic matter biochemistry were not correlated at any of

the three decomposition stages.

The abundances of 28 bacterial OTUs (1.6% of total OTUs) were significantly higher under

experimental N deposition as compared to the ambient treatment (Figure S5, Table S11). These

OTUs included members of the phylum Bacteroidetes (39%), Proteobacteria (29%),

Actinobacteria (14%), Patescibacteria (11%), and Armatimonadetes (7%). Under ambient

conditions, 13 bacterial OTUs (0.73% of total OTUs) were present in significantly greater

abundance. The phyla included Bacteroidetes (54%), Proteobacteria (38%), and Acidobacteria

(8%). The abundances of 14 fungal OTUs (2.6% of total OTUs) were significantly higher under

experimental N deposition as compared to the ambient treatment (Figure S5, Table S12). These

OTUs included members of the fungal classes Tremellomycetes (42%), Eurotiomycetes (17%),

Leotiomycetes (17%), Agaricomycetes (8%), Dothideomycetes (8%), and Sordariomycetes (8%).

Under ambient conditions, 2 fungal OTUs (0.43% of total OTUs) were present in significantly

greater abundance as compared to the N deposition treatment. The fungal classes include

Agaricomycetes (50%) and Sordariomycetes (50%).

4. Discussion

4.1 High resolution molecular analyses reveal N deposition elicits changes to SOM

biochemistry that were previously undetectable.

It is common to assess SOM biochemistry by grouping individual biochemical compounds

into classes, such as lipids or lignin-derived compounds, as a means to simplify biochemical

heterogeneity and diversity by grouping compounds that are suspected to behave similarly based

on the relative similarity of their structure or other physical properties or factors[12,13,23,68,69]. In doing

so, the abundances of individual compounds within a compound class are combined, leading to a

loss in the potential resolution of SOM biochemical information, and potentially making impacts

of environmental change on SOM biochemical constituents undetectable. As the resolution at

which we can examine SOM biochemistry has increased with technological advances, so has the

importance in identifying and elucidating the complexity within these traditional compound

classes. While it has been common to assume that individual compounds within a class of

compounds behave uniformly, there is increasing evidence for contrasting and compounding

origins within a compound class and the different potential pathways of mineralization will have

implications for the formation and persistence of mineral SOM[2,7,19,20]. As such, the implementation

of more refined analytical analyses that aid in elucidating these molecular-level differences can

increase understanding of mechanisms governing SOM persistence; a necessary step toward

increasing organic matter in soil as a viable climate change mitigation strategy[15,20].

To demonstrate this assertion, previous research efforts in this long-term field experiment

have detected increases in lignin-derived compounds in mineral SOM under experimental N

deposition, with no detectable changes to other components of mineral SOM biochemistry[23,24].

However, high resolution analysis used in this study revealed numerous individual compounds

changed in mineral SOM within multiple additional compound classes in response to the

experimental N deposition that may have consequences for the process of organic matter decay

and soil C storage (Figure 3; Table S7). For example, the abundance of lipids in mineral SOM

exhibited no effect of experimental N deposition at the compound class-level. However, at the

individual compound-level, in the ambient treatment, two molecules that are produced from the

combustion of lipids (propene and isocrotonic acid) were 15-times and over 200-times more

abundant, respectively, compared to the experimental N deposition treatment (Figure 3; Table S7).

Whereas under experimental N deposition, one lipid molecule (n-decane) was over 100-times more

abundant compared to in the ambient treatment (Figure 3; Table S7), ultimately leading to no

observable effect of the experimental N deposition on the abundance of lipids as a class of

compounds (Table S2; Figure S1). This reorganization of individual compounds within compound

classes is undetectable by less refined analysis and is critical for understanding the fate of these

substrates. Lipids are a highly heterogenous class of compounds with significant input from both

plants and soil microbes[70,71]. Though there is rising evidence that stable mineral SOM is largely

microbial-derived, it remains unclear and highly context dependent how much is plant-

derived[13,19,20,72–75]. Estimations of microbial necromass C in forest soils suggest that a substantial

portion (~70%) of C in stabilized SOM could be plant-derived[76,77], and that the contribution of

plant-derived lipids to this C pool can be greater than that of lignin[20]. In this study, all three of the

differentially abundant lipid molecules across the ambient and experimental N deposition

treatments are involved in both plant and microbial metabolic processes[78–80]. As such, more

detailed knowledge of the specific metabolic reactions, which plants or microbes are producing

them, and how these compounds end up in mineral SOM is necessary to understand the longevity

of these substrates in the system[81].

Proteins and other N-containing compounds exhibited similar trends at these differing

analytical scales. Despite no statistical difference in the abundance of proteins at the compound

class-level, at the individual compound-level one protein molecule (benzenepropanenitrile) was

almost 8-times more abundant in the ambient treatment compared under experimental N deposition

(Figure 3; Table S7). The hydrolysis of benzenepropanenitrile (i.e., 3-phenylpropionitrile) is

catalyzed by the nitrilase enzyme (EC 3.5.5.1)[82–85], yielding a carboxylic acid and ammonium

(NH_4^+). The gene which encodes the enzyme that ultimately carries out this reaction is found in

the genomes of two soil bacteria, one of which being *Paraburkholderia phymatum*[83,86–89]. In this

study, *Paraburkholderia spp.* ($n = 3$; Table S11) were in significantly greater abundance under

experimental N deposition compared to the ambient treatment (Figure S4). *Paraburkholderia spp.*

are known decomposers in forests, particularly sugar maple forests like the ones in this study[90].

Thus, it is plausible that increased enzymatic activity produced by *Paraburkholderia spp.* under

experimental N deposition catalyzed this reaction, leading to significantly lower

benzenepropanenitrile abundance in the N deposition treatment. These findings suggest that

changes to microbial community composition in response to environmental stimuli may have

functional consequences and elicit changes to the biochemical composition of organic matter that

eventually becomes stable mineral SOM.

4.2 High resolution analysis reveals novel insights into previous findings on lignin

accumulation in response to N deposition that may underlie resistance to microbial

decomposition.

In addition to the novel findings detailed above, high resolution analysis of SOM

biochemistry revealed additional compositional insight into lignin dynamics in response to the

experimental N deposition. Previous work in this long-term experiment showed that anthropogenic

N deposition slowed the rate and extent of organic matter decomposition, ultimately resulting in

the accumulation of lignified material in mineral SOM under experimental N deposition (+53%,

$P = 0.09$; Figure S1; Table S2)[23–25,28]. However, at the individual compound-level, changes in the

composition of lignin compounds in mineral SOM were detected under both ambient and

experimental N deposition conditions. In the ambient treatment, one lignin-derived molecule

(benzene, 1-ethenyl-4-methoxy-) was almost 8-times more abundant compared to under

experimental N deposition (Figure 3; Table S7). Whereas under experimental N deposition, three

different lignin-derived molecules (benzene, 1-methoxy-4-methyl-, homovanillic acid, and 4-

guaiacylpropane) were 8-times, 14-times, and 29-times more abundant than in the ambient

treatment, respectively, ultimately leading to the overall increase observed at the compound class-

level (Figure 3; Figure S1; Table S7).

At the relatively less-refined compound class-level, it is not possible to detect this

reorganization of different lignin molecules, as observed here, that ultimately affects the

interpretability of the stability of the compound class as a whole. Lignin is a complex

macromolecule that is highly variable in terms of its primary structure. Which lignin monomers

and functional groups are present, in what quantities, and what linkages connect them affect the

decomposability of lignin polymers by soil microbes[91–94]. For example, lignin polymers containing

higher quantities of monomers of the guaiacyl (G) subunit are more resistant to decomposition,

resulting in significantly lower mass loss, than lignin polymers composed of other subunits[16,93].

The 29-times greater abundance of 4-guaiacylpropane and 14-times greater abundance of

homovanillic acid, a G subunit decomposition product[95], under experimental N deposition suggests

that the SOM that has accumulated under anthropogenic N deposition is particularly

compositionally resistant to microbial decomposition in terms of its lignin content. Thus, it is

plausible that in this system, differences in the quality of lignin, in addition to the increases in

lignin quantity[23], may be a factor contributing to the increased SOM observed under high N

deposition conditions.

Differences in the composition of decomposing microbial communities observed under

experimental N deposition support the claim that lignin in mineral SOM under increased

anthropogenic N deposition is particularly resistant to decomposition or is simply not being

metabolized by the microbial community in this treatment. While fungi are widely considered as

primary lignin decomposers, evidence has shown that bacteria may play a non-inconsequential

role in lignin decay[96,97]. Of the bacterial OTUs in higher abundance under N deposition conditions,

just under half (43%) belong to the phyla Proteobacteria and Actinobacteria (Figure S4; Table

S11). This finding aligns with previous work in this system that observed an increase in the

abundance of bacteria associated with lignin decay under experimental N deposition[26,27]. The

majority of lignin-degrading bacteria identified to date represent three bacterial classes, all which

are within the Proteobacteria and Actinobacteria phyla: Actinomycetes, α-Proteobacteria, and γ-

Proteobacteria[98–100]. Such bacteria capable of lignin degradation do so in an incomplete manner,

such that smaller aromatic lignin-derived compounds are produced[101–103]. Thus, it is plausible that

the accumulation of decomposition resistant lignin material under experimental N deposition was

due, in part, to the incomplete bacterial metabolism of lignin and the inability to break down these

compounds specifically.

4.3 Molecular complexity and diversity are not drivers of SOM persistence in this system.

In this study, despite the above-described differences observed at the individual compound-

level, we did not detect overall difference in the molecular complexity of SOM between the

undecomposed root litter, 1-year decomposed root litter, and mineral SOM (Figure S3). This

finding indicates that although the composition of the individual compounds changed during decay

(Figure 2), the molecular complexity (i.e., structural [dis]similarity) of the compounds that

constitute SOM at a given stage of decomposition did not (Figure 2; Figure S3). This finding

differs from both what has been proposed conceptually about organic matter complexity

underlying SOM persistence[15] and from recent studies that have found the opposite: that persistent

SOM is less complex than its initial inputs[12,21,104]. In this system, the lack of relationship between

SOM molecular complexity and persistence may be attributable to site-specific factors or

differences in the physiological capacity of the active microbial community. The soils in our

experiment are sandy (~85%) spodosols and thus chemical sorption to mineral soil particles is a

less prominent mechanism of SOM persistence[20]. Molecular complexity may be a more important

indicator for soils with greater clay content and thus higher matrix stabilization potential.

In contrast to SOM molecular complexity, SOM diversity (i.e., molecular richness and

evenness) did change throughout decomposition (Figure 4). Both molecular richness and evenness

were higher after one year of decomposition compared to initial undecomposed inputs (Figure 4;

Table S9). This is likely the result of the production of microbially-derived compounds during the

initial phases of root decay. When microbes break down plant litter, new byproducts are

produced[7,13,105], which likely explains the observed increases in molecular richness, i.e., the

number of compounds present, after one year of decomposition. Molecular richness of mineral

SOM returning to the same levels as initial inputs may indicate the complete decay of more

accessible components of litter or that the microbially processed C substrates that were produced

during one year of decomposition were similarly consumed, returning to the same number of

compounds as initial inputs, though the composition of those compounds are distinct (Figure 2;

Figure 4).

4.4 N deposition decouples relationships between SOM and decomposing microbes,

potentially reflecting interactions between energetics and nutrient availability.

The bacterial and saprotrophic fungal communities exhibit strong relationships with the

biochemistry of the organic matter in the undecomposed litter and mineral SOM under ambient

conditions (Figure 6; Table S10). Yet, under experimental N deposition these relationships are

lost. The expression of microbial genes and the subsequent production of extracellular enzymes

that govern the breakdown of SOM are highly sensitive to N additions[28,106–109]. Under high N

conditions, genes responsible for the production of extracellular enzymes that break down complex

macromolecules, such as lignin, are downregulated[28]. Additional studies have observed a shift in

microbial community composition under N deposition conditions towards a community whose

primary niche is the decay of simpler compounds, such as cellulose[106,110,111]. Taken together, it is

plausible that when N is readily available, microbes do not need to expend the energy producing

extracellular enzymes to break down organic material and this decreased activity may result in a

shift in the community composition. This lack of microbial activity may explain why we observed

a decoupling of SOM biochemistry and the microbial community composition under experimental

N deposition.

4.5 Conclusions and implications

Our results demonstrate that N deposition altered the abundance of specific compounds in

mineral SOM, a reorganization of SOM biochemistry that is not detectable when assessing changes

at the less-refined compound class-level, as well as decoupled the relationship between microbial

composition and SOM biochemistry. Whether these observed changes have functional

implications for the long-term storage of SOM is unknown and warrants further investigation.

Though these molecular-level differences in SOM biochemistry under anthropogenic N deposition

may appear subtle, the effect that these differences have on the C stored in the system long-term

may be disproportionately large. For example, in this system, upon termination of the experimental

N deposition, a globally ecologically relevant environmental change in response to Clean Air Act

Amendments, microbial activity significantly increased and the C that built up under experimental

N deposition was lost from the system within 5-years of treatment termination[38]. Studies have

shown that molecular-level changes in SOM biochemistry, in response to various environmental

change conditions, impact the capacity of temperate forests to sequester C[112]. A microbial

mechanism may underlie this possibility as differences in the structural properties of the molecules

in greater abundance under anthropogenic N deposition may have implications for how and by

whom these products are transformed, what products are produced, and, ultimately, the

decomposability of their byproducts[80,113,114]. Taken together, it is plausible that the SOM that

accumulated under experimental N deposition was molecularly distinct as a result of functional

changes in the microbial community elicited by the N deposition treatment[23,28]. Thus, upon

termination of experimental N deposition, the accumulated, molecularly distinct SOM may have

been readily decomposed by the active microbial community upon return to low (i.e., ambient) N

conditions. By analyzing SOM biochemistry at more refined analytical scales, we were able to

elucidate the mechanisms underlying this system's fluctuating soil C dynamics in response to

ongoing environmental change.

Author Contributions

D.R.Z. designed the study. R.A.U. and W.A.A. performed field and laboratory analyses.

A.S.G. performed biochemical analyses. B.E.P. analyzed the data with the help of G.A.C. B.E.P

wrote the first draft of the manuscript. D.R.Z., W.A.A., G.A.C., and Z.B.F. provided significant

feedback on subsequent versions and analyses.

Acknowledgements

We thank Annalisa Stevenson for her help in ArcGIS. We also wish to thank Thea Whitman

(and lab), Erika Marín-Spiotta (and lab), the Department of Soil and Environmental Sciences

graduate students, and the anonymous reviewers for their thoughtful feedback on this work. This

material is based upon work supported by the National Science Foundation Graduate Research

Fellowship Program under Grant No. DGE-2137424 awarded to B.E.P. and from the United States

Department of Energy Biological Environmental Research program. The maintenance of our long-

term field experiment is supported by the National Science Foundation Long-Term Research in

Environmental Biology program.

References

1. Bowden, R. D. *et al.* Litter Input Controls on Soil Carbon in a Temperate Deciduous Forest.

Soil Sci. Soc. Am. J. **78**, (2014).

2. Cotrufo, M. F. & Lavallee, J. M. Soil organic matter formation, persistence, and functioning:

A synthesis of current understanding to inform its conservation and regeneration. in *Advances*

in Agronomy vol. 172 1–66 (Elsevier, 2022).

3. Jenkinson, D. S. & Rayner, J. H. THE TURNOVER OF SOIL ORGANIC MATTER IN

SOME OF THE ROTHAMSTED CLASSICAL EXPERIMENTS: *Soil Sci.* **123**, 298–305

(1977).

4. Parton, W. J., Schimel, D. S., Cole, C. V. & Ojima, D. S. Analysis of Factors Controlling Soil

Organic Matter Levels in Great Plains Grasslands. *Soil Sci. Soc. Am. J.* **51**, 1173–1179 (1987).

5. Adair, E. C. *et al.* Simple three-pool model accurately describes patterns of long-term litter

decomposition in diverse climates. *Glob. Change Biol.* **14**, 2636–2660 (2008).

6. Berg, B. & Meentemeyer, V. Litter quality in a north European transect versus carbon storage

potential. *Plant Soil* **242**, 83–92 (2002).

7. Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Denef, K. & Paul, E. The Microbial

Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with

soil organic matter stabilization: do labile plant inputs form stable soil organic matter? *Glob.*

Change Biol. **19**, 988–995 (2013).

8. Prescott, C. E. Litter decomposition: what controls it and how can we alter it to sequester

more carbon in forest soils? *Biogeochemistry* **101**, 133–149 (2010).

9. Zhang, D., Hui, D., Luo, Y. & Zhou, G. Rates of litter decomposition in terrestrial

ecosystems: global patterns and controlling factors. *J. Plant Ecol.* **1**, 85–93 (2008).

10.   Berg, B. & McClaugherty, C. *Plant Litter*. (Springer Berlin Heidelberg, Berlin,

Heidelberg, 2008). doi:10.1007/978-3-540-74923-3.

11.   Waksman, S. A. Humus Origin, Chemical Composition, and Importance in Nature. *Soil*

Science **41**, (1936).

12.   Jones, A. R. *et al.* Molecular complexity and diversity of persistent soil organic matter.

Soil Biol. Biochem. **184**, 109061 (2023).

13.   Kallenbach, C. M., Frey, S. D. & Grandy, A. S. Direct evidence for microbial-derived

soil organic matter formation and its ecophysiological controls. *Nat. Commun.* 10 (2016).

14.   Lehmann, J. & Kleber, M. The contentious nature of soil organic matter. *Nature* **528**, 60–

68 (2015).

15.   Lehmann, J. *et al.* Persistence of soil organic carbon caused by functional complexity.

Nat. Geosci. **13**, 529–534 (2020).

16.   Talbot, J. M., Yelle, D. J., Nowick, J. & Treseder, K. K. Litter decay rates are determined

by lignin chemistry. *Biogeochemistry* **108**, 279–295 (2012).

17.   Wickings, K., Grandy, A. S., Reed, S. C. & Cleveland, C. C. The origin of litter chemical

complexity during decomposition. *Ecol. Lett.* **15**, 1180–1188 (2012).

18.   Sugai, S. F. & Schimel, J. P. Decomposition and biomass incorporation of 14c-labeled

glucose and phenolics in taiga forest floor: effect of substrate quality, successional state, and

season. *Soil Biol. Biochem.* **25**, 1379–1389 (1993).

19.   Whalen, E. D. *et al.* Clarifying the evidence for microbial- and plant-derived soil organic

matter, and the path toward a more quantitative understanding. *Glob. Change Biol.* gcb.16413

(2022) doi:10.1111/gcb.16413.

20.   Angst, G., Mueller, K. E., Nierop, K. G. J. & Simpson, M. J. Plant- or microbial-derived?

A review on the molecular composition of stabilized soil organic matter. *Soil Biol. Biochem.*

156, 108189 (2021).

21.   Davenport, R. *et al.* Decomposition decreases molecular diversity and ecosystem

similarity of soil organic matter. *Proc. Natl. Acad. Sci.* **120**, e2303335120 (2023).

22.   Yang, S. *et al.* Soil organic carbon content and mineralization controlled by the

composition, origin and molecular diversity of organic matter: A study in tropical alpine

grasslands. *Soil Tillage Res.* **215**, 105203 (2022).

23.   Argiroff, W. A., Zak, D. R., Upchurch, R. A., Salley, S. O. & Grandy, A. S.

Anthropogenic N deposition alters soil organic matter biochemistry and microbial

communities on decaying fine roots. *Glob. Change Biol.* **25**, 4369–4382 (2019).

24.   Zak, D. R., Holmes, W. E., Burton, A. J., Pregitzer, K. S. & Talhelm, A. F.

SIMULATED ATMOSPHERIC NO_3^- DEPOSITION INCREASES SOIL ORGANIC

MATTER BY SLOWING DECOMPOSITION. *Ecol. Appl.* **18**, 2016–2027 (2008).

25.   Whittinghill, K. A., Currie, W. S., Zak, D. R., Burton, A. J. & Pregitzer, K. S.

Anthropogenic N Deposition Increases Soil C Storage by Decreasing the Extent of Litter

Decay: Analysis of Field Observations with an Ecosystem Model. *Ecosystems* **15**, 450–461

(2012).

26.   Freedman, Z. & Zak, D. R. Atmospheric N Deposition Increases Bacterial Laccase-Like

Multicopper Oxidases: Implications for Organic Matter Decay. *Appl. Environ. Microbiol.* **80**,

4460–4468 (2014).

27.   Freedman, Z. B., Upchurch, R. A. & Zak, D. R. Microbial Potential for Ecosystem N

Loss Is Increased by Experimental N Deposition. *PLOS ONE* **11**, e0164531 (2016).

28.   Zak, D. R. *et al.* Anthropogenic N deposition, fungal gene expression, and an increasing

soil carbon sink in the Northern Hemisphere. *Ecology* **100**, (2019).

29.   Angst, G., Mueller, K. E., Kögel-Knabner, I., Freeman, K. H. & Mueller, C. W.

Aggregation controls the stability of lignin and lipids in clay-sized particulate and mineral

associated organic matter. *Biogeochemistry* **132**, 307–324 (2017).

30.   Bol, R., Poirier, N., Balesdent, J. & Gleixner, G. Molecular turnover time of soil organic

matter in particle-size fractions of an arable soil. *Rapid Commun. Mass Spectrom.* **23**, 2551–

2558 (2009).

31.   Geng, J. *et al.* Different Molecular Characterization of Soil Particulate Fractions under N

Deposition in a Subtropical Forest. *Forests* **10**, 914 (2019).

32.   Grandy, A. S. & Neff, J. C. Molecular C dynamics downstream: The biochemical

decomposition sequence and its impact on soil organic matter structure and function. *Sci.*

Total Environ. **404**, 297–307 (2008).

33.   Petrén, H., Köllner, T. G. & Junker, R. R. Quantifying chemodiversity considering

biochemical and structural properties of compounds with the R package CHEMODIV. *New*

Phytol. **237**, 2478–2492 (2023).

34.   Burton, A. J., Ramm, C. W., Pregitzer, K. S. & Reed, D. D. Use of multivariate methods

in forest research site selection. *Can. J. For. Res.* **21**, 1573–1580 (1991).

35.   Xia, M., Talhelm, A. F. & Pregitzer, K. S. Long-Term Simulated Atmospheric Nitrogen

Deposition Alters Leaf and Fine Root Decomposition. *Ecosystems* **21**, 1–14 (2018).

36.   Pregitzer, K. S. *et al.* FINE ROOT ARCHITECTURE OF NINE NORTH AMERICAN

TREES. *Ecol. Monogr.* **72**, 293–309 (2002).

37. Xia, M., Talhelm, A. F. & Pregitzer, K. S. Fine roots are the dominant source of

recalcitrant plant litter in sugar maple-dominated northern hardwood forests. *New Phytol.* **208**,

715–726 (2015).

38. Propson, B. E., Zak, D. R., Classen, A. T., Burton, A. J. & Freedman, Z. B. Gains in soil

carbon storage under anthropogenic nitrogen deposition are rapidly lost following its

cessation. *Ecology* **105**, e4444 (2024).

39. Rose, R. E. & Bailey, R. W. Ethylene Oxide for Soil Sterilization. *Nature* **169**, 716–716

(1952).

40. Gennari, M., Negre, M. & Ambrosoli, R. Effects of ethylene oxide on soil microbial

content and some soil chemical characteristics. *Plant Soil* **102**, 197–200 (1987).

41. Berns, A. E. *et al.* Effect of gamma-sterilization and autoclaving on soil organic matter

structure as studied by solid state NMR, UV and fluorescence spectroscopy. *Eur. J. Soil Sci.*

59, 540–550 (2008).

42. Hobbie, S. E. Contrasting Effects of Substrate and Fertilizer Nitrogen on the Early Stages

of Litter Decomposition. *Ecosystems* **8**, 644–656 (2005).

43. Grandy, A. S., Strickland, M. S., Lauber, C. L., Bradford, M. A. & Fierer, N. The

influence of microbial communities, management, and soil texture on soil organic matter

chemistry. *Geoderma* **150**, 278–286 (2009).

44. Pold, G., Grandy, A. S., Melillo, J. M. & DeAngelis, K. M. Changes in substrate

availability drive carbon cycle response to chronic warming. *Soil Biol. Biochem.* **110**, 68–78

(2017).

45.   Vilgalys, R. & Hester, M. Rapid genetic identification and mapping of enzymatically

641   amplified ribosomal DNA from several Cryptococcus species. *J. Bacteriol.* **172**, 4238–4246

642   (1990).

643   46.   Lane, D. J. 16S/23S rRNA sequencing. *Nucleic acid techniques in bacterial systematics*

644   (1991).

645   47.   Schloss, P. D. *et al.* Introducing mothur: Open-Source, Platform-Independent,

646   Community-Supported Software for Describing and Comparing Microbial Communities.

647   *Appl. Environ. Microbiol.* **75**, 7537–7541 (2009).

648   48.   Mueller, R. C., Balasch, M. M. & Kuske, C. R. Contrasting soil fungal community

649   responses to experimental nitrogen addition using the large subunit RRNA taxonomic marker

650   and cellobiohydrolase I functional marker. *Mol. Ecol.* **23**, 4406–4417 (2014).

651   49.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data

652   processing and web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2012).

653   50.   Cole, J. R. *et al.* Ribosomal Database Project: data and tools for high throughput rRNA

654   analysis. *Nucleic Acids Res.* **42**, D633–D642 (2014).

655   51.   Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naïve Bayesian Classifier for Rapid

656   Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Appl. Environ. Microbiol.*

657   **73**, 5261–5267 (2007).

658   52.   Anderson, M. J. PERMANOVA_2factor: a FORTRAN computer program for

659   permutational multivariate analysis of variance (for any two-factor ANOVA design) using

660   permutation tests. *Department of Statistics, University of Auckland, Auckland* 35 (2004).

53.   Langley, S. R. & Mayr, M. Comparative analysis of statistical methods used for detecting

differential expression in label-free mass spectrometry proteomics. *J. Proteomics* **129**, 83–92

(2015).

54.   Yang, L. & Chen, J. A comprehensive evaluation of microbial differential abundance

analysis methods: current status and potential solutions. *Microbiome* **10**, 130 (2022).

55.   Oksanen, J. *et al.* vegan: Community Ecology Package. 2.7–1

https://doi.org/10.32614/CRAN.package.vegan (2001).

56.   Yandell, B. S. *Practical Data Analysis for Designed Experiments*. (Routledge, 1997).

doi:10.1201/9780203742563.

57.   Bertz, S. H. The first general index of molecular complexity. *J. Am. Chem. Soc.* **103**,

3599–3601 (1981).

58.   Hendrickson, J. B., Huang, P. & Toczko, A. G. Molecular complexity: a simplified

formula adapted to individual atoms. *J. Chem. Inf. Comput. Sci.* **27**, 63–67 (1987).

59.   Kim, S. *et al.* PubChem 2025 update. *Nucleic Acids Res.* **53**, D1516–D1525 (2025).

60.   Petrén, H., Köllner, T. G. & Junker, R. R. chemodiv: Analysing Chemodiversity of

Phytochemical Data. 0.3.1 https://doi.org/10.32614/CRAN.package.chemodiv (2022).

61.   Nguyen, N. H. *et al.* FUNGuild: An open annotation tool for parsing fungal community

datasets by ecological guild. *Fungal Ecol.* **20**, 241–248 (2016).

62.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. *Journal of the Royal statistical society: series B* **57**,

289–300 (1995).

63.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

64.   R Core Team. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing (2022).

65.   RStudio Team. RStudio: Integrated development environment for R. (2022).

66.   Arnhold, E. easyanova: Analysis of Variance and Other Important Complementary

Analyses. 11.0 https://doi.org/10.32614/CRAN.package.easyanova (2012).

67.   Wickham, H. *et al.* Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

68.   Geng, J., Fang, H., Cheng, S. & Pei, J. Effects of N deposition on the quality and quantity

of soil organic matter in a boreal forest: Contrasting roles of ammonium and nitrate. *CATENA*

198, 104996 (2021).

69.   Sanderman, J. & Grandy, A. S. Ramped thermal analysis for isolating biologically

meaningful soil organic matter fractions with distinct residence times. *SOIL* **6**, 131–144

(2020).

70.   Arteaga-Clemente, G., García-González, M. A. & González-González, M. Soil lipid

analysis by chromatography: A critical review of the current state in sample preparation. *J.*

Chromatogr. Open **6**, 100173 (2024).

71.   Dinel, H., Schnitzer, M. & Mehuys, G. R. Soil Lipids: Origin, Nature, Content,

Decomposition, and Effect on Soil Physical Properties. in *Soil Biochemistry* 397–430

(Routledge, 2017). doi:10.1201/9780203739389-8.

72.   Kindler, R., Miltner, A., Thullner, M., Richnow, H.-H. & Kästner, M. Fate of bacterial

biomass derived fatty acids in soil and their contribution to soil organic matter. *Org.*

Geochem. **40**, 29–37 (2009).

73.   Miltner, A., Bombach, P., Schmidt-Brücken, B. & Kästner, M. SOM genesis: microbial

biomass as a significant source. *Biogeochemistry* **111**, 41–55 (2012).

74.   Olivelli, M. S. *et al.* Unraveling Mechanisms behind Biomass–Clay Interactions Using

Comprehensive Multiphase Nuclear Magnetic Resonance (NMR) Spectroscopy. *ACS Earth*

Space Chem. **4**, 2061–2072 (2020).

75.   Rempfert, K. R. *et al.* Biomolecular budget of persistent, microbial-derived soil organic

carbon: The importance of underexplored pools. *Sci. Total Environ.* **932**, 172916 (2024).

76.   Khan, K. S., Mack, R., Castillo, X., Kaiser, M. & Joergensen, R. G. Microbial biomass,

fungal and bacterial residues, and their relationships to the soil organic matter C/N/P/S ratios.

Geoderma **271**, 115–123 (2016).

77.   Liang, C., Amelung, W., Lehmann, J. & Kästner, M. Quantitative assessment of

microbial necromass contribution to soil organic matter. *Glob. Change Biol.* **25**, 3578–3590

(2019).

78.   Fahy, E. *et al.* Update of the LIPID MAPS comprehensive classification system for

lipids. *J. Lipid Res.* **50**, S9–S14 (2009).

79.   Liebisch, G. *et al.* Update on LIPID MAPS classification, nomenclature, and shorthand

notation for MS-derived lipid structures. *J. Lipid Res.* **61**, 1539–1555 (2020).

80.   Tian, Z. *et al.* PMhub 1.0: a comprehensive plant metabolome database. *Nucleic Acids*

Res. **52**, D1579–D1587 (2024).

81.   Klier, K. M. & Anantharaman, K. An updated view of metabolic handoffs in

microbiomes. *Trends Microbiol.* S0966842X25002203 (2025) doi:10.1016/j.tim.2025.07.009.

82.   Harper, D. B. Microbial metabolism of aromatic nitriles. Enzymology of C–N cleavage

by *Nocardia* sp. (*Rhodochrous* group) N.C.I.B. 11216. *Biochem. J.* **165**, 309–319 (1977).

83.   Howden, A. J. M., Rico, A., Mentlak, T., Miguet, L. & Preston, G. M. *Pseudomonas*

syringae pv. *syringae* B728a hydrolyses indole-3-acetonitrile to the plant hormone indole-3-

acetic acid. *Mol. Plant Pathol.* **10**, 857–865 (2009).

84.   Thimann, K. V. & Mahadevan, S. Nitrilase. *Arch. Biochem. Biophys.* **105**, 133–141

(1964).

85.   Pace, H. C. & Brenner, C. The nitrilase superfamily: classification, structure and

function. *Genome Biol.* **2**, reviews0001.1 (2001).

86.   Bessonnet, T. *et al.* Purification and Characterization of Nitphym, a Robust Thermostable

Nitrilase From Paraburkholderia phymatum. *Front. Bioeng. Biotechnol.* **9**, 686362 (2021).

87.   Moulin, L. *et al.* Complete Genome sequence of Burkholderia phymatum STM815T, a

broad host range and efficient nitrogen-fixing symbiont of Mimosa species. *Stand. Genomic*

Sci. **9**, 763–774 (2014).

88.   Teepakorn, C. *et al.* Nitrilase immobilization and transposition from a micro-scale batch

to a continuous process increase the nicotinic acid productivity. *Biotechnol. J.* **16**, 2100010

(2021).

89.   Vergne-Vaxelaire, C. *et al.* Nitrilase Activity Screening on Structurally Diverse

Substrates: Providing Biocatalytic Tools for Organic Synthesis. *Adv. Synth. Catal.* **355**, 1763–

1779 (2013).

90.   Wilhelm, R. C., DeRito, C. M., Shapleigh, J. P., Madsen, E. L. & Buckley, D. H.

Phenolic acid-degrading *Paraburkholderia* prime decomposition in forest soil. *ISME*

Commun. **1**, 4 (2021).

91.   Bhatnagar, J. M., Peay, K. G. & Treseder, K. K. Litter chemistry influences

decomposition through activity of specific microbial functional guilds. *Ecol. Monogr.* **88**,

429–444 (2018).

92.   Pisani, O., Frey, S. D., Simpson, A. J. & Simpson, M. J. Soil warming and nitrogen

deposition alter soil organic matter composition at the molecular-level. *Biogeochemistry* **123**,

391–409 (2015).

93.   Thirunavukkarasu, A. *et al.* Unraveling the dynamics of lignin chemistry on

decomposition to understand its contribution to soil organic matter accumulation. *Plant Soil*

(2024) doi:10.1007/s11104-024-07066-y.

94.   vandenEnden, L. *et al.* Molecular-level changes in soil organic matter composition after

10 years of litter, root and nitrogen manipulation in a temperate forest. *Biogeochemistry* **141**,

183–197 (2018).

95.   Alruwaili, A. *et al.* Elucidation of microbial lignin degradation pathways using synthetic

isotope-labelled lignin. *RSC Chem. Biol.* **4**, 47–55 (2023).

96.   DeAngelis, K. M. *et al.* Characterization of Trapped Lignin-Degrading Microbes in

Tropical Forest Soil. *PLoS ONE* **6**, e19306 (2011).

97.   DeAngelis, K. M. *et al.* Evidence supporting dissimilatory and assimilatory lignin

degradation in Enterobacter lignolyticus SCF1. *Front. Microbiol.* **4**, (2013).

98.   Bugg, T. D. H., Ahmad, M., Hardiman, E. M. & Rahmanpour, R. Pathways for

degradation of lignin in bacteria and fungi. *Nat. Prod. Rep.* **28**, 1883 (2011).

99.   Janusz, G. *et al.* Lignin degradation: microorganisms, enzymes involved, genomes

analysis and evolution. *FEMS Microbiol. Rev.* **41**, 941–962 (2017).

100. Lee, S., Kang, M., Bae, J.-H., Sohn, J.-H. & Sung, B. H. Bacterial Valorization of Lignin:

Strains, Enzymes, Conversion Pathways, Biosensors, and Perspectives. *Front. Bioeng.*

Biotechnol. **7**, 209 (2019).

101. Bugg, T. D., Ahmad, M., Hardiman, E. M. & Singh, R. The emerging role for bacteria in

lignin degradation and bio-product formation. *Curr. Opin. Biotechnol.* **22**, 394–400 (2011).

102. Granja-Travez, R. S., Persinoti, G. F., Squina, F. M. & Bugg, T. D. H. Functional

genomic analysis of bacterial lignin degraders: diversity in mechanisms of lignin oxidation

and metabolism. *Appl. Microbiol. Biotechnol.* **104**, 3305–3320 (2020).

103. Gu, J. *et al.* Bacterial transformation of lignin: key enzymes and high-value products.

Biotechnol. Biofuels Bioprod. **17**, 2 (2024).

104. Zhang, Z. *et al.* Changes in molecular composition, diversity, and network complexity of

soil organic carbon along the elevation in Changbai Mountain, Northeast China. *Geoderma*

443, 116842 (2024).

105. Prescott, C. E. & Vesterdal, L. Decomposition and transformations along the continuum

from litter to soil organic matter in forest soils. *For. Ecol. Manag.* **498**, 119522 (2021).

106. Frey, S. D., Knorr, M., Parrent, J. L. & Simpson, R. T. Chronic nitrogen enrichment

affects the structure and function of the soil microbial community in temperate hardwood and

pine forests. *For. Ecol. Manag.* **196**, 159–171 (2004).

107. Hesse, C. N. *et al.* Forest floor community metatranscriptomes identify fungal and

bacterial responses to N deposition in two maple forests. *Front. Microbiol.* **6**, (2015).

108. Li, X. *et al.* Nitrogen addition restricts key soil ecological enzymes and nutrients by

reducing microbial abundance and diversity. *Sci. Rep.* **15**, 5560 (2025).

109.   Moore, J. A. M. *et al.* Fungal community structure and function shifts with atmospheric

nitrogen deposition. *Glob. Change Biol.* **27**, 1349–1364 (2021).

110.   Frey, S. D. *et al.* Chronic nitrogen additions suppress decomposition and sequester soil

carbon in temperate forests. *Biogeochemistry* **121**, 305–316 (2014).

111.   Van Diepen, L. T. A., Frey, S. D., Landis, E. A., Morrison, E. W. & Pringle, A. Fungi

exposed to chronic nitrogen enrichment are less able to decay leaf litter. *Ecology* **98**, 5–11

(2017).

112.   Castañeda-Gómez, L. *et al.* Soil organic matter molecular composition with long-term

detrital alterations is controlled by site-specific forest properties. *Glob. Change Biol.*

gcb.16456 (2022) doi:10.1111/gcb.16456.

113.   Bossio, D. A. *et al.* The role of soil carbon in natural climate solutions. *Nat. Sustain.* **3**,

391–398 (2020).

114.   Crow, S. E. *et al.* Sources of plant-derived carbon and stability of organic matter in soil:

implications for global change. *Glob. Change Biol.* **15**, 2003–2019 (2009).

Figure 1. The distribution of study sites in Michigan, USA that span the geographic range of northern hardwood forests. At each site ($n = 4$), there are six 30 m x 30 m plots, three of which received annual experimental nitrogen (N) additions in the form of 30 kg $NO_3{-}N$ ha^{-1} yr^{-1} beginning in 1994 and three of which receive ambient atmospheric N deposition (i). Maps created using ArcGIS Pro: release 3.3.1 (2024; Dewitz, 2021). Cores were collected ($n = 3$ per plot) to yield the undecomposed and mineral SOM samples. Subsets of undecomposed material were deployed in litter bags and left to decompose for 1-year to yield the decomposed samples (ii). The biochemistry of all three sample types was assessed using pyrolysis-GC/MS and bacterial and fungal community composition was determined on the decomposed samples (iii).

830   **Figure 2.** First two axes of non-metric multidimensional scaling (NMDS) ordination of

831   Euclidean distances of log_2 transformed py-GC/MS biochemical composition of individual

832   samples ($n = 72$, $k = 3$, stress = 0.10). Data points represent the biochemical profile of each

833   sample ($n = 6$ samples per site) and are colored by decomposition stage. The shape of the point

834   represents if it received the ambient or the 30 kg $NO_3{-}N$ ha^{-1} yr^{-1} nitrogen deposition treatment.

Figure 3. MA plot of the Log_2 Fold Change over the Mean of Normalized Counts for the mineral

SOM py-GC/MS compounds comparing compound abundance between the ambient and

nitrogen deposition treatments. Compounds with a $P_{adj} < 0.1$ are colored by their compound

source class. Colored compounds above the 0 line indicate significantly higher abundance in the

nitrogen deposition treatment, as compared to the ambient treatment, and vice versa for below

the 0 line. For significant compound test statistics, see Table S7.

Figure 4. The effect of the nitrogen deposition treatment and decomposition stage on molecular richness (left) and evenness (right). Within each figure, ambient values are displayed on the left (solid) and nitrogen deposition treatment values are displayed on the right (dashed) for each decomposition stage ($n = 3$). Individual sample values ($n = 72$) are displayed. Two-way ANOVA was used to test the effect of Decomposition Stage, Treatment, and Decomposition Stage x Treatment interaction. The effect of Decomposition Stage was significant for both richness ($F_{decomp} = 9.49$, $P < 0.001$) and evenness ($F_{decomp} = 12.22$, $P < 0.001$). Decomposition Stage pairwise comparisons (Tukey's HSD) are displayed as letters (see Supplementary Table S9).

Figure 5. Molecular network showing relationships between the biochemical compounds present

in the mineral SOM under ambient (left) and nitrogen deposition (right) conditions. Nodes

represent individual molecules (ambient: $n = 197$; nitrogen treatment: $n = 189$) and edges

represent the quantity of interactions and relationships between those molecules (ambient: $n =$

2450; nitrogen treatment: $n = 2502$). Thickness of the edges is determined by the molecular

similarity of which the cut off value was set to 0.7. Nodes are colored by the compound class to

which that individual molecule belongs. Enlarged nodes represent the molecules in greater

abundance in that treatment from differential abundance analysis (Fig. 4).

Figure 6. Mantel statistic (R) for the relationships between the bacterial (left), fungal (middle),

and saprotrophic (right) communities and the biochemistry of the organic matter at each

decomposition stage. Higher R values indicate stronger relationships (red). See Table S10 for

full Mantel statistics.

Table S1. Site, climate, vegetation, and soil characteristics of the four northern hardwood forest

stands in Michigan, USA (see Figure 2). The sites exist across a natural north-south climatic

gradient, yet are similar in age, plant composition, and soil development.

Characteristic	Site A	Site B	Site C	Site D
Location				
Latitude (N)	46°52'	45°33'	44°23'	43°40'
Longitude (W)	88°53'	84°52'	85°50'	86°09'
Climate				
Mean annual temperature (°C)	4.7	6.0	6.9	7.6
Mean annual precipitation (mm)	873	871	888	812
Ambient wet + dry total N deposition (g m^{-2} yr^{-1})	0.68	0.91	1.17	1.18
Growing season length (days)	134	150	154	157
Vegetation				
Overstory biomass (Mg ha^{-1})	261	261	274	234
Acer saccharum biomass (Mg ha^{-1})	237	224	216	201
Soil				
Soil texture (0-10cm) (%sand-%silt-%clay)	75:22:3	89:9:2	89:9:2	87:10:3

875   **Table S2.** Two-way ANOVA results for the effect of Site, Treatment, and Site x Treatment

876   interaction on $\log_2$-transformed relative abundances of the 8 compounds classes in Mineral Soil

877   Organic Matter (SOM). Significant P-values ($P \leq 0.1$) are presented in bold.

Compound class	Factor	Numerator df	Denominator df	F-value	P-value
Aromatic	Site	3	20	2.5041	**0.0962**
	Treatment	1	22	1.4847	0.2407
	Site x Treatment	3	20	0.5880	0.6316
Lignin	Site	3	20	6.3903	**0.0047**
	Treatment	1	22	3.2040	**0.0924**
	Site x Treatment	3	20	0.4467	0.7230
Lipid	Site	3	20	1.2469	0.3257
	Treatment	1	22	1.5861	0.2259
	Site x Treatment	3	20	0.1884	0.9027
N-Bearing	Site	3	20	7.3233	**0.0026**
	Treatment	1	22	0.8199	0.3786
	Site x Treatment	3	20	2.9805	**0.0626**
Phenol	Site	3	20	0.6791	0.5775
	Treatment	1	22	2.9435	0.1055
	Site x Treatment	3	20	0.4292	0.7349
Polysaccharide	Site	3	20	0.5502	0.6552
	Treatment	1	22	0.4042	0.5339
	Site x Treatment	3	20	4.5208	**0.0177**
Protein	Site	3	20	1.7234	0.2024
	Treatment	1	22	0.5559	0.4667
	Site x Treatment	3	20	0.3023	0.8233
Unspecified	Site	3	20	1.1552	0.3574
	Treatment	1	22	0.4160	0.5281
	Site x Treatment	3	20	0.8866	0.4691

Figure S1. Stacked abundance chart of the relative abundances of the 8 compound classes across

the three decomposition stages from least (left, Undecomposed) to most (right, Mineral SOM)

decomposed under ambient and simulated nitrogen deposition conditions. Across the three

decomposition stages, the nitrogen deposition treatment only significantly affected the relative

abundance of lignin in the mineral SOM (+53% in nitrogen treatment, $P = 0.09$).

Table S3. Two-way ANOVA results for the effect of Site, Treatment, and Site x Treatment

interaction on log$_2$-transformed relative abundances of the 8 compounds classes in

Undecomposed fine roots. Significant P-values ($P \leq 0.1$) are presented in bold.

Compound class	Factor	Numerator df	Denominator df	F-value	P-value
Aromatic	Site	3	20	0.9337	0.4473
	Treatment	1	22	0.2609	0.6165
	Site x Treatment	3	20	2.1511	0.1338
Lignin	Site	3	20	0.0870	0.9662
	Treatment	1	22	0.4924	0.4929
	Site x Treatment	3	20	1.2213	0.3343
Lipid	Site	3	20	1.1512	0.3588
	Treatment	1	22	0.1842	0.6735
	Site x Treatment	3	20	1.1179	0.3711
N-Bearing	Site	3	20	2.7075	**0.0799**
	Treatment	1	22	0.4525	0.5108
	Site x Treatment	3	20	1.1713	0.3516
Phenol	Site	3	20	1.5606	0.2378
	Treatment	1	22	0.3278	0.5749
	Site x Treatment	3	20	4.5536	**0.0173**
Polysaccharide	Site	3	20	4.3391	**0.0203**
	Treatment	1	22	1.5337	0.2334
	Site x Treatment	3	20	3.9589	**0.0275**
Protein	Site	3	20	0.7584	0.5337
	Treatment	1	22	0.0003	0.9860
	Site x Treatment	3	20	0.5795	0.6369
Unspecified	Site	3	20	1.5001	0.2526
	Treatment	1	22	0.3898	0.5412
	Site x Treatment	3	20	1.3106	0.3054

889   **Table S4.** Two-way ANOVA results for the effect of Site, Treatment, and Site x Treatment

890   interaction on log$_2$-transformed relative abundances of the 8 compounds classes in the 1-yr

891   Decomposed fine roots. Significant P-values ($P \leq 0.1$) are presented in bold.

Compound class	Factor	Numerator df	Denominator df	F-value	P-value
Aromatic	Site	3	20	3.5241	**0.0393**
	Treatment	1	22	0.7394	0.4026
	Site x Treatment	3	20	1.4160	0.2747
Lignin	Site	3	20	0.3242	0.8078
	Treatment	1	22	0.0587	0.8116
	Site x Treatment	3	20	1.4453	0.2668
Lipid	Site	3	20	1.1059	0.3757
	Treatment	1	22	0.0909	0.7669
	Site x Treatment	3	20	1.5248	0.2464
N-Bearing	Site	3	20	2.2232	0.1250
	Treatment	1	22	0.1254	0.7279
	Site x Treatment	3	20	0.2880	0.8334
Phenol	Site	3	20	2.2061	0.1270
	Treatment	1	22	0.4187	0.5267
	Site x Treatment	3	20	0.6662	0.5849
Polysaccharide	Site	3	20	2.5255	**0.0943**
	Treatment	1	22	1.3879	0.2560
	Site x Treatment	3	20	0.5719	0.6416
Protein	Site	3	20	2.4745	**0.0988**
	Treatment	1	22	2.6923	0.1203
	Site x Treatment	3	20	4.2287	**0.0222**
Unspecified	Site	3	20	0.1201	0.9469
	Treatment	1	22	0.3110	0.5848
	Site x Treatment	3	20	0.2234	0.8787

893   **Table S5.** Three-way PERMANOVA results for the effect of Decomposition Stage, Site,

894   Treatment, and their interactions on $\log_2$-transformed Euclidean distance matrices of individual

895   compound abundances. Below the three-way PERMANOVA results are pairwise two-way

896   PERMANOVA results for the effect of Site, Treatment, and their interaction at the three

897   decomposition stages. Significant P-values ($P \leq 0.1$) are presented in bold.

Decomp Stage	Factor	Df	R^2	F-value	P-value
	Decomp Stage	2	0.35930	20.707	**0.001**
	Site	3	0.01930	0.7416	0.645
	Treatment	1	0.01351	1.5572	0.193
	Decomp Stage x Site	6	0.06012	1.1548	0.312
	Decomp Stage x Treatment	2	0.02397	1.3816	0.211
	Site x Treatment	3	0.01440	0.5531	0.843
	Decomp Stage x Site x Treatment	6	0.09296	1.7858	**0.048**
Undecomposed	Site	3	0.12377	0.9610	0.444
	Treatment	1	0.00765	0.1783	0.934
	Site x Treatment	3	0.17265	1.4107	0.212
Decomposed	Site	3	0.13225	1.0294	0.405
	Treatment	1	0.00869	0.2028	0.924
	Site x Treatment	3	0.17384	1.3530	0.244
Mineral SOM	Site	3	0.11883	1.0912	0.362
	Treatment	1	0.15467	4.2612	**0.004**
	Site x Treatment	3	0.14573	1.3382	0.197

899 **Table S6.** Permutation test for homogeneity of multivariate dispersion results for the effect of

900 Treatment on $\log_2$-transformed Euclidean distance matrices of individual compound abundances

901 at each of the three decomposition stages.

Decomp Stage	Factor	Df	F-value	P-value
Undecomposed	Treatment	1	0.0949	0.756
Decomposed	Treatment	1	0.2859	0.612
Mineral SOM	Treatment	1	1.7029	0.213

903   **Table S7.** Mineral SOM compounds with significant differential abundance (Benjamini-

904   Hochberg $P_{adj} \leq 0.1$) and Log2 fold change values that are plotted in Figure 4. Treatment refers

905   to the treatment that the compound exhibits significantly greater abundance in.

Compound	Compound Class	Treatment	Log$_2$ Fold Change	Standard Error	Wald Statistic	BH P$_{adj}$
Phenol, 3-methyl-	Aromatic	Nitrogen	3.72780	1.12256	3.32081	0.01017
Anthracene	Aromatic	Nitrogen	3.38703	0.94245	3.59388	0.00462
Phenol, 2-methoxy-4-propyl- (4-Guaiacylpropane)	Lignin	Nitrogen	4.90243	0.97139	5.04680	0.00003
Benzeneacetic acid, 4-hydroxy-3-methoxy- (Homovanillic acid)	Lignin	Nitrogen	3.81029	0.94943	4.01324	0.00127
Benzene, 1-methoxy-4-methyl-	Lignin	Nitrogen	3.11383	0.90764	3.43070	0.00787
Benzene, 1-ethenyl-4-methoxy-	Lignin	Ambient	-2.97359	1.01665	-2.92489	0.03446
n-Decane	Lipid	Nitrogen	6.77821	1.05588	6.41951	<0.0001
Propene	Lipid	Ambient	-3.95030	1.01705	-3.88408	0.00183
Isocrotonic acid	Lipid	Ambient	-7.93390	1.07392	-7.38778	<0.0001
3-Pyridinol	N-Bearing	Nitrogen	4.62658	1.06348	4.35040	0.00039
3-Methylpyridazine	N-Bearing	Ambient	-2.74810	1.00508	-2.73422	0.05905
Ethanone, 1-(1-methyl-1H-pyrrol-2-yl)-	N-Bearing	Ambient	-3.52273	0.92806	-3.79581	0.00227
2,5-Furandione, 3-methyl-	N-Bearing	Ambient	-3.94198	1.01805	-3.87210	0.00183
2(5H)-Furanone, 5-methyl-	Polysaccharide	Ambient	-2.75395	1.07045	-2.57270	0.09029
1,2-Cyclohexanedione	Polysaccharide	Ambient	-4.35563	1.04552	-4.16602	0.00075
Acetic anhydride	Polysaccharide	Ambient	-4.98762	1.09738	-4.54501	0.00019
Benzenepropanenitrile	Protein	Ambient	-2.92576	0.86588	-3.37895	0.00884
Fluoranthene	Unspecified	Ambient	-2.66957	0.83642	-3.19167	0.01503
(E)-1,3-Butadien-1-ol	Unspecified	Ambient	-5.03650	1.09046	-4.61868	0.00016

907   **Table S8.** Two-way ANOVA results for the effect of Decomposition Stage, Treatment, and

908   Decomposition Stage x Treatment interaction on molecular richness (top) and molecular

909   evenness (bottom) diversity metrics. Significant P-values ($P \leq 0.1$) are presented in bold.

Diversity Metric	Factor	Numerator df	Denominator df	F-value	P-value
Molecular Richness	Decomp Stage	2	69	9.4898	**<0.001**
	Treatment	1	70	0.0016	0.9686
	Decomp Stage x Treatment	2	69	0.3979	0.6733
Molecular Evenness	Decomp Stage	2	69	12.224	**<0.001**
	Treatment	1	70	0.3925	0.5331
	Decomp Stage x Treatment	2	69	0.6333	0.5340

Figure S2. The effect of the nitrogen deposition treatment on the molecular complexity of
individual biochemical compounds present in the Mineral SOM ($n = 386$). No treatment effect
was detected by ANOVA ($F_{1,384} = 0$, $P = 0.997$).

930   **Table S9.** Tukey's HSD pairwise comparisons where two-way ANOVA of molecular diversity

931   metrics revealed significant ($P \leq 0.1$) effect of Decomposition Stage on molecular richness (top)

932   and evenness (bottom) (see Supplementary Table S8). Significant P-values ($P \leq 0.1$) are

933   presented in bold.

Diversity Metric	Factor	Post-hoc Comparison	Mean Difference	P-value
Molecular Richness	Decomp Stage	Undecomposed – Decomposed	-12.8750	**0.0011**
		Decomposed – Mineral SOM	13.0833	**<0.001**
		Undecomposed – Mineral SOM	0.2083	0.9980
	Decomp Stage x Treatment	Ambient Undecomposed – Ambient Decomposed	-12.5833	**0.0316**
		Ambient Decomposed – Ambient Mineral SOM	3.0000	0.8116
		Ambient Undecomposed – Ambient Mineral SOM	15.5833	**0.0059**
		Nitrogen Undecomposed – Nitrogen Decomposed	-13.1667	**0.0232**
		Nitrogen Decomposed – Nitrogen Mineral SOM	10.5834	**0.0830**
		Nitrogen Undecomposed – Nitrogen Mineral SOM	-2.5833	0.8565
Molecular Evenness	Decomp Stage	Undecomposed – Decomposed	-0.0580	**0.0026**
		Decomposed – Mineral SOM	-0.0220	0.3902
		Undecomposed – Mineral SOM	-0.0800	**<0.001**
	Decomp Stage x Treatment	Ambient Undecomposed – Ambient Decomposed	-0.0624	**0.0274**
		Ambient Decomposed – Ambient Mineral SOM	-0.0040	0.9843
		Ambient Undecomposed – Ambient Mineral SOM	-0.0664	**0.0176**
		Nitrogen Undecomposed – Nitrogen Decomposed	-0.0536	**0.0673**
		Nitrogen Decomposed – Nitrogen Mineral SOM	-0.0401	0.2136
		Nitrogen Undecomposed – Nitrogen Mineral SOM	-0.0937	**0.0005**

Figure S3. Molecular network showing relationships between the biochemical compounds at

each decomposition stage ($n = 3$). Nodes represent individual molecules (Undecomposed: $n = $

209; Decomposed: $n = 225$, Mineral SOM: $n = 216$) and edges represent the quantity of

interactions and relationships between those molecules (Undecomposed: $n = 2950$; Decomposed:

$n = 3372$, Mineral SOM: $n = 3020$). Thickness of the edges is determined by the molecular

similarity of which the cut off value was set to 0.7. Nodes are colored by their relative

abundance.

Figure S4. The effect of decomposition and the nitrogen deposition treatment on the molecular

complexity of individual biochemical compounds present at each decomposition stage (n_{total} =

1182). ANOVA detected no difference in decomposition stage (F = 0.01, P = 0.989), treatment

(F = 0.03, P = 0.852), or their interaction (F = 0.02, P = 0.976).

Table S10. Mantel test results comparing log$_2$-transformed Euclidean distance matrices of

individual compound abundances and Hellinger-transformed Bray-Curtis distance matrices of

microbial community OTU abundances. Significant P-values ($P \leq 0.1$) are presented in bold.

Type	Treatment	Decomp Stage	Mantel Statistic (R)	P-value	Upper quantiles of permutations (null model)			
					90%	95%	97.5%	99%
Bacteria (16S)	Ambient	Undecomposed	**0.288700**	**0.069**	0.255	0.314	0.360	0.406
		Decomposed	-0.001153	0.476	0.245	0.339	0.428	0.474
		Mineral SOM	0.2607	0.123	0.296	0.375	0.426	0.467
	Nitrogen	Undecomposed	-0.2065	0.912	0.227	0.278	0.328	0.371
		Decomposed	0.117	0.211	0.278	0.376	0.441	0.518
		Mineral SOM	-0.1325	0.720	0.261	0.321	0.384	0.425
Fungi (28S)	Ambient	Undecomposed	0.02193	0.455	0.203	0.251	0.299	0.341
		Decomposed	0.1921	0.107	0.200	0.237	0.273	0.326
		Mineral SOM	**0.1981**	**0.091**	0.187	0.239	0.292	0.328
	Nitrogen	Undecomposed	0.002091	0.473	0.161	0.210	0.243	0.290
		Decomposed	**0.2292**	**0.041**	0.175	0.208	0.246	0.298
		Mineral SOM	0.1662	0.105	0.170	0.216	0.243	0.298
Saprotrophs	Ambient	Undecomposed	0.1829	0.148	0.230	0.275	0.316	0.359
		Decomposed	0.09608	0.290	0.229	0.284	0.363	0.413
		Mineral SOM	**0.3031**	**0.064**	0.240	0.319	0.368	0.411
	Nitrogen	Undecomposed	-0.01454	0.518	0.229	0.281	0.335	0.382
		Decomposed	0.1114	0.242	0.256	0.336	0.408	0.467
		Mineral SOM	-0.1032	0.681	0.281	0.347	0.382	0.449

968 **Figure S4.** Plot of the Log$_2$ Fold Change over the Mean of Normalized Counts for the bacteria

969 (left) and fungi (right) genera comparing species abundance between the ambient and nitrogen

970 deposition treatments in the "decomposed" samples. Species with a $p_{adj} < 0.1$ are colored by

971 phylum. Colored species above the 0 line indicate a significantly higher abundance in the

972 nitrogen deposition treatment, as compared to the ambient treatment, and vice versa for below

973 the 0 line.

974 **Table S11.** Bacteria with significant differential abundance (Benjamini-Hochberg $P_{\mathrm{adj}} \leq 0.1$) and Log2 fold change values that are

975 plotted in Figure S4 (left). Treatment refers to the treatment that the OTU (species) exhibits significantly greater abundance in.

Kingdom	Phylum	Class	Order	Family	Genus	Treatment	Log2 Fold Change	Standard Error	Wald Statistic	BH P$_{adj}$
Bacteria	Patescibacteria	Saccharimonadia	Saccharimonadales	Unclassified	Unclassified	Nitrogen	5.39412	1.50297	3.58898	0.01505
Bacteria	Patescibacteria	Saccharimonadia	Saccharimonadales	Unclassified	Unclassified	Nitrogen	5.15561	1.19097	4.32891	0.00322
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Unclassified	Nitrogen	4.88380	1.29715	3.76504	0.01029
Bacteria	Bacteroidetes	Bacteroidia	Sphingobacteriales	Sphingobacteriaceae	Mucilaginibacter	Nitrogen	4.70389	0.84858	5.54322	0.00001
Bacteria	Actinobacteria	Actinobacteria	Micrococcales	Microbacteriaceae	Unclassified	Nitrogen	3.95287	1.29021	3.06375	0.05308
Bacteria	Actinobacteria	Actinobacteria	Micrococcales	Microbacteriaceae	Amnibacterium	Nitrogen	3.62117	1.18129	3.06544	0.05308
Bacteria	Actinobacteria	Actinobacteria	Frankiales	Nakamurellaceae	Nakamurella	Nitrogen	3.48676	1.03957	3.35404	0.02843
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Ferruginibacter	Nitrogen	3.47086	1.25134	2.77371	0.09754
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Unclassified	Nitrogen	3.46616	1.15698	2.99588	0.06041
Bacteria	Armatimonadetes	Armatimonadia	Armatimonadales	Unclassified	Unclassified	Nitrogen	3.42300	0.93394	3.66513	0.01401
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Uncultured	Nitrogen	3.38620	0.95645	3.54038	0.01688
Bacteria	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	Acidiphilium	Nitrogen	3.30733	0.93798	3.52600	0.01688
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Unclassified	Nitrogen	3.16901	1.14426	2.76950	0.09754
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Unclassified	Nitrogen	3.02005	1.05278	2.86863	0.08495
Bacteria	Bacteroidetes	Bacteroidia	Sphingobacteriales	Sphingobacteriaceae	Mucilaginibacter	Nitrogen	2.99831	1.07783	2.78179	0.09754
Bacteria	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	Unclassified	Nitrogen	2.75872	0.95259	2.89603	0.08031
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Unclassified	Nitrogen	2.72105	0.90886	2.99392	0.06041
Bacteria	Armatimonadetes	Armatimonadia	Armatimonadales	Unclassified	Unclassified	Nitrogen	2.56260	0.74517	3.43895	0.02206
Bacteria	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	Burkholderia-Caballeronia-Paraburkholderia	Nitrogen	2.34356	0.62196	3.76802	0.01029
Bacteria	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	Acidisoma	Nitrogen	2.31260	0.59484	3.88776	0.00860
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Uncultured	Nitrogen	3.38620	0.95645	3.54038	0.01688
Bacteria	Proteobacteria	Alphaproteobacteria	Acetobacterales	Acetobacteraceae	Acidiphilium	Nitrogen	3.30733	0.93798	3.52600	0.01688
Bacteria	Bacteroidetes	Bacteroidia	Sphingobacteriales	Sphingobacteriaceae	Mucilaginibacter	Nitrogen	2.01288	0.51097	3.93931	0.00794
Bacteria	Actinobacteria	Actinobacteria	Frankiales	Nakamurellaceae	Nakamurella	Nitrogen	1.96852	0.60164	3.27192	0.03310
Bacteria	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	Burkholderia-Caballeronia-Paraburkholderia	Nitrogen	1.95134	0.62661	3.11412	0.04825

Bacteria	Patescibacteria	Saccharimonadia	Saccharimonadales	Unclassified	Unclassified	Nitrogen	1.89504	0.68840	2.75283	0.09799
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Unclassified	Nitrogen	1.89319	0.67628	2.79941	0.09754
Bacteria	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	Burkholderia-Caballeronia-Paraburkholderia	Nitrogen	1.61355	0.44282	3.64382	0.01405
Bacteria	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Unclassified	Unclassified	Nitrogen	1.58444	0.48965	3.23585	0.03436
Bacteria	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	Rhizobacter	Nitrogen	1.12582	0.29837	3.77320	0.01029
Bacteria	Bacteroidetes	Bacteroidia	Sphingobacteriales	Sphingobacteriaceae	Mucilaginibacter	Ambient	-1.28789	0.40941	-3.14574	0.04506
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Unclassified	Ambient	-1.39623	0.42964	-3.24976	0.03415
Bacteria	Proteobacteria	Deltaproteobacteria	Myxococcales	Polyangiaceae	Aetherobacter	Ambient	-1.46319	0.40563	-3.60718	0.01504
Bacteria	Proteobacteria	Gammaproteobacteria	Xanthomonadales	Rhodanobacteraceae	Rhodanobacter	Ambient	-1.53376	0.38863	-3.94662	0.00794
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Unclassified	Ambient	-1.80558	0.45541	-3.96475	0.00794
Bacteria	Acidobacteria	Acidobacteriia	Acidobacteriales	Acidobacteriaceae_	Edaphobacter	Ambient	-1.83235	0.64683	-2.83280	0.09228
Bacteria	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Burkholderiaceae	AAP99	Ambient	-1.94233	0.45411	-4.27721	0.00322
Bacteria	Proteobacteria	Alphaproteobacteria	Micropepsales	Micropepsaceae	Uncultured	Ambient	-1.95804	0.65177	-3.00419	0.06041
Bacteria	Bacteroidetes	Bacteroidia	Sphingobacteriales	Sphingobacteriaceae	Mucilaginibacter	Ambient	-2.02805	0.60709	-3.34058	0.02843
Bacteria	Bacteroidetes	Bacteroidia	Chitinophagales	Chitinophagaceae	Unclassified	Ambient	-2.52434	0.44489	-5.67414	0.00001
Bacteria	Proteobacteria	Gammaproteobacteria	Betaproteobacteriales	Nitrosomonadaceae	IS-44	Ambient	-2.67385	0.95610	-2.79662	0.09754
Bacteria	Bacteroidetes	Bacteroidia	Sphingobacteriales	Sphingobacteriaceae	Mucilaginibacter	Ambient	-3.12362	1.13077	-2.76239	0.09754
Bacteria	Bacteroidetes	Bacteroidia	Sphingobacteriales	Sphingobacteriaceae	Mucilaginibacter	Ambient	-3.85663	1.17895	-3.27124	0.03310

978   **Table S12.** Fungi with significant differential abundance (Benjamini-Hochberg $P_{adj} \leq 0.1$) and Log_2 fold change values that are

979   plotted in Figure S4 (right). Treatment refers to the treatment that the OTU (species) exhibits significantly greater abundance in.

Kingdom	Phylum	Class	Order	Family	Genus	Treatment	Log_2 Fold Change	Standard Error	Wald Statistic	BH P_{adj}
Fungi	Basidiomycota	Agaricomycetes	Agaricales	Schizophyllaceae	Henningsomyces	Nitrogen	21.90978	2.94575	7.43776	<0.00001
Fungi	Ascomycota	Leotiomycetes	Unclassified	Unclassified	Unclassified	Nitrogen	4.69786	1.43680	3.26967	0.03446
Fungi	Basidiomycota	Tremellomycetes	Tremellales	Unclassified	Unclassified	Nitrogen	4.60797	1.72523	2.67094	0.09310
Fungi	Basidiomycota	Tremellomycetes	Tremellales	Unclassified	Unclassified	Nitrogen	4.36903	1.65903	2.63349	0.09659
Fungi	Ascomycota	Dothideomycetes	Pleosporales	Unclassified	Unclassified	Nitrogen	4.33137	1.34815	3.21282	0.03505
Fungi	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	Unclassified	Nitrogen	3.67041	1.26887	2.89266	0.06651
Fungi	Basidiomycota	Tremellomycetes	Tremellales	Unclassified	Unclassified	Nitrogen	3.57464	1.02756	3.47875	0.02015
Fungi	Basidiomycota	Tremellomycetes	Tremellales	Tremellaceae	Asterotremella	Nitrogen	3.37403	1.20852	2.79186	0.06987
Fungi	Ascomycota	Eurotiomycetes	Chaetothyriales	Herpotrichiellaceae	Exophiala	Nitrogen	3.14462	1.09721	2.86602	0.06651
Fungi	Ascomycota	Leotiomycetes	Helotiales	Unclassified	Unclassified	Nitrogen	2.48835	0.79990	3.11083	0.04264
Fungi	Ascomycota	Sordariomycetes	Hypocreales	Nectriaceae	Gibberella	Nitrogen	2.44656	0.87541	2.79476	0.06987
Fungi	Basidiomycota	Tremellomycetes	Tremellales	Tremellaceae	Unclassified	Nitrogen	1.39111	0.38410	3.62174	0.01808
Fungi	Ascomycota	Sordariomycetes	Unclassified	Unclassified	Unclassified	Ambient	-4.22650	1.47236	-2.87056	0.06651
Fungi	Basidiomycota	Agaricomycetes	Agaricales	Clavariaceae	Ramariopsis	Ambient	-7.48238	2.08804	-3.58345	0.01808

982   Supplemental References:

984   S Xu, M Chen, T Feng, L Zhan, L Zhou, G Yu. Use ggbreak to effectively utilize plotting space

985   to deal with large datasets and outliers. Frontiers in Genetics. 2021, 12:774846.

986   https://doi:10.3389/fgene.2021.774846