1   Anthropogenic nitrogen deposition restructures decomposing microbial communities, altering

2   SOM molecular composition, but not molecular complexity or diversity

5   Brooke E. Propson[1,2], William A. Argiroff[3], Grace A. Cagle[1], Rima A. Upchurch[3], Donald R.

6   Zak[3,4], A. Stuart Grandy[5], and Zachary B. Freedman[1*]

8   [1]Department of Soil and Environmental Sciences

9   University of Wisconsin, Madison, WI, USA 53706

10   [2]University of Michigan Biological Station

11   9133 Biological Rd, Pellston, MI, USA 49769

12   [3]School for Environment and Sustainability

13   [4]Ecology and Evolutionary Biology

14   University of Michigan, Ann Arbor, MI, USA 48109

15   [5]Department of Natural Resources and the Environment

16   University of New Hampshire, Durham, NH, USA 03824

18   *Corresponding author: Zachary B. Freedman, zfreedman@wisc.edu (ORCID: 0000-0001-9160-

19   7470)

Abstract

Soil organic matter (SOM) consists of diverse biochemical constituents, spanning a spectrum of chemical complexity, and the relative abundance of these substrates influences microbial metabolism and soil carbon persistence. However, mechanistic controls governing these processes and how they may be affected by environmental change remains incomplete. This study aims to assess (1) the molecular-level changes that occur across stages of root decomposition, from undecayed plant root litter to 1-year decomposed root litter, to mineral SOM and (2) how these changes are altered by anthropogenic nitrogen (N) deposition by using SOM biochemical and microbiome datasets and a long-term field experiment. N deposition did not significantly alter undecomposed or 1-year decomposed root litter, but did alter decomposing microbial communities and mineral SOM biochemical composition, specifically in lignin- and lipid-derived compounds. Taken together, this restructuring of microbial communities and alteration of SOM biochemistry likely contributed to the previously observed reduction in SOM decomposition.

Keywords: SOM, biochemistry, bacteria, fungi, py-GCMS, decomposition

1. Introduction

Soil organic matter (SOM) constitutes the largest terrestrial organic carbon reservoir, and altered SOM decomposition dynamics under fluctuating environmental conditions could intensify or mitigate impacts of climate change. Inputs from plants, through the deposition of above- and below-ground plant residues and the release of exudates from plant roots represents the largest organic matter input to soils[1,2]. Litter chemical composition, which is often estimated in terms of its nitrogen (N) or lignin contents[3,4], is regarded as one of the primary controls over decomposition rates and SOM accumulation and the fate of these inputs is governed by soil microbial activity[5–9]. In recent decades, there has been a shift from traditional theory of SOM chemical recalcitrance, where biochemically recalcitrant lignified material was thought to constitute the majority of SOM because of its theorized resistance to microbial decomposition[10,11], towards a continuum model of SOM complexity that proposes microbial-derived products as the primary constituents of SOM and the potential importance of molecular diversity in enabling SOM persistence[12–15].

Biochemical compounds that comprise litter exist across a spectrum of chemical complexity from simple to complex substrates and the relative abundance of these potential substrates for microbial metabolism can have consequences for the fate of C stored as mineral SOM[7,14,16]. When plant litter undergoes microbially mediated breakdown during decomposition, the resulting products will differ depending on the initial biochemical quality of the plant litter, as well as the composition of the microbial community decomposing that material[7,17]. The potential importance of biochemical complexity and diversity for SOM persistence lies in the different microbial requirements for metabolizing different biochemical compounds, both for compounds that are structurally dissimilar, as well as ones that are largely similar but may

exhibit minor differences, such as differences in the position of functional groups (for example,

ortho- versus *para-* benzoic acid)[15,18]. This rise in the acknowledgement and potential

importance of a higher resolution understanding of SOM is a deviation from the standard

practice in which compounds are typically grouped together as classes (e.g., lipids, etc.) that are

assumed to behave similarly through the decomposition process. This results in individual

compounds being grouped into classes that are inherently less refined. However, with the

conceptual move away from stable SOM largely consisting of recalcitrant material towards a

spectrum of which biochemical complexity and diversity enable SOM persistence, research has

shown that certain SOM biochemical constituents are resistant to decomposition despite being in

a compound class that was traditionally viewed as "labile" (ex: long-chain lipids)[19]. As such,

there is increasing emphasis on the potential importance of SOM molecular complexity and

diversity in driving SOM persistence[12,15,20–22]. Critically, there is a need to determine how this

proposed relationship between SOM molecular diversity and persistence responds to changing

climatic conditions[23].

To better understand how anthropogenic N deposition, a pervasive driver of global

environmental change, affects the cycling and storage of SOM, we conducted a 37-year field

experiment to simulate future rates of anthropogenic N deposition. After 20 years of

experimental N deposition inputs, we found that anthropogenic N deposition slowed the rate and

extent of organic matter decomposition, resulting in the accumulation of lignified material in

mineral soil derived primarily from fine root litter[24–26]. This change in SOM biochemistry in

response to N deposition was associated with decreases in lignolytic fungi and positively

associated with saprotrophic bacteria[24,27–29]. However, lignin, and its identifiable derivatives,

often comprises $\leq 10\%$ mineral SOM[19,30–33], meaning a substantial majority of SOM

biochemistry has been considered unaltered by N deposition. A higher resolution analysis of

SOM chemistry (i.e., at the level of individual compounds) and the soil microbiome through

decomposition and N deposition remains a critical gap in our knowledge on the persistence of

SOM that built up under anthropogenic N deposition.

By using both broad (compound classes) and more refined (individual compounds) SOM

biochemical and soil microbiome datasets, in the context of a long-term field study, the primary

objective of this study was to assess the effect of experimental N deposition on the trajectory of

organic matter decomposition and stabilization from undecayed plant root litter to mineral SOM.

We assessed the effect of experimental N deposition on SOM molecular abundance, diversity,

and complexity during microbial decay by establishing a decomposition gradient from

undecomposed fine roots to mineral SOM. Here, we define SOM molecular diversity analogous

to species richness and evenness, where molecular richness represents individual biochemical

compound counts and molecular evenness describes their relative abundance. We define SOM

molecular complexity as how structurally (dis)similar biochemical compounds that constitute

SOM are to one another[34]. More molecularly complex SOM would thus have fewer structural

similarities between its compounds. We hypothesized that experimental N deposition will elicit

changes in organic matter biochemistry at the individual compound-level that are not detected

when assessed at the less refined compound class-level, and these compound substitutions may

have functional implications for storage of SOM in the system. We further hypothesized that

changes in SOM biochemistry will be associated with changes in microbial community

composition that can explain the observed biochemical changes. Finally, we hypothesize that

individual compound substitutions result in altered soil organic matter biochemical diversity and

molecular complexity, potentially leading to long-term changes in SOM persistence.

2. Methods

2.1. Description of study sites

The Michigan Nitrogen Deposition Gradient Study was established in 1987 to examine the effects of atmospheric N deposition on ecosystem biogeochemical processes. Spanning a 500-km distance, the study area in Michigan, USA, consists of four replicate sugar maple (*Acer saccharum*) dominated northern hardwood forest stands on sandy spodosol soil (Typic Haplorthods of the Kalkaska series). The stands contain a thick organic horizon that includes a mat of fine roots (Oe/Oa) at its boundary with the A horizon. The stands are floristically and edaphically matched and encompass the full latitudinal range of northern hardwood forests in eastern North America and the Upper Great Lakes Region[35], allowing us to generalize our findings across this widespread ecologically and economically important ecosystem (Table S1). At each site ($n = 4$), there are six 30-m by 30-m plots, three of which received experimental N additions (30 kg NO_3^-–N ha^{-1} yr^{-1}) in the form of $NaNO_3$ pellets applied annually in six equal applications throughout the growing season beginning in 1994 and three of which receive ambient atmospheric N deposition (Figure 1).

Figure 1. The distribution of study sites in Michigan, USA that span the geographic range of northern hardwood forests. At each site ($n = 4$), there are six 30 m x 30 m plots, three of which received annual experimental nitrogen (N) additions in the form of 30 kg NO_3^-N ha^{-1} yr^{-1} beginning in 1994 and three of which receive ambient atmospheric N deposition (i). Maps created using ArcGIS Pro: release 3.3.1[36]. Cores were collected ($n = 3$ per plot) to yield the undecomposed and mineral SOM samples. Subsets of undecomposed material were deployed in litter bags and left to decompose for 1-year to yield the decomposed samples (ii). The biochemistry of all three sample types was assessed using pyrolysis-GC/MS and bacterial and fungal community composition was determined on the decomposed samples (iii).

2.2. Field-based experiment, sampling, and sample processing

This experiment was designed to produce samples that constitute a gradient of fine root decomposition, spanning undecomposed fine roots, 1-year decomposed fine roots, and more highly decomposed mineral soil organic matter. To do this, samples were collected in September and October 2013 by collecting soil cores to a depth of 10cm (5-cm-diameter), which included both Oe/Oa and A horizons[24,37]. Cores were transported on ice to the University of Michigan and stored at -20°C. Cores were thawed and passed through a 2mm sieve, and first- through third-order fine roots were collected and pooled at the plot level[38,39]. Roots were rinsed to remove mineral soil and dried at 60°C for 24 hours. The outcome of this processing yielded the "undecomposed" and "mineral SOM" sample fractions (Figure 1). For this purpose, roots from fresh Oe/Oa horizon samples are considered undecomposed in that relative to the other sample types in this experiment, they have undergone decomposition to a lesser degree as indicated by a light coloration and lack of observable decay. Organic material in the mineral soil represents organic matter that has undergone the greatest degree of decomposition and exists as a relatively stable form of SOM compared to organic horizon material[40].

To obtain the intermediate, "decomposed" samples, a subset of the undecomposed fine roots from each plot were sterilized and placed in mesh litter bags, deployed in the field, and collected after 12 months. Prior to field incubation, fine roots were sterilized to remove soil microbes using ethylene oxide[41], a process that minimally alters organic matter biochemistry[42], as compared to other methods such as autoclaving[43]. Sterilized fine roots (~2g dry weight) were placed into 15x15cm mesh litter bags constructed with 300 µm polyester mesh on top and 20µm polyester mesh on bottom to allow for microfauna and fungal hyphae to enter the bags[37,44]. Litter bags were deployed in their respective plots to their original location in the soil profile at the

boundary of the Oe/Oa and A horizons ($n = 3$ litter bags/plot at 6 plots/site; $n = 72$ litter bags

total). Litter bags were placed in the field in June 2014 and collected in June 2015, immediately

storing each bag on ice. The contents of each litter bag were homogenized by hand and

composited at the plot level. The decomposed fine root samples were subsampled for

biochemical analyses, dried at 60°C for 24 hours, and the remaining decomposed fine root

material was stored at -80°C for microbial community analyses. Supporting justification and

additional details on sample collection and processing can be found in Argiroff et al. (2019)[24].

2.3. Organic matter biochemistry

Subsets of each of the three sample types (undecomposed fine roots, decomposed fine

roots, and mineral soil; herein referred to as three different stages of decomposition) were ground

with a ball mill and assessed for organic matter biochemistry using pyrolysis-GC/MS[45]. Samples

were pyrolyzed at 600°C in quartz tubes for 20 seconds using a DS Pyroprobe 5150 pyrolyzer

and analyzed using a ThermoTrace GC Ultra gas chromatograph (Thermo Fisher Scientific,

Walthan, MA, USA) and ITQ 900 mass spectrometer (Thermo Fisher Scientific)[24,46] at the

University of New Hampshire. Compound peaks were identified and assigned using the

Automated Mass Spectral Deconvolution and Identification Systems (AMDIS v2.65) software,

the National Institute of Standards and Technology (NIST) compounds library, and published

literature[13,45]. Organic matter compounds are expressed as the % relative abundance of total

sample peak area and classified at the individual compound-level and grouped by compound

source origin (e.g., lignin-derivatives, phenolics, aromatics, proteins, N-bearing, polysaccharides,

and lipids). Critically, individual compounds observed in pyrolysis-GC/MS analysis are

pyrolysis products of a source compound and are classified according to the nature of their

source compound. Identifiable compounds with potentially multiple or unknown origins were

classified as unspecified[13]. In total, this effort yielded 260 identifiable compounds, each assigned

to one of the 8 compound source classes described above (Figure 1). In this study, we refer to

analyses of individual pyrolysis-GC/MS compounds as "high-resolution" though we emphasize

that, in this context, high-resolution refers strictly to the increased molecular resolution and

biochemical information provided at the individual compound level as compared to at the

broader compound class level. High-resolution analysis in this context is therefore not referring

to the depth or analytical capabilities of pyrolysis-GC/MS itself relative to other high- and ultra-

high resolution mass spectrometry techniques, such as FTICR-MS, used in elucidating soil

organic matter biochemistry.

2.4. Fungal and bacterial community composition

Fungal and bacterial community composition was determined from the remaining subset of

decomposed root litter and characterized using ribosomal DNA sequence abundances.

Supporting justification and additional details on DNA isolation, PCR amplification, amplicon

sequencing, and sequence quality control can be found in Argiroff et al. (2019)[24]. Briefly, total

genomic DNA from three replicate subsamples taken from each litter bag was isolated using the

DNeasy Plant Mini Kit (Qiagen, Germantown, MD) and checked for quality with a NanoDrop

8000 Spectrophotometer (Thermo Scientific) and gel electrophoresis. Replicate extractions from

each litter bag were pooled and stored at -80°C prior to PCR amplification. PCR amplification

was performed using the LROR and LR3 primers[47] that target the D1-D2 region of the 28S

rRNA gene. The V1-V3 regions of the bacterial 16S rRNA gene were targeted using the primers

27f and 519r[48]. PCR products were purified using the MinElute PCR Purification Kit (Qiagen)

and the quality of the purified PCR products was assessed as described above. Sequencing was

performed at the University of Michigan DNA Sequencing Core on 16 SMRT chips with a

PacBio RS II system (Pacific Biosciences, Menlo Park, CA). Mean amplicon lengths were 688

bp for fungal 28S and 525 bp for bacterial 16S rRNA genes. Sequences were processed using

mothur v1.40.5[49]. Fungal sequences were aligned against a 28S reference alignment from the

RDP LSU training set[50] and bacterial sequences were aligned against the 16S SILVA v132

reference alignment[51]. Sequences were clustered into operational taxonomic units (OTUs) for

fungal and bacterial sequences at 99% and 97% sequence similarity, respectively. The most

abundant sequence for each OTU was used as the representative for that OTU, and taxonomic

assignments were made using the RDP classifier with the LSU training set v11 for fungi[52] and

the SILVA v132 reference alignment with the naïve Bayesian classifier[53] in mothur for bacteria.

Raw sequences are available in GenBank under the accession numbers SRR8591550 (16S) and

SRR8591551 (28S).

2.5. Statistical Analyses

To assess the effect of experimental N deposition on the trajectory of organic matter

biochemistry during decay, we assessed compositional changes elicited by the treatment in a

three-step approach at two increasing levels of biochemical resolution. At the least resolved

level, herein referred to as the compound class-level, we used two-way analysis of variance

(ANOVA) to test the effect of experimental N deposition, site, and their interaction on $\log_2$-

transformed compound class ($n = 8$) abundances for each of the three decomposition stages. At

the higher resolution level of analysis, herein referred to as the individual compound-level, we

performed PERMANOVA on Euclidean distance matrices of $\log_2$-transformed relative

abundances of individual compounds at the three decomposition stages testing for the effect of

decomposition stage, treatment, site, and their interactions on organic matter biochemistry. In the

case of a significant PERMANOVA outcome, we performed PERMDISP[54] to determine whether

this effect was driven by differences in composition, heterogenous variance (i.e., dispersion), or

both. The third step in our approach, at the individual compound-level, we additionally

performed differential abundance analysis (Wald test)[55,56] to test for the effect of N deposition on

individual compound abundances in decomposition stages with a significant PERMANOVA

treatment effect.

To assess how N deposition altered microbial community composition, we tested the

effects of experimental N deposition, site, and their interaction on bacterial, fungal, and

saprotroph community composition (as defined using FUNGuild in R)[57] using two-way

PERMANOVA and Bray-Curtis dissimilarity matrices calculated from Hellinger-transformed

bacterial phylum, fungal genus, and saprotrophic family abundances. We performed PERMDISP

to evaluate whether experimental N deposition (i.e., treatment) effects detected by

PERMANOVA were driven by true differences in community composition. To determine the

taxa that most differed in relative abundance across experimental N deposition, differential

abundance analysis was performed on the bacterial and fungal communities using DESeq2 with

P-value adjustment for multiple comparisons using the false discovery rate method[58,59]. To

assess how N deposition altered associations between microbial communities and organic matter

biochemistry, we performed Mantel tests on $\log_2$-transformed Euclidean distances of organic

matter biochemistry and Hellinger-transformed Bray-Curtis distances of bacterial, fungal, and

saprotroph communities. To determine whether changes in organic matter biochemistry were

associated with changes in microbial community composition elicited by experimental N

deposition, distance-based redundancy analysis (db-RDA) was performed.

We measured organic matter molecular diversity of each sample ($n = 24$ per

decomposition stage; 72 total) by calculating molecular species (i.e., individual compounds)

richness and evenness using the vegan package (v2.6.4)[60] in R and performed two-way ANOVA

to test the effect of decomposition stage, N deposition, and their interaction on each molecular

diversity metric. Molecular richness represents individual compound counts and molecular

evenness was determined by calculating Pielou's Eveness Index[61], which measures how equally

individual compound abundances are distributed within a sample. When applicable, pairwise

comparisons were made using Tukey's honest significant difference test (HSD) to control for

multiple comparisons. In contrast to molecular diversity, which assess the number of individual

compounds present (i.e., richness) and their distributions within a sample (i.e., evenness), we

define molecular complexity as a measure of how complex each individual compounds' physical

molecular structure is. To determine this, we calculated molecular complexity through two

approaches. First, using the chemodiv package (v0.3.0)[34] in R, we used molecular fingerprints

(PubChem Fingerprint) to quantify compound dissimilarities based on the structural properties of

the molecules[34]. The fingerprints were then used to calculate Jaccard dissimilarities between

compounds as a measure of their structural dissimilarity[34]. We created molecular structural

(dis)similarity networks to visually represent changes in organic matter molecular complexity

through decomposition and between the ambient and experimental N deposition treatment. Using

this method of visualization, more complex SOM would thus have fewer structural similarities

between compounds and depict a less connected network. In addition to the molecular structural

(dis)similarity networks, we computed molecular complexity values for each individual

compound using the Bertz/Hendrickson/Ihlenfeldt formula[62,63] computed by Cactvs 3.4.8.18

(PubChem release 2021.10.14)[64] and performed ANOVA to test the effect of N deposition on

molecular complexity. These values are estimates of how complicated an individual compound's

molecular structure is in terms of both the elements contained and structural symmetry and

provide statistical significance to support network visualizations.

Due to the broad geographic expanse of our long-term field experiment and inherent

heterogeneity of the soil environment, we accepted statistical significance at $\alpha = 0.1$[24,27,29]. All

data processing, statistical analyses, and visualizations were performed and produced in R

(v4.2.2)[65] and RStudio (version 2022.12.0+353)[66], relying extensively on R packages such as

vegan (v2.6.4)[60], easyanova (v11.0)[67], and the tidyverse (v1.3.1)[68] packages.

3. Results

3.1 Compound class biochemistry

At the less-refined compound class-level, experimental N deposition significantly

increased the relative abundance of lignin-derived compounds in the mineral SOM (+53%, $P =$

0.09; Figure S1; Table S2) across all sites. However, the experimental N deposition did not affect

the abundance of any other compound class at any stage of decomposition (Table S3 and S4).

3.2 Individual compound biochemistry

When assessing the effect of experimental N deposition on the higher resolution,

individual compound relative abundances across the three decomposition stages, the composition

of individual compounds differed dependent on decomposition stage, site, and the N deposition

treatment (decomposition stage × site × treatment; F = 1.29, $P = 0.05$; Figure 2; Table S5; Figure

S2). Pairwise PERMANOVA at each decomposition stage revealed a significant effect of

experimental N deposition on mineral SOM biochemistry (F = 4.26, $P = 0.004$; Figure 2; Figure

S2), and no effect on the organic matter biochemistry at the undecomposed (F = 0.18, $P = 0.93$)

or 1-yr decomposed (F = 0.20, $P = 0.92$) stages of decomposition (Table S5; Figure S2).

Permutation tests of homogeneity of multivariate dispersion on the organic matter biochemistry

at each of the three decomposition stages were nonsignificant ($P > 0.1$; Table S6), indicating that

PERMANOVA detected differences in mineral SOM biochemical composition at the individual

compound-level between the ambient and experimental N deposition treatment, rather than

detecting changes in variability among samples (i.e., multivariate dispersion). Differential

abundance tests of individual compound abundances in mineral SOM revealed that experimental

N deposition led to the differential abundance of 19 out of 216 total compounds detected in the

mineral SOM (Figure 3). Of these differentially abundant compounds, 7 (3.2% of the total

number of compounds) were significantly more abundant in the N deposition treatment and 12

compounds (5.6% of total) were significantly more abundant in the ambient treatment (all $P \leq$

0.1; Table S7). Compounds more abundant under experimental N deposition compared to

ambient conditions included: phenol, 3-methyl- (aromatic combustion product), anthracene

(aromatic), 4-guaiacylpropane (lignin), benzeneacetic acid, 4-hydroxy-3-methoxy- (lignin),

benzene 1-methoxy-4-methyl- (lignin), n-decane (lipid), and 3-pyridinol (N-bearing).

Compounds more abundant in the ambient treatment included: benzene 1-ethenyl-4-methoxy-

(lignin), propene (lipid), isocrotonic acid (lipid), 3-methylpyridazine (N-bearing), ethanone, 1-(1-

methyl-1H-pyrrol-2-yl)- (N-bearing), 2,5-furandione, 3-methyl- (N-bearing), 2(5H)-furanone, 5-

methyl- (polysaccharide), 1,2-cyclohexanedione (polysaccharide), acetic anhydride

(polysaccharide), benzenepropanenitrile (protein), fluoranthene (unspecified), and (E)-1,3-

butadien-1-ol (unspecified).

Figure 2. First two axes of non-metric multidimensional scaling (NMDS) ordination of
Euclidean distances of $\log_2$ transformed py-GC/MS biochemical composition of individual
samples ($n = 72$, $k = 3$, stress = 0.10). Data points represent the biochemical profile of each
sample ($n = 6$ samples per site) and are colored by decomposition stage. The shape of the point
represents if it received the ambient or the 30 kg NO_3—N ha^{-1} yr^{-1} nitrogen deposition treatment.

Figure 3. MA (Minus vs. Average) plot of mineral SOM py-GC/MS $\log_2$ fold changes of

individual compound abundances between the ambient and nitrogen deposition treatments over

the mean of normalized counts, calculated as the average normalized compound abundances

taken across all samples. Compounds with a $P_{adj} < 0.1$ are colored by their compound source

class. Colored compounds above the 0 line (represented by the black dashed line) indicate

significantly higher abundance in the nitrogen deposition treatment, as compared to the ambient

treatment, and vice versa for below the 0 line. See Table S7 for significant compound test

statistics.

3.3 Microbial community composition

Experimental N deposition significantly altered bacterial (F = 3.20, P = 0.009), fungal (F = 1.90, P = 0.017), and fungal saprotrophic community composition (F = 2.77, P = 0.006; Table S8; Table S9). Permutation tests of homogeneity of multivariate dispersion on each community were nonsignificant (P > 0.1; Table S10), indicating that PERMANOVA detected differences in microbial community composition between the ambient and experimental N deposition treatment for bacterial, fungal, and saprotrophic communities, rather than detecting changes in variability among samples (i.e., multivariate dispersion). Specifically, using the Log_2 fold change approach in DeSeq2, the abundances of 28 bacterial OTUs (1.6% of total OTUs) were significantly higher under experimental N deposition as compared to the ambient treatment (Figure S3, Table S11). These OTUs included members of the phylum Bacteroidetes (39%), Proteobacteria (29%), Actinobacteria (14%), Patescibacteria (11%), and Armatimonadetes (7%). Under ambient conditions, 13 bacterial OTUs (0.73% of total OTUs) were present in significantly greater abundance. The phyla included Bacteroidetes (54%), Proteobacteria (38%), and Acidobacteria (8%). The abundances of 14 fungal OTUs (2.6% of total OTUs) were significantly higher under experimental N deposition as compared to the ambient treatment (Figure S3, Table S12). These OTUs included members of the fungal classes Tremellomycetes (42%), Eurotiomycetes (17%), Leotiomycetes (17%), Agaricomycetes (8%), Dothideomycetes (8%), and Sordariomycetes (8%). Under ambient conditions, 2 fungal OTUs (0.43% of total OTUs) were present in significantly greater abundance as compared to the N deposition treatment. The fungal classes include Agaricomycetes (50%) and Sordariomycetes (50%). However, the composition of bacterial (F = 1.70; P = 0.06) and fungal communities (F = 2.88; P < 0.001) differed among sites (Table S8; Table S9) and the effect of experimental N deposition on bacterial and fungal

363   communities was not uniform across sites (site × treatment $P \leq 0.1$; Table S8). Specifically,

364   experimental N deposition did not elicit significant differences in bacterial community

365   composition at site B (F = 2.65; $P = 0.20$) or fungal community composition at site D (F = 1.00;

366    $P = 0.30$; Table S9).

367 3.4 Microbial community associations with SOM biochemistry

Fungal and bacterial communities were differentially associated with organic matter

biochemistry at the three decomposition stages in the ambient and N deposition treatments

(Figure S4). In the ambient treatment, bacterial community composition was significantly

correlated with organic matter biochemistry in the undecomposed root litter (R = 0.29, $P = 0.07$)

and weakly associated in the mineral SOM (R = 0.26, $P = 0.12$) but exhibited no relationship

with the organic matter biochemistry in the 1-yr decomposed root litter (Table S13). Under

experimental N deposition, bacterial community composition and organic matter biochemistry

were not correlated at any of the three decomposition stages. In the ambient treatment, fungal

community composition was also not correlated with the 1-yr decomposed root litter (Table S13)

and was significantly correlated with the organic matter biochemistry in the mineral SOM (R =

0.20, $P = 0.09$). Under experimental N deposition, fungal community composition was

significantly correlated with the 1-yr decomposed root litter biochemistry (R = 0.23, $P = 0.04$)

and marginally associated in the mineral SOM (R = 0.17, $P = 0.11$) but exhibited no relationship

with the organic matter biochemistry in the undecomposed root litter (Figure S4; Table S13). In

the ambient treatment, saprotroph community composition was marginally associated with

organic matter biochemistry in the undecomposed root litter (R = 0.18, $P = 0.15$) and

significantly correlated with the mineral SOM (R = 0.30, $P = 0.06$) but exhibited no relationship

with the organic matter biochemistry in the 1-yr decomposed root litter (Figure S4; Table S13).

Under experimental N deposition, saprotroph community composition and organic matter

biochemistry were not correlated at any of the three decomposition stages.

To directly link changes in organic matter biochemistry with changes in microbial

community composition elicited by experimental N deposition, we fit vectors for mineral SOM

individual compound abundances to bacterial ($R^2 = 0.27$; $P = 0.001$), fungal ($R^2 = 0.10$; $P =$

0.09), and saprotrophic ($R^2 = 0.27$; $P = 0.001$) distance-based redundancy analysis (db-RDA)

ordinations (Figure 4). Mineral SOM individual compound abundances were selected given the

observed differences in organic matter biochemistry in the mineral SOM in response to

experimental N deposition (Figure 2; Table S5; Figure 3; Table S7). The shift in bacterial

community composition driven by experimental N deposition (F = 3.20, $P = 0.009$) was

associated with shifts in the abundance of 28 compounds in mineral SOM (Figure 4; Table S14).

The three most represented compound classes include lignin ($n = 10$; 35.7%), unspecified ($n = 7$;

25.0%), and lipids ($n = 3$; 10.7%). The longest vectors include Pyrazolo[5,1-

c][1,2,4]benzotriazin-8-ol (N-bearing; vector length = 0.698), Phenol, 2,6-dimethoxy-4-(2-

propenyl)- (Methoxyeugenol) (lignin; length = 0.649), Benzene, 1,4-dimethoxy- (phenol; length

= 0.604), and Benzeneacetic acid, 4-hydroxy-3-methoxy- (Homovanillic acid) (lignin; length =

0.599). The shift in fungal community composition driven by experimental N deposition (F =

1.90, $P = 0.017$) was associated with shifts in the abundance of 42 compounds in mineral SOM

(Figure 4; Table S15). The three most represented compound classes include lignin ($n = 10$;

23.8%), unspecified ($n = 9$; 21.4%), and lipids ($n = 8$; 19.1%). The longest vectors include

Phenol, 2,6-dimethoxy-4-(2-propenyl)- (Methoxyeugenol) (lignin; length = 1.370), Ethanone, 1-

(3,4-dimethoxyphenyl)- (lignin; length = 1.268), Benzoic acid, 4-hydroxy-3,5-dimethoxy-,

hydrazide (unspecified; length = 1.262), and Ethanone, 1-(4-hydroxy-3,5-dimethoxyphenyl)-

(Acetosyringone) (lignin; length = 1.186). The shift in saprotrophic fungal community

composition under experimental N deposition (F = 2.77, P = 0.006) was associated with shifts in

the abundance of 21 compounds in mineral SOM (Figure 4; Table S16). The three most

represented compound classes include lignin (n = 7; 33.3%), unspecified (n = 7; 33.3%), and N-

bearing (n = 3; 14.3%). The longest vectors include Trimethylphenol (unspecified; length =

1.370), Ethanone, 1-(4-hydroxy-3,5-dimethoxyphenyl)- (Acetosyringone) (lignin; length =

1.360), n-Decane (lipid; 1.350), and Benzoic acid, 4-hydroxy-3,5-dimethoxy-, hydrazide

(unspecified; 1.350).

Figure 4. Biplots of distance-based redundancy analysis (db-RDA) ordinations of bacterial

phyla, fungal genera, and saprotrophic families (points) and SOM compounds (vectors). The db-

RDA was performed on Bray-Curtis dissimilarity matrices calculated from Hellinger-

transformed taxon abundances. Ordinations were constrained by experimental N deposition and

site, which together accounted for 31.8% of variation in Bray-Curtis dissimilarity for bacteria,

38.1% for fungi, and 33.3% for saprotrophs. Points represent individual samples (n = 6 samples

per site) and are colored by site. Point shape indicates nitrogen treatment (ambient or 30 kg

NO_3—N ha^{-1} yr^{-1} deposition). Treatment centroids are shown in black as "A" (ambient) or "N"

(nitrogen deposition). Vectors represent mineral SOM compounds fitted to the ordination.

Compounds with goodness of fit P-values $\leq$ 0.1 are displayed and are colored by their source

compound class.

3.5 Biochemical diversity and complexity

Experimental N deposition did not affect molecular species richness (F = 0, P = 0.97) or

evenness (F = 0.39, P = 0.53) at any of the three decomposition stages (Figure 5; Table S17).

Given the observed treatment differences in biochemical composition in the mineral SOM only,

we assessed the effect of experimental N deposition on molecular complexity at only the mineral

SOM decomposition stage. To determine how experimental N deposition affects molecular

complexity (i.e., how structurally similar compounds that constitute SOM are to one another),

compound co-occurrence and structural dissimilarity matrices were generated for mineral SOM

biochemistry in both treatments. Under ambient conditions in the mineral SOM, molecular

network analysis revealed 197 nodes, representing compounds, and 2450 edges, representing the

structural similarities present between those compounds (Figure 6). Molecular network analysis

of the mineral SOM biochemistry in experimental N deposition revealed it did not significantly

alter the number of compounds present (n_{nodes} = 187) nor the structural similarities between

compounds (n_{edges} = 2502; Figure 6). Molecular complexity values calculated using the

Bertz/Hendrickson/Ihlenfeldt formula affirmed that the N deposition treatment did not

significantly alter individual compound molecular complexity in the mineral SOM (F = 0, P =

0.997; Figure S5).

Decomposition stage had a significant effect on molecular richness (F = 9.49, P<0.001)

and evenness (F = 12.22, P<0.001; Figure 5; Table S17). HSD-adjusted pairwise comparisons

revealed significantly higher molecular richness at the 1-yr decomposed stage compared to the

undecomposed (post hoc P = 0.001) and mineral SOM (post hoc P<0.001; Figure 5; Table S18).

Molecular evenness was significantly higher in the 1-yr decomposed (post hoc P = 0.002) and

mineral SOM (post hoc P<0.001) stages of decomposition as compared to the undecomposed

root litter (Figure 5; Table S18). There was no difference in molecular evenness between the 1-yr

decomposed and mineral SOM decomposition stages (post hoc $P = 0.39$; Figure 5; Table S18).

In the undecomposed root litter, molecular network analysis revealed 209 nodes and 2095 edges,

while the 1-yr decomposed root litter network had 225 nodes and 3372 edges, and the mineral

SOM network had 216 nodes and 3022 edges (Figure S6). Molecular complexity values

calculated using the Bertz/Hendrickson/Ihlenfeldt formula affirmed that decomposition stage did

not significantly alter individual compound molecular complexity at any point (F = 0.01, $P =$

0.99, Figure S7).

Figure 5. The effect of the nitrogen deposition treatment and decomposition stage on molecular richness (left) and evenness (right). Within each figure, ambient values are displayed on the left (solid) and nitrogen deposition treatment values are displayed on the right (dashed) for each decomposition stage ($n = 3$). Individual sample values ($n = 72$) are displayed. Two-way ANOVA was used to test the effect of Decomposition Stage, Treatment, and Decomposition Stage x Treatment interaction. The effect of Decomposition Stage was significant for both richness ($F_{decomp} = 9.49$, $P < 0.001$) and evenness ($F_{decomp} = 12.22$, $P < 0.001$). Decomposition Stage pairwise comparisons (Tukey's HSD) are displayed as letters (see Supplementary Table S18).

Figure 6. Molecular networks showing relationships between the biochemical compounds

present in the mineral SOM under ambient (left) and nitrogen deposition (right) conditions.

Nodes represent individual molecules (ambient: $n = 197$; nitrogen treatment: $n = 189$) and edges

represent the quantity of interactions and relationships between those molecules (ambient: $n =$

2450; nitrogen treatment: $n = 2502$). Thickness of the edges was determined by the molecular

similarity with a cut off value of 0.7. Nodes are colored by the compound class to which that

individual molecule belongs. Enlarged nodes represent the molecules in greater abundance in

each respective treatment derived from differential abundance analysis (Figure 3; Table S7).

4. Discussion

High resolution molecular analyses reveal N deposition elicits changes to SOM biochemistry that were previously undetectable, supporting our first hypothesis. It is common to assess SOM biochemistry by grouping individual biochemical compounds into broader classes, such as lipids or lignin-derived compounds, as a means to simplify biochemical heterogeneity and diversity by grouping compounds that are suspected to behave similarly based on the relative similarity of their structure or other physical properties or factors[12,13,24,69,70]. In doing so, the abundances of individual compounds within a compound class are combined, leading to a loss in the potential resolution of SOM biochemical information, and potentially making impacts of environmental change on SOM biochemical constituents undetectable. As the resolution at which we can examine SOM biochemistry has increased with technological advances, so has the importance of identifying and elucidating the complexity within these traditional compound classes. While it has been common to assume that individual compounds within a class of compounds behave uniformly, there is increasing evidence for contrasting and compounding origins within a compound class and the different potential pathways of mineralization will have implications for the formation and persistence of mineral SOM[2,7,19,20]. As such, the implementation of more refined analytical analyses (i.e., high[er] resolution) that aid in elucidating these molecular-level differences can increase understanding of mechanisms governing SOM persistence; a necessary step toward increasing organic matter in soil as a viable climate change mitigation strategy[15,20].

To demonstrate this assertion, previous research efforts in this long-term field experiment have detected increases in lignin-derived compounds in mineral SOM under experimental N deposition, with no detectable changes to other components of mineral SOM biochemistry[24,25].

However, high resolution analysis used in this study revealed numerous individual compounds

changed in mineral SOM within multiple additional compound classes in response to

experimental N deposition that may have consequences for the processes of organic matter decay

and soil C storage (Figure 3; Table S7). For example, the abundance of lipids in mineral SOM

exhibited no effect of experimental N deposition at the compound class-level (Figure S1; Table

S2). However, at the individual compound-level, in the ambient treatment, two molecules that

are produced from the combustion of lipids (propene and isocrotonic acid) were 15-times and

over 200-times more abundant, respectively, compared to the experimental N deposition

treatment (Figure 3; Table S7). Whereas under experimental N deposition, one lipid molecule (n-

decane) was over 100-times more abundant compared to in the ambient treatment (Figure 3;

Table S7) and was one of the most significant compounds associated with the shift in

saprotrophic community composition driven by experimental N deposition (Figure 4; Table

S16). This reorganization of individual compounds within compound classes is undetectable by

less refined analysis, ultimately leading to no observable effect of experimental N deposition on

the abundance of lipids as a class of compounds (Figure S1; Table S2), but is critical for

understanding the fate of these substrates in SOM. Lipids are a highly heterogenous class of

compounds with significant input from both plants and soil microbes[71,72]. Though there is rising

evidence that stable mineral SOM is largely microbial-derived, it remains unclear and highly

context dependent how much is plant-derived[13,19,20,73–76]. Thus, it is likely that observed changes

in the microbial community can account for a portion of the variance observed in SOM lipid

biochemistry, supporting our second hypothesis. Microbial taxa that are relatively more abundant

under experimental N deposition and exhibit fast growing (i.e., copiotrophic) traits may

contribute to increased lipid signatures in mineral SOM through enhanced turnover, resulting in

necromass production. In this study, members of the phyla Bacteroidetes and Acidobacteria, and

the genus *Paraburkholderia* were in significantly greater abundance under experimental N

deposition compared to the ambient treatment (Figure S3), all of which harbor taxa that exhibit

copiotrophic characteristics[77,78]. However, estimations of microbial necromass C in forest soils

suggest that a substantial portion (~70%) of C in persistent SOM could be plant-derived[79,80], and

that the contribution of plant-derived lipids to this C pool can be greater than that of lignin[20]. All

three of the differentially abundant lipid molecules across the ambient and experimental N

deposition treatments are involved in both plant and microbial metabolic processes[81–83]. As such,

more detailed knowledge of the specific metabolic reactions, which plants or microbes are

producing them, and how these compounds end up in mineral SOM is necessary to understand

the longevity of these substrates in the system[84].

Proteins and other N-containing compounds exhibited similar trends at these differing

analytical scales. Despite no statistical difference in the abundance of proteins at the compound

class-level, at the individual compound-level one protein molecule (benzenepropanenitrile) was

almost 8-times more abundant in the ambient treatment compared under experimental N

deposition (Figure 3; Table S7). The hydrolysis of benzenepropanenitrile (i.e., 3-

phenylpropionitrile) is catalyzed by the nitrilase enzyme (EC 3.5.5.1)[85–88], yielding a carboxylic

acid and ammonium (NH_4^+). The gene which encodes the enzyme that ultimately carries out this

reaction is found in the genomes of two soil bacteria, one of which being *Paraburkholderia*

phymatum[86,89–92]. In this study, *Paraburkholderia spp.* ($n = 3$; Table S11) were in significantly

greater abundance under experimental N deposition compared to the ambient treatment (Figure

S3). *Paraburkholderia spp.* are known decomposers in forests, particularly sugar maple forests

like the ones in this study[93]. Thus, it is plausible that increased enzymatic activity produced by

Paraburkholderia spp. under experimental N deposition catalyzed this reaction, leading to

significantly lower benzenepropanenitrile abundance in the N deposition treatment. These

findings suggest that changes to microbial community composition in response to environmental

stimuli may have functional consequences and elicit changes to the biochemical composition of

organic matter that eventually becomes stable mineral SOM.

In addition to the novel findings detailed above, high resolution analysis of SOM

biochemistry revealed additional compositional insight into lignin dynamics in response to the

experimental N deposition that may underlie resistance to microbial decomposition, further

supporting our first hypothesis. Previous work in this long-term experiment showed that

anthropogenic N deposition slowed the rate and extent of organic matter decomposition,

ultimately resulting in the accumulation of lignified material in mineral SOM under experimental

N deposition (+53%, $P = 0.09$; Figure S1; Table S2)[24–26,29]. However, at the individual

compound-level, changes in the composition of lignin compounds in mineral SOM were detected

under both ambient and experimental N deposition conditions. In the ambient treatment, one

lignin-derived molecule (benzene, 1-ethenyl-4-methoxy-) was almost 8-times more abundant

compared to under experimental N deposition (Figure 3; Table S7). Whereas under experimental

N deposition, three different lignin-derived molecules (benzene, 1-methoxy-4-methyl-,

homovanillic acid, and 4-guaiacylpropane) were 8-times, 14-times, and 29-times more abundant

than in the ambient treatment, respectively, ultimately leading to the overall increase observed at

the compound class-level (Figure 3; Figure S1; Table S7).

At the relatively less-refined compound class-level, it is not possible to detect this

reorganization of different lignin molecules, as observed here, that ultimately affects the

interpretability of the stability of the compound class as a whole. Lignin is a complex

macromolecule that is highly variable in terms of its primary structure. Which lignin monomers

and functional groups are present, in what quantities, and what linkages connect them affect the

decomposability of lignin polymers by soil microbes[94–97]. For example, lignin polymers

containing higher quantities of monomers of the guaiacyl (G) subunit are more resistant to

decomposition, resulting in significantly lower mass loss, than lignin polymers composed of

other subunits[16,96]. The 29-times greater abundance of 4-guaiacylpropane and 14-times greater

abundance of homovanillic acid, a G subunit decomposition product[98], under experimental N

deposition suggests that the SOM that has accumulated under anthropogenic N deposition is

particularly compositionally resistant to microbial decomposition in terms of its lignin content.

Thus, it is plausible that in this system, differences in the quality of lignin, in addition to the

increases in lignin quantity[24], may be a factor contributing to the increased SOM observed under

high N deposition conditions. This possibility has been suggested from earlier modeling studies

in this system[26], and this is the first experimental observation to support this modeled result.

Differences in the composition of decomposing microbial communities observed under

experimental N deposition support the claim that lignin in mineral SOM under increased

anthropogenic N deposition is particularly resistant to decomposition or is simply not being

metabolized by the microbial community in this treatment, further supporting our second

hypothesis. While fungi are widely considered as primary lignin decomposers, evidence has

shown that bacteria may play a non-inconsequential role in lignin decay[99,100]. Of the bacterial

OTUs in higher abundance under N deposition conditions, just under half (43%) belong to the

phyla Proteobacteria and Actinobacteria (Figure S3; Table S11). This finding aligns with

previous work in this system that observed an increase in the abundance of bacteria associated

with lignin decay under experimental N deposition[27,28]. The majority of lignin-degrading bacteria

identified to date represent three bacterial classes, all of which are within the Proteobacteria and

Actinobacteria phyla: Actinomycetes, α-Proteobacteria, and γ-Proteobacteria[101–103]. Such

bacteria capable of lignin degradation do so in an incomplete manner, such that smaller aromatic

lignin-derived compounds are produced[104–106]. For example, bacterial species belonging to the

class γ-Proteobacteria produce the intermediate phenolic compound homovanillic acid during

plant litter and lignin decomposition[107]. In this study, homovanillic acid was one of the most

significant compounds associated with the shift in bacterial community composition driven by

experimental N deposition (Figure 4; Table S14) and was in significantly higher abundance in

the N deposition treatment compared to the ambient treatment (Figure 3; Table S7).

Additionally, benzoic acid, 4-hydroxy-3,5-dimethoxy-, hydrazide, a decomposition derivative of

syringic acid, which is a key intermediate of bacterial catabolism of lignin compounds[108], was

one of the most significant compounds associated with the shifts in fungal and saprotrophic

community composition driven by experimental N deposition (Figure 4; Table S15; Table S16).

Taken together, it is plausible that the accumulation of decomposition resistant lignin material

under experimental N deposition was due, in part, to the incomplete bacterial metabolism of

lignin and the inability to break down these compounds specifically.

Further, both bacterial and saprotrophic fungal communities exhibit strong relationships

with the biochemistry of the organic matter in the undecomposed litter and mineral SOM under

ambient conditions (Figure S4; Table S13). Yet, under experimental N deposition these

relationships are lost. The expression of microbial genes and the subsequent production of

extracellular enzymes that govern the breakdown of SOM are highly sensitive to N

additions[29,109–112]. Under high N conditions, genes responsible for the production of extracellular

enzymes that break down complex macromolecules, such as lignin, are downregulated[29].

Additional studies have observed a shift in microbial community composition under N

deposition conditions towards a community whose primary niche is the decay of simpler

compounds, such as cellulose[109,113,114]. Taken together, it is plausible that when N is readily

available, microbes do not need to expend the energy producing extracellular enzymes to break

down organic material and this decreased activity may result in a shift in the community

composition. The lack of microbial activity observed in previous studies, in consideration with

SOM biochemical relationships to microbial taxa observed here, may explain why we observed a

decoupling of SOM biochemistry and microbial community composition under experimental N

deposition.

Despite the above-described differences in individual compound biochemical

composition being distinct under N deposition conditions in the most highly decomposed mineral

SOM (Figure 3; Table S7), the biochemical diversity and molecular complexity of mineral SOM

as a whole was not altered by N deposition (Figure 5; Figure 6), contradicting our third

hypothesis. However, SOM biochemical diversity (i.e., molecular richness and evenness) did

change throughout decomposition (Figure 5). Both molecular richness and evenness were higher

after one year of decomposition compared to initial undecomposed inputs (Figure 5; Table S17;

Table S18). This is likely the result of the production of microbially-derived compounds during

the initial phases of root decay. When microbes break down plant litter, new byproducts are

produced[7,13,115], which likely explains the observed increases in molecular richness, i.e., the

number of compounds present, after one year of decomposition. Molecular richness of mineral

SOM returning to the same levels as initial inputs may indicate the complete decay of more

accessible components of litter or that the microbially processed C substrates that were produced

during one year of decomposition were similarly consumed, returning to the same number of

compounds as initial inputs, though the composition of those compounds are distinct (Figure 2;

Figure 5). Despite differences in biochemical diversity through decomposition, we did not detect

differences in the molecular complexity of SOM between the undecomposed root litter, 1-year

decomposed root litter, and mineral SOM (Figure S6; Figure S7). Ultimately these findings

indicate that although the composition of the individual compounds changed during decay

(Figure 2; Table S5), and were ultimately distinct under N deposition conditions in the most

highly decomposed mineral SOM (Figure 2; Figure 3; Table S7), the molecular complexity (i.e.,

structural [dis]similarity) of the compounds that constitute SOM at any given stage of

decomposition did not (Figure S4). This finding differs from both what has been proposed

conceptually about organic matter complexity underlying SOM persistence[15] and from recent

studies that have found the opposite: that persistent SOM is less complex than its initial

inputs[12,21,116]. In similar studies, SOM persistence was estimated across wider ecosystem ranges

(e.g., across soil depths or elevational gradients that span different ecosystem types[12,21,116]) than

in this study, which utilized a long-term experiment in sugar maple dominated forests. Thus, in

this system, the lack of relationship between SOM molecular complexity and persistence may be

attributable to relatively similar site-specific edaphic factors and vegetation composition, which

may not be variable enough to detect differences in SOM decay using this method that detects

large-scale molecular changes (e.g., terpenes vs. benzenoids). Taken together, our findings

suggest that SOM biochemical diversity and molecular complexity are not drivers of SOM

persistence in this system.

Collectively, our results demonstrate that N deposition altered the abundance of specific

compounds in mineral SOM, a reorganization of SOM biochemistry that is not detectable when

assessing changes at the less-refined compound class-level, as well as decoupled the relationship

between microbial composition and SOM biochemistry. Whether these observed changes have

functional implications for the long-term storage of SOM is unknown and warrants further

investigation. Though these molecular-level differences in SOM biochemistry under

anthropogenic N deposition may appear subtle, the effect that these differences have on the C

stored in the system long-term may be disproportionately large. For example, in this system,

upon termination of the experimental N deposition, a globally ecologically relevant

environmental change in response to Clean Air Act Amendments, microbial activity significantly

increased and the C that built up under experimental N deposition was lost from the system

within 5-years of treatment termination[40]. Studies have shown that molecular-level changes in

SOM biochemistry, in response to various environmental change conditions, impact the capacity

of temperate forests to sequester C[117]. A microbial mechanism may underlie this possibility as

differences in the structural properties of the molecules in greater abundance under

anthropogenic N deposition may have implications for how and by whom these products are

transformed, what products are produced, and, ultimately, the decomposability of their

byproducts[83,118,119]. Taken together, it is plausible that the SOM that accumulated under

experimental N deposition was molecularly distinct as a result of functional changes in the

microbial community elicited by the N deposition treatment[24,29]. Thus, upon termination of

experimental N deposition, the accumulated, molecularly distinct SOM may have been readily

decomposed by the active microbial community upon return to low (i.e., ambient) N conditions.

While empirically untested, this response is plausible due to the relative increase in lignin-like

compounds, yet observed decrease in the expression of genes that encode lignolytic enzymes[29],

as previously observed under experimental N deposition. As such, lignolytic fungal gene

expression may rapidly increase in response to reduced N deposition, supporting a rapid decay of

the relatively lignin-rich SOM that accumulated during the active experimental N deposition

treatment in this long-term study. Future work should explore the relative decomposition

potential of different lignin- and lipid-related compounds that were differentially abundant under

ambient or the experimental N deposition treatment to deepen our understanding of the potential

ecosystem-level consequences of the reorganization of SOM chemistry observed here.

Acknowledgements

We thank Annalisa Stevenson for her help in ArcGIS. We also wish to thank Thea

Whitman (and lab), Erika Marín-Spiotta (and lab), and the Department of Soil Science graduate

students for their thoughtful feedback on this work.

Funding Declaration

This material is based upon work supported by the National Science Foundation Graduate

Research Fellowship Program under Grant No. DGE-2137424 awarded to B.E.P. and from the

United States Department of Energy Biological Environmental Research Program. The

maintenance of our long-term field experiment is supported by the National Science Foundation

Long-Term Research in Environmental Biology Program.

Author Contributions Statement

D.R.Z. designed the study. R.A.U. and W.A.A. performed field and laboratory analyses.

A.S.G. performed biochemical analyses. B.E.P. analyzed the data with the help of G.A.C and

Z.B.F. B.E.P wrote the first draft of the manuscript. D.R.Z., W.A.A., G.A.C., A.S.G., and Z.B.F.

provided significant feedback on subsequent versions and analyses.

References

1. Bowden, R. D. *et al.* Litter Input Controls on Soil Carbon in a Temperate Deciduous Forest.

Soil Sci. Soc. Am. J. **78**, (2014).

2. Cotrufo, M. F. & Lavallee, J. M. Soil organic matter formation, persistence, and functioning:

A synthesis of current understanding to inform its conservation and regeneration. in *Advances*

in Agronomy vol. 172 1–66 (Elsevier, 2022).

3. Jenkinson, D. S. & Rayner, J. H. The Turnover of Soil Organic Matter in Some of the

Rothamsted Classical Experiments: *Soil Sci.* **123**, 298–305 (1977).

4. Parton, W. J., Schimel, D. S., Cole, C. V. & Ojima, D. S. Analysis of Factors Controlling Soil

Organic Matter Levels in Great Plains Grasslands. *Soil Sci. Soc. Am. J.* **51**, 1173–1179 (1987).

5. Adair, E. C. *et al.* Simple three-pool model accurately describes patterns of long-term litter

decomposition in diverse climates. *Glob. Change Biol.* **14**, 2636–2660 (2008).

6. Berg, B. & Meentemeyer, V. Litter quality in a north European transect versus carbon storage

potential. *Plant Soil* **242**, 83–92 (2002).

7. Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Denef, K. & Paul, E. The Microbial

Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with

soil organic matter stabilization: do labile plant inputs form stable soil organic matter? *Glob.*

Change Biol. **19**, 988–995 (2013).

8. Prescott, C. E. Litter decomposition: what controls it and how can we alter it to sequester

more carbon in forest soils? *Biogeochemistry* **101**, 133–149 (2010).

9. Zhang, D., Hui, D., Luo, Y. & Zhou, G. Rates of litter decomposition in terrestrial

ecosystems: global patterns and controlling factors. *J. Plant Ecol.* **1**, 85–93 (2008).

10.   Berg, B. & McClaugherty, C. *Plant Litter*. (Springer Berlin Heidelberg, Berlin,

Heidelberg, 2008). doi:10.1007/978-3-540-74923-3.

11.   Waksman, S. A. Humus Origin, Chemical Composition, and Importance in Nature. *Soil*

Science **41**, (1936).

12.   Jones, A. R. *et al.* Molecular complexity and diversity of persistent soil organic matter.

Soil Biol. Biochem. **184**, 109061 (2023).

13.   Kallenbach, C. M., Frey, S. D. & Grandy, A. S. Direct evidence for microbial-derived

soil organic matter formation and its ecophysiological controls. *Nat. Commun.* 10 (2016).

14.   Lehmann, J. & Kleber, M. The contentious nature of soil organic matter. *Nature* **528**, 60–

68 (2015).

15.   Lehmann, J. *et al.* Persistence of soil organic carbon caused by functional complexity.

Nat. Geosci. **13**, 529–534 (2020).

16.   Talbot, J. M., Yelle, D. J., Nowick, J. & Treseder, K. K. Litter decay rates are determined

by lignin chemistry. *Biogeochemistry* **108**, 279–295 (2012).

17.   Wickings, K., Grandy, A. S., Reed, S. C. & Cleveland, C. C. The origin of litter chemical

complexity during decomposition. *Ecol. Lett.* **15**, 1180–1188 (2012).

18.   Sugai, S. F. & Schimel, J. P. Decomposition and biomass incorporation of 14c-labeled

glucose and phenolics in taiga forest floor: effect of substrate quality, successional state, and

season. *Soil Biol. Biochem.* **25**, 1379–1389 (1993).

19.   Whalen, E. D. *et al.* Clarifying the evidence for microbial- and plant-derived soil organic

matter, and the path toward a more quantitative understanding. *Glob. Change Biol.* gcb.16413

(2022) doi:10.1111/gcb.16413.

20. Angst, G., Mueller, K. E., Nierop, K. G. J. & Simpson, M. J. Plant- or microbial-derived?

A review on the molecular composition of stabilized soil organic matter. *Soil Biol. Biochem.*

156, 108189 (2021).

21. Davenport, R. *et al.* Decomposition decreases molecular diversity and ecosystem

similarity of soil organic matter. *Proc. Natl. Acad. Sci.* **120**, e2303335120 (2023).

22. Yang, S. *et al.* Soil organic carbon content and mineralization controlled by the

composition, origin and molecular diversity of organic matter: A study in tropical alpine

grasslands. *Soil Tillage Res.* **215**, 105203 (2022).

23. Tang, X., Chang, C. & Li, M. Pyrolysis gas chromatography mass spectrometry (py-

GC/MS) revealed independent effects of precipitation and temperature on diversity and

molecular composition of soil organic matter. *Environ. Earth Sci.* **84**, 164 (2025).

24. Argiroff, W. A., Zak, D. R., Upchurch, R. A., Salley, S. O. & Grandy, A. S.

Anthropogenic N deposition alters soil organic matter biochemistry and microbial

communities on decaying fine roots. *Glob. Change Biol.* **25**, 4369–4382 (2019).

25. Zak, D. R., Holmes, W. E., Burton, A. J., Pregitzer, K. S. & Talhelm, A. F. Simulated

Atmospheric NO_3^- Deposition Increases Soil Organic Matter By Slowing Decomposition.

Ecol. Appl. **18**, 2016–2027 (2008).

26. Whittinghill, K. A., Currie, W. S., Zak, D. R., Burton, A. J. & Pregitzer, K. S.

Anthropogenic N Deposition Increases Soil C Storage by Decreasing the Extent of Litter

Decay: Analysis of Field Observations with an Ecosystem Model. *Ecosystems* **15**, 450–461

(2012).

27.   Freedman, Z. & Zak, D. R. Atmospheric N Deposition Increases Bacterial Laccase-Like

768   Multicopper Oxidases: Implications for Organic Matter Decay. *Appl. Environ. Microbiol.* **80**,

769   4460–4468 (2014).

770   28.   Freedman, Z. B., Upchurch, R. A. & Zak, D. R. Microbial Potential for Ecosystem N

771   Loss Is Increased by Experimental N Deposition. *PLOS ONE* **11**, e0164531 (2016).

772   29.   Zak, D. R. *et al.* Anthropogenic N deposition, fungal gene expression, and an increasing

773   soil carbon sink in the Northern Hemisphere. *Ecology* **100**, (2019).

774   30.   Angst, G., Mueller, K. E., Kögel-Knabner, I., Freeman, K. H. & Mueller, C. W.

775   Aggregation controls the stability of lignin and lipids in clay-sized particulate and mineral

776   associated organic matter. *Biogeochemistry* **132**, 307–324 (2017).

777   31.   Bol, R., Poirier, N., Balesdent, J. & Gleixner, G. Molecular turnover time of soil organic

778   matter in particle-size fractions of an arable soil. *Rapid Commun. Mass Spectrom.* **23**, 2551–

779   2558 (2009).

780   32.   Geng, J. *et al.* Different Molecular Characterization of Soil Particulate Fractions under N

781   Deposition in a Subtropical Forest. *Forests* **10**, 914 (2019).

782   33.   Grandy, A. S. & Neff, J. C. Molecular C dynamics downstream: The biochemical

783   decomposition sequence and its impact on soil organic matter structure and function. *Sci.*

784   *Total Environ.* **404**, 297–307 (2008).

785   34.   Petrén, H., Köllner, T. G. & Junker, R. R. Quantifying chemodiversity considering

786   biochemical and structural properties of compounds with the R package CHEMODIV. *New*

787   *Phytol.* **237**, 2478–2492 (2023).

788   35.   Burton, A. J., Ramm, C. W., Pregitzer, K. S. & Reed, D. D. Use of multivariate methods

789   in forest research site selection. *Can. J. For. Res.* **21**, 1573–1580 (1991).

36.   Jon Dewitz. National Land Cover Database (NLCD) 2019 Products (ver. 3.0, February

2024). U.S. Geological Survey https://doi.org/10.5066/P9KZCM54 (2024).

37.   Xia, M., Talhelm, A. F. & Pregitzer, K. S. Long-Term Simulated Atmospheric Nitrogen

Deposition Alters Leaf and Fine Root Decomposition. *Ecosystems* **21**, 1–14 (2018).

38.   Pregitzer, K. S. *et al.* Fine Root Architecture of Nine North American Trees. *Ecol.*

Monogr. **72**, 293–309 (2002).

39.   Xia, M., Talhelm, A. F. & Pregitzer, K. S. Fine roots are the dominant source of

recalcitrant plant litter in sugar maple-dominated northern hardwood forests. *New Phytol.* **208**,

715–726 (2015).

40.   Propson, B. E., Zak, D. R., Classen, A. T., Burton, A. J. & Freedman, Z. B. Gains in soil

carbon storage under anthropogenic nitrogen deposition are rapidly lost following its

cessation. *Ecology* **105**, e4444 (2024).

41.   Rose, R. E. & Bailey, R. W. Ethylene Oxide for Soil Sterilization. *Nature* **169**, 716–716

(1952).

42.   Gennari, M., Negre, M. & Ambrosoli, R. Effects of ethylene oxide on soil microbial

content and some soil chemical characteristics. *Plant Soil* **102**, 197–200 (1987).

43.   Berns, A. E. *et al.* Effect of gamma-sterilization and autoclaving on soil organic matter

structure as studied by solid state NMR, UV and fluorescence spectroscopy. *Eur. J. Soil Sci.*

59, 540–550 (2008).

44.   Hobbie, S. E. Contrasting Effects of Substrate and Fertilizer Nitrogen on the Early Stages

of Litter Decomposition. *Ecosystems* **8**, 644–656 (2005).

45.   Grandy, A. S., Strickland, M. S., Lauber, C. L., Bradford, M. A. & Fierer, N. The

influence of microbial communities, management, and soil texture on soil organic matter

chemistry. *Geoderma* **150**, 278–286 (2009).

46.   Pold, G., Grandy, A. S., Melillo, J. M. & DeAngelis, K. M. Changes in substrate

availability drive carbon cycle response to chronic warming. *Soil Biol. Biochem.* **110**, 68–78

(2017).

47.   Vilgalys, R. & Hester, M. Rapid genetic identification and mapping of enzymatically

amplified ribosomal DNA from several Cryptococcus species. *J. Bacteriol.* **172**, 4238–4246

(1990).

48.   Lane, D. J. 16S/23S rRNA sequencing. *Nucleic acid techniques in bacterial systematics*

(1991).

49.   Schloss, P. D. *et al.* Introducing mothur: Open-Source, Platform-Independent,

Community-Supported Software for Describing and Comparing Microbial Communities.

Appl. Environ. Microbiol. **75**, 7537–7541 (2009).

50.   Mueller, R. C., Balasch, M. M. & Kuske, C. R. Contrasting soil fungal community

responses to experimental nitrogen addition using the large subunit ʀRNA taxonomic marker

and cellobiohydrolase I functional marker. *Mol. Ecol.* **23**, 4406–4417 (2014).

51.   Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2012).

52.   Cole, J. R. *et al.* Ribosomal Database Project: data and tools for high throughput rRNA

analysis. *Nucleic Acids Res.* **42**, D633–D642 (2014).

53.   Wang, Q., Garrity, G. M., Tiedje, J. M. & Cole, J. R. Naïve Bayesian Classifier for Rapid

Assignment of rRNA Sequences into the New Bacterial Taxonomy. *Appl. Environ. Microbiol.*

73, 5261–5267 (2007).

54.   Anderson, M. J. PERMANOVA_2factor: a FORTRAN computer program for

permutational multivariate analysis of variance (for any two-factor ANOVA design) using

permutation tests. *Department of Statistics, University of Auckland, Auckland* 35 (2004).

55.   Langley, S. R. & Mayr, M. Comparative analysis of statistical methods used for detecting

differential expression in label-free mass spectrometry proteomics. *J. Proteomics* **129**, 83–92

(2015).

56.   Yang, L. & Chen, J. A comprehensive evaluation of microbial differential abundance

analysis methods: current status and potential solutions. *Microbiome* **10**, 130 (2022).

57.   Nguyen, N. H. *et al.* FUNGuild: An open annotation tool for parsing fungal community

datasets by ecological guild. *Fungal Ecol.* **20**, 241–248 (2016).

58.   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. *Journal of the Royal statistical society: series B* **57**,

289–300 (1995).

59.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

60.   Oksanen, J. *et al.* vegan: Community Ecology Package. 2.7–1

https://doi.org/10.32614/CRAN.package.vegan (2001).

61.   Pielou, E. C. The measurement of diversity in different types of biological collections. *J.*

Theor. Biol. **13**, 131–144 (1966).

62.   Bertz, S. H. The first general index of molecular complexity. *J. Am. Chem. Soc.* **103**,

3599–3601 (1981).

63.   Hendrickson, J. B., Huang, P. & Toczko, A. G. Molecular complexity: a simplified

formula adapted to individual atoms. *J. Chem. Inf. Comput. Sci.* **27**, 63–67 (1987).

64.   Kim, S. *et al.* PubChem 2025 update. *Nucleic Acids Res.* **53**, D1516–D1525 (2025).

65.   R Core Team. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing (2022).

66.   RStudio Team. RStudio: Integrated development environment for R. (2022).

67.   Arnhold, E. easyanova: Analysis of Variance and Other Important Complementary

Analyses. 11.0 https://doi.org/10.32614/CRAN.package.easyanova (2012).

68.   Wickham, H. *et al.* Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

69.   Geng, J., Fang, H., Cheng, S. & Pei, J. Effects of N deposition on the quality and quantity

of soil organic matter in a boreal forest: Contrasting roles of ammonium and nitrate. *CATENA*

198, 104996 (2021).

70.   Sanderman, J. & Grandy, A. S. Ramped thermal analysis for isolating biologically

meaningful soil organic matter fractions with distinct residence times. *SOIL* **6**, 131–144

(2020).

71.   Arteaga-Clemente, G., García-González, M. A. & González-González, M. Soil lipid

analysis by chromatography: A critical review of the current state in sample preparation. *J.*

Chromatogr. Open **6**, 100173 (2024).

72.   Dinel, H., Schnitzer, M. & Mehuys, G. R. Soil Lipids: Origin, Nature, Content,

Decomposition, and Effect on Soil Physical Properties. in *Soil Biochemistry* 397–430

(Routledge, 2017). doi:10.1201/9780203739389-8.

73.   Kindler, R., Miltner, A., Thullner, M., Richnow, H.-H. & Kästner, M. Fate of bacterial

878   biomass derived fatty acids in soil and their contribution to soil organic matter. *Org.*

879   *Geochem.* **40**, 29–37 (2009).

880   74.   Miltner, A., Bombach, P., Schmidt-Brücken, B. & Kästner, M. SOM genesis: microbial

881   biomass as a significant source. *Biogeochemistry* **111**, 41–55 (2012).

882   75.   Olivelli, M. S. *et al.* Unraveling Mechanisms behind Biomass–Clay Interactions Using

883   Comprehensive Multiphase Nuclear Magnetic Resonance (NMR) Spectroscopy. *ACS Earth*

884   *Space Chem.* **4**, 2061–2072 (2020).

885   76.   Rempfert, K. R. *et al.* Biomolecular budget of persistent, microbial-derived soil organic

886   carbon: The importance of underexplored pools. *Sci. Total Environ.* **932**, 172916 (2024).

887   77.   Cyle, K. T., Klein, A. R., Aristilde, L. & Martínez, C. E. Ecophysiological Study of

888   *Paraburkholderia* sp. Strain 1N under Soil Solution Conditions: Dynamic Substrate

889   Preferences and Characterization of Carbon Use Efficiency. *Appl. Environ. Microbiol.* **86**,

890   e01851-20 (2020).

891   78.   Stone, B. W. G. *et al.* Life history strategies among soil bacteria—dichotomy for few,

892   continuum for many. *ISME J.* https://doi.org/10.1038/s41396-022-01354-0 (2023)

893   doi:10.1038/s41396-022-01354-0.

894   79.   Khan, K. S., Mack, R., Castillo, X., Kaiser, M. & Joergensen, R. G. Microbial biomass,

895   fungal and bacterial residues, and their relationships to the soil organic matter C/N/P/S ratios.

896   *Geoderma* **271**, 115–123 (2016).

897   80.   Liang, C., Amelung, W., Lehmann, J. & Kästner, M. Quantitative assessment of

898   microbial necromass contribution to soil organic matter. *Glob. Change Biol.* **25**, 3578–3590

899   (2019).

81.   Fahy, E. *et al.* Update of the LIPID MAPS comprehensive classification system for

lipids. *J. Lipid Res.* **50**, S9–S14 (2009).

82.   Liebisch, G. *et al.* Update on LIPID MAPS classification, nomenclature, and shorthand

notation for MS-derived lipid structures. *J. Lipid Res.* **61**, 1539–1555 (2020).

83.   Tian, Z. *et al.* PMhub 1.0: a comprehensive plant metabolome database. *Nucleic Acids*

Res. **52**, D1579–D1587 (2024).

84.   Klier, K. M. & Anantharaman, K. An updated view of metabolic handoffs in

microbiomes. *Trends Microbiol.* S0966842X25002203 (2025) doi:10.1016/j.tim.2025.07.009.

85.   Harper, D. B. Microbial metabolism of aromatic nitriles. Enzymology of C–N cleavage

by *Nocardia* sp. (*Rhodochrous* group) N.C.I.B. 11216. *Biochem. J.* **165**, 309–319 (1977).

86.   Howden, A. J. M., Rico, A., Mentlak, T., Miguet, L. & Preston, G. M. *Pseudomonas*

syringae pv. *syringae* B728a hydrolyses indole-3-acetonitrile to the plant hormone indole-3-

acetic acid. *Mol. Plant Pathol.* **10**, 857–865 (2009).

87.   Thimann, K. V. & Mahadevan, S. Nitrilase. *Arch. Biochem. Biophys.* **105**, 133–141

(1964).

88.   Pace, H. C. & Brenner, C. The nitrilase superfamily: classification, structure and

function. *Genome Biol.* **2**, reviews0001.1 (2001).

89.   Bessonnet, T. *et al.* Purification and Characterization of Nitphym, a Robust Thermostable

Nitrilase From Paraburkholderia phymatum. *Front. Bioeng. Biotechnol.* **9**, 686362 (2021).

90.   Moulin, L. *et al.* Complete Genome sequence of Burkholderia phymatum STM815T, a

broad host range and efficient nitrogen-fixing symbiont of Mimosa species. *Stand. Genomic*

Sci. **9**, 763–774 (2014).

91.   Teepakorn, C. *et al.* Nitrilase immobilization and transposition from a micro-scale batch

to a continuous process increase the nicotinic acid productivity. *Biotechnol. J.* **16**, 2100010

(2021).

92.   Vergne-Vaxelaire, C. *et al.* Nitrilase Activity Screening on Structurally Diverse

Substrates: Providing Biocatalytic Tools for Organic Synthesis. *Adv. Synth. Catal.* **355**, 1763–

1779 (2013).

93.   Wilhelm, R. C., DeRito, C. M., Shapleigh, J. P., Madsen, E. L. & Buckley, D. H.

Phenolic acid-degrading *Paraburkholderia* prime decomposition in forest soil. *ISME*

Commun. **1**, 4 (2021).

94.   Bhatnagar, J. M., Peay, K. G. & Treseder, K. K. Litter chemistry influences

decomposition through activity of specific microbial functional guilds. *Ecol. Monogr.* **88**,

429–444 (2018).

95.   Pisani, O., Frey, S. D., Simpson, A. J. & Simpson, M. J. Soil warming and nitrogen

deposition alter soil organic matter composition at the molecular-level. *Biogeochemistry* **123**,

391–409 (2015).

96.   Thirunavukkarasu, A. *et al.* Unraveling the dynamics of lignin chemistry on

decomposition to understand its contribution to soil organic matter accumulation. *Plant Soil*

https://doi.org/10.1007/s11104-024-07066-y (2024) doi:10.1007/s11104-024-07066-y.

97.   vandenEnden, L. *et al.* Molecular-level changes in soil organic matter composition after

10 years of litter, root and nitrogen manipulation in a temperate forest. *Biogeochemistry* **141**,

183–197 (2018).

98.   Alruwaili, A. *et al.* Elucidation of microbial lignin degradation pathways using synthetic

isotope-labelled lignin. *RSC Chem. Biol.* **4**, 47–55 (2023).

99.   DeAngelis, K. M. *et al.* Characterization of Trapped Lignin-Degrading Microbes in

Tropical Forest Soil. *PLoS ONE* **6**, e19306 (2011).

100.   DeAngelis, K. M. *et al.* Evidence supporting dissimilatory and assimilatory lignin

degradation in Enterobacter lignolyticus SCF1. *Front. Microbiol.* **4**, (2013).

101.   Bugg, T. D. H., Ahmad, M., Hardiman, E. M. & Rahmanpour, R. Pathways for

degradation of lignin in bacteria and fungi. *Nat. Prod. Rep.* **28**, 1883 (2011).

102.   Janusz, G. *et al.* Lignin degradation: microorganisms, enzymes involved, genomes

analysis and evolution. *FEMS Microbiol. Rev.* **41**, 941–962 (2017).

103.   Lee, S., Kang, M., Bae, J.-H., Sohn, J.-H. & Sung, B. H. Bacterial Valorization of Lignin:

Strains, Enzymes, Conversion Pathways, Biosensors, and Perspectives. *Front. Bioeng.*

Biotechnol. **7**, 209 (2019).

104.   Bugg, T. D., Ahmad, M., Hardiman, E. M. & Singh, R. The emerging role for bacteria in

lignin degradation and bio-product formation. *Curr. Opin. Biotechnol.* **22**, 394–400 (2011).

105.   Granja-Travez, R. S., Persinoti, G. F., Squina, F. M. & Bugg, T. D. H. Functional

genomic analysis of bacterial lignin degraders: diversity in mechanisms of lignin oxidation

and metabolism. *Appl. Microbiol. Biotechnol.* **104**, 3305–3320 (2020).

106.   Gu, J. *et al.* Bacterial transformation of lignin: key enzymes and high-value products.

Biotechnol. Biofuels Bioprod. **17**, 2 (2024).

107.   Kim, D., Kim, H.-W., & Hyoungseok Lee. Kraft Lignin Decomposition by Forest Soil

Bacterium *Pseudomonas kribbensis* CHA-19. *J. Microbiol. Biotechnol.* **34**, 1867–1875

(2024).

108. Ookawa, Z. *et al.* Catabolic System of Syringic Acid, a Key Intermediate of Lignin-

Derived Aromatic Compounds, via a Novel Linear Pathway in *Pseudomonas* sp. NGC7. *J.*

Agric. Food Chem. **73**, 18899–18913 (2025).

109. Frey, S. D., Knorr, M., Parrent, J. L. & Simpson, R. T. Chronic nitrogen enrichment

affects the structure and function of the soil microbial community in temperate hardwood and

pine forests. *For. Ecol. Manag.* **196**, 159–171 (2004).

110. Hesse, C. N. *et al.* Forest floor community metatranscriptomes identify fungal and

bacterial responses to N deposition in two maple forests. *Front. Microbiol.* **6**, (2015).

111. Li, X. *et al.* Nitrogen addition restricts key soil ecological enzymes and nutrients by

reducing microbial abundance and diversity. *Sci. Rep.* **15**, 5560 (2025).

112. Moore, J. A. M. *et al.* Fungal community structure and function shifts with atmospheric

nitrogen deposition. *Glob. Change Biol.* **27**, 1349–1364 (2021).

113. Frey, S. D. *et al.* Chronic nitrogen additions suppress decomposition and sequester soil

carbon in temperate forests. *Biogeochemistry* **121**, 305–316 (2014).

114. Van Diepen, L. T. A., Frey, S. D., Landis, E. A., Morrison, E. W. & Pringle, A. Fungi

exposed to chronic nitrogen enrichment are less able to decay leaf litter. *Ecology* **98**, 5–11

(2017).

115. Prescott, C. E. & Vesterdal, L. Decomposition and transformations along the continuum

from litter to soil organic matter in forest soils. *For. Ecol. Manag.* **498**, 119522 (2021).

116. Zhang, Z. *et al.* Changes in molecular composition, diversity, and network complexity of

soil organic carbon along the elevation in Changbai Mountain, Northeast China. *Geoderma*

443, 116842 (2024).

117.   Castañeda-Gómez, L. *et al.* Soil organic matter molecular composition with long-term

detrital alterations is controlled by site-specific forest properties. *Glob. Change Biol.*

gcb.16456 (2022) doi:10.1111/gcb.16456.

118.   Bossio, D. A. *et al.* The role of soil carbon in natural climate solutions. *Nat. Sustain.* **3**,

391–398 (2020).

119.   Crow, S. E. *et al.* Sources of plant-derived carbon and stability of organic matter in soil:

implications for global change. *Glob. Change Biol.* **15**, 2003–2019 (2009).