Multi-omics-driven kinetic modeling reveals metabolic vulnerabilities and differential drug-response dynamics in ovarian cancer

Ilias Toumpe[1], Maria Masid[2,3], Vassily Hatzimanikatis[1,3], Ljubisa Miskovic[1,3]

[1]Laboratory of Computational Systems Biotechnology, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

[2]Ludwig Institute for Cancer Research, Department of Oncology, University of Lausanne and Lausanne University Teaching Hospital (CHUV), Lausanne, Switzerland

[3]Correspondence : maria.masidbarcon@unil.ch, vassily.hatzimanikatis@epfl.ch, ljubisa.miskovic@epfl.ch

Summary

Cancer cells reprogram metabolism to fuel aggressive growth and resist therapies, yet mapping these metabolic changes remains a major challenge. While constraint-based genome-scale models describe steady states, kinetic models that resolve time-dependent fluxes and metabolite concentrations in human cancers are rare. Here, we integrate multi-omics data and physicochemical constraints to construct large-scale kinetic models of ovarian cancer metabolism that capture how tumor cells adapt their nutrient use and energy production. Comparing ovarian cancer cells with and without *BRCA1* loss revealed distinct metabolic strategies driven by transcriptional regulation. The models reproduce hallmark phenotypes, correctly predict known metabolic drug targets, and identify previously uncharacterized vulnerabilities in nucleotide and lipid synthesis. Simulations of drug treatments mirror clinical responses and consistently reveal a ceramide-linked stress signature common to many chemotherapies. Using the same framework, we show that *BRCA1* loss redirects metabolic pathway activity through enzyme activity changes linked to transcription factors interacting with *BRCA1*, suggesting regulatory routes for network-level rewiring. By capturing dynamic fluxes and concentrations, these models bridge molecular insight with therapeutic action, guiding biomarker discovery and dosing strategies. This open-access resource provides a mechanistic foundation for testing metabolic interventions, deepening our understanding of cancer metabolism across tumor types, and advancing the promise of precision oncology.

Keywords: cancer metabolism, ovarian cancer, *BRCA1* loss of function, drug targets, large-scale kinetic models, multi-omics, drug-response simulation

Introduction

Altered metabolism has long been accepted as one of the hallmarks[1–3] of cancer physiology. Cancer cells undergo metabolic rewiring, which allows them to achieve elevated growth rates, survive anticancer treatments, and adapt to nutrient-deprived environments. Studies have shown common metabolic alterations across different types of cancer cells, including the Warburg effect, increased biosynthetic production, and elevated energy requirements.[4–6] These conserved metabolic traits have garnered considerable interest from the scientific community, as they may reveal metabolic vulnerabilities that can be therapeutically targeted to halt tumor growth and proliferation.[7] For example, folate antagonists, among the first successful chemotherapeutic treatments, specifically target metabolic enzymes in the nucleotide synthesis pathway, inspiring the development of many subsequent anti-metabolite drugs.[8]

Given the complexity of cellular metabolism, a wide range of experimental and computational technologies have been developed to study its structure and function, resulting in a vast amount of data across various biological contexts.[9–11] One of the most effective tools for systematically studying cancer metabolism is Genome-Scale Metabolic Models (GEMs).[12,13] GEMs provide a computational platform for integrating omics data and estimating intracellular metabolic fluxes using constraint-based modeling methods, such as Flux Balance Analysis (FBA).[14–16] These modeling pipelines have been instrumental in capturing key characteristics

of human health[17–22] and cancer metabolism, enabling the identification of candidate anticancer drugs[23–25], exploration of cancer phenotypes[26,27], and forming hypotheses about the mechanisms underlying cellular processes.[28] Importantly, GEMs developed using tissue- and patient-specific data have revealed distinct metabolic responses and vulnerabilities, indicating the importance of patient-specific approaches in cancer metabolism studies.[29]

However, the growing availability of proteomics and metabolomics has exposed core limitations of conventional GEM-based approaches. GEMs typically operate under steady-state flux assumptions and do not explicitly account for time-dependent features, such as metabolite concentrations, enzyme abundances, or regulatory effects. Efforts have been made to integrate multi-omics data and add more physiological constraints[20,21,30–34] to enhance the predictive capabilities of the models, but they still rely on simplifying assumptions and lack mechanistic enzyme kinetics and transient dynamics.[35] These gaps are particularly significant in cancer, where metabolic reprogramming and drug responses are inherently dynamic.[36] FBA-centric analyses typically emphasize gene knockouts and growth. As a result, they fail to capture how shifts in metabolite levels or enzyme activities shape drug resistance[37], toxicity[38,39], and tumor-microenvironment interactions[28,40,41], where clinically meaningful phenotypes often arise.

While GEMs provide a versatile framework for steady-state analysis, kinetic models go beyond this approach by explicitly representing enzyme kinetics and regulation through systems of ordinary differential equations (ODEs) [35], and have been recognized for their mechanistic depth and applications in human metabolism.[12,35,36,42] Small-scale applications to glycolysis and central carbon metabolism have revealed rate-limiting steps where enzyme activity or metabolite availability constrain pathway function.[43–46] Personalized kinetic models of red blood cell metabolism have further shown how patient-specific parameters can refine mechanistic predictions and support precision medicine.[36]

To address the complexity of tumor metabolic reprogramming, however, it is necessary to scale kinetic modeling to the systems level, developing large- and near-genome-scale models that integrate multi-omics data and capture tissue- or patient-specific transient metabolic behavior. Furthermore, a single kinetic model with a single parameter set offers a narrow snapshot of cancer metabolism and cannot represent the diversity observed within cell lines or among patients. Therefore, there is a need to construct populations of kinetic models that vary slightly in parameters and initial metabolic states to approximate this heterogeneity and capture tissue- or patient-specific behavior. Studying these ensembles preserves variability while showing that distinct metabolic states can still reveal common biological mechanisms and global therapeutic strategies.

In this work, we present a resource of curated large-scale kinetic models of ovarian cancer metabolism, built upon multi-omics and physicochemical data for *BRCA1* wild-type (*BRCA1*^WT) and mutant (*BRCA1*^MUT) cell lines.[47] We employed these physiologically relevant model populations to identify metabolic drug targets, accurately predicting known targets and identifying novel ones. The models were also used to investigate the network-wide transient responses of cancer metabolism to different drug administration simulations, capturing both experimentally observed and previously uncharacterized metabolic shifts. We further explored potential regulatory mechanisms underlying the distinct physiologies of *BRCA1*^WT and *BRCA1*^MUT cancer cells, providing a core set of regulatory structures for further research.

Overall, this resource offers a mechanistic foundation for *in silico* studies aimed at understanding cancer metabolism and guiding cancer prognosis, treatment response, and therapeutic strategies.

Results

Overview of the reconstruction framework for physiologically relevant kinetic models of cancer metabolism

We present a population of near-genome-scale kinetic models of cancer metabolism that capture the distinct physiologies of ovarian cancer cells with and without a *BRCA1* mutation, and enable dynamic interrogation of enzyme-level vulnerabilities and network-wide adaptations. Kinetic model construction relies on a context-specific metabolic network derived from model reduction[48,49] and, when available, integrated multi-omics data[30,31]. Here, we used and slightly extended the cancer-specific stoichiometric model of Masid et al. (unpublished work), derived from Recon3D[50] through systematic reduction to ovarian cancer pathways and integrated with gene expression data from two isogenic cell lines (*BRCA1*[WT] and *BRCA1*[MUT]) together with exofluxomic and exometabolomic datasets under thermodynamic constraints. The extension included incorporating measured adenylate levels (ATP, ADP, and AMP) and enforcing minimal enzymatic activity across all reactions to ensure global network connectivity. The resulting steady-state models capture distinct flux distributions consistent with differential gene activity and served as the scaffold for subsequent kinetic model construction (Figure 1A).

We next generated flux and concentration samples from each steady-state model to form a population of feasible metabolic states (Figure 1B). Sampling revealed multiple feasible metabolic phenotypes, highlighting the inherent metabolic diversity found among genetically similar cancer cells.[51,52]

Using the reduced metabolic network as the starting point, we constructed two large-scale kinetic model ensembles, one for *BRCA1*[WT] and one for *BRCA1*[MUT] ovarian cancer cells. Each model comprises more than 710 mass balances (ODEs) and more than 2,000 reactions governed by diverse kinetic rate laws that reflect the underlying enzymatic mechanisms (Figure 1C).[53] In total, each model contains over 9,500 kinetic parameters. To parameterize the models, we used flux and concentration profiles obtained from the steady-state models and sampled kinetic parameters consistent with the observed metabolic behavior (Figure 1D, Methods). Model instances exhibiting non-physiological dynamics were discarded, ensuring that the final model populations capture biologically meaningful metabolic behavior beyond what steady-state models alone can reveal.

The curated model populations reproduce the metabolic diversity observed across cancer cells, providing physiologically grounded, dynamic representations of cancer cell metabolism. Validation against experimental data (Figure 1) confirms that the models provide a reliable representation of ovarian cancer metabolism. The curated *BRCA1*[WT] and *BRCA1*[MUT] model populations now provide a platform to investigate (i) which enzymes are most critical for sustaining tumor growth and thus represent promising drug targets, (ii) how the metabolic network adapts over time when exposed to anticancer drugs, and (iii) which specific enzyme

activity changes underlie metabolic differences between *BRCA1*[WT] and *BRCA1*[MUT] cells. Together, these applications establish the model ensembles as a systematic framework for exploring therapeutic vulnerabilities and regulatory rewiring in ovarian cancer metabolism.

Figure 1. **Overview of the reconstruction workflow, model characteristics, and applications of large-scale kinetic models of *BRCA1*[WT] and *BRCA1*[MUT] ovarian cancer metabolism.** (A) The cancer-specific metabolic network was derived from the human genome-scale model Recon3D through a systematic reduction process. Transcriptomic, exometabolomic, and thermodynamic data were integrated to tailor the model to ovarian cancer physiology. (B) Sampling steady-state fluxes and metabolite concentrations produced physiologically relevant metabolic states that served as the basis for the kinetic models. (C) Each reaction was assigned a kinetic rate law mechanism, resulting in a

model scaffold comprising hundreds of mass balances, rate laws, and kinetic parameters. (D) Parameter sets were sampled and filtered for physiological relevance, yielding populations of $BRCA1^{WT}$ and $BRCA1^{MUT}$ kinetic models that capture the observed metabolic diversity. These ensembles were used to identify drivers of proliferation, simulate drug administration, quantify pharmacodynamic responses and network-wide shifts, and link enzyme alterations to upstream transcription factors associated with $BRCA1$-dependent metabolic rewiring.

Drug target analysis predicts known and novel cytostatic targets

A central challenge in cancer research is identifying strategies to slow or halt tumor cell proliferation. Kinetic models of metabolism provide a quantitative means to assess how changes in enzymatic activity reshape the metabolic state and impact cancer cell growth. Using the $BRCA1^{WT}$ and $BRCA1^{MUT}$ kinetic model populations, we applied Metabolic Control Analysis (MCA)[54,55] to quantify how changes in enzyme activity propagate through the network and alter reaction fluxes and metabolite concentrations. We focused on the control of cellular growth rate to identify enzymes whose activity most strongly determines proliferation, and thus represents potential metabolic drug targets. The MCA workflow yielded distributions of control coefficients, capturing the mean effect and variability of each enzyme's impact on growth.

We identified the top 20 enzymes with the strongest effect on cancer proliferation in each of the two cell types ($BRCA1^{WT}$ and $BRCA1^{MUT}$), resulting in a combined set of 28 unique enzymes (Figure 2A). Whether enzymes exhibited positive or negative control over proliferation was generally consistent between $BRCA1^{WT}$ and $BRCA1^{MUT}$ models, although differences in magnitude revealed context-specific sensitivities. For example, most enzymes involved in central carbon metabolism have a greater effect on the growth rate of $BRCA1^{MUT}$, indicating a stronger dependence on glycolytic activity (Figure 2A). On the other hand, mitochondrial complex I and multiple enzymes from lipid metabolism exert greater control over $BRCA1^{WT}$ proliferation, reflecting a higher reliance on oxidative phosphorylation and lipid biosynthesis (Figure 2A). Most high-control enzymes have a positive effect on cancer cell proliferation, indicating that their inhibition would attenuate tumor growth, e.g., PGI, TMDS, and mitochondrial complex I. Conversely, a smaller subset exhibits negative control, suggesting that enhancing their activity could suppress cancer progression, including PPAP, NTD1, and MI1PP. Such inverse relationships have been observed in cancer metabolism. For instance, GLS2 is downregulated in hepatocellular carcinoma, and its overexpression has been shown to markedly impair tumor cell colony formation.[56]

To place these findings in the context of the broader cellular metabolic network, we mapped the functional roles of the 28 targets across major metabolic subsystems (Figure 2A), revealing their participation in anabolic (nucleotide and lipid biosynthesis), catabolic (glycolysis, TCA cycle, urea cycle), and bioenergetic (oxidative phosphorylation) pathways. These results align with therapeutic directions highlighted in recent reviews, which emphasize that targeting anabolic, catabolic, bioenergetic, and signaling processes represents a core strategy in cancer metabolism.[8,57] Our models encompass the first three domains, while signaling remains beyond the scope of this study.

Validation with known drug targets and implications for repurposing

Having established a set of enzymes with strong control over cancer cell proliferation, we next examined which of these predicted targets are already associated with existing drugs and which represent novel predictions of the kinetic models. Of the 28 identified enzyme targets, 24 are linked to experimental or market-approved drugs in the U.S., Canada, and the EU, as cataloged in DrugBank[58] (Figure 2B; Data S1). More importantly, 11 out of the 28 are targeted by at least one anticancer drug (Figure 2B). Notably, there are 20 known antimetabolites, which are structural analogs of natural metabolites that disrupt key metabolic pathways (ATC code L01B), and the models identified associated targets for 16 of these (Data S2). The remaining four include: (i) Carmofur, a fluorouracil derivative without documented targets and withdrawn from the market; (ii) one drug without associated enzymes in Recon3D; and (iii) two drugs whose targets appear in the reduced model but were not predicted here.

Many additional predicted targets correspond to drugs initially developed for non-cancer indications, with several already in clinical trials for the treatment of cancer. Overall, ~60% of all documented drugs linked to the 28 targets are in experimental stages or clinical evaluation (Figure 2B), reflecting sustained therapeutic interest across multiple diseases, including cancer.

We also recorded each drug's mode of action (inhibitory, activating, or antagonistic) (Data S1). Most act as inhibitors, consistent with our finding that most enzymes have a positive influence on growth. For example, thymidylate synthase (TMDS) is associated with 23 drugs, 10 of which act as inhibitors (Figure 2B, Data S1). In a few cases, however, drug effects diverged from model predictions. For example, IDH2 (ICDHxm) and CYTK1 were predicted to suppress growth when their activity increases (Figure 2A), yet both are inhibited by anticancer agents. However, it is worth noting that the documented anticancer drugs for IDH2 target mutant isoforms of the enzyme, which acquire neomorphic activity, producing the oncometabolite D-2-hydroxyglutarate.[59,60] These inhibitors have been produced to selectively target mutant activity rather than wild-type IDH2. The apparent discrepancy thus reflects the specificity of current therapies for mutant IDH2 rather than a limitation of the modeling framework.

Novel targets and mechanistic insights

Among the enzymes highlighted by our models, most align with experimentally validated mechanisms, whereas the four with no associated drugs represent previously unexplored metabolic vulnerabilities. These novel candidates may offer promising avenues for future metabolic drug development, and we further evaluated their biological relevance and potential as therapeutic targets:

i. UPRT, a uracil phosphoribosyltransferase (HMR_4343). This enzyme catalyzes the salvage of uracil to UMP, a key precursor in pyrimidine nucleotide biosynthesis. In our model, inhibition of URPT reduces nucleotide pools and slows predicted growth rates, underscoring the enzyme's potential role in sustaining rapid proliferation. Although the gene has been annotated in the human genome, its enzymatic activity has not been validated *in vitro*, and no studies have established a link between it and tumor biology.[61] Nevertheless, pyrimidine metabolism is a well-established therapeutic axis, with multiple drugs targeting *de novo* or salvage pathways. The model positions URPT as a potential bottleneck in nucleotide supply

and broadens the scope of established targeting strategies to an uncharacterized salvage enzyme, highlighting a gap where experimental validation could confirm a new metabolic vulnerability in cancer.

ii. TRIOK, a triokinase. TRIOK activity participates in the production of glyceraldehyde 3-phosphate. Perturbation of TRIOK alters flux through glycolytic and fructose metabolism pathways, affecting predicted growth rates, and suggests that cancer cells may rely on this enzyme when rewiring central carbon metabolism. While glycolysis and fructose metabolism are known to undergo rewiring in tumors, TRIOK itself has received little attention in cancer research.[62] These findings raise the possibility that TRIOK supports metabolic flexibility in tumors, a role that remains to be tested experimentally.

iii. KDSR, a 3-dehydrosphinganine reductase (3DSPHR). Our model predicts that reduced KDSR activity slows growth by limiting the supply of structural sphingolipids required for membrane biosynthesis, leading to the accumulation of its substrate, 3-ketodihydrosphingosine. Blocking the activity of KDSR is toxic to cancer cells but not to normal cells[63], providing a mechanistic link between the model prediction and selective cytotoxicity. Indeed, recent studies have demonstrated that many tumors rely on increased *de novo* sphingolipid biosynthesis to sustain their signaling and structural demands, and that targeting KDSR results in the buildup of 3-ketodihydrosphingosine and cancer cell death.[63,64] Thus, while the model provides a direct metabolic explanation for slower growth, it also highlights the accumulation of 3-ketodihydrosphingosine as a measurable signature that can mechanistically connect to the downstream cytotoxic effects observed experimentally.

iv. PPAP, a phosphatidic acid phosphatase. Kinetic modeling predicts that increased PPAP activity suppresses growth by redirecting flux through lipid metabolism, specifically by converting phosphatidic acid to diacylglycerol and lowering the levels of the signaling lipid lysophosphatidic acid (LPA). This prediction aligns with experimental observations showing that phosphate phosphatase enzymes are significantly downregulated across multiple cancers, including ovarian cancer.[65–67] Experimental studies have shown that introducing hLPP-3[68] (which hydrolyzes phosphatidic acid, LPA, and sphingosine) or LPP-1[67] into ovarian cancer cells reduces colony formation, promotes apoptosis, and inhibits tumor growth, suggesting that targeting LPA metabolism or signaling may be a promising therapeutic approach for ovarian cancer.[68] Thus, while the model predicts growth inhibition through shifts in lipid metabolic balance, experimental evidence connects this mechanism to apoptosis and tumor suppression.

These findings demonstrate the model's capacity to uncover both well-established and previously uncharacterized metabolic vulnerabilities. In some cases, the model directly predicts slower growth or altered metabolite levels, while in others, external experimental evidence links these metabolic changes to downstream phenotypes such as toxicity, apoptosis, or tumor suppression. Importantly, none of these targets were explicitly biased in the modeling process. Rather, their identification emerged naturally from the model architecture, kinetic formulation, and integrated multi-omics constraints.

Metabolic Enzyme Targets Disrupt Nucleotide and Lipid Supply

We interpreted the effect of each enzyme on cellular growth through its impact on the synthesis of biomass precursors: inhibiting an enzyme that depletes or limits the synthesis of

a growth-limiting precursor is expected to decrease growth; conversely, increased precursor synthesis could promote it. In Recon3D, cellular growth is represented as the net synthesis of amino acids, nucleotides, deoxynucleotides, lipids, and glucose-6-phosphate. For each enzyme, we quantified its mean effect on the synthesis rate (Figure 2C) and intracellular concentration (Figure 2D) of each biomass precursor.

Most enzymes had minimal to no effect on either the concentration or the synthesis rate of amino acids, suggesting that cancer cells are robust to perturbations in amino acid metabolism. This finding aligns with prior reports[69,70] showing that therapies targeting amino acid metabolism often have limited efficacy *in vivo*. Whereas in vitro assays typically probe isolated pathways, our kinetic models capture system-wide metabolic interactions that compensate for such perturbations, providing a mechanistic explanation for the reduced therapeutic response observed *in vivo*. Targeting the metabolism of essential amino acids is likely to cause systemic toxicity, while tumors can reroute fluxes around inhibitors of non-essential amino acid synthesis.[69]

In contrast, two precursors, sphingomyelin and deoxythymidine triphosphate (dTTP), were consistently sensitive to most enzyme targets for both physiologies (Figure 2C, 2D):

i. Sphingomyelin, which accounts for roughly 10–20% of cell membrane lipids, supports membrane rigidity and lipid raft formation through tight packing with cholesterol. It also serves as a source of signaling molecules such as ceramide and S1P, both regulators of apoptosis and proliferation.[71] In our models, sphingomyelin is explicitly required in the biomass composition, ensuring that its contribution to growth is quantitatively represented. The models provide a non-obvious mechanistic insight by predicting that triokinase (TRIOK) and 3-dehydrosphinganine reductase (3DSPHR) exert the strongest control over sphingomyelin levels (Figure 2C, 2D). Specifically, increased TRIOK activity and decreased 3DSPHR activity are predicted to divert carbon away from sphingolipid metabolism, leading to sphingomyelin depletion.

ii. Deoxythymidine triphosphate (dTTP) is essential for DNA synthesis and repair, making it critical for rapidly proliferating cells. Cancer cells are particularly sensitive to disruptions in dTTP synthesis since limiting its availability causes uracil misincorporation, leading to DNA damage and cell death.[72] Its *de novo* synthesis largely depends on a single enzyme, thymidylate synthase (TMDS), a well-established therapeutic target associated with many clinically approved anticancer drugs (as also predicted by the models; Figure 2B).[72] The biomass formulation explicitly requires balanced pools of nucleotides, including dTTP, and the models predict that thymidylate synthase (TMDS) and 5'-nucleotidase (NTD1) are the most significant regulators of dTTP homeostasis (Figure 2C, 2D). TMDS is the only enzyme that produces *de novo* dTMP from dUMP, while NTD1 diverts dUMP away from dTTP production by converting it into deoxyuridine. Thus, the models show how dTTP balance can be controlled by both established and less obvious enzymes, reinforcing known strategies and revealing new vulnerabilities.

Overall, our framework successfully identified a set of enzymes with strong and specific control over cancer cell proliferation, many of which correspond to existing or emerging drug targets. Beyond growth control, the models uncovered systemic nodes that regulate the balance of key biomass precursors, providing mechanistic insight into how cellular homeostasis can be maintained or disrupted. In doing so, the models not only recapitulate well-validated

therapeutic strategies but also highlight less obvious enzymes whose influence would not be apparent from single pathways alone. As this approach is extended to incorporate patient-specific omics data, it holds promise for guiding personalized interventions and precision oncology strategies.

Figure 2. Metabolic Control Analysis of kinetic model populations identifies enzymes with strong control over cancer proliferation, their associated drug targets, and the impacts on biomass precursors. (A) Mean flux control coefficients (FCCs), quantifying the relative effect of a small

change in enzyme activity on growth, for $BRCA1^{WT}$ (blue) and $BRCA1^{MUT}$ (red) kinetic model populations. Error bars denote the interquartile range (25th–75th percentiles). Displayed are the 28 enzymes with the highest absolute FCCs, grouped by metabolic subnetworks. Positive FCCs indicate that enzyme inhibition suppresses growth, and negative FCCs indicate that inhibition enhances proliferation. (B) Stacked bar chart showing the number of documented drugs targeting each of the 28 control-coefficient enzymes, classified by Anatomical Therapeutic Chemical (ATC) categories. Drug–target associations were curated from the DrugBank database. Pie charts illustrate the number of market-approved (blue) and experimental (red) drugs associated with each enzyme, with the larger pie chart displaying the totals across all targets. Heatmaps of (C) average flux control coefficients (FCC) and (D) concentration control coefficients (CCCs), quantifying how enzyme activity affects synthesis rates and metabolite concentrations, on biomass precursor classes: nucleotides, deoxynucleotides, lipids, and glucose-6-phosphate. Essential and nonessential amino acids are omitted due to minimal control. Color scale denotes the direction and magnitude of the CCCs (blue: negative effect; red: positive effect). Lists of enzyme and biomass precursor abbreviations are provided in Tables S1, S2.

Simulations uncover transient metabolic shifts during cancer drug treatment

Despite decades of effort to target cancer metabolism, the mechanisms underlying many metabolic therapies remain poorly understood across the complex metabolic networks of both cancerous and healthy cells.[8] Clinical efficacy is often inconsistent, and unintended side effects can emerge due to system-wide metabolic shifts.[73] Dissecting these diverse outcomes is particularly challenging in human tumors, where dynamic microenvironments are experimentally complicated to probe. Kinetic models offer a systems-level solution to address these limitations by simulating drug administration and tracking how interventions impact metabolic fluxes and intracellular concentrations over time. Such simulations can reveal mechanisms and adaptive responses that are inaccessible through experiments alone.

We characterized intracellular metabolic responses to drug therapies by simulating pharmacological interventions against three functionally distinct metabolic enzyme targets in $BRCA1^{WT}$ and $BRCA1^{MUT}$ cancer models: (1) the hexokinases (HEX1, r0354, r0355), which catalyze the rate-limiting step of glycolysis and will serve as our primary case study; (2) thymidylate synthase (TMDS), a clinically validated anticancer target; and (3) triokinase (TRIOK), a novel growth-associated enzyme predicted by our models. All three were previously identified by Metabolic Control Analysis (MCA) as exerting strong control over cancer cell proliferation (Figure 2A).

Differential Drug Response Across Metabolic States

As a representative case, we first examined hexokinase inhibition in $BRCA1^{WT}$ ovarian cancer (Figure 3), with analogous analyses performed across all targets and physiologies (Figure S1-S5). To capture distinct metabolic states and examine their differential responses to treatment, we selected a representative subset of kinetic models through stratified sampling based on their glycolytic and mitochondrial ATP production levels (Figures S6, S7). This sampling design mirrors patient stratification in clinical trials. It allows us to assess how cancer cells that regulate glycolytic and mitochondrial ATP production independently respond to targeted inhibitors.[5] By doing so, the analysis captures metabolic heterogeneity, which is known to dictate therapeutic outcomes.[37,39]

Figure 3. Hexokinase inhibition elicits dose-dependent growth arrest and network-wide metabolic adaptation in *BRCA1*[WT] ovarian cancer models. (A) Dose–response curves of cell viability from 972 kinetic models, plotted against residual hexokinase activity. Despite variability in baseline states and parameters, all converge toward a common threshold: approximately two-thirds inhibition induces a 50% growth reduction ($EC_{50} = 0.33 \pm 0.18$). Grey dots represent individual model responses; orange-shaded ranges denote fitted logistic curves, with the average curve highlighted. (B) Transient growth trajectories following a 10-fold reduction in hexokinase activity, stratified by glycolytic vs. mitochondrial ATP production initial states (colored boxes). Shaded areas indicate interquartile ranges,

and solid lines trace the model closest to each group's mean. Transient responses of (C) oxygen consumption rate (OCR), (D) mitochondrial ATP synthesis, (E) lactate secretion rate, and (F) glycolytic ATP synthesis following the same inhibition. (G) Final growth rate distribution across the population following 10-fold inhibition. Nearly half of the models cease proliferation, underscoring pronounced heterogeneity in drug response. (H) Changes in biomass precursor pools after inhibition. Glucose-6-phosphate emerges as the principal bottleneck, accompanied by depletion of lipid species and dATP. Bars denote average percent changes, and the error ranges indicate the interquartile range for the five most affected precursors. (I-J) Volcano plots of steady-state changes in (I) reaction fluxes and (J) metabolite concentrations. Dots are color-coded by network distance from the inhibited hexokinases. Central carbon intermediates generally decrease, whereas several nucleotide and sphingolipid species accumulate, revealing pathway-specific rerouting upon glycolytic blockade.

We simulated simultaneous inhibition of all hexokinase isoforms by progressively reducing their enzymatic activity over an 8-hour window (Figure S8) and tracking the cancer cells' metabolism until it reached a new metabolic state. The 8-hour timeframe was chosen as a conservative approximation of *in vivo* drug delivery, considering the limitations of tumor penetration and cellular uptake. In our framework, drug action was modeled by directly reducing the enzyme activity (Figure S8). However, if the mechanism of action is known, more mechanistic pharmacokinetic and pharmacodynamic (PK/PD) descriptions could be integrated, enabling simulation of competitive inhibition, allosteric modulation, or other regulatory effects.

Dose–response curves were generated for all selected models to assess cell viability, measured as changes in final growth rate across progressively reduced enzyme activity (Figure 3A). Because the models differ in their kinetic parameters and in the initial metabolic states at which inhibition is applied, the simulated responses varied substantially, mirroring the diversity of patient-to-patient physiology. Nevertheless, a clear pattern emerged: modest reductions in hexokinase activity had a limited impact, while complete inhibition reliably suppressed proliferation across all models. Logistic regression analysis was used to derive pharmacodynamic metrics such as EC_{50} (as performed in other studies[74]), revealing that, on average, a two-thirds reduction in enzyme activity was required to achieve 50% growth inhibition ($EC_{50} = 0.33 \pm 0.18$). These simulations provide a mechanistic estimate of the degree of inhibition required to effectively impair cancer proliferation.

To investigate how cells adapt at the network level, we examined the effects of a 10-fold reduction in hexokinase activity on both fluxes and metabolite concentrations. The resulting responses varied across models with different initial glycolytic and mitochondrial activities, as shown in the apparent differences in final growth rates (Figure 3B) and internal metabolic states (Figure 3C-F). Models with lower glycolytic flux were generally more sensitive, showing reduced proliferation regardless of mitochondrial ATP output. Some models compensated by increasing the oxygen consumption rate (OCR) (Figure 3C) and mitochondrial ATP production (Figure 3D), while lactate secretion rate remained largely unchanged for all models (Figure 3E), even if glycolytic ATP generation decreased (Figure 3F). This compensatory increase in mitochondrial respiration following suppression of glycolysis has also been observed in cancer cells subjected to hexokinase inhibition.[75,76] These results confirm that distinct metabolic states exhibit differential responses to the same enzymatic inhibition.

Population-level analysis showed that nearly half of the models ceased proliferation entirely (Figure 3D). Because the simulations track all intracellular metabolite concentrations and

fluxes, we identified which biomass precursors were most depleted (Figure 3E). Glucose-6-phosphate was the principal bottleneck, consistent with its position as the direct product of the hexokinase-catalyzed reactions. Lipid species and dATP were also depleted, indicating growth arrest from the combined shortage of multiple precursors. This contrasts with the TMDS inhibition, where only dTTP was significantly depleted (Figures S1 and S4).

We next examined how hexokinase inhibition propagates across the metabolic network by comparing the initial and final metabolic states through the fold changes of fluxes and concentrations (Figures 3I, J). Central carbon intermediates generally declined, while several nucleotides and sphingolipid-related metabolites accumulated, revealing broad metabolic reprogramming (Figure 3J). Flux changes followed a similar pattern, with substantial decreases immediately downstream of hexokinase (one to four reaction steps) and a selective reduction in distal reaction fluxes, reflecting adaptive network rerouting (Figure 3I). Overall, these results demonstrate how kinetic modeling can capture the intracellular consequences of targeted therapy and inform the selection of metabolic biomarkers for follow-up studies.

Shared and Distinct Effects Across Drug Targets and Physiologies

Having established the workflow in the hexokinase inhibition case, we next applied the same workflow to TMDS and TRIOK inhibition in both *BRCA1*[WT] and *BRCA1*[MUT] model populations (Figures S2–S6). Our goal was to assess whether *BRCA1* status is associated with distinct drug-response patterns and whether shared metabolic shifts occur independently of the specific therapy. To this end, we summarized the concentration and flux changes across models using pathway-level enrichment analysis, mapping them to pathways annotated in Recon3D (Figure 4). As anticipated from their distinct biological roles, each target produced characteristic responses. Hexokinase and triokinase inhibition resulted in system-wide shifts in energy and biosynthetic fluxes, while thymidylate synthase inhibition led to nucleotide-specific depletions (Figure 4A, 4B). Interestingly, TRIOK inhibition triggered distinct metabolic rewiring depending on the *BRCA1* physiology: the pentose phosphate pathway was disrupted in *BRCA1*[WT], while cholesterol and lipid metabolism were significantly affected in *BRCA1*[MUT] (Figure 4A).

Across all targets and physiologies, sphingolipid metabolism was consistently deregulated, characterized by accumulation of ceramide, dihydroceramide, sphingomyelin, and sphinganine. This pattern aligns with the established role of ceramide in stress responses. Ceramide is a key bioactive lipid that engages signaling pathways governing growth arrest, differentiation, and senescence.[77] Interestingly, similar to our results, ceramide levels have been found to increase during various drug therapies and across different types of cancer.[78–80] In drug-resistant cancer cells, ceramide is converted into sphingomyelin, which is suggested to help prevent pro-apoptotic signals.[81,82] While our current models lack the S1P branch, the consistent accumulation of ceramide and its derivatives recapitulates metabolic contexts that mirror experimentally observed drug-induced shifts. Importantly, the model itself does not simulate signaling cascades; however, the models suggest that ceramide accumulation represents a direct metabolic signature of growth arrest, rather than a downstream effect specific to individual drug mechanisms. This shared metabolic buildup provides a mechanistic explanation for why diverse therapies trigger apoptosis through ceramide signaling.

In summary, our population of kinetic models captures how metabolic therapies reshape cancer metabolism, revealing vulnerabilities that depend on cellular context and reproducing known drug-induced shifts in metabolism. By linking dose–response behavior to pathway-level flux and metabolite changes, the models provide mechanistic insights often inaccessible experimentally, offering a robust platform for identifying therapeutic targets, characterizing metabolic biomarkers, and guiding strategies for precision metabolic interventions.

Figure 4. Distinct targets and *BRCA1* physiologies produce characteristic subsystem signatures with a common stress response. (A) Pathway-level enrichment analysis of steady-state metabolite concentration changes predicted across kinetic model populations. Enrichment represents the fraction of consistently deregulated metabolites per pathway (hypergeometric test), with circle size denoting this fraction and color indicating the corresponding p-value. Each drug produces a distinct subsystem signature: for example, TRIOK blockade selectively perturbs the pentose phosphate pathway (PPP) and cholesterol/lipid metabolism in *BRCA1*MUT cells, whereas TMDS inhibition primarily affects nucleotide metabolism in both physiologies. All three interventions consistently deregulate sphingolipid metabolism, reflecting the ceramide-mediated stress response characteristic of diverse chemotherapeutics. (B) Schematic overview of the effects of hexokinases (HEX), triokinase (TRIOK), and thymidylate synthase (TMDS) inhibition on the core metabolic pathways. Red arrows denote increased pathway activity, green arrows denote decreased activity, and equality symbols indicate minimal or no change.

Cancer phenotype comparisons reveal metabolic and regulatory shifts

BRCA1 loss-of-function (as in *BRCA1*[MUT]) reshapes ovarian cancer metabolism, manifesting as elevated glycolysis and other malignant traits compared to *BRCA1*[WT], which retains a less aggressive metabolic state. Yet the extent to which these changes reflect direct regulatory consequences of *BRCA1* loss remains largely unexplored.

To address this, we used the kinetic models to identify the minimal set of enzyme activity changes that can account for the metabolic differences between *BRCA1*[WT] and *BRCA1*[MUT] cells. Defining these enzymatic changes provides a mechanistic basis to link altered metabolic states back to upstream regulation by connecting the affected enzymes to transcription factors and ultimately to *BRCA1*.

We quantified metabolic differences by comparing pathway fluxes between *BRCA1*[MUT] and *BRCA1*[WT] across 12 major subsystems of central and peripheral metabolism (Figure 5). These pathways, comprising 298 reactions, span commonly studied subnetworks such as glycolysis and mitochondrial energy production[83–85] , but also extend into additional subsystems, providing a more holistic view of the metabolic rewiring that underlies the *BRCA1*[MUT] phenotype.

As we allowed progressively more enzymes to change activity, the metabolic state of *BRCA1*[MUT] became increasingly similar to that of *BRCA1*[WT] cells (Figure 5A, Figure S9). Strikingly, adjusting 250 enzymes was almost as effective as altering all candidates, capturing most of the differences between the two physiologies (Figure S9). Many pathways showed substantial improvement in flux alignment, and in nucleotide metabolism, specifically purine and pyrimidine biosynthesis, the mutant state could be fully reconciled with the *BRCA1*[WT] profile (Figure 5A). These findings suggest that a coordinated, yet limited set of enzyme activity shifts is sufficient to account for most of the metabolic reprogramming that distinguishes *BRCA1*[MUT] from *BRCA1*[WT] ovarian cancer cells.

The way these flux alignments were achieved depended on the pathway. In glycolysis, alignment with the wild-type state required distributed adjustments across multiple connected pathways, consistent with its role as a central hub in cellular metabolism (Figure 5B). By contrast, pyrimidine biosynthesis was largely brought into alignment through local changes within the pathway itself (Figure 5C). Local regulation is observed in other distal pathways too (Figure S10). This contrast reflects an emergent stoichiometric control pattern: core metabolic pathways are modulated by broad network-level interactions, while terminal pathways are locally regulated. Together, these results suggest that extensive, coordinated adjustments to enzyme activity are required to reconcile the metabolic states of *BRCA1*[MUT] and *BRCA1*[WT] cells.

Because metabolism is highly interconnected, there are numerous possible ways to adjust enzyme activity to perturb the *BRCA1* mutant state so that it resembles the wild-type state. Across all possible ways, a consistent group of 145 enzymes always appeared among the top candidates (Figure 5D). These enzymes represent a core set of metabolic regulators that seem essential for understanding how *BRCA1* loss alters ovarian cancer metabolism. Most of them showed lower activity when shifting from the mutant to the wild-type state, which aligns with the observation that *BRCA1* mutant cells grow faster and behave more aggressively,

Figure 5. Enzyme level shifts capture the metabolic differences between *BRCA1*^MUT^ and *BRCA1*^WT^, suggesting a *BRCA1*-linked network-wide reprogramming. (A) The residual flux deviation of the 12 core metabolic pathways progressively decreases as more enzyme fold changes are imposed. Allowing 250 enzymes recapitulates most of the *BRCA1*^WT^ flux profile, with little additional gain when all 877 candidate enzymes are utilized, highlighting a limited set of drivers of metabolic shift. (B) Glycolytic fluxes are redirected from the *BRCA1*^MUT^ to the *BRCA1*^WT^ state through coordinated adjustments across neighboring pathways, illustrating distributed network-wide control. (C) Flux deviations in pyrimidine biosynthesis are corrected predominantly by enzymes within the pathway itself, reflecting localized regulatory control. (D) We identified a core set of 145 enzymes that consistently require a fold change. They are grouped by pathway and categorized by up- or downregulation. The

predominance of downregulated enzymes is consistent with an overall reduction in metabolic activity in $BRCA1^{WT}$ compared to $BRCA1^{MUT}$. (E) Sankey diagram illustrates the relationship between the core set of enzymes and their regulating transcription factors (TFs). A subset of $BRCA1$-interacting TFs exerts widespread influence, suggesting that $BRCA1$ loss propagates through specific TF hubs to drive network-level metabolic rewiring.

typically relying on higher enzyme activity to meet their demands.[86] These core enzymes are distributed across multiple pathways (Figure 5D), but amino acid metabolism is especially enriched. This finding is consistent with our broader observation that disturbances in amino acid metabolism account for a significant portion of the metabolic differences between $BRCA1^{MUT}$ and $BRCA1^{WT}$ cells (Figure 5A).

To link the predicted enzyme-level changes and $BRCA1$-mediated transcriptional regulation, we investigated transcription factors (TFs) known to regulate the expression of the 145 core enzymes. Using the TFLink database[87], we identified 108 TFs regulating 60 of these enzymes through 182 distinct genes (Data S3). Focusing on the top 20 TFs with the broadest regulatory coverage, we identified six (MYC, STAT1, HIF1A, AR, EGR1, TP53) that are known to interact directly with $BRCA1$. A complementary literature search identified three more $BRCA1$-interacting TFs (SP1, RelA, and NFKB1) that engage $BRCA1$ through protein-protein interactions. Specifically, $BRCA1$ physically interacts with SP1, which interferes with SP1's ability to transactivate IGF-IR.[88] This may have novel downstream effects stemming from $BRCA1$ action. $BRCA1$ has also been shown to bind the p65/RelA subunit of NF-κB[89] and act as a transcriptional co-activator. Interestingly, it has been demonstrated that the absence of functional $BRCA1$ leads to an increase in the expression of NF-κB target genes.[89,90]

Collectively, these nine $BRCA1$-associated TFs regulate 46 of the 145 core enzymes, suggesting a possible transcriptional mechanism through which $BRCA1$ could exert control over a significant fraction of the metabolic network (Figure 5E). Notably, MYC and STAT1 could mediate strong effects on the electron transport chain, while glycolysis and several amino acid biosynthesis pathways are also prominently regulated by this TF set.

Although $BRCA1$-linked transcription factors explained only part of the predicted enzyme activity changes, this is not unexpected, given the gaps in current interaction maps, the missing gene annotations for 24 of the 145 enzymes, and indirect influences such as feedback loops or post-transcriptional regulation. Together, these findings point to a broader and more complex regulatory network that connects $BRCA1$ to ovarian cancer metabolism. Importantly, the modeling condenses this complexity into a clear set of enzyme activity shifts that can now be tested experimentally. Approaches such as ^{13}C-tracing, targeted proteomics, or CRISPR-based transcription factor screens provide a practical path to uncover how $BRCA1$ loss drives regulatory and metabolic reprogramming in ovarian cancer.

Discussion

Large-scale kinetic modeling constrained by multi-omics and physicochemical data transcends the steady-state perspective of genome-scale metabolic models, capturing transient, system-wide dynamics in ovarian cancer cells with and without $BRCA1$ loss-of-function. Populations of physiologically consistent kinetic models captured the observed heterogeneity of the two isogenic cell lines, linking flux and metabolite concentration

dynamics. Metabolic control analysis (MCA) identified 28 enzymes exerting the strongest influence over proliferation; 24 of these correspond to market-approved or experimental drugs, validating the framework, while the remaining four represent unexplored metabolic vulnerabilities. Compared to gene knockout simulations in steady-state models, which identified hundreds of potential targets (417 for $BRCA1^{WT}$ and 416 for $BRCA1^{MUT}$, Data S4), MCA identified a smaller, mechanistically interpretable set of enzymes with strong control over proliferation. These results highlight the enhanced resolution and mechanistic depth of kinetic modeling, which captures nonlinear and regulatory effects not accessible through stoichiometry alone.

Simulations of drug therapies captured experimentally observed hallmark responses. The model tracks all intracellular fluxes and metabolite levels, enabling us to monitor how drug therapies impact the entire metabolic network. On a macroscopic scale, the simulations reproduced hallmark behaviors that have been reported experimentally, such as the compensatory activation of mitochondrial respiration following hexokinase inhibition and the ceramide-centered sphingolipid accumulation associated with stress responses. These simulations demonstrate that the models accurately capture known treatment responses, while also providing a better understanding of how therapies alter cancer cell metabolism.

Finally, a minimal set of enzyme fold changes explained the differences in the metabolic states between $BRCA1^{MUT}$ and $BRCA1^{WT}$ physiologies, and we proposed a core set of TFs that are potentially mediated by $BRCA1$ to achieve the observed metabolic shifts. This allows us to focus on a much narrower set of TFs, especially because $BRCA1$ interacts with dozens of transcription factors (TFs) that control hundreds of metabolic genes. According to TFLink[87], $BRCA1$-associated TFs regulate 872 Recon3D genes encoding more than 2,000 enzymes. However, such associations alone cannot reveal which enzyme activity changes are responsible for the distinct metabolic behavior of $BRCA1^{MUT}$ and $BRCA1^{WT}$ cells. This framework could be further extended to explain the enzymatic changes required between healthy cells and tumor cells, thereby deciphering the mechanisms of cancer initiation and progression.

Overall, these results establish kinetic modeling as a dynamic, mechanistic complement to conventional genome-scale models, providing a resource for hypothesis generation, target prioritization, and pharmacological exploration.

This study has several limitations that should be considered when interpreting the results. First, the current models are based on multi-omics data from an isogenic pair of ovarian cancer cell lines ($BRCA1^{WT}$ and $BRCA1^{MUT}$). This design enables isolation of the effects of $BRCA1$ loss but restricts generalizability, as broader validation with patient-derived samples and other tumor contexts may be needed to extend the predictions. Second, although the models incorporate detailed mechanistic rate laws and are constrained by multi-omics and physicochemical data, they do not explicitly account for enzyme regulation through allosteric interactions or signaling cascades, both key components in mammalian physiology. Nonetheless, despite not including these layers of regulation, the models reliably reproduced hallmark drug responses and physiological phenotypes, indicating that much of the observed system behavior can be explained by enzyme abundances and metabolic network structure alone. Finally, the models quantify $BRCA1$-dependent differences in ovarian cancer metabolism across nearly 300 reactions spanning 12 subsystems. In contrast, most prior

studies have focused on glycolytic and oxidative pathways, often describing *BRCA1*-associated shifts simply as enhanced glycolysis. This difference may complicate direct comparison with earlier reports, but it provides an unprecedented mechanistic resolution and generates experimentally testable hypotheses for future work.

Future and implications

Kinetic models provide a platform for generating mechanistic predictions that guide experimental and clinical investigations. Incorporating explicit pharmacokinetic and pharmacodynamic (PK/PD) modules, such as expressions that track drug absorption, distribution, metabolism, and clearance, will enhance the relevance of the predictions, facilitate dose optimization, and support the time-consuming and expensive drug development process.

Integrating patient-specific data could enable personalized models, tailored to each patient's physiology, linking predicted metabolic vulnerabilities to customized drug therapies. Efforts are underway to accelerate model construction and parameterization using machine learning[91–93], streamlining the workflow from data acquisition to actionable predictions.

It is also worth noting that cancer physiology is significantly influenced by oncogenic signaling cascades, which regulate metabolic pathway activity, cell growth, and proliferation. Key signaling pathways frequently activated in tumors, such as the PI3K–AKT–mTORC1 and MYC pathways, increase metabolic flux through numerous pathways to support cell proliferation.[4,5] Metabolites themselves can feed back onto signaling cascades, thereby creating an interconnected network that facilitates tumor progression.[4,5] Despite this, current models remain mostly small-scale or cascade-specific, highlighting the need for a robust workflow to develop dynamic, large-scale kinetic models of signaling that can integrate with metabolism.[94]

Kinetic models also offer the opportunity to capture how the dynamic availability of nutrients shapes cancer metabolism. Cancer cells utilize various carbon sources, including glucose, lactate, amino acids, and proteins, to sustain proliferation.[5,57] Modeling how fluxes adjust to shifts in primary carbon sources or dietary regimes could reveal previously unrecognized metabolic vulnerabilities and inform the development of novel pharmacological interventions.[95,96]

Simulating interactions between cancer cells and the tumor microenvironment (TME) could further clarify context-dependent metabolic adaptations. Numerous, yet incompletely understood, mechanisms enable cancer cells to exploit their neighboring cells to increase their chances of survival.[40,97,98] Cancer cells compete with immune cells, such as activated T cells, for key nutrients such as glucose and amino acids.[40] They also co-opt the metabolism of neighboring stromal cells, which secrete numerous metabolites, including lactate and glutamine.[40] Incorporating the metabolism of these primary TME cell types into future kinetic models would open new opportunities to design therapies that target tumor metabolism without compromising immune function.[41,47]

Conclusions

Near-genome-scale kinetic models constrained by tissue- or patient-specific multi-omics data extend conventional genome-scale frameworks by capturing the dynamic behavior of metabolic networks. The ovarian cancer models presented here, validated against diverse

experimental and clinical observations, provide a robust platform for mechanistic interrogation, hypothesis generation, and rational therapeutic design. They hold strong potential for advancing precision oncology and personalized medicine by enabling dynamic, network-level insights into cancer metabolism.

Materials and Methods

Reduced metabolic network and multi-omics data integration

The first step in our ORACLE workflow[99] for constructing kinetic models involves defining the reduced metabolic network that contains the pathways of interest for a given biological question. The next step is the integration of available omics data by applying methods adapted for integrating transcriptomics (REMI[31], iMAT[100,30,101], INIT[102], ΔFBA[103]), metabolomics (TFA[104], INTEGRATE[105]), and fluxomics[106] data. In this study, we employed the reduced stoichiometric model of ovarian cancer metabolism with the integrated omics data developed by Masid et al. (unpublished work). Starting from the Recon3D[50] genome-scale model of human metabolism, those authors applied the redGEM[107], lumpGEM[108], and redGEMX[48] tools to generate a reduced network tailored to ovarian cancer. They then integrated transcriptomic data from $BRCA1^{WT}$ and $BRCA1^{MUT}$ ovarian cancer cell lines using the iMSEA workflow (Masid et al., unpublished work), which consistently merges transcriptomics, metabolomics, and fluxomics information through TFA-[104] and REMI[31]-based constraints. The resulting stoichiometric model comprised 2,369 reactions and 974 metabolites distributed across eight cellular compartments and 64 metabolic subsystems.

In the present study, we further integrated experimentally measured metabolite levels of ATP, ADP, and AMP from Park et al.[109] We additionally imposed a basal nonzero flux constraint (10^{-6} mmol·gDW^{-1}·h^{-1}) for all reactions to enforce minimal enzyme activity and ensure global network connectivity. As a result, some reactions lost their ability to carry flux and were removed from the models, yielding final network sizes of 2,359 reactions for the $BRCA1^{WT}$ physiology and 2,344 for the $BRCA1^{MUT}$ physiology.

Determining reference metabolic steady states

Many intracellular reactions can operate in both directions. Even after integrating multi-omics and physicochemical data along with expert knowledge, a significant number of reactions remained bidirectional (BDRs), resulting in considerable flux uncertainty. To reduce the feasible solution space, we applied an iterative, criteria-driven procedure to assign reaction directionalities:

1. Exchange-related reactions. For BDRs connecting intracellular metabolites to the extracellular medium, directionality was set based on the growth medium: metabolites absent from the medium were assumed to be secreted.
2. Futile-cycle elimination. We identified reaction pairs whose opposing directions could result in artificial ATP dissipation, e.g., ATP-driven import opposed by passive export. When such futile cycles were identified, we aligned their directions to eliminate them and maintain energetic consistency.

3. Flexibility-based selection. For the remaining BDRs, we performed flux variability analysis (FVA) and, wherever possible, assigned the direction with the larger feasible range, thereby favoring greater flexibility.

It is important to note that assigning a directionality to a reaction does not restrict it to operate exclusively in that direction. The specified directionality defines only the reference orientation of fluxes at the sampled steady state. For reactions governed by reversible kinetic rate laws (e.g., Reversible Michaelis-Menten, Convenience kinetics[110]), the net flux can reverse when the substrate-to-product concentration ratio changes sufficiently. Thus, although a reference direction is chosen for computing the steady-state solution, reaction directionalities remain dynamically adaptable to environmental or pharmacological perturbations.

Once the reference reaction directionalities were established, we sampled steady states consistent with the integrated experimental constraints, thereby reproducing the observed metabolic heterogeneity. To achieve this, we utilized the Artificial Centering Hit-and-Run (ACHR) sampler implemented in the pyTFA toolbox[33] to generate 5,000 steady-state samples with a thinning step of 500. Each sample included flux and metabolite concentration values across the network, as well as thermodynamic information such as ΔG and ΔG^o.

Generation of the kinetic model scaffold

The reduced metabolic network served as the scaffold for building the kinetic model. Each reaction was assigned an appropriate rate law mechanism from a comprehensive library implemented in the SKiMpy toolbox.[53] By default, SKiMpy assigns the enzyme mechanism based on the reaction's stoichiometry. The produced kinetic model consists of 716 mass balances (and 25 pool-conservation equations for dependent metabolites) governed by 2,131 rate laws. In total, the model required values for 9,594 kinetic parameters, including 7,463 Michaelis-Menten constants (K_m), 2,131 maximum velocity constants (v_{max}), and 2,193 thermodynamic equilibrium constants (K_{eq}).

Generation of a population of kinetic models

To generate kinetic parameter values, we applied the ORACLE workflow.[99] Using a steady-state sample of fluxes and concentrations, we generated 100 independent parameter sets, each of which was consistent with the imposed steady state. To evaluate their physiological relevance, we analyzed the local dynamics of each model by computing the eigenvalues of the system's Jacobian. We retained models based on physiologically relevant dynamics: models were considered physiologically relevant if their dominant time constant (τ_d) was at least three times faster than the doubling time of the modeled cancer cell.[92,93,111,112]

$$\tau_d = \frac{1}{\min|\mathrm{Re}(\lambda_i)|} \text{ (Equation 1)}$$

Because some steady-state samples did not yield physiologically relevant kinetic models, we repeated this process for each physiology separately until the desired distributions were obtained, as described in the next section. The final number of kinetic models was 13,731 for the *BRCA1*[WT] physiology, derived from 556 steady-state samples, and 6,732 for the *BRCA1*[MUT] physiology, derived from 396 steady-state samples.

Metabolic Control Analysis

Metabolic Control Analysis (MCA) was performed using the SKiMpy toolbox to quantify how changes in enzyme activity propagate throughout the network. The sensitivity of fluxes and metabolite concentrations to enzyme perturbations is expressed through the control coefficients:[55,113]

$$FCC_i^j = \frac{\partial ln v_i}{\partial ln E_j} = \frac{E_j}{v_i}\frac{\partial v_i}{\partial E_j} \text{ (Equation 2)}$$

$$CCC_k^j = \frac{\partial ln x_k}{\partial ln E_j} = \frac{E_j}{x_k}\frac{\partial k}{\partial E_j} \text{ (Equation 3)}$$

where FCC_i^j and CCC_i^j denote the flux and concentration control coefficient, respectively. v_i is the flux of reaction i, E_j the enzyme concentration j, and x_k is the concentration of metabolite k at the reference steady state. A positive control coefficient indicates that the flux or concentration is expected to increase with an increase in enzyme concentration. For example, the flux control coefficient of the growth rate with respect to triokinase (TRIOK) for the *BRCA1*^WT physiology is 0.72 (Figure 1A), meaning that a 2-fold increase in TRIOK concentration would result in an approximately 1.65 times increase in the growth rate relative to the reference state.

MCA predictions are accurate only for small perturbations around the reference state. Nevertheless, the magnitude and sign of the control coefficients provide valuable insight into enzymes that affect cancer proliferation. The actual impact of larger perturbations was further assessed by performing drug simulations. Different steady-state samples and kinetic parameter values will yield different values for the FCCs and CCCs. In the analysis, we report the population averages and the interquartile ranges.

To evaluate the effect of top enzymes on the biomass precursors, we calculated their CCCs (Eq. 3) and the control coefficients of their synthesis rate:

$$FCC_{synthesis}^j = \frac{\sum_n v_{n^{ref}}FCC_n^j}{\sum_n v_{n^{ref}}} \text{ (Equation 4)}$$

where n denotes the number of reactions that are producing the biomass building block, and the $v_{n^{ref}}$ is the flux value at the reference state.

For every kinetic model that satisfied the local dynamic criteria, we calculated all the flux and concentration control coefficients across the network. To identify statistically significant trends, we evaluated the convergence of the distributions of the flux control coefficients of growth rate with respect to each enzyme. To ensure robustness, outlier control coefficient values were removed using the interquartile range (IQR) method, applied independently to each enzyme's flux control coefficient distribution. Convergence was considered achieved when the first three moments (mean, variance, skewness) of each distribution changed by less than 10% upon addition of a new kinetic model to the population. This allowed us to discuss the averages and the standard deviations of the key enzymes for cancer growth, which were picked based on the absolute average values of the control coefficients. For these key enzymes, we further calculated the average flux and concentration control coefficients for all network variables (fluxes or concentrations), allowing us to obtain a system-level understanding of their metabolic effects.

DrugBank data extraction

For each of the 28 drug targets predicted by the MCA workflow, we retrieved associated drugs based on information from DrugBank (version 5.1.13, released on January 2, 2025).[58] The 28 enzymes were mapped to their Uniprot identifiers, and checked which drugs are associated with them. For each drug, we also extracted its ATC code, approval status (approved or experimental), and documented mode of action, whenever available.

In silico drug administration simulations

We performed the drug administration simulations on a representative subset of the original kinetic model population, selected through a stratified sampling approach. Analogous to clinical studies, where participants are recruited to capture variability across defined strata, we divided the models according to their initial glycolytic and mitochondrial ATP yields. Glycolytic ATP yield is calculated as the ratio of the sum of PYK and PGK fluxes to the glucose uptake rate. Similarly, the mitochondrial ATP yield was calculated as the ratio of the mitochondrial ATP synthase (ATPS4mi) flux to the glucose uptake rate. We then divided the observed ranges into three strata for each yield, resulting in nine possible combinations, and randomly selected 50 steady-state samples in total, spread across the nine metabolic subtypes. Importantly, because the number of steady-state samples varied across strata, we ensured that the 50 representative metabolic states reflected the relative density of models within each stratum. This resulted in 972 kinetic models for the *BRCA1*[WT] physiology and 1,134 for the *BRCA1*[MUT] physiology.

To simulate the effects of anticancer drugs, we assumed that the drug action directly impacts the enzymatic activity of the target enzyme. This effect was modeled by introducing an additional ODE that exponentially reduces the enzyme concentration toward a specified target value:

$$\frac{dE}{dt} = k\left(E_{target} - E\right) \text{ (Equation 5)}$$

where k is the rate constant of the effect, and E_{target} is the target enzyme concentration achieved due to the drug action. Since the models are formulated in terms of V_{max} values rather than explicit enzyme concentrations, we assumed an initial enzyme concentration of 1 M. Under this assumption, the catalytic rate constant, k_{cat}, is equal to the sampled V_{max} value.

We subsequently integrated all kinetic models using as initial conditions the steady state sample around which each model was built. Numerical integration was performed using CVODE[114] over a simulation time of 25 model days, providing ample time for the system to reach a new steady state. In practice, most models settled into a new metabolic state within one doubling time. This procedure yielded the time evolution of all the concentrations and fluxes in the metabolic network across the subpopulation of kinetic models.

To estimate EC_{50} values for each drug, we performed drug administration simulations by progressively decreasing E_{target} from 0.999 (effectively no perturbation) to 10^{-9} (complete inhibition). Cell viability was calculated as the ratio of the initial to the final growth rate. The resulting data points were used to fit a 4-parameter logistic model, following the format described by Corsello et al.[74]

$$Cell\ viability(a) = E_{\infty} + \frac{1 - E_{\infty}}{1 + e^{HS(a - EC_{50})}} \text{ (Equation 6)}$$

where a denotes the target enzyme concentration (E_{target}), E_∞ the residual effect when the enzyme is completely inhibited, EC_{50} the final enzyme concentration that produces a half-maximal effect (the cell viability will be halfway between 1 and E_∞), and HS is the hill slope that determines the steepness of the dose-response curves.

This analysis across all kinetic models yielded a distribution of pharmacodynamic parameters, capturing the expected ranges of drug-induced growth inhibition. The reported values for each parameter represent the average and standard deviation after removing outliers, which were excluded using the interquartile range method.

Volcano plots

To assess the system-level effects of drug target inhibition, we created volcano plots that show the significantly perturbed fluxes and concentrations within the network. For each model, we calculated $\log_2$ fold changes in metabolite concentrations and $\log_{10}$ fold changes in reaction fluxes between the initial and final metabolic states. A one-sample t-test was applied across all the models to identify significantly perturbed concentrations and fluxes, and p-values were corrected for multiple testing with the Benjamini–Hochberg method.[115] To visualize the distance of each metabolite and flux from the target enzyme, we utilized the adjacency matrices generated by the redGEM workflow[107] and calculated the minimum distance between a reaction or metabolite from the metabolites participating in the inhibited reaction.

Pathway enrichment analysis

To compile and compare the changes in metabolic pathways upon drug treatment, we conducted pathway enrichment analysis. Metabolites with an absolute $\log_2$ concentration fold change ≥ 2 and q ≤ 0.01 were considered significantly perturbed and were mapped to pathways through the Recon3D GEM. We calculated the fraction of significantly altered metabolites within each metabolic pathway and used a hypergeometric statistical test to determine whether the number of perturbed metabolites exceeded what would be expected by chance. To determine whether each pathway was predominantly up- or downregulated following treatment, we examined both the direction and magnitude of metabolite fold changes within each pathway. In nearly all cases, the majority of metabolites and fluxes within a pathway changed in the same direction, allowing us to assign an overall qualitative trend of up- or downregulation. These summarized effects were then visualized on a simplified metabolic map, providing an intuitive overview of how drug treatment reshaped intracellular metabolism.

Comparison of *BRCA1*^{WT} and *BRCA1*^{MUT} phenotypes

To identify enzyme fold changes that will minimize flux deviations between *BRCA1*^WT and *BRCA1*^MUT phenotypes, we used the NRA formulation[116,117] implemented in the NOMAD framework.[111] The NRA requires as inputs the reference *BRCA1*^MUT flux state (serving as the initial metabolic state) and a kinetic parameter set (used to calculate the control coefficients of the network), which together enable the estimation of the new metabolic state after applying specified enzyme fold changes. For comparison, the target flux state was defined from *BRCA1*^WT physiology.

We calculated the flux deviations for each pair of *BRCA1*[WT] and *BRCA1*[MUT] steady states that were sampled for the MCA workflow, resulting in a distribution of deviation values. From this distribution, we selected the representative pair that was closest to the average of the distribution.

We then selected a physiologically relevant kinetic parameter set and used it to construct a kinetic model centered on the selected *BRCA1*[MUT] steady-state sample. A series of robustness tests was performed to determine whether the chosen kinetic parameter set was suitable for the subsequent analysis. Specifically, we randomly perturbed each metabolite concentration by up to ±10% and simulated the system's response. A response was considered robust if the model returned to its original flux and concentration profile within one doubling time. This procedure was performed 100 times, and in all cases the model was able to return to its original steady state. Having passed this test, the kinetic model was deemed robust and was subsequently used in the NRA formulation.

NRA was used to predict the new flux distribution of the metabolic network following perturbations in enzyme activity. For this purpose, we optimized the fluxes of 298 reactions belonging to the 12 core pathways, representing key features of cancer metabolism activity. We allowed 877 enzymes to vary in activity, comprising all enzymes catalyzing biotransformations in our reduced model and 112 enzymes catalyzing the transport of metabolic species, and whose genes exhibited significant expression fold changes between the ovarian cancer cell lines. Enzyme upregulation was constrained to a maximum 50-fold increase, based on the maximum gene expression fold change observed in our reference gene expression data, while downregulation was allowed to effectively knock out the enzyme. To guide NRA toward a solution producing a flux distribution closest to the *BRCA1*[WT] physiology, we used the following objective function:

$$\min \sum_i \left| lnv_{NRA}^i - lnv_{WT}^i \right| \text{ (Equation 7)}$$

where v_{NRA} is the new flux solution produced by NRA, and v_{WT} is the reference *BRCA1*[WT] flux profile. The fluxes are represented on a log scale, as NRA operates in a log-linear space. We introduced additional constraints to limit the number of enzymes allowed to have a fold change, enabling the identification of the minimal set of perturbations required to minimize the distance between the *BRCA1*[WT] and *BRCA1*[MUT] flux distributions. This analysis revealed that applying a fold change to 250 of the 877 enzymes produced results nearly indistinguishable from those obtained when all enzymes were permitted to vary, indicating that a subset of enzymes exerts dominant control over the phenotypic differences between the two physiologies.

To quantify the contribution of NRA-predicted enzyme fold changes to pathway-level alignment between physiologies, we defined pathway effect scores. For a given target pathway, the effect from a source (contributing) pathway was calculated as:

$$Effect = \sum_i \sum_j FCC_i^j \ln\left(\frac{E_j}{E_{j_{ref}}}\right) \text{ (Equation 8)}$$

with *i* indexing all fluxes in the target pathway and *j* indexing all enzymes in the source pathway. Positive effect scores indicate that enzymes in the source pathway collectively push upward fluxes in the target pathway, whereas negative scores indicate a downward influence. Central pathways, such as glycolysis, typically integrate contributions by multiple source

pathways, whereas biosynthetic pathways, like pyrimidine synthesis, are predominantly driven by enzymes within the same pathway.

Identifying a minimal core set of enzyme fold changes

Due to the strong interconnectivity of the metabolic network, multiple combinations of 250 perturbed enzymes could reproduce similar alignment between the two phenotypes. To identify the enzymes that were consistently required, we developed an iterative workflow. Starting with one solution set of 250 enzymes, we systematically blocked the fold change of each enzyme and re-optimized for Equation 7 using the NRA formulation. If the new solution achieved an objective value within 5% of the original, the removed enzyme was considered non-essential for reproducing the flux alignment. Conversely, if removing the enzyme prevented the optimization from meeting this criterion, the enzyme was deemed necessary and considered active across all alternative solutions. This procedure identified a robust core of 145 enzymes that consistently appeared across solutions, highlighting them as key determinants of how BRCA1 loss reshapes ovarian cancer metabolism.

TFLink data extraction

We used the TFLink database[87] to retrieve information on transcription factor (TF) interactions with the metabolic genes of the network, as well as with *BRCA1*. We used information flagged as *small-scale evidence* to ensure that the interactions reported are well-characterized.

Acknowledgements

This work was supported by funding from the Swiss National Science Foundation grants 200021_188623 and CRSII5_198543, the European Union's Horizon 2020 research and innovation programme under grant agreement 814408, and the Ecole Polytechnique Fédérale de Lausanne (EPFL).

Data and Code availability

The data supporting this study's findings are publicly available in the Zenodo repository at https://zenodo.org/records/17304777 and the links therein.

The code needed to reproduce this study's results is publicly available at https://github.com/EPFL-LCSB/human-cancer-kinetic-models. The models were built and analyzed using the SKiMpy toolbox, which can be accessed at https://github.com/EPFL-LCSB/skimpy. NRA is implemented within the NOMAD workflow, which can be accessed at https://github.com/EPFL-LCSB/NOMAD.

Abbreviations

All metabolite, reaction, and enzyme abbreviations used in this study are consistent with those defined in the human genome-scale reconstruction (Recon3D) and are retained in the curated kinetic models. The full list of abbreviations can be accessed directly within the model files

provided with this publication. The same standardized abbreviations are also available through the BiGG/VMH database (https://vmh.life/#home).

Contributions

Conceptualization and methodology, I.T., M.M., V.H., and L.M.; software, I.T.; data curation, I.T.; data analysis, I.T., M.M., V.H., and L.M.; visualization, I.T. and L.M.; writing — original draft, I.T.; writing — review & editing, I.T., M.M., V.H., and L.M.; supervision, M.M., V.H., and L.M.; funding acquisition, V.H., and L.M.

Declaration of interests

The authors declare no financial or commercial conflict of interest.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used ChatGPT and Grammarly to improve the readability and language of the manuscript. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the published article.

Supplemental information

Document S1. Figures S1-S10 and Tables S1-S2

Data S1. Predicted metabolic drug targets and their associated drugs, as cataloged in DrugBank

Data S2. Annotated list of antimetabolites cataloged in DrugBank that were not predicted by the models

Data S3. Core set of enzyme fold changes predicted by the NRA formulation, and their connections to transcription factors and BRCA1

Data S4. Essential genes identified for both physiologies

References

1. Hanahan, D., and Weinberg, R.A. (2000). The Hallmarks of Cancer. Cell *100*, 57–70. https://doi.org/10.1016/S0092-8674(00)81683-9.

2. Hanahan, D., and Weinberg, R.A. (2011). Hallmarks of Cancer: The Next Generation. Cell *144*, 646–674. https://doi.org/10.1016/j.cell.2011.02.013.

3. Hanahan, D. (2022). Hallmarks of Cancer: New Dimensions. Cancer Discovery *12*, 31–46. https://doi.org/10.1158/2159-8290.CD-21-1059.

4. Martínez-Reyes, I., and Chandel, N.S. (2021). Cancer metabolism: looking forward. Nat Rev Cancer *21*, 669–680. https://doi.org/10.1038/s41568-021-00378-6.

5. DeBerardinis, R.J., and Chandel, N.S. (2016). Fundamentals of cancer metabolism. Science Advances *2*, e1600200. https://doi.org/10.1126/sciadv.1600200.

6. Intlekofer, A.M., and Finley, L.W.S. (2019). Metabolic signatures of cancer cells and stem cells. Nat Metab *1*, 177–188. https://doi.org/10.1038/s42255-019-0032-0.

7. Bergers, G., and Fendt, S.-M. (2021). The metabolism of cancer cells during metastasis. Nat Rev Cancer *21*, 162–180. https://doi.org/10.1038/s41568-020-00320-2.

8. Stine, Z.E., Schug, Z.T., Salvino, J.M., and Dang, C.V. (2022). Targeting cancer metabolism in the era of precision oncology. Nat Rev Drug Discov *21*, 141–162. https://doi.org/10.1038/s41573-021-00339-6.

9. Faubert, B., Tasdogan, A., Morrison, S.J., Mathews, T.P., and DeBerardinis, R.J. (2021). Stable isotope tracing to assess tumor metabolism in vivo. Nat Protoc *16*, 5123–5145. https://doi.org/10.1038/s41596-021-00605-2.

10. Lewis, N.E., Nagarajan, H., and Palsson, B.O. (2012). Constraining the metabolic genotype–phenotype relationship using a phylogeny of in silico methods. Nat Rev Microbiol *10*, 291–305. https://doi.org/10.1038/nrmicro2737.

11. Patti, G.J., Yanes, O., and Siuzdak, G. (2012). Metabolomics: the apogee of the omics trilogy. Nat Rev Mol Cell Biol *13*, 263–269. https://doi.org/10.1038/nrm3314.

12. Yizhak, K., Chaneton, B., Gottlieb, E., and Ruppin, E. (2015). Modeling cancer metabolism on a genome scale. Molecular Systems Biology *11*, 817. https://doi.org/10.15252/msb.20145307.

13. Nielsen, J. (2017). Systems Biology of Metabolism: A Driver for Developing Personalized and Precision Medicine. Cell Metabolism *25*, 572–579. https://doi.org/10.1016/j.cmet.2017.02.002.

14. Bordbar, A., Monk, J.M., King, Z.A., and Palsson, B.O. (2014). Constraint-based models predict metabolic and associated cellular functions. Nature Reviews Genetics *15*, 107–120. https://doi.org/10.1038/nrg3643.

15. Orth, J.D., Thiele, I., and Palsson, B.Ø. (2010). What is flux balance analysis? Nat Biotechnol *28*, 245–248. https://doi.org/10.1038/nbt.1614.

16. Gu, C., Kim, G.B., Kim, W.J., Kim, H.U., and Lee, S.Y. (2019). Current status and applications of genome-scale metabolic models. Genome Biol *20*, 121–121. https://doi.org/10.1186/s13059-019-1730-3.

17. Stanway, R.R., Bushell, E., Chiappino-Pepe, A., Roques, M., Sanderson, T., Franke-Fayard, B., Caldelari, R., Golomingi, M., Nyonda, M., Pandey, V., et al. (2019). Genome-Scale Identification of Essential Metabolic Processes for Targeting the Plasmodium Liver Stage. Cell *179*, 1112-1128.e26. https://doi.org/10.1016/j.cell.2019.10.030.

18. Mardinoglu, A., Wu, H., Bjornson, E., Zhang, C., Hakkarainen, A., Räsänen, S.M., Lee, S., Mancina, R.M., Bergentall, M., Pietiläinen, K.H., et al. (2018). An Integrated Understanding of the Rapid Metabolic Benefits of a Carbohydrate-Restricted Diet on Hepatic Steatosis in Humans. Cell Metabolism *27*, 559-571.e5. https://doi.org/10.1016/j.cmet.2018.01.005.

19. Shoaie, S., Ghaffari, P., Kovatcheva-Datchary, P., Mardinoglu, A., Sen, P., Pujos-Guillot, E., de Wouters, T., Juste, C., Rizkalla, S., Chilloux, J., et al. (2015). Quantifying Diet-Induced Metabolic Changes of the Human Gut Microbiome. Cell Metabolism *22*, 320–331. https://doi.org/10.1016/j.cmet.2015.07.001.

20. Thiele, I., Sahoo, S., Heinken, A., Hertel, J., Heirendt, L., Aurich, M.K., and Fleming, R.M. (2020). Personalized whole-body models integrate metabolism, physiology, and the gut microbiome. Molecular Systems Biology *16*, e8982. https://doi.org/10.15252/msb.20198982.

21. Zaunseder, E., Mütze, U., Okun, J.G., Hoffmann, G.F., Kölker, S., Heuveline, V., and Thiele, I. (2024). Personalized metabolic whole-body models for newborns and infants predict growth and biomarkers of inherited metabolic diseases. Cell Metabolism *36*, 1882-1897.e7. https://doi.org/10.1016/j.cmet.2024.05.006.

22. Morotti, M., Grimm, A.J., Hope, H.C., Arnaud, M., Desbuisson, M., Rayroux, N., Barras, D., Masid, M., Murgues, B., Chap, B.S., et al. (2024). PGE2 inhibits TIL expansion by disrupting IL-2 signalling and mitochondrial function. Nature *629*, 426–434. https://doi.org/10.1038/s41586-024-07352-w.

23. Agren, R., Mardinoglu, A., Asplund, A., Kampf, C., Uhlen, M., and Nielsen, J. (2014). Identification of anticancer drugs for hepatocellular carcinoma through personalized genome-scale metabolic modeling. Molecular Systems Biology *10*, 721. https://doi.org/10.1002/msb.145122.

24. Folger, O., Jerby, L., Frezza, C., Gottlieb, E., Ruppin, E., and Shlomi, T. (2011). Predicting selective drug targets in cancer through metabolic networks. Molecular Systems Biology *7*, 501. https://doi.org/10.1038/msb.2011.35.

25. Lim, J., Jung, H.D., Park, S.Y., Jeon, M., Kim, D.S., Cho, R., Han, D., Ryu, H.S., Kim, Y., and Kim, H.U. (2025). Genome-scale knockout simulation and clustering analysis of drug-resistant breast cancer cells reveal drug sensitization targets. Proc. Natl. Acad. Sci. U.S.A. *122*, e2425384122. https://doi.org/10.1073/pnas.2425384122.

26. Aurich, M.K., Fleming, R.M.T., and Thiele, I. (2017). A systems approach reveals distinct metabolic strategies among the NCI-60 cancer cell lines. PLOS Computational Biology *13*, e1005698. https://doi.org/10.1371/journal.pcbi.1005698.

27. Meeson, K.E., and Schwartz, J.-M. (2024). Constraint-based modelling predicts metabolic signatures of low and high-grade serous ovarian cancer. npj Syst Biol Appl *10*, 96. https://doi.org/10.1038/s41540-024-00418-5.

28. Gustafsson, J., Roshanzamir, F., Hagnestål, A., Patel, S.M., Daudu, O.I., Becker, D.F., Robinson, J.L., and Nielsen, J. (2024). Metabolic collaboration between cells in the tumor microenvironment has a negligible effect on tumor growth. The Innovation *5*, 100583. https://doi.org/10.1016/j.xinn.2024.100583.

29. Bidkhori, G., Benfeitas, R., Klevstig, M., Zhang, C., Nielsen, J., Uhlen, M., Boren, J., and Mardinoglu, A. (2018). Metabolic network-based stratification of hepatocellular carcinoma reveals three distinct tumor subtypes. Proceedings of the National Academy of Sciences *115*, E11874–E11883. https://doi.org/10.1073/pnas.1807305115.

30. Zur, H., Ruppin, E., and Shlomi, T. (2010). iMAT: an integrative metabolic analysis tool. Bioinformatics *26*, 3140–3142. https://doi.org/10.1093/bioinformatics/btq602.

31. Pandey, V., Hadadi, N., and Hatzimanikatis, V. (2019). Enhanced flux prediction by integrating relative expression and relative metabolite abundance into thermodynamically consistent metabolic models. PLoS Comput Biol *15*, e1007036. https://doi.org/10.1371/journal.pcbi.1007036.

32. Salvy, P., and Hatzimanikatis, V. (2020). The ETFL formulation allows multi-omics integration in thermodynamics-compliant metabolism and expression models. Nat Commun *11*, 30. https://doi.org/10.1038/s41467-019-13818-7.

33. Salvy, P., Fengos, G., Ataman, M., Pathier, T., Soh, K.C., and Hatzimanikatis, V. (2019). pyTFA and matTFA: a Python package and a Matlab toolbox for Thermodynamics-based Flux Analysis. Bioinformatics *35*, 167–169. https://doi.org/10.1093/bioinformatics/bty499.

34. Chen, Y., Gustafsson, J., Tafur Rangel, A., Anton, M., Domenzain, I., Kittikunapong, C., Li, F., Yuan, L., Nielsen, J., and Kerkhoven, E.J. (2024). Reconstruction, simulation and analysis of enzyme-constrained metabolic models using GECKO Toolbox 3.0. Nat Protoc *19*, 629–667. https://doi.org/10.1038/s41596-023-00931-7.

35. Toumpe, I., Choudhury, S., Hatzimanikatis, V., and Miskovic, L. (2025). The Dawn of High-Throughput and Genome-Scale Kinetic Modeling: Recent Advances and Future Directions. ACS Synth. Biol. https://doi.org/10.1021/acssynbio.4c00868.

36. Bordbar, A., McCloskey, D., Zielinski, D.C., Sonnenschein, N., Jamshidi, N., and Palsson, B.O. (2015). Personalized Whole-Cell Kinetic Models of Metabolism for Discovery in Genomics and Pharmacodynamics. Cell Systems *1*, 283–292. https://doi.org/10.1016/j.cels.2015.10.003.

37. Schuhknecht, L., Ortmayr, K., Jänes, J., Bläsi, M., Panoussis, E., Bors, S., Dorčáková, T., Fuhrer, T., Beltrao, P., and Zampieri, M. (2025). A human metabolic map of pharmacological perturbations reveals drug modes of action. Nat Biotechnol, 1–13. https://doi.org/10.1038/s41587-024-02524-5.

38. Cherkaoui, S., Durot, S., Bradley, J., Critchlow, S., Dubuis, S., Masiero, M.M., Wegmann, R., Snijder, B., Othman, A., Bendtsen, C., et al. (2022). A functional analysis of 180 cancer cell lines reveals conserved intrinsic metabolic programs. Molecular Systems Biology *18*, e11033. https://doi.org/10.15252/msb.202211033.

39. Sinkala, M., Mulder, N., and Patrick Martin, D. (2019). Metabolic gene alterations impact the clinical aggressiveness and drug responses of 32 human cancers. Commun Biol *2*, 414. https://doi.org/10.1038/s42003-019-0666-1.

40. Elia, I., and Haigis, M.C. (2021). Metabolites and the tumour microenvironment: from cellular mechanisms to systemic metabolism. Nat Metab *3*, 21–32. https://doi.org/10.1038/s42255-020-00317-z.

41. Zou, W., and Green, D.R. (2023). Beggars banquet: Metabolism in the tumor immune microenvironment and cancer therapy. Cell Metabolism *35*, 1101–1113. https://doi.org/10.1016/j.cmet.2023.06.003.

42. Pillai, M., Hojel, E., Jolly, M.K., and Goyal, Y. (2023). Unraveling non-genetic heterogeneity in cancer with dynamical models and computational tools. Nat Comput Sci *3*, 301–313. https://doi.org/10.1038/s43588-023-00427-0.

43. Jaiswal, A., and Singh, R. (2024). A negative feedback loop underlies the Warburg effect. npj Syst Biol Appl *10*, 55. https://doi.org/10.1038/s41540-024-00377-x.

44. Liberti, M.V., Dai, Z., Wardell, S.E., Baccile, J.A., Liu, X., Gao, X., Baldi, R., Mehrmohamadi, M., Johnson, M.O., Madhukar, N.S., et al. (2017). A Predictive Model for Selective Targeting of the Warburg Effect through GAPDH Inhibition with a Natural Product. Cell Metabolism *26*, 648-659.e8. https://doi.org/10.1016/j.cmet.2017.08.017.

45. Marín-Hernández, Á., and Saavedra, E. (2023). Metabolic control analysis as a strategy to identify therapeutic targets, the case of cancer glycolysis. Biosystems *231*, 104986. https://doi.org/10.1016/j.biosystems.2023.104986.

46. Lee, W.D., Weilandt, D.R., Liang, L., MacArthur, M.R., Jaiswal, N., Ong, O., Mann, C.G., Chu, Q., Hunter, C.J., Ryseck, R.-P., et al. (2025). Lactate homeostasis is maintained through regulation of glycolysis and lipolysis. Cell Metabolism *0*. https://doi.org/10.1016/j.cmet.2024.12.009.

47. Bruand, M., Barras, D., Mina, M., Ghisoni, E., Morotti, M., Lanitis, E., Fahr, N., Desbuisson, M., Grimm, A., Zhang, H., et al. (2021). Cell-autonomous inflammation of BRCA1-deficient ovarian cancers drives both tumor-intrinsic immunoreactivity and immune resistance via STING. Cell Reports *36*. https://doi.org/10.1016/j.celrep.2021.109412.

48. Masid, M., Ataman, M., and Hatzimanikatis, V. (2020). Analysis of human metabolism by reducing the complexity of the genome-scale models using redHUMAN. Nat Commun *11*, 2821. https://doi.org/10.1038/s41467-020-16549-2.

49. Küken, A., Wendering, P., Langary, D., and Nikoloski, Z. (2021). A structural property for reduction of biochemical networks. Sci Rep *11*, 17415. https://doi.org/10.1038/s41598-021-96835-1.

50. Brunk, E., Sahoo, S., Zielinski, D.C., Altunkaya, A., Dräger, A., Mih, N., Gatto, F., Nilsson, A., Preciat Gonzalez, G.A., Aurich, M.K., et al. (2018). Recon3D enables a three-dimensional view of gene variation in human metabolism. Nat Biotechnol *36*, 272–281. https://doi.org/10.1038/nbt.4072.

51. Simões, R.V., Serganova, I.S., Kruchevsky, N., Leftin, A., Shestov, A.A., Thaler, H.T., Sukenick, G., Locasale, J.W., Blasberg, R.G., Koutcher, J.A., et al. (2015). Metabolic Plasticity of Metastatic Breast Cancer Cells: Adaptation to Changes in the Microenvironment. Neoplasia *17*, 671–684. https://doi.org/10.1016/j.neo.2015.08.005.

52. Chen, P.-H., Cai, L., Huffman, K., Yang, C., Kim, J., Faubert, B., Boroughs, L., Ko, B., Sudderth, J., McMillan, E.A., et al. (2019). Metabolic Diversity in Human Non-Small Cell Lung Cancer Cells. Molecular Cell *76*, 838-851.e5. https://doi.org/10.1016/j.molcel.2019.08.028.

53. Weilandt, D.R., Salvy, P., Masid, M., Fengos, G., Denhardt-Erikson, R., Hosseini, Z., and Hatzimanikatis, V. (2023). Symbolic kinetic models in python (SKiMpy): intuitive modeling of large-scale biological kinetic models. Bioinformatics *39*, btac787. https://doi.org/10.1093/bioinformatics/btac787.

54. Kacser, H., and Burns, J.A. (1973). The control of flux. Symp Soc Exp Biol *27*, 65–104.

55. Hatzimanikatis, V., and Bailey, J.E. (1996). MCA has more to say. J Theor Biol *182*, 233–242. https://doi.org/10.1006/jtbi.1996.0160.

56. Hu, W., Zhang, C., Wu, R., Sun, Y., Levine, A., and Feng, Z. (2010). Glutaminase 2, a novel p53 target gene regulating energy metabolism and antioxidant function. Proceedings of the National Academy of Sciences *107*, 7455–7460. https://doi.org/10.1073/pnas.1001006107.

57. Martinez-Outschoorn, U.E., Peiris-Pagés, M., Pestell, R.G., Sotgia, F., and Lisanti, M.P. (2017). Cancer metabolism: a therapeutic perspective. Nat Rev Clin Oncol *14*, 11–31. https://doi.org/10.1038/nrclinonc.2016.60.

58. Knox, C., Wilson, M., Klinger, C.M., Franklin, M., Oler, E., Wilson, A., Pon, A., Cox, J., Chin, N.E. (Lucy), Strawbridge, S.A., et al. (2024). DrugBank 6.0: the DrugBank Knowledgebase for 2024. Nucleic Acids Research *52*, D1265–D1275. https://doi.org/10.1093/nar/gkad976.

59. Yen, K., Travins, J., Wang, F., David, M.D., Artin, E., Straley, K., Padyana, A., Gross, S., DeLaBarre, B., Tobin, E., et al. (2017). AG-221, a First-in-Class Therapy Targeting Acute Myeloid Leukemia Harboring Oncogenic IDH2 Mutations. Cancer Discovery *7*, 478–493. https://doi.org/10.1158/2159-8290.CD-16-1034.

60. Pirozzi, C.J., and Yan, H. (2021). The implications of IDH mutations for cancer development and therapy. Nat Rev Clin Oncol *18*, 645–661. https://doi.org/10.1038/s41571-021-00521-0.

61. Li, J., Huang, S., Chen, J., Yang, Z., Fei, X., Zheng, M., Ji, C., Xie, Y., and Mao, Y. (2007). Identification and characterization of human uracil phosphoribosyltransferase (UPRTase). J Hum Genet *52*, 415–422. https://doi.org/10.1007/s10038-007-0129-2.

62. Nakagawa, T., Lanaspa, M.A., Millan, I.S., Fini, M., Rivard, C.J., Sanchez-Lozada, L.G., Andres-Hernando, A., Tolan, D.R., and Johnson, R.J. (2020). Fructose contributes to the Warburg effect for cancer growth. Cancer & Metabolism *8*, 16. https://doi.org/10.1186/s40170-020-00222-9.

63. Spears, M.E., Lee, N., Hwang, S., Park, S.J., Carlisle, A.E., Li, R., Doshi, M.B., Armando, A.M., Gao, J., Simin, K., et al. (2022). *De novo* sphingolipid biosynthesis necessitates detoxification in cancer cells. Cell Reports *40*, 111415. https://doi.org/10.1016/j.celrep.2022.111415.

64. Liu, Q., Chan, A.K.N., Chang, W.-H., Yang, L., Pokharel, S.P., Miyashita, K., Mattson, N., Xu, X., Li, M., Lu, W., et al. (2022). 3-Ketodihydrosphingosine reductase maintains ER homeostasis and unfolded protein response in leukemia. Leukemia *36*, 100–110. https://doi.org/10.1038/s41375-021-01378-z.

65. Tang, X., Benesch, M.G.K., Dewald, J., Zhao, Y.Y., Patwardhan, N., Santos, W.L., Curtis, J.M., McMullen, T.P.W., and Brindley, D.N. (2014). Lipid phosphate phosphatase-1 expression in cancer cells attenuates tumor growth and metastasis in mice [S]. Journal of Lipid Research *55*, 2389–2400. https://doi.org/10.1194/jlr.M053462.

66. Benesch, M.G.K., Wu, R., Tang, X., Brindley, D.N., Ishikawa, T., and Takabe, K. (2023). Decreased Lipid Phosphate Phosphatase 1/3 and Increased Lipid Phosphate Phosphatase 2 Expression in the Human Breast Cancer Tumor Microenvironment Promotes Tumor Progression and Immune System Evasion. Cancers *15*, 2299. https://doi.org/10.3390/cancers15082299.

67. Tanyi, J.L., Hasegawa, Y., Lapushin, R., Morris, A.J., Wolf, J.K., Berchuck, A., Lu, K., Smith, D.I., Kalli, K., Hartmann, L.C., et al. (2003). Role of Decreased Levels of Lipid Phosphate Phosphatase-1 in Accumulation of Lysophosphatidic Acid in Ovarian Cancer1. Clinical Cancer Research *9*, 3534–3545.

68. Tanyi, J.L., Morris, A.J., Wolf, J.K., Fang, X., Hasegawa, Y., Lapushin, R., Auersperg, N., Sigal, Y.J., Newman, R.A., Felix, E.A., et al. (2003). The Human Lipid Phosphate Phosphatase-3 Decreases the Growth, Survival, and Tumorigenesis of Ovarian Cancer Cells: Validation of the Lysophosphatidic Acid Signaling Cascade as a Target for Therapy in Ovarian Cancer1. Cancer Research *63*, 1073–1082.

69. Lieu, E.L., Nguyen, T., Rhyne, S., and Kim, J. (2020). Amino acids in cancer. Exp Mol Med *52*, 15–30. https://doi.org/10.1038/s12276-020-0375-3.

70. Chen, J., Cui, L., Lu, S., and Xu, S. (2024). Amino acid metabolism in tumor biology and therapy. Cell Death Dis *15*, 42. https://doi.org/10.1038/s41419-024-06435-w.

71. Slotte, J.P. (2013). Biological functions of sphingomyelins. Progress in Lipid Research *52*, 424–437. https://doi.org/10.1016/j.plipres.2013.05.001.

72. Mullen, N.J., and Singh, P.K. (2023). Nucleotide metabolism: a pan-cancer metabolic dependency. Nat Rev Cancer *23*, 275–294. https://doi.org/10.1038/s41568-023-00557-7.

73. Xiao, Y., Yu, T.-J., Xu, Y., Ding, R., Wang, Y.-P., Jiang, Y.-Z., and Shao, Z.-M. (2023). Emerging therapies in cancer metabolism. Cell Metabolism *35*, 1283–1303. https://doi.org/10.1016/j.cmet.2023.07.006.

74. Corsello, S.M., Nagari, R.T., Spangler, R.D., Rossen, J., Kocak, M., Bryan, J.G., Humeidi, R., Peck, D., Wu, X., Tang, A.A., et al. (2020). Discovering the anticancer potential of non-oncology drugs by systematic viability profiling. Nat Cancer *1*, 235–248. https://doi.org/10.1038/s43018-019-0018-6.

75. Shiratori, R., Furuichi, K., Yamaguchi, M., Miyazaki, N., Aoki, H., Chibana, H., Ito, K., and Aoki, S. (2019). Glycolytic suppression dramatically changes the intracellular metabolic profile of multiple cancer cell lines in a mitochondrial metabolism-dependent manner. Sci Rep *9*, 18699. https://doi.org/10.1038/s41598-019-55296-3.

76. DeWaal, D., Nogueira, V., Terry, A.R., Patra, K.C., Jeon, S.-M., Guzman, G., Au, J., Long, C.P., Antoniewicz, M.R., and Hay, N. (2018). Hexokinase-2 depletion inhibits glycolysis and induces oxidative phosphorylation in hepatocellular carcinoma and sensitizes to metformin. Nat Commun *9*, 446. https://doi.org/10.1038/s41467-017-02733-4.

77. Ogretmen, B. (2018). Sphingolipid metabolism in cancer signalling and therapy. Nat Rev Cancer *18*, 33–50. https://doi.org/10.1038/nrc.2017.96.

78. Swanton, C., Marani, M., Pardo, O., Warne, P.H., Kelly, G., Sahai, E., Elustondo, F., Chang, J., Temple, J., Ahmed, A.A., et al. (2007). Regulators of Mitotic Arrest and Ceramide Metabolism Are Determinants of Sensitivity to Paclitaxel and Other Chemotherapeutic Drugs. Cancer Cell *11*, 498–512. https://doi.org/10.1016/j.ccr.2007.04.011.

79. Gong, L., Yang, B., Xu, M., Cheng, B., Tang, X., Zheng, P., Jing, Y., and Wu, G. (2014). Bortezomib-induced apoptosis in cultured pancreatic cancer cells is associated with ceramide production. Cancer Chemother Pharmacol *73*, 69–77. https://doi.org/10.1007/s00280-013-2318-3.

80. Lai, M., Amato, R., La Rocca, V., Bilgin, M., Freer, G., Spezia, P., Quaranta, P., Piomelli, D., and Pistello, M. (2021). Acid ceramidase controls apoptosis and increases autophagy in human melanoma cells treated with doxorubicin. Sci Rep *11*, 11221. https://doi.org/10.1038/s41598-021-90219-1.

81. Jung, J.H., Taniguchi, K., Lee, H.M., Lee, M.Y., Bandu, R., Komura, K., Lee, K.Y., Akao, Y., and Kim, K.P. (2020). Comparative lipidomics of 5-Fluorouracil–sensitive and –resistant colorectal cancer cells reveals altered sphingomyelin and ceramide controlled by acid sphingomyelinase (SMPD1). Sci Rep *10*, 6124. https://doi.org/10.1038/s41598-020-62823-0.

82. Zheng, K., Chen, Z., Feng, H., Chen, Y., Zhang, C., Yu, J., Luo, Y., Zhao, L., Jiang, X., and Shi, F. (2019). Sphingomyelin synthase 2 promotes an aggressive breast cancer phenotype by disrupting the homoeostasis of ceramide and sphingomyelin. Cell Death Dis *10*, 157. https://doi.org/10.1038/s41419-019-1303-0.

83. Chiyoda, T., Hart, P.C., Eckert, M.A., McGregor, S.M., Lastra, R.R., Hamamoto, R., Nakamura, Y., Yamada, S.D., Olopade, O.I., Lengyel, E., et al. (2017). Loss of BRCA1 in the Cells of Origin of Ovarian Cancer Induces Glycolysis: A Window of Opportunity for Ovarian Cancer Chemoprevention. Cancer Prevention Research *10*, 255–266. https://doi.org/10.1158/1940-6207.CAPR-16-0281.

84. Kanakkanthara, A., Kurmi, K., Ekstrom, T.L., Hou, X., Purfeerst, E.R., Heinzen, E.P., Correia, C., Huntoon, C.J., O'Brien, D., Wahner Hendrickson, A.E., et al. (2019). BRCA1 Deficiency Upregulates NNMT, Which Reprograms Metabolism and Sensitizes Ovarian Cancer Cells to Mitochondrial Metabolic Targeting Agents. Cancer Research *79*, 5920–5929. https://doi.org/10.1158/0008-5472.CAN-19-1405.

85. Privat, M., Radosevic-Robin, N., Aubel, C., Cayre, A., Penault-Llorca, F., Marceau, G., Sapin, V., Bignon, Y.-J., and Morvan, D. (2014). BRCA1 Induces Major Energetic Metabolism Reprogramming in Breast Cancer Cells. PLOS ONE *9*, e102438. https://doi.org/10.1371/journal.pone.0102438.

86. Lakhani, S.R., van de Vijver, M.J., Jacquemier, J., Anderson, T.J., Osin, P.P., McGuffog, L., and Easton, D.F. (2002). The Pathology of Familial Breast Cancer: Predictive Value of Immunohistochemical Markers Estrogen Receptor, Progesterone Receptor, HER-2, and p53 in Patients With Mutations in BRCA1 and BRCA2. JCO *20*, 2310–2318. https://doi.org/10.1200/JCO.2002.09.023.

87. Liska, O., Bohár, B., Hidas, A., Korcsmáros, T., Papp, B., Fazekas, D., and Ari, E. (2022). TFLink: an integrated gateway to access transcription factor–target gene interactions for multiple species. Database *2022*, baac083. https://doi.org/10.1093/database/baac083.

88. Abramovitch, S., Glaser, T., Ouchi, T., and Werner, H. (2003). BRCA1–Sp1 interactions in transcriptional regulation of the IGF-IR gene. FEBS Letters *541*, 149–154. https://doi.org/10.1016/S0014-5793(03)00315-6.

89. Benezra, M., Chevallier, N., Morrison, D.J., MacLachlan, T.K., El-Deiry, W.S., and Licht, J.D. (2003). BRCA1 Augments Transcription by the NF-κB Transcription Factor by Binding to the Rel Domain of the p65/RelA Subunit*. Journal of Biological Chemistry *278*, 26333–26341. https://doi.org/10.1074/jbc.M303076200.

90. Sau, A., Lau, R., Cabrita, M.A., Nolan, E., Crooks, P.A., Visvader, J.E., and Pratt, M.A.C. (2016). Persistent Activation of NF-κB in BRCA1-Deficient Mammary Progenitors Drives Aberrant Proliferation and Accumulation of DNA Damage. Cell Stem Cell *19*, 52–65. https://doi.org/10.1016/j.stem.2016.05.003.

91. Li, F., Yuan, L., Lu, H., Li, G., Chen, Y., Engqvist, M.K.M., Kerkhoven, E.J., and Nielsen, J. (2022). Deep learning-based kcat prediction enables improved enzyme-constrained model reconstruction. Nature Catalysis *5*, 662–672. https://doi.org/10.1038/s41929-022-00798-z.

92. Choudhury, S., Moret, M., Salvy, P., Weilandt, D., Hatzimanikatis, V., and Miskovic, L. (2022). Reconstructing Kinetic Models for Dynamical Studies of Metabolism using Generative Adversarial Networks. Nat. Mach. Intell. *4*, 710-+. https://doi.org/10.1038/s42256-022-00519-y.

93. Choudhury, S., Narayanan, B., Moret, M., Hatzimanikatis, V., and Miskovic, L. (2024). Generative machine learning produces kinetic models that accurately characterize intracellular metabolic states. Nat Catal, 1–13. https://doi.org/10.1038/s41929-024-01220-6.

94. Garrido-Rodriguez, M., Zirngibl, K., Ivanova, O., Lobentanzer, S., and Saez-Rodriguez, J. (2022). Integrating knowledge and omics to decipher mechanisms via large-scale models of signaling networks. Molecular Systems Biology *18*, e11036. https://doi.org/10.15252/msb.202211036.

95. Tajan, M., and Vousden, K.H. (2020). Dietary Approaches to Cancer Therapy. Cancer Cell *37*, 767–785. https://doi.org/10.1016/j.ccell.2020.04.005.

96. Apaolaza, I., José-Enériz, E.S., Valcarcel, L.V., Agirre, X., Prosper, F., and Planes, F.J. (2022). A network-based approach to integrate nutrient microenvironment in the prediction of synthetic lethality in cancer metabolism. PLOS Computational Biology *18*, e1009395. https://doi.org/10.1371/journal.pcbi.1009395.

97. Zhang, X., Fam, K.T., Dai, T., and Hang, H.C. (2025). Microbiota mechanisms in cancer progression and therapy. Cell Chemical Biology *32*, 653–677. https://doi.org/10.1016/j.chembiol.2025.04.005.

98. Vitale, I., Manic, G., Coussens, L.M., Kroemer, G., and Galluzzi, L. (2019). Macrophages and Metabolism in the Tumor Microenvironment. Cell Metabolism *30*, 36–50. https://doi.org/10.1016/j.cmet.2019.06.001.

99. Miskovic, L., and Hatzimanikatis, V. (2010). Production of biofuels and biochemicals: in need of an ORACLE. Trends Biotechnol *28*, 391–397. https://doi.org/10.1016/j.tibtech.2010.05.003.

100. Shlomi, T., Cabili, M.N., Herrgård, M.J., Palsson, B.Ø., and Ruppin, E. (2008). Network-based prediction of human tissue-specific metabolism. Nat Biotechnol *26*, 1003–1010. https://doi.org/10.1038/nbt.1487.

101. Baloni, P., Funk, C.C., Yan, J., Yurkovich, J.T., Kueider-Paisley, A., Nho, K., Heinken, A., Jia, W., Mahmoudiandehkordi, S., Louie, G., et al. (2020). Metabolic Network Analysis Reveals Altered Bile Acid Synthesis and Metabolism in Alzheimer's Disease. CR Med *1*. https://doi.org/10.1016/j.xcrm.2020.100138.

102. Agren, R., Bordel, S., Mardinoglu, A., Pornputtapong, N., Nookaew, I., and Nielsen, J. (2012). Reconstruction of Genome-Scale Active Metabolic Networks for 69 Human Cell Types and 16 Cancer Types Using INIT. PLOS Computational Biology *8*, e1002518. https://doi.org/10.1371/journal.pcbi.1002518.

103. Ravi, S., and Gunawan, R. (2021). ΔFBA—Predicting metabolic flux alterations using genome-scale metabolic models and differential transcriptomic data. PLOS Computational Biology *17*, e1009589. https://doi.org/10.1371/journal.pcbi.1009589.

104. Henry, C.S., Broadbelt, L.J., and Hatzimanikatis, V. (2007). Thermodynamics-based metabolic flux analysis. Biophys J *92*, 1792–1805. https://doi.org/10.1529/biophysj.106.093138.

105. Filippo, M.D., Pescini, D., Galuzzi, B.G., Bonanomi, M., Gaglio, D., Mangano, E., Consolandi, C., Alberghina, L., Vanoni, M., and Damiani, C. (2022). INTEGRATE: Model-based multi-omics data integration to characterize multi-level metabolic regulation. PLOS Computational Biology *18*, e1009337. https://doi.org/10.1371/journal.pcbi.1009337.

106. Antoniewicz, M.R. (2018). A guide to 13C metabolic flux analysis for the cancer biologist. Exp Mol Med *50*, 1–13. https://doi.org/10.1038/s12276-018-0060-y.

107. Ataman, M., Hernandez Gardiol, D.F., Fengos, G., and Hatzimanikatis, V. (2017). redGEM: Systematic reduction and analysis of genome-scale metabolic reconstructions for development of consistent core metabolic models. PLoS Comput Biol *13*, e1005444. https://doi.org/10.1371/journal.pcbi.1005444.

108. Ataman, M., and Hatzimanikatis, V. (2017). lumpGEM: Systematic generation of subnetworks and elementally balanced lumped reactions for the biosynthesis of target metabolites. PLoS Comput Biol *13*, e1005513. https://doi.org/10.1371/journal.pcbi.1005513.

109. Park, J.O., Rubin, S.A., Xu, Y.-F., Amador-Noguez, D., Fan, J., Shlomi, T., and Rabinowitz, J.D. (2016). Metabolite concentrations, fluxes and free energies imply efficient enzyme usage. Nat Chem Biol *12*, 482–489. https://doi.org/10.1038/nchembio.2077.

110. Liebermeister, W., and Klipp, E. (2006). Bringing metabolic networks to life: convenience rate law and thermodynamic constraints. Theoretical Biology and Medical Modelling *3*, 41. https://doi.org/10.1186/1742-4682-3-41.

111. Narayanan, B., Weilandt, D., Masid, M., Miskovic, L., and Hatzimanikatis, V. (2024). Rational strain design with minimal phenotype perturbation. Nat Commun *15*, 723. https://doi.org/10.1038/s41467-024-44831-0.

112. Narayanan, B., Jiang, W., Wang, S., Sáez-Sáez, J., Weilandt, D., Barcon, M.M., Hesselberg-Thomsen, V., Borodina, I., Hatzimanikatis, V., and Miskovic, L. (2025). Kinetic-model-guided engineering of multiple *S. cerevisiae* strains improves *p*-coumaric acid production. Metabolic Engineering *91*, 430–441. https://doi.org/10.1016/j.ymben.2025.06.008.

113. Wang, L., Birol, I., and Hatzimanikatis, V. (2004). Metabolic Control Analysis under Uncertainty: Framework Development and Case Studies. Biophysical Journal *87*, 3750–3763. https://doi.org/10.1529/biophysj.104.048090.

114. Gardner, D.J., Reynolds, D.R., Woodward, C.S., and Balos, C.J. (2022). Enabling New Flexibility in the SUNDIALS Suite of Nonlinear and Differential/Algebraic Equation Solvers. ACM Trans. Math. Softw. *48*, 31:1-31:24. https://doi.org/10.1145/3539801.

115. Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B (Methodological) *57*, 289–300.

116. Hatzimanikatis, V., Floudas, C.A., and Bailey, J.E. (1996). Analysis and design of metabolic reaction networks via mixed-integer linear optimization. AIChE Journal *42*, 1277–1292. https://doi.org/10.1002/aic.690420509.

117. Tsouka, S., Ataman, M., Hameri, T., Miskovic, L., and Hatzimanikatis, V. (2021). Constraint-based metabolic control analysis for rational strain engineering. Metab. Eng. *66*, 191–203. https://doi.org/10.1016/j.ymben.2021.03.003.