Title: Time-Restricted Feeding Extends Healthspan in Both Sexes and Lifespan in Male C57BL/6J Mice

Authors: Samantha E. Iiams[1] [†], Nathan J. Skinner[1] [†], Mary Wight-Carter[2], Victoria A. Acosta-Rodríguez[1,4*], Carla B. Green[1*], and Joseph S. Takahashi[1,3*]

Affiliations:
[1]Department of Neuroscience, Peter O'Donnell Jr. Brain Institute, University of Texas Southwestern Medical Center, Dallas, TX, USA
[2]Animal Resource Center, University of Texas Southwestern Medical Center, Dallas, TX, USA
[3]Howard Hughes Medical Institute, University of Texas Southwestern Medical Center, Dallas, TX, USA
[4]Present address: Circadian Biology of Aging Unit, Translational Gerontology Branch, National Institute on Aging (NIA/NIH), Baltimore, MD, 21224, USA

[†] These two authors contributed equally to this work and are co-first authors

[*] Corresponding authors:
Victoria A. Acosta- Rodríguez (VAR); victoria.acosta-rodriguez@nih.gov
Carla B. Green (CBG); carla.green@utsouthwestern.edu,
Joseph S. Takahashi (JST); joseph.takahashi@utsouthwestern.edu

Abstract: Time-restricted feeding (TRF) aligned with an organism's circadian rhythm has been shown to improve health, but its long-term effects on healthspan and lifespan in mammals, especially under normal dietary conditions, remain unclear. Here, we examined the impact of 12-hour (h) and 8h nightly TRF windows in male and female mice fed regular chow. TRF improved multiple health measures, including behavioral rhythmicity, body weight and composition, frailty, and disease onset. These effects were most pronounced in the 8h-TRF group, which exhibited voluntary caloric restriction in addition to time restriction. A composite Healthspan Index revealed that TRF extended healthspan in both sexes, though the benefits were more prolonged in females relative to their total lifespan. Median lifespan was significantly extended in males under 8h-TRF by 12%, whereas females showed no significant lifespan extension, highlighting sex-specific responses to TRF.

Main Text:

While human life expectancy has nearly doubled since the early 20th century (*1*), healthspan remains limited, with many spending over 10% of their lives in poor health due to age-related functional declines and disease development (*2*). Caloric restriction (CR), a potent aging intervention involving a 30-50% reduction in food intake without malnutrition, has been extensively studied for its ability to delay the onset of age-associated diseases, including cancer, cardiovascular disorders, and neurodegeneration, while extending lifespan across multiple model organisms (*3-5*). Incorporating daily feeding/fasting cycles with CR enhances these geroprotective effects (*6-9*), and aligning feeding with an organism's active circadian phase can further extend lifespan (*10*). This synergy likely reflects both the synchronization of nutrient intake with internal metabolic processes and the protective effects of temporal compartmentalization, driven by circadian clock-regulated gene expression in the liver and other peripheral tissues (*11-18*).

Aging triggers a progressive damping of circadian rhythmicity in both humans and animal models. This decline, evident in behavior and physiology, stems from a reduction in amplitude and shifted phases of the molecular clock's transcriptional output (*19*). Interventions that restore circadian rhythms can improve health and extend longevity in model organisms (*20*), suggesting that optimizing meal timing is a promising strategy to counteract age-related rhythm disruptions and promote healthy aging (*21*).

Given that CR is challenging to maintain in people (*22*), meal timing alone via time-restricted feeding (TRF) in animals, or time-restricted eating (TRE) in humans, has emerged as an alternative strategy to delay aging (*23, 24*). TRF, which restricts feeding to specific daily windows without limiting overall caloric intake, can confer similar benefits as timed CR, including improved rhythmic gene expression, cognitive function, and cardiometabolic health, even in obesity models

fed a high-fat, high-sucrose diet (*25-32*). Recent clinical TRE trials report high adherence and

improved cardiometabolic outcomes in subjects with pre-existing conditions, demonstrating its

potential as a feasible intervention to enhance human healthspan (*33-37*).

However, few studies have examined the benefits of TRF/TRE under typical metabolic conditions

without diet- or genetically-induced obesity (*38-40*), and even fewer have compared impacts

between the sexes (*41*). The long-term impact of TRF on healthspan and lifespan in mammalian

models also remains unclear (*42, 43*). In this study, we sought to address these key knowledge

gaps and determine whether TRF could serve as an effective aging intervention for healthy male

and female mice.

Results:

Early Onset Circadian Aligned TRF Has Sex-Specific Impacts on Longitudinal

Feeding and Wheel-Running Activity Levels

To determine how time-restricted feeding (TRF) impacts healthspan and lifespan in mammals, we

individually housed 264 female and 264 male C57BL/6J mice in wheel cages with automated

feeders at 2 months of age. To measure impacts under normal nutritional conditions, mice were

fed purified precision pellets (Bioserv F0075). For the first 8-weeks, all mice were fed *ad libitum*

(free-feeding; AL) with access to a maximum of 22 pellets or 6.6g (23.8 Kcal) per day. At 4 months

of age, N=78/sex were switched to a 12-hour (h) TRF window (Zeitgeber time (ZT) 12-24) and

another N=78/sex to an 8h-TRF window (ZT14-22) both circadian aligned to the night, the active

phase of nocturnal rodents. The remaining N=108/sex were maintained AL to serve as the control.

Across all feeding groups, mice were provided with a daily allotment of food that exceeded their

typical intake, and no group ever consumed the full amount, ensuring that no caloric restriction

was imposed (Fig. 1, A and B, and Fig. S1) (*44*). These feeding regimens were maintained for the

duration of the study. Any food hoarded within the cage was removed and excluded from Fig. 1B

and Figs. S2-3 as described in the methods.

Food Intake

Measurements of daily food intake throughout each mouse's lifespan showed that feeding

increased over time in both sexes, then declined near the end of life in males, similar to previous

mouse studies (Fig. 1, A and B, and Fig. S2-3) (*10, 44*). In line with other rodent longitudinal

recordings (*45*), females in our study consumed more food relative to their body weight at older

ages than males. We also observed sex-specific responses to TRF where 12h-TRF females

consumed the same amount as AL, while 12h-TRF males significantly reduced intake compared

to AL by 8-14% from 140-539 days (~5-18 months) (Full details of ANOVA results for all figures

can be found in Table S1). In contrast, 8h-TRF self-imposed caloric restriction (CR) for the

majority of life in both sexes, with intake in females reduced by 10-22% between 140-728 days

(~5-24 months) and males reduced by 9-23% between 140-980 days (~5-33 months) relative to

sex- and age-matched AL controls. This 8h-TRF schedule, which narrows the daily eating window

without explicitly limiting calorie intake as seen in prior mouse and human studies (*46, 47*),

presents an alternative strategy for achieving long-term CR in both sexes.

Although actograms (Fig. 1A) show occasional pellet-taking events outside the defined TRF

windows, these largely reflect a technical limitation of the feeding system: pellets dispensed at the

very end of the dark phase cannot be retracted, allowing a single pellet to remain available and be

retrieved during the light period. Importantly, this represents at most one 300 mg pellet per day,

accounting for ≤10% of daily intake in 8h-TRF females (lifetime intake 3.0–5.7 g), ≤8% in 12h-

TRF females (3.6–5.9 g), ≤9% in 8h-TRF males (3.2–4.7 g), and ≤8% in 12h-TRF males (3.6–5.4

g). We quantitatively assessed this behavior across the full lifespan and found the true average

proportion of pellets taken outside the feeding window was low, ranging from 5.17–6.34% across

feeding groups and sexes (Fig. S1B). In addition, food hoarded within the cage accounted for no

more than ~10–12% of the amount removed from the feeder (Fig. S1C). Together, these data

suggest that the majority of food intake in both TRF schedules occurs within the intended feeding

window. In contrast, some daytime pellet-taking in AL mice is expected, as C57BL/6J mice

naturally consume a portion of their food during the inactive phase, with ~75% of intake occurring

at night under AL conditions (*9, 10*).

Wheel-Running

Analysis of daily wheel-running recordings for each mouse revealed that individuals in all feeding

groups, of both sexes, consistently displayed nocturnal activity, which subsequently declined with

age, as anticipated (Fig. 1, A and C, and Figs. S4-5) (*48, 49*). Only 8h-TRF males had increased

levels of wheel activity from 224-833 days of age (~7-28 months) with the enhanced activity

ranging +36-82% from 329-833 days (~11-28 months) relative to AL. This male-specific

difference further underscores the sex-specific effects of TRF on behavior. Additionally, in line

with previous work (*10, 50*), the sustained increases in activity under 8h-TRF suggest a healthspan-

enhancing effect in male mice.

TRF Improves the Amplitude of Daily Rhythmic Behaviors

We next examined age-related changes in the diurnal profiles of feeding and wheel-running, two

well-established rhythmic behaviors, as high-amplitude behavioral rhythms are closely associated

with improved health outcomes (Fig. 2, A and B) (*19, 20, 51, 52*). As intended, TRF reduced the

feeding window, shown by longitudinal measurements of the time required for each group to

retrieve 90% of their daily food intake from the feeder. Across both sexes, mice under 12h-TRF

took 90% of their food within 12h, whereas those under 8h-TRF did so within 8h (Fig. 2C). The

time required to reach this threshold declined with age across groups, particularly in females,

indicating that feeding became more consolidated in older animals (Fig. 2, A and C). Consistent

with this pattern, both TRF groups showed significantly longer daily maximum fasting durations

than AL controls, exceeding 10h in the 12h-TRF group and 14h in the 8h-TRF group across both

sexes (Fig. 2D).

A Fast Fourier Transform (FFT) analysis of diurnal amplitude, reflecting the strength of daily

rhythms, confirmed that TRF increased feeding amplitude and led to a preservation of enhanced

rhythmicity with age, particularly in the 8h-TRF group. (Fig. 2, A and E). While the diurnal

amplitude of wheel-running declined with age across all groups, 8h-TRF males maintained

stronger rhythms throughout life compared to controls (Fig. 2, B and F). In females, TRF modestly

increased wheel-running amplitude, with intermittent enhancements relative to controls from ~249

days onward and a sustained elevation in the 8h-TRF group between 543–669 days of age. As

mice age, TRF, particularly an 8h window, counteracted the natural decline in behavioral

rhythmicity by consolidating food intake and preserving higher-amplitude wheel running activity

compared with AL controls.

TRF Improves Body Composition with Limited Effects on Systemic Metabolic

Markers

TRF improves body weight and composition

Body weights were recorded every 21 days (Fig. 3A, and Fig. S6), and body composition was

analyzed every 6 months using nuclear magnetic resonance imaging (EchoMRI, Houston, TX)

(Fig. 3, B and C). In females, 12h-TRF without CR lessened age-related gains in body weight by

5-8% from 350-686 days of age (~12-23 months) compared with AL controls. This feeding

restriction also improved fat vs lean body composition, reducing fat mass while increasing lean

mass by 3-4% at 12 and 18 months of age. Unexpectedly, 8h-TRF females with self-imposed CR

did not exhibit any further improvements beyond 12h-TRF, in weight or fat/lean composition. In

males, 12h-TRF reduced body weight gain by 5-7% from 224-476 days (~7-16 months) and

improved fat vs lean composition by 3% at 12 months. 8h-TRF in males enhanced these benefits,

lowering body weight gain by 5-16% from 161-749 days (~5-25 months) and improving fat vs

lean composition by 3-7% from 6-12 months. Ultimately, these results demonstrate that while 12h-

TRF is sufficient to attenuate weight gain and optimize body composition in both sexes, the

benefits are more prolonged in females; conversely, males derive significantly greater long-term

physiological advantages from a more restrictive 8h-TRF schedule combined with self-imposed

CR.

TRF reshapes diurnal RER and EE rhythms

Metabolic changes under TRF were assessed using indirect calorimetry at 19 months of age. Mice

were randomly selected and individually housed in Promethion metabolic cages equipped with

automated feeders for 5-6 days at room temperature (25°C), followed by 5 days at thermoneutrality

(30°C), to assess metabolic parameters independent of the energetic demands of thermoregulation

(Fig. S7-8). As expected, 24h respiratory exchange ratio (RER) profiles were significantly

influenced by time of day, reflecting normal circadian variation in substrate utilization, and were

significantly impacted by the interaction of TRF x time of day (Fig S7A). While TRF did not have

a significant main effect on mean 24h RER, post-hoc analyses revealed significantly lower RER

values during the delayed onset of feeding in the 8h-TRF group compared with AL controls in

both sexes, consistent with extended lipid oxidation during the prolonged fasting period.

Additionally, in males housed at room temperature, post-hoc analyses identified higher nocturnal

RER values at discrete time points under 8h-TRF, suggesting a shift toward increased carbohydrate

utilization during the night and an enhanced diurnal amplitude of RER.

24h energy expenditure (EE) profiles also exhibited expected diurnal variation and were

significantly influenced by TRF in females, with 8h-TRF trending lower during the daytime/rest

phase (Fig. S7B). Post-hoc analyses further revealed that in 8h-TRF females housed at

thermoneutrality, daytime EE was significantly reduced at several time points, whereas in males,

the minimal impact of TRF was largely restricted to light/dark transitions and feeding/fasting

periods across both temperature conditions. These results indicate that TRF modulates EE in a

time-dependent manner, with effects shaped by sex and ambient temperature. Impacts on EE were

further analyzed by ANCOVA to account for body composition (Fig. S8). Homogeneity of

regression slopes was confirmed, and while lean mass significantly influenced outcomes in males,

TRF effects on EE remained largely time-specific rather than altering overall 24h EE in either sex.

Glucose homeostasis is moderately improved by 8h-TRF in males

We measured 12h fasting glucose levels and performed glucose tolerance tests at 6, 12, and 18

months of age at ZT10 to assess glucose homeostasis. Despite being under circadian control and

strongly influenced by feeding (*53*), neither 12h-TRF nor 8h-TRF produced consistent reductions

in fasting glucose across all ages as seen in our previous study (Fig. S9A) (*9*). Interestingly, 8h-

TRF females exhibited lower fasting glucose at 12 months, but at 18 months levels were

significantly higher. In mice, unlike humans, fasting glucose typically declines with age, so the

higher levels observed at 18 months in 8h-TRF females may in fact reflect a healthier glucose state

(*54*). Glucose tolerance tests also revealed sex-specific effects of TRF. 8h-TRF females showed a

modest improvement in glucose homeostasis at 12 months of age, but no impacts were found at 6

or 18 months of age in either TRF regimen (Fig. S9, B and C). Males exhibited significant

improvements at 6 months in both 12h and 8h-TRF, with modest benefits still apparent at 12

months in the 8h-TRF group. Fasting insulin levels from plasma were not significantly altered by

either TRF regimen at any age examined (Fig. S10).

Little to no impact on circulating metabolic and inflammatory markers

To determine whether TRF broadly altered systemic metabolic or inflammatory signaling, we also

measured circulating levels of leptin, brain derived neurotrophic factor (BDNF), monocyte c

chemoattractant protein-1 (MCP-1), tumor necrosis factor α (TNFα), interleukins (IL)-1β, -6, and

-10 from 12h fasted plasma collected at ZT10 at 6, 12 and 18 months of age. Overall, TRF had

minimal impact on these adipokines, with no sustained differences observed between feeding

groups across ages (Fig. S10). An exception was observed at 6 months of age, where males

subjected to 8h-TRF exhibited higher plasma BDNF levels than 12h-TRF, suggesting an increased

neuroprotective benefit, along with elevated levels of the inflammation-associated chemokine

MCP-1. However, these differences were not maintained with aging. Another exception was found

at 18 months of age where 12h-TRF males had higher levels of IL-1β and 8h-TRF females had

higher levels of MCP-1 suggesting that TRF may have age- and sex-specific effects on

inflammatory markers in older animals.

TRF Slows Age-Related Increases in Frailty Index Values

Frailty, a widely used measure of healthspan in aging mice, integrates 31 physiological and

behavioral parameters into a composite index (*55*). Frailty assessments were conducted every 6 months and revealed that both 12h and 8h-TRF significantly reduced frailty index scores at specific ages compared to AL controls (Fig. 4, A-C, and Figs. S11-12). In females, 12h-TRF significantly reduced frailty index scores at 18 months, with hearing loss and body condition assessments defining the key differences between the two groups. 8h-TRF further extended reductions in frailty index from 12 to 24 months, with specific improvements observed in grimace, piloerection, forelimb grip strength, coat condition, ulcerative dermatitis, and body condition scores. Males showed similar trends: 12h-TRF reduced frailty indices between 12 and 18 months, primarily improving grimace and body condition, while 8h-TRF sustained these benefits through 24 months and improved additional parameters, including hearing, piloerection, menace reflex, fur color loss, and coat condition. These results show that TRF significantly lowers age-related frailty in a dose- and sex-dependent manner, with 8h-TRF providing prolonged benefits and improving the greatest number of health measures.

The Age-Related Onset of Health Reports Reveals Sex-Specific Differences and A Delayed Onset in 8h-TRF Males

As mice developed health issues they were reported to veterinary staff for more frequent monitoring. These reports revealed that TRF did not reduce the relative frequency of commonly reported ailments compared to AL (Fig. 4D). Indeed, 8h-TRF males had significantly more reports of body condition scores declining to 2 (Scored 1-5 with: 1 being emaciated, 2 underconditioned, 3 ideal condition, 4 overconditioned, and 5 obese) (Fisher's exact test 8h-TRF vs AL: Body Condition Score=2, $P < 0.05$). However, we observed a clear sex-dependent pattern in the types

and frequencies of reported conditions. Females, regardless of feeding group, were significantly more likely to be reported for ulcerative dermatitis (Fisher's exact test females vs males: $P < 0.0001$), rectal prolapse ($P < 0.0001$), and hind limb paralysis (often associated with spinal tumors, $P < 0.001$). In contrast, males were more frequently reported for external tumors and abdominal masses (Fisher's exact test females vs males: $P < 0.0001$), as well as low Body Condition Score=2 ($P < 0.0001$). The onset of age-related health issues also differed by sex, with 8h-TRF delaying the median health report onset by 52 days in males only compared to AL controls (Log-rank Mantel-Cox, $P = 0.0018$) (Fig. 4E). Additionally, while there were no significant differences under TRF, overall, females overall spent significantly fewer days on health report prior to death, than males (Fig. 4F). Altogether this demonstrates there is sex-specificity in the development of common husbandry related-disease.

TRF Has Minimal Impacts on Hematological Markers

To determine whether lifelong TRF impacted hematological markers, which are emerging as predictors of systemic health and aging (*56*), we assessed complete blood counts from whole blood at 6, 12 and 18 months of age. Overall, TRF had minimal effects on circulating blood parameters in either sex. Red blood cell counts (RBC), hemoglobin, hematocrit, red cell indices including mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC), and platelet counts were largely unchanged across feeding regimens (Figs. S13-14). TRF also had no significant impacts on total white blood cell counts or leukocyte subtypes (Fig. S15). The only notable exception was observed in females at 18 months of age, where 8h-TRF mice displayed higher RBC counts accompanied by lower MCV (smaller RBCs; typically indicate older, longer maintained cells) and MCH compared to AL controls (Fig.

261   S13). Given that old age in mice is commonly associated with reduced RBC survival (*57*), this

262   pattern might reflect improved maintenance of RBC homeostasis rather than a pathological shift.

263   Importantly, MCHC, hemoglobin levels, and hematocrit were unchanged in 8h-TRF females,

264   indicating preserved hemoglobin concentration within cells and maintained oxygen-carrying

265   capacity.

8h-TRF Extends Lifespan in Males Only

For the first time, we were able to investigate the impact of circadian aligned TRF with regular

chow diet, on lifespan in mice (Fig. 5A, and Tables S2-3). In females, no significant extension of

median lifespan was observed with either 12h-TRF or 8h-TRF. Median survival was 715 days and

729 days, respectively, compared to 694 days in AL controls. In males, 12h-TRF also failed to

extend lifespan, with a median survival of 839 days compared to 818 days in AL, a finding now

reproduced across two independent experiments (Fig. S16). Interestingly, 8h-TRF with lifelong

self-imposed CR, significantly improved overall survival for males (Log-rank Mantel-Cox, P =

0.0001). Median lifespan increased by approximately 12%, reaching 916 days, and maximal

lifespan was prolonged by ~3% (Fisher's exact test, P = 0.0038). Though TRF is beneficial for

improving health in both sexes, its capacity to extend lifespan appears to be feeding window- and

sex-dependent.

Sex-Dependent Differences in Disease Types and Incidence at Death

After death, each mouse underwent gross necropsy and histopathological analysis by a veterinary

pathologist to determine the types of diseases that contributed to its debilitation (Fig. 5B, and

Tables S4-6). As shown previously (*10*), the types and relative frequency of diseases were similar

for males across all feeding conditions with the most common cause of death or debilitation being

neoplasia (Fig. 5B, and Table S7). Along with the delayed onset of health reports, overall, this

supports that rather than preventing disease, 8h-TRF delays the development of age-related

diseases in males.

However, a few specific neoplastic subtypes were differentially affected by TRF. In males,

bronchioloalveolar adenomas, benign tumors, were significantly increased in both 12h-TRF and

8h-TRF groups compared to AL controls. (Fisher's exact tests AL vs TRF: 12h and 8h, $P < 0.05$).

In females, hepatocellular adenomas, were significantly more frequent in the 8h-TRF group

(Fisher's exact tests AL vs TRF: 8h $P < 0.05$). While this indicates increased growth of tumors

with malignant potential, the absence of a corresponding increase in carcinomas suggests that TRF

may shift the spectrum of pathology toward less aggressive neoplastic lesions rather than

completely preventing tumorigenesis.

TRF did not reduce the overall occurrence of disease in females either. However, females

developed neoplasms far less frequently than males (Fisher's exact tests females vs males:

Adenoma in liver, $P < 0.05$; Carcinoma in liver and lung, each $P < 0.0001$; Histiocytic sarcoma in

liver and lung, each $P < 0.0001$; Histiocytic sarcoma in spleen, $P < 0.001$; Acidophilic macrophage

pneumonia in lung, $P < 0.0001$; Adenoma in lung, $P < 0.0001$). Instead, most females were

debilitated by glomerulonephritis in the kidneys and had a higher incidence of lymphoma than

males (Fisher's exact tests females vs males: Lymphoma in liver, kidney, spleen, each $P < 0.05$;

Glomerulonephritis, $P < 0.0001$). These results have revealed pronounced differences in neoplasia

and development of other age-related morbidities between the sexes.

TRF Extends Healthspan, With More Prolonged Benefits in Females

While TRF improved and attenuated age-related changes in several markers of health, we next aimed to quantify both the magnitude and duration of its overall impact by measuring healthspan. Frailty index is commonly used to determine healthspan (*55, 58*), but it cannot incorporate the impact of behavioral changes in wheel-activity and food consumption and can miss important physiological changes in the time between assessments. To generate a continuous healthspan index, we integrated frailty with 12 additional health measures by first normalizing each data set (mean = 0, standard deviation = 1), then scaling the parameter directionally by providing context based on whether higher or lower values reflected better health (e.g., high wheel-running activity was scaled positively, while low frailty scores were inverted and also scaled positively) (Fig. 6A), and finally summing the values across age points (Fig. 6B). With higher values reflecting better health, we found that both females and males under TRF had significantly improved healthspan indices throughout life compared to AL controls (Fig. 6, B and C). In 12h-TRF females, this improvement in health was maintained from the onset of TRF until 774 days of age (~26 months). In 8h-TRF females, health benefits persisted until 879 days of age (~29 months), and these mice remained healthier than the 12h-TRF group until 858 days (~29 months). In males, 12h-TRF also improved the healthspan index through 879 days of age (~29 months), whereas 8h-TRF extended these improvements to 963 days (~32 months) and were significantly healthier than the 12h-TRF group until 942 days (~31 months). Importantly, these healthspan indices strongly correlate with lifespan in both sexes, demonstrating that this integrated metric captures biologically meaningful health outcomes (Fig. 6D).

We also found that health declines were more delayed in females than in males, particularly under TRF. Using subjective thresholds of '0' or '-5' (black dashed lines), AL females crossed the '-5'

threshold more than 100 days later than AL males, indicating that females maintain better health

for longer, especially relative to their total lifespan. Under 12h-TRF, females not only had

prolonged healthspan relative to controls at these thresholds, but also crossed the '0' threshold

more than 200 days later than males on the same TRF regimen. While 8h-TRF extended health

above the '0' threshold by approximately 500 days relative to AL in both sexes, this also made up

a greater portion of the females lifespan. Together, these results demonstrate that TRF extends

healthspan in both sexes and further suggest that, despite the absence of lifespan extension in

females, females in both TRF regimens remain healthier for a greater proportion of their lifespan

than males.

To validate our findings of TRF-induced healthspan extension further, we also applied a previously

published summary metric, Frailty-Adjusted Mouse Years (FAMY) (*59*). FAMY, which relies

solely on frailty scores, similarly detected a significant healthspan increase in 8h-TRF males (Fig.

S17A). The lower frequency of frailty assessments in our study, performed only once every six

months, likely contributes to the banded distribution of FAMY scores and may explain

discrepancies between the two methods. Nevertheless, FAMY showed strong correlations with

lifespan across all groups (Fig. S17B). These results support the robustness of our healthspan

measurements and suggest that our more comprehensive, higher-resolution index may be better

suited for detecting healthspan changes in females.

Discussion:

Previous studies have established that TRF improves circadian rhythms, ameliorates metabolic

syndrome, and improves overall health in obesity models consuming a 'Western diet' high in fats

and sugars (*28-32*). We expand on these findings by revealing that circadian aligned TRF confers

significant, long-term health benefits even under normal nutritional conditions. We show that,

depending on the feeding window and sex, TRF prolonged higher amplitudes of daily rhythms in

feeding and wheel-running, which are known to feed back on the circadian clock to drive robust

molecular and behavioral oscillations (*14, 15, 17, 47, 60-62*). Both TRF paradigms were sufficient

to significantly reduce age-related increases in weight and fat mass, having the largest impact

during midlife. TRF similarly slowed the age-related decline in lean composition, which may also

contribute to the observed improvements in adiposity and suggests that TRF can delay the onset

of functional disabilities (*63-65*). We also present novel findings that both TRF windows reduce

frailty index in females and males, as in CR and intermittent fasting studies (*66-70*). Veterinary

health reports were largely sex-dependent, and overall TRF did not alter the types of deficits

observed. Notably, 8h-TRF males displayed a higher proportion of low body condition scores

(score of 2), but this occurred alongside a delayed onset of health issues and overall increased

lifespan and likely reflects the survival of more mice to advanced ages rather than a detrimental

effect of TRF. Consistently, no differences in body condition score were observed at 30 months in

males within the individual frailty assessments. Altogether, this demonstrates that TRF does not

merely act as a remedial strategy to 'rescue' poor health but can actively improve behavioral and

physiological markers of health for both sexes under normal metabolic aging conditions.

Despite these robust improvements in health, TRF-induced reductions in body weight and

adiposity were not accompanied by consistent improvements in glucose tolerance or sustained

alterations in fasting glucose, insulin, leptin, or circulating inflammatory markers. These findings

indicate that the metabolic benefits of TRF do not require large shifts in systemic endocrine or

inflammatory signaling. Instead, they suggest that TRF may promote health through alternative

mechanisms, such as enhanced circadian organization, behavioral rhythmicity, or tissue-specific

metabolic adaptations. Consistent with this, 24h EE and RER profiles revealed that TRF primarily

reshaped temporal patterns of substrate utilization and EE, rather than altering overall 24h mean

values. Importantly, these benefits occurred without detrimental effects on hematological health.

TRF had no impact on red blood cell indices, platelet counts, or immune cell abundance across

aging, and the modest changes observed in 8h-TRF females were not associated with impaired

oxygen-carrying capacity, supporting the long-term physiological safety of circadian aligned TRF.

By integrating measures of frailty, physiology, and behavior into a novel, comprehensive

healthspan index, we demonstrate that TRF significantly enhances healthspan in both sexes, with

proportionally greater benefits in females relative to their lifespan. We also found that 12h-TRF

strongly delayed the initial healthspan decline in females during middle-age, whereas its effect in

males at the same threshold is minimal. However, this approach may oversimplify sex-specific

differences, as males and females exhibit distinct patterns of health deficits that associate

differently with longevity. A single healthspan scale as used here might not adequately reflect

these complex, sex-dependent impacts on aging. Although females generally show better health

indices, we found they had shorter lifespans than males, highlighting a potential disconnect

between this healthspan index and ultimate survival outcomes. It is also important to note, that sex

differences observed in mice may not fully translate to humans, as differences in circadian

rhythms, hormonal cycles and reproduction, and metabolic rate can alter the magnitude and timing

of sex-specific responses to TRF (*71-74*). Regardless, this sex-specific response in healthspan

likely reflects a combination of biological factors. Estrogen is known to confer broad protective

effects on health and aging (*75*), which may explain the overall female advantage even under AL

feeding, and is known to feedback onto the circadian clock (*76-78*). With recent literature showing

that females possess a more robust circadian system (*71*), this may underlie their enhanced

responsiveness to circadian-based interventions. Recent studies have found that females express a

greater number of rhythmic genes, display higher amplitude rhythms, and are more sensitive to

external cues (*79-81*). As such, TRF may more effectively enhance rhythms at the molecular level

to promote health in females (*19, 20*).

For the first time, we show that aligning feeding to the active phase under 12h-TRF does not extend

lifespan. However, a narrower feeding window of 8h-TRF along with self-imposed CR conferred

strong lifespan extension in males. These results contrast with previous studies showing that

lifelong 20% CR, adjusted relative to age-related changes in AL feeding, extends lifespan by

40.6% in female and 24.4% in male C57BL/6J mice (*82*). While our 8h-TRF mice maintained

~20% CR for nearly half of their lifespan, both males and females gradually lessened the degree

of CR relative to AL feeding shortly before reaching median lifespan. Despite this, 8h-TRF males

gained a 12% extension in median lifespan, indicating that even a lower degree of CR is still

impactful in males. In contrast, females appear to require a sustained degree of CR ≥20% to see

any lifespan benefits. Alternatively, the duration of fasting may be more crucial for female

lifespan, as longer fasting periods with CR have enhanced lifespan effects in females (*8, 82*), but

not in males (*10*). The mechanisms underlying these sex-dependent differences in response to CR

and TRF are not fully understood, especially between different strains of mice (*5, 82*), but are

likely influenced by a complex interplay of factors such as differences in nutrient sensing and

metabolic pathways, DNA repair, and the effects of sex hormones (*41, 71, 83*).

Survival curves for AL mice unexpectedly showed an 18% increase in median survival for males

compared to females, contrary to the common trend in mammalian studies where females typically

live longer due to the genetic advantage of a backup X chromosome and the protective effects of

estrogen (*84, 85*). However, in C57BL/6J mice, lifespan outcomes are inconsistent possibly due to

genetic drift within the isogenic line as well as environmental variations between experiments (*85-*

88). While genetic factors may play a role, the shorter lifespan in females could stem from

enhanced cold sensitivity and higher thermoregulatory energy demands (*89-92*). The smaller

bodied females, needing more energy to maintain body temperature as evidenced by the

pronounced increase in food intake with age, may have been adversely affected by the harsher

individual housing conditions without nesting material. Although lower body temperatures can

extend lifespan (*93, 94*), cold stress can also divert energy away from and impair metabolic,

immune, and reproductive functions (*95, 96*). Future studies will be needed to directly evaluate

thermogenic pathways underlying these sex-specific effects, including gene expression in brown

adipose tissue (*47*). In addition, evaluating TRF and lifespan outcomes under thermoneutral

housing conditions or with access to nesting material will help disentangle the relative roles of

thermoregulation and feeding timing in shaping healthspan and longevity.

Health report monitoring and post-death histopathological analyses revealed significant

differences between the sexes in the progression of age-related diseases, particularly in neoplasia.

With the exception of lymphoma, female mice had significantly lower occurrences of neoplasms

in the liver, lung, kidney, and spleen which supports a growing body of data that females have

greater resistance to cancer (*97-99*). This resistance is largely attributed to sex differences in

immunity, mutational burden, and DNA repair, as well as the protective role of X-linked tumor

suppressor genes and their interaction with the p53 pathway in females. Although mechanistic

experiments are beyond the scope of this study, RNA-sequencing and proteomic analysis of the

liver, the tissue most affected by neoplasia in males and least in females, could help to dissect the

molecular mechanisms underlying these sex-specific differences in cancer development under

TRF. These mechanisms are likely to include pathways regulating the cell cycle and cellular

proliferation. Females in this cohort were instead most susceptible to kidney failure due to

glomerulonephritis, similar to a previous study (*100*). This condition has been linked to menopause

in women, with its progression often accelerating after the decline in estrogen levels (*101*). In

mice, this disease can be further exacerbated in the absence of the *estrogen receptor alpha* gene

(*102*), emphasizing the protective role of estrogen signaling and uncovering renal disease as a

potential biomarker of reproductive aging.

One limitation of this study was the inability of the automated feeders to retract or block a food

pellet that goes uneaten beyond the restricted food intake window. This meant that often, a single

pellet could be taken by the mouse, outside of its 12h or 8h-TRF window. Additionally, some mice

had the tendency to hoard pellets within their cage, leading again to food availability outside of

the TRF window, a trait that has already been shown in aged mouse populations (*44*). However, a

single daytime pellet accounted for $\leq 6.34\%$ of total caloric intake, and after subtracting hoarded

food, mice consumed $\geq 90\%$ of the pellets taken from the feeder during the correct feeding

windows. These findings, along with evidence that TRF 5 out of 7 days per week regimens still

improves metabolic health in mice (*39*), and lifespan in flies (*103*), suggest that moderate TRE

adherence may still prolong healthspan and be compatible with modern human lifestyles.

Our study focused on TRF regimens initiated in early adulthood and maintained throughout life,

leaving the efficacy of late-life implementation yet to be determined. Short-term studies indicate

that TRF in the elderly can reduce high-fat diet-induced inflammation and metabolic dysfunction,

and studies with CR in aged mice still provided health and lifespan benefits (*104-106*). However,

these benefits provide diminishing returns, dependent on the age at which an intervention is started.

Furthermore, as most of the observed benefits occurred through mid-life and the onset of old age,

this raises the question of whether a lifelong intervention is necessary, or if stopping TRF before

old age might be sufficient, or even preferable, to avoid potential late-life trade-offs. This

possibility is supported by other models, where short-term TRF applied in flies earlier in life

increases longevity and functional health even after returning to AL feeding (*42, 103, 107*), and

by some evidence that TRF in aged mice had negative impacts on lean mass and cardiovascular

health (*108*).

Beyond the life stage at which TRF is initiated, the specific timing of the feeding window relative

to the internal circadian clock represents a critical variable in metabolic optimization. In this study,

the 8h-TRF window was intentionally positioned in the middle of the active period (ZT14–ZT22),

instead of the onset (ZT12), as this is consistent with several previous mouse studies (*29, 109*),

and prior work in humans suggests that delaying breakfast and/or advancing the final meal can

confer metabolic benefits (*110*). Thus, it remains possible that the enhanced healthspan and

lifespan benefits observed with 8h-TRF reflect not only feeding duration, but also a shift in the

midpoint of daily caloric intake, an aspect of circadian alignment that warrants further

investigation. While we observed robust benefits of 8h-TRF in mice, it is also important to note

that direct translation to humans is limited by differences in metabolic rate and circadian timing

(*73*). In practice, fasting intervals in mice correspond to approximately ~2.7–3.6 times longer

relative windows in humans, so caution is warranted when extrapolating the duration and timing

of TRF across species (*73, 74*).

The absence of direct cardiovascular assessments remains a limitation, particularly as

cardiovascular disease is a prevalent age-associated pathology in mice. Given the limited data from

long-term studies directly evaluating cardiovascular effects of TRF/TRE (*111-113*), future studies

incorporating dedicated measures of cardiac function, structure, and blood pressure will be

essential to determine how timed feeding influences cardiovascular health across the lifespan,

particularly under conditions of normal metabolic aging.

In conclusion, our study provides compelling evidence that the early initiation of circadian aligned

TRF either via a 12h or 8h window significantly improves healthspan in both sexes, more so in

females proportional to lifespan, and that 8h-TRF robustly extends lifespan in males. While this

still warrants further exploration in additional mouse strains and at varying ages of onset, taken

together this highlights the potential of TRF as a practical, non-pharmacological intervention to

prolong the health quality of human lifespans.

 Figures and Legends:

Figure 1: Circadian aligned TRF reduces feeding and improves wheel-running activity in a feeding window- and sex-dependent manner. (**A**) In females 2-26 months of age and males 2-33 months of age, representative actograms of individual mice show wheel-running (black mark) and food pellet taking (pink dot) activity over the course of the 24-hour (h) light:dark cycle. Black horizontal line shows ends of *ad libitum* (AL) baseline feeding for the TRF groups at 4 months of age. (**B**) (*Top*) Longitudinal profiles of average daily food intake and energy intake with age and (*Bottom*) percent differences in CR relative to AL controls with age. Two-way ANOVA: Females: Age=****, Feeding=****, Interaction=****; Males: Age=****, Feeding=****, Interaction=**** (Full details of ANOVA results for each figure can be found in Table S1). (**C**) Longitudinal profiles of average daily wheel revolutions with age. Two-way ANOVA: Females: Age=****, Feeding=**, Interaction=ns; Males: Age=****, Feeding=****, Interaction=ns. B and C: AL baseline feeding (Gray area). TRF feeding conditions (White area). Each point represents mean of 21 days ± SEM (shaded regions). Asterisks/percents represent significant differences in TRF vs AL (*$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$) as determined by two-way ANOVA and Tukey's post-hoc. Females N=71-103 per group. Males N=71-97 per group.

Figure 2: TRF improves the amplitude of daily feeding and wheel-running rhythms. (**A**) 24h

profiles of feeding and (**B**) wheel-running starting at 3 and then every 6 months of age. 12h light

(white area), 12h dark (gray area). (**C**) Longitudinal plots for daily time it takes each group of mice

to take 90% of their daily intake from the feeder. Type III ANOVA (Wald χ^2): Females: Age=****,

Feeding=****, Interaction=****; Males: Age=****, Feeding=****, Interaction=****. (**D**)

Longitudinal plots for daily maximum fasting times achieved. Type III ANOVA (Wald χ^2):

Females: Age=****, Feeding=****, Interaction=****; Males: Age=****, Feeding=****,

Interaction=****. To note: Large, transient dips in fasting due to interruptions in data acquisition

(e.g., fasting blood collection attempt and computer outage). (**E**) Longitudinal FFT plots of the

diurnal amplitude of feeding with age. Type III ANOVA (Wald χ^2): Females: Age=****,

Feeding=****, Interaction=****; Males: Age=****, Feeding=****, Interaction=****. (**F**)

Longitudinal FFT plots of the diurnal amplitude of wheel-running with age. Type III ANOVA

(Wald χ^2): Females: Age=****, Feeding=*, Interaction=****; Males: Age=****, Feeding=****,

Interaction=***. C-F: AL baseline feeding (Gray area). TRF feeding conditions (White area). Each

point represents mean of 14 days ± SEM (shaded regions). Asterisks represent significant

differences in TRF vs AL (*$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$) as determined

by type III ANOVA (Wald χ^2) and Holm's post-hoc. For all panels: Females N=71-103 per group.

Males N=71-97 per group.

Figure 3: A 12h TRF window is sufficient to reduce body weight and fat mass in both sexes.

(**A**) (*Top*) Longitudinal profiles of body weight and (*Bottom*) percent differences in body weight relative to AL controls with age. Two-way ANOVA: Females: Age=****, Feeding=****, Interaction=****; Males: Age=****, Feeding=****, Interaction=****. AL baseline feeding (Gray area). TRF feeding conditions (White area). Each point represents mean of 21 days (cage change) ± SEM (bars). Asterisks represent significant differences in TRF vs AL (*P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤0.0001) as determined by two-way ANOVA and Tukey's post-hoc. Females

N=76-103 per group. Males N=76-102 per group. (**B**) Total fat mass as a percentage of body

weight with age. (**C**) Total lean mass as a percentage of body weight with age. <u>B and C:</u> Means

every 6 months ± SEM (bars). Asterisks represent significant differences in TRF vs AL as

determined by two-way ANOVA and Tukey's post-hoc. Females 6-24 months: AL, N=43-101.

12h-TRF, N=32-71. 8h-TRF, N=38-72. Males 6-30 months: AL, N=23-102. 12h-TRF, N=19-77.

8h-TRF, N=36-75.

Figure 4: TRF reduces frailty index in both sexes but only delays onset of health reports in males. (**A**) Boxplots of frailty index scores every 6 months. Two-way ANOVA: Females: Age=****, Feeding=****, Interaction=ns; Males: Age=****, Feeding=****, Interaction=**.

Asterisks represent significant differences in TRF vs AL (*$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$;

****$P \leq 0.0001$) as determined by two-way ANOVA and Tukey's post-hoc. Females: AL, N=44-

79. 12h-TRF, N=34-74. 8h-TRF, N=40-74. Males: AL, N=23-79. 12h-TRF, N=22-78. 8h-TRF,

N=39-76. (**B**) One example parameter contributing to reduced frailty index in 8h-TRF females:

coat condition, $P < 0.05$ at 24 months compared to AL. (Full list see Fig. S11). (**C**) One example

parameter contributing to reduced frailty index in 8h-TRF males: body condition score, $P < 0.05$

12-18 months compared to AL. (Full list see Fig. S12). (**D**) Percentage of mice in each feeding

group and sex reported to veterinary staff for Body Condition Score=2, Tumor/Mass, Ulcerative

dermatitis, Rectal prolapses, Eye ulcerations/swelling, Hind end paralysis, Malocclusions, and

Head tilts. Black asterisks represent significant differences in TRF vs AL, and all other colors

represent male vs female comparisons as determined by contingency tables and Fisher's exact

tests. (**E**) Cumulative curve showing the percentage of veterinary health reports filed in each group

with age. Log-Rank Mantel-Cox Test. (**F**) Time spent on health report until death. Two-way

ANOVA: Sex=**, Feeding=ns, Interaction=ns. Data shown as mean ± SEM.

Figure 5: 8h-TRF extends lifespan in males. (**A**) Kaplan-Meier survival curves and day median lifespan reached. Log-Rank Mantel-Cox Test for significant difference in overall survival TRF vs AL. Fisher's exact test for maximal survival TRF vs AL. Females N=72-99 per group. Males N=73-99 per group. (**B**) (*Left*) Histopathology results showing the percentage incidence of any disease type in the most impacted tissue types between females and males and (*Right*) incidence by specific disease type within these tissues separated by sex and feeding group (Full table with all tissues and diseases shown in Tables S4-6). Significant differences in females vs males highlighted by blue boxes and shown with blue asterisks (*P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤0.0001) as determined by contingency tables and Fisher's exact tests.

Figure 6: TRF improves healthspan in both sexes. (A) Ridgeline plots for each of the 13 health measures recorded throughout the experiment. All parameters were standardized (mean = 0, SD =

1) and, where appropriate, oriented in the positive direction to reflect better health. (**B**) Healthspan

index summing the 13 health measures for each group from panel A. Higher values indicate better

health and lower values poorer health. Dashed lines (black) mark the age at which each group's

index crosses subjective thresholds of 0 and –5. Type III ANOVA (Wald χ^2): Females: Age=****,

Feeding=****, Interaction=****; Males: Age=****, Feeding=****, Interaction=ns. Each point

represents mean of 21 days ± SEM (shaded regions). Asterisks/percents represent significant

differences (*P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤0.0001) as determined by type III

ANOVA and Holm's post-hoc. (**C**) Area under the curve (AUC) for healthspan index in females

and males. Type III ANOVA: Sex=****, Feeding=****, Interaction=***. Mean ± SEM (bars).

Asterisks represent significant differences as determined by type III ANOVA and Holm's post-

hoc. (**D**) Correlation plots of healthspan AUC vs lifespan in years. Correlation coefficients (r) and

coefficients of determination (r^2) shown.

References and notes:

1.   D. Costa, Health and the Economy in the United States, from 1750 to the Present. *J Econ Lit* **53**, 503-570 (2015).

2.   J. W. Vaupel, Biodemography of human ageing. *Nature* **464**, 536-542 (2010).

3.   C. L. Green, D. W. Lamming, L. Fontana, Molecular mechanisms of dietary restriction promoting health and longevity. *Nat Rev Mol Cell Biol* **23**, 56-73 (2022).

4.   A. McKay *et al.*, An automated feeding system for the African killifish reveals the impact of diet on lifespan and allows scalable assessment of associative learning. *Elife* **11**, (2022).

5.   A. Di Francesco *et al.*, Dietary restriction impacts health and lifespan of genetically diverse mice. *Nature* **634**, 684-692 (2024).

6.   S. J. Mitchell *et al.*, Daily Fasting Improves Health and Survival in Male Mice Independent of Diet Composition and Calories. *Cell Metab* **29**, 221-228 e223 (2019).

7.   H. H. Pak *et al.*, Fasting drives the metabolic, molecular and geroprotective effects of a calorie-restricted diet in mice. *Nat Metab* **3**, 1327-1341 (2021).

8.   E. Duregon *et al.*, Prolonged fasting times reap greater geroprotective effects when combined with caloric restriction in adult female mice. *Cell Metab* **35**, 1179-1194 e1175 (2023).

9. V. A. Acosta-Rodriguez, M. H. M. de Groot, F. Rijo-Ferreira, C. B. Green, J. S. Takahashi,
Mice under Caloric Restriction Self-Impose a Temporal Restriction of Food Intake as
Revealed by an Automated Feeder System. *Cell Metab* **26**, 267-277 e262 (2017).

10. V. Acosta-Rodriguez *et al.*, Circadian alignment of early onset caloric restriction promotes
longevity in male C57BL/6J mice. *Science* **376**, 1192-1202 (2022).

11. H. Reinke, G. Asher, Crosstalk between metabolism and circadian clocks. *Nat Rev Mol*
Cell Biol **20**, 227-241 (2019).

12. I. Laothamatas, E. S. Rasmussen, C. B. Green, J. S. Takahashi, Metabolic and chemical
architecture of the mammalian circadian clock. *Cell Chem Biol* **30**, 1033-1052 (2023).

13. J. S. Takahashi, Transcriptional architecture of the mammalian circadian clock. *Nat Rev*
Genet **18**, 164-179 (2017).

14. G. Manella *et al.*, The liver-clock coordinates rhythmicity of peripheral tissues in response
to feeding. *Nat Metab* **3**, 829-842 (2021).

15. B. J. Greenwell *et al.*, Rhythmic Food Intake Drives Rhythmic Gene Expression More
Potently than the Hepatic Circadian Clock in Mice. *Cell Rep* **27**, 649-657 e645 (2019).

16. C. Vollmers *et al.*, Time of feeding and the intrinsic circadian clock drive rhythms in
hepatic gene expression. *Proc Natl Acad Sci U S A* **106**, 21453-21458 (2009).

17. K. A. Stokkan, S. Yamazaki, H. Tei, Y. Sakaki, M. Menaker, Entrainment of the circadian
clock in the liver by feeding. *Science* **291**, 490-493 (2001).

18. L. Palluth, J. S. Takahashi, C. B. Green, Keeping up with the nicotinamides: NADP(H),
the forgotten circadian cofactor that keeps metabolic time. *Life Metabolism*, loaf034
(2025).

19. S. Hood, S. Amir, The aging clock: circadian rhythms and later life. *J Clin Invest* **127**, 437-
446 (2017).

20. V. A. Acosta-Rodriguez, F. Rijo-Ferreira, C. B. Green, J. S. Takahashi, Importance of
circadian timing for aging and longevity. *Nat Commun* **12**, 2862 (2021).

21. A. Chaix, E. N. C. Manoogian, G. C. Melkani, S. Panda, Time-Restricted Eating to Prevent
and Manage Chronic Metabolic Diseases. *Annu Rev Nutr* **39**, 291-315 (2019).

22. W. E. Kraus *et al.*, 2 years of calorie restriction and cardiometabolic risk (CALERIE):
exploratory outcomes of a multicentre, phase 2, randomised controlled trial. *Lancet*
Diabetes Endocrinol **7**, 673-683 (2019).

23. V. D. Longo, S. Panda, Fasting, Circadian Rhythms, and Time-Restricted Feeding in
Healthy Lifespan. *Cell Metab* **23**, 1048-1059 (2016).

24. M. E. Parrotta *et al.*, Time Restricted Eating: A Valuable Alternative to Calorie Restriction
for Addressing Obesity? *Curr Obes Rep* **14**, 17 (2025).

25. D. M. Arble, J. Bass, A. D. Laposky, M. H. Vitaterna, F. W. Turek, Circadian timing of
food intake contributes to weight gain. *Obesity (Silver Spring)* **17**, 2100-2102 (2009).

26. E. N. C. Manoogian, L. S. Chow, P. R. Taub, B. Laferrere, S. Panda, Time-restricted Eating for the Prevention and Management of Metabolic Diseases. *Endocr Rev* **43**, 405-436 (2022).

27. P. Balasubramanian *et al.*, Time-restricted feeding (TRF) for prevention of age-related vascular cognitive impairment and dementia. *Ageing Res Rev* **64**, 101189 (2020).

28. S. Deota *et al.*, Diurnal transcriptome landscape of a multi-tissue response to time-restricted feeding in mammals. *Cell Metab* **35**, 150-165 e154 (2023).

29. M. Hatori *et al.*, Time-restricted feeding without reducing caloric intake prevents metabolic diseases in mice fed a high-fat diet. *Cell Metab* **15**, 848-860 (2012).

30. H. Sherman *et al.*, Timed high-fat diet resets circadian metabolism and prevents obesity. *FASEB J* **26**, 3493-3502 (2012).

31. P. Regmi *et al.*, Early or delayed time-restricted feeding prevents metabolic impact of obesity in mice. *J Endocrinol* **248**, 75-86 (2021).

32. C. Hepler *et al.*, Time-restricted feeding mitigates obesity through adipocyte thermogenesis. *Science* **378**, 276-284 (2022).

33. E. N. C. Manoogian *et al.*, Feasibility of time-restricted eating and impacts on cardiometabolic health in 24-h shift workers: The Healthy Heroes randomized control trial. *Cell Metab* **34**, 1442-1456 e1447 (2022).

34. K. L. Haganes *et al.*, Time-restricted eating and exercise training improve HbA1c and body composition in women with overweight/obesity: A randomized controlled trial. *Cell Metab* **34**, 1457-1471 e1454 (2022).

35. T. Che *et al.*, Time-restricted feeding improves blood glucose and insulin sensitivity in overweight patients with type 2 diabetes: a randomised controlled trial. *Nutr Metab (Lond)* **18**, 88 (2021).

36. M. Dote-Montero *et al.*, Effects of early, late and self-selected time-restricted eating on visceral adipose tissue and cardiometabolic health in participants with overweight or obesity: a randomized controlled trial. *Nat Med* **31**, 524-533 (2025).

37. V. Pavlou *et al.*, Effect of Time-Restricted Eating on Weight Loss in Adults With Type 2 Diabetes: A Randomized Clinical Trial. *JAMA Netw Open* **6**, e2339337 (2023).

38. Z. Xie *et al.*, Randomized controlled trial for time-restricted eating in healthy volunteers without obesity. *Nat Commun* **13**, 1003 (2022).

39. A. Chaix, A. Zarrinpar, P. Miu, S. Panda, Time-restricted feeding is a preventative and therapeutic intervention against diverse nutritional challenges. *Cell Metab* **20**, 991-1005 (2014).

40. D. S. Whittaker *et al.*, Circadian modulation by time-restricted feeding rescues brain pathology and improves memory in mouse models of Alzheimer's disease. *Cell Metab* **35**, 1704-1721 e1706 (2023).

41. A. Chaix, S. Deota, R. Bhardwaj, T. Lin, S. Panda, Sex- and age-dependent outcomes of 9-hour time-restricted feeding of a Western high-fat high-sucrose diet in C57BL/6J mice. *Cell Rep* **36**, 109543 (2021).

42. M. Ulgherait *et al.*, Circadian autophagy drives iTRF-mediated longevity. *Nature* **598**, 353-358 (2021).

43. K. Xie *et al.*, Every-other-day feeding extends lifespan but fails to delay many symptoms of aging in mice. *Nat Commun* **8**, 155 (2017).

44. M. E. Starr, H. Saito, Age-related increase in food spilling by laboratory mice may lead to significant overestimation of actual food consumption: implications for studies on dietary restriction, metabolism, and dose calculations. *J Gerontol A Biol Sci Med Sci* **67**, 1043-1048 (2012).

45. Y. D. Rathod, M. Di Fulvio, The feeding microstructure of male and female mice. *PLoS One* **16**, e0246569 (2021).

46. K. Gabel *et al.*, Effects of 8-hour time restricted feeding on body weight and metabolic disease risk factors in obese adults: A pilot study. *Nutr Healthy Aging* **4**, 345-353 (2018).

47. V. A. Acosta-Rodriguez *et al.*, Misaligned feeding uncouples daily rhythms within brown adipose tissue and between peripheral clocks. *Cell Rep* **43**, 114523 (2024).

48. G. Manzanares, G. Brito-da-Silva, P. G. Gandra, Voluntary wheel running: patterns and physiological effects in mice. *Braz J Med Biol Res* **52**, e7830 (2018).

49. V. S. Valentinuzzi, K. Scarbrough, J. S. Takahashi, F. W. Turek, Effects of aging on the circadian rhythm of wheel-running activity in C57BL/6 mice. *Am J Physiol* **273**, R1957-1964 (1997).

50. R. Garcia-Valles *et al.*, Life-long spontaneous exercise does not prolong lifespan but improves health span in mice. *Longev Healthspan* **2**, 14 (2013).

51. O. Froy, Circadian aspects of energy metabolism and aging. *Ageing Res Rev* **12**, 931-940 (2013).

52. C. Lopez-Otin, L. Galluzzi, J. M. P. Freije, F. Madeo, G. Kroemer, Metabolic Control of Longevity. *Cell* **166**, 802-821 (2016).

53. D. Gatfield, U. Schibler, Circadian glucose homeostasis requires compensatory interference between brain and liver clocks. *Proc Natl Acad Sci U S A* **105**, 14753-14754 (2008).

54. D. L. Palliyaguru *et al.*, Fasting blood glucose as a predictor of mortality: Lost in translation. *Cell Metab* **33**, 2189-2200 e2183 (2021).

55. J. C. Whitehead *et al.*, A clinical frailty index in aging mice: comparisons with frailty index data in humans. *J Gerontol A Biol Sci Med Sci* **69**, 621-632 (2014).

56. J. Martinez-Romero *et al.*, A hematology-based clock derived from the Study of Longitudinal Aging in Mice to estimate biological age. *Nat Aging* **4**, 1882-1896 (2024).

57. M. Magnani *et al.*, Effect of age on some properties of mice erythrocytes. *Mechanisms of Ageing and Development* **42**, 37-47 (1988).

58. D. L. Palliyaguru, J. M. Moats, C. Di Germanio, M. Bernier, R. de Cabo, Frailty index as a biomarker of lifespan and healthspan: Focus on pharmacological interventions. *Mech Ageing Dev* **180**, 42-48 (2019).

59. D. W. Lamming, Quantification of healthspan in aging mice: introducing FAMY and GRAIL. *Geroscience* **46**, 4203-4215 (2024).

60. R. Huang *et al.*, Multi-omics profiling reveals rhythmic liver function shaped by meal timing. *Nat Commun* **14**, 6086 (2023).

61. J. S. Pendergast, K. L. Branecky, R. Huang, K. D. Niswender, S. Yamazaki, Wheel-running activity modulates circadian organization and the daily rhythm of eating behavior. *Front Psychol* **5**, 177 (2014).

62. A. M. Schroeder *et al.*, Voluntary scheduled exercise alters diurnal rhythms of behaviour, physiology and gene expression in wild-type and vasoactive intestinal peptide-deficient mice. *J Physiol* **590**, 6213-6226 (2012).

63. I. Janssen, S. B. Heymsfield, R. Ross, Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc* **50**, 889-896 (2002).

64. E. Marzetti, C. Leeuwenburgh, Skeletal muscle apoptosis, sarcopenia and frailty at old age. *Exp Gerontol* **41**, 1234-1238 (2006).

65. C. Livelo, Y. Guo, G. C. Melkani, A Skeletal Muscle-Centric View on Time-Restricted Feeding and Obesity under Various Metabolic Challenges in Humans and Animals. *Int J Mol Sci* **24**, (2022).

66. Y. O. Henderson *et al.*, Late-life intermittent fasting decreases aging-related frailty and increases renal hydrogen sulfide production in a sexually dimorphic manner. *Geroscience* **43**, 1527-1554 (2021).

67. Y. Yamada *et al.*, Caloric Restriction and Healthy Life Span: Frail Phenotype of Nonhuman Primates in the Wisconsin National Primate Research Center Caloric Restriction Study. *J Gerontol A Biol Sci Med Sci* **73**, 273-278 (2018).

68. O. Arum, Z. A. Rasche, D. J. Rickman, A. Bartke, Prevention of neuromusculoskeletal frailty in slow-aging ames dwarf mice: longitudinal investigation of interaction of longevity genes and caloric restriction. *PLoS One* **8**, e72255 (2013).

69. A. E. Kane *et al.*, Impact of Longevity Interventions on a Validated Mouse Clinical Frailty Index. *J Gerontol A Biol Sci Med Sci* **71**, 333-339 (2016).

70. A. E. Kane *et al.*, Animal models of frailty: current applications in clinical research. *Clin Interv Aging* **11**, 1519-1529 (2016).

71. R. Lok, J. Qian, S. L. Chellappa, Sex differences in sleep, circadian rhythms, and metabolism: Implications for precision medicine. *Sleep Med Rev* **75**, 101926 (2024).

72. G. K. Paschos, R. Lordan, G. A. FitzGerald, Intersection of sex and circadian biology. *Current Opinion in Physiology* **45**, 100834 (2025).

73. Z. Bahadoran, P. Mirmiran, K. Kashfi, A. Ghasemi, Effects of time-restricted feeding (TRF)-model of intermittent fasting on adipose organ: a narrative review. *Eat Weight Disord* **29**, 77 (2024).

74. P. Regmi, L. K. Heilbronn, Time-Restricted Eating: Benefits, Mechanisms, and Challenges in Translation. *iScience* **23**, 101161 (2020).

75. P. Chen, B. Li, L. Ou-Yang, Role of estrogen receptors in health and disease. *Front*
Endocrinol (Lausanne) **13**, 839005 (2022).

76. V. M. Alvord, E. J. Kantra, J. S. Pendergast, Estrogens and the circadian system. *Semin*
Cell Dev Biol **126**, 56-65 (2022).

77. K. M. Hatcher, S. E. Royston, M. M. Mahoney, Modulation of circadian rhythms through
estrogen receptor signaling. *Eur J Neurosci* **51**, 217-228 (2020).

78. L. Zhu *et al.*, Estrogens prevent metabolic dysfunctions induced by circadian disruptions
in female mice. *Endocrinology* **156**, 2114-2123 (2015).

79. L. Talamanca, C. Gobet, F. Naef, Sex-dimorphic and age-dependent organization of 24-
hour gene expression rhythms in humans. *Science* **379**, 478-483 (2023).

80. J. A. Krizo, E. M. Mintz, Sex differences in behavioral circadian rhythms in laboratory
rodents. *Front Endocrinol (Lausanne)* **5**, 234 (2014).

81. A. A. Astafev, V. Mezhnina, A. Poe, P. Jiang, R. V. Kondratov, Sexual dimorphism of
circadian liver transcriptome. *iScience* **27**, 109483 (2024).

82. S. J. Mitchell *et al.*, Effects of Sex, Strain, and Energy Intake on Hallmarks of Aging in
Mice. *Cell Metab* **23**, 1093-1112 (2016).

83. A. E. Kane, D. A. Sinclair, J. R. Mitchell, S. J. Mitchell, Sex differences in the response to
dietary restriction in rodents. *Curr Opin Physiol* **6**, 28-34 (2018).

84. J. F. Lemaitre *et al.*, Sex differences in adult lifespan and aging rates of mortality across
wild mammals. *Proc Natl Acad Sci U S A* **117**, 8546-8553 (2020).

85. S. N. Austad, K. E. Fischer, Sex Differences in Lifespan. *Cell Metab* **23**, 1022-1033 (2016).

86. D. E. Harrison *et al.*, Acarbose, 17-alpha-estradiol, and nordihydroguaiaretic acid extend
mouse lifespan preferentially in males. *Aging Cell* **13**, 273-282 (2014).

87. R. A. Miller *et al.*, Rapamycin, but not resveratrol or simvastatin, extends life span of
genetically heterogeneous mice. *J Gerontol A Biol Sci Med Sci* **66**, 191-201 (2011).

88. R. Strong *et al.*, Nordihydroguaiaretic acid and aspirin increase lifespan of genetically
heterogeneous male mice. *Aging Cell* **7**, 641-650 (2008).

89. B. N. Gaskill *et al.*, Impact of nesting material on mouse body temperature and physiology.
Physiol Behav **110-111**, 87-95 (2013).

90. K. Kaikaew, J. Steenbergen, A. P. N. Themmen, J. A. Visser, A. Grefhorst, Sex difference
in thermal preference of adult mice does not depend on presence of the gonads. *Biol Sex*
Differ **8**, 24 (2017).

91. K. Kaikaew *et al.*, Sex difference in cold perception and shivering onset upon gradual cold
exposure. *J Therm Biol* **77**, 137-144 (2018).

92. C. Fernandez-Pena, A. Reimundez, F. Viana, V. M. Arce, R. Senaris, Sex differences in
thermoregulation in mammals: Implications for energy homeostasis. *Front Endocrinol*
(Lausanne) **14**, 1093376 (2023).

93. Z. Zhao *et al.*, Body temperature is a more important modulator of lifespan than metabolic
rate in two small mammals. *Nat Metab* **4**, 320-326 (2022).

94. B. Conti *et al.*, Transgenic mice with a reduced core body temperature have an increased life span. *Science* **314**, 825-828 (2006).

95. A. W. Fischer, B. Cannon, J. Nedergaard, Optimal housing temperatures for mice to mimic the thermal environment of humans: An experimental study. *Mol Metab* **7**, 161-170 (2018).

96. C. L. Karp, Unstressing intemperate models: how cold stress undermines mouse modeling. *J Exp Med* **209**, 1069-1074 (2012).

97. A. L. Ray *et al.*, The role of sex in the innate and adaptive immune environment of metastatic colorectal cancer. *Br J Cancer* **123**, 624-632 (2020).

98. E. G. Vichaya *et al.*, Sex differences in the behavioral and immune responses of mice to tumor growth and cancer therapy. *Brain Behav Immun* **98**, 161-172 (2021).

99. J. B. Rubin, The spectrum of sex differences in cancer. *Trends Cancer* **8**, 303-315 (2022).

100. M. K. E. Ougaard *et al.*, Murine Nephrotoxic Nephritis as a Model of Chronic Kidney Disease. *Int J Nephrol* **2018**, 8424502 (2018).

101. United States Renal Data System. (National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD).

102. G. J. Shim, L. L. Kis, M. Warner, J. A. Gustafsson, Autoimmune glomerulonephritis with spontaneous formation of splenic germinal centers in mice lacking the estrogen receptor alpha gene. *Proc Natl Acad Sci U S A* **101**, 1720-1724 (2004).

103. J. H. Catterson *et al.*, Short-Term, Intermittent Fasting Induces Long-Lasting Gut Health and TOR-Independent Lifespan Extension. *Curr Biol* **28**, 1714-1724 e1714 (2018).

104. Y. Yang, D. Liu, Impacts of time-restricted feeding on middle-aged and old mice with obesity. *J Physiol*, (2024).

105. P. Domaszewski *et al.*, Effect of a six-week times restricted eating intervention on the body composition in early elderly men with overweight. *Sci Rep* **12**, 9816 (2022).

106. O. Hahn *et al.*, A nutritional memory effect counteracts benefits of dietary restriction in old mice. *Nat Metab* **1**, 1059-1073 (2019).

107. A. C. Hofacker, M. Knop, S. Krauss-Etschmann, T. Roeder, Time-Restricted Feeding Promotes Longevity and Gut Health Without Fitness Trade-Offs. *FASEB J* **39**, e70627 (2025).

108. M. J. Schafer *et al.*, Late-life time-restricted feeding and exercise differentially alter healthspan in obesity. *Aging Cell* **18**, e12966 (2019).

109. M. Das *et al.*, Time-Restricted Feeding Attenuates Metabolic Dysfunction-Associated Steatohepatitis and Hepatocellular Carcinoma in Obese Male Mice. *Cancers (Basel)* **16**, (2024).

110. E. B. Parr, B. L. Devlin, B. E. Radford, J. A. Hawley, A Delayed Morning and Earlier Evening Time-Restricted Feeding Protocol for Improving Glycemic Control and Dietary Adherence in Men with Overweight/Obesity: A Randomized Controlled Trial. *Nutrients* **12**, (2020).

111. K. Gabel, S. Cienfuegos, F. Kalam, M. Ezpeleta, K. A. Varady, Time-Restricted Eating to
Improve Cardiovascular Health. *Curr Atheroscler Rep* **23**, 22 (2021).

112. P. Pati *et al.*, Time-restricted feeding reduces cardiovascular disease risk in obese mice.
JCI Insight **10**, (2025).

113. M. Milan *et al.*, Time-restricted feeding improves aortic endothelial relaxation by
enhancing mitochondrial function and attenuating oxidative stress in aged mice. *Redox Biol*
73, 103189 (2024).

114. R. A. Miller *et al.*, Glycine supplementation extends lifespan of male and female mice.
Aging Cell **18**, e12953 (2019).

115. N. Ogiso *et al.*, Biological characteristics of age-related changes in C57BL/6 mice sub-
strains in the national center for geriatrics and gerontology aging farm. *Exp Anim* **74**, 229-
238 (2025).

116. W. J. Schwartz, P. Zimmerman, Circadian timekeeping in BALB/c and C57BL/6 inbred
mouse strains. *J Neurosci* **10**, 3685-3694 (1990).

117. M. Pfeffer, C. von Gall, H. Wicht, H. W. Korf, The Role of the Melatoninergic System in
Circadian and Seasonal Rhythms-Insights From Different Mouse Strains. *Front Physiol*
13, 883637 (2022).

118. G. Banks *et al.*, Genetic background influences age-related decline in visual and nonvisual
retinal responses, circadian rhythms, and sleep. *Neurobiol Aging* **36**, 380-393 (2015).

119. S. M. Siepka, J. S. Takahashi, Methods to record circadian rhythm wheel running activity
in mice. *Methods Enzymol* **393**, 230-239 (2005).

120. A. A. Bachmanov, D. R. Reed, G. K. Beauchamp, M. G. Tordoff, Food intake, water
intake, and drinking spout side preference of 28 mouse strains. *Behav Genet* **32**, 435-443
(2002).

121. D. R. Reed, A. A. Bachmanov, M. G. Tordoff, Forty mouse strain survey of body
composition. *Physiol Behav* **91**, 593-600 (2007).

122. J. W. Cooley, J. W. Tukey, An Algorithm for the Machine Calculation of Complex Fourier
Series. *Mathematics of Computation* **19**, 297-301 (1965).

123. R. B. Blackman, J. W. Tukey, The measurement of power spectra from the point of view
of communications engineering — Part I. *The Bell System Technical Journal* **37**, 185-282
(1958).

124. C. Wang, Q. Li, D. T. Redden, R. Weindruch, D. B. Allison, Statistical methods for testing
effects on "maximum lifespan". *Mech Ageing Dev* **125**, 629-632 (2004).

Acknowledgements: Thanks to G. Martinez, K. Brown, C. Joseph, G. Flowers, A. Torres, A.

Herrera, R. Lewis, and L. Thomas, for assisting with the mouse husbandry and experiments. To S.

McPherson, K. Goedde, K. Auldridge, and S. Elias Perryman of the Animal Resource Center

Diagnostics Lab for their assistance with the gross necropsies. To veterinarians Dr. S. Lewis and

Dr. C. Jones for providing guidance on mouse health care. To S. Dixon for providing

administrative support. Additional thanks to M. Wellems at Phenome Technologies and D. Ferster

at Actimetrics for the feeders and software system. This work was supported by the Howard

Hughes Medical Institute awarded to JST, the National Institute of Health (NIH/NIA AG045795

to JST and CBG; NIH/NIA AG072736 to JST and CBG; NIH/NIGMS GM127122 to CBG; NIH

1T32HL138438 to JST and SEI), and the Milky Way Research Foundation MWRF210823

awarded to JST and CBG.

This research was supported in part by the Intramural Research Program of the National Institutes

of Health (NIH). The contributions of the NIH author(s) were made as part of their official duties

as NIH federal employees, are in compliance with agency policy requirements, and are considered

Works of the United States Government. However, the findings and conclusions presented in this

paper are those of the author(s) and do not necessarily reflect the views of the NIH or the U.S.

Department of Health and Human Services.

Author Contributions: SEI, NJS, VAR, CBG, and JST conceptualized the study, developed the

experimental design, and oversaw all aspects of project execution and supervision. MW-C

performed all histopathological assessments and interpretations. SEI, NJS, and VAR conducted all

remaining experimental work, including animal handling, data curation, and statistical analyses.

SEI, NJS, VAR, CBG, and JST wrote and revised the manuscript with additional edits from MW-

C. Funding acquisition was provided by CBG and JST.

Competing interests: The authors declare no competing interests.

Data and materials availability: Data are available in the main text or the supplementary materials

Supplementary Materials:

Materials and Methods

Figures S1 to S17

Tables S1 to S7

Materials and Methods:

Animals and Housing Conditions

For a minimum detection of 10% life extension with 80% power a minimum of 96 animals were needed in the control group and 72 in each of the two TRF groups (*114*). Extra mice were included in each group to account for any that may not adapt to the feeder/diet or for non-aging-related deaths.

264 male and 264 female C57BL/6J mice 6 weeks of age were ordered from Jackson Laboratories, Bar Harbor, ME. We selected C57BL/6J mice because they are widely used research model which exhibit robust circadian rhythms and well-known aging phenotypes (*115-118*). Further, using a single, well-characterized strain maximized statistical power and lifespan resolution in our large cohort (*114*). As done previously, starting at 2 months of age the mice were: **(i)** individually housed in standard polycarbonate cages with stainless steel running wheels inside isolation cabinets under

light:dark (LD) of 12:12 hours and ambient building temperature 72-78°F, **(ii)** fed 300mg pellets

of purified diet (F0075, Bio-Serv) using automated feeders with water provided *ad libitum* (AL),

and **(iii)** cage changed every 21 days (*9, 10, 119*). Nesting material and igloos were excluded,

besides shavings, to prevent wheel blockages. The Institutional Animal Care and Use Committee

(IACUC) of the University of Texas Southwestern Medical Center approved the animal protocol

(APN 2015-100925), which has been renewed every 3 years (in 2018, 2021, and 2024).

The purified diet (F0075, Bio-Serv) is composed of 18.7% protein, 5.6% fat, 4.7% fiber, and 59.1%

carbohydrates, similar to regular mouse chows. Major ingredients include Sucrose. Dextrose,

Casein, Corn Oil, Mineral Mix, Cellulose, Corn Syrup, Calcium Silicate, Vitamin Mix,

Magnesium Stearate, Choline Bitartrate, DL-Methionine, L-Cystine, Ascorbic Acid, Vitamin E

Acetate, and tBHQ.

An additional cohort of C57BL/6J animals (N=48/feeding condition/sex) were included in this

study in order to undergo more invasive assays. These mice were randomly selected (N=8-

12/group/age point) for analysis of metabolism in metabolic chambers, fasting blood glucose

levels, glucose tolerance, biomarkers of inflammation and metabolism and hematology. These

mice were not included in the longitudinal health measurements or the assessment of lifespan.

Time-Restricted Feeding (TRF) Regimens

At 2 months of age, unmated mice were placed in individual cages as described above and fed *ad*

libitum (AL) for 8 weeks using the automated feeder system (*9, 10*). At 4 months of age, the mice

were divided into three feeding groups for the remainder of the experiment: **(i)** 108 of each sex

continued in AL as a control group, **(ii)** 78 of each sex in a 12-hour TRF group where the automated

feeder restricted food dispensing to the 12-hour night (12h-TRF, ZT12-24), and **(iii)** 78 of each

sex in an 8-hour TRF group with food dispensing restricted to the middle 8 hours of the night (8h-

TRF, ZT14-22). These feeding windows were chosen as they have been previously validated and

demonstrated to confer health benefits in mice (*25, 29*). In addition to the timing of the feeding

window, the feeders were also programmed with a 10-minute delay between pellet dispensing to

reduce hoarding behavior. Regardless of the feeding group, all mice had access to a maximum of

22 pellets/day or 6.6 g of food. In this study, no group exceeded 6.6 g daily consumption. Previous

studies have shown that C57BL/6J mice consume 4-5g of food per day (*120*), so this amount of

food was chosen to ensure mice were not calorically restricted.

Exclusion of Hoarded Food

Mice exhibit a natural tendency to hoard food within the cage, especially later in life or when ill

(*44*). In all TRF groups, mice identified as hoarders were flagged, and hoarded food was removed

from the cage on a weekly basis. For mice on veterinary health report, hoarded food was removed

at their twice weekly full health check. The total amount of hoarded food was quantified by weight

(g) at each 21-day cage change. For AL mice, hoarded food was allowed to remain in the cage

until the scheduled cage change, at which point it was collected and quantified. In all groups, the

total hoarded food weight was divided by 21 days to estimate an average daily hoarding amount,

which was then subtracted from the average daily food retrieval recorded by the feeder.

Daily Monitoring of Feeding and Wheel-Running Activity

As before, the feeders were set and continuously recorded quantity and timing of food intake via ClockLab Chamber Control Software v3.401 (Actimetrics Inc., Wilmette, IL, USA) (*9, 10*). The feeding data presented in Figure 1B and Supplemental figures S2 and S3 represent food consumption with hoarding removed. All other feeding figures represent the quantity and pattern of pellet taking from the feeder, not excluding hoarded food. Wheel-running activity was also continuously recorded as before using an updated version of ClockLab Data Acquisition System v3.604 (Actimetrics Inc., Wilmette, IL, USA).

Body Weight and Composition Measurements

The body weight of each mouse was measured every 21 days during cage change (morning, ZT4-7) throughout its lifespan. Body composition measurements of lean and fat mass (g) were assessed every 6 months during cage change (ZT4-7) using a 100H-EchoMRI Body Composition Analyzer (EchoMRI, Houston, TX, USA).

Promethion Metabolic Chambers

These tests were performed in a follow-up cohort of mice. Mice were randomly selected.

At 19 months of age, mice received body composition measurements (EchoMRI) and then were individually housed in metabolic cages (Promethion, Sable Systems International) modified with automated feeders to maintain the same feeding regimens. These cages were housed in light-tight, temperature-controlled cabinets. Mice were acclimatized for 2-3 days at room temperature (25°C) before recording for an additional 5-6 days at room temperature. Cabinet temperature was then raised to thermoneutrality (30°C) for 5 days.

Fasting Glucose and Glucose Tolerance Testing

These tests were performed in a follow-up cohort of mice. Mice were randomly selected from the cohort at each age point.

For fasting glucose, cages for all mice to be tested were changed out fresh the day before. Feeders for all groups were shut off at ZT21.5 to allow the mice 30 minutes to consume the last pellet dropped and officially begin their 12h fast at ZT22. Mice were checked again at ZT0 to ensure no pellets remained in the cage or available in the pellet chute. At ZT10, a drop of blood was measured from the tail using an Accu-Check Glucose Meter.

For glucose tolerance testing, cages were changed fresh and mice fasted as described above. At ZT10, mice were injected with a 20% glucose solution at a dose of 0.1mL per 25g of body weight. A drop of blood from the tail was measured for glucose at the time of injection (0 min) and then at 15-, 30-, 45-, 60-, and 120-min post-injection.

Biomarkers of Metabolism and Inflammation

These markers were measured in a follow-up cohort of mice. Mice were randomly selected from the cohort at each age point.

Mice were switched to fresh cages and were fasted for 12h as described above for fasting glucose. At ZT10, 70-100μL of blood was collected via submandibular bleed into K_2 EDTA coated collection tubes (RAM Scientific Safe-T-Fill™ Capillary Blood Collection Systems) with cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail (Roche) and DPP-IV (Millipore).

1001 Blood was immediately kept on ice and then spun down at 2000g for 10 min at 4°C. The upper

1002 layer of plasma was transferred to a new tube and stored at -80°C until use.

1003 20-25µL of plasma was then measured for leptin, insulin, IL-1β, IL-6, IL-10, TNFα, MCP-1, and

1004 BDNF using a U-PLEX Adipokine Combo 1 (mouse) assay kit (Meso Scale Discovery).

1005

Hematology

Measurements for red blood cells, white blood cells, and platelets were performed in a follow-up

cohort of mice. Mice were randomly selected from the cohort at each age point.

Mice were switched to fresh cages and were fasted for 12h as described above for fasting glucose.

At ZT10, 25-30µL of blood was collected via submandibular bleed into K_2 EDTA coated

collection tubes (RAM Scientific Safe-T-Fill™ Capillary Blood Collection Systems) with

cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail (Roche) and DPP-IV (Millipore). 20µL

of whole blood was then measured in a Hemavet 950FS (Drew Scientific) at room temperature

within 2-4 hours of collection.

1015

Frailty Scoring

Every 6 months the mice were assessed for frailty, or physical vulnerabilities, according to 31

parameters of aging modified from Whitehead et al. 2014 (*55*). These parameters include a range

of physiological evaluations, scoring coat condition, ocular and auditory responses, physical and

musculoskeletal state, respiration rate, body weight, and temperature. Each assessment was scored

either 0=absent, 0.5=mild, or 1=severe. For scoring Body Condition Scores, 0.5 was used for either

a BCS of 2 or 4, and 1 was for a BCS of 1 or 5. For body weight and body temperature scores only,

these were determined by averaging the group measurements and then determining how many

standard deviations (SD) each individual mouse was from their age-, sex- and feeding-matched

group mean. If <1 SD, mouse scored as 0 for that parameter, 1-1.99 SD = 0.25, 2-2.99 SD = 0.5,

3-3.99 = 0.75, and >4 SD = 1. The scores were totaled to calculate an overall index score.

1027

Health Monitoring and Survival Study

Mice were physically health checked every 21 days during cage change and visually inspected

every 10 days during refilling of the automated feeders. Using the feeding and wheel monitoring

software, mice were also checked virtually every day and only physically health checked if feeding

fell to <5 pellets per/day and/or wheel running fell below set thresholds (1.14 counts/min within

24hr, reduced by 10% every 6 months as the mice age) as performed previously (*10*).

If any mouse was found to have a mild/moderate health ailment (e.g. dermatitis, external tumor,

abdominal mass, head tilt) or a body condition score < 3 (Scored 1-5 with: 1 being emaciated, 3

ideal condition, and 5 obese) it was placed on veterinary health report. These records were also

used to generate Figures 4D-F. Mice on health report were physically health checked weekly by

the UT Southwestern Animal Resource Facility (ARC) veterinary staff and twice a week by our

own laboratory staff. In order to obtain accurate but humane lifespan data, mice were monitored

carefully for euthanasia criteria that would indicate imminent death (moribund) or unacceptable

levels of pain (analgesics were not used to avoid confounding lifespan results). Criteria were

determined in conjunction with the ARC veterinarians according to AAALAC guidelines and

include: Non-responsive to touch, hypothermic or cold to touch, slow or labored breathing, failure

or inability to eat/drink after moist chow is given, a body condition score of 1, body weight loss

of >20% from baseline or between cage changes, a broken/fractured/dislocated limb, head tilt that

is resulting in an animal being unable to maintain sternal body position which would affect their

ability to consume food and water, severe eye protrusion/infection, severe dermatitis, grade 3 rectal

or penile prolapse, urinary obstruction, or a subcutaneous tumor >2cm.

The number and date of those euthanized or found dead across the feeding groups and sexes were

recorded and used to generate the Kaplan-Meier survival curves. A total of 35 animals (35/528)

were censored from the survival curve if death occurred earlier than 6 months or due to non-aging

related injuries or death: 9 AL females, 6 12h-TRF females, 6 8h-TRF females, 9 AL males and 5

8h-TRF males.

1054

1055   **Necropsy and Histopathology**

1056   Mice euthanized or found dead were submitted for gross necropsy and blinded histopathological

1057   analysis was performed, as done previously (*10*). Less than 10% of the animals submitted were

1058   too autolyzed to analyze. Disease types were tallied together by tissue type, sex, and feeding group.

1059

Healthspan Index

Throughout the experiment we recorded thirteen health measures including: wheel-running

activity to determine overall activity level and diurnal amplitude, food consumption to determine

overall level and diurnal amplitude, 90% feeding duration, maximum fasting times, body weight,

composition of lean and fat mass as a percentage of body weight, body condition score, frailty

index, and the onset and duration of veterinary health reports.

Healthspan index was calculated by summing the calculated scaled values of each of the thirteen

health measures. In the case of body condition score, an absolute difference from BCS=3 was

calculated. For body fat percentage, an absolute difference in value from 24.45% was calculated.

The value 24.45% was used based on previous literature showing the average body fat percentage

across multiple healthy mouse strains at the age of 16 weeks (*121*). All measures were then

normalized to a mean of 0 and a standard deviation of 1, within each measure. Certain values were

then inverted, given their context, in order to have positive numbers representing a healthier state,

and negative values representing an unhealthy state. For example, "Days on Health Report" is

inverted, as higher values would be less healthy, and lower values are healthier. All normalized

values were then summed together at 21-day intervals.

1076

1077 **FAMY**

1078 FAMY was calculated using the lifetime of frailty scores collected for each mouse and as described

1079 in Lamming 2024 (*59*).

1080

Statistical Analyses

All measurements represent repeated measures from the same animals over time, with the

exception of data shown in Figs. S7–S10 and S13–S15, which were collected from a separate

cohort using different mice at each age point. Sample sizes (N) for each experiment are indicated

in the corresponding figure legends.

As performed previously, all feeding and wheel-running activity was collected using Chamber

Control and ClockLab, respectively (Actimetrics, Inc) (*10*). Each point on the longitudinal profiles

represents a 21-day mean. 24h profiles were binned by hour, with each age point reflecting a 14-

day mean (taken from the middle of each 21-day cage cycle to avoid behavioral changes due to

cage change stress). For each mouse, each 14-day bin was then analyzed by 1-dimensional Discrete

Fast Fourier Transform (FFT) using the mixed-radix algorithm for real-valued input (SciPy

v1.12.0; scipy.fft.rfft) (*122*). The Blackman window function is applied to the data prior to FFT

analysis to inhibit spectral leakage (*123*). The power of each frequency component is then

normalized, with the total normalized power of all frequency components equaling 100. The

normalized power from the 24h frequency component is then taken as the diurnal amplitude of

either food intake behavior or running wheel activity. This value represents the percentage of

contribution of the 24h frequency component, to the entire 14-day behavioral period. For example,

if the diurnal amplitude for food intake is 21, it indicates that 21% of the observed variation in

food intake can be directly attributed to the animal's consistent 24-hour daily cycle. Group-wise

statistics were then performed to determine the overarching impact of each intervention, as

described below.

For the calculation of the 90% feeding window, food intake data was similarly partitioned into 3-

week intervals, with each sampling point consisting of a 14-day window. Only windows containing

a minimum of 84 feeding events (averaging 6 events per day, or ~1.8g of intake) were included in

the final density analysis to ensure robust statistical estimation. Because time-of-day is a periodic

variable, we employed a von Mises distribution (the circular analogue of the Normal distribution)

to estimate the probability density of feeding activity. We utilized the vonmises.pdf method (SciPy

v1.12.0), to compute the circular distribution, where the concentration parameter κ, was defined

by a fixed bandwidth of 0.04363 radians, representing approximately 10 minutes of smoothing.

To define the primary feeding window, we identified the shortest continuous arc on the circle

containing 90% of the total probability mass. This method accounts for feeding windows that cross

over the ZT24/ZT0 boundary.

Cumulative health report and survival curves were analyzed using Log-rank Mantel Cox and

Fisher's exact test, health report type and necropsy report disease type percentages analyzed by

contingency tables and Fisher's exact test and, and all other plots analyzed using two-way ANOVA

or type III ANOVA with Tukey's or Holm's post-hoc analyses (*124*). All statistical tests were two-

sided. Normality was assumed but not formally tested. Data plots were generated and statistically

analyzed by Prism 10 (Graphpad Software Inc), Python, and R software, using custom-written

scripts and standard libraries.

 Supplemental Figures and Legends:

Figure S1: Food taking and hoarding behavior: 300mg pellets are dispensed from the feeder

one at a time for the mouse to retrieve. In females and males: (**A**) Longitudinal profiles of daily

food taken from the feeder including uneaten (hoarded) food found in the cage (g/day) and the

equivalent energy intake (kcal/day). (**B**) Lifetime averages of the percent (%) of food taken from

the feeder within or outside of the TRF window. (**C**) Longitudinal profiles of hoarded food

removed from cages as a % of the total food taken from the feeder. A and C: *Ad libitum* (AL)

baseline feeding (Gray area). Time-restricted feeding (TRF) (White area). Mean of 21 days ± SEM

(shaded regions). Females N=71-103 per group. Males N=71-97 per group.

Figure S2: Food intake by monthly age epochs: (**A**) In females and (**B**) males, box plots of food intake with hoarding removed (g/day) at 3 and 6 months of age and then every 6 months following. Food intake per mouse averaged every 21 days. Any caloric restriction (CR) relative to the AL control at each age epoch shown as %. Two-way ANOVA, Tukey's post-hoc. *P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤0.0001.

Figure S3: Food intake relative to body weight by monthly age epochs: (**A**) In females and (**B**) males, boxplots of food intake (g/day) with hoarding removed relative to body weight (g) at 6 months of age and then every 6 months following. Two-way ANOVA, Tukey's post-hoc. *P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤ 0.0001.

Figure S4: Wheel activity by monthly age epochs: (A) In females and (B) males, boxplots of daily wheel revolutions (counts/min) at 3 and 6 months of age and then every 6 months following. Wheel running per mouse averaged across 21 days at each age point. Two-way ANOVA, Tukey's post-hoc. *P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤0.0001.

Figure S5: Wheel activity relative to body weight by monthly age epochs: (**A**) In females and

(**B**) males, boxplots of wheel revolutions (counts/min) relative to body weight (g) at 6 months of

age and then every 6 months following. Two-way ANOVA, Tukey's post-hoc. *P ≤ 0.05; **P ≤

0.01; ***P ≤ 0.001; ****P ≤0.0001.

Figure S6: Body weight by monthly age epochs: (A) In females and **(B)** males, boxplots of body

weight (g) at 3 and 6 months of age and then every 6 months following. Two-way ANOVA,

Tukey's post-hoc. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$.

A

Respiratory Exchange Ratio - Across Experiment

Respiratory Exchange Ratio - 24h Profiles, 3-day average

B

Energy Expenditure - Across Experiment

Energy Expenditure - 24h Profiles, 3-day average

Figure S7: Measures of metabolism at 19 months of age in a follow-up cohort of mice: (A)

(*Top*) Respiratory exchange ratios (RER; VCO2/VO2) in females and males throughout a 11-day

recording in Promethion metabolic chambers. 6 days recorded at room temperature (25°C; white

shaded area), and 5 days were recorded at thermoneutrality (30°C; red shaded area). (*Bottom*) 24h

profiles of RER averaged from the last 3 days of recording in each temperature. 12h light (white

shaded area), 12h dark (gray shaded area). Means ± SEM (shading) presented in 30-minute bins.

(**B**) (*Top*) Energy expenditure (EE; kcal/h) in females and males throughout the 11-day recording.

6 days recorded at room temperature (25°C; white shaded area), and 5 days were recorded at

thermoneutrality (30°C; red shaded area). Measurements are presented in 30-minute bins. (*Bottom*)

24h profiles of EE from the last 3 days of recording in each temperature. 12h light (white shaded

area), 12h dark (gray shaded area). Means ± SEM (shading) presented in 30-minute bins. A and

B: Type III ANOVA (Wald χ^2), and Holm's post-hoc. *P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P

≤0.0001.

	Female Room Temperature					
1. INTERACTION MODEL (Slope Consistency)						
	Df	Sum Sq	Mean Sq	F value	Pr(>F)	signif
Lean	1	0.02322	0.02322	5.97928	0.02729	*
feeding	2	0.02204	0.01102	2.83776	0.09011	ns
Lean:feeding	2	0.02754	0.01377	3.54642	0.0548	ns
Residuals	15	0.05825	0.00388			

	Male Room Temperature					
1. INTERACTION MODEL (Slope Consistency)						
	Df	Sum Sq	Mean Sq	F value	Pr(>F)	signif
Lean	1	0.06713	0.06713	12.7309	0.00182	**
feeding	2	0.02161	0.01081	2.04916	0.15383	ns
Lean:feedin	2	0.00929	0.00464	0.88045	0.42935	ns
Residuals	21	0.11074	0.00527			

2. ANCOVA MODEL (Type III SS)					
	Sum Sq	Df	F value	Pr(>F)	signif
(Intercept)	0.00554	1	1.09845	0.30928	ns
Lean	0.01599	1	3.16844	0.09295	ns
feeding	0.02204	2	2.1836	0.14321	ns
Residuals	0.08579	17			

2. ANCOVA MODEL (Type III SS)					
	Sum Sq	Df	F value	Pr(>F)	signif
(Intercept)	2.8E-05	1	0.00528	0.94269	ns
Lean	0.04917	1	9.42346	0.00542	**
feeding	0.02161	2	2.07069	0.14897	ns
Residuals	0.12002	23			

	Female Thermoneutrality					
1. INTERACTION MODEL (Slope Consistency)						
	Df	Sum Sq	Mean Sq	F value	Pr(>F)	signif
Lean	1	0.02085	0.02085	8.61881	0.01022	*
feeding	2	0.02061	0.0103	4.25862	0.0343	*
Lean:feeding	2	0.01696	0.00848	3.50538	0.05635	ns
Residuals	15	0.03629	0.00242			

	Male Thermoneutrality					
1. INTERACTION MODEL (Slope Consistency)						
	Df	Sum Sq	Mean Sq	F value	Pr(>F)	signif
pre_lean	1	0.02215	0.02215	9.1224	0.00651	**
feeding	2	0.00477	0.00239	0.98324	0.39067	ns
pre lean:fee	2	0.00572	0.00286	1.17758	0.32755	ns
Residuals	21	0.05099	0.00243			

2. ANCOVA MODEL (Type III SS)					
	Sum Sq	Df	F value	Pr(>F)	signif
(Intercept)	0.00416	1	1.32941	0.26487	ns
Lean	0.01125	1	3.59171	0.0752	ns
feeding	0.02061	2	3.28915	0.06201	ns
Residuals	0.05325	17			

2. ANCOVA MODEL (Type III SS)					
	Sum Sq	Df	F value	Pr(>F)	signif
(Intercept)	0.00285	1	1.1577	0.2931	ns
pre lean	0.01371	1	5.55981	0.02726	*
feeding	0.00477	2	0.96829	0.39468	ns
Residuals	0.05671	23			

Figure S8: ANCOVA of TRF effects on energy expenditure: 24h mean EE plotted vs lean mass for females and males. Homogeneity of regression slopes was confirmed for all groups (test of Feeding × Lean Mass interaction, all p > 0.05, ns), validating the use of ANCOVA to test for significance of intercepts.

Figure S9: Fasting glucose levels and glucose tolerance tests in follow-up mouse cohort: In females and males: (**A**) Bar graphs of 12h fasting glucose levels (mg/dL) collected at zeitgeber

time (ZT) 10, (**B**) glucose tolerance tests (GTT) performed starting at ZT10, and (**C**) area under the curve (AUC) for the GTTs tested at 6, 12, and 18 months of age. Means ± SEM (bars). Two-way ANOVA, Tukey's post-hoc. *P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤0.0001. N=9-12 per group and sex.

Figure S10: Circulating levels of adipokines from blood plasma in follow-up mouse cohort.

In females and males, dot plots of circulating levels of interleukin (IL)-1β, IL-6, IL-10, tumor

1176   necrosis factor α (TNFα), brain-derived neurotrophic factor (BDNF), monocyte chemoattractant

1177   protein (MCP)-1, leptin, and insulin. Plasma was collected at ZT10 from 12h fasted blood at 6, 12,

1178   and 18 months of age. Median with interquartile range. Two-way ANOVA, Tukey's post-hoc. *P

1179 ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤ 0.0001. N=8-12 per group and sex.

AL
12h-TRF
8h-TRF

Nasal Discharge vs Age (Months)

Diarrhea vs Age (Months)

Microphthalmia vs Age (Months)

Cataracts vs Age (Months)

Body Condition Score vs Age (Months)

***
**
****
***

1 1

22 8 5
43 26 20
19 14 17

Body Temperature Score vs Age (Months)

1 1 1
1 3
2 2 3
3 1 3
1 2 4
2 1 1
22 16 15
22 21 10
20 22 17
11 10 11

Body Weight Score vs Age (Months)

2 4 3
3 4
1 3 1
1 2
28 20 15
20 26 19
18 12 15
13 15 13

Figure S11: Individual frailty parameter scores in females. Dot plots of frailty scores every 6

months of age in females for each of the 31 parameters modified from (*55*). Most parameters

scored as 0 for no, 0.5 for mild, and 1 for severe deficit. Body temperature and body weight

scores were scored 0, .25, .5, .75, or 1 with higher values representing more severe deviation of

the parameter from the group mean. N values shown in graph for any mice with scores >0. Mean

shown as dashed line. Two-way ANOVA, Tukey's post-hoc. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq$

0.001; ****$P \leq 0.0001$. Total N: AL, N=44-79. 12h-TRF, N=34-74. 8h-TRF, N=40-74.

1193

Figure S12: Individual frailty parameter scores in males. Dot plots of frailty scores every 6

months of age in males for each of the 31 parameters modified from (*55*). Most parameters scored

as 0 for no, 0.5 for mild, and 1 for severe deficit. Body temperature and body weight scores were

scored 0, .25, .5, .75, or 1 with higher values representing more severe deviation of the parameter

from the group mean. N values shown in graph for any mice with scores >0. Mean shown as dashed

line. Two-way ANOVA, Tukey's post-hoc. *P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤0.0001.

Total N: AL, N=23-79. 12h-TRF, N=22-78. 8h-TRF, N=39-76.

Figure S13: Red blood cell measures with age in follow-up mouse cohort. (A) In females and

(**B**) males, bar graphs of red blood cell (RBC) counts (M/µL), hemoglobin levels (g/dL),

hematocrit (HCT, percentage of RBCs in total blood volume) (%), mean corpuscular volume

(MCV, average size of RBCs) (fL), mean corpuscular hemoglobin (MCH, average amount of

hemoglobin per RBC) (pg), and mean corpuscular hemoglobin concentration (MCHC, average

concentration of hemoglobin within the volume of the RBC) (g/dL). Measured at ZT10 from 12h

fasted blood at 6, 12, and 18 months of age. Mean ± SEM (bars). Two-way ANOVA, Tukey's

post-hoc. *$P \leq 0.05$; **$P \leq 0.01$; ***$P \leq 0.001$; ****$P \leq 0.0001$. N=11-12 per group and sex.

Figure S14: Platelet counts with age in follow-up mouse cohort. In females and males,

platelet counts (K/μL). Measured at ZT10 from 12h fasted blood at 6, 12, and 18 months of age.

Mean ± SEM (bars). Two-way ANOVA, Tukey's post-hoc. *P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001;

****P ≤0.0001. N=11-12 per group and sex.

Figure S15: White blood cells measures with age in follow up mouse cohort. (A) In females

1215   and (**B**) males, bar graphs of total white blood cell (WBC) counts (K/µL) and counts for each

1216   WBC sub-type including neutrophils, lymphocytes, monocytes, eosinophils, and basophils.

1217   Measured at ZT10 from 12h fasted blood at 6, 12, and 18 months of age. Mean ± SEM (bars).

1218   Two-way ANOVA, Tukey's post-hoc. *P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤0.0001.

1219   N=11-12 per group and sex.

Figure S16: Repeat experiments show 12h-TRF does not extend lifespan in males: Kaplan-Meier survival curves and day median lifespan reached in a pilot experiment vs our current experiment. Log-Rank Mantel-Cox Test for significant difference in overall survival TRF vs AL. Pilot experiment: AL, N=43. 12h-TRF, N=33. Current experiment: AL, N=99. 12h-TRF, N=78. 8h-TRF, N=73.

A

B

Figure S17: Healthspan as measured by Frailty-Adjusted Mouse Years (FAMY). In females

and males: (**A**) FAMY (years) is another method to measure healthspan which utilizes frailty

scores as described in (*59*). Higher FAMY = more healthy years. Mean ± SEM (bars). Type III

ANOVA, Holm's post-hoc. *P ≤ 0.05; **P ≤ 0.01; ***P ≤ 0.001; ****P ≤0.0001. (**B**) Correlation

plots of FAMY vs lifespan in years. Correlation coefficients (r) and coefficients of determination

shown (r^2).