1   Covert infections with a honeybee virus induce fitness costs in ant colonies
2

3   Lai Ka Lo[1,2], Ashmita Baruah[4], Robert J. Paxton[2,3], Yuko Ulrich[1,2]
4
Affiliations:
[1] Lise Meitner Research Group Social Behaviour, Max Planck Institute for Chemical Ecology,
Jena, Germany
[2] German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig,
Germany
[3] General Zoology, Institute for Biology, Martin Luther University Halle-Wittenberg, Halle
(Saale), Germany
[4] Indian Institute of Science Education and Research, Pune, India

Keywords: clonal raider ant, ABPV, honey bee virus, virulence, spillover

Abstract:
Acute Bee Paralysis Virus (ABPV) is a key driver of honey bee colony losses that has been
increasingly reported in non-bee hosts in the past decades. Ants have long been hypothesised
to act as viral reservoirs, but most evidence comes from field surveys, and experimental tests
are still scarce. Here, we combined survival and transmission assays, viral load
measurements, viral replication assays, and host immune gene expression analyses to test
whether the clonal raider ant *Ooceraea biroi* can harbour and transmit ABPV. ABPV-injected
ants showed delayed development and elevated mortality. Although we found no evidence of
viral replication, the virus was able to spread among colony members and viral particles
persisted in colonies for several days. These results show that ants can acquire ABPV, incur
fitness costs, and pass the virus within their colonies, suggesting that they may act as
incidental viral reservoirs. By maintaining and disseminating honey bee viruses, even without
supporting replication, ants could contribute to their environmental persistence and spillover
across species.

30
31

1. Introduction

Honey bees are vital pollinators for ecosystems and agriculture, yet highly vulnerable to human activities and environmental changes. Over the past two decades, substantial winter colony losses of western honeybees, *Apis mellifera*, have been reported worldwide [1]. Several iflaviruses and dicistroviruses have emerged as major drivers of colony losses in both managed and wild populations [1–5]. While these viruses can persist covertly at low titres without causing apparent symptoms, the introduction of the ectoparasitic mite *Varroa destructor* (henceforth: varroa) has facilitated viral spread and turned asymptomatic infections into lethal diseases [6] by injecting viruses into the bee haemolymph [7].

Acute Bee Paralysis Virus (ABPV), a +ssRNA virus in the family *Dicistroviridae* [8,9], has been detected repeatedly in collapsing and *Varroa*-infested colonies and is considered a significant driver of honey bee mortality [9–14]. It occurs across all host life stages and castes, with pupae being especially permissive to viral replication [9,14–16]. Experimental infections have shown that the injection of fewer than 100 viral particles into pupae or adults can induce tremors, paralysis, and death within days (Bailey et al., 1963). ABPV has been detected in bumble bees [17–20] and ants [21–27] and is considered a multi-host virus, with cross-species transmission facilitated by shared floral resources and nesting sites [5].

In recent years, ants have been proposed as viral reservoirs for ABPV based on repeated field detections [21–26,28]. With an estimated 20 quadrillion individuals and a biomass exceeding that of all wild birds and mammals, ants are globally dominant insects [29,30] whose foraging and nesting ranges overlap with those of many taxa, making them potential pathogen vectors [31,32]. ABPV has been reported in ants and honey bees from ant-infested hives, suggesting ants as viral vectors within apiaries [28]. Yet, most studies remain observational, and whether ants can sustain viral replication, transmit infections within colonies, or mount effective immune responses is largely unknown. Experimental work is limited to one study, which showed that oral exposure to ABPV impaired locomotion and colony growth in *Lasius niger* [33].

Here, we investigate ABPV infections in the clonal raider ant *Ooceraea biroi*, a powerful model for studying disease dynamics under controlled laboratory conditions. Workers in *O. biroi* colonies reproduce clonally and synchronously [34], allowing precise experimental control over genotype and age at the individual and colony levels. In addition, the predatory lifestyle of *O. biroi*, whose diet includes ants and other insects [35], makes them prone to pathogen spillovers.

To investigate the consequences of ABPV exposure in this system, we measured the effects of controlled inoculations on host survival, assessed the ability of the virus to replicate within hosts and transmit within colonies, and measured host immune responses by quantifying antiviral immune gene expression [36–39].

2. Materials & Methods

(a) Ant rearing

All ants were sourced from stock colonies of genetic lineage B [40], and maintained at 28 °C in airtight plastic containers (8 x 12 x 5 cm) with a humidified plaster of Paris floor (Brand). The nests were regularly cleaned, re-humidified, and fed with frozen ant brood (*Tetramorium bicarinatum* and *Oecophylla* sp.), as well as frozen houseflies. For each experiment, all ants were clonally related.

(b) Infection assay in pupae and adults

    (i)   Preparation of viral inoculum

The ABPV inoculum was prepared as described by Tehel et al. (2019). Briefly, 1 µL of homogenate from a heavily ABPV-infected *A. mellifera* worker was injected into white-eyed pupae obtained from virus-free colonies. The injected pupae were incubated for three to five days to allow viral replication and systemic infection, after which they were homogenised in 0.5 M cold potassium phosphate buffer (PPB, pH 8.0). The resulting homogenates were screened using RT-qPCR to confirm the absence of other common bee viruses, such as deformed wing virus (DWV), black queen cell virus (BQCV), chronic bee paralysis virus (CBPV), Israeli acute paralysis virus (IAPV), sacbrood virus (SBV), and slow bee paralysis virus (SBPV). Only homogenates confirmed to contain ABPV exclusively were used in the experiments. Control inoculum was prepared from uninjected white-eyed pupae processed in the same way.

    (ii)   Inoculation via microinjection

10-day old ant pupae were placed on glass slides and injected with viral inoculum (ABPV at 10^5 genome equivalents, GEs) laterally at the posterior abdomen using a FemtoJet microinjector (Pi = 200, Pc = 30; Eppendorf AG, Hamburg, Germany) and borosilicate glass capillaries (8.89 cm length, 0.530 mm ± 25 µm inner diameter, 1.14 mm outer diameter; World Precision Instruments). Inoculum was injected until the pupal body visibly stretched. Control pupae were injected with the control inoculum following the same procedure. Following inoculation, eight pupae of either treatment were placed together with six paint-marked, uninjected cohabiting adults in plastic cups (diameter 4.5 cm) with a humidified plaster of Paris floor. For each treatment (control and ABPV-injected) and each sampling time point (0, 3, and 5 days post-inoculation, d.p.i.), four replicate colonies were established. At time point 0, pupae were flash-frozen immediately after injection. At later time points (3 and 5 d.p.i.). Both pupae and cohabiting adults were frozen to assess viral transmission. Pupae survival and development were assessed in the colonies sampled 5 d.p.i. (four control and four infected colonies). Pupae that disappeared, turned dark, or were partly cannibalized were classified as dead.

Adult ants (mixed age) were immobilised by placing them in slits cut in pencil erasers (Läufer Plast 0120) and injected with 10 nL of viral inoculum (ABPV 5×10^4 GEs) or control inoculum at a rate of 5 nL/s under the largest abdominal tergite using a NanoFil 36 GA bevelled needle connected to a NanoFil syringe and an electronic microinjection pump (UMP3 pump with Micro2T SMARTouch controller, World Precision Instruments, USA). Ten injected adults from either treatment were housed with five paint-marked, uninjected cohabiting adults in plastic cups (diameter 4.5 cm) with a humidified plaster of Paris floor. For each treatment (control and ABPV-injected) and each sampling time point (0, 1, 3, and 5 d.p.i.), four replicate colonies were established. At time point 0, focal ants were flash-frozen immediately after the injection. At later time points (1, 3, and 5 d.p.i.), both focal ants and cohabiting adults were flash-frozen. The colonies sampled 5 d.p.i. were monitored for survival over the full period before sampling.

(c) Viral load quantification

126 For each colony, five injected adults or three injected pupae were flash-frozen in liquid nitrogen
127 and stored at -80 °C. In the ABPV-injected groups, five cohabiting adults were also sampled.
128 Total RNA was extracted from pooled, whole-body samples using a combined Trizol/phenol-
129 chloroform-column protocol [41] and eluted in 20 µL of nuclease-free water. RNA
130 concentration and purity were assessed using a BioPhotometer D30 (Eppendorf, Germany).
131
Viral load quantification was performed according to a protocol modified from Tehel et al.
(2019). For cDNA synthesis, 200 ng of total RNA was reverse-transcribed using oligo(dT)18
primers (Thermo Scientific) and reverse transcriptase (M-MLV and Revertase, Promega,
Mannheim, Germany) according to the manufacturer's instructions in a Veriti Thermal Cycler
(Applied Biosystems). The resulting cDNA was diluted 1:3 and stored at -20 °C. qPCR was
performed using the KAPA SYBR FAST Universal kit (Kapa Biosystems, Sigma-Aldrich) on a
LightCycler 480 (Roche), with two technical replicates per sample, and the following thermal
cycle: three min at 95 °C; 40 cycles of 10 s at 95 °C, 30 s at 57 °C, 20 s at 72 °C; final
elongation for 1 min at 95 °C; melt curve analysis was performed from 55 °C to 95 °C with 0.5
°C per second increments to check for amplification specificity. Forward primer (5′-
CATGGCTCAAGACACTTCATCG-3′) and reverse primer (5′-CCAGCAATGACCTCAATGTG
G-3′) were used to amplify a 105 bp fragment specific to ABPV capsid protein gene, were
designed with Primer-BLAST [42]. Each plate included negative controls (control inoculum
and no-template control) and a positive control (viral inoculum). Samples with Ct values >35
or with high variation between technical duplicates (Ct standard deviation >0.5) were re-
assayed, except for negative controls and control inoculum, with actin (host reference gene)
amplified using primers Ob-actinF (5′-GCCGATCGTATGCAGAAGGA-3′) and Ob-actinR (5′-
ACCCTTGGCGATGTGTCTTT-3′) designed with Primer-BLAST. To quantify the number of viral
GEs in each sample, external DNA standards were generated from the viral inoculum using
primers ABPV-F6531 (5′-TCATACCTGCCGATCAAG-3′) (Locke et al., 2012) and ABPV-
R6774 (5′- CCAGCAATGACCTCAATGTGG-3′). The PCR product size (244 bp) was verified
by electrophoresis on a 2% agarose gel, and the corresponding band was excised and purified
using the NucleoSpin Gel and PCR Purification Kit (Macherey-Nagel, Düren, Germany)
according to the manufacturer's instructions. The purified PCR product was quantified using
a spectrophotometre and used to generate ten-fold serial dilutions. Viral loads were expressed
as $\log_{10}$ GEs per individual ant by dividing total GEs by the number of individuals in each
sample.
159

(d) Negative-sense strand detection
In a separate experiment, three-week-old adult ants were injected with 10 nL of viral inoculum
(ABPV 10^4 GEs), control inoculum, or left uninjected; five focal adults from each treatment
were housed with five paint-marked, uninjected cohabiting adults, as described above. A lower
dose was chosen to increase survival and extend the observation period. For each treatment
and time point (0, 6, 12, 24, and 48 hours post-injection, h.p.i.), four replicate colonies were
established. At each time point, focal and cohabiting ants from each replicate were pooled
separately, flash-frozen in liquid nitrogen, and stored at -80 °C. RNA was extracted as
described above, and the extracted RNA was divided, with a portion used for the negative-
strand assay and the remaining portion used to quantify immune gene expression (see below).
170

Focal and cohabiting ants sampled 24 h.p.i were used to assess viral replication via negative-
strand detection. This method relies on RT-qPCR amplification of negative-strand RNA
intermediates, which serve as markers of active replication of positive-strand RNA viruses
within host cells, rather than the presence of inert genomic RNA [9,43]. Two-step RT-PCR
was performed using a tagged forward primer designed to specifically anneal to the
complementary DNA (cDNA) product of the negative-sense RNA strand of ABPV. Superscript
III reverse transcriptase (Invitrogen, USA) was used for cDNA synthesis. In preliminary tests,
all ant and bee samples produced non-specific PCR products using previously published
primers (Schläppi et al., 2020). To avoid false positives, we designed new primers for the

assay. A total of 200 ng RNA was diluted to 10 µL with water and combined with 1 µL of the tagged forward primer tag-ABPV(-)F5185 (5'-AGCCTGCGCACCGTGG-TTGGTTTGGTGCA GAAGGTG-3') and 1 µL 10 mM dNTPs. The mixture was incubated at 65 °C for 5 min and then placed on ice for at least 1 min. Subsequently, 4 µL of 5X First-Strand Buffer, 1 µL of 0.1 M DTT, and 1 µL of Superscript III Reverse Transcriptase were added, bringing the total reaction volume to 20 µL. The reaction was incubated at 55 °C for 50 min, followed by incubation at 70 °C for 15 min to inactivate the enzyme. cDNA was purified using the NucleoSpin Gel and PCR Clean-up Kit (Macherey-Nagel, Düren, Germany) and eluted in 20 µL of elution buffer. The purified cDNA (diluted 1:3 with nuclease-free water) served as the template for PCR amplification with Taq polymerase (New England Biolabs), using a tag oligonucleotide (5'-AGCCTGCGCACCGTGG-3') as the forward primer and the corresponding reverse primer, ABPV(-)R5380 (5'-AGAAAAGTCCATAGGCCCGT-3'). PCR reactions without the tag primer were performed for all samples to verify the complete removal of residual tagged forward primers using the following cycling conditions: initial denaturation at 95 °C for 3 min; 35 cycles of denaturation at 95 °C for 20 s, annealing at 56 °C for 20 s, and extension at 68 °C for 30 s; and final extension at 68 °C for 2 min. The resulting PCR products were analysed by electrophoresis on a 2% agarose gel. A 196 bp PCR product was expected in samples containing negative-sense RNA, indicating active viral replication. Bee pupae injected with control inoculum or ABPV inoculum served as negative and positive controls, respectively.

(e) Immune gene expression

RNA extracted from adults injected with 10^4 GEs ABPV, which was also used for the negative-strand assay, was used to quantify the expression of the immune gene *Dcr-2* in ABPV-injected ants, control-injected ants, uninjected ants, and cohabiting ants. qPCR were performed as for viral load quantification, with different primers. Primers specific for *Dcr-2* were designed using Primer-BLAST: Dcr-2F (5'-CTCGGTGGATCGGCGAAATA-3') and Dcr-2R (5'-TGAGGTTCACGATCATCGCC-3'), and were used for qPCR analysis. Relative expression levels were normalised to the housekeeping gene ribosomal protein L13a (RPL-13a), amplified using primers RPL-13aF (5'-GCAAACAAAAGCGTGTCAAG-3') and RPL-13aR (5'-TCGGCCAGAGTAAAACCTCT-3') [44].

(f) Statistical analyses

All data analyses were performed using R version 4.4.1 (R Core Team, 2024). Models were selected based on Akaike's Information Criterion (AIC) [45] and compared using anova() function to test if added complexity significantly improved model fit. The final model assumptions and residual diagnostics were assessed using the *DHARMa* package [46]. Post-hoc pairwise comparisons were performed with the *emmeans* package [47] with Tukey's adjustment for multiple comparisons.

Survival

Due to complete separation in the pupal development data (no deaths in the control group and no eclosion in the ABPV-inoculated group), Wilcoxon-Mann-Whitney tests were used to compare the proportion of pupae that hatched into adults or died between treatment groups. For adult survival, the data violated the proportional hazard assumption, and endpoint mortality was therefore analysed using a generalised linear model (GLM) with a binomial error structure and logit link, with Treatment as a fixed factor and Colony as a random factor.

Viral load

The viral load in focal and cohabiting ants was quantified by converting the Ct values to GEs using a standard curve, followed by a $\log_{10}$ transformation. These transformed titres were analysed using a linear model (*lm* function, base R) with one focal and one cohabiting sample per colony. To assess how viral load changed over time in focal individuals, models included Treatment (control-injected vs. ABPV-injected) and Timepoint (0, 3, and 5 d.p.i. for pupae; 0, 1, 3, and 5 d.p.i. for adults), and their interaction (Treatment × Timepoint) as fixed

effects. To examine the viral load differences between focal and cohabiting individuals, a
separate model was fitted for overlapping timepoints between the two groups
including Individual type (focal vs. cohabiting) as a fixed factor, along with Treatment and
Timepoint. For pupae infection assay, comparisons between focal and cohabiting individuals
were performed only within the virus-inoculated treatment group as cohabiting adults were
only sampled in colonies of virus-inoculated pupae; thus, only Individual type and Timepoint
were included as fixed effects.
241
<u>Immune gene expression</u>
For visualisation, *Dcr-2* expression was normalised to the reference gene (*RPL-13a*), and fold
change ($2^{-\Delta\Delta Cq}$) relative to uninjected controls was calculated for injected focal ants and
cohabiting ants from control and ABPV-injected groups. The effects of Treatment (control-
injected vs. ABPV-injected), Time point (0, 12, 24 and 48 h.p.i.) were analyzed separately for
focal and cohabiting individuals using linear models (*lm* function, base R) on $\log_2$-
transformed $-\Delta\Delta Cq$ values as each sample was a pooled ants from the single colony.
249

3. Results

ABPV-injected ant pupae had drastically reduced eclosion rates compared to control-injected pupae 5 d.p.i. (mean ± SD: 0 ± 0% in ABPV, 75 ± 10.2% in controls; Wilcoxon–Mann–Whitney test: z = 2.477, p = 0.0286; Figure 1A). Similarly, ABPV-injected pupae had higher mortality compared to controls (mean ± SD: 31.3 ± 16.1% mortality in ABPV, 0 ± 0% mortality in controls; Wilcoxon–Mann–Whitney test; z = -2.461, p-value = 0.0286).

ABPV-injected adult ants had higher mortality than control-injected adults (mean ± SD: 47.5 ± 31% in ABPV; 15 ± 12.9% in controls; $\chi^2(1)$ = 4.628, p = 0.031), a 5.1-fold increase in the odds of death, mostly occurring 1 d.p.i (Figure 1B). Although virus-injected focal pupae and adults showed increased mortality, no deaths were observed among the cohabiting adults.

Figure 1. ABPV affects *O. biroi* development and survival. (A) Development and survival of ABPV-injected (10^5 GEs, n=4 colonies) and control-injected pupae (n=4 colonies), shown as the mean proportion of pupae (yellow) that eclosed into adults (orange) or died (gray) five days post-injection. **(B)** Survival of ABPV-injected (5×10^4 GEs, red, n=40 individuals) and control-injected (blue) adult ants (n=40 individuals). Kaplan–Meier curves with 95% confidence intervals.

Viral load differed between ABPV-injected and control-injected ant pupae (ANOVA: $F_{1,18}$ = 51.802, p < 0.001) and across time points ($F_{2,18}$ = 5.207, p = 0.016) (Figure 2A). As expected, ABPV-injected pupae had higher viral loads than control-injected pupae at all time points (Tukey's post-hoc; Control vs. ABPV at 0 d.p.i.: t = -7.197, p < 0.0001; 3 d.p.i.: t = -5.726, p <.0001; 5 d.p.i: t = -3.500, p = 0.003). In ABPV-injected pupae, viral load remained consistently high up to 5 d.p.i. (Tukey's post-hoc; 0 vs. 5 d.p.i.: t = 1.2237, p = 0.447).

In adult ants, viral differed between treatments (ANOVA: $F_{1,16}$ = 140.252, p < 0.001) the difference between the control and ABPV-injected groups was not constant across all time points (ANOVA; treatment × timepoint interaction: $F_{3,16}$ = 3.87, p = 0.030, Figure 2B). This effect was driven by a sharp decline in viral load in ABPV-injected adults within the first day post-injection (Tukey's post-hoc; 0 vs. 1 d.p.i.: t = 4.306, p = 0.003), decreasing from approximately $10^{3.08}$ to $10^{1.42}$ GEs per individual. Thereafter, viral titres remained at a similar level up to day 5 (Tukey's post-hoc; 1 vs. 5 d.p.i.: t = 0.795, p = 0.856).

We consistently detected ABPV in uninjected adults that had cohabited with ABPV-injected pupae (Figure 2A) and adults (Figure 2B) for three to five days. Strikingly, adults cohabiting

with ABPV-injected pupae carried viral loads as high as those of the injected pupae
themselves (ANOVA; focal pupae vs. cohabiting adults, $F_{1,\,13}$= 1.922, p = 0.189), suggesting
efficient transmission from pupae to adults. In contrast, adults cohabiting with ABPV-injected
adults had viral loads lower than those in the injected adults themselves (Tukey's post-hoc;
cohabiting vs. focal: t = -5.381, p <0.0001), yet they still had significantly higher viral loads
than controls (Tukey's post-hoc; control-cohabiting vs. ABPV-cohabiting: t = -6.615, p <
0.0001). Thus, this data indicates rapid and efficient horizontal transmission of the virus, both
from the brood to adults and between adults.
295

296
Figure 2. ABPV transmits within ant colonies. (A) Viral load in control-injected pupae (blue
triangles), ABPV-injected pupae (red triangles), and cohabiting adults (red circles). Data points
represent pools of three pupae or five cohabiting adults. Grey: mean ± SD. **(B)** Viral load in
control-injected adults (blue triangles), ABPV-injected adults (red triangles), and cohabiting
adults (blue and red circles). Data points represent pools of five adults (n = 3-4 per treatment
and time point). d.p.i.: days post-injection. Grey: mean ± SD.
303
304   In line with the lack of increase in viral load over time, we found no evidence of viral replication
305   in adult ants. Viral negative-strand replication intermediates were detected in positive controls
306   (ABPV-injected honey bee pupae), but not in ABPV-injected, control-injected, and cohabiting
307   adult ants 24 h post-injection (Figure 3).
308

309
Figure 3. ABPV does not replicate in adult ants. Top row: ABPV-injected and control-
injected bee pupae served as positive and negative controls, respectively. Red: ABPV-injected
focals and cohabiting ants; blue: control-injected focals and cohabiting ants; white: uninjected.
The expected PCR product size was 196 bp. L: DNA ladder. Bottom row: cDNA purification
controls corresponding to the samples in the top row.

We investigated host immune responses by measuring the expression of the antiviral gene
Dcr-2 in response to ABPV injection. In focal ants, *Dcr-2* expression changed with time
(ANOVA; time point: $F_{3,35}$= 4.759, p = 0.007; Figure 4A) regardless of the treatment (Treatment:
F1,35 = 0.030, p= 0.864). Similarly, in cohabiting adults, *Dcr-2* gene expression varied with
time (ANOVA; time point: $F_{2,27}$= 23.134, p <0.001) but not with treatment (Figure 4B; ANOVA;
Treatment: F1,27 = 2.7126, p= 0.111). Thus, *Dcr-2* did not respond to ABPV inoculation under
the conditions tested here.

Figure 4. The expression of *Dcr-2* is not affected by ABPV injection. *Dcr-2* expression
over time (hours post-injection, h.p.i.) in (A) injected and (B) cohabiting ants. Expression levels
are normalised to the reference gene (RPL-13a) and presented as fold changes relative to
uninjected ants. Data points represent pools of five adults; orange: ABPV-injected and
cohabiting adults; blue: control-injected and cohabiting adults. Large circles and error bars
represent the mean ± SEM. The horizontal line at 1 indicates the expression level of *Dcr-2* in
uninjected ants.

4. Discussion

Our study shows that injection of ABPV into *O. biroi* pupae and adults reduced host fitness by delaying development and decreasing survival. ABPV persisted over days in inoculated adults and pupae and was efficiently transmitted to adults, likely through social interactions or the nest environment. Strikingly, uninoculated nestmates acquired ABPV titres comparable to those of directly inoculated ants. These results demonstrate that ABPV particles can persist in ants, spread among nestmates, and impact host fitness, even in the absence of detectable viral replication or canonical antiviral RNAi response, as indicated by stable *Dcr-2* expression.

The higher mortality observed in *O. biroi* injected with ABPV is in line with previous studies demonstrating that bee viruses such as ABPV, can impair fitness in non-honey bee hosts, with outcomes depending on the inoculation methods and dosage. For example, black garden ants (*Lasius niger*) fed on ABPV-infected bee pupae for ten weeks showed impaired locomotion and reduced adult emergence [33]. Similarly, IAPV, a close relative of ABPV [9], reduces host reproductive success and survival when fed to bumblebees at very high doses (2×10^7 and 1×10^8 particles), [38,48,49], while relatively low-dose injections of IAPV (ca. 10^2 particles) were sufficient to cause acute infection, paralysis, and mortality in bumblebee workers [50,51].

Despite host fitness impairment, we found no evidence of viral replication, as shown by stable or decreasing viral loads over time and the lack of detectable replicative viral strands. This suggests that fitness costs arise from host immunopathology rather than active infection [52]. One plausible mechanism is immune-mediated self-damage: ABPV particles may trigger antiviral defences such as apoptosis and subsequent phagocytosis, but excessive apoptosis near vital organs can harm the host [53,54]. Similarly, activation of the phenoloxidase cascade and the release of cytotoxic intermediates or reactive oxidative species confer broad antimicrobial activity, but can cause host tissue damage and early senescence [55–59].

We speculate that the lack of ABPV replication in *O. biroi* is due to host specificity limiting viral infectivity and pathogenicity in a novel host [20,60]. Host phylogeny strongly predicts infection outcomes, as shown for *Drosophila* C virus and Cricket paralysis virus, where changes in viral load correlate with relatedness between natural and novel hosts [61,62]. Successful infection typically requires compatible host cell surface receptors for viral attachment and entry (Maginnis et al., 2018; Weaver et al., 2021) and ABPV may lack the receptor-binding domains required to interact with *O. biroi* cells. Additionally, viral host shifts are often constrained by host-specific physiological and immune barriers, requiring substantially higher inoculation doses to overcome immunity and initiate replication in a novel host [19,33,63,64], a threshold potentially not reached in our experiments. Taken together with cross-species infection experiments in ants [33,65] and bumblebees [19,49,63], our results indicate that although honey bee viruses can spill over into novel hosts and impair fitness, onwards interspecific transmission or spillback to honey bees is not guaranteed and greatly depends on exposure dosage and host-virus compatibility [27,28].

Consistent with the absence of replication, *Dcr-2* expression was not induced by ABPV injection, indicating no activation of the RNAi antiviral pathway. Both ABPV- and control-injected ants showed similar temporal shifts in *Dcr-2* expression, likely reflecting a general response to injection-induced wounding rather than virus-specific immunity. Alternatively, other immune pathways might be involved, as insect antiviral responses can vary across species [66,67].

Adult ants cohabiting with ABPV-injected pupae or adults acquired detectable viral loads within one to three days. Transmission likely occurred through social contact, such as mutual grooming and brood care, or via environmental contamination of the shared nest space [63,68]. Infected carcasses may also have contributed to viral uptake by cohabiting individuals, as honey bee viruses can remain viable and infectious after host death[69]. Other possible uptake

387 routes include the consumption of pupal fluid [70] or brood cannibalism, which we observed
388 in our experiment and is used by other ants to reduce infection risk [71]. The unexpectedly
389 high viral loads in cohabiting adults may therefore reflect repeated recontamination or
390 incomplete immune clearance, similar to the observation that direct contact and cannibalism
391 facilitate efficient intraspecific virus transmission [72].
392
Our work adds to evidence (Valizadeh et al., 2025) showing that ants can acquire and carry
bee viruses without supporting active replication. Their role as incidental hosts or mechanical
reservoirs may facilitate viral maintenance and transmission across insect communities [73].
Positive-sense RNA viruses, such as ABPV, replicate rapidly and with high error rates, which
can accelerate their adaptation to novel hosts [9]. Consequently, even transient uptake by
ants may promote low-level viral persistence that, under selective pressure, may give rise to
variants capable of overcoming interspecific barriers and initiating host shifts, a process
recognised as a key driver of emerging infectious diseases [60]. Environmental stressors, such
as pesticide exposure [74,75] and climate change [76], which affect host immune function and
physiology, can potentially amplify the risk of adaptation to novel hosts and cross-species
transmission of bee viruses.
404
Our findings highlight the potential of honey bee viruses to impose fitness costs and transmit
in non-Apis hosts, such as ants, even without active replication. Such hosts can potentially act
as reservoirs of infection and contribute to future spillover or spillback events, thereby shaping
the epidemiology of bee pathogens in insect communities. Understanding these dynamics is
critical for predicting viral emergence and designing strategies to safeguard pollinator health
and enhance ecosystem resilience.
411

Author contributions:
L.K.L.: conceptualisation, experimental design, data collection and analysis, manuscript writing, manuscript reviewing, and editing; A.B.: experimental design, data collection and analysis, manuscript reviewing; R.J.P.: conceptualisation, manuscript reviewing and editing, funding; Y.U.: conceptualisation, manuscript writing, reviewing and editing, funding.

Conflict of interest declaration:
The authors declare no competing interests.

Acknowledgments:
We thank Paxton lab members Anja Tehel, Tabea Streicher, Patrycja Pluta, and Anja Manigk for supplying the viral inoculum and providing training, as well as members of the Lise Meitner Research Group for Social Behaviour for their valuable input on the experimental design and statistics.

Funding:
This research was funded by the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig and the German Research Foundation (DFG) under the Flexpool programme (Grant number: W47XEF004).

References

1. Neumann P, Carreck NL. 2010 Honey bee colony losses. *J. Apic. Res.* **49**, 1–6. (doi:10.3896/ibra.1.49.1.01)

2. Carreck NL, Ball BV, Martin SJ. 2010 Honey bee colony collapse and changes in viral prevalence associated with Varroa destructor. *J. Apic. Res.* **49**, 93–94. (doi:10.3896/ibra.1.49.1.13)

3. Genersch E, Aubert M. 2010 Emerging and re-emerging viruses of the honey bee (Apis mellifera L.). *Vet. Res.* **41**, 54. (doi:10.1051/vetres/2010027)

4. McMenamin AJ, Genersch E. 2015 Honey bee colony losses and associated viruses. *Curr. Opin. Insect Sci.* **8**, 121–129. (doi:10.1016/j.cois.2015.01.015)

5. Yañez O *et al.* 2020 Bee viruses: Routes of infection in Hymenoptera. *Front. Microbiol.* **11**, 943. (doi:10.3389/fmicb.2020.00943)

6. Wiegers FP. 2021 Transmission of honeybee viruses by Varroa jacobsoni Oud. In *European Research on Varroatosis Control*, pp. 99–104. CRC Press. (doi:10.1201/9781003211464-25)

7. Ryabov EV, Wood GR, Fannon JM, Moore JD, Bull JC, Chandler D, Mead A, Burroughs N, Evans DJ. 2014 A virulent strain of deformed wing virus (DWV) of honeybees (Apis mellifera) prevails after Varroa destructor-mediated, or in vitro, transmission. *PLoS Pathog.* **10**, e1004230. (doi:10.1371/journal.ppat.1004230)

8. Benjeddou M, Leat N, Allsopp M, Davison S. 2001 Detection of acute bee paralysis virus and black queen cell virus from honeybees by reverse transcriptase pcr. *Appl. Environ. Microbiol.* **67**, 2384–2387. (doi:10.1128/AEM.67.5.2384-2387.2001)

9. de Miranda JR, Cordoni G, Budge G. 2010 The Acute bee paralysis virus-Kashmir bee virus-Israeli acute paralysis virus complex. *J. Invertebr. Pathol.* **103 Suppl 1**, S30-47. (doi:10.1016/j.jip.2009.06.014)

10. Ball BV. 1985 Acute paralysis virus isolates from honeybee colonies infested withVarroa jacobsoni. *J. Apic. Res.* **24**, 115–119. (doi:10.1080/00218839.1985.11100658)

11. Ball BV, Allen MF. 1988 The prevalence of pathogens in honey bee (*Apis mellifera*) colonies infested with the parasitic mite *Varroa jacobsoni. Ann. Appl. Biol.* **113**, 237–244. (doi:10.1111/j.1744-7348.1988.tb03300.x)

12. Cox-Foster DL *et al.* 2007 A metagenomic survey of microbes in honey bee colony collapse disorder. *Science* **318**, 283–287. (doi:10.1126/science.1146498)

13. Berthoud H, Imdorf A, Haueter M, Radloff S, Neumann P. 2010 Virus infections and winter losses of honey bee colonies (Apis mellifera). *J. Apic. Res.* **49**, 60–65. (doi:10.3896/ibra.1.49.1.08)

14. Žvokelj L, Bakonyi T, Korošec T, Gregorc A. 2020 Appearance of acute bee paralysis virus, black queen cell virus and deformed wing virus in Carnolian honey bee (Apis mellifera carnica) queen rearing. *J. Apic. Res.* **59**, 53–58. (doi:10.1080/00218839.2019.1681115)

15. Bailey L, Milne RG. 1969 The multiplication regions and interaction of acute and chronic bee-paralysis viruses in adult honey bees. *J. Gen. Virol.* **4**, 9–14. (doi:10.1099/0022-1317-4-1-9)

16. Azzami K, Ritter W, Tautz J, Beier H. 2012 Infection of honey bees with acute bee paralysis virus does not trigger humoral or cellular immune responses. *Arch. Virol.* **157**, 689–702. (doi:10.1007/s00705-012-1223-0)

17. McMahon DP, Fürst MA, Caspar J, Theodorou P, Brown MJF, Paxton RJ. 2015 A sting in the spit: widespread cross-infection of multiple RNA viruses across wild and managed bees. *J. Anim. Ecol.* **84**, 615–624. (doi:10.1111/1365-2656.12345)

18. Pascall DJ, Tinsley MC, Clark BL, Obbard DJ, Wilfert L. 2021 Virus prevalence and genetic diversity across a wild bumblebee community. *Front. Microbiol.* **12**, 650747. (doi:10.3389/fmicb.2021.650747)

19. Gusachenko O, Woodford L, Balbirnie-Cumming K, Ryabov E, Evans DJ. 2020 Evidence for and against deformed wing virus spillover from honey bees to bumble bees: a reverse genetic analysis. *Sci. Rep.* **10**. (doi:10.1038/s41598-020-73809-3)

20. Doublet V, Doyle TD, Carvell C, Brown MJF, Wilfert L. 2025 Host ecology and phylogeny shape the temporal dynamics of social bee viromes. *Nat. Commun.* **16**, 2207. (doi:10.1038/s41467-025-57314-7)

21.   Celle O, Blanchard P, Olivier V, Schurr F, Cougoule N, Faucon J-P, Ribière M. 2008 Detection of Chronic
bee paralysis virus (CBPV) genome and its replicative RNA form in various hosts and possible ways of spread.
Virus Res. **133**, 280–284. (doi:10.1016/j.virusres.2007.12.011)

22.   Levitt AL, Singh R, Cox-Foster DL, Rajotte E, Hoover K, Ostiguy N, Holmes EC. 2013 Cross-species
transmission of honey bee viruses in associated arthropods. *Virus Res.* **176**, 232–240.
(doi:10.1016/j.virusres.2013.06.013)

23.   Sébastien A, Lester PJ, Hall RJ, Wang J, Moore NE, Gruber MAM. 2015 Invasive ants carry novel viruses in
their new range and form reservoirs for a honeybee pathogen. *Biol. Lett.* **11**, 20150610.
(doi:10.1098/rsbl.2015.0610)

24.   Gruber MAM, Cooling M, Baty JW, Buckley K, Friedlander A, Quinn O, Russell JFEJ, Sébastien A, Lester PJ.
2017 Single-stranded RNA viruses infecting the invasive Argentine ant, Linepithema humile. *Sci. Rep.* **7**,
3304. (doi:10.1038/s41598-017-03508-z)

25.   Lin C-Y, Lee C-C, Nai Y, Hsu H-W, Lee C, Tsuji K, Yang C. 2020 Deformed wing virus in two widespread
invasive ants: Geographical distribution, prevalence, and phylogeny. *Viruses* **12**. (doi:10.3390/v12111309)

26.   Payne AN, Shepherd TF, Rangel J. 2020 The detection of honey bee (Apis mellifera)-associated viruses in
ants. *Sci. Rep.* **10**. (doi:10.1038/s41598-020-59712-x)

27.   Tiritelli R, Cilia G, Grasso DA. 2024 Occurrence of honey bee (Apis mellifera) pathogens in ants
(Hymenoptera: Formicidae): a systematic review. *Myrmecol. News* **34**, 91–104.
(doi:10.25849/MYRMECOL.NEWS_034:091)

28.   Rivas Fontan I, Gonzalez FN, Moja J, Ferrufino C, Zanola D, Scally B, Calcaterra L, Dus Santos MJ, Josens
R. 2025 Assessing potential viral transmission between honeybees and hive-infesting ants using novel per-
hive co-detection indices. *Apidologie (Celle)* **56**. (doi:10.1007/s13592-025-01190-z)

29.   Schultheiss P, Nooten SS, Wang R, Wong MKL, Brassard F, Guénard B. 2022 The abundance, biomass,
and distribution of ants on Earth. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2201550119.
(doi:10.1073/pnas.2201550119)

30.   Parker J, Kronauer DJC. 2021 How ants shape biodiversity. *Curr. Biol.* **31**, R1208–R1214.
(doi:10.1016/j.cub.2021.08.015)

31.   Tiritelli R, Giannetti D, Schifani E, Grasso DA, Cilia G. 2025 Neighbors sharing pathogens: the intricate
relationship between Apis mellifera and ants (Hymenoptera: Formicidae) nesting in hives. *Insect Sci.* **32**, 943–
956. (doi:10.1111/1744-7917.13433)

32.   Dobelmann J, Felden A, Lester PJ. 2023 An invasive ant increases deformed wing virus loads in honey bees.
Biol. Lett. **19**, 20220416. (doi:10.1098/rsbl.2022.0416)

33.   Schläppi D, Chejanovsky N, Yañez O, Neumann P. 2020 Foodborne transmission and clinical symptoms of
honey bee viruses in ants Lasius spp. *Viruses* **12**. (doi:10.3390/v12030321)

34.   Ravary F, Jahyny B, Jaisson P. 2006 Brood stimulation controls the phasic reproductive cycle of the
parthenogenetic ant Cerapachys biroi. *Insectes Soc.* **53**, 20–26. (doi:10.1007/s00040-005-0828-7)

35.   Kronauer DJ. 2009 Recent advances in army ant biology (Hymenoptera: Formicidae). *Myrmecological News*
12, 51–65.

36.   Deddouche S *et al.* 2008 The DExD/H-box helicase Dicer-2 mediates the induction of antiviral activity in
drosophila. *Nat. Immunol.* **9**, 1425–1432. (doi:10.1038/ni.1664)

37.   Niu J, Meeus I, Cappelle K, Piot N, Smagghe G. 2014 The immune response of the small interfering RNA
pathway in the defense against bee viruses. *Curr. Opin. Insect Sci.* **6**, 22–27. (doi:10.1016/j.cois.2014.09.014)

38.   Piot N, Snoeck S, Vanlede M, Smagghe G, Meeus I. 2015 The effect of oral administration of dsRNA on viral
replication and mortality in Bombus terrestris. *Viruses* **7**, 3172–3185. (doi:10.3390/v7062765)

39.   Poirier EZ *et al.* 2018 Dicer-2-dependent generation of viral DNA from defective genomes of RNA viruses
modulates antiviral immunity in insects. *Cell Host Microbe* **23**, 353-365.e8. (doi:10.1016/j.chom.2018.02.001)

40.   Kronauer DJC, Pierce NE, Keller L. 2012 Asexual reproduction in introduced and native populations of the
523   ant Cerapachys biroi. *Mol. Ecol.* **21**, 5221–5235. (doi:10.1111/mec.12041)

524   41.   Libbrecht R, Oxley PR, Keller L, Kronauer DJC. 2016 Robust DNA methylation in the clonal raider ant brain.
525   *Curr. Biol.* **26**, 391–395. (doi:10.1016/j.cub.2015.12.040)

526   42.   Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. 2012 Primer-BLAST: a tool to design
527   target-specific primers for polymerase chain reaction. *BMC Bioinformatics* **13**, 134. (doi:10.1186/1471-2105-
528   13-134)

529   43.   Yue C, Genersch E. 2005 RT-PCR analysis of Deformed wing virus in honeybees (Apis mellifera) and mites
530   (Varroa destructor). *J. Gen. Virol.* **86**, 3419–3424. (doi:10.1099/vir.0.81401-0)

531   44.   Oxley PR, Ji L, Fetter-Pruneda I, McKenzie SK, Li C, Hu H, Zhang G, Kronauer DJC. 2014 The genome of
532   the clonal raider ant Cerapachys biroi. *Curr. Biol.* **24**, 451–458. (doi:10.1016/j.cub.2014.01.018)

533   45.   Akaike H. 1976 Canonical correlation analysis of time series and the use of an information criterion.
534   *Mathematics in science and engineering* **126**, 27–96. (doi:10.1016/S0076-5392(08)60869-3)

535   46.   Hartig F. 2020 DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models. *R
536   package version 0.3*

537   47.   Lenth RV, Singmann H, Love J, Buerkner P, Herve M. 2022 Emmeans: estimated marginal means. *AKA
538   least-squares means*

539   48.   Meeus I, de Miranda JR, de Graaf DC, Wäckers F, Smagghe G. 2014 Effect of oral infection with Kashmir
540   bee virus and Israeli acute paralysis virus on bumblebee (Bombus terrestris) reproductive success. *J.
541   Invertebr. Pathol.* **121**, 64–69. (doi:10.1016/j.jip.2014.06.011)

542   49.   Wang H, Meeus I, Piot N, Smagghe G. 2018 Systemic Israeli acute paralysis virus (IAPV) infection in
543   bumblebees (Bombus terrestris) through feeding and injection. *J. Invertebr. Pathol.* **151**, 158–164.
544   (doi:10.1016/j.jip.2017.11.015)

545   50.   Wang H, Meeus I, Smagghe G. 2016 Israeli acute paralysis virus associated paralysis symptoms, viral tissue
546   distribution and Dicer-2 induction in bumblebee workers (Bombus terrestris). *J. Gen. Virol.* **97**, 1981–1989.
547   (doi:10.1099/jgv.0.000516)

548   51.   Niu J, Cappelle K, de Miranda JR, Smagghe G, Meeus I. 2014 Analysis of reference gene stability after Israeli
549   acute paralysis virus infection in bumblebees Bombus terrestris. *J. Invertebr. Pathol.* **115**, 76–79.
550   (doi:10.1016/j.jip.2013.10.011)

551   52.   González-González A, Wayne ML. 2020 Immunopathology and immune homeostasis during viral infection in
552   insects. *Adv. Virus Res.* **107**, 285–314. (doi:10.1016/bs.aivir.2020.06.001)

553   53.   Nainu F, Shiratsuchi A, Nakanishi Y. 2017 Induction of apoptosis and subsequent phagocytosis of virus-
554   infected cells as an antiviral mechanism. *Front. Immunol.* **8**, 1220. (doi:10.3389/fimmu.2017.01220)

555   54.   Clarke TE, Clem RJ. 2003 Insect defenses against virus infection: the role of apoptosis. *Int. Rev. Immunol.*
556   **22**, 401–424. (doi:10.1080/08830180305215)

557   55.   Nappi AJ, Christensen BM. 2005 Melanogenesis and associated cytotoxic reactions: applications to insect
558   innate immunity. *Insect Biochem. Mol. Biol.* **35**, 443–459. (doi:10.1016/j.ibmb.2005.01.014)

559   56.   Sadd BM, Siva-Jothy MT. 2006 Self-harm caused by an insect's innate immunity. *Proc. Biol. Sci.* **273**, 2571–
560   2574. (doi:10.1098/rspb.2006.3574)

561   57.   Wu S-C, Liao C-W, Pan R-L, Juang J-L. 2012 Infection-induced intestinal oxidative stress triggers organ-to-
562   organ immunological communication in Drosophila. *Cell Host Microbe* **11**, 410–417.
563   (doi:10.1016/j.chom.2012.03.004)

564   58.   Mikonranta L, Mappes J, Kaukoniitty M, Freitak D. 2014 Insect immunity: oral exposure to a bacterial
565   pathogen elicits free radical response and protects from a recurring infection. *Front. Zool.* **11**, 23.
566   (doi:10.1186/1742-9994-11-23)

567   59.   Dowling DK, Simmons LW. 2009 Reactive oxygen species as universal constraints in life-history evolution.
568   *Proc. Biol. Sci.* **276**, 1737–1745. (doi:10.1098/rspb.2008.1791)

60. Woolhouse ME, Haydon DT, Antia R. 2005 Emerging pathogens: the epidemiology and evolution of species
jumps. *Trends in ecology & evolution* **20**, 238–244.

61. Longdon B, Hadfield JD, Webster CL, Obbard DJ, Jiggins FM. 2011 Host phylogeny determines viral
persistence and replication in novel hosts. *PLoS Pathog.* **7**, e1002260. (doi:10.1371/journal.ppat.1002260)

62. Imrie RM, Roberts KE, Longdon B. 2021 Between virus correlations in the outcome of infection across host
species: Evidence of virus by host species interactions. *Evol. Lett.* **5**, 472–483. (doi:10.1002/evl3.247)

63. Tehel A, Streicher T, Tragust S, Paxton RJ. 2022 Experimental cross species transmission of a major viral
pathogen in bees is predominantly from honeybees to bumblebees. *Proc. Biol. Sci.* **289**, 20212255.
(doi:10.1098/rspb.2021.2255)

64. Streicher T, Tehel A, Tragust S, Paxton RJ. 2023 Experimental viral spillover can harm *Bombus terrestris*
workers under field conditions. *Ecol. Entomol.* **48**, 81–89. (doi:10.1111/een.13203)

65. Valizadeh B, Hardy J, Chen J, Amiri E. 2025 Dynamics of Israeli acute paralysis virus in the red imported fire
ant (Solenopsis invicta Buren) colonies. *J. Invertebr. Pathol.* **211**, 108310. (doi:10.1016/j.jip.2025.108310)

66. Lester PJ, Buick KH, Baty J, Felden A, Haywood J. 2019 Different bacterial and viral pathogens trigger distinct
immune responses in a globally invasive ant. *Sci. Rep.* **9**. (doi:10.1038/s41598-019-41843-5)

67. Ray AM, Tehel A, Rasgon JL, Paxton RJ, Grozinger CM. 2025 The intensity of the transcriptional response
varies across infection with distinct viral strains in an insect host. *BMC Genomics* **26**, 175.
(doi:10.1186/s12864-025-11365-8)

68. Dalmon A, Diévart V, Thomasson M, Fouque R, Vaissière BE, Guilbaud L, Le Conte Y, Henry M. 2021
Possible spillover of pathogens between bee communities foraging on the same floral resource. *Insects* **12**,
122. (doi:10.3390/insects12020122)

69. Payne AN, Prayugo V, Dolezal AG. 2025 A honey bee-associated virus remains infectious and quantifiable
in postmortem hosts. *J. Invertebr. Pathol.* **209**, 108258. (doi:10.1016/j.jip.2024.108258)

70. Snir O, Alwaseem H, Heissel S, Sharma AK, Valdés-Rodríguez S, Carroll T, Jiang CS, Razzauti J, Kronauer
D. 2022 The pupal moulting fluid has evolved social functions in ants. *Nature* **612**, 488–494.
(doi:10.1038/s41586-022-05480-9)

71. Bizzell F, Pull CD. 2024 Ant queens cannibalise infected brood to contain disease spread and recycle
nutrients. *Curr. Biol.* **34**, R848–R849. (doi:10.1016/j.cub.2024.07.062)

72. Posada-Florez F, Lamas ZS, Hawthorne DJ, Chen Y, Evans JD, Ryabov EV. 2021 Pupal cannibalism by
worker honey bees contributes to the spread of deformed wing virus. *Sci. Rep.* **11**, 8989.
(doi:10.1038/s41598-021-88649-y)

73. Dobelmann J, Manley R, Wilfert L. 2024 Caught in the act: the invasion of a viral vector changes viral
prevalence and titre in native honeybees and bumblebees. *Biol. Lett.* **20**, 20230600.
(doi:10.1098/rsbl.2023.0600)

74. Al Naggar Y, Paxton RJ. 2021 The novel insecticides flupyradifurone and sulfoxaflor do not act synergistically
with viral pathogens in reducing honey bee (Apis mellifera) survival but sulfoxaflor modulates host
immunocompetence. *Microb. Biotechnol.* **14**, 227–240. (doi:10.1111/1751-7915.13673)

75. Schläppi D, Kettler N, Glauser G, Straub L, Yañez O, Neumann P. 2021 Varying impact of neonicotinoid
insecticide and acute bee paralysis virus across castes and colonies of black garden ants, Lasius niger
(Hymenoptera: Formicidae). *Sci. Rep.* **11**, 20500. (doi:10.1038/s41598-021-98406-w)

76. Roberts KE, Hadfield JD, Sharma MD, Longdon B. 2018 Changes in temperature alter the potential outcomes
of virus host shifts. *PLoS Pathog.* **14**, e1007185. (doi:10.1371/journal.ppat.1007185)