OCA-B/Pou2af1 Expression in Mouse T Cells Promotes PD-1 Blockade-Induced Autoimmunity but is Dispensable for Anti-Tumor Immunity

Junhong Du[1,2], Asit K. Manna[1,2], Erik P. Hughes[1,2], Paulla Bisoma[1,2], Kimberley J. Evason[1,2], Arabella Young[1,2], W. David Wilson[3], Abdelbasset A. Farahat[4,5], Dean Tantin[1,2]

[1]Department of Pathology, University of Utah School of Medicine, Salt Lake City, UT, 84112, USA.

[2]Huntsman Cancer Institute, University of Utah School of Medicine, Salt Lake City, UT, 84112, USA.

[3]Department of Chemistry, Georgia State University, Atlanta, GA 30303, USA.

[4]Master of Pharmaceutical Science Program, California North State University, Elk Grove CA 95757, USA.

[5]Department of Pharmaceutical Organic Chemistry, Faculty of Pharmacy, Mansoura University, Mansoura 35516, Egypt

Correspondence to Dean Tantin: dean.tantin@path.utah.edu

Short title: OCA-B Promotes PD-1 Blockade-Triggered Diabetes

Keywords OCA-B/Pou2af1; Type-1 diabetes; Non-obese diabetic mouse; tumor immunity; MC38

Conflicts of interest statement: The authors have declared that no conflict of interest exists

Words in abstract: 225

Words main text: 11,201

Total main display items: 6

Abstract

The transcription coregulator OCA-B promotes CD4[+] T cell memory recall responses and
autoimmunity. OCA-B T cell deletion blocks spontaneous T1D onset in non-obese diabetic (NOD)
mice and blunts T1D in a subset of more aggressive models. However, the role of OCA-B in diabetes
induced by treatment with checkpoint blockade therapies, and the role of OCA-B in the control of
tumors with and without α-PD-1 treatment, has not been studied. We show that deletion of OCA-B in
T cells fully insulates 8-week-old non-obese diabetic (NOD) mice against α-PD-1 antibody-induced
diabetes induction and partially protects 12-week-old mice. Salivary/lacrimal gland infiltration and
inflammation were also reduced. Protection was associated with a block in the differentiation of
progenitor exhausted CD8[+] T cells (T_{PEX}) into terminally exhausted CD8[+] T cells (T_{EX}). We show that
OCA-B T cell loss preserves anti-tumor immune responses following PD-1 blockade therapy in a
variety of tumor cell lines. These findings point to a potential therapeutic window in which
pharmaceuticals that target OCA-B could be used to block the emergence of both spontaneous and
checkpoint blockade-induced autoimmunity while minimally affecting anti-tumor immune responses.
We develop first-in-class small molecule inhibitors of Oct1/OCA-B transcription complexes and show
that administration into NOD mice also blocks diabetes emergence following α-PD-1 treatment. These
results identify OCA-B as a promising therapeutic target for the prevention of autoimmunity and
immune-related adverse events (irAEs).

Introduction

Type 1 diabetes (T1D) affects 1.8 million people in the United States and is increasing worldwide [1-3]. It is characterized by autoimmune destruction of pancreatic β cells and consequent inability to produce insulin [4]. While T1D symptoms can be managed with insulin injections, they must be taken for life. Transplantation of islets of Langerhans can restore insulin-independence to T1D patients, but results in reactivation of islet-specific memory cells T cells, β cell destruction and transplant rejection [5,6]. Similarly, engraftment of islets from non-obese diabetic (NOD) mice with severe combined immunodeficiency (NOD.SCID) into diabetic NODs briefly restores normoglycemia, but results in robust memory autoimmune responses and complete loss of grafted islet function in 10-14 days [7]. Human therapies aim to minimize graft rejection or block the initial emergence of T1D in at-risk patients altogether while sparing normal immune function [8].

In humans, α-PD-1 therapy as well as therapies directed against a number of other inhibitory T cell receptors (immune checkpoint inhibitors or ICIs) are now used for treatment of a variety of cancer indications [9-11]. The development and wide adoption of ICIs has resulted in a concomitant increase in associated unwanted immune-related adverse events, or irAEs, affecting a variety of tissues (e.g., skin, rheumatoid, pulmonary, gut, cardiac), some of which can be permanent [9,12-19]. IrAEs include endocrinopathies such as thyroiditis [20], and less frequently hypophysitis, autoimmune diabetes and other diseases [15,17]. A critical gap in the field is the identification of treatment strategies that block the emergence of irAEs while preserving anti-tumor immunity.

Polymorphisms in the *HLA-DR* and *HLA-DQ* genes encoding MHC class II have the strongest genetic influence on T1D, accounting for 40-50% of familial risk [21]. These findings implicate CD4+ T cells as primary drivers of pathogenesis. CD4+ T cells direct other cell types such as CD8+ T cells and macrophages to infiltrate pancreatic islets and cause β-cell destruction [22]. Consistent with these findings, transient T cell depletion using α-CD3ε antibodies (Teplizumab-mzwv) significantly delays T1D onset in at-risk individuals in human clinical trials [23,24]. Though promising, such treatments are limited by the resulting global immunosuppression. An ideal alternative would selectively deplete pathogenic T cell function while preserving beneficial immune function, including immunity against tumors.

OCA-B is a B cell- and T cell-restricted transcription coactivator [25,26]. In CD4+ T cells, OCA-B docks with the transcription factor Oct1 (gene symbol *Pou2f1*) to regulate targets including *Il2*, *Ifng*, *Icos*, *Csf2* (*Gmcsf*) and *Tbx21* (*Tbet*) [27,28]. Unlike transcription factors such as NFAT, AP-1 and NF-κB, OCA-B controls the expression of these genes specifically during repeated antigen exposures [27,28]. Agents targeting this pathway would selectively inhibit over-active T cell responses to antigen re-exposure but preserve acute immune response. Consistent with this prediction, in the absence of either OCA-B or its cognate transcription factor Oct1/Pou2f1, primary T cell responses to systemic and glial-tropic viruses (LCMV and JHMV, respectively) and viral clearance are normal [26,29], however fewer central memory (T_{CM}) CD4+ T cells form, and those that do respond poorly to antigenic re-challenge [26,28]. Three studies directly link polymorphisms in the human *OCAB* (*POU2AF1*) locus with autoimmune disease [30-32]. Human polymorphisms affecting Oct1 (and therefore OCA-B) binding to the *TNFA* and *CTLA4* loci are associated with inflammatory bowel disease and systemic lupus erythematosus, respectively [33,34]. More recent work identified additional polymorphisms in Oct1 binding sites within the human *IL13*, *IL18* and *HLA-C* loci [35-37]. Additionally, two meta-analyses show a strong association between Oct1/OCA-B binding site polymorphisms and autoimmunity including T1D[38,39]. Collectively, these studies link OCA-B with human autoimmunity.

OCA-B loss in T cells protects mice against spontaneous T1D onset [40]. Single-cell RNA-sequencing
(scRNA-seq) reveals that protection is associated with loss of autoreactive clones but maintenance of
the overall pancreatic islet immune architecture [40]. These findings suggest that targeting OCA-B could
be used to treat T1D early in the disease course while sparing baseline immunity. OCA-B's lymphoid-
restricted expression pattern [25,26], its dispensability for primary antiviral responses [26,28,29], and its
specificity for Oct1/Pou2f1 and Oct2/Pou2f2 complexes with DNA [41-43] make it a promising therapeutic
target that could be used to specifically block autoreactive CD4$^+$ T cells while sparing normal
immunity. OCA-B peptide mimics linked to membrane-penetrating peptides can reduce inflammation
and hyperglycemia in recent-onset T1D NOD mice, but also show significant toxicity [40]. An
outstanding question is whether OCA-B can be successfully targeted by more drug-like small
molecules. It also remains to be determined if OCA-B promotes diabetes driven by ICI therapy, and/or
if OCA-B promotes anti-tumor immunity, including that promoted by ICI.

Here, we show that OCA-B in T cells promotes the emergence of autoimmune diabetes following ICI
administration in NOD mice, while being dispensable for anti-tumor immune responses or tumor
regression following ICI administration. We develop small molecules that effectively disrupt
Oct1:OCA-B complexes in vitro and disrupt OCA-B's transcription functions in cultured cells. We show
that administration of these compounds to NOD mice blocks the emergence of diabetes and other
endocrinopathies following ICI administration, with in vivo efficacy tracking with activity in vitro.
Collectively, the findings indicate that OCA-B promotes diabetes development following ICI treatment,
and that targeting OCA-B could be used to block irAEs following ICI while sparing anti-tumor
immunity.

Results

OCA-B T cell knockout protects NOD mice against diabetes induced by α-PD-1 antibody treatment

T cell conditional OCA-B loss protects NOD mice against spontaneous T1D [40]. Modulating the PD-1

pathway induces diabetes and other endocrinopathies in NOD mice [44-47]. To test the effect of OCA-B T

cell loss in diabetes induced by PD-1 blockade, we treated 8-week-old male NOD mice with four

injections of α-PD-1 antibody. OCA-B loss in T cells completely blocked α-PD-1-induced diabetes

development (Figure 1A-B). Treatment of 12-week-old female NOD mice with α-PD-1 antibodies

resulted in less complete protection, with half the mice failing to develop diabetes and the other half

showing delayed onset (Supp. Figure 1A-B). Flow cytometry revealed a significant reduction in $CD45^+$

leukocytes in α-PD-1 antibody treated knockout mice (Figure 1C-D). We also observed a skewing

from $CD8^+$ to $CD4^+$ T cells in the islets of 8-week-old knockouts (Figure 1E-F), consistent with

changes in immune profiles reflecting diabetes progression [22]. At 12 weeks, islet $PD-1^+TIM-3^+$ $CD8^+$ T

cells were decreased in the OCA-B deficient condition (Supp. Figure 1C-D). PD-1 blockade is known

to normally induce exhausted $CD8^+$ T cell proliferation and differentiation in the pancreas, promoting

the expansion of both T_{PEX} and T_{EX} cells but also skewing the islet total population in favor of T_{EX} [22].

The finding of decreased TIM-3 expression could therefore be explained if OCA-B loss blocks this

pathogenic maturation. To test this, we used SLAMF6 and CD39 to distinguish T_{PEX} and T_{EX} $CD8^+$ T

cells at both 8 and 12 weeks. OCA-B loss resulted in a marked retardation in the ability of $PD-1^+CD8^+$

T_{PEX} cells within the islets of 8-week-old mice to differentiate into T_{EX} (Figure 1G-H). Similar findings

were made at 12 weeks (Supp. Figure 1E-F). Cumulatively, these results indicate that OCA-B loss in

T cells blocks the induction of diabetes by α-PD-1 treatment, and blocks the pathogenic maturation of

a T_{PEX}-like T cell population in the pancreatic islets.

OCA-B T cell knockout maintains anti-tumor response and tumor control by ICI

To determine if OCA-B T cell loss has a corresponding effect on anti-tumor immunity, we used

C57BL/6 and NOD mice OCA-B conditional knockout mice [26,40] with syngeneic tumor lines. In the

C57BL/6 strain background we used two lines, B16F10 and MC38. Compared to littermate controls

lacking the CD4-Cre driver, OCA-B loss had no effect on tumor growth of B16F10 tumors (Supp.

Figure 2). This result was expected because B16 cells are "immune-cold" and do no elicit robust

immune responses unless engineered to express specific antigens or unless multiple checkpoint-

inhibiting antibodies are combined [48]. In contrast, "immune-hot" MC38 colon carcinoma cells exhibit

significant control by T cells that can be augmented by α-PD-1 treatment [49]. MC38 tumor control was

also preserved with OCA-B T cell loss either in the absence or presence of α-PD-1 antibody treatment

(Figure 2A-B). Flow cytometry of cells from the tumor microenvironment (TME) showed equivalent

$CD45.2^+$ leukocyte infiltration and ratios of $CD8^+$ to $CD4^+$ T cells with α-PD-1 treatment (Figure 2C-D).

Similar percentages of $CD4^+$ and $CD8^+$ $SLAMF6^+CD39^-$ and $SLAMF6^-CD39^+$ cells were also observed

in the TME regardless of the status of OCA-B following α-PD-1 antibody treatment (Figure 2E-F). TIM-

3 expression in tumor-infiltrating $CD4^+$ and $CD8^+$ was also equivalent (Figure 2G-H). Despite the lack

of difference in T cell infiltration, exhaustion and tumor control, within the tumors there were

interesting differences in both the $CD4^+$ and $CD8^+$ T_{CM}-like ($CD62L^{hi}CD44^{hi}$) and naïve-like

($CD62L^{hi}CD44^{lo}$) compartments (Supp. Figure 3). The significance of these differences for tumor

control is currently unclear.

Because the protection from PD-1-induced diabetes was observed using NOD background mice, we
also assessed tumor responses using a syngeneic "immune-hot" 9495 sarcoma cell line [50], allowing
us to simultaneously measure diabetes incidence and tumor control in the same animals. α-PD-1
antibody treatment is effective in controlling 9495 tumor growth in syngeneic NOD mice [50]. To
minimize complications from induced diabetes and hyperglycemia, mice were administered insulin
pellets when glucose levels reached ≥250 mg/dL for two consecutive days to minimize complications
from induced diabetes and hyperglycemia. In these tumors, α-PD-1 antibody treatment was effective,
as expected [50]. Again, tumor control either with or without α-PD-1 treatment was equivalent regardless
of OCA-B status (Figure 3A-B). Simultaneously, in another cohort of mice with the same experiment
procedures performed, the OCA-B deficient group were again fully protected against diabetes (Supp.
Figure 4A). Similar to published work [50], control mice with induced diabetes exhibited superior tumor
control compared to uninvolved mice (Supp. Figure 4B-C). As with MC38 tumors in C57BL/6 mice,
CD45$^+$ leukocyte tumor infiltration, CD8$^+$ to CD4$^+$ T cell ratio, CD8$^+$ T cell activation, (Figure 3B-E) and
T_{PEX} to T_{EX} differentiation (Figure 3F-I) were equivalent. These findings indicate minimal differences in
anti-tumor immune responses to multiple syngeneic tumor lines in using mice with OCA-B T cell
knockout, indicating a divergence between anti-tumor immunity and the precipitation of irAEs that
could be leveraged therapeutically.

Development of small molecule inhibitors of Oct1:OCA-B complexes

The Oct1:OCA-B:DNA co-crystal structure [43] suggests multiple possible targeting strategies, including
rationally designed OCA-B peptide mimics (Figure 4A, upper lightning bolt) which are functional but
associated with toxicity in cells and mice due to the membrane penetrating peptide [40]. To identify novel
small molecule inhibitors, we used another strategy in which DNA minor groove-binding compounds
with specificity for the consensus octamer motif (5'ATGCAAAT) would disrupt Oct1:OCA-B
complexes. These inhibitors would dock on the underside of the DNA in the structure shown in Figure
4A (bottom lightning bolt) and would interfere with the unstructured Oct1 linker domain that also spans
the minor groove. We synthesized and tested 9 candidate minor groove-binding compounds with
specificity for the octamer motif. To test efficacy, we developed two cell-based assay systems. The
first is a transcription-based platform in which we stably transfected the human T cell line SupT1 to
express secreted nanoluciferase (nLuc) under the control of five Oct1/OCA-B binding sites. SupT1 T
cells express OCA-B at higher levels than, e.g. Jurkat cells, and at levels comparable to primary
thymocytes (Figure 4B). We generated a stable-transfected SupT1 clone expressing nLuc under the
control of five wild-type octamer sequences ("5×Oct") and a second using mutant sites incapable of
binding Oct1 and OCA-B ("5×Mut", Figure 4C). The 5×Oct clone produced substantial levels of
luciferase activity relative to empty vector-transduced controls, while the 5×Mut clone expressed lower
but still measurable activity (Figure 4D). Timecourse experiments show that 12- to 18-hr incubations
have not yet saturated nLuc activity and capture a dynamic range allowing both activators and
inhibitors to be detected (Figure 4E). We therefore chose 18-20 hr for all further experiments.
AlamarBlue readings are also taken to measure cell viability, using 2.5 μM staurosporine a positive
control for maximum toxicity. The treatment kills >98% of SupT1 cells within 18 hr and
correspondingly inhibits luciferase expression (Supp. Figure 5). Of the 9 candidate compounds two –
DB1033 (IC$_{50}$~25 mM, Figure 4F) and DB2232 (IC$_{50}$~20 mM, Supp. Figure 6A) – inhibited luciferase
activity without significantly affecting luciferase expression from the mutant cell line and without
significant toxicity. Other compounds such as DB2658 showed significant toxicity in the same screen
(Supp. Figure 6B) or were less active (e.g., DB2385, Supp. Figure 6C-H). The DB1033 structure is

shown in Figure 4G. Molecular docking indicated that DB1033 preferentially interacts with DNA minor
groove in the AAAT portion of the consensus octamer sequence, overlapping with the location of the
unstructured Oct1 linker domain (Figure 4H-I).

The reduce the possibility of off-target hits, we tested DB1033 compound activity using an orthogonal
cell-based method. We developed a nano-bioluminescence resonance energy transfer (NanoBRET)
assay that measures OCA-B's interaction with its cognate transcription factor Oct1. In this assay,
293T cells are co-transfected with an OCA-B construct carrying a HaloTag and an Oct1 construct
carrying an nLuc tag (Figure 5A). DB1033 again showed dose-dependent inhibition (Figure 5B),
indicating that DB1033 disrupts the interaction between Oct1 and OCA-B. To test this prediction at a
mechanism level, we used electrophoretic mobility shift assays (EMSA) with recombinant Oct1 DNA
binding domain and recombinant full-length OCA-B to measure formation of Oct1:OCA-B:DNA
complexes (Figure 5C). We used Cy5-conjugated double-stranded DNA to visualize complexes with
cognate DNA in native polyacrylamide gels. Labeled DNA was in excess relative to protein such that
changes in the ability to form a complex can be readily observed. Because of "caging effects",
changes in off-rates are not efficiently captured within a running native gel. Instead, in this assay
effects of an inhibitor on complex formation are likely generated during the incubation prior to gel
loading. We assembled the protein, DNA and buffer components on ice for 25 minutes in stringent
conditions (150 mM KCl), and incubated the mixture for a further 5 min at room temperature
±compound prior to loading. Oct1 readily formed a complex with DNA that was augmented and
slightly supershifted in the presence of OCA-B, indicating the formation of a ternary complex (Figure
5D, lane 3). Inclusion of DB1033 efficiently disrupted complex formation in a dose-dependent manner
(lanes 4-7).

A complimentary in vitro assay uses differential scanning flurimetry (nanoDSF) to measure
Oct1:OCA-B:DNA complex thermal stability. NanoDSF measures the gradual exposure of UV-
absorbing tyrosine side chains as proteins unfold with increasing temperature. We developed a
single-polypeptide chimeric molecule containing the Oct1 DNA binding domain fused to the first 44
amino acids of OCA-B, encompassing the structured part of OCA-B that contacts Oct1 (Figure 5E).
The first 15 amino acids of OCA-B do not appear to be well-structured and presumably serve as a
linker between the two molecules. AlphaFold 3 [51] predicts that the chimeric molecule folds similarly to
natural Oct1 and OCA-B on DNA (Figure 5F). The protein was expressed in *E. coli* with N-terminal 6-
His, consecutive C-terminal Twinstrep and FLAG tags. Following purification, the protein was found to
be mostly pure as assessed by Coomassie staining SDS-PAGE (Supp. Figure 7A). Size exclusion
chromatography indicated that the purified protein was mostly monodisperse, and EMSA indicated
that it was able to interact with consensus octamer DNA (Supp. Figure 7B-C). The presence of target
DNA strongly increased the stability of the chimeric protein in nanoDSF, as expected (Figure 5G),
while adding DB1033 in the presence of DNA destabilized the complex (Figure 5H). These findings
identify DB1033 as the best compound tested, indicate that DB1033 acts to disrupt Oct1:OCA-B
complexes on DNA, and establishes a pipeline for assessing molecules that modulate Oct1 and/or
OCA-B's interactions with one another and with DNA, their overall stability, and their transcriptional
functionality.

Small molecule Oct1:OCA-B inhibitors block α-*PD-1-indcued diabetes and other endocrinopathies in*
NOD mice

We treated 8-week-old male NOD mice with five *i.p.* injections of 15 mg/kg DB1033 or DB2232
simultaneously with α-PD-1 antibody treatment. 8 week-old male NOD.*Ocab^{fl/fl}*;CD4-Cre mice were

used as a positive control for protection. As with genetic OCA-B T cell knockout, DB1033 treatment
completely protected mice against diabetes emergence (Figure 6A-B). These results indicate 1) that
DB1033 is effective in protecting mice against α-PD-1 antibody-induced diabetes, and that 2) acute
inhibition at the same time as α-PD-1 antibody delivery is sufficient to protect mice against diabetes
development. DB1033 toxicity was negligible or low-level, as there was a trending but statistically
nonsignificant loss of weight associated with treated animals at endpoint (Figure 6C). Protection was
not only confined to the pancreatic islets but was also noted with salivary and lacrimal gland tissue,
where leukocyte infiltration was strongly reduced (Figure 6D-E). Importantly, the structurally related
DB2232 compound, which was inferior in cell-based assays, was also functionally inferior at
preventing diabetes (Supp. Figure 7D-F). These findings provide evidence of a structure-activity
relationship linking compound activity to OCA-B complex inhibition. In summary, the findings show
that OCA-B promotes the development of autoimmunity in NOD mice administered ICI while being
dispensable for anti-tumor immunity on multiple strain backgrounds, and identify small molecule
inhibitors that can be administered acutely and phenocopy the genetic knockout.

Discussion

Here we show that OCA-B expression in T cells promotes induced irAEs following ICI treatment while minimally affecting anti-tumor responses. We also, for the first time, show that a small molecule Oct1:OCA-B inhibitor phenocopies the protection against irAEs observed using the OCA-B T cell knockouts. The latter findings indicate that acute OCA-B inhibition is sufficient to protect against irAEs and suggest the existence of a therapeutic window in which OCA-B pharmacological targeting can be used to block autoimmunity while sparing baseline immune function, including immunity to tumors.

OCA-B is one of three tissue-restricted transcription coregulators known as OCA proteins: OCA-B, OCA-T1 and OCA-T2 [52]. Like the other two proteins OCA-B docks with the sequence-specific DNA binding POU transcription factors to regulate transcription [41-43]. The OCA-Ts are master regulators of tuft cells, and a therapeutic vulnerability in small cell lung cancers originating from this cell type [52]. OCA-B has important B cell-intrinsic functions including in the formation of germinal centers [53,54]. In T cells, OCA-B docks with the POU factor Oct1 to regulate target genes encoding cytokines and other immunomodulatory proteins [28]. Because OCA-B expression is 50-100-fold higher in B compared to T cells[26], a calibrated dose of competitive inhibitor could selectively block T cell functions while sparing B cell-intrinsic OCA-B functions. For all three proteins, whole-body null mice are healthy, viable and fertile [52,55,56], suggesting that they a good candidate drug targets, however no small molecules targeting OCA proteins have been described.

Multiple lines of evidence show that pathogenic human and mouse progenitor-like T cells promote and sustain autoimmunity including T1D [57-60]. In mice, OCA-B promotes pathogenesis in experimental autoimmune encephalomyelitis (EAE), a model of multiple sclerosis, through maturation of stem-like cells into encephalitogenic effectors [29]. These properties are T cell intrinsic. In a relapsing-remitting NOD.EAE model, OCA-B T cell-deficiency specifically protects mice from relapse [29]. On the NOD strain background, both OCA-B whole body and T cell-specific knockout block the onset of spontaneous T1D [40]. Using more aggressive T1D mouse models including CD25-mediated T_{reg} depletion and the CD8-restricted, autoreactive NY8.3 transgenic line revealed partial protection, while using the CD4-restricted BDC2.5 and CD8-restricted RIP-mOVA-OT-1 models showed no differences [40]. The protection conferred by OCA-B loss is therefore strongest in more chronic polyclonal models. These findings also reveal that OCA-B T cell loss does not result in global immunosuppression, which would result in broader protection. In female NOD mice with newly arisen T1D, intravenous administration of membrane-penetrating OCA-B peptide mimics briefly reverses elevated blood glucose, as well as pancreatic islet T cell infiltration and pro-inflammatory cytokine production [40]. In contrast to the pancreas, pLN showed no change in T cell numbers or percentages, but did show reduced T cell cytokine responses following self-peptide stimulation [40]. However, greater than three peptide injections was toxic, with mice exhibiting ruffled skin, moribund appearance and weight loss [40]. The Tat membrane-penetrating peptide is associated with acute toxicity that can be mitigated by spacing the dose[61-63].

The ability to trigger endocrine irAEs using α-PD-1 treatment in NOD mice presented an opportunity to test the OCA-B T cell loss of function model in different ways. First, to test the effect of OCA-B loss in an aggressive but physiologically relevant model of autoimmune diabetes. Second, to use the diabetes and other endocrinopathies as models for ICI-triggered irAEs. Third, to determine the role of OCA-B in anti-tumor immunity. We found that OCA-B T cell loss protected NOD mice against ICI-triggered diabetes. The effect of OCA-B loss was partially dependent on the age at which ICI was administered, with 8-week-old mice showing more complete protection compared to 12-week-old

mice. This finding is consistent with a model in which OCA-B loss protects NOD mice at multiple

levels, one of which spans the 8-week timepoint and is PD-1 independent, and one of which spans

the 12-week timepoint and is partially PD-1 dependent. Profiling the immune cells in these mice flow

cytometrically indicated significant reduction in T_{EX}, and a concomitant increase in cells with a

progenitor-like (SLAMF6^{hi}CD39^{lo}) phenotype.

In sharp contrast to the autoimmune protection conferred by OCA-B loss, immune responses and

tumor control were preserved either with or without α-PD-1 treatment. No differences were observed

using immune-cold B16F10 melanoma cells, as well as immune-hot MC38 colon carcinoma cells and

9495 basal cell carcinoma cells. Similar to the preservation of tumor control, loss of OCA-B in T cells

preserves normal primary response and viral control following infection with either the acute strain of

lymphocytic choriomeningitis virus (LCMV) or the glial-tropic β-coronavirus JHMV [26,28,29]. The normal T

cell primary immune response phenotypes observed with Oct1 and OCA-B genetic deletion suggest

the existence of a therapeutic window in which an OCA-B pharmaceutical could be used not just to

block autoimmunity, but could be co-administered with ICIs to block irAEs while sparing immunity to

viruses and tumors. A likely side effect of therapies targeting this pathway would be an inability of

newly-generated memory CD4$^+$ T cells to mount robust recall responses [26,28]. Our findings indicate

that OCA-B promotes a memory-like population of CD4$^+$ T cells important for immune responses to

tumor re-challenge and relapse (manuscript in prep). An OCA-B inhibitor may therefore need to be

withdrawn prior to any potential tumor relapse.

Despite the prevailing but erroneous idea that transcription factors are "undruggable," there are

multiple historical[61,64-66], and recent[67-81] counterexamples, the latter enabled by technologies such as

PROTAC[82,83]. We show that systemic administration a small molecule inhibitor, DB1033, into NOD

mice blocks the emergence of diabetes following the administration of α-PD-1 antibodies. These

molecules disrupt Oct1:OCA-B:DNA complexes in cell culture and in vitro. The finding that DB1033

administration concomitant with α-PD-1 antibody treatment is sufficient to block the induction of

diabetes and other endocrinopathies suggests that OCA-B actions close to the time of injection, and

therefore spatially proximate to the pLNs and pancreatic islets, rather than development of auto-

reactive T cells in an OCA-B deficient environment from the double-positive thymocyte stage onwards,

mediates autoimmune protection. Moreover, we show that a structurally related compound, DB2232,

that is inferior in attenuating OCA-B-dependent transcription in cell culture and in disrupting

Oct1:OCA-B complexes in vitro is also functionally inferior at protecting NOD mice against ICI-

triggered diabetes, strongly suggesting that these compounds act through OCA-B disruption. These

reagents provide a conceptual milestone as a first-in-class small molecule Oct1/OCA-B inhibitor that

can reduce autoimmunity. More work is necessary to develop superior and more drug-like small

molecule inhibitors.

Materials and Methods

Laboratory mice. All mice used in this study were non-obese diabetic (NOD) background, with the exception of B16-F10 and MC38 tumor injections which used C57BL/6 mice. *Pou2af1* (*Ocab*) conditional (floxed) mice on the C57BL/6 and NOD backgrounds were generated as described previously [26,40]. All animal experiments were approved by the University of Utah Institutional Animal Care and Use Committee (IACUC approval 00001553).

α-PD-1-induced diabetes. 250 μg α-PD-1 (clone RMP1-14, Cat# BE0146, Bio X Cell) or rat IgG2a isotype control (Cat# BE0089, Bio X Cell) was interperitoneally (*i.p.*) administered into NOD.*Ocab*[fl/fl] or *Ocab*[fl/fl];CD4-Cre on day 0, 4, 8, and 12 (or on day 0, 4, 8, 12, and 16 as indicated). Antibody was diluted in PBS for injection of 200 μL per mouse.

Histology. Salivary glands and lacrimal glands were isolated and fixed in 10% formalin. After 24-72 hr, tissues were transferred to 70% ethanol and embedded in paraffin, sectioned at 5~7 μm, and H&E staining was performed. Slides were examined and imaged using an Olympus BX53 microscope with UPlanFL objectives, Olympus DP73 camera, and Olympus cellSens imaging software.

Secondary lymphoid organ and tumor harvest. Spleens and lymph nodes were ground and filtered through a 70 μm nylon cell strainer (VWR #76327-100). Red blood cells were removed by Ammonium-Chloride-Potassium (ACK) lysis buffer (10 mM $KHCO_3$, 150 mM NH_4Cl, and 0.1 mM Na_2EDTA). MC38 tumors were manually disrupted by grinding between the frosted end of slide glass in a 6-well plate with 3 mL of PBS plus 4% FBS, and filtered through a 40 μm nylon cell strainer (VWR #76327-098). 9495 tumors were mechanically disrupted in 6-well-plate and digest in RPMI media containing 16mg/mL collagenase D (Sigma-Aldrich #11088858001) and 10mg/mL Dnase (Sigma-Aldrich #10104159001) at 37°C in CO_2 incubator for 45 minutes and filtered through a 40 μm nylon cell strainer, spined in 37.5% percoll gradient, and red blood cells (RBCs) were lysed to produce single-cell suspensions.

Pancreatic islet harvest. Lymphocytes were isolated based on a protocol from Jing et al. [84]. Pancreata were harvested and perfused with collagenase 4 (Worthington #LS004188) at 37°C, and islets were picked using a 10 μL pipette. Islet single-cell suspensions were obtained by vortexing in cell dissociation buffer (ThermoFisher Scientific #13151014).

Flow cytometry. Single-cell samples were stained with antibodies (1:100 diluted in FACS buffer) for flow cytometric analysis. For intracellular staining, cells were cultured in RPMI1640 (ThermoFisher Scientific #11875093) supplemented with 10% FBS, 1% penicillin/streptomycin (ThermoFisher Scientific #15140122) and 1% GlutaMAX (ThermoFisher Scientific #35050061), 1 μL/mL Brefeldin A (BD Biosciences #555029), 50 ng/mL phorbol 12-myristate 13-acetate (Sigma-Aldrich #P1585) and 1 ug/mL ionomycin (Sigma-Aldrich #I0634) for 4 hr. Cells were stained for surface markers and fixed using Cytofix/Cytoperm (BD Biosciences #554714) following the manufacture's protocol, and resuspended in Perm/Wash buffer for intracellular cytokine staining. For intranuclear staining, cells were stained for surface markers, fixed using the Foxp3/Transcription Factor Staining Buffer Kit (ThermoFisher Scientific #00-5523-00) and stained for intranuclear markers. Flow cytometry was performed using a BD Fortessa (BD Biosciences). Data analysis was performed using FlowJo software (BD Biosciences).

Cell lines. Mouse MC38 cell line was a gift from Ryan O'Connell (University of Utah). B16-F10 cells were a gift from the lab of Matthew A. Williams (University of Utah). MC38 cells were grown in DMEM

(ThermoFisher Scientific #11965092) supplemented with 10% FBS, 1% penicillin/streptomycin, 1%
MEM-NEAA (ThermoFisher Scientific #11140050) and 1% HEPES-Cl pH 7.3 (ThermoFisher Scientific
#15630130). B16-F10 cells were grown in DMEM supplemented with 10% FBS and 1%
penicillin/streptomycin. NOD 9495 cells were grown similar to a prior report [50] in DMEM (no glutamine,
no sodium pyruvate, ThermoFisher Scientific #11960044) supplemented with 10% FBS
(ThermoFisher Scientific #A5256701) 1% penicillin/streptomycin, 1% sodium pyruvate, 1% 100×
MEM-NEAA, 1% GlutaMAX (ThermoFisher Scientific #35050061). SupT1 human T lymphoblast cells
were grown in RPMI1640 supplemented with 5% FBS, 5% calf serum (Sigma), 1%
penicillin/streptomycin and 1% GlutaMAX at 37°C with 5% CO_2.

Tumor cell line engraftment. All tumor experiments used 6- to 10-week-old C57BL/6 or 6~8-week-old
NOD background mice. For injections of B16-F10 and MC38, after trypsinization, cultured tumor cells
were resuspended in PBS and 1×10^6 and 5×10^5 tumor cells were *s.c.* injected into the hind flank of
mice, respectively. For the injection of NOD 9495 tumor line, after trypsinization, cultured tumor cells
were resuspended in PBS and 1.5×10^6 tumor cells were *s.c.* injected into the hind flank of mice as
described [50].

Small molecules. Synthesis of DB1033, DB2232, DB2658, DB2379, DB2559, and DB2385 has been
previously reported [85]. All compounds were dissolved in 100% DMSO and stored at 5mM at -80°C for
later use. For cell-based assays, stock solutions or DMSO vehicle controls were diluted in culture
media. For EMSA, compounds were diluted in 0.6X buffer D [86,87]. For mouse injections, sterile PBS
was used to dissolve the compounds.

Generation of SupT1 luciferase reporter cell lines. Reporter constructs were generated using a
modified secreted nano-luciferase vector, pNL2.3 (Promega #N108A), with the hygromycin resistance
cassette replaced with puromycin resistance cassette (to generate pNL2.3-Puro). Gene blocks
(Integrated DNA Technologies) were synthesized containing 5 canonical Oct1 binding motifs
(ATGCAAAT, 5×Oct) or mutant motifs (CAAACTAC, 5×Mut) upstream of a minimal CMV promoter.
Gene blocks were subcloned into the multiple cloning site of pNL2.3-Puro using EcoRV and HindIII
restriction sites. To generate stable transfectants, human SupT1 cells were electroporated using Neon
(ThermoFisher Scientific) with 1 μg of either the pNL2.3-Puro empty-vector (EV), pNL2.3-Puro 5×Oct
(5×Oct), or pNL2.3-Puro 5×Mut (5×Mut) construct. Electroporation parameters were: 1700 V, 10
milliseconds, ×3 pulses. Electroporated cells were allowed to recover for 2 days, then selected for
stable DNA integration and expanded in complete media containing 1 μg/mL puromycin
(ThermoFisher Scientific #A1113803) for 2 weeks. Cells were then single cell-cloned by limited dilution
in complete media containing 1 μg/mL puromycin.

SupT1 OCA-B activity assays. 1.0×10^5 5×Oct and 5×Mut reporter cells were seeded into wells of 96-
well plates were cultured in 100 μL complete RPMI in complete medium with serially diluted
compounds for 18-20 hr. Supernatant was collected and mixed with 1:50 buffer-diluted Nano-Glo
luciferase substrate (Cat# N1110, Promega). Luminescence was measured using a Modulus
Microplate Multimode Reader (Promega).

In vivo treatment of PD-1 blockade-induced diabetes. NOD.*Ocab*^*fl/fl* and NOD.*Ocab*^*fl/fl*;CD4-Cre mice
were *i.p.* administered with α-PD-1 and 15 mg/kg of DB1033 was administered *i.p.* to the mice at day
0, 4, 8, 11, and 12. Mice were sacrificed on day 24.

Statistical analyses. Two-tailed student's t-tests were used to determine all statistical differences. All
error bars represent ±standard deviations unless otherwise noted. All error bars tumor size represents
±SEM unless otherwise denoted. For all figures, ns, $p > 0.05$; *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$;
****, $p \leq 0.0001$.

Construction and purification of recombinant Oct1 DNA binding domain Oct1-fusion proteins. A cDNA
for the Oct1 DNA binding domain (comprising POU_S and POU_H subdomains) fused to OCA_{-B1-44} was
ordered from IDT and included an N-terminal 6-His, consecutive C-terminal Twinstrep and FLAG tags.
the cDNAs were cloned into the pET28a vector (Invitrogen) between the *Nco*I and *Hind*III sites. The
plasmids were transformed into Rosetta 2 (DE3) pLysS competent cells (Sigma) using 50 µg/ml
kanamycin as a selectable marker. Protein was expressed using 1 mM IPTG (Sigma) at 37°C for
three hr. The cells were lysed in buffer containing 20 mM HEPES, pH 7.4 (Sigma), 300 mM NaCl
(Sigma), 5% glycerol (Sigma) and 1 mM DTT (ThermoFisher Scientific) containing EDTA-free
protease inhibitor cocktail (Sigma) at 2X concentration. Protein was bound to a Histrap column
(Cytiva) using the same buffer and eluted with buffer containing 400 mM imidazole (Sigma). The
protein was further purified by size exclusion chromatography on a Superdex increase 75 10/300 GL
column (Cytiva.inc). The column was loaded using PBS containing 1 mM DTT. The protein was
analyzed on 12% SDS-PAGE gel visualized by Coomassie staining.

NanoBRET. The NanoBRET donor plasmid was made by cloning nLuc in-frame with the N-terminus of
full-length human Oct1 between the *Nco*I and *Xba*I restriction sites of pACE-MAM2 (Geneva Biotech).
The acceptor plasmid was made by cloning full-length human OCA-B containing a C-terminal HaloTag
into pHTC vector (Promega) between the *Eco*RI and *Xho*I sites. nanoBRET was performed using the
NanoBRET protein:protein interaction kit (Promega) following the manufacturer protocol. Briefly, 2×10^6
293T cells were plated in 6-well plates were transfected with a total of 2 µg of donor plus acceptor
plasmid at a ratio of 1:100. After 24 hr, cells were harvested and counted. 2×10^5 resuspended cells
were seeded into the wells of 96-well plates using DMEM lacking phenol red (ThermoFisher Scientific)
containing 10% FBS. After 1 hr, DB1033 or DMSO vehicle control were also added at the indicated
concentrations in biological triplicate. After 24 hr, the Nanuck substrate and Halo-ligand were added
before measuring BRET ratio using plate reader (Spectromax iD5, Molecular Devices Inc.). The
miliBRET ratio was plotted with respect to the DB1033 concentration.

EMSA. Published protocols [86,87] were used for EMSA. The octamer sequence probe was 5' labeled
with Cy-5 and used at a final concentration of 40 nM. The probe was incubated with 10 nM Oct1-OCA
B chimeric protein ±equimolar OCA B (1-44) peptide over ice for 15 min prior to the addition of 100
nM-400 nM DB1033, then further incubated over ice for another 15 min. Reaction mixtures were
electrophoresed through 6% native polyacrylamide gels. Images were collected using a Molecular
Dynamics Typhoon scanner using the Cy-5 fluorescence filter.

NanoDSF. Tryptophan fluorescence at 330 nm and 350 nm were simultaneously monitored over a
temperature range from 25°C to 95°C using a NanoTemper Technologies nanoDSF instrument with an
excitation wavelength of 280 nm based on published procedures [88]. The Oct1-OCA B chimeric protein
complexed with double-stranded DNA containing an Oct1-OCA-B binding site was incubated on ice
for 15 min. Protein lacking DNA was used as a control. The effect of DB1033 compound or DMSO
control on the protein DNA complex were measured by comparing the melting point of the
protein/DNA complex ±DB1033 compound. 10 µl from each sample was loaded into each capillary
avoiding any air bubbles. The fluorescence intensity ratio and its first derivative were calculated with

the manufacturer's software (PR.ThermControl, version 2.1.2). The smoothened curves and the first
derivative were plotted in using Microsoft Excel software. The peak from the first-order derivative
indicates the melting temperature of the sample protein sample.

Figure Legends

Figure 1. OCA-B deletion in T cells confers protection against PD-1 blockade-induced diabetes
in NOD mice. (A) Kaplan-Meier plot of diabetes-free mice in 8-week-old male $Ocab^{fl/fl}$ (n=7) and
$Ocab^{fl/fl}$;CD4-Cre (n=7) NOD mice. 250 µg α-PD-1 was administered by intraperitoneal (*i.p.*) injection
at day 0, 4, 8 and 12. Blood glucose levels were measured during different stages of diabetes
development. **(B)** Blood glucose levels of individual mice from (A). **(C)** Islets were prepared from 8-
week-old treated female knockout and littermate control mice 5 days after one dose of α-PD-1
antibody and used in flow cytometry. Representative flow cytometry plots depicting total infiltrating
CD45$^+$ leukocytes for control and knockout mice. **(D)** Quantification of total infiltrating CD45$^+$ islet
leukocyte flow cytometry data from NOD.$Ocab^{fl/fl}$ (n=4) and NOD.$Ocab^{fl/fl}$;CD4-Cre mice (n=3). **(E)**
Representative flow cytometry plots depicting CD4$^+$ and CD8$^+$ islet T cells. Cells were pre-gated using
α-CD3 staining. **(F)** Quantification of flow cytometry data showing the percentages of CD4$^+$ and CD8$^+$
T cells in islets from NOD.$Ocab^{fl/fl}$ (n=4) and NOD.$Ocab^{fl/fl}$;CD4-Cre mice (n=3). **(G)** Representative
flow cytometry plots showing the percentage of PD-1$^+$SLAMF6$^+$CD39$^-$CD8$^+$ T$_{PEX}$ and PD-1$^+$SLAMF6$^-$
CD39$^+$CD8$^+$ T$_{EX}$ in islets (pre-gated on PD-1$^+$CD8$^+$ T cells). **(H)** Quantification of flow cytometry data
for the percentage of T$_{PEX}$ and T$_{EX}$ in islets from NOD.$Ocab^{fl/fl}$ (n=4) and NOD.$Ocab^{fl/fl}$;CD4-Cre mice
(n=3).

Figure 2. OCA-B in T cells is dispensable for anti-tumor immunity in C57BL/6 mice. (A) 5×10^5
MC38 tumor cells were *s.c.* implanted into the hind flanks of female C57BL/6 $Ocab^{fl/fl}$ and
$Ocab^{fl/fl}$;CD4-Cre mice. 300 µg α-PD-1 (or PBS) was administered *i.p.* at day 4, 7, 10. Tumor sizes
were measured every 3-4 days, and mice were euthanized at end point (day 23) or upon tumor size
reaching 2 cm in any direction. $Ocab^{fl/fl}$+PBS (n=8), $Ocab^{fl/fl}$;CD4-Cre+PBS (n=11), $Ocab^{fl/fl}$+α-PD-1
(n=8), and $Ocab^{fl/fl}$;CD4-Cre+α-PD-1 (n=8). **(B)** Mean tumor sizes from (A) are shown. $Ocab^{fl/fl}$+PBS
(n=8), $Ocab^{fl/fl}$;CD4-Cre+PBS (n=11), $Ocab^{fl/fl}$+α-PD-1 (n=8), and $Ocab^{fl/fl}$;CD4-Cre+α-PD-1 (n=8). **(C)**
Quantification of flow cytometry data showing the percentage of CD45.2$^+$ tumor-infiltrating leukocytes
(TILs) in MC38 TMEs from α-PD-1 antibody-treated $Ocab^{fl/fl}$ (n=6) and $Ocab^{fl/fl}$;CD4-Cre (n=6) mice on
day 23 or upon tumor size reaching 2 cm in any direction. **(D)** Quantification of flow cytometry data
showing the ratio of CD8$^+$ to CD4$^+$ T cells in the TME. **(E)** Representative flow cytometry plots of
endpoint TME CD4$^+$ and CD8$^+$ T cell SLAMF6 and CD39 expression. $Ocab^{fl/fl}$ (n=6) and $Ocab^{fl/fl}$;CD4-
Cre (n=5) tumor-bearing mice were treated with α-PD-1 antibodies. **(F)** Quantification of flow
cytometry data for endpoint TME CD4$^+$ and CD8$^+$ T cell SLAMF6 and CD39 expression from $Ocab^{fl/fl}$
(n=6) and $Ocab^{fl/fl}$;CD4-Cre (n=5) mice. **(G)** Representative flow cytometry data for endpoint TME
CD8$^+$ and CD4$^+$ T cell TIM-3 expression. **(H)** Quantification of flow cytometry data at endpoint showing
TME CD4$^+$ and CD8$^+$ T cell TIM-3 expression from $Ocab^{fl/fl}$ (n=6) and $Ocab^{fl/fl}$;CD4-Cre (n=6) mice.

Figure 3. OCA-B in T cells is dispensable for anti-tumor immunity in NOD mice. (A) 1.5×10^6
9495 tumor cells were *s.c.* implanted into the hind flanks of male NOD.$Ocab^{fl/fl}$ and NOD.$Ocab^{fl/fl}$;CD4-
Cre mice. Three doses of 200 µg α-PD-1 (or PBS vehicle control) were *i.p.* administered at day 10, 14
and 18. Tumor sizes were measured during different stages of tumor growth. $Ocab^{fl/fl}$+PBS (n=3),
$Ocab^{fl/fl}$;CD4-Cre+PBS (n=4), $Ocab^{fl/fl}$+α-PD-1 (n=8), and $Ocab^{fl/fl}$;CD4-Cre+α-PD-1 (n=12). **(B)**
Quantification of flow cytometry data for the percentage of CD45$^+$ infiltrating cells in the TME from α-
PD-1 antibody-treated control $Ocab^{fl/fl}$ (n=10) and littermate experimental $Ocab^{fl/fl}$;CD4-Cre (n=6) mice.

(C) Representative flow cytometry plots depicting CD4$^+$ and CD8$^+$ TME T cells from α-PD-1 antibody-treated control NOD.*Ocab*$^{fl/fl}$ and littermate experimental NOD.*Ocab*$^{fl/fl}$;CD4-Cre mice. Cells were pre-gated using α-CD3 staining. (**D**) Quantification of flow cytometry data depicting the percentage of CD4$^+$ and CD8$^+$ T cells in the TME from NOD.*Ocab*$^{fl/fl}$ (n=9) and NOD.*Ocab*$^{fl/fl}$;CD4-Cre mice (n=4). (**E**) Quantification of flow cytometry data showing the percentage of activated (CD62L^{lo}CD44^{hi}) CD8$^+$ T cells in the TME of α-PD-1-treated *Ocab*$^{fl/fl}$ (n=9) and littermate experimental *Ocab*$^{fl/fl}$;CD4-Cre (n=5) mice at endpoint. (**F**) Representative flow cytometry plots depicting relative percentages of SLAMF6- and CD39-expressing infiltrating CD8$^+$PD-1$^+$ T cells. (**G**) Quantification of PD-1$^+$SLAMF6$^+$CD39$^-$CD8$^+$ T$_{PEX}$ and PD-1$^+$SLAMF6$^-$CD39$^+$CD8$^+$ T$_{EX}$ in the TME from NOD.*Ocab*$^{fl/fl}$ (n=9) and NOD.*Ocab*$^{fl/fl}$;CD4-Cre mice (n=5). (**H**) Representative flow cytometry plots depicting TIM-3 levels in infiltrating CD8$^+$ T cells. (**I**) Quantification of TIM-3$^+$ cells in the TME from NOD.*Ocab*$^{fl/fl}$ (n=9) and NOD.*Ocab*$^{fl/fl}$;CD4-Cre mice (n=5).

Figure 4. OCA-B cellular inhibition by small molecule DNA minor groove binders (**A**) Oct1:OCA-B:DNA co-crystal structure [43] showing two targeting approaches: a peptide that mimics the key OCA-B alpha helix involved in downstream effector functions (top lightning bolt) [40], and a minor groove DNA binding molecule with potential to disrupt the unstructured Oct1 linker domain between the two DNA binding subdomains (bottom lightning bolt). (**B**) OCA-B immunoblots from SupT1 and Jurkat T cell lines (lanes 3-4). Isolated mouse primary thymocyte and 293T cell lysates were also used as positive and negative controls, respectively (lanes 1-2). Histone H3 is shown as a loading control. (**C**) Schematic for OCA-B luciferase reporter cell lines. SupT1 cells were stably transfected with reporter plasmids containing five tandem canonical Oct1:OCA-B binding sites linked to a CMV promoter upstream of secreted nLuc (5×Oct) or an identical plasmid except containing mutant binding sites used to determine nonspecific effects (5×Mut). Cells were selected with puromycin, single-cell cloned, and individual clones used subsequently. (**D**) Luciferase activity of pNL2.3 empty vector ("EV"), 5×Oct, or 5×Mut reporter lines. Cells were incubated for 18 hr after plating. (**E**) Luciferase expression kinetics of cultured 5×Oct cells. A timecourse of 2-46 hr is shown. Averages are shown from n=5 biological replicate platings. Trendline shows logistic growth fit. (**F**) Inhibition of cellular OCA-B activity by an escalating dose of the small molecule DB1033. Values from 5×Oct and 5×Mut clones were normalized to one another in the absence of DB1033. Averages are shown across n=6 biological replicate platings. Red curve shows the "5×Oct" fit to the Hill equation. Asterisks denote results from two-tailed student T-test comparing the two lines at the highest concentration. (**G**) Chemical structure of DB1033. (**H**) Molecular docking of DB1033 to canonical octamer B-form DNA. Red shows water molecules. DB1033 is shown in purple. Red/underlined nucleotides show the minor groove bases contacted by DB1033 docked with the lowest free energy. (**I**) Wireframe model showing DB1033 bound to the minor groove AAAT site. Hydrogen bond interactions formed between DB1033 and the minor groove AAAT site are shown with dashed lines.

Figure 5. DB1033 inhibits OCA-B-Oct1-DNA complex formation. (**A**) Schematics of full-length Oct1 tagged with N-terminal nLuc and full-length OCA-B with a C-terminal HaloTag used for nanoBRET. (**B**) Quantification of nanoBRET signal obtained by co-transfecting the 293T cells with the constructs shown in A in the presence of increasing concentrations of DB1033. An average of n=3 biological replicate platings is shown. DB1033 was applied at the indicated concentrations 1 hr after seeding the cells and cells were collected after a further 24 hr.(**C**) Schematics of the C-terminally

Twinstep-tagged Oct1 DNA-binding domain (Oct1$_{DBD}$) and full-length OCA-B with a C-terminal FLAG

tag used in EMSA. (**D**) Cy5-DNA scan of native PAGE EMSA showing DNA-Oct1$_{DBD}$ complex

formation with OCA-B, and concentration-dependent inhibition by DB1033. (**E**) A schematic of the

Oct1$_{DBD}$:OCA-B$_{1-44}$ fusion protein engineered for nanoDSF. (**F**) Structure of the OCA-B$_{1-44}$-Oct1$_{DBD}$

complex predicted using Alphafold 3. Cyan: OCA-B$_{1-44}$ peptide. Green: Oct1$_{DBD}$ peptide. Dark blue

shows the Oct1:DNA structure derived from X-ray crystallography [43]. (**G**) nanoDSF measurement of

the thermal stability of 10 μM Oct1$_{DBD}$-OCA-B$_{1-44}$ chimeric protein with or without added canonical

octamer DNA. First derivative is shown below. (**H**) First derivative nanoDSF data in the presence of

DNA and the presence (above) or absence (below) of 100 μM DB1033.

Figure 6. DB1033 inhibits PD-1 blockade-induced diabetes in NOD mice. (**A**) Kaplan-Meier plot of

diabetes-free mice in 8-week-old male NOD.*Ocab*$^{fl/fl}$ treated with PBS (n=4), NOD.*Ocab*$^{fl/fl}$ treated with

DB1033 (n=4) and NOD.*Ocab*$^{fl/fl}$;CD4-Cre mice treated with PBS (n=4). Black arrows: 250 µg α-PD-1

and 15 mg/kg *i.p.* injections at day 0, 4, 8, 12. Grey arrow: an additional *i.p.* injection of 15 mg/kg

DB1033 was administered on day 11. (**B**) Average blood glucose levels of mice in (A). (**C**) Mouse

body weight on day 24 of the experiment described in (A). (**D**) Example H&E images of lacrimal and

salivary gland sections harvested from 8-week-old male NOD mice injected with α-PD-1 antibodies in

(A). (**E**) Quantification of the histological scoring described in (D). NOD.*Ocab*$^{fl/fl}$ treated with PBS

(n=4), NOD.*Ocab*$^{fl/fl}$ treated with DB1033 (n=4) and NOD.*Ocab*$^{fl/fl}$;CD4-Cre mice treated with PBS

(n=4). Histologic scoring was conducted in a blinded fashion.

Supplemental Figure Legends

Figure S1. OCA-B deletion in T cells protects NOD mice from spontaneous and PD-1 blockade-

induced diabetes. (**A**) Kaplan-Meier plot of diabetes-free mice in 12-week-old female littermate

control NOD.*Ocab*$^{fl/fl}$ (n=5) or experimental NOD.*Ocab*$^{fl/fl}$;CD4-Cre (n=6) mice receiving 250 µg α-PD-1

antibody injections at day 0, 4, 8, 12, and 16. Spontaneous T1D was monitored in NOD.*Ocab*$^{fl/fl}$ (n=5)

and NOD.*Ocab*$^{fl/fl}$;CD4-Cre (n=5) mice receiving PBS. (**B**) Blood glucose levels were plotted from

individual mice receiving α-PD-1 antibody in (A). (**C**) Flow cytometry plots showing percentages of

CD8$^+$PD-1$^+$TIM-3$^+$ islet T cells collected from 12-week-old mice 3 days post-α-PD-1 treatment. (**D**)

Quantification of the flow cytometry data from *Ocab*$^{fl/fl}$ (n=4) and *Ocab*$^{fl/fl}$;CD4-Cre mice (n=3). (**E**) 12

week-old NOD.*Ocab*$^{fl/fl}$ or *Ocab*$^{fl/fl}$;CD4-Cre mice were injected once with α-PD-1 antibodies. Islet CD8$^+$

T cells were assessed for SLAMF6 and CD39 by flow cytometry. An example *Ocab*$^{fl/fl}$ and example

Ocab$^{fl/fl}$;CD4-Cre mouse are shown. (**F**) Quantification of flow cytometry data from *Ocab*$^{fl/fl}$ (n=4) and

Ocab$^{fl/fl}$;CD4-Cre mice (n=3).

Figure S2. OCA-B deletion in T cells leaves B16F10 tumor growth and immune response

unaltered. Growth of *s.c.* B16F10 tumors in C57BL/6 mice sufficient (blue, *Ocab*$^{fl/fl}$) and deficient (red,

Ocab$^{fl/fl}$;CD4-Cre) for OCA-B in T cells.

Figure S3. OCA-B deletion in T cells decreases the T$_{CM}$- and naïve-like CD4$^+$ and CD8$^+$ T cell

populations in MC38 TME. (A) Representative plots of flow cytometric analysis performed on tumors

harvested from 6-10-week-old female α-PD-1-treated C57BL/6.*Ocab*^{fl/fl} and C57BL/6.*Ocab*^{fl/fl};CD4-Cre
mice on day 23 post-tumor inoculation (or upon tumor size reaching 2 cm in any direction). CD4$^+$ and
CD8$^+$ T$_{CM}$-like (CD62L$^+$CD44$^+$) and T$_N$-like (CD62L$^+$CD44$^-$) cells in the TME are shown. (**B**)
Quantification of the percentages of T$_{CM}$-like (CD62L$^+$CD44$^+$) and T$_N$-like (CD62L$^+$CD44$^-$) in TME
CD4$^+$ cells (n=5 *Ocab*^{fl/fl} and n=5 *Ocab*^{fl/fl};CD4-Cre) and CD8$^+$ cells from *Ocab*^{fl/fl} controls and
Ocab^{fl/fl};CD4-Cre (n=5 *Ocab*^{fl/fl} and n=5 *Ocab*^{fl/fl};CD4-Cre for T$_N$-like and n=5 *Ocab*^{fl/fl} and n=6
Ocab^{fl/fl};CD4-Cre for T$_{CM}$-like).

Figure S4. NOD tumor additional data. (A) Kaplan-Meier plot of another cohort of diabetes-free
mice performed with the same experiment procedures as in Figure 3. **(B-C)** Average blood glucose
levels of mice in panel A.

Figure S5. Effect of staurosporine on luciferase-based transcription activity in SupT1 cells. A
dose curve of staurosporine is shown using both the 5×Oct and 5×Mut cell lines. Trendlines show the
fit to the Hill equation.

Figure S6. OCA-B cellular inhibition by additional small molecule DNA minor groove binders.
(**A**) Inhibition of cellular OCA-B activity by DB2232. Blue datapoints represent control measurements
of nonspecific transcription measured in 5×Mut cells. Red datapoints represent experimental
measurements from the 5×Oct line. Luciferase values from both cells lines in were normalized to 1.0
in the absence of DB2232. Trendlines show the fit to the Hill equation. (**B-H**) Similar data for additional
molecules, except cytotoxicity was measured using AlamarBlue.

Figure S7. DB1033 and DB2232 effect on PD-1 blockade-induced diabetes in NOD mice. (A)
Image of a 12% SDS-polyacrylamide gel loaded with the purified recombinant chimeric Oct1 DBD-
OCA-B protein and stained with Coomassie blue. Molecular weight makers are shown at left as a
standard. **(B)** Size exclusion chromatography elution profile using the Oct1-OCA-B chimera. **(C)**
EMSA using the Oct1-OCA-B chimera and Cy5-labeled prefer octamer consensus DNA. **(D)** Kaplan-
Meier plot of diabetes-free mice in 8-week-old male NOD.*Ocab*^{fl/fl} mice treated with PBS (n=7),
NOD.*Ocab*^{fl/fl} mice treated with DB2232 (n=4) and NOD.*Ocab*^{fl/fl};CD4-Cre mice treated with PBS
(n=7). Black arrows: 250 µg α-PD-1 and 15 mg/kg *i.p.* injections at day 0, 4, 8, 12. Grey arrow: an
additional *i.p.* injection of 15 mg/kg DB2232 was administered on day 11. **(E)** Average blood glucose
levels of mice in (D). **(F)** Body weight of mice at day 24 of the experiment described in (D).

Acknowledgements

We thank R. O'Connell and M. Williams for MC38 and B16F10 cells, and A. Young for the 9495 cells. We thank O. Allen, B. Dalley and the University of Utah High-Throughput Genomics Core facility. We thank J. Marvin and the University of Utah Flow Cytometry Core facility. We acknowledge the Huntsman Cancer Institute Cancer Center Support Grant (CCSG). This work was supported by grants from the National Institutes of Health/National Institute of Allergy and Infectious Diseases (R01AI162929), the Praespero Autoimmune Fund (Canada), and an internal seed grant from the Huntsman Cancer Institute Genomics, Epigenetics and Metabolism (GEM) program to DT.

Author contributions

JD, AKM, EPH and PB performed experiments and analyzed data. AY provided key reagents and methods. WDW and AAF synthesized and purified DNA minor groove binding small molecules. DT conceptualized and supervised the study. All authors contributed to the writing of the manuscript.

References

1.   National Diabetes Statistics Report. (2024). Centers for Disease Control and Prevention.
2.   Ogrotis, I., Koufakis, T., and Kotsa, K. (2023). Changes in the Global Epidemiology of Type 1
Diabetes in an Evolving Landscape of Environmental Factors: Causes, Challenges, and
Opportunities. Medicina (Kaunas) *59*. 10.3390/medicina59040668.
3.   Gregory, G.A., Robinson, T.I.G., Linklater, S.E., Wang, F., Colagiuri, S., de Beaufort, C.,
Donaghue, K.C., Magliano, D.J., Maniam, J., Orchard, T.J., et al. (2022). Global incidence,
prevalence, and mortality of type 1 diabetes in 2021 with projection to 2040: a modelling study.
Lancet Diabetes Endocrinol *10*, 741-760. 10.1016/S2213-8587(22)00218-2.
4.   Cooke, D.W., and Plotnick, L. (2008). Type 1 diabetes mellitus in pediatrics. Pediatr Rev *29*,
374-384; quiz 385.
5.   Sutherland, D.E., Sibley, R., Xu, X.Z., Michael, A., Srikanta, A.M., Taub, F., Najarian, J., and
Goetz, F.C. (1984). Twin-to-twin pancreas transplantation: reversal and reenactment of the
pathogenesis of type I diabetes. Trans Assoc Am Physicians *97*, 80-87.
6.   Vendrame, F., Pileggi, A., Laughlin, E., Allende, G., Martin-Pagola, A., Molano, R.D.,
Diamantopoulos, S., Standifer, N., Geubtner, K., Falk, B.A., et al. (2010). Recurrence of type 1
diabetes after simultaneous pancreas-kidney transplantation, despite immunosuppression, is
associated with autoantibodies and pathogenic autoreactive CD4 T-cells. Diabetes *59*, 947-
957. 10.2337/db09-0498.
7.   Wang, Y., Pontesilli, O., Gill, R.G., La Rosa, F.G., and Lafferty, K.J. (1991). The role of CD4+
and CD8+ T cells in the destruction of islet grafts by spontaneously diabetic mice. Proc Natl
Acad Sci U S A *88*, 527-531. 10.1073/pnas.88.2.527.
8.   Lernmark, A., and Larsson, H.E. (2013). Immune therapy in type 1 diabetes mellitus. Nat Rev
Endocrinol *9*, 92-103.
9.   Tawbi, H.A., Schadendorf, D., Lipson, E.J., Ascierto, P.A., Matamala, L., Castillo Gutierrez, E.,
Rutkowski, P., Gogas, H.J., Lao, C.D., De Menezes, J.J., et al. (2022). Relatlimab and
Nivolumab versus Nivolumab in Untreated Advanced Melanoma. N Engl J Med *386*, 24-34.
10.1056/NEJMoa2109970.
10.   Hamid, O., Robert, C., Daud, A., Hodi, F.S., Hwu, W.J., Kefford, R., Wolchok, J.D., Hersey, P.,
Joseph, R.W., Weber, J.S., et al. (2013). Safety and tumor responses with lambrolizumab
(anti-PD-1) in melanoma. N Engl J Med *369*, 134-144. 10.1056/NEJMoa1305133.
11.   Cortes, J., Rugo, H.S., Cescon, D.W., Im, S.A., Yusof, M.M., Gallardo, C., Lipatov, O., Barrios,
C.H., Perez-Garcia, J., Iwata, H., et al. (2022). Pembrolizumab plus Chemotherapy in
Advanced Triple-Negative Breast Cancer. N Engl J Med *387*, 217-226.
10.1056/NEJMoa2202809.
12.   Postow, M.A., Callahan, M.K., and Wolchok, J.D. (2015). Immune Checkpoint Blockade in
Cancer Therapy. J Clin Oncol *33*, 1974-1982. 10.1200/JCO.2014.59.4358.
13.   Berti, A., Bortolotti, R., Dipasquale, M., Kinspergher, S., Prokop, L., Grandi, G., Inchiostro, S.,
Paolazzi, G., Caffo, O., and Veccia, A. (2021). Meta-analysis of immune-related adverse
events in phase 3 clinical trials assessing immune checkpoint inhibitors for lung cancer. Crit
Rev Oncol Hematol *162*, 103351. 10.1016/j.critrevonc.2021.103351.
14.   Chen, T.W., Razak, A.R., Bedard, P.L., Siu, L.L., and Hansen, A.R. (2015). A systematic review
of immune-related adverse event reporting in clinical trials of immune checkpoint inhibitors.
Ann Oncol *26*, 1824-1829. 10.1093/annonc/mdv182.
15.   Postow, M.A., Sidlow, R., and Hellmann, M.D. (2018). Immune-Related Adverse Events
Associated with Immune Checkpoint Blockade. N Engl J Med *378*, 158-168.
10.1056/NEJMra1703481.
16.   Young, A., Quandt, Z., and Bluestone, J.A. (2018). The Balancing Act between Cancer
Immunity and Autoimmunity in Response to Immunotherapy. Cancer Immunol Res *6*, 1445-
1452. 10.1158/2326-6066.CIR-18-0487.
17.   Molina, G.E., Zubiri, L., Cohen, J.V., Durbin, S.M., Petrillo, L., Allen, I.M., Murciano-Goroff,
Y.R., Dougan, M., Thomas, M.F., Faje, A.T., et al. (2021). Temporal Trends and Outcomes

691   Among Patients Admitted for Immune-Related Adverse Events: A Single-Center Retrospective
692   Cohort Study from 2011 to 2018. Oncologist *26*, 514-522. 10.1002/onco.13740.

693 18. Stamatouli, A.M., Quandt, Z., Perdigoto, A.L., Clark, P.L., Kluger, H., Weiss, S.A., Gettinger, S.,
694   Sznol, M., Young, A., Rushakoff, R., et al. (2018). Collateral Damage: Insulin-Dependent
695   Diabetes Induced With Checkpoint Inhibitors. Diabetes *67*, 1471-1480. 10.2337/dbi18-0002.

696 19. Quandt, Z., Young, A., Perdigoto, A.L., Herold, K.C., and Anderson, M.S. (2021). Autoimmune
697   Endocrinopathies: An Emerging Complication of Immune Checkpoint Inhibitors. Annu Rev Med
698   *72*, 313-330. 10.1146/annurev-med-050219-034237.

699 20. Osorio, J.C., Ni, A., Chaft, J.E., Pollina, R., Kasler, M.K., Stephens, D., Rodriguez, C.,
700   Cambridge, L., Rizvi, H., Wolchok, J.D., et al. (2017). Antibody-mediated thyroid dysfunction
701   during T-cell checkpoint blockade in patients with non-small-cell lung cancer. Ann Oncol *28*,
702   583-589. 10.1093/annonc/mdw640.

703 21. Noble, J.A., and Valdes, A.M. (2011). Genetics of the HLA region in the prediction of type 1
704   diabetes. Curr Diab Rep *11*, 533-542. 10.1007/s11892-011-0223-x.

705 22. Hu, H., Zakharov, P.N., Peterson, O.J., and Unanue, E.R. (2020). Cytocidal macrophages in
706   symbiosis with CD4 and CD8 T cells cause acute diabetes following checkpoint blockade of
707   PD-1 in NOD mice. Proc Natl Acad Sci U S A *117*, 31319-31330. 10.1073/pnas.2019743117.

708 23. Sims, E.K., Bundy, B.N., Stier, K., Serti, E., Lim, N., Long, S.A., Geyer, S.M., Moran, A.,
709   Greenbaum, C.J., Evans-Molina, C., et al. (2021). Teplizumab improves and stabilizes beta
710   cell function in antibody-positive high-risk individuals. Sci Transl Med *13*.
711   10.1126/scitranslmed.abc8980.

712 24. Ramos, E.L., Dayan, C.M., Chatenoud, L., Sumnik, Z., Simmons, K.M., Szypowska, A.,
713   Gitelman, S.E., Knecht, L.A., Niemoeller, E., Tian, W., et al. (2023). Teplizumab and beta-Cell
714   Function in Newly Diagnosed Type 1 Diabetes. N Engl J Med *389*, 2151-2161.
715   10.1056/NEJMoa2308743.

716 25. Zwilling, S., Dieckmann, A., Pfisterer, P., Angel, P., and Wirth, T. (1997). Inducible expression
717   and phosphorylation of coactivator BOB.1/OBF.1 in T cells. Science *277*, 221-225.

718 26. Sun, W., Hughes, E.P., Kim, H., Perovanovic, J., Charley, K.R., Perkins, B., Du, J., Ibarra, A.,
719   Syage, A.R., Hale, J.S., et al. (2024). OCA-B/Pou2af1 is sufficient to promote CD4+ T cell
720   memory and prospectively identifies memory precursors. Proc Natl Acad Sci U S A *121*,
721   e2309153121. 10.1073/pnas.2309153121.

722 27. Shakya, A., Kang, J., Chumley, J., Williams, M.A., and Tantin, D. (2011). Oct1 is a switchable,
723   bipotential stabilizer of repressed and inducible transcriptional states. J Biol Chem *286*, 450-
724   459. 10.1074/jbc.M110.174045.

725 28. Shakya, A., Goren, A., Shalek, A., German, C.N., Snook, J., Kuchroo, V.K., Yosef, N., Chan,
726   R.C., Regev, A., Williams, M.A., and Tantin, D. (2015). Oct1 and OCA-B are selectively
727   required for CD4 memory T cell function. J Exp Med *212*, 2115-2131. 10.1084/jem.20150363.

728 29. Hughes, E.P., Syage, A.R., Mirzaei Mehrabad, E., Lane, T.E., Spike, B.T., and Tantin, D.
729   (2025). OCA-B promotes pathogenic maturation of stem-like CD4+ T cells and autoimmune
730   demyelination. J Clin Invest. 10.1172/JCI187862.

731 30. Nakamura, M., Nishida, N., Kawashima, M., Aiba, Y., Tanaka, A., Yasunami, M., Nakamura, H.,
732   Komori, A., Nakamuta, M., Zeniya, M., et al. (2012). Genome-wide association study identifies
733   TNFSF15 and POU2AF1 as susceptibility loci for primary biliary cirrhosis in the Japanese
734   population. Am J Hum Genet *91*, 721-728.

735 31. Goris, A., Sawcer, S., Vandenbroeck, K., Carton, H., Billiau, A., Setakis, E., Compston, A., and
736   Dubois, B. (2003). New candidate loci for multiple sclerosis susceptibility revealed by a whole
737   genome association screen in a Belgian population. J Neuroimmunol *143*, 65-69.
738   10.1016/j.jneuroim.2003.08.013.

739 32. Ban, M., Booth, D., Heard, R., Stewart, G., Goris, A., Vandenbroeck, K., Dubois, B.,
740   Laaksonen, M., Ilonen, J., Alizadeh, M., et al. (2006). Linkage disequilibrium screening for
741   multiple sclerosis implicates JAG1 and POU2AF1 as susceptibility genes in Europeans. J
742   Neuroimmunol *179*, 108-116.

33. van Heel, D.A., Udalova, I.A., De Silva, A.P., McGovern, D.P., Kinouchi, Y., Hull, J., Lench,
N.J., Cardon, L.R., Carey, A.H., Jewell, D.P., and Kwiatkowski, D. (2002). Inflammatory bowel
disease is associated with a TNF polymorphism that affects an interaction between the OCT1
and NF(-kappa)B transcription factors. Hum Mol Genet *11*, 1281-1289.
34. Cunninghame Graham, D.S., Wong, A.K., McHugh, N.J., Whittaker, J.C., and Vyse, T.J.
(2006). Evidence for unique association signals in SLE at the CD28-CTLA4-ICOS locus in a
family-based study. Hum Mol Genet *15*, 3195-3205.
35. Leon Rodriguez, D.A., Carmona, F.D., Echeverria, L.E., Gonzalez, C.I., and Martin, J. (2016).
IL18 Gene Variants Influence the Susceptibility to Chagas Disease. PLoS Negl Trop Dis *10*,
e0004583.
36. Kiesler, P., Shakya, A., Tantin, D., and Vercelli, D. (2009). An Allergy-Associated Polymorphism
in a Novel Regulatory Element Enhances IL13 Expression. Hum Molec Genet *18*, 4513-4520.
37. Vince, N., Li, H., Ramsuran, V., Naranbhai, V., Duh, F.M., Fairfax, B.P., Saleh, B., Knight, J.C.,
Anderson, S.K., and Carrington, M. (2016). HLA-C Level Is Regulated by a Polymorphic Oct1
Binding Site in the HLA-C Promoter Region. Am J Hum Genet *99*, 1353-1358.
38. Farh, K.K., Marson, A., Zhu, J., Kleinewietfeld, M., Housley, W.J., Beik, S., Shoresh, N.,
Whitton, H., Ryan, R.J., Shishkin, A.A., et al. (2015). Genetic and epigenetic fine mapping of
causal autoimmune disease variants. Nature *518*, 337-343.
39. Maurano, M.T., Humbert, R., Rynes, E., Thurman, R.E., Haugen, E., Wang, H., Reynolds, A.P.,
Sandstrom, R., Qu, H., Brody, J., et al. (2012). Systematic localization of common disease-
associated variation in regulatory DNA. Science *337*, 1190-1195. 10.1126/science.1222794.
40. Kim, H., Perovanovic, J., Shakya, A., Shen, Z., German, C.N., A., I., Jafek, J.L., Lin, N.-P.,
Evavold, B.D., Chou, D.H., et al. (2021). Targeting OCA-B/Pou2af1 blocks activated,
autoreactive T cells in the pancreas and type-1 diabetes. J Exp Med *218*, e20200533.
10.1084/jem.20200533.
41. Strubin, M., Newell, J.W., and Matthias, P. (1995). OBF-1, a novel B cell-specific coactivator
that stimulates immunoglobulin promoter activity through association with octamer-binding
proteins. Cell *80*, 497-506.
42. Gstaiger, M., Knoepfel, L., Georgiev, O., Schaffner, W., and Hovens, C.M. (1995). A B-cell
coactivator of octamer-binding transcription factors. Nature *373*, 360-362.
43. Chasman, D., Cepek, K., Sharp, P.A., and Pabo, C.O. (1999). Crystal structure of an OCA-B
peptide bound to an Oct-1 POU domain/octamer DNA complex: specific recognition of a
protein-DNA interface. Genes Dev. *13*, 2650-2657.
44. Ansari, M.J., Salama, A.D., Chitnis, T., Smith, R.N., Yagita, H., Akiba, H., Yamazaki, T., Azuma,
M., Iwai, H., Khoury, S.J., et al. (2003). The programmed death-1 (PD-1) pathway regulates
autoimmune diabetes in nonobese diabetic (NOD) mice. J Exp Med *198*, 63-69.
10.1084/jem.20022125.
45. Fife, B.T., Guleria, I., Gubbels Bupp, M., Eagar, T.N., Tang, Q., Bour-Jordan, H., Yagita, H.,
Azuma, M., Sayegh, M.H., and Bluestone, J.A. (2006). Insulin-induced remission in new-onset
NOD mice is maintained by the PD-1-PD-L1 pathway. J Exp Med *203*, 2737-2747.
10.1084/jem.20061577.
46. Zhou, J., Jin, J.O., Kawai, T., and Yu, Q. (2016). Endogenous programmed death ligand-1
restrains the development and onset of Sjögren's syndrome in non-obese diabetic mice. Sci
Rep *6*, 39105. 10.1038/srep39105.
47. Ippolito, S., Di Dalmazi, G., Pani, F., Sabini, E., and Caturegli, P. (2021). Distinct Cytokine
Signatures in Thyroiditis Induced by PD-1 or CTLA-4 Blockade: Insights from a New Mouse
Model. Thyroid *31*, 1839-1849. 10.1089/thy.2021.0165.
48. Martinez-Usatorre, A., Donda, A., Zehn, D., and Romero, P. (2018). PD-1 Blockade Unleashes
Effector Potential of Both High- and Low-Affinity Tumor-Infiltrating T Cells. J Immunol *201*, 792-
803. 10.4049/jimmunol.1701644.
49. Homet Moreno, B., Zaretsky, J.M., Garcia-Diaz, A., Tsoi, J., Parisi, G., Robert, L., Meeth, K.,
Ndoye, A., Bosenberg, M., Weeraratna, A.T., et al. (2016). Response to Programmed Cell

Death-1 Blockade in a Murine Melanoma Syngeneic Model Requires Costimulation, CD4, and
CD8 T Cells. Cancer Immunol Res *4*, 845-857. 10.1158/2326-6066.CIR-16-0060.

50.   Young, A., Medrano, R., Wynn, P., Kang, J.H., Hansen, C., Cina, M., Fadlullah, M.C.H.,
Sheehan, K.C.F., Quandt, Z., Nahar, K., et al. (2025). Improving preclinical modeling of irAEs
and anti-tumor immunity using syngeneic tumor models in autoimmune-prone NOD mice. in
prep.

51.   Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O.,
Willmore, L., Ballard, A.J., Bambrick, J., et al. (2024). Accurate structure prediction of
biomolecular interactions with AlphaFold 3. Nature *630*, 493-500. 10.1038/s41586-024-07487-
w.

52.   Wu, X.S., He, X.Y., Ipsaro, J.J., Huang, Y.H., Preall, J.B., Ng, D., Shue, Y.T., Sage, J.,
Egeblad, M., Joshua-Tor, L., and Vakoc, C.R. (2022). OCA-T1 and OCA-T2 are coactivators of
POU2F3 in the tuft cell lineage. Nature *607*, 169-175. 10.1038/s41586-022-04842-7.

53.   Betzler, A.C., Fiedler, K., Hoffmann, T.K., Fehling, H.J., Wirth, T., and Brunner, C. (2022).
BOB.1/OBF.1 is required during B-cell ontogeny for B-cell differentiation and germinal center
function. Eur J Immunol *52*, 404-417. 10.1002/eji.202149333.

54.   Shi, W., Liao, Y., Willis, S.N., Taubenheim, N., Inouye, M., Tarlinton, D.M., Smyth, G.K.,
Hodgkin, P.D., Nutt, S.L., and Corcoran, L.M. (2015). Transcriptional profiling of mouse B cell
terminal differentiation defines a signature for antibody-secreting plasma cells. Nat Immunol
16, 663-673.

55.   Kim, U., Qin, X.F., Gong, S., Stevens, S., Luo, Y., Nussensweig, M., and Roeder, R.G. (1996).
The B-cell-specific transcription coactivator OCA-B/OBF-1/Bob-1 is essential for normal
production of immunoglobulin isotypes. Nature *383*, 542-547.

56.   Schubart, D.B., Rolink, A., Kosco-Vilbois, R.G., Botteri, F., and Matthias, P. (1996). B-cell-
specific coactivator OBF-1/OCA-B/Bob1 required for immune response and germinal centre
formation. Nature *383*, 538-542.

57.   Chee, J., Ko, H.J., Skowera, A., Jhala, G., Catterall, T., Graham, K.L., Sutherland, R.M.,
Thomas, H.E., Lew, A.M., Peakman, M., et al. (2014). Effector-memory T cells develop in islets
and report islet pathology in type 1 diabetes. J Immunol *192*, 572-580.

58.   Gearty, S.V., Dundar, F., Zumbo, P., Espinosa-Carrasco, G., Shakiba, M., Sanchez-Rivera,
F.J., Socci, N.D., Trivedi, P., Lowe, S.W., Lauer, P., et al. (2022). An autoimmune stem-like
CD8 T cell population drives type 1 diabetes. Nature *602*, 156-161. 10.1038/s41586-021-
04248-x.

59.   Li, Y., Ramirez-Suastegui, C., Harris, R., Castaneda-Castro, F.E., Ascui, G., Perez-Jeldres, T.,
Diaz, A., Morong, C., Giles, D.A., Chai, J., et al. (2024). Stem-like T cells are associated with
the pathogenesis of ulcerative colitis in humans. Nat Immunol *25*, 1231-1244.
10.1038/s41590-024-01860-7.

60.   Hughes, E.P., Syage, A.R., and Tantin, D. (2024). Durable CD4+ immunity: cherchez la stem.
Trends Imm *45*, 158-166. 10.1016/j.it.2024.01.004.

61.   Polo, J.M., Dell'Oso, T., Ranuncolo, S.M., Cerchietti, L., Beck, D., Da Silva, G.F., Prive, G.G.,
Licht, J.D., and Melnick, A. (2004). Specific peptide interference reveals BCL6 transcriptional
and oncogenic mechanisms in B-cell lymphoma cells. Nat Med *10*, 1329-1335.

62.   Caron, N.J., Quenneville, S.P., and Tremblay, J.P. (2004). Endosome disruption enhances the
functional nuclear delivery of Tat-fusion proteins. Biochem Biophys Res Commun *319*, 12-20.

63.   Krautwald, S., Ziegler, E., Tiede, K., Pust, R., and Kunzendorf, U. (2004). Transduction of the
TAT-FLIP fusion protein results in transient resistance to Fas-induced apoptosis in vivo. J Biol
Chem *279*, 44005-44011.

64.   Wright, H.M., Clish, C.B., Mikami, T., Hauser, S., Yanagi, K., Hiramatsu, R., Serhan, C.N., and
Spiegelman, B.M. (2000). A synthetic antagonist for the peroxisome proliferator-activated
receptor gamma inhibits adipocyte differentiation. J Biol Chem *275*, 1873-1877.

65.   Vassilev, L.T., Vu, B.T., Graves, B., Carvajal, D., Podlaski, F., Filipovic, Z., Kong, N., Kammlott,
U., Lukacs, C., Klein, C., et al. (2004). In vivo activation of the p53 pathway by small-molecule
antagonists of MDM2. Science *303*, 844-848. 10.1126/science.1092472.

66. Cerchietti, L.C., Yang, S.N., Shaknovich, R., Hatzi, K., Polo, J.M., Chadburn, A., Dowdy, S.F.,
and Melnick, A. (2009). A peptomimetic inhibitor of BCL6 with potent antilymphoma effects in
vitro and in vivo. Blood *113*, 3397-3405. 10.1182/blood-2008-07-168773.
67. Green, D.R. (2016). A BH3 Mimetic for Killing Cancer Cells. Cell *165*, 1560.
68. Skwarczynska, M., and Ottmann, C. (2015). Protein-protein interactions as drug targets.
Future Med Chem *7*, 2195-2219.
69. Lu, G., Middleton, R.E., Sun, H., Naniong, M., Ott, C.J., Mitsiades, C.S., Wong, K.K., Bradner,
J.E., and Kaelin, W.G., Jr. (2014). The myeloma drug lenalidomide promotes the cereblon-
dependent destruction of Ikaros proteins. Science *343*, 305-309. 10.1126/science.1244917.
70. Fischer, E.S., Böhm, K., Lydeard, J.R., Yang, H., Stadler, M.B., Cavadini, S., Nagel, J.,
Serluca, F., Acker, V., Lingaraju, G.M., et al. (2014). Structure of the DDB1-CRBN E3 ubiquitin
ligase in complex with thalidomide. Nature *512*, 49-53. 10.1038/nature13527.
71. Fiskus, W., Mill, C.P., Nabet, B., Perera, D., Birdwell, C., Manshouri, T., Lara, B., Kadia, T.M.,
DiNardo, C., Takahashi, K., et al. (2021). Superior efficacy of co-targeting GFI1/KDM1A and
BRD4 against AML and post-MPN secondary AML cells. Blood Cancer J *11*, 98.
10.1038/s41408-021-00487-3.
72. Issa, G.C., Aldoss, I., DiPersio, J., Cuglievan, B., Stone, R., Arellano, M., Thirman, M.J., Patel,
M.R., Dickens, D.S., Shenoy, S., et al. (2023). The menin inhibitor revumenib in KMT2A-
rearranged or NPM1-mutant leukaemia. Nature *615*, 920-924. 10.1038/s41586-023-05812-3.
73. Antony-Debré, I., Paul, A., Leite, J., Mitchell, K., Kim, H.M., Carvajal, L.A., Todorova, T.I.,
Huang, K., Kumar, A., Farahat, A.A., et al. (2017). Pharmacological inhibition of the
transcription factor PU.1 in leukemia. J Clin Invest *127*, 4297-4313. 10.1172/jci92504.
74. Taylor, S.J., Stauber, J., Bohorquez, O., Tatsumi, G., Kumari, R., Chakraborty, J., Bartholdy,
B.A., Schwenger, E., Sundaravel, S., Farahat, A.A., et al. (2024). Pharmacological restriction
of genomic binding sites redirects PU.1 pioneer transcription factor activity. Nat Genet *56*,
2213-2227. 10.1038/s41588-024-01911-7.
75. Gibson, S.A., Yang, W., Yan, Z., Qin, H., and Benveniste, E.N. (2018). CK2 Controls Th17 and
Regulatory T Cell Differentiation Through Inhibition of FoxO1. J Immunol *201*, 383-392.
10.4049/jimmunol.1701592.
76. Wang, F., Demir, S., Gehringer, F., Osswald, C.D., Seyfried, F., Enzenmüller, S., Eckhoff, S.M.,
Maier, T., Holzmann, K., Debatin, K.M., et al. (2018). Tight regulation of FOXO1 is essential for
maintenance of B-cell precursor acute lymphoblastic leukemia. Blood *131*, 2929-2942.
10.1182/blood-2017-10-813576.
77. Klase, Z., Yedavalli, V.S., Houzet, L., Perkins, M., Maldarelli, F., Brenchley, J., Strebel, K., Liu,
P., and Jeang, K.T. (2014). Activation of HIV-1 from latent infection via synergy of RUNX1
inhibitor Ro5-3335 and SAHA. PLoS Pathog *10*, e1003997. 10.1371/journal.ppat.1003997.
78. Zhu, J., Singh, M., Selivanova, G., and Peuget, S. (2020). Pifithrin-α alters p53 post-
translational modifications pattern and differentially inhibits p53 target genes. Sci Rep *10*,
1049. 10.1038/s41598-020-58051-1.
79. Struntz, N.B., Chen, A., Deutzmann, A., Wilson, R.M., Stefan, E., Evans, H.L., Ramirez, M.A.,
Liang, T., Caballero, F., Wildschut, M.H.E., et al. (2019). Stabilization of the Max Homodimer
with a Small Molecule Attenuates Myc-Driven Transcription. Cell Chem Biol *26*, 711-723.e714.
10.1016/j.chembiol.2019.02.009.
80. Frangoul, H., Altshuler, D., Cappellini, M.D., Chen, Y.S., Domm, J., Eustace, B.K., Foell, J., de
la Fuente, J., Grupp, S., Handgretinger, R., et al. (2021). CRISPR-Cas9 Gene Editing for
Sickle Cell Disease and β-Thalassemia. N Engl J Med *384*, 252-260.
10.1056/NEJMoa2031054.
81. Jonasch, E., Donskov, F., Iliopoulos, O., Rathmell, W.K., Narayan, V.K., Maughan, B.L.,
Oudard, S., Else, T., Maranchie, J.K., Welsh, S.J., et al. (2021). Belzutifan for Renal Cell
Carcinoma in von Hippel-Lindau Disease. N Engl J Med *385*, 2036-2046.
10.1056/NEJMoa2103425.
82. Bushweller, J.H. (2019). Targeting transcription factors in cancer - from undruggable to reality.
Nat Rev Cancer *19*, 611-624. 10.1038/s41568-019-0196-7.

83.   Bekes, M., Langley, D.R., and Crews, C.M. (2022). PROTAC targeted protein degraders: the
902   past is prologue. Nat Rev Drug Discov *21*, 181-200. 10.1038/s41573-021-00371-6.
903   84.   Jing, Y., Kong, Y., Allard, D., Liu, B., Kolawole, E., Sprouse, M., Evavold, B., Bettini, M., and
904   Bettini, M. (2023). Increased TCR signaling in regulatory T cells is disengaged from TCR
905   affinity. bioRxiv. 10.1101/2023.01.17.523999.
906   85.   Liu, Y., Chai, Y., Kumar, A., Tidwell, R.R., Boykin, D.W., and Wilson, W.D. (2012). Designed
907   compounds for recognition of 10 base pairs of DNA with two at binding sites. J Am Chem Soc
908   *134*, 5290-5299. 10.1021/ja211628j.
909   86.   Tantin, D., Gemberling, M., Callister, C., and Fairbrother, W. (2008). High-throughput
910   Biochemical Analysis of in-vivo Location Data Reveals Novel Distinct Classes of
911   POU5F1(Oct4)/DNA complexes. Genome Res *18*, 631-639.
912   87.   Wang, V.E., Schmidt, T., Chen, J., Sharp, P.A., and Tantin, D. (2004). Embryonic Lethality,
913   Decreased Erythopoiesis, and Defective Octamer-Dependent Promoter Activation in Oct-1-
914   Deficient Mice. Mol Cell Biol *24*, 1022-1032.
915   88.   Magnusson, A.O., Szekrenyi, A., Joosten, H.J., Finnigan, J., Charnock, S., and Fessner, W.D.
916   (2019). nanoDSF as screening tool for enzyme libraries and biotechnology development.
917   FEBS J *286*, 184-204. 10.1111/febs.14696.

A

B

40 kDa — OCA-B
~15 kDa — H3

Thymocytes 293T SupT1 Jurkat
1   2   3   4

C

5×Oct

OCA-B OCA-B OCA-B OCA-B OCA-B
Oct1   Oct1   Oct1   Oct1   Oct1
CMV   Nano-Luc

5×Mut

Oct1
CMV   Nano-Luc

D

SupT1, 18 hr

Luminescence (×10⁵)

EV   5×Oct   5×Mut

E

5×Oct

linear range

Luminescence / cell

Time (hours)

F

5×Mut   5×Oct

Luminescence / cell

IC₅₀=26.11µM

[DB1033], µM

****

G

H₂N   NH₂
H₂N   NH₂
DB1033

H

5'-G¹ C² A³ T⁴ G⁵ C⁶ A⁷ A⁸ A⁹ T¹⁰ C¹¹ G¹²
3'-C²⁴G²³T²²A²¹C²⁰G¹⁹T¹⁸T¹⁷T¹⁶A¹⁵G¹⁴C¹³

I

2.9 Å N5 N3 N4 C20 2.7 Å
3.0 Å 3.2 Å 2.8 Å T17 T18 2.9 Å 3.0 Å 2.9 Å
C11 T10 N7 3.2 Å 2.7 Å
A7 C6 3.2 Å