Title: Canonical decision computations underlie behavioral and neural signatures of cooperation in primates

Short title: Canonical decision computations underlie cooperation in primates

Authors:

Weikang Shi[1,2,3], Olivia C. Meisner[1,2], Monika P. Jadi[1,3,4,*], Steve W. C. Chang[1,2,3,5,*], Anirvan S. Nandy[1,2,3,5,*]

Affiliations:

1 Department of Neuroscience, Yale University School of Medicine, New Haven, CT 06510, USA

2 Department of Psychology, Yale University, New Haven, CT 06510, USA

3 Wu Tsai Institute, Yale University, New Haven, CT 06510, USA

4 Department of Psychiatry, Yale University School of Medicine, New Haven, CT 06510, USA

5 Kavli Institute for Neuroscience, Yale University School of Medicine, New Haven, CT 06510, USA

* co-senior authors

Figure Information: 4 main figures, 4 supplemental figures, 1 supplemental movie

Correspondences:

Anirvan S. Nandy / Steve W. C. Chang / Monika P. Jadi

Yale University

P.O. Box 208047

New Haven, CT 06520

Email: anirvan.nandy@yale.edu / steve.chang@yale.edu / monika.jadi@yale.edu

Tel: 1-314-307-0498 / 1-203-785-3269

Acknowledgments: This work was supported by the 'Biological Sciences Training Program (BSTP)' training grant (T32 MH014276 to W.S.), the National Institute of Mental Health (R21 MH126072, RF1 MH138396 to S.W.C.C., A.S.N., M.P.J.), the Simons Foundation Autism Research Initiative (SFARI 875855 to S.W.C.C., A.S.N., M.P.J.), Wu Tsai Institute at Yale University (to S.W.C.C., A.S.N., M.P.J., W.S.), a National Eye Institute core grant for vision research (P30 EY026878 to Yale University) and the National Science Foundation Graduate Research Fellowship (DGE2139841 to O.C.M.). We thank the veterinary and husbandry staff at Yale for their excellent animal care, and Anthony DeSimone from the Yale Machine Shop for advice on the neural recording system design.

Author contributions:

Conceptualization: WS, MPJ, SWCC, ASN; Data Collection: WS, OCM; Data Analysis: WS; Computational Modeling: WS; Supervision: MPJ, SWCC, ASN; Writing: WS, MPJ, SWCC, ASN

Conflict of Interest: None

Abstract

Successful cooperation requires dynamic integration of social cues. However, the neural mechanisms supporting this complex process remain unknown. Here, we reveal that the primate dorsomedial prefrontal cortex (dmPFC) implements a gaze-dependent social evidence accumulation process to guide cooperative decisions in freely moving marmoset dyads. A drift-diffusion process in which the partner's action variability is accumulated by social gaze best explains the cooperative actions of the actor. Single-neuron recordings in dmPFC revealed a direct neural correlate: the slope of predictive ramping activity mapped directly onto the rate of evidence accumulation, while baseline firing, modulated by prior outcomes, mapped onto the initial bias. At the population level, the geometry of dmPFC neural trajectories reflected the strength of social evidence and was linked to cooperative success. Together, these findings establish a multi-level neural mechanism for transforming active sensing into a decision variable, linking a canonical computation to cooperative behavior in a naturalistic setting.

Introduction

Cooperation is a defining feature of advanced social behavior. To cooperate successfully, individuals must monitor the actions of their partners, infer intentions, and decide whether to act in a coordinated manner (*1*). A central challenge lies in transforming social cues—particularly through gaze, which reveals attention and intention (*2, 3*)—into decision variables that shape action selection.

Evidence accumulation frameworks provide a powerful explanation for how the brain integrates information to form a decision (*4, 5*). Such mechanisms are well-established in domains such as value-based choice (*6-8*) and, most notably, perceptual decision-making (*5, 9*). In perceptual tasks, neural correlates of evidence accumulation have been identified in regions like the dorsolateral prefrontal cortex (dlPFC) (*10*) and the lateral intraparietal area (LIP) (*11*), where the neural activity encodes the strength of incoming sensory evidence. However, two fundamental challenges remain. First, it is unclear whether evidence accumulation is a core, conserved computational mechanism that extends to the domain of complex social cognition. Second, because prior studies have relied on highly constrained, head-restrained task structures, it is unknown if these accumulation processes operate during naturalistic, unconstrained social interactions.

To address these challenges, we designed a cooperation task where freely moving pairs of marmosets coordinate their actions for mutual reward (*12*), allowing us to investigate the process of social decision-making in a naturalistic setting. While recent work has identified the use of gaze-dependent strategies during these cooperative interactions (*13*), the precise manner in which gaze is used to integrate available social information and the neural mechanisms underlying this integration process remain unknown. We hypothesized that during these naturalistic social interactions, the animals self-organize their behavior into discrete, self-paced episodes (henceforth "trials"), each consisting of a period of gaze-dependent social evidence accumulation culminating in a cooperative decision. Conceptually, we defined each trial as a complete behavioral epoch, beginning with the initiation of movement leading to a pull action and ending after the outcome of that action was delivered and evaluated. To investigate the neural underpinnings of this process, we recorded from the dorsomedial prefrontal cortex (dmPFC), a key node in the social brain network (*14-16*). Given the dmPFC's involvement in encoding others' actions and outcomes (*17*), we hypothesized that it plays a key role in the accumulation of social evidence. By combining behavioral modeling with single-neuron and population-level analyses, we provide the first evidence demonstrating that the dmPFC implements a social gaze-dependent evidence accumulation process to transform social information into decision variables guiding cooperation **(Fig. 1A)**.

Results

A quantitative framework for studying naturalistic cooperation

To quantify the animals' behavior in the mutual cooperation task, we first used markerless facial tracking to extract eight continuous behavior variables for each marmoset (*18, 19*), capturing gaze direction, face position, and movement dynamics (**Fig. 1B; Supp. Mov. S1**). Analysis of the intervals between consecutive pulls by each animal revealed a rhythmic pattern, supporting the characterization of the interactions as a series of self-paced trials (**Fig. 1C**). For both animals from which we recorded the neural activity (hereafter, the 'actors'), the majority of these inter-pull intervals were longer than 4 seconds, a timescale that informed the functional window for our subsequent analyses. Within each trial, we defined social gaze as the actor looking at the partner's face, and social gaze accumulation as the area under the curve of the social gaze probability in this functional window preceding a pull action (**Fig. 1D**).

To objectively define the start of each trial, we sought the single most powerful and unique behavioral predictor of a future pull. Using a Generalized Linear Model (GLM), we found that the linear speed (LS) of the actor's face was a consistently high-importance predictor in both animals (**Supp. Fig. S1A**). Furthermore, a Variance Inflation Factor (VIF) analysis confirmed that LS had low collinearity with the other variables, indicating it provided unique predictive information (**Supp. Fig. S1B**). Based on this evidence, we used the ramping onset of face linear speed to define the start of each trial, and operationalized the duration from this starting point to the pull action as the response time (RT) for that trial (**Fig. 1E**).

With this quantitative framework in place, we hypothesized that the actor animal utilizes a holistic representation of the partner's actions that would collectively allow it to infer the partner's cooperative intention. To define such a representation, we performed Principal Component Analysis (PCA) on the partner's eight behavioral variables. The first principal component (PC1) captured a large fraction of the behavioral variance and thus served as a summary of the partner's actions (**Fig. 1F**). This pipeline thus allowed us to deconstruct continuous behavior into a structured, trial-based format suitable for studying the social evidence accumulation process.

Gaze-dependent drift-diffusion model explains cooperative decisions

To formalize the decision-making process underlying cooperation, we fit the animals' response times and choices using a drift-diffusion model (DDM) (*20*), which decomposes behavior into several key parameters: drift rate (v, the speed of accumulation), the starting point (z, the initial bias), and non-decision time (t_{ND}) (**Fig. 1G; see Methods**). Critically, we formulated four competing models to explicitly test how social gaze mediates the integration of partner information (**Supp. Fig. S2A**). Model comparison using the Deviance Information Criterion (DIC) revealed that a model which allows partner evidence to be integrated only during period of social gaze (M1; **Fig. 1H**) provided the best fit compared to alternate models assuming either no gaze modulation (M2), or an additive (M3), or multiplicative (M4) effect of gaze accumulation (**Supp. Fig. S2B**). This result supports a gaze-dependent integration mechanism, where information about the partner's actions is specifically sampled during periods of social attention.

Analysis of the coefficients from the best fitting model (M1) provided crucial insight into the nature of the social evidence being accumulated. We found that the coefficient for the variability of the partner's actions (β_4) was consistently and significantly negative in both animals (**Fig. 1H**).

This indicates that the variability of the partner's action (PC1) was the major social information
being accumulated, where a noisier or more ambiguous action (higher standard deviation) leads to
a smaller drift rate, slowing the decision process. A control model using a single partner variable
(partner's linear speed, LS), instead of the holistic PC1, did not perform better and its key
coefficients were not significant, demonstrating that the decision process relies on a rich, holistic
representation of the partner's actions (**Supp. Fig. S2C-D**). We therefore defined social evidence
as a measure of variability in partner's action (PC1), with lower variability (less noise)
corresponding to stronger evidence (**Fig. 1I**; **see Methods**).

Figure 1. Conceptual hypothesis, behavioral metrics, and drift diffusion model. (**A**) Conceptual hypothesis. In
the mutual cooperation (MC) task, marmoset dyads are required to pull their levers within a 1-second time window
to receive a juice reward for both. We hypothesize that the social gaze of the actor animal collects social evidence
about a partner's cooperative intention, which is accumulated by the dorsomedial prefrontal cortex to guide action
decisions. (**B**) Top: Definition of eight continuous behavioral variables. Bottom: Example traces for one animal in
one session. (**C**) Distribution of inter-pull intervals. (**D**) Definition of the social gaze accumulation as the area under
curve (AUC) of the social gaze probability. Dashed rectangle shows 4 s functional window of analysis. Pink shaded
areas highlight social gaze duration (social gaze probability> 0). (**E**) Definition of the trial start and response time
(RT). (**F**) Definition of the holistic representation of the partner's actions via principal component analysis (PCA).
(**G**) Structure of the drift diffusion model. (**H**) Coefficient estimates from the main model (M1) showing that partner
action (PC1) contributes only when social gaze is present. (**I**) Definition of the social evidence. Wilcoxon signed-
rank test for panel H (* p < 0.05).

dmPFC neurons show predictive, mixed selectivity for social variables

To investigate the neural basis of this social gaze-dependent process, we recorded single-unit
activity from the dmPFC of the actor marmosets using a custom-built, wireless recording system
(**Fig. 2A; Supp. Mov. S1**). We implemented a GLM (**Fig. 2B; see Methods**) to formally
characterize the diversity of observed neural responses (**Supp. Fig. S3A**). Many dmPFC neurons
exhibited mixed selectivity, including a core subpopulation of "triple-encoding" neurons
simultaneously represented all three key task variables: the actor's pull, the actor's gaze, and the
partner's actions (**Fig. 2C**).

Figure 2. dmPFC neurons show mixed selectivity. dmPFC neurons integrate self, partner, pull and gaze variables,
and encode both predictive and reactive components of the actions. (**A**) Wireless recording system design: (**A1**)
illustration of the headcap design; (**A2**) example video frame highlighting the wireless recording apparatus on one
of the animals during task performance; (**A3**) estimation of the dmPFC recording sites; (**A4**) example of action

potentials (black ticks) from the 64-channel probe aligned at a pull action. (**B**) GLM pipeline for defining tuning
properties and encoding types. (**C**) Distribution of selectivity across neurons reveals mixed coding for pull, gaze,
and partner action variables, supporting flexible social computations. (**D**) Distribution of encoding categories for
pull and gaze neurons. R, reactive; P, predictive. (**E**) Subset of "triple-encoding" neurons' coding categories for pull
and gaze actions. (**F**) Predictive encoding is enriched among triple-encoding neurons. Chi-squared test for panel C
and Fisher's exact test for panel F (* p < 0.05, ** p <0.01, *** p< 0.001).

We next investigated the temporal dynamics of this neural encoding, refining our GLM to classify
tuning as either "predictive" (neural spikes preceding behaviors) or "reactive" (neural spikes
reacting to behaviors) (**Fig. 2B; see Methods**). We found distinct subpopulations with purely
predictive or reactive profiles, as well as many that exhibited both (**Fig. 2D-E; Supp. Fig. S3B**).
Critically, we reasoned that the triple-encoding neurons, encoding the most comprehensive
information, would be primarily involved in integrating social information, thereby leading to the
actor's own actions. A statistical comparison confirmed this intuition: the triple-encoding
subpopulation was significantly enriched for predictive encoding of pull and gaze actions relative
to the broader population of task-responsive neurons (**Fig. 2F**). Together, these findings show that
dmPFC neurons integrate social information and self-actions, with the most informative signals
emerging before cooperative actions rather than in response to them.

Figure 3. Neural firing rate slopes encode both response time and social evidence. (**A**) Mean normalized firing
rate of all dmPFC neurons aligned at the pull action and separating into three quantiles of the RTs (left), and
correlations between dmPFC firing rate slopes and RT (right). (**B**) Mean normalized firing rate of dmPFC neurons
aligned at the trial start and separating into three quantiles of the social evidence (left), and correlations between
dmPFC firing rate slopes and RT social evidence (right). (**C**) Significant proportions of neurons encode both factors
simultaneously. (**D**) Neuron numbers with and without significant correlations between firing rate slopes and social
evidence in successful vs failed pulls. (**E**) Neural DDM linking firing rate slope to drift rate (v) and baseline firing

195   to starting point (z). (**F**) The difference in the mean baseline firing rates following successful pulls vs. failed pulls.
196   (**G**) Behavioral DDM model incorporating outcome-dependent starting points. Wilcoxon signed-rank test for panels
197   A, B, E, F and G, Chi-squared test for panels C and E, and McNemar test for panel D (* $p < 0.05$, ** $p < 0.01$, ***
198     $p < 0.001$). For the red datapoints, Pearson's correlation test for panels A and B, 95% Credible Interval (CI) of the
199   posterior distributions for panel E, and Mann-Whitney U Test for panel F.

Neural ramping activity reflects evidence accumulation

Having established that dmPFC neurons show predictive, mixed selectivity, we next asked whether
their activity exhibits the dynamic signatures of the evidence accumulation process itself. A key
feature of a neural accumulator is ramping activity, wherein the slope is inversely related to RT.
We found precisely this relationship in the dmPFC population: the average slope of the pre-pull
firing rate was steeper for faster responses and shallower for slower ones (**Fig. 3A**). Critically,
these firing rate slopes scaled with the strength of social evidence (**Fig. 3B**). A significant
subpopulation of neurons conjunctively encoded both RT and social evidence in their firing rate
slopes (**Fig. 3C**). Crucially, this encoding of social evidence by differential ramping was strongly
linked to behavioral outcome: this relationship was robust on successful trials but significantly
weaker on failed trials (**Fig. 3D**). We further found a strong correspondence between this dynamic
slope-based encoding and the static tuning properties of the neurons (**Supp. Fig. S4A**). Moreover,
this slope-encoding population showed a trend of being enriched for predictive tuning, consistent
with an anticipatory, rather than reactive, process (**Supp. Fig. S4B**).

Neural activity maps directly onto computational model parameters

To formally connect the observed neural dynamics to the computational model, we constructed a
neural DDM that incorporated both the firing rate slopes and baseline firing rates as predictors.
The results revealed a direct mapping: trial-by-trial firing rate slopes were a significant positive
predictor of the DDM's drift rate, while baseline firing rates quantified during the non-decision
time predicted the starting point (**Fig. 3E**). This starting point bias was itself influenced by recent
history, in which a failed prior trial was associated with higher baseline firing (**Fig. 3F**). To test
the hypothesis suggested by this neural result—that post-error changes in baseline activity have a
direct behavioral consequence—we returned to our best behavioral model (M1) and added the
previous trial's outcome (1 for success pulls and 0 for failed pull) as a predictor of the starting
point. This updated model (M1+) confirmed the hypothesis, revealing a significant negative
coefficient for this relationship (**Fig. 3G**). This finding provides a direct behavioral correlate for
the neural activity, showing that the outcome of the prior trial systematically biased the starting
point of the current decision process, potentially reflecting a post-error increase in motivation.
These analyses provide a direct bridge between the firing rate and computational levels,
demonstrating that the dynamic and static components of dmPFC activity map onto the core
parameters of the evidence accumulation process.

Population dynamics link neural processing to cooperative success

235   While single-neuron activity correlates with key decision variables, cognitive functions are
236   ultimately implemented by the coordinated, dynamic activity of large neural populations.
237   Therefore, to understand how the dmPFC as a whole processes social evidence, we examined the

collective neural activity by analyzing population trajectories at the level of single trials (**Fig. 4A**). We found that the geometry of these neural population trajectories, specifically their path variability, defined as the local tortuosity of the trajectory (*21*), was significantly negatively correlated with the level of social evidence (**Fig. 4B,C**). This effect was most prominent early in the trial during the initial phase of the decision process (**Fig. 4B,E**). Crucially, this relationship between the geometry of the neural population and the strength of social evidence was also tightly linked to behavioral outcome. At the population level, the strong relationship between path variability and social evidence was present only on successful trials and was absent on failed trials (**Fig. 4D**). This finding at the population level mirrored our observations with single neurons, where the encoding of social evidence by firing rate slopes was robust on successful trials and significantly weaker on failed trials (**Fig. 3D**). Together, these results demonstrate that the effective neural processing of social cues in dmPFC is critical for guiding successful cooperation.

Figure 4. Population dynamics reflect social gaze accumulation. (**A**) Definition of path variability (tortuosity) of the single-trial neural population trajectory, example session. (**B**) Heatmaps of the temporal structure of the path variability showing that social evidence shapes dmPFC population dynamics during social decisions, averaged across sessions. (**C**) Correlation between the strength of social evidence and path variability of the neural trajectory across sessions. (**D**) Correlation between social evidence and path variability of the neural trajectory in successful and failed pulls. (**E**) Average change of path variability over time in individual trials. Wilcoxon signed-rank test for

Discussion

Our study provides a multi-level account of how the primate dmPFC orchestrates cooperation in a
naturalistic behavioral setting. We show that the dmPFC implements a social gaze-dependent
evidence accumulation process, transforming actively sampled social cues into a decision variable.
By identifying this canonical computation in freely moving animals without any imposed task
structure, our work bridges the gap between classic decision neuroscience and the complexity of
real-world social interactions.

Our findings connect behavior, computation, and neural activity across multiple scales. A gaze-
dependent DDM best explained cooperative decisions, establishing that evidence about a partner's
actions is integrated only when actively sampled. We then identified a direct neural correlate for
this process. At the single-neuron level, the predictive ramping of dmPFC neurons encoded both
the response time and the strength of social evidence. These neural dynamics mapped directly onto
the DDM's core parameters, with firing rate slopes predicting the drift rate and baseline activity—
updated by recent trial outcomes—predicting the starting point bias. Finally, this mechanism was
reflected at the population level, where the geometry of neural trajectories tracked the level of
social evidence. Crucially, this signature was present mostly on successful trials, cementing the
link between the neural mechanism and the behavioral outcome.

Evidence accumulation has long been established as a canonical computation for perceptual
decisions (5), with clear neural signatures in sensorimotor areas such as LIP (11) and dlPFC (10).
Our findings extend this framework into the social domain, identifying the dmPFC as a key locus
for this more complex form of accumulation. Unlike perceptual tasks, where evidence is often
externally driven, social evidence about another's intentions is uncertain and must be actively
sampled via gaze (2). The dmPFC is uniquely suited for this computation (22), integrating
information about the partner's actions with the actor's own social attention.

A significant advance of our study is the application of a detailed, mechanistic approach to neural
function in freely moving, socially interacting primates. While the field has begun to embrace the
study of more naturalistic behaviors (23), as exemplified by recent work in rodents (24), tree
shrews (25) and macaques (26), research on the detailed computational mechanism has been
lacking, facing both technical hurdles in recording from unconstrained animals and analytical
challenges in interpreting complex, continuous behavior. We addressed the technical challenges
by developing a novel modular head chamber (27), which improved surgical success rates. We
also used movable probes, in contrast to fixed chronic implants (28), to sample diverse neural
populations across sessions. To overcome the analytical challenges, we employed a sophisticated
computational framework, including the GLM pipelines (29-32), to deconstruct the continuous
behavioral stream into meaningful, quantifiable variables. By identifying computational signatures
of decision-making in this unconstrained paradigm, we show that even without the rigid structure
of a classic task, marmosets spontaneously organize their behavior in a way that is amenable to an
evidence accumulation process. This demonstrates the robustness of this neural algorithm,
suggesting it is a fundamental process that is flexibly deployed to make decisions in dynamic
environments.

Our results also highlight several avenues for future research. The influence of the previous trial's
outcome on the DDM starting point, mediated by baseline dmPFC firing, suggests a rapid,

experience-dependent plasticity that could be interpreted as an alerting signal or a shift in
motivation following failure (*33-35*). Future studies employing neural manipulation could also
directly test the causal role of this mechanism; for instance, microstimulation delivered during the
gaze accumulation period could potentially disrupt the evidence integration process and alter the
cooperative decision (*36*).

In conclusion, by linking behavior, modeling, and neural dynamics across multiple scales, we
established a comprehensive framework for how the brain transforms active sensing into a decision
variable to guide cooperation. This neurocomputational framework provides a mechanistic
grounding for economic models of strategic interaction (*37*), a potential biomarker for social
deficits in psychiatry (*38*), and offers new biological insights for designing the next generation of
cooperative multi-agent artificial systems (*39*).

Materials and Methods

Animal subjects

Two common marmosets (*Callithrix jacchus*) participated in the study: a 7-year-old female (Animal K) and a 6.5-year-old male (Animal D). Animal K participated in 21 sessions with four different partners (2 females and 2 males), and Animal D participated in 29 sessions with three different partners (3 females). A detailed comparison of behavior and neural activity across different partners is beyond the scope of this study. However, to confirm that our findings were not confounded by partner identity, we performed alternative analyses using linear mixed-effects models, which included partner identity as a random effect. These confirmatory models did not change the statistical conclusions of the primary analyses reported throughout the manuscript. The animals were pair-housed, maintained on a 12-hour light-dark cycle, and food and water were removed from their home cages for 1–3 hours prior to testing. All experimental procedures were approved by the Yale Institutional Animal Care and Use Committee and complied with the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Experimental task

The experimental apparatus and training procedures were identical to those described in previous studies (*12, 13*). The animals had been previously trained to perform a mutual cooperation (MC) task. Both marmosets in the dyads were required to pull their respective levers within a 1-second time window to successfully receive a reward. A pull was registered when the lever passed a predefined position threshold, and its timestamp was recorded by the task-control computer at a 10 kHz sampling rate. A successful, coordinated trial resulted in the delivery of a 0.2 mL liquid reward (marshmallow fluff solution) to both animals, which was immediately preceded by an auditory tone (2 KHz) to signal success. A failed trial occurred if the pulls were not synchronized within the 1-second window. This task was designed to be performed in a naturalistic setting, with no explicitly defined or implemented temporal structure; instead, the animals engaged in a continuous series of self-paced cooperative decisions. The data analyzed in this study were collected after the animals had been fully trained on this task, representing established cooperative performance.

Behavioral Data Acquisition and Processing

Continuous behavior was recorded at 30 frames/sec from three synchronized cameras (GoPro Hero10) positioned at the front of the apparatus. The cameras were controlled via a shared remote control. To align the video data with the event timestamps from the task-control computer, we used two redundant signals, both controlled by the task computer: an audible tone at the start of the session for coarse alignment, and an LED that flashed every 20 seconds for fine alignment, providing a reliable and recurring synchronization signal captured by the cameras. We used the multi-animal version of DeepLabCut 2.0 to perform markerless pose estimation (*19, 40*). Six facial key points were labeled on 500 frames: two ears, two eyes, the central blaze, and the mouth. The DeepLabCut neural network was trained on these labels for 150,000 iterations until the tracking error reached a stable minimum. The 2D coordinates from the three cameras were then used to triangulate the 3D position of each key point using Anipose (*18*). This 3D reconstruction allowed for the precise calculation of each animal's head position and orientation in space throughout the

session. From the 3D facial coordinates, we derived the eight continuous behavioral variables used for all subsequent analyses (**Fig. 1B**):

- ***A1-A3***: Three angular variables measuring the direction of gaze relative to the other animal, the reward tube, and the lever.
- ***D_{AA}, D_{AT}, D_{AL}***: Three distance variables measuring the distance to the other animal, the reward tube, and the lever.
- ***LS***: The face linear speed, calculated from the movement of the head's 3D position.
- ***AS***: The gaze vector angular speed, capturing the rate of change in head orientation.

Similar to earlier work (*13*), head gaze was defined as a 15-degree cone extending perpendicularly from the facial plane formed by the tracked key points. Social gaze was determined to have occurred whenever one animal's gaze cone intersected with the other animal's face.

Implant Design and Surgical Processing

We designed a novel, multi-component head implant to facilitate wireless neural recordings in freely moving marmosets (**Fig. 2A**). The system was custom-designed in CAD software (Shapr3D) to fit each animal's cranium (obtained from pre-operative CT scan) and consists of two main parts. The first is a 3D-printed medical-grade titanium base (Materialise NV) that protects the surgical site where the skull is exposed. An integrated headpost on the base allows for brief periods of stable head-fixation. The second part is a 3D-printed recording chamber (nylon; Protolabs), which consists of two sealed and separated chambers. The probe chamber houses and protects a miniature screw microdrive (nanodrive; Cambridge Neurotechnologies) and the probe, while the electronics chamber protects the headstage chips needed for digitization and signal amplification. A silicone-based sealant (Kwik-Sil; World Precision Instruments, LLC) separates the chambers, allowing the probe to remain sterile while the data logger is connected to the headstage electronics. This modular design is advantageous for animal welfare, as it permits a multi-step surgical approach with shorter individual procedures and simplifies implant cleaning and maintenance.

Enabled by this modular implant design, the surgical implantation followed a multi-step process. First, the titanium base was implanted. In a subsequent procedure, a small opening was made in the skull within the protected base, and a nanodrive holding a 64-channel linear array probe (NeuroNexus) was mounted inside the probe chamber. The nanodrive allowed movement of the probe along the superior-inferior axis, while the mounting apparatus holding the nanodrive allowed repositioning along the anterior-posterior and mediolateral axes. The target cortical area, the dorsomedial prefrontal cortex (dmPFC), was identified using pre-operative CT scans and stereotaxic coordinates to guide the initial placement. The probe was lowered to a site within the dmPFC during the surgery, and the probe track location was confirmed post-operatively by co-registering a CT scan with the probe implanted against a digital marmoset brain atlas (*41*).

Neural Recording and Signal Processing

During each experimental session, which lasted approximately 15 minutes, we used a wireless eCube headstage system (White Matter LLC) to record single-unit activity while animals were freely moving and performing the Mutual Cooperation (MC) task (Animal K: 490 neurons, Animal D: 464 neurons). The data logger was attached at the start and removed at the end of each session

while the animals were briefly head-fixed, allowing for untethered social interaction during the
task. Synchronization between the wireless recordings and the task-control computer was managed
by a DAQ board (National Instruments Corp.) connected to the logger's transceiver. At the start of
each recording, the logger sent a pulse that was captured by the NI board and timestamped by the
task computer for initial alignment. To correct for any temporal drift, the task computer also sent
a continuous 10 kHz signal through the NI board to the logger for auto-correction.
Electrophysiological signals were acquired at a 20 kHz sampling rate. The raw data was processed
offline: action potentials were first automatically detected and clustered using Kilosort4 (*42*), and
the resulting clusters were then manually curated using phy, an open-source Python library, to
isolate well-defined single units for all subsequent analyses.

Following spike-sorting, all subsequent analyses were performed on neural data that was first
downsampled from 20 kHz to 30 Hz to match the camera's frame rate. Continuous firing rates were
estimated by counting the number of spikes in 33.3 ms (1/30 s) bins and then smoothing the
resulting histogram with a $\sigma = 50\ ms$ Gaussian kernel. This continuous firing rate trace was then
z-scored using the mean and standard deviation of that neuron's activity across the entire session
to obtain a normalized firing rate. For the neural DDM, the trial-by-trial slope of firing rate was
calculated by fitting a linear regression model to the normalized (z-scored) firing rate trace over
the trial's duration. For the neural GLM (see below), the model was fit directly to the 30 Hz spike
count train for each neuron.

Computational Modeling of Decision-Making

To prepare the behavioral data for modeling, we first defined trial epochs and key predictor
variables. Continuous behavioral data were segmented into discrete trials centered on pull actions.
The start of each trial was marked by the onset of the actor's face linear speed (LS), with response
time (RT) defined as the duration until the subsequent pull. The justification for this marker is
detailed in the supplementary materials (**Supp. Fig S1**). To create a holistic measure of the
partner's actions, we performed Principal Component Analysis (PCA) on their eight behavioral
variables. These variables were then used to fit a hierarchical Bayesian drift-diffusion model
(DDM) to formalize the evidence accumulation process. The DDM decomposes behavior into
distinct cognitive parameters: the drift rate (v), representing the speed of evidence accumulation;
the starting point (z), representing an initial bias; and the non-decision time (t_{ND}). To explicitly
test our hypothesis about how social gaze mediates the integration of partner information, we
formulated and compared four competing models (**Supp. Fig S2A**):

- **M1 (Gaze-gated model):** A model in which partner information influences the drift rate only
during periods of social gaze. Periods of social gaze were defined as the time points in the
functional window when social gaze probability was greater than zero.

$$v \sim \beta_1 * E(LS_{actor}) + \beta_2 * \sigma(LS_{actor}) + \beta_3 * E(action_{partner}|gaze_{actor}) + \beta_4$$
$$* \sigma(action_{partner}|gaze_{actor})$$

- **M2 (No gaze modulation):** A model where partner information is integrated regardless of
gaze.

$$v \sim \beta_1 * E(LS_{actor}) + \beta_2 * \sigma(LS_{actor}) + \beta_3 * E(action_{partner}) + \beta_4$$
$$* \sigma(action_{partner})$$

- **M3 (Additive gaze effect):** A model where social gaze (measured as social gaze accumulation, $gaze_{auc}$) has a separate, additive effect on the drift rate.

$$v \sim \beta_1 * E(LS_{actor}) + \beta_2 * \sigma(LS_{actor}) + \beta_3 * E(action_{partner}) + \beta_4 * \sigma(action_{partner}) + \beta_5 * gaze_{auc}$$

- **M4 (Multiplicative gaze effect):** A model where social gaze multiplicatively scales the influence of partner information.

$$v \sim \beta_1 * E(LS_{actor}) + \beta_2 * \sigma(LS_{actor}) + \beta_3 * [E(action_{partner}) * gaze_{auc}] + \beta_4 * [\sigma(action_{partner}) * gaze_{auc}]$$

- **M1+ (Gaze-gated model with history effect):** An extension of M1 where the starting point (z) is modulated by the outcome of the previous trial, which is 1 if the previous trial is successful, and 0 if it failed. The drift rate (v) is formulated identically to M1.

$$z \sim \beta_Z *< previous\ trial\ outcome >$$

- **M5 (Gaze-gated model with single behavioral variable):** A control model with the same gaze-gated structure as M1, but using the partner's linear speed ($LS_{partner}$) instead of the holistic action variable (PC1, $action_{partner}$).

$$v \sim \beta_1 * E(LS_{actor}) + \beta_2 * \sigma(LS_{actor}) + \beta_3 * E(LS_{partner}|gaze_{actor}) + \beta_4 * \sigma(LS_{partner}|gaze_{actor})$$

To link the cognitive model to neural activity, we constructed a **neural DDM**. This model used trial-by-trial neural features to predict the DDM parameters. The drift rate (v) was predicted by the slope of a neuron's firing rate, while the starting point (z) was predicted by the neuron's mean baseline firing rate. This baseline activity was calculated during the non-decision time period (t_{ND}), using the duration estimated for each trial from the fitted parameters of model M1. The specific relationships were:

$$v \sim \beta_1 *< slope\ of\ firing\ rate >$$

$$z \sim \beta_2 *< mean\ firing\ rate\ during\ t_{ND} >$$

All DDM models were fit using the HDDM (Hierarchical Drift-Diffusion Model) Python package (*20*), which uses Markov Chain Monte Carlo (MCMC) sampling to estimate posterior distributions of the model parameters. For each model, we generated 2000 samples from the posterior distribution following a burn-in period of 1000 iterations, which were discarded. We confirmed model convergence by visually inspecting the MCMC chains. The final coefficient values reported are the mean of the posterior distributions from the collected samples.

Social Evidence Estimation

We defined the social evidence for a given trial by subtracting the standard deviation of the partner's action (PC1) during periods of social gaze from the maximum value observed across all trials, a linear transformation where lower variability (less noise) corresponds to stronger evidence:

475 $$Social\ evidence = max\langle\sigma(action_{partner}|gaze_{actor})\rangle - \sigma(action_{partner}|gaze_{actor})$$

Statistical Modeling of Behavior and Neural Tuning

Behavioral GLM for Trial Start Identification

To objectively identify the behavioral variable that most reliably predicted the initiation of a pull
action, we used a Generalized Linear Model (GLM) with a logistic link function, fit to the eight
continuous behavioral variables prepared at a 30 Hz sampling rate. The model was fit separately
for each behavioral session. Within each session, the data was split into an 80% training set and a
20% testing set. The model was trained to predict the probability of a pull event from the eight
continuous behavioral variables. To capture temporal dynamics, each variable trace was convolved
with a basis set of 10 temporal kernels spanning the 4-second window preceding each pull action
(-4s to 0s). To determine the importance of each variable, we used a leave-one-out (LOO)
procedure. The variable importance was quantified as the drop in the Area Under the Receiver
Operating Characteristic curve (AUROC), calculated on the 20% testing set, when that variable
was removed from the model. This entire procedure was repeated for 1000 bootstrap iterations to
assess the statistical significance of the drop in AUROC for each variable.

Neural GLM for Single-Neuron Tuning Properties

We used a Poisson GLM to characterize the tuning of individual dmPFC neurons by fitting the
model to the 30 Hz downsampled spike count train for each neuron. The model was fit separately
for each neuron using an 80/20 train/test split. The model predicted the number of spikes in each
30 Hz time bin based on four main predictors: the actor's pull action, social gaze, the partner's
action (PC1), and reward delivery. The reward delivery variable was included as a control to ensure
that observed neural activity was not confounded by the motor action of drinking. Each of the four
predictors was convolved with a basis set of 10 temporal kernels spanning an 8-second window
around each event (-4s to +4s), creating a total of 40 weighted features for the full model.

To determine a neuron's overall selectivity, we performed an LOO comparison where all 10 kernels
for a given variable were excluded. A neuron was considered significantly tuned if the AUROC of
the full model on the testing set was significantly better than that of the reduced model.
Furthermore, to distinguish between predictive and reactive activity, we assessed them separately
based on their temporal relationship to the event under consideration. This was achieved by
splitting the 10 temporal kernels into a predictive set (spanning -4s to 0s pre-event) and a reactive
set (spanning 0s to +4s post-event). For instance, to test for predictive tuning, we compared the
full model to a reduced model where only the 5 predictive kernels for a given variable were
excluded. A neuron was classified as having predictive tuning if the full model's AUROC on the
testing set was significantly better than that of this specific reduced model.

Population Dynamics

To examine the collective dynamics of the dmPFC population, we carried out single-trial
population analyses based on existing methods (*21*). This analysis was restricted to sessions with
more than 5 simultaneously recorded neurons (up to 40 neurons per session; 19 sessions in Animal
K, 26 sessions in Animal D). For each of these sessions, the normalized (z-scored) firing rates of
the population were arranged into an activity matrix. We then performed PCA on each session's

data separately and focused on the neural state space defined by the first three principal
components (PCs).

Before calculating geometric measures, each 3D trajectory was preprocessed. First, because trials
had different durations, each trajectory was normalized by being reparameterized by its arc length.
This process resamples the trajectory so that each path consists of a uniform 100 time points,
making them comparable. Second, these reparameterized trajectories were smoothed using a
rolling Gaussian window ($\sigma = 3$) to isolate the large-scale geometry of the path from small-scale
neural jitter.

On these preprocessed trajectories, we calculated the following two local geometric measures (**Fig.**
4A):

• **Path Direction (Curvature, κ)**: Curvature measures how quickly the trajectory deviates from
a straight line. It is formally defined as the magnitude of the rate of change of the curve's unit
tangent vector *T(s)* with respect to arc length *s*.

$$\kappa(s) = \left\| \frac{dT(s)}{ds} \right\|$$

531 • **Path Variability (Tortuosity, τ):** Tortuosity captures the "wiggleness" or complexity of the
532 trajectory. We defined it as the squared ratio of the rate of change of curvature with respect to
533 arc length ($d\kappa/ds$) to the curvature itself.

$$\tau(s) = \left(\frac{d\kappa/ds}{\kappa(s)} \right)^2$$

These continuous measures were estimated numerically from the discrete, preprocessed
trajectories. Derivatives were approximated using a rolling window of 5 time points, and the term
$d\kappa/ds$ was calculated using the gradient of the curvature trace. Both the final curvature and
tortuosity traces were themselves smoothed with a Gaussian kernel ($\sigma = 3$) for final analysis.

Statistical Analysis

All statistical analyses were performed using custom scripts written in Python with the SciPy and
statsmodels libraries. All tests were two-sided, and results were considered significant at $p < 0.05$.
Shaded regions in plots represent the standard error of the mean (SEM).

• **Comparison of Proportions:** A Fisher's exact test was used to assess the enrichment of
predictive tuning in triple-encoding neurons (Fig. 2F). A Chi-squared test was used for other
comparisons of categorical data (Fig. 2C, 3C, 3E).
• **Paired Comparisons:** To compare the presence of significant slope encoding on successful
versus failed trials within the same neuronal population (Fig. 3D), we used the McNemar test
for paired nominal data.
• **One-Sample Tests:** Distributions of coefficients and correlation values were tested for
significance against a median of zero using the non-parametric Wilcoxon signed-rank test (Fig.
1H, 3A-B, 3F-G, 4C-D).
• **Time-Series Analysis:** To assess the significance of the group-average time course of path
variability (Fig. 4E), we used a permutation test with cluster-based correction. The null
hypothesis data was generated by creating one "shuffled" trace for each session. To do this, we

first took the real firing rate trace for each neuron in each trial and performed a circular shuffle. The PCA and path variability calculations were then re-run on this shuffled single-trial data to generate the final shuffled trace for each session. The group of real session traces was then compared to the group of shuffled traces using a paired t-test at each time point, with cluster-based correction to control for multiple comparisons.

- **Confirmatory Analyses:** To ensure that results were not confounded by the use of different partners across sessions, all primary statistical analyses were confirmed using linear mixed-effects models with partner identity included as a random effect. These confirmatory models yielded the same statistical conclusions, and therefore, we kept the original results.

- **Successful and failed pull comparison:** Since the successful and failed pulls had different numbers of occurrences, to match the statistical power, we used all failed pull trials and randomly picked the same number of successful pull trials for each session, and performed pair-wise comparisons.

References

1. J. M. Burkart, C. P. van Schaik, Cognitive consequences of cooperative breeding in primates? *Anim Cogn* **13**, 1-19 (2010).

2. N. J. Emery, The eyes have it: the neuroethology, function and evolution of social gaze. *Neurosci Biobehav R* **24**, 581-604 (2000).

3. A. Frischen, A. P. Bayliss, S. P. Tipper, Gaze cueing of attention: Visual attention, social cognition, and individual differences. *Psychol Bull* **133**, 694-724 (2007).

4. R. Ratcliff, P. L. Smith, S. D. Brown, G. McKoon, Diffusion Decision Model: Current Issues anc History. *Trends in Cognitive Sciences* **20**, 260-281 (2016).

5. J. I. Gold, M. N. Shadlen, The neural basis of decision making. *Annual Review of Neuroscience* **30**, 535-574 (2007).

6. J. R. Busemeyer, S. Gluth, J. Rieskamp, B. Turner, Cognitive and Neural Bases of Multi-Attribute, Multi-Alternative, Value-based Decisions. *Trends in Cognitive Sciences* **23**, 251-263 (2019).

7. I. Krajbich, C. Armel, A. Rangel, Visual fixations and the computation and comparison of value in simple choice. *Nature Neuroscience* **13**, 1292-1298 (2010).

8. I. Krajbich, A. Rangel, Multialternative drift-diffusion model predicts the relationship between visual fixations and choice in value-based decisions. *P Natl Acad Sci USA* **108**, 13852-13857 (2011).

9. T. D. Hanks, C. Summerfield, Perceptual Decision Making in Rodents, Monkeys, and Humans. *Neuron* **93**, 15-31 (2017).

10. J. N. Kim, M. N. Shadlen, Neural correlates of a decision in the dorsolateral prefrontal cortex of the macaque. *Nat Neurosci* **2**, 176-185 (1999).

11. M. N. Shadlen, W. T. Newsome, Neural basis of a perceptual decision in the parietal cortex (area LIP) of the rhesus monkey. *Journal of Neurophysiology* **86**, 1916-1936 (2001).

12. O. C. Meisner *et al.*, Development of a Marmoset Apparatus for Automated Pulling to study cooperative behaviors. *Elife* **13**, (2024).

13. O. C. Meisner *et al.*, Diverse and flexible strategies enable successful cooperation in marmoset dyads. *Curr Biol*, (2025).

14. C. C. Ruff, E. Fehr, The neurobiology of rewards and values in social decision making. *Nature Reviews Neuroscience* **15**, 549-562 (2014).

15. P. Gangopadhyay, M. Chawla, O. Dal Monte, S. W. C. Chang, Prefrontal-amygdala circuits in social decision-making. *Nat Neurosci* **24**, 5-18 (2021).

16. M. Isoda, The Role of the Medial Prefrontal Cortex in Moderating Neural Representations of Self and Other in Primates. *Annual Review of Neuroscience, Vol 44, 2021* **44**, 295-313 (2021).

17. A. Noritake, T. Ninomiya, M. Isoda, Social reward monitoring and valuation in the macaque brain. *Nature Neuroscience* **21**, 1452-+ (2018).

18. P. Karashchuk *et al.*, Anipose: A toolkit for robust markerless 3D pose estimation. *Cell Reports* **36**, (2021).

19. J. Lauer *et al.*, Multi-animal pose estimation, identification and tracking with DeepLabCut. *Nat Methods* **19**, 496-504 (2022).

20.   T. V. Wiecki, I. Sofer, M. J. Frank, HDDM: Hierarchical Bayesian estimation of the Drift-
Diffusion Model in Python. *Front Neuroinform* **7**, (2013).

21.   M. M. Monsalve-Mercado, G. M. Stine, M. N. Shadlen, K. D. Miller, The geometry of the neural
state space of decisions. *bioRxiv*, (2025).

22.   O. Dal Monte *et al.*, Widespread implementations of interactive social gaze neurons in the
primate prefrontal-amygdala networks. *Neuron* **110**, 2183-+ (2022).

23.   S. Q. Fan, O. Dal Monte, S. W. C. Chang, Levels of naturalism in social neuroscience research.
Iscience **24**, (2021).

24.   M. Jiang *et al.*, Neural basis of cooperative behavior in biological and artificial intelligence
systems. *Science*, eadw8151 (2025).

25.   M. Jiang *et al.*, Evolution and neural representation of mammalian cooperative behavior. *Cell Rep*
37, 110029 (2021).

26.   M. Franch *et al.*, Visuo-frontal interactions during social learning in freely moving macaques.
Nature, (2024).

27.   P. Jendritza, F. J. Klein, P. Fries, Multi-area recordings and optogenetics in the awake, behaving
marmoset. *Nat Commun* **14**, 577 (2023).

28.   J. D. Walker *et al.*, Chronic wireless neural population recordings with common marmosets. *Cell
Rep* **36**, 109379 (2021).

29.   A. J. Calhoun, J. W. Pillow, M. Murthy, Unsupervised identification of the internal states that
shape natural behavior. *Nature Neuroscience* **22**, 2040-+ (2019).

30.   P. Coen, M. Xie, J. Clemens, M. Murthy, Sensorimotor Transformations Underlying Variability
in Song Intensity during Courtship. *Neuron* **89**, 629-644 (2016).

31.   P. Coen *et al.*, Dynamic sensory cues shape song structure in. *Nature* **507**, 233-+ (2014).

32.   J. W. Li, M. C. Aoi, C. T. Miller, Representing the dynamics of natural marmoset vocal behaviors
in frontal cortex. *Neuron* **112**, (2024).

33.   B. A. Purcell, R. Kiani, Neural Mechanisms of Post-error Adjustments of Decision Policy in
Parietal Cortex. *Neuron* **89**, 658-671 (2016).

34.   J. I. Gold, C. T. Law, P. Connolly, S. Bennur, The Relative Influences of Priors and Sensory
Evidence on an Oculomotor Decision Variable During Perceptual Learning. *Journal of
Neurophysiology* **100**, 2653-2668 (2008).

35.   O. Odoemene, S. Pisupati, H. Nguyen, A. K. Churchland, Visual Evidence Accumulation Guides
Decision-Making in Unrestrained Mice. *J Neurosci* **38**, 10143-10155 (2018).

36.   S. Q. Fan, O. Dal Monte, A. R. Nair, N. A. Fagan, S. W. C. Chang, Closed-loop
microstimulations of the orbitofrontal cortex during real-life gaze interaction enhance dynamic
social attention. *Neuron* **112**, (2024).

37.   C. Camerer, *Behavioral game theory : experiments in strategic interaction*. The Roundtable
series in behavioral economics (Russell Sage Foundation; Princeton University Press, New York;
Princeton, N.J., 2003), pp. xv, 550 p.

38.   D. P. Kennedy, R. Adolphs, The social brain in psychiatric and neurological disorders. *Trends in
Cognitive Sciences* **16**, 559-572 (2012).

39.   C. X. Zhu, M. Dastani, S. H. Wang, A survey of multi-agent deep reinforcement learning with
communication. *Auton Agent Multi-Ag* **38**, (2024).

40.   A. Mathis *et al.*, DeepLabCut: markerless pose estimation of user-defined body parts with deep
learning. *Nature Neuroscience* **21**, 1281-+ (2018).

41.   A. Woodward *et al.*, The Brain/MINDS 3D digital marmoset brain atlas (vol 5, 180009, 2018).
Sci Data **9**, (2022).

42.   M. Pachitariu, S. Sridhar, J. Pennington, C. Stringer, Spike sorting with Kilosort4. *Nat Methods*
21, (2024).

Supplemental Figures and Movies

Figure S1. Identifying behavioral variables defining trial start; related to Figure 1. Trial start
is best defined by the linear speed (LS) variable. (**A**) Generalized linear model (GLM) pipeline to
identify which behavioral variables predict pull actions. (**B**) Collinearity analysis of continuous
variables, quantified with variance inflation factors (VIF). Wilcoxon signed-rank test for panel B
(* $p < 0.05$)

Figure S2. Control drift diffusion models; related to Figure 1. (**A**) Four models testing effects
of partner actions and gaze (M1: main model; M2-4: control models). (**B**) Model comparison using
Deviance Information Criterion (DIC). (**C**) Comparison of DIC values between the main model
(M1, partner PC1 with gaze filter) and control model (M5, partner linear speed, LS, with gaze
filter). (**D**) Coefficients from M5, showing non-significant effects. Wilcoxon signed-rank test for
panels B-D (ns: no significance, * $p < 0.05$).

677
678 **Figure S3. Heatmaps of dmPFC neural activities; related to Figure 2.** (**A**) Heatmaps of
679 normalized (z-scored) firing rates aligned to pull and gaze actions. (**B**) Heatmaps of normalized
680 (z-scored) firing rates for different encoding types.
681

Figure S4. Predictive encoding in slope neurons; related to Figure 3. Slope-encoding neurons
are enriched for predictive coding of partner actions. (**A**) Slope-encoding neurons show mixed
selectivity for pull, gaze, and partner action variables. (**B**) Proportion of predictive vs reactive
encoding among slope and non-slope neurons for pull and gaze. Animal D showed a significant
enrichment of predictive coding among slope neurons. Chi-squared test for panel A, Fisher's exact
test for panel B (** $p < 0.01$).

Movie S1. Example clip of the animal tracking and the multi-channel wireless recording
form dmPFC.

692