Title: *Corylus avellana* disease management: using metagenomics to illuminate the rhizosphere

microbiome of *Corylus*

Authors: John K. Robinson[1], John Steele[1], Thomas J. Molnar[2], Sharon Regan[1], George C.

diCenzo[1*]

Affiliations: [1] Department of Biology, Queen's University, Kingston, Ontario, Canada

[2] Department of Plant Biology, Rutgers University, Newark, New Jersey, USA

* Corresponding author: George C. diCenzo, george.dicenzo@queensu.ca

Keywords: Hazelnut, Rhizosphere, Metagenomics, Microbiome, Plant-Microbe Interaction

ABSTRACT

The European hazelnut, *Corylus avellana*, is one of the most economically important tree nut crops

globally. The biotrophic ascomycete pathogen *Anisogramma anomala*, found naturally associated

with wild *C. americana*, continues to pose a significant threat to European hazelnut production

across North America. Rhizosphere microorganisms have demonstrated the capacity to impact

diverse plant features and serve as an important resource for the improvement of plant disease

outcomes. Here, metagenomics was used to examine the taxonomic and functional features of the

rhizosphere microbial communities of hazelnut trees differing in their levels of resistance to *A.*

anomala: highly tolerant *Corylus americana*, and resistant and susceptible *Corylus avellana*. No

statistically significant differences in microbial alpha diversity or beta diversity were noted

between the three rhizosphere groups. Compared to bulk soil, all three rhizosphere groups were

enriched for the fungal phylum Basidiomycota and bacterial phylum "*Candidatus*

Rokubacteriota". At the genus level, the bacterial genera *Actinospica*, *Occallatibacter*, and

"*Candidatus* Sulfotelmatobacter" were under-represented, while the genus *Rhizobacter* was over-

represented, in the resistant and susceptible *C. avellana* rhizosphere samples compared to the bulk

soil. A total of 45 dereplicated, high-quality metagenome-assembled genomes were generated,

corresponding to 41 bacteria and 4 archaea. Many of the metagenome-assembled genomes carried

multiple biosynthetic gene clusters, including MAGs corresponding to the genera *Lysobacter* and

Actinospica. Overall, the low differentiation of the rhizosphere microbiomes of the three hazelnut

rhizosphere groups suggest that differences in *A. anomala* disease expression are likely not

associated with differences in the rhizosphere microbiome. Nevertheless, the results shed new light

on the rhizosphere communities of two species of hazelnut, and woody perennials more broadly,

and identify potential avenues for future research into the development of microbial inoculants for

Corylus spp..

INTRODUCTION

Hazelnut cultivation has a rich history and is still of economic importance in contemporary societies. The European hazelnut *Corylus avellana* has been cultivated since Roman times (Boccacci and Botta 2009) and remains one of the world's principal tree nut crops (Helmstetter et al. 2020). Approximately 70% percent of the world's hazelnuts are presently grown in Turkey (Food and Agricultural Organization of the United Nations 2021), but the future of Turkey's hazelnut industry remains uncertain. To stabilize existing hazelnut production and supply growing global demand, the world's hazelnut orchards must expand across suitable growing regions. Historically, commercial production of *C. avellana* in North America outside of the Pacific Northwest was limited due to the devastating impact of the biotrophic ascomycete pathogen *Anisogramma anomala* (Molnar 2011; Revord et al. 2020; Thompson et al. 1996). Unlike the native but not-commercially amenable American hazelnut species, *Corylus americana*, that is native to North America and resistant or highly tolerant of *A. anomala* infection, European hazelnut is generally highly susceptible. In susceptible hosts, *A. anomala* infection results in the formation of perennial cankers that girdle branches, culminating in tree death (Pinkerton 1995). Current management practices are both costly and labour intensive relying on pruning and large quantities of fungicides (Jeger et al. 2018). Further, it is documented that the *C. avellana* host protection imparted through widely disseminated resistance (R) genes from *C. avellana* 'Gasaway' in Oregon and other R genes mapped to the same linkage group, have been overcome in certain regions (Jacobs et al. 2024a). Consequently, there is a pressing need to identify alternate strategies, such as biocontrol, to reduce the impact of *A. anomala* on hazelnut cultivation.

A potential source of biocontrol agents is the native microbial community associated with a plant species of interest. The plant microbiome is typically divided into three major regions: the

phyllosphere, representing the above-ground surface regions occupied by microbes; the

endosphere, representing the regions comprised of the plant's inner tissues; and the rhizosphere,

which refers to the thin layer of soil surrounding plant roots (Gupta et al. 2021; Santoyo 2022;

Zhang et al. 2021). The rhizosphere ecosystem is particularly complex (Jones and Hinsinger 2008)

laden with an exceptional diversity of life (Dlamini et al. 2022; Zhang et al. 2020). These plant

root-associated microbial communities have been referred to as the plant's second genome

(Berendsen et al. 2012), as they are capable of profoundly extending the plant's functional

potential (Bakker et al. 2013a). Notable examples of plant benefits conferred through these

associations include enhanced nutrient availability and improved tolerance to both biotic and

abiotic stress (Bakker et al. 2013b; Hakim et al. 2021; Pantigoso et al. 2022).

Importantly, rhizosphere microbial populations can be influenced by both the plant

genotype and its disease state (Liu et al. 2019; Liu et al. 2021; Liu et al. 2023a). Though conflicting

evidence has been reported (Fitzpatrick et al. 2018), increasing host plant phylogenetic distance

has been associated with less similar bacterial rhizosphere composition (Bouffaud et al. 2014; Lei

et al. 2019; Liu et al. 2023b), while two genotypes of the same plant species can possess similar

but distinguishable microbial populations (Peiffer et al. 2013). Likewise, domestication has been

found to influence microbiome assembly and metabolic functions (Yue et al. 2023).

At the same time, rhizosphere microbial communities may be modified based on plant root

secretions, with plants potentially recruiting and stimulating disease suppressive microbial taxa in

response to pathogens as a means of improving disease outcomes (Liu et al. 2023a). Rhizosphere

microbes may mitigate plant disease through antagonistic microbe-microbe interactions (Hassani

et al. 2018), priming of the plant immune system via induced systemic resistance (ISR) (Yu et al.

2022), and enhanced nutrient acquisition (Dordas 2008; Dotaniya and Meena 2015). Microbes that

exhibit these properties may therefore prove useful in disease management, through serving as

biopesticides, enhancing systemic plant disease tolerance through eliciting ISR, or by bolstering

overall plant vigour through facilitating nutrient acquisition. The rhizosphere microbiome is

therefore worth examining as a promising reservoir of disease attenuating agents, and

characterizing interactions between plant roots and their surrounding organisms has demonstrated

extensive promise for the improvement of economically valuable crops (Orozco-Mosqueda et al.

2022).

Unfortunately, the rhizosphere of woody perennial plants remains understudied relative to

annual crops. Despite the relative paucity of research in this area, the diverse utility of woody

perennial associated rhizosphere microbes has been validated by numerous studies. For example,

phosphate solubilizing microbes with plant growth promoting potential have been reported in the

peach rhizosphere (Li et al. 2025), highlighting the prospect of leveraging rhizosphere microbial

communities to enhance woody perennial vigour. In addition, the promise of certain walnut-

associated plant growth-promoting rhizobacteria (PGPR) for enhancing drought tolerance (Lotfi

et al. 2022) illustrates the abiotic stress ameliorating potential of microbes in the woody perennial

rhizosphere. Further, examples of the capacity of rhizosphere microbes to aid in combating biotic

stress have been reported for woody perennial crops including apple (Berdeni et al. 2018;

Derikvand et al. 2023), pear (Zhang et al. 2023), plum (Ali et al. 2018), and citrus (Ezrari et al.

2021). Collectively, these studies suggest woody perennial rhizosphere microbes may aid in

addressing several significant challenges facing woody perennial cultivation.

In the current study, shotgun metagenomics was used to investigate the rhizosphere

microbiome of North American grown hazelnut plants. To investigate the effect of genotype and

disease resistant phenotypes, samples were collected from an orchard infected with *A. anomala*

and were taken from the rhizospheres of native *C. americana* plants and *C. avellana* genotypes

that are resistant or susceptible to *A. anomala* infection. By gaining insight into the taxonomic and

functional features of hazelnut trees, this study broadens the scope of available genomic

information relevant to hazelnut improvement, including for disease management, and may help

guide future identification of biocontrol agents.

112   **RESULTS**

Hazelnut rhizosphere sampling

Rhizosphere samples were collected from 15 hazelnut trees situated at Rutgers University Hort

Farm 3, East Brunswick, New Jersey, USA, in October 2022. Five samples were collected from

each of the following tree types: native *C. americana* plants (henceforth referred to as "native"),

C. avellana resistant to *A. anomala* (henceforth referred to as "resistant), and susceptible *C.*

avellana (henceforth referred to as "susceptible"). The *C. americana* plants were derived from

open pollinated seeds obtained from wild plants in Dearborn, Michigan, whereas the *C. avellana*

plants were the product of a controlled cross of a H4AR21P05 (Jacobs et al. 2024b) female (open

pollinated seed from Warsaw, Poland) with a OSU 474.084 ('Lewis' x 'Tonda di Giffoni') x OSU

540.084 (a full sibling of 'Sacajawea') male pollen source, obtained from Shawn Mehlenbacher

of Oregon State University. These *C. avellana* plants were expected to segregate for a resistance

(R) gene originating from Poland (Capik et al., 2013). The two groups of *C. avellana* trees used

for this study, therefore, represent full siblings with the resistant group expressing no signs or

symptoms of disease and the susceptible expressing obvious signs of infection with pronounced

cankers and stem dieback predicated on segregation of the major R gene. At the time of collection,

the trees were ~6 years of age and have maintained consistent disease phenotypes for ~8 years as

of 2024. All rhizosphere samples were collected from soils adjacent to tree roots, at a depth of ~ 6

inches and a radius of ~ 6 inches from the trunk of the tree. For comparison, five bulk soil samples

(not near tree roots) were also collected from a depth of ~ 6 inches around the orchard; however,

the DNA isolation from one bulk soil sample failed, leaving us with four bulk soil samples.

Metagenomic sequencing and assembly statistics

Illumina 150 bp paired-end sequencing was used to characterize the microbial community of each

of the soil samples, generating between 35,138,424 and 118,478,008 paired-end reads per sample

(**Table S1**), of which between 31,340,823 and 113,588,930 survived quality control (**Table S1**).

Metagenome assembly using MEGAHIT resulted in assemblies of 544 Mb to 4007 Mb in size,

with N50 values between 446 kb and 2260 kb (**Table S1**). When limiting the metagenomes to

contigs of at least 0.5 kb in size, the assemblies averaged 612 Mb and contained an average of

768,075 contigs (**Table S1**). Likewise, when limiting the metagenomes to contigs of at least 1 kb

in size, the assemblies averaged 187 Mb and contained an average of 115,156 contigs (**Table S1**).

Taxonomic characterization of the hazelnut rhizosphere

Initially, the community composition of each rhizosphere and bulk soil sample was evaluated

based on the raw Illumina reads using Kraken2 and Bracken, with a database consisting of the

bacteria, archaea, fungi, and virus reference libraries. However, the rate of classification by

Kraken2 was low, with just 0.80% to 1.77% of reads assigned a taxonomy (**Table S2**).

Consequently, we instead used MMSeqs2 with the UniRef90 database, to taxonomically classify

the assembled contigs (minimum contig size of 0.5 kb); using this approach, contigs encompassing

between 6% and 31% of reads were assigned a taxonomy (**Table S2**). Overall, the metagenomes

were dominated by bacteria, with 82% to 96% of reads mapping to the classified contigs

corresponding to bacteria, compared to 1.0% to 9.5% and 0.3% to 3.8% for archaea and

eukaryotes, respectively.

Alpha diversity (Shannon diversity index and Simpson diversity index [1-D]) was

calculated from the MMSeqs2 output (**Figures 1A, 1B**). No statistically significant differences

were observed between sample groups using either of the diversity measures. However, both

diversity metrics displayed a trend of the bulk soil samples having lower diversity than the

rhizosphere samples; indeed, statistically significant differences were detected for both diversity

metrics when comparing bulk soil versus all rhizosphere samples grouped as a single treatment

(Kruskal-Wallis p-value < 0.05 for both metrics). In addition, although not statistically significant,

there appeared to be greater between-replicate variance in the diversity metrics for the non-native

C. avellana compared to native *C. americana* samples, perhaps suggesting that the rhizosphere

microbial community of the non-native plants was less stably assembled than that of the native

plants.

Beta diversity was calculated at the species level using Bray-Curtis dissimilarity (**Figure**

1C). Statistically significant differences (PERMANOVA p-value < 0.001) in beta diversity were

observed between the control and each of the three rhizosphere sample groups, providing support

that the rhizosphere samples were correctly collected and were reflective of the rhizosphere

community. However, no significant differences were observed between the native, resistant, and

susceptible rhizosphere samples, suggesting that plant genotype or disease state had little to no

impact on the rhizosphere microbiome at a global level.

To ensure that the diversity metrics were not being skewed by differences in read depth,

the sequencing reads were subsampled to achieve equivalent read numbers (following QC), after

which metagenomes were reassembled and MMSeqs2 rerun. The alpha and beta diversity metrics

were then recalculated from these data. This process produced qualitatively similar results for the

alpha and beta diversity metrics (**Figure S1**), indicating that the observed diversity values were

not heavily skewed by differences in read counts between samples.

Differentially abundant taxa

The relative abundances of major taxa were investigated across samples at the phylum and genus

levels (**Figures 2 and 3**). Initial, qualitative examination of the data revealed that all samples were

dominated by the bacterial phyla *Actinomycetota*, *Pseudomonadota*, and *Acidobacteriota* (**Figure**

2A). At the genus level, the three most abundant taxa were the common soil bacteria

Bradyrhizobium, *Gaiella*, and *Nocardioides* (**Figure 3A**). Although the archaea represented only

a minority of the metagenomes, the archaeal component was dominated by the phylum

Thermoproteota (syn. *Nitrosophaerota*) (**Figure 2B**), driven by the abundance of the genera

Nitrososphaera and "*Candidatus* Nitrosocosmicus" (**Figure 3B**). Lastly, examination of the

eukaryotic taxa revealed the phylum Basidiomycota was substantially enriched in the rhizosphere

samples relative to the bulk soil control, while the phylum Ascomycota also appeared somewhat

enriched in many of the rhizosphere samples (**Figure 2C**). The differential abundance of the

phylum Basidiomycota was driven mostly by the genera *Hebeloma*, *Laccaria*, *Galerina*,

Scleroderma, and *Pisolithus* while the differential abundance of the phylum Ascomycota was

driven mostly by the genus *Morchella* and to a lesser extent the genus *Terfezia* (**Figure 3C**). It is

important to note, however, that some unexpected eukaryotic taxa were identified in relatively

high abundance, such as the shark genus *Chiloscyllium* (**Figure 3C**). This may be the result of

poor taxonomic classification of the eukaryotic contigs and/or the overall low abundance of

eukaryotes in the data. Regardless, it does suggest that caution should be taken in interpreting the

abundance data for the eukaryotic taxa.

Following qualitative assessment of the abundance data, ANCOMBC2 was used for

quantitative assessment of differential abundances of the taxa accounting for at least 1% of the

overall relative abundances of the communities. At the phylum level, the eukaryotic phylum

Basidiomycota and the bacterial phylum "*Candidatus* Rokuibacteriota" appeared enriched in all

three rhizosphere groups relative to the bulk soil controls with (log2[fold change] $\geq$ 3.9 and 2.3,

for Basidiomycota and "*Candidatus* Rokuibacteriota", respectively; **Figure 2A**). On the other

hand, the bacterial phylum *Verrucomicrobiota* appeared underrepresented specifically in the

rhizosphere communities of the "resistant" *C. avellana* plants relative to the bulk soil controls

(log2[fold change] of -1.2; **Figure 2A**). At the genus level, the three rhizosphere groups appeared

depleted in the bacterial genus *Trebonia* relative to the bulk soil control (**Figure 3A**). Compared

to the bulk soil control, the resistant and susceptible *C. avellana* samples (but not the *C. americana*

samples) both had an underrepresentation of the bacterial genera *Actinospica* (log2[fold change]

$\leq$ -1.4), *Occallatibacter* (log2[fold change] $\leq$ -2.8) and "*Candidatus* Sulfotelmatobacter"

(log2[fold change] $\leq$ -2.9), while the bacterial genus *Rhizobacter* appeared enriched (log2[fold

change] $\geq$ 1.5). The resistant *C. avellana* samples also appeared enriched in the bacterial genus

Illumatobacter relative to the bulk soil control. Lastly, the bacterial genus *Microbacterium* was

enriched in the resistant *C. avellana* samples relative to the native variety (log2[fold change] of

2.0). *Microbacterium* also visually appeared to be enriched in the susceptible *C. avellana*

compared to the native variety although this difference was not statistically significant.

Functional overview of the metagenomes

DRAM was used for functional annotation of all contigs of at least 1 kb in length, and the relative abundance of genes assigned KEGG annotations were used to construct a PCoA plot using Bray-Curtis dissimilarities as a general overview of the functional differentiation of each sample (**Figure 4**). Although global differences between samples were detected (PERMANOVA p-value of 0.031), no statistically significant differences were noted for any of the pairwise comparisons between sample groups. However, although not statistically significant, there did appear to be a trend towards the native *C. americana* and resistant *C. avellana* samples grouping together and separating from the sensitive *C. avellana* and bulk soil control. Likewise, no clear differences were noted between sample types when comparing the presence of various functions and pathways related to carbohydrate metabolism, methanogenesis and methanotrophy, nitrogen metabolism, short-chain fatty acid and alcohol conversions, and sulfur metabolism (**Figure S2**). Overall, these results indicate that there was little functional differentiation between the sample types at a global level.

In addition, prokaryotic biosynthetic gene clusters (BGCs) were annotated in all metagenome assemblies (limited to contigs of at least 0.5 kb) using the program antiSMASH. BiG-SCAPE was then used to group the antiSMASH annotations based on the class and predicted products of the BGCs (**Figure 5**). The distribution of BGCs across samples was largely similar, regardless of the sample group, with BGC classes associated with nonribosomal peptide synthetases (NRPS), ribosomally synthesized and post-translationally modified peptides (RiPPs), and terpenes generally present with the highest relative abundance.

Reconstruction and characterization of metagenome-assembled genomes

Following dereplication using an average nucleotide identity (ANI) threshold of 99%, 45 high-quality (completeness ≥ 70%, contamination ≤ 10%) metagenome-assembled genomes (MAGs) were constructed from the dataset and taxonomically categorized with GTDB-Tk, including 41 bacterial and 4 archaeal MAGs (**Figure 6, Table S3**). Interestingly, only one of the 45 MAGs could be taxonomically classified as the species level when using the Genome Taxonomy Database Toolkit (GTDB-Tk) with database R220, while an additional nine were classified to the genus level. The four archaeal MAGs represent four distinct species from three genera within the family *Nitrososphaeraceae* (phylum *Thermoproteota*). On the other hand, nine phyla are represented amongst the 41 bacterial MAGs, with the majority of MAGs belonging to the phyla *Actinomycetota* (21 MAGs), *Pseudomonadota* (9 MAGs), and *Acidobacteriota* (6 MAGs). MAGs were functionally annotated with DRAM, and a summary of the results are provided as **Figure S3**. In addition, antiSMASH and BiG-SCAPE were used to detect and summarize the abundance of BGCs in the MAGs, and these results are summarized in **Figure S4**.

DISCUSSION

The hazelnut rhizosphere microbiome showed only minor differences compared to bulk soil

Previous studies examining bacterial community structure in the roots of the woody plants (*Quercus*, *Populus*, and *Pinus*), found that bacterial community structure was shaped more strongly by host tree species than by soil origin (Bonito et al. 2014). We therefore expected to see differences between the microbial community of the rhizosphere samples compared to the bulk soil samples. As expected, the bulk soil samples were clearly delineated from the rhizosphere samples based on the alpha and beta diversity measures. This differentiation appeared to be mainly

driven by changes in frequency of low abundance taxa as we observed few differences in the

relative abundances of the primary taxa (those with relative abundances ≥ 1%). However, the

current study examined only a single time point. Considering that another study observed that the

prokaryotic component of the maize rhizosphere only differed from bulk soil at specific stages of

plant development (Wattenburger et al. 2019), it may be that sampling at a different time of year,

or sampling deeper roots, would reveal more dramatic changes in the hazelnut rhizosphere

microbial community structure than observed here.

Few differences were observed between the microbial communities of the different

hazelnut genotypes, contrary to our expectation (Lopes et al. 2023). Likewise, we observed little

difference between *C. avellana* plants that were or were not infected with *A. anomala*, despite

previous studies observing that foliar pathogens can indirectly impact rhizosphere microbial

communities in ways that could potentially have positive impacts on plant disease tolerance

(Berendsen et al. 2018; Jiang et al. 2023; Luo et al., 2022; Yang et al. 2020). While these results

could suggest that hazelnut genotype and disease status have little impact on the rhizosphere

microbiome, it may also be that differences would be more pronounced at other times of the year,

or that deeper sequencing and/or taxonomic classification of a higher portion of the contigs would

reveal additional differences missed here.

Composition of the hazelnut microbial community

We found that the fungal phylum Basidiomycota was enriched in all rhizosphere samples relative

to bulk soil. Likewise, the fungal phyla Ascomycota was slightly enriched in the rhizosphere

samples, although this enrichment was not statistically significant. Interestingly, a separate study

reported Ascomycota, Basidiomycota, and Mortierellomycota as the dominant fungal phyla in the

rhizosphere of hazelnut varieties grown in Beijing, China (Ma et al. 2021). Similarly, the phyla

Basidiomycota and Ascomycota were the dominant fungal phyla in the rhizosphere of the

deciduous shrub *Paeonia ludlowii*, with Basidiomycota being the most dominant in the autumn

(Qiao et al. 2023). Overall, the consistency in these results provides support that our rhizosphere

samples accurately capture the hazelnut rhizosphere microbiome.

Although there is limited data on the effects of bacteria on hazelnut tree health and growth,

one study reported that inoculation of *C. avellana* seedlings with the bacteria *Pseudomonas putida*,

B. subtilis, and *Enterobacter cloacae* can improve growth (Rostamikia et al. 2017). Therefore, we

wished to use our metagenomics data to help identify microbes in the hazelnut rhizosphere

microbiome that could potentially be targets to develop into biocontrol agents to limit the impact

of *A. anomala* infection. To this end, we focused on the microbes that were either enriched or

depleted in the hazelnut microbiome. One taxon of particular interest based on the relative and

differential abundance data was the bacterial phylum "*Candidatus* Rokuibacteriota", which was

enriched in all three rhizosphere groups relative to bulk soil. "*Candidatus* Rokuibacteriota" have

large genomes with high GC content, with the capacity for multifaceted mixotrophic metabolism

(Becraft et al. 2017). Bacteria from this phylum have also been reported to encode numerous

biosynthetic loci in their genomes (Sharrar et al. 2020). Consequently, members of this phylum

exhibit characteristics of relevance for biocontrol applications, and we suggest that their utility

should be explored within this context.

On the other hand, the bacterial genera *Actinospica*, *Occallatibacter*, and "*Candidatus*

Sulfotelmatobacter" were depleted in the susceptible and tolerant varieties of *C. avellana* relative

to bulk soil, unlike the native *C. americana* variety. *Actinospica* may be of particular note for *A.*

anomala disease management, as there is an ongoing interest in the efficacy of acidophilic

actinomycetes as biocontrol agents against phytopathogenic fungi (Poomthongdee et al. 2015).

Supplementing the abundance of this genus in the *C. avellana* microbiome may therefore be a

strategy worth exploring for *A. anomala* disease management.

In terms of eukaryotic microbes, associations with mycorrhizal fungi have been shown to

enhance disease tolerance in woody perennials (Dong et al. 2021; Kebert et al. 2022), and a variety

of additional benefits have been described for mycorrhizal associated hazelnut trees. In *C.*

avellana, association with the ectomycorrhizal forming fungus *Tuber melanosporum* was reported

to regulate important plant defense genes, which may favourably impact the host plant's response

to stress (Sillo et al. 2022). Inoculation of hybrid hazel (*C. heterophylla* x *C. avellana*) seedlings

with the ectomycorrhizal fungus *Scleroderma bovista* was reported to promote growth, enhancing

photosynthetic efficiency and total plant weight (Cheng et al. 2024). In addition, inoculation with

the arbuscular mycorrhizal fungus *Glomus iranicum* var. *tenuihypharum* was found to increase the

number of female flowers and overall yield in the Tonda di Giffoni variety of *C. avellana* by 60%

and 38%, respectively, relative to the uninoculated control (Luciani et al. 2019). In our study, the

fungal genera *Hebeloma*, *Galerina*, *Laccaria*, *Morchella*, *Pisolithus*, and *Scleroderma* appeared

enriched in the three groups of rhizosphere samples, relative to the control group; however, as the

relative abundances of these taxa were < 1%, the differences in their abundances was not tested

using ANCOMBC2. Isolates of species from these fungal taxa from the hazelnut rhizosphere may

therefore represent interesting candidates to explore for plant growth promotion and/or biocontrol

against *A. anomala* in North America.

Primary metabolism of the hazelnut rhizosphere microbiome

Rhizosphere microbes play important roles in nutrient cycling and availability (Thepbandit and

Athinuwat 2024) and consequently, they can impact plant nutrition and thus disease tolerance. For

example, nitrogen availability can impact both physical and biochemical defence mechanisms in

plants, yet the consequences of enhanced endogenous nitrogen can be difficult to anticipate, as

they may be favourable or detrimental to plant pathogen tolerance depending on context (Sun et

al. 2020). In our study, global differences in the KEGG metabolic annotations (PERMANOVA p-

value of 0.031) were observed between samples, however no pairwise differences were detected

between sample groups.

Pathways involved in nitrogen metabolism were well represented across most samples

including ammonia oxidation, conversion of nitrate to nitrite, nitrite to nitrate, nitrite to nitric

oxide, and nitric oxide to nitrous oxide. Though less common, some samples possessed the

capacity for nitrite reduction to ammonia as observed in Bulk soil 1, Bulk soil 2, Native 3,

Susceptible 1, Susceptible 2, and Susceptible 4 samples. As nitrate serves as the major source of

nitrogen for most land plants in aerobic soils (Ho and Tsay 2010), the presence of pathways

involved in the production of nitrate across samples highlights the capacity of microbes within

these systems to facilitate the production of bioavailable nitrogen for hazelnut. Bioavailable sulfur

can also have drastic impacts on plant disease outcomes. For example, inorganic sulfate salts can

significantly enhance the resistance of plants to certain fungal pathogens, through a process

deemed sulfur-induced resistance (SIR) (Künstler et al. 2020). Here, pathways responsible for the

conversion of thiosulfate to sulfate were well represented across all samples besides the

Susceptible 3 sample. As both bioavailable nitrogenous and sulfur-containing compounds have

been demonstrated to alter plant disease tolerance (Bloem et al. 2015; Mur et al. 2017) the

application of certain microbial communities that alter the abundance of bioavailable forms of

these compounds within the soil may be explored as an alternative means of broadly improving

plant health.

Many of the MAGs generated in this study, particularly those within the orders *Gaiellales*

and *Hyphomicrobiales* (syn. *Rhizobiales*), encoded pathways involved in the conversion of nitrite

to nitric oxide (NO). NO can be generated endogenously in plants and performs important

functions in various physiological responses and developmental stages (Hancock 2020; Sun et al.

2021), with roles in plant tolerance to abiotic and biotic stress (Fancy et al. 2017, 201; Mur et al.

2017). Exogenous application of NO, through the use of NO donors, has been shown to improve

plant disease tolerance in several instances (Nasir et al. 2020; Khan et al. 2021; Bakhoum et al.

2023). Furthermore, studies suggest that leveraging NO-producing organisms to enhance disease

tolerance in plants is a promising avenue to explore. For example, a study examining tomato

infected with *Ralstonia solanacearum* found that *Pseudomonas fluorescens* strains overproducing

NO displayed better biocontrol efficacy than those that don't overproduce NO (Wang et al. 2005).

Consequently, it would be interesting to isolate diverse strains from promising genera within the

Gaiellales and *Hyphomicrobiales* orders, which we found commonly encode the pathways for NO

production, and assess them for their impact on hazelnut disease tolerance. Of particularly interest

would be isolates that additionally have the potential to convert thiosulfate to sulfate.

Secondary metabolism of the hazelnut rhizosphere microbiome

Soil microbes represent a rich source of secondary metabolites (Sharrar et al. 2020) that perform

diverse functions in ecological systems and can mediate symbioses between microbes and plants

in certain contexts (Demain and Fang 2000) and ameliorate biotic stress in plants (Ruparelia et al.

2022). Analyzing the biosynthetic gene clusters (BGCs) involved in the production of secondary

metabolites within the hazelnut rhizosphere, therefore, provides an opportunity to identify

compounds and microbial taxa that can be screened as biological control agents in hazelnut.

In our study, no substantial differences in the relative abundance of specific BGC classes

were evident between conditions, although all conditions had an abundance of BGCs encoding the

production of non-ribosomal peptides (NRPs), ribosomally synthesized and post-translationally

modified peptides (RiPPs), and terpenes. The production of secondary metabolites of these classes

has repeatedly been shown to contribute to the biocontrol potential of plant growth-promoting

rhizobacteria (PGPRs). For example, *Bacillus* species produce NRPs that can serve as biocontrol

agents, such as iturin, surfactin and fengycin (Ranjan et al. 2023). Likewise, species of the fungal

genus *Trichoderma* have demonstrated potential as biocontrol agents (Sood et al. 2020), which

could relate to the diversity of terpenoids and complex terpenes produced by these organisms (Bai

et al. 2022; Lee et al. 2016).

Many of the MAGs assembled in this study have the genetic potential to produce diverse

secondary metabolites. One such example is MAG-30, classified as belonging to the species

"*Lysobacter yananisis*". Species of the genus *Lysobacter* have potential as biocontrol agents

(Kilic-Ekici and Yuen 2003; Puopolo et al. 2014), and they produce a range of antibiotics active

against numerous plant pathogens (Gómez Expósito et al. 2015; Hayward et al. 2010). Likewise,

our *Lysobacter* MAG has the genetic potential to produce various secondary metabolites of

relevance for biocontrol, including NRPs, lantipeptides, polyketides, and RiPPs. Another MAG

with high biosynthetic potential is MAG-07, classified as belonging to the genus *Actinospica*.

Species of this genus have been isolated from diverse ecosystems including forest soils (Kusuma

et al. 2022; Cavaletti et al. 2006), and studies suggest that members of this genus have strong

biocontrol potential (Qi et al. 2019; Rodriguez-Mena et al. 2022). Indeed, members of the genus

Actinospica have high genetic potential to produce a wide array of secondary metabolites (Majer

et al. 2021). Likewise, our *Actinospica* MAG contains several genomic regions potentially

involved in the production of NRPs, lantipeptides, RIPPs, polyketides, and terpenes. Considering

these results, it would be interesting to isolate strains of the genera *Lysobacter* and *Actinospica*

from hazelnut rhizosphere soils and investigate their potential to serve as biocontrol strains and

reduce the susceptibility of hazelnut to *A. anomala* infection.

Conclusions

The co-evolution of plants and their resident microbial communities has resulted in a complex

dynamic where the microbial genetic complement can drastically impact important plant features.

Soil microbes participate in important biogeochemical cycles that impact plant nutrient

acquisition, while plant-associated microbes can also influence plant immunity. Here,

metagenomics was used to examine the microbial communities of three hazelnut rhizosphere

sample groups obtained from trees that displayed differing levels of resistance to the pathogen *A.*

anomala: *C. americana* (resistant), *C. avellana* (susceptible) and *C. avellana* (resistant). While the

rhizosphere samples were differentiated from the bulk soil samples, little taxonomic or functional

variation was seen between the different rhizosphere groups. A total of 45 dereplicated, high-

quality MAGs were generated, many of which carried multiple BGCs. This included two MAGs

representing organisms from the bacterial genera *Actinospica* and *Lysobacter* that have previously

been identified as potential biocontrol agents. This study serves to inform future efforts aimed at

leveraging hazelnut-associated microbes toward the improvement of diverse hazelnut features

including disease outcomes, identifying promising candidate taxa that could be isolated and

screened for their capacity to manage *A. anomala* infection. In addition, this study provides

insights into the taxonomic diversity and functional potential of the rhizosphere microbial

communities of hazelnut, expanding our understanding of the rhizosphere microbiome of woody

perennials.

431   **METHODS**

Soil sample collection

Bulk soil and rhizosphere samples were collected from Hort Farm 3, 67 Ryders Lane, East

Brunswick, NJ 08816, between 1:00 PM and 4:00 PM on 19 October 2022. Soils were collected

using a small shovel that was sterilized with 70% ethanol before each sample collection, and

transferred to sterile 50 mL tubes. Soils from highly tolerant *C. americana* trees were collected

from row 1 of the farm. Soils from *C. avellana* trees collected from row 3 were divided into two

groups segregating for disease resistance, designated resistant and susceptible, based on the

presence or absence of pronounced *A. anomala* cankers. Five soil samples were collected for each

of the three hazelnut varieties; samples were collected at a depth of approximately 15 cm,

approximately 15 cm away from the trunk of the tree. In addition, five bulk soil samples were

collected at a depth of approximately 15 cm from five sites around the periphery of the hazelnut

orchard.

DNA isolation and sequencing

Approximately one hour after sample collection was complete, ~0.42 g subsamples of soil were

taken from each soil sample. DNA extraction was performed using a Qiagen DNeasy PowerSoil

Pro Kit, according to the manufacturer's instructions. One of the bulk soil samples was omitted

from downstream analysis due to failure of the DNA extraction. All other DNA samples were

subsequently refrigerated until they were shipped to Génome Québec (Montréal, Quebec, Canada)

for library preparation and Illumina sequencing. For samples B2, B3, B5, R1P5, R1P9, R1P14,

R3P3, R3P15, R3P20, R3P9-, R3P10-, R3P12- (**Table 1**), library preparation was performed using

the NxSeq® AmpFREE Low DNA Library Preparation Kit (Lucigen) using the Xp protocol, as

suggested by the manufacturer. These samples were sequenced using 150-bp paired-end

technology on an Illumina NovaSeq 6000 instrument. For samples B4, R1P1, R1P18, R3P5,

R3P11, R3P14-, R3P18- (**Table 1**), library preparation was performed using the (Illumina®) DNA

PCR-Free Prep, Tagmentation Kit following the manufacturer's recommendations. These samples

were likewise sequenced using 150-bp paired-end technology, but on an Illumina NovaSeq X Plus

instrument.

Taxonomic classification of the raw Illumina reads

Illumina reads were taxonomically classified using Kraken (v2.1.3) (Wood et al. 2019) using a

database consisting of the bacteria, archaea, fungi, and virus reference libraries, and post-processed

using Bracken (v2.7) (Lu et al. 2017). Alpha diversity and beta diversity measurements were

determined based on the species-level output of Bracken. Diversity measurements were

determined, analyzed, and visualized in R (v4.5.1) (R Core Team 2025) using tidyverse (v2.0.0)

(Wickham et al. 2019), dplyr (v1.1.4) (Wickham et al. 2018) , phyloseq (v1.52.0) (McMurdie and

Holmes 2013), stats (v4.5.1) (R Core Team 2025), ggpubr (v0.6.1) (Kassambara 2025), vegan

(v2.7.1) (Oksanen et al. 2024), pairwiseAdonis (v0.4.1) (Martinez Arbizu 2017), ggplot2 (v4.0.0)

(Wickham 2009), and patchwork (v1.3.2) (Pedersen 2019) packages. KronaTools (v2.8) (Ondov

et al. 2011) was used for generation of krona plots.

Metagenome assembly and binning

Illumina reads were filtered using BBduk (v39.06) (Bushnell 2014) and trimmed using Trimmomatic (v0.39) (Bolger et al. 2014) with the parameters: LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36. Following trimming, metagenomes were assembled using MEGAHIT (v1.2.9) (Li et al. 2015) with the "presets" option set to "meta-large". Metagenome assembly statistics were calculated using the stats.sh script of BBMap (v37.36) (Bushnell 2014). In addition, to confirm that the alpha and beta diversity metrics were not confounded by differences in sequencing depth, seqtk (v1.3) (https://github.com/lh3/seqtk) was used to filter the trimmed reads to obtain a subset corresponding to 90% of the number of reads from the sample with the fewest reads. These reduced read sets were then used for metagenome assembly using MEGAHIT with the "presets" option set to "meta-large".

Assembly of metagenome-assembled genomes

Prior to binning, contigs less than 1500 bp were discarded using pullseq (v1.0.2) (https://github.com/bcthomas/pullseq). Next, an all-against-all mapping of the Illumina reads against each metagenome was performed using bowtie2 (v2.4.5) (Langmead and Salzberg 2012) and sorted using samtools (v1.15.1) (Li et al. 2009). Binning was then performed independently for each sample using MetaBAT2 (v2.14) (Kang et al. 2019), CONCOCT (v1.1.0) (Alneberg et al. 2014), MaxBin2 (v2.2.7) (Wu et al. 2016), and SemiBin2 (v2.1.0) (Pan et al. 2023). From each of the metagenomes, an optimized set of MAGs was produced using DAS_Tool (v1.1.4) (Sieber et al. 2018) and the output of the four binning algorithms. In addition, bins were generated with VAMB (v4.1.3) (Nissen et al. 2021, 20), using all metagenomes and with read mapping performed using minimap2 (v2.24-r1122) (Li 2018). Finally, a non-redundant set of high-quality MAGs (>

70% completeness; < 10% contamination) was generated from all the produced MAGs using dRep

(v3.4.1) (Olm et al. 2017) and a secondary clustering threshold of 99%. The dependencies for dRep

included Mash (v2.3) (Ondov et al. 2016) MUMmer (v4.0.0rc1) (Kurtz et al. 2004), CheckM

(v1.2.2) (Parks et al. 2015), Prodigal (v2.6.3) (Hyatt et al. 2010), and pplacer (v1.1.alpha19)

(Matsen et al. 2010).

Taxonomic classification of the assembled metagenomes

Prior to taxonomic classification, contigs less than 500 bp were discarded using pullseq. Contigs

were taxonomically classified using MMSeqs2 (v0a6b8c) (Steinegger and Söding 2017) with the

UniRef90 database (downloaded 18 July 2024). Custom scripts were used to convert the MMSeqs2

output data into taxa abundance tables, with taxa abundances calculated as the sum of contig length

multiplied by contig sequencing depth for all contigs with a given taxonomic classification. Alpha

diversity and beta diversity metrics were determined, analyzed, and visualized in R using the

tidyverse, ggplot2, dplyr, phyloseq, stats, ggpubr, vegan, pairwiseAdonis, and patchwork

packages. The relative abundances of the taxa were visualized as bubble plots and stacked bar plots

with R's ggplot2, patchwork, and randomcoloR (v1.1.0.1) (Ammar 2019) packages. Lastly, the R

packages ANCOMBC2 (v2.10.1) (Lin and Peddada 2024) and microbiome (v1.30.0) (Lahti and

Shetty 2019) were used to identify statistically significant differences in taxa abundances for taxa

with a relative abundance of at least 1% in at least one sample. The relevant data from these

analyses were visualized in R using the ggplot2 and patchwork packages.

Taxonomic classification of metagenome-assembled genomes

MAGs were taxonomically classified using GTDB-Tk (v2.4.0) with the R220 database and the

classify_wf workflow (Chaumeil et al. 2020). To construct a phylogeny of the bacterial MAGs,

the identify function of GTDB-Tk was used to identify homologs of the bac120 gene set, following

which the align function of GTDB-Tk was used to align each marker set and then concatenate the

alignments. Lastly, the concatenated alignment was used to construct a maximum likelihood

phylogeny using IQ-TREE2 (v2.2.2.4) (Minh et al. 2020). First, ModelFinder (Kalyaanamoorthy

et al. 2017), was used to identify the best scoring model based on Bayesian information criterion

(BIC). Then, IQ-TREE2 was run using the best-scoring model (LG+F+R5) with branch support

assessed using the Shimodaira–Hasegawa-like approximate likelihood ratio test (SH-aLRT) and

ultrafast jackknife analysis with a subsampling proportion of 40%, with both metrics calculated

from 1,000 replicates. Phylogenies were visualized using iTOL (Letunic and Bork 2024).

Characterization of the metabolic potential of the metagenomes and MAGs

Prior to functional annotation, contigs less than 1000 bp were discarded from the metagenomes

using pullseq. Next, DRAM (v0.1.2) (Shaffer et al. 2020), as implemented on the KBase platform

(Arkin et al. 2018), was used to annotate all metagenomes and MAGs. For the metagenomes, the

KEGG annotations were extracted from the annotation file, and a Bray-Curtis distance matrix was

produced and visualized using a Principal Coordinate Analysis in R with the ggplot2, dplyr,

phyloseq, and patchwork packages. Statistical analyses were performed in R using the vegan and

pairwiseAdonis packages. Group specific functional differences were determined and visualized

using the dplyr, stats, dunn.test (v1.3.6) (Dinno 2024), and ggplot2 packages.

Prior to annotation of BGCs, contigs less than 500 bp were discarded from the

metagenomes using pullseq. BGCs were identified in the reduced metagenomes using antiSMASH

(v7.0) (Blin et al. 2023) with default parameters, after which BiG-SCAPE (v1.1.8) (Navarro-

Muñoz et al. 2020) was run with default parameters to examine BGC diversity across the

metagenomes. Likewise, BGCs were identified in the MAGs using antiSMASH and BiG-SCAPE.

The output files were processed and visualized as heatmaps in R using the dplyr, ggplot2 and

patchwork packages.

Data availability

All raw Illumina sequencing data, metagenome assemblies, and MAGs are available from the

National Center for Biotechnology Information (NCBI) under BioProject PRJNA1338491

Accessions for individual datasets are provided in **Table S3**. All code to repeat the analyses

described in this manuscript is available through GitHub (github.com/diCenzo-

Lab/016_2025_hazelnut_rhizosphere_metagenomics).

555   **ACKNOWLEDGEMENTS**

We thank Sharen Roland and other personnel at the Centre d'Expertise et de Services (CES) of

Génome Québec for the advice on sequencing strategies and for performing all the Illumina

sequencing reported in this study. We would also like to thank Rutgers University Department of

Plant Biology for providing access to their facilities for sample collection and preparation. This

work was also supported by the USDA-NIFA Hatch project NJ12171 through the New Jersey

Agricultural Experiment Station. This work was funded by the Natural Sciences and Engineering

Research Council of Canada (NSERC) through Discovery Grants to S.R. and G.C.D. and the

NSERC Collaborative Research and Training Experience Grant "Genome Editing for

Food Security and Environmental Sustainability" to S.R. This research was enabled, in part,

through computational resources provided by Compute Ontario (computeontario.ca) and the

Digital Research Alliance of Canada (alliancecan.ca).

568   **REFERENCES**

Ali, I., Sultan, T., Subhan, F., Haleem, K. S., Sultana, N., and Tauseef, I. 2018. PGPRs of plum

(*Prunus domestica*) rhizosphere enhance plant growth and antagonize fungal activity in

vitro. **Acta Agric. Scand. Sect. B — Soil Plant Sci.** 68:367–378.

https://doi.org/10.1080/09064710.2017.1410565.

Alneberg, J., Bjarnason, B. S., de Bruijn, I., Schirmer, M., Quick, J., Ijaz, U. Z., Lahti, L., Loman,

N. J., Andersson, A. F., and Quince, C. 2014. Binning metagenomic contigs by coverage and

composition. **Nat. Methods** 11:1144–1146. https://doi.org/10.1038/nmeth.3103.

Ammar, R. 2019. **randomcoloR: generate attractive random colors.** https://CRAN.R-

project.org/package=randomcoloR.

Arkin, A. P., Cottingham, R. W., Henry, C. S., Harris, N. L., Stevens, R. L., Maslov, S., Dehal, P.,

Ware, D., Perez, F., Canon, S., Sneddon, M. W., Henderson, M. L., Riehl, W. J., Murphy-

Olson, D., Chan, S. Y., Kamimura, R. T., Kumari, S., Drake, M. M., Brettin, T. S., Glass, E.

M., Chivian, D., Gunter, D., Weston, D. J., Allen, B. H., Baumohl, J., Best, A. A., Bowen,

B., Brenner, S. E., Bun, C. C., Chandonia, J.-M., Chia, J.-M., Colasanti, R., Conrad, N.,

Davis, J. J., Davison, B. H., DeJongh, M., Devoid, S., Dietrich, E., Dubchak, I., Edirisinghe,

J. N., Fang, G., Faria, J. P., Frybarger, P. M., Gerlach, W., Gerstein, M., Greiner, A.,

Gurtowski, J., Haun, H. L., He, F., Jain, R., Joachimiak, M. P., Keegan, K. P., Kondo, S.,

Kumar, V., Land, M. L., Meyer, F., Mills, M., Novichkov, P. S., Oh, T., Olsen, G. J., Olson,

R., Parrello, B., Pasternak, S., Pearson, E., Poon, S. S., Price, G. A., Ramakrishnan, S.,

Ranjan, P., Ronald, P. C., Schatz, M. C., Seaver, S. M. D., Shukla, M., Sutormin, R. A., Syed,

M. H., Thomason, J., Tintle, N. L., Wang, D., Xia, F., Yoo, H., Yoo, S., and Yu, D. 2018. KBase: The United States Department of Energy Systems Biology Knowledgebase. **Nat. Biotechnol.** 36:566–569. https://doi.org/10.1038/nbt.4163.

Bai, B., Liu, C., Zhang, C., He, X., Wang, H., Peng, W., and Zheng, C. 2022. *Trichoderma* species from plant and soil: an excellent resource for biosynthesis of terpenoids with versatile bioactivities. **J. Adv. Res.** 49:81–102. https://doi.org/10.1016/j.jare.2022.09.010.

Bakhoum, G. Sh., Sadak, M. Sh., and Thabet, M. S. 2023. Induction of tolerance in groundnut plants against drought stress and *Cercospora* leaf spot disease with exogenous application of arginine and sodium nitroprusside under field conditions. **J. Soil Sci. Plant Nutr.** 23:6612–6631. https://doi.org/10.1007/s42729-023-01514-x.

Bakker, P. A. H. M., Berendsen, R. L., Doornbos, R. F., Wintermans, P. C. A., and Pieterse, C. M. J. 2013a. The rhizosphere revisited: root microbiomics. **Front. Plant Sci.** 4. https://doi.org/10.3389/fpls.2013.00165.

Bakker, P. A. H. M., Doornbos, R. F., Zamioudis, C., Berendsen, R. L., and Pieterse, C. M. J. 2013b. Induced systemic resistance and the rhizosphere microbiome. **Plant Pathol. J.** 29:136–143. https://doi.org/10.5423/PPJ.SI.07.2012.0111.

Becraft, E. D., Woyke, T., Jarett, J., Ivanova, N., Godoy-Vitorino, F., Poulton, N., Brown, J. M., Brown, J., Lau, M. C. Y., Onstott, T., Eisen, J. A., Moser, D., and Stepanauskas, R. 2017. Rokubacteria: genomic giants among the uncultured bacterial phyla. **Front. Microbiol.** 8. https://doi.org/10.3389/fmicb.2017.02264.

Berdeni, D., Cotton, T. E. A., Daniell, T. J., Bidartondo, M. I., Cameron, D. D., and Evans, K. L. 2018. The effects of arbuscular mycorrhizal fungal colonization on nutrient status, growth, productivity, and canker resistance of apple (*Malus pumila*). **Front. Microbiol.** 9:1461.

https://doi.org/10.3389/fmicb.2018.01461.

Berendsen, R. L., Pieterse, C. M. J., and Bakker, P. A. H. M. 2012. The rhizosphere microbiome

and plant health. **Trends Plant Sci.** 17:478–486.

https://doi.org/10.1016/j.tplants.2012.04.001.

Berendsen, R. L., Vismans, G., Yu, K., Song, Y., de Jonge, R., Burgman, W. P., Burmølle, M.,

Herschend, J., Bakker, P. A. H. M., and Pieterse, C. M. J. 2018. Disease-induced assemblage

of a plant-beneficial bacterial consortium. **ISME J.** 12:1496–1507.

https://doi.org/10.1038/s41396-018-0093-1.

Blin, K., Shaw, S., Augustijn, H. E., Reitz, Z. L., Biermann, F., Alanjary, M., Fetter, A., Terlouw,

B. R., Metcalf, W. W., Helfrich, E. J. N., van Wezel, G. P., Medema, M. H., and Weber, T.

2023. antiSMASH 7.0: new and improved predictions for detection, regulation, chemical

structures and visualization. **Nucleic Acids Res.** 51:W46–W50.

https://doi.org/10.1093/nar/gkad344.

Bloem, E., Haneklaus, S., and Schnug, E. 2015. Milestones in plant sulfur research on sulfur-

induced-resistance (SIR) in Europe. **Front. Plant Sci.** 5.

https://doi.org/10.3389/fpls.2014.00779.

Boccacci, P., and Botta, R. 2009. Investigating the origin of hazelnut (*Corylus avellana* L.)

cultivars using chloroplast microsatellites. **Genet. Resour. Crop Evol.** 56:851–859.

https://doi.org/10.1007/s10722-009-9406-6.

Bolger, A. M., Lohse, M., and Usadel, B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. **Bioinformatics** 30:2114–2120.

https://doi.org/10.1093/bioinformatics/btu170.

Bonito, G., Reynolds, H., Robeson II, M. S., Nelson, J., Hodkinson, B. P., Tuskan, G., Schadt, C.

W., and Vilgalys, R. 2014. Plant host and soil origin influence fungal and bacterial

assemblages in the roots of woody plants. **Mol. Ecol.** 23:3356–3370.

https://doi.org/10.1111/mec.12821.

Bouffaud, M., Poirier, M., Muller, D., and Moënne-Loccoz, Y. 2014. Root microbiome relates to

plant host evolution in maize and other Poaceae. **Environ. Microbiol.** 16:2804–2814.

https://doi.org/10.1111/1462-2920.12442.

Bushnell, B. 2014. BBMap: a fast, accurate, splice-aware aligner.

https://www.osti.gov/biblio/1241166.

Capik, J. M., Muehlbauer, M., Novy, A., Honig, J. A., & Molnar, T. J. 2013. Eastern Filbert Blight-

resistant hazelnuts from Russia, Ukraine, and Poland. **HortScience** 48:466–

473. https://doi.org/10.21273/HORTSCI.48.4.466.

Cavaletti, L., Monciardini, P., Schumann, P., Rohde, M., Bamonte, R., Busti, E., Sosio, M., and

Donadio, S. 2006. *Actinospica robiniae* gen. nov., sp. nov. and *Actinospica acidiphila* sp.

nov.: proposal for *Actinospicaceae* fam. nov. and *Catenulisporinae* subord. nov. in the order

Actinomycetales. **Int. J. Syst. Evol. Microbiol.** 56:1747–1753.

https://doi.org/10.1099/ijs.0.63859-0.

Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P., and Parks, D. H. 2020. GTDB-Tk: a toolkit to

classify genomes with the Genome Taxonomy Database. **Bioinformatics** 36:1925–1927.

https://doi.org/10.1093/bioinformatics/btz848.

Cheng, Y., Sun, S., Lou, H., Dong, Y., He, H., Mei, Q., and Liu, J. 2024. The ectomycorrhizal

fungus *Scleroderma bovista* improves growth of hazelnut seedlings and plays a role in auxin

signaling and transport. **Front. Microbiol**. 15. https://doi.org/10.3389/fmicb.2024.1431120.

Demain, A. L., and Fang, A. 2000. The natural functions of secondary metabolites. **Adv. Biochem.**

Eng. Biotechnol. 69:1–39. https://doi.org/10.1007/3-540-44964-7_1.

Derikvand, F., Bazgir, E., El Jarroudi, M., Darvishnia, M., Mirzaei Najafgholi, H., Laasli, S.-E.,

and Lahlali, R. 2023. Unleashing the potential of bacterial isolates from apple tree

rhizosphere for biocontrol of *Monilinia laxa*: a promising approach for combatting brown

rot disease. **J. Fungi** 9:828. https://doi.org/10.3390/jof9080828.

Dinno, A. 2024. **dunn.test: dunn's test of multiple comparisons using rank sums**.

https://CRAN.R-project.org/package=dunn.test.

Dlamini, S. P., Akanmu, A. O., and Babalola, O. O. 2022. Rhizospheric microorganisms: the

gateway to a sustainable plant health. **Front. Sustain. Food Syst.** 6:925802.

https://doi.org/10.3389/fsufs.2022.925802.

Dong, F., Wang, Y., and Tang, M. 2021. Effects of *Laccaria bicolor* on gene expression of *Populus*

trichocarpa root under poplar canker stress**. J. Fungi** 7:1024.

https://doi.org/10.3390/jof7121024.

Dordas, C. 2008. Role of nutrients in controlling plant diseases in sustainable agriculture. A review.

Agron. Sustain. Dev. 28:33–46. https://doi.org/10.1051/agro:2007051.

Dotaniya, M. L., and Meena, V. D. 2015. Rhizosphere effect on nutrient availability in soil and its

uptake by plants: a review. **Proc. Natl. Acad. Sci. India Sect. B Biol. Sci.** 85:1–12.

https://doi.org/10.1007/s40011-013-0297-0.

Ezrari, S., Mhidra, O., Radouane, N., Tahiri, A., Polizzi, G., Lazraq, A., and Lahlali, R. 2021.

Potential role of rhizobacteria isolated from citrus rhizosphere for biological control of citrus

dry root rot. **Plants Basel Switz.** 10:872. https://doi.org/10.3390/plants10050872.

Fancy, N. N., Bahlmann, A.-K., and Loake, G. J. 2017. Nitric oxide function in plant abiotic stress.

Plant Cell Environ. 40:462–472. https://doi.org/10.1111/pce.12707.

Fitzpatrick, C. R., Copeland, J., Wang, P. W., Guttman, D. S., Kotanen, P. M., and Johnson, M. T.

J. 2018. Assembly and ecological function of the root microbiome across angiosperm plant

species. **Proc. Natl. Acad. Sci.** 115:E1157–E1165.

https://doi.org/10.1073/pnas.1717617115.

Food and Agricultural Organization of the United Nations. 2021. **FAOSTAT**.

https://www.fao.org/faostat/en/#data/QCL [Accessed 15 Oct. 2024]

Gómez Expósito, R., Postma, J., Raaijmakers, J. M., and De Bruijn, I. 2015. Diversity and activity

of *Lysobacter* species from disease suppressive soils. **Front. Microbiol.** 6:1243.

https://doi.org/10.3389/fmicb.2015.01243.

Gupta, R., Anand, G., Gaur, R., and Yadav, D. 2021. Plant–microbiome interactions for sustainable

agriculture: a review. **Physiol. Mol. Biol. Plants** 27:165–179.

https://doi.org/10.1007/s12298-021-00927-1.

Hakim, S., Naqqash, T., Nawaz, M. S., Laraib, I., Siddique, M. J., Zia, R., Mirza, M. S., and Imran,

A. 2021. Rhizosphere engineering with plant growth-promoting microorganisms for

agriculture and ecological sustainability. **Front. Sustain. Food Syst.** 5:617157.

https://doi.org/10.3389/fsufs.2021.617157.

Hancock, J. T. 2020. Nitric oxide signaling in plants. **Plants** 9:1550.

https://doi.org/10.3390/plants9111550.

Hassani, M. A., Durán, P., and Hacquard, S. 2018. Microbial interactions within the plant

holobiont. **Microbiome** 6:58. https://doi.org/10.1186/s40168-018-0445-0.

Hayward, A. C., Fegan, N., Fegan, M., and Stirling, G. R. 2010. *Stenotrophomonas* and

Lysobacter: ubiquitous plant-associated gamma-proteobacteria of developing significance in

applied microbiology. **J. Appl. Microbiol.** 108:756–770. https://doi.org/10.1111/j.1365-

2672.2009.04471.x.

Helmstetter, A. J., Oztolan-Erol, N., Lucas, S. J., and Buggs, R. J. A. 2020. Genetic diversity and

domestication of hazelnut (*Corylus avellana* L.) in Turkey. **Plants, People, Planet** 2:326–

339. https://doi.org/10.1002/ppp3.10078.

Ho, C. H., and Tsay, Y. F. 2010. Nitrate, ammonium, and potassium sensing and signaling. **Curr.**

Opin. Plant Biol. 13:604–610. https://doi.org/10.1016/j.pbi.2010.08.005.

Hyatt, D., Chen, G.-L., LoCascio, P. F., Land, M. L., Larimer, F. W., and Hauser, L. J. 2010.

Prodigal: prokaryotic gene recognition and translation initiation site identification. **BMC**

Bioinformatics 11:119. https://doi.org/10.1186/1471-2105-11-119.

Jacobs, D. C., Revord, R. S., Capik, J. M., Mehlenbacher, S. A., and Molnar, T. J. 2024a. Variable

response of eastern filbert blight resistance sources in New Jersey. **Front. Plant Sci.** 15.

https://doi.org/10.3389/fpls.2024.1419265.

Jacobs, D. C., Revord, R. S., Capik, J. M., and Molnar, T. J. 2024b. Eastern filbert blight

resistant *Corylus avellana* identified from 20 years of germplasm introduction and

evaluation at Rutgers University, New Jersey, USA. **Front. Plant Sci.** 15.

https://doi.org/10.3389/fpls.2024.1502392.

Jeger, M., Bragard, C., Caffier, D., Candresse, T., Chatzivassiliou, E., Dehnen-Schmutz, K.,

Gilioli, G., Grégoire, J., Jaques Miret, J. A., MacLeod, A., Navajas Navarro, M., Niere, B.,

Parnell, S., Potting, R., Rafoss, T., Rossi, V., Urek, G., Van Bruggen, A., Van der Werf, W.,

West, J., Winter, S., Boberg, J., Gonthier, P., and Pautasso, M. 2018. Pest categorization of

Anisogramma anomala. **EFSA J.** 16:e05184. https://doi.org/10.2903/j.efsa.2018.5184.

Jiang, H., Xu, X., Fang, Y., Ogunyemi, S. O., Ahmed, T., Li, X., Yang, Y., Yan, C., Chen, J., and

Li, B. 2023. Metabarcoding reveals response of rice rhizosphere bacterial community to rice

bacterial   leaf   blight.   **Microbiol.**   **Res.**   270:127344.

https://doi.org/10.1016/j.micres.2023.127344.

Jones, D. L., and Hinsinger, P. 2008. The rhizosphere: complex by design. **Plant Soil** 312:1–6.

https://doi.org/10.1007/s11104-008-9774-2.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., and Jermiin, L. S. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. **Nat. Methods**

14:587–589. https://doi.org/10.1038/nmeth.4285.

Kang, D. D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., and Wang, Z. 2019. MetaBAT 2: an

adaptive binning algorithm for robust and efficient genome reconstruction from metagenome

assemblies. **PeerJ.** 7:e7359. https://doi.org/10.7717/peerj.7359.

Kassambara, A. 2025. **ggpubr: 'ggplot2' based publication ready plots**. https://CRAN.R-

project.org/package=ggpubr.

Kebert, M., Kostić, S., Zlatković, M., Stojnic, S., Čapelja, E., Zorić, M., Kiprovski, B., Budakov,

D., and Orlović, S. 2022. Ectomycorrhizal fungi modulate biochemical response against

powdery   mildew   disease   in   *Quercus*   *robur*   L.   **Forests**   13:1491.

https://doi.org/10.3390/f13091491.

Khan, M., Al Azawi, T. N. I., Pande, A., Mun, B.-G., Lee, D.-S., Hussain, A., Lee, B.-H., and Yun,

B.-W. 2021. The role of nitric oxide-induced ATILL6 in growth and disease resistance in

Arabidopsis thaliana. **Front. Plant Sci.** 12. https://doi.org/10.3389/fpls.2021.685156.

Kilic-Ekici, O., and Yuen, G. Y. 2003. Induced resistance as a mechanism of biological control by

Lysobacter   *enzymogenes*   strain   C3**.**   **Phytopathology**   93:1103–1110.

https://doi.org/10.1094/PHYTO.2003.93.9.1103.

Künstler, A., Gullner, G., Ádám, A. L., Kolozsváriné Nagy, J., and Király, L. 2020. The versatile

roles of sulfur-containing biomolecules in plant defense—a road to disease resistance. **Plants**

9:1705. https://doi.org/10.3390/plants9121705.

Kurtz, S., Phillippy, A., Delcher, A. L., Smoot, M., Shumway, M., Antonescu, C., and Salzberg, S.

L. 2004. Versatile and open software for comparing large genomes. **Genome Biol.** 5:R12.

https://doi.org/10.1186/gb-2004-5-2-r12.

Kusuma, A. B., Putra, K. E., Vanggy, L. R., Loh, J., Nouioui, I., and Goodfellow, M. 2022.

Actinospica acidithermotolerans sp. nov., a novel actinomycete isolated from sediment from

an Indonesian hot spring. **Arch. Microbiol.** 204:518. https://doi.org/10.1007/s00203-022-

03058-7.

Lahti, L., and Shetty, S. 2019. **microbiome R package**. http://microbiome.github.io

Langmead, B., and Salzberg, S. L. 2012. Fast gapped-read alignment with Bowtie 2. **Nat. Methods**

9:357–359. https://doi.org/10.1038/nmeth.1923.

Lee, S., Yap, M., Behringer, G., Hung, R., and Bennett, J. W. 2016. Volatile organic compounds

emitted by *Trichoderma* species mediate plant growth. **Fungal Biol. Biotechnol.** 3:7.

https://doi.org/10.1186/s40694-016-0025-7.

Lei, S., Xu, X., Cheng, Z., Xiong, J., Ma, R., Zhang, L., Yang, X., Zhu, Y., Zhang, B., and Tian, B.

2019. Analysis of the community composition and bacterial diversity of the rhizosphere

microbiome across different plant taxa. **MicrobiologyOpen** 8:e00762.

https://doi.org/10.1002/mbo3.762.

Letunic, I., and Bork, P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the

phylogenetic tree display and annotation tool. **Nucleic Acids Res.** 52:W78–W82.

https://doi.org/10.1093/nar/gkae268.

Li, D., Liu, C. M., Luo, R., Sadakane, K., and Lam, T.-W. 2015. MEGAHIT: an ultra-fast single-

node solution for large and complex metagenomics assembly via succinct de Bruijn graph.

Bioinformatics 31:1674–1676. https://doi.org/10.1093/bioinformatics/btv033.

Li, H. 2018. Minimap2: pairwise alignment for nucleotide sequences. **Bioinformatics** 34:3094–

3100. https://doi.org/10.1093/bioinformatics/bty191.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., and 1000 Genome Project Data Processing Subgroup. 2009. The sequence

alignment/map format and SAMtools. **Bioinformatics** 25:2078–2079.

https://doi.org/10.1093/bioinformatics/btp352.

Li, Z., Li, J., Liu, G., Li, Y., Wu, X., Liang, J., Wang, Z., Chen, Q., and Peng, F. 2025. Isolation,

characterization and growth-promoting properties of phosphate-solubilizing bacteria (PSBs)

derived from peach tree rhizosphere. **Microorganisms** 13:718.

https://doi.org/10.3390/microorganisms13040718.

Lin, H., and Peddada, S. D. 2024. Multigroup analysis of compositions of microbiomes with

covariate adjustments and repeated measures. **Nat. Methods** 21:83–91.

https://doi.org/10.1038/s41592-023-02092-7.

Liu, F., Hewezi, T., Lebeis, S. L., Pantalone, V., Grewal, P. S., and Staton, M. E. 2019. Soil

indigenous microbiome and plant genotypes cooperatively modify soybean rhizosphere

microbiome assembly. **BMC Microbiol.** 19:201. https://doi.org/10.1186/s12866-019-1572-

x.

Liu, H., Li, J., Carvalhais, L. C., Percy, C. D., Prakash Verma, J., Schenk, P. M., and Singh, B. K.

2021. Evidence for the plant recruitment of beneficial microbes to suppress soil-borne

pathogens. **New Phytol.** 229:2873–2885. https://doi.org/10.1111/nph.17057.

Liu, L., Ma, L., Zhu, M., Liu, B., Liu, X., and Shi, Y. 2023b. Rhizosphere microbial community

assembly and association networks strongly differ based on vegetation type at a local

environment   scale.   **Front.   Microbiol.**   14:1129471.

https://doi.org/10.3389/fmicb.2023.1129471.

Liu, S., Tao, C., Zhang, L., Wang, Z., Xiong, W., Xiang, D., Sheng, O., Wang, J., Li, R., Shen, Z.,

Li, C., Shen, Q., and Kowalchuk, G. A. 2023a. Plant pathogen resistance is mediated by

recruitment   of   specific   rhizosphere   fungi.   **ISME   J.**   17:931–942.

https://doi.org/10.1038/s41396-023-01406-z.

Lopes, A. C. de A., Mendes, L. W., Brito, K. A. da S. B., da Silva, J. L., Rocha, S. M. B., Antunes,

J. E. L., Oliveira, L. M. de S., Melo, V. M. M., Oliveira, F. A. S., Pereira, A. P. de A., Costa,

G. do N., da Silva, V. B., Gomes, R. L. F., de Alcantara Neto, F., and Araujo, A. S. F. 2023.

Rhizospheric microbial community in plant species from the *Phaseolus* genus. **Appl. Soil**

Ecol. 182:104731. https://doi.org/10.1016/j.apsoil.2022.104731.

Lotfi, N., Soleimani, A., Çakmakçı, R., Vahdati, K., and Mohammadi, P. 2022. Characterization of

plant growth-promoting rhizobacteria (PGPR) in Persian walnut associated with drought

stress tolerance. **Sci. Rep.** 12:12725. https://doi.org/10.1038/s41598-022-16852-6.

Lu, J., Breitwieser, F., Thielen, P., and Salzberg, S. 2017. Bracken: estimating species abundance

in metagenomics data. **PeerJ Comput. Sci.** 3:e104. https://doi.org/10.7717/peerj-cs.104.

Luciani, E., Palliotti, A., Tombesi, S., Gardi, T., Micheli, M., Berrios, J. G., Zadra, C., and Farinelli,

D. 2019. Mitigation of multiple summer stresses on hazelnut (*Corylus avellana* L.): effects

of the new arbuscular mycorrhiza *Glomus iranicum tenuihypharum* sp. *nova*. **Sci. Hortic.**

257:108659. https://doi.org/10.1016/j.scienta.2019.108659.

Luo, L., Zhang, J., Ye, C., Li, S., Duan, S., Wang, Z., Huang, H., Liu, Y., Deng, W., Mei, X.,

He, X., Yang, M., Zhu, S. 2022. Foliar pathogen infection manipulates soil health through

root exudate-modified rhizosphere microbiome. **Microbiol Spectr** 10:e02418-22.

https://doi.org/10.1128/spectrum.02418-22

Ma, W., Yang, Z., Liang, L., Ma, Q., Wang, G., and Zhao, T. 2021. Characteristics of the fungal

communities and co-occurrence networks in hazelnut tree root endospheres and rhizosphere

soil. **Front. Plant Sci.** 12. https://doi.org/10.3389/fpls.2021.749871.

Majer, H. M., Ehrlich, R. L., Ahmed, A., Earl, J. P., Ehrlich, G. D., and Beld, J. 2021. Whole

genome sequencing of *Streptomyces actuosus* ISP-5337, *Streptomyces sioyaensis* B-5408,

and *Actinospica acidiphila* B-2296 reveals secondary metabolomes with antibiotic potential.

Biotechnol. Rep. 29:e00596. https://doi.org/10.1016/j.btre.2021.e00596.

Martinez Arbizu, P. 2017. **pairwiseAdonis: pairwise multilevel comparison using adonis**.

https://github.com/pmartinezarbizu/pairwiseAdonis.

Matsen, F. A., Kodner, R. B., and Armbrust, E. V. 2010. pplacer: linear time maximum-likelihood

and Bayesian phylogenetic placement of sequences onto a fixed reference tree. **BMC**

Bioinformatics 11:538. https://doi.org/10.1186/1471-2105-11-538.

McMurdie, P. J., and Holmes, S. 2013. phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. **PLOS ONE** 8:e61217.

https://doi.org/10.1371/journal.pone.0061217.

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von Haeseler, A.,

and Lanfear, R. 2020. IQ-TREE 2: new models and efficient methods for phylogenetic

inference in the genomic era. **Mol. Biol. Evol.** 37:1530–1534.

https://doi.org/10.1093/molbev/msaa015.

Molnar, T. J. 2011. Corylus. Pages 15–48 in: ***Wild Crop Relatives: Genomic and Breeding***

Resources: Forest Trees. C. Kole, ed. Springer, Heidelberg, Germany.

842 https://doi.org/10.1007/978-3-642-21250-5_2.

843 Mur, L. A. J., Simpson, C., Kumari, A., Gupta, A. K., and Gupta, K. J. 2017. Moving nitrogen to

844 the centre of plant defence against pathogens. **Ann. Bot.** 119:703–709.

845 https://doi.org/10.1093/aob/mcw179.

846 Nasir, N. N. M., Ho, C.-L., Lamasudin, D. U., and Saidi, N. B. 2020. Nitric oxide improves

847 tolerance to *Fusarium oxysporum* f. sp. cubense tropical race 4 in banana. **Physiol. Mol.**

848 **Plant Pathol.** 111:101503. https://doi.org/10.1016/j.pmpp.2020.101503.

849 Navarro-Muñoz, J. C., Selem-Mojica, N., Mullowney, M. W., Kautsar, S. A., Tryon, J. H.,

850 Parkinson, E. I., De Los Santos, E. L. C., Yeong, M., Cruz-Morales, P., Abubucker, S.,

851 Roeters, A., Lokhorst, W., Fernandez-Guerra, A., Cappelini, L. T. D., Goering, A. W.,

852 Thomson, R. J., Metcalf, W. W., Kelleher, N. L., Barona-Gomez, F., and Medema, M. H.

853 2020. A computational framework to explore large-scale biosynthetic diversity. **Nat. Chem.**

854 **Biol.** 16:60–68. https://doi.org/10.1038/s41589-019-0400-9.

855 Nissen, J. N., Johansen, J., Allesøe, R. L., Sønderby, C. K., Armenteros, J. J. A., Grønbech, C. H.,

856 Jensen, L. J., Nielsen, H. B., Petersen, T. N., Winther, O., and Rasmussen, S. 2021. Improved

857 metagenome binning and assembly using deep variational autoencoders. **Nat. Biotechnol.**

858 39:555–560. https://doi.org/10.1038/s41587-020-00777-4.

859 Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R.

860 B., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M.,

861 Bolker, B., Borcard, D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S.,

862 Evangelista, H. B. A., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M.

863 O., Lahti, L., McGlinn, D., Ouellette, M.-H., Ribeiro Cunha, E., Smith, T., Stier, A., Ter

864 Braak, C. J. F., and Weedon, J. 2024. **vegan: Community Ecology Package**.

https://CRAN.R-project.org/package=vegan

Olm, M. R., Brown, C. T., Brooks, B., and Banfield, J. F. 2017. dRep: a tool for fast and accurate

genomic comparisons that enables improved genome recovery from metagenomes through

de-replication. **ISME J.** 11:2864–2868. https://doi.org/10.1038/ismej.2017.126.

Ondov, B. D., Bergman, N. H., and Phillippy, A. M. 2011. Interactive metagenomic visualization

in a web browser. **BMC Bioinformatics** 12:385. https://doi.org/10.1186/1471-2105-12-385.

Ondov, B. D., Treangen, T. J., Melsted, P., Mallonee, A. B., Bergman, N. H., Koren, S., and

Phillippy, A. M. 2016. Mash: fast genome and metagenome distance estimation using

MinHash. **Genome Biol.** 17:132. https://doi.org/10.1186/s13059-016-0997-x.

Orozco-Mosqueda, Ma. D. C., Fadiji, A. E., Babalola, O. O., Glick, B. R., and Santoyo, G. 2022.

Rhizobiome engineering: unveiling complex rhizosphere interactions to enhance plant

growth and health. **Microbiol. Res.** 263:127137.

https://doi.org/10.1016/j.micres.2022.127137.

Pan, S., Zhao, X.-M., and Coelho, L. P. 2023. SemiBin2: self-supervised contrastive learning leads

to better MAGs for short- and long-read sequencing. **Bioinformatics** 39:i21–i29.

https://doi.org/10.1093/bioinformatics/btad209.

Pantigoso, H. A., Newberger, D., and Vivanco, J. M. 2022. The rhizosphere microbiome: plant–

microbial interactions for resource acquisition. **J. Appl. Microbiol.** 133:2864–2876.

https://doi.org/10.1111/jam.15686.

Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., and Tyson, G. W. 2015. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. **Genome Res.** 25:1043–1055. https://doi.org/10.1101/gr.186072.114.

Pedersen, T. L. 2019. **patchwork: The Composer of Plots**. http://CRAN.R-

project.org/package=patchwork

Peiffer, J. A., Spor, A., Koren, O., Jin, Z., Tringe, S. G., Dangl, J. L., Buckler, E. S., and Ley, R. E.

2013. Diversity and heritability of the maize rhizosphere microbiome under field conditions.

Proc. Natl. Acad. Sci. 110:6548–6553. https://doi.org/10.1073/pnas.1302837110.

Pinkerton, J. N. 1995. Infection of European hazelnut by *Anisogramma anomala*: ascospore

adhesion, mode of penetration of immature shoots, and host response. **Phytopathology**

85:1260. https://doi.org/10.1094/Phyto-85-1260.

Poomthongdee, N., Duangmal, K., and Pathom-aree, W. 2015. Acidophilic actinomycetes from

rhizosphere soil: diversity and properties beneficial to plants. **J. Antibiot.** 68:106–114.

https://doi.org/10.1038/ja.2014.117.

Puopolo, G., Giovannini, O., and Pertot, I. 2014. *Lysobacter capsici* AZ78 can be combined with

copper to effectively control *Plasmopara viticola* on grapevine. **Microbiol. Res.** 169:633–

642. https://doi.org/10.1016/j.micres.2013.09.013.

Qi, G., Ma, G., Chen, S., Lin, C., and Zhao, X. 2019. Microbial network and soil properties are

changed in bacterial wilt-susceptible soil. **Appl. Environ. Microbiol.** 85:e00162-19.

https://doi.org/10.1128/AEM.00162-19.

Qiao, H., Gao, D., and Yuan, T. 2023. Differences in rhizosphere soil fungal communities of wild

and cultivated *Paeonia ludlowii* species. **Front. Plant Sci.** 14.

https://doi.org/10.3389/fpls.2023.1194598.

Ranjan, A., Rajput, V. D., Prazdnova, E. V., Gurnani, M., Bhardwaj, P., Sharma, S., Sushkova, S.,

Mandzhieva, S. S., Minkina, T., Sudan, J., Zargar, S. M., Chauhan, A., and Jindal, T. 2023.

Nature's antimicrobial arsenal: non-ribosomal peptides from PGPB for plant pathogen

biocontrol. **Fermentation** 9:597. https://doi.org/10.3390/fermentation9070597.

R Core Team. 2025. The R Project for Statistical Computing. https://www.R-project.org/

Revord, R. S., Lovell, S. T., Capik, J. M., Mehlenbacher, S. A., and Molnar, T. J. 2020. Eastern

filbert blight resistance in American and interspecific hybrid hazelnuts. **J. Am. Soc. Hortic.**

Sci. 145:162–173. https://doi.org/10.21273/JASHS04732-19.

Rodriguez-Mena, S., Camacho, M., de los Santos, B., Miranda, L., and Camacho-Sanchez, M.

2022. Microbiota modulation in blueberry rhizosphere by biocontrol bacteria. **Microbiol.**

Res. 13:809–824. https://doi.org/10.3390/microbiolres13040057.

Rostamikia, Y., Tabari Kouchaksaraei, M., Asgharzadeh, A., and Rahmani, A. 2017. Effect of

growth promoting rhizobacteria on growth and nutrient elements of common hazelnut

(*Corylus avellana* L.) seedlings in Ardabil Fandoqlou nursery. **Iran. J. For. Poplar Res.**

25:126–116. https://doi.org/10.22092/ijfpr.2017.109781.

Ruparelia, J., Rabari, A., Mitra, D., Panneerselvam, P., Das-mohapatra, P. K., and Jha, C. K. 2022.

Efficient applications of bacterial secondary metabolites for management of biotic stress in

plants. **Plant Stress** 6:100125. https://doi.org/10.1016/j.stress.2022.100125.

Santoyo, G. 2022. How plants recruit their microbiome? New insights into beneficial interactions.

J. Adv. Res. 40:45–58. https://doi.org/10.1016/j.jare.2021.11.020.

Shaffer, M., Borton, M. A., McGivern, B. B., Zayed, A. A., La Rosa, S. L., Solden, L. M., Liu, P.,

Narrowe, A. B., Rodríguez-Ramos, J., Bolduc, B., Gazitúa, M. C., Daly, R. A., Smith, G. J.,

Vik, D. R., Pope, P. B., Sullivan, M. B., Roux, S., and Wrighton, K. C. 2020. DRAM for

distilling microbial metabolism to automate the curation of microbiome function. **Nucleic**

Acids Res. 48:8883–8900. https://doi.org/10.1093/nar/gkaa621.

Sharrar, A. M., Crits-Christoph, A., Méheust, R., Diamond, S., Starr, E. P., and Banfield, J. F. 2020.

Bacterial secondary metabolite biosynthetic potential in soil varies with phylum, depth, and

vegetation type. **mBio** 11:e00416-20. https://doi.org/10.1128/mBio.00416-20.

Sieber, C. M. K., Probst, A. J., Sharrar, A., Thomas, B. C., Hess, M., Tringe, S. G., and Banfield,

J. F. 2018. Recovery of genomes from metagenomes via a dereplication, aggregation and

scoring strategy. **Nat. Microbiol.** 3:836–843. https://doi.org/10.1038/s41564-018-0171-1.

Sillo, F., Brunetti, C., Marroni, F., Vita, F., dos Santos Nascimento, L. B., Vizzini, A., Mello, A.,

and Balestrini, R. 2022. Systemic effects of *Tuber melanosporum* inoculation in two *Corylus*

avellana genotypes. **Tree Physiol.** 42:1463–1480. https://doi.org/10.1093/treephys/tpac012.

Sood, M., Kapoor, D., Kumar, V., Sheteiwy, M. S., Ramakrishnan, M., Landi, M., Araniti, F., and

Sharma, A. 2020. Trichoderma: The "secrets" of a multitalented biocontrol agent. **Plants**

9:762. https://doi.org/10.3390/plants9060762.

Steinegger, M., and Söding, J. 2017. MMseqs2 enables sensitive protein sequence searching for

the analysis of massive data sets. **Nat. Biotechnol.** 35:1026–1028.

https://doi.org/10.1038/nbt.3988.

Sun, C., Zhang, Y., Liu, L., Liu, X., Li, B., Jin, C., and Lin, X. 2021. Molecular functions of nitric

oxide and its potential applications in horticultural crops. **Hortic. Res.** 8:1–17.

https://doi.org/10.1038/s41438-021-00500-7.

Sun, Y., Wang, M., Mur, L. A. J., Shen, Q., and Guo, S. 2020. Unravelling the roles of nitrogen

nutrition in plant disease defenses. **Int. J. Mol. Sci.** 21:572.

https://doi.org/10.3390/ijms21020572.

Thepbandit, W., and Athinuwat, D. 2024. Rhizosphere microorganisms supply availability of soil

nutrients and induce plant defense. **Microorganisms** 12:558.

https://doi.org/10.3390/microorganisms12030558.

Thompson, M. M., Lagerstedt, H. B., and Mehlenbacher, S. A. 1996. Hazelnuts. Pages 125–184

in: **Fruit Breeding, vol. III Nuts.** J. Janick, and N. Moore, eds. John Wiley and Sons Inc,

New York, U.S.A.

Wang, Y., Yang, Q., Tosa, Y., Nakayashiki, H., and Mayama, S. 2005. Nitric oxide-overproducing

transformants of *Pseudomonas fluorescens* with enhanced biocontrol of tomato bacterial

wilt. J. **Gen. Plant Pathol.** 71:33–38. https://doi.org/10.1007/s10327-004-0157-0.

Wattenburger, C. J., Halverson, L. J., and Hofmockel, K. S. 2019. Agricultural management affects

root-associated microbiome recruitment over maize development. **Phytobiomes J.** 3:260–

272. https://doi.org/10.1094/PBIOMES-03-19-0016-R.

Wickham, H. 2009. **ggplot2: Elegant Graphics for Data Analysis**. Springer, New York, U.S.A.

https://doi.org/10.1007/978-0-387-98141-3

Wickham, H., François, R., Henry, L., and Müller, K. 2018. **dplyr: A Grammar of Data**

Manipulation. https://CRAN.Rproject.org/package=dplyr.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan L. D., François, R., Grolemund, G.,

Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T. L., Miller, E., Bache S. M.,

Müller, K., Ooms, J., Robinson, D., Seidel, D. P., Spinu, V., Takahashi, K., Vaughan, D.,

Wilke, C., Woo, K., Yutani, H. 2019. Welcome to the tidyverse. **Journal of Open Source**

Software 4:1686. https://doi.org/10.21105/joss.01686>

Wood, D. E., Lu, J., and Langmead, B. 2019. Improved metagenomic analysis with Kraken

Genome Biol. 20:257. https://doi.org/10.1186/s13059-019-1891-0.

Wu, Y.-W., Simmons, B. A., and Singer, S. W. 2016. MaxBin 2.0: an automated binning algorithm

to recover genomes from multiple metagenomic datasets. **Bioinformatics** 32:605–607.

https://doi.org/10.1093/bioinformatics/btv638.

Yang, J., Wei, S., Su, D., Zhang, Z., Chen, S., Luo, Z., Shen, X., Lai, Y., Jamil, A., Tong, J., and

Cui, X. 2020. Comparison of the rhizosphere soil microbial community structure and

diversity between powdery mildew-infected and noninfected strawberry plants in a

greenhouse by high-throughput sequencing technology. **Curr. Microbiol.** 77:1724–1736.

https://doi.org/10.1007/s00284-020-01948-x.

Yu, Y., Gui, Y., Li, Z., Jiang, C., Guo, J., and Niu, D. 2022. Induced systemic resistance for

improving plant immunity by beneficial microbes. **Plants** 11:386.

https://doi.org/10.3390/plants11030386.

Yue, H., Yue, W., Jiao, S., Kim, H., Lee, Y.-H., Wei, G., Song, W., and Shu, D. 2023. Plant

domestication shapes rhizosphere microbiome assembly and metabolic functions.

Microbiome 11:70. https://doi.org/10.1186/s40168-023-01513-1.

Zhang, F., Wen, S., Wang, B., Zhang, Z., Liu, F., Ye, T., Wang, K., Hu, H., Yang, X., and Fang, W.

2023. Biocontrol potential of *Streptomyces odonnellii* SZF-179 toward *Alternaria alternata*

to control pear black spot disease. **Int. J. Mol. Sci.** 24:17515.

https://doi.org/10.3390/ijms242417515.

Zhang, J., Cook, J., Nearing, J. T., Zhang, J., Raudonis, R., Glick, B. R., Langille, M. G. I., and

Cheng, Z. 2021. Harnessing the plant microbiome to promote the growth of agricultural

crops. **Microbiol. Res.** 245:126690. https://doi.org/10.1016/j.micres.2020.126690.

Zhang, M., Chai, L., Huang, M., Jia, W., Guo, J., and Huang, Y. 2020. Deciphering the archaeal

communities in tree rhizosphere of the Qinghai-Tibetan plateau. **BMC Microbiol.** 20:235.

https://doi.org/10.1186/s12866-020-01913-5.

1001

 FIGURE LEGENDS

Figure 1. Diversity metrics of the metagenomes. (A, **B)** Alpha diversity values calculated using

the (A) Shannon Diversity Index (Kruskal-Wallis rank sum test, p-value = 0.16) or (B) Simpson

Diversity Index [1-D] (Kruskal-Wallis rank sum test, p-value = 0.06) are shown. Alpha diversity

was calculated based on species level taxonomic classification of assembled contigs (minimum

length of 500 bp). (**C**) A Principal Coordinate Analysis (PCoA) plot based on Bray-Curtis

dissimilarity values is shown (PERMANOVA, p-value < 0.001), with pairwise adonis tests

indicating that the bulk soil was statistically different from the rhizospheres of the native

(p.adjusted = 0.048), susceptible (p.adjusted = 0.047), and resistant (p.adjusted = 0.047) varieties.

1011

Figure 2. Relative abundance of phyla across the metagenomes. The abundance of (**A**) bacterial

phyla, (**B**) archaeal phyla, and (**C**) eukaryotic phyla, displayed as a percentage of the total bacteria,

archaea, and eukaryotes, respectively. The rows represent each of the metagenomic samples,

colour coded by group, while the columns represent taxa with an average relative abundance of at

least 1% of bacteria, archaea, or eukaryotes in at least one sample. The legend to the right of the

figure applies to all three panels.

1018

Figure 3. Relative abundance of genera across the metagenomes. The abundance of (**A**)

bacterial genera, (**B**) archaeal genera, and (**C**) eukaryotic genera, displayed as a percentage of the

total bacteria, archaea, and eukaryotes, respectively. The rows represent each of the metagenomic

samples, colour coded by group, while the columns represent taxa with an average relative

abundance of at least 1% of bacteria, archaea, or eukaryotes in at least one sample. The size of the

dots corresponds is proportional to the relative abundance of the taxon in a given sample. The

legend below the figure applies to all three panels.

1026

1027   **Figure 4. Global comparison of the metabolic potential of the metagenomes.** A Principal

1028   Coordinate Analysis (PCoA) plot based on Bray-Curtis dissimilarity values calculated from the

1029   KEGG annotations produced for each metagenome using DRAM. While statistically significant

1030   global differences were observed (PERMANOVA, p-value = 0.031), no statistically significant

1031   differences between groups were detected using pairwise adonis tests.

1032

Figure 5. Putative biosynthetic gene clusters detected in the metagenomes. (**A**) A heatmap

showing the number of predicted biosynthetic gene clusters summarized by biosynthetic gene

cluster classification. (**B**) A heatmap showing the number of predicted biosynthetic gene clusters

summarized by predicted product. (**A**, **B**) The absence of a box indicates that no genes predicted

to be associated with a particular biosynthetic gene cluster class or product were detected in that

metagenome. The colour of a box represents the number of biosynthetic gene clusters detected as

belonging to a given biosynthetic gene cluster class or product, with deeper red indicating higher

counts.

1041

Figure 6. Unrooted phylogeny of the bacterial MAGs generated in this study. An unrooted,

maximum likelihood phylogeny of the 41 high-quality metagenome-assembled genomes (MAGs)

generated in this study, prepared from a concatenated alignment of the bac120 gene set of the

Genome Taxonomy Database. Taxa names and branches are colour coded based on phylum. The

scale represents the mean number of nucleotide substitutions per site. The numbers on the nodes

indicate the SH-aLRT support values (top numbers) and the ultra-fast jackknife values using a

40% resampling rate (bottom numbers), both calculated from 1000 replicates; values are only

shown at nodes where at least one value is below 95.

1050

1051

Figure 1. Diversity metrics of the metagenomes. (**A**, **B**) Alpha diversity values calculated using
the (A) Shannon Diversity Index (Kruskal-Wallis rank sum test, p-value = 0.16) or (B) Simpson
Diversity Index [1-D] (Kruskal-Wallis rank sum test, p-value = 0.06) are shown. Alpha diversity
was calculated based on species level taxonomic classification of assembled contigs (minimum
length of 500 bp). (**C**) A Principal Coordinate Analysis (PCoA) plot based on Bray-Curtis
dissimilarity values is shown (PERMANOVA, p-value < 0.001), with pairwise adonis tests
indicating that the bulk soil was statistically different from the rhizospheres of the native
(p.adjusted = 0.048), susceptible (p.adjusted = 0.047), and resistant (p.adjusted = 0.047) varieties.

1060

Figure 2. Relative abundance of phyla across the metagenomes. The abundance of (**A**) bacterial phyla, (**B**) archaeal phyla, and (**C**) eukaryotic phyla, displayed as a percentage of the total bacteria, archaea, and eukaryotes, respectively. The rows represent each of the metagenomic samples, colour coded by group, while the columns represent taxa with an average relative abundance of at least 1% of bacteria, archaea, or eukaryotes in at least one sample. The legend to the right of the figure applies to all three panels.

Figure 3. Relative abundance of genera across the metagenomes. The abundance of (**A**) bacterial genera, (**B**) archaeal genera, and (**C**) eukaryotic genera, displayed as a percentage of the total bacteria, archaea, and eukaryotes, respectively. The rows represent each of the metagenomic samples, colour coded by group, while the columns represent taxa with an average relative abundance of at least 1% of bacteria, archaea, or eukaryotes in at least one sample. The size of the dots corresponds is proportional to the relative abundance of the taxon in a given sample. The legend below the figure applies to all three panels.

1075

1076   **Figure 4. Global comparison of the metabolic potential of the metagenomes.** A Principal
1077   Coordinate Analysis (PCoA) plot based on Bray-Curtis dissimilarity values calculated from the
1078   KEGG annotations produced for each metagenome using DRAM. While statistically significant
1079   global differences were observed (PERMANOVA, p-value = 0.031), no statistically significant
1080   differences between groups were detected using pairwise adonis tests.

Figure 5. Putative biosynthetic gene clusters detected in the metagenomes. (**A**) A heatmap showing the number of predicted biosynthetic gene clusters summarized by biosynthetic gene cluster classification. (**B**) A heatmap showing the number of predicted biosynthetic gene clusters summarized by predicted product. (**A**, **B**) The absence of a box indicates that no genes predicted to be associated with a particular biosynthetic gene cluster class or product were detected in that metagenome. The colour of a box represents the number of biosynthetic gene clusters detected as belonging to a given biosynthetic gene cluster class or product, with deeper red indicating higher counts.

1098

Figure 6. Unrooted phylogeny of the bacterial MAGs generated in this study. An unrooted, maximum likelihood phylogeny of the 41 high-quality metagenome-assembled genomes (MAGs) generated in this study, prepared from a concatenated alignment of the bac120 gene set of the Genome Taxonomy Database. Taxa names and branches are colour coded based on phylum. The scale represents the mean number of nucleotide substitutions per site. The numbers on the nodes indicate the SH-aLRT support values (top numbers) and the ultra-fast jackknife values using a 40% resampling rate (bottom numbers), both calculated from 1000 replicates; values are only shown at nodes where at least one value is below 95.