Aβ-42 sidechain deamidation at Q15 and N27 modulate protein aggregation and microglial responses via altered cytokine production and CD68 expression

Martin N. Griffin[1+], Daniel M. Dinakarapandian[2+], Chenxing Li[3], Pranay Ramaswamy[4], Sia J. Saheba[3], Janie Lee[3], Tarunya Rao Sudarshan[2], Maria Sajimon[5], Christopher J. Wheeler[5,6,7,8], Jevgenij A. Raskatov[5], Anant K. Paravastu[2,9], Levi B. Wood[1,3,9*]

[1] George W. Woodruff School of Mechanical Engineering, Georgia Institute of Technology, Atlanta, GA, USA

[2] School of Chemical and Biomolecular Engineering, Georgia Institute of Technology, Atlanta, GA, USA

[3] Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of Technology and Emory University, Atlanta, GA, USA

[4] School of Chemistry and Biochemistry, Georgia Institute of Technology, Atlanta, GA, USA

[5] Department of Chemistry and Biochemistry, University of California Santa Cruz, Santa Cruz, CA, USA

[6] T-NeuroDx, Albuquerque, NM, USA

[7] StemVax Therapeutics, a NovAccess Global Company, Chagrin Falls, OH, USA

[8] World Brain Mapping Foundation, Society for Brain Mapping & Therapeutics, Pacific Palisades, CA, USA

[9] Parker H. Petit Institute for Bioengineering and Bioscience, Georgia Institute of Technology, Atlanta, GA, USA

+ Equally contributing authors

* Corresponding Author

Running Title: Aβ-42 Deamidation Alters Aggregation and Microglial Function

Keywords: Amyloid beta, Alzheimer's disease, protein deamidation, aggregation, inflammation, cytokines

Abstract

The progressive aggregation of amyloid beta (Aβ) monomers into oligomers is a critical factor in Alzheimer's disease (AD) pathogenesis. Although mutated forms of Aβ have been shown to display altered aggregation dynamics, the specific effects of deamidated Aβ on microglial function remain understudied. Our research group previously found that the deamidated variant Aβ-42-N27D modified Aβ aggregation, reduced neurotoxicity, and reduced microglial reactivity, but the impact of Aβ-42 side chain deamidation in general on such parameters remained unclear. Here, we expanded on our prior work by investigating how two site-specific Aβ-42 mutations (Q15E & N27D), where neutral amide side chains are replaced with negatively charged carboxylic acids, affect aggregation and microglial immune response using a mouse microglial cell line. Size exclusion chromatography revealed that Aβ-42-Q15E and Aβ-42-N27D exhibit distinct aggregation profiles compared to Aβ-42 wild type (WT). Multiplexed analysis of 8 cytokines secreted into the culture medium revealed that Aβ-42-Q15E and Aβ-42-N27D

decrease the expression of inflammatory cytokines such as IL-6, IP-10, and MIP-1α relative to
Aβ-42-WT. Immunocytochemistry revealed that Aβ-42-Q15E and Aβ-42-N27D decrease CD68
expression relative to Aβ-42-WT. These findings demonstrate that deamidation significantly
alters Aβ-42 aggregation and microglial activation, suggesting structural modifications to Aβ-42
modulate inflammatory signaling in AD. This work provides a foundation for future studies on
Aβ-42 post-translational modifications as potential therapeutic targets in AD.
51
Introduction
Alzheimer's disease (AD) is a progressive neurodegenerative disorder that affects millions of
people each year and is the leading cause of dementia globally [1]. Accumulation of Aβ peptides
into extracellular plaques is a defining hallmark of AD pathology [2]. While it remains unclear
whether Aβ plaques directly drive pathology, Aβ oligomers as well as chronic
neuroinflammation have both been strongly implicated in AD pathogenesis [3, 4]. Microglia are
the resident immune cells of the brain and function to clear cellular debris, prune synapses, and
maintain homeostasis under healthy conditions [5]. While microglia are believed to provide early
protection during AD, microglial function becomes dysregulated during disease progression,
reducing their ability to clear Aβ aggregates and contributing to a state of chronic
neuroinflammation linked to cognitive decline [6, 7].
63
Given that Aβ aggregation state influences microglial activity [8], understanding how specific
structural modifications to Aβ impact microglial activation is critical for elucidating mechanisms
of inflammation in AD. In the healthy brain, microglia secrete neurotrophic factors that support
neuronal health [9]. However, in AD, oligomeric forms of Aβ deregulate signaling pathways
vital for cellular function and gene expression [10]. This dysregulation exacerbates chronic
inflammation and accelerates neurodegeneration in AD patients [11]. Aβ interacts with
microglial surface receptors, such as TREM2, CD36, and toll-like receptors, resulting in
dysregulated phospho-protein signaling cascades that drive expression of inflammatory
cytokines, including IL-1β, IL-6, and TNF-α [12].
73
In this study, we chose to focus on Aβ-42 as this form of Aβ is known to form more oligomers
and be more neurotoxic than Aβ-40 [13]. Although limited, previous studies have explored how
site-specific modification to Aβ variants influence aggregation and the resulting cellular
responses [14]. Our recent study examined Aβ-42 with a substitution at position 27, where
asparagine was replaced with aspartic acid (Aβ-42-N27D) and found that this variant exhibited
reduced aggregation compared to wild-type Aβ-42, and notably produced substantially lower
quantities of soluble oligomers [15]. In the current study, this result was successfully reproduced
within a distinct laboratory environment, further validating the impact of Aβ-42-N27D on Aβ
aggregation behavior. Our previous study also reported changes in microglial cytokine
expression in response to the N27D variant. However, aspects of microglial activation and

phagocytic function remained uncharacterized, providing the basis for our investigation into how
deamidated Aβ-42 affects aggregation and broader microglial response.
86
In this study, we investigated how site-specific deamidation of Aβ-42 influences its aggregation
kinetics and microglial immune response. We found that Q15E and N27D mutants of Aβ-42
exhibited reduced aggregation, lower expression of inflammatory cytokines, and diminished
CD68 expression compared to Aβ-42-WT. Our data indicate that Aβ-42 deamidation at either
N27 or Q15 modifies its aggregation and immune activation, highlighting deamidation as a
potentially general mechanism for attenuating Aβ-induced inflammation.
93
94 **Results**
95
Deamidation of Aβ-42 at Q15 and N27 Alters Aggregation
We began the current study by investigating how site-specific deamidation affects Aβ-42
aggregation. To do so, we used size exclusion chromatography (SEC) to monitor the aggregation
profiles of Aβ-42-Q15E, Aβ-42-N27D, and Aβ-42-WT over 24 hours. Freshly prepared
monomers of each variant were isolated using SEC (**Fig. 1A**), diluted to 100 μM, and incubated
at 37°C for 24 hours to allow aggregation to occur. The resulting aggregation profiles were then
analyzed by re-injection onto the SEC column.
The SEC elution profiles revealed distinct aggregation behaviors among the three variants (**Fig.
1B**). Aβ-42-WT displayed a pronounced oligomeric peak at approximately 9 ml elution volume,
indicating substantial formation of higher-order aggregates after 24 hours of incubation. The
deamidated variants, Aβ-42-N27D and Aβ-42-Q15E, showed reduced oligomeric peak
formation, with peak heights approximately 75% lower than the wild-type variant. The
monomeric peaks (eluting at ~14 - 15 ml) remained prominent for both mutants, suggesting that
deamidation inhibits the progression from monomeric to oligomeric states.

Figure 1. Deamidation of Aβ-42 at Q15 and N27 Leads to Less Oligomer Formation
A. Aβ-42 monomers isolation using Size Exclusion Chromatography. Crude Aβ-42 peptides were dissolved in 50mM NaOH and injected into SEC column at 0.5 mL/min flow rate. The monomers eluted at 14-15 mL (highlighted area in blue) were collected and diluted to the final concentration of 100 uM. The collected monomers were incubated at 37°C for 24 hours. **B**. SEC profiles showing difference in aggregates formed from incubated monomers of Aβ-42 WT, Aβ-42 N27D, and Aβ-42 Q15E after 24 hours.

Deamidation of Aβ-42 modulates cytokine expression in mouse microglial cell line.

Chronic microglial activation is a hallmark of AD pathology and is closely linked to Aβ oligomerization [16]. We therefore next investigated whether deamidated Aβ-42 variants would elicit distinct cytokine responses compared to Aβ-42-WT. To do so, we applied a physiologically relevant concentration (50nM [17, 18]) of Aβ-42-WT, Aβ-42-N27D, and Aβ-42-Q15E to mouse

microglial cells (MMCs) for 24 hours. We then quantified 8 inflammatory cytokines associated with chronic inflammation secreted into the culture medium using a Luminex multiplexed immunoassay [19] (**Fig. 2A**).

Principal component analysis (PCA) identified principal components (PC1 and PC2) that captured the greatest variance in cytokine expression across samples, revealing clear separation between deamidated variants and Aβ-42-WT (**Fig. 2B**). Scores on PC1 showed that Aβ-42-WT clustered separately on the far right, both Aβ-42-N27D and Aβ-42-Q15E clustered in the middle, while Vehicle clustered on the left. Similarly, PC1 scores showed significant upregulation in Aβ-42-WT compared to Vehicle, Aβ-42-N27D, and Aβ-42-Q15E (**Fig. 2C**), indicating the effect of Aβ-42-WT was normalized with Aβ deamidation. PC1 consisted of a cytokine profile and the loadings on PC1 reflected the relative influence of each cytokine. MIP-1β, MIP-1α and IP-10 had the highest absolute loadings on PC1, indicating their most significant contribution to the PC1 score. Among the top contributors to PC1 (**Fig. 2D**), MIP-1β, MIP-1α, IP-10, and IL-6 were significantly upregulated in Aβ-42-WT compared to vehicle, Aβ-42-N27D, and Aβ-42-Q15E (**Fig. 2E**). Together, both Aβ-42-N27D and Aβ-42-Q15E elicited lower cytokine expression compared to Aβ-42-WT, suggesting that Aβ deamidation may mitigate microglial dysregulation in AD.

Figure 2. Deamidated Aβ modulates cytokine expression in mouse microglial cell line.
A. Multiplexed Luminex analysis of 8 cytokines (columns) expressed into the culture medium after 24 hours of incubation with 50nM of each Aβ variant (n=6-8 wells/group). Each row in the z-scored heat map denotes an individual culture sample well. **B**. Scores plot of principle component analysis (PCA). Each triangle represents a sample well. PC1 explained 54% of the variance, and PC2 explained an additional 17%. **C**. Bar plot of PC1 scores for each group. In all bar plots, dots denote individual samples.

151 **D**. PCA identified a variable, PC1, that captured the greatest axis of variation across experimental groups.
152 PC1 consisted of a profile of cytokine that were upregulated with Aβ-42-WT. **E**. Dot plots for individual
153 cytokines (mean ± SEM Wilcoxon rank sum test). In all barplots, dots denote individual samples. Each
154 dot denotes an individual sample well. mean±SEM. All p-values reflect Wilcoxon rank sum tests with
155 Bonferroni adjustment for multiple comparisons. a.u., arbitrary units.
156
157 **Aβ-42 deamidation attenuates CD68 expression in mouse microglial cell line.**
158
Aβ-42 oligomers have been shown to increase microglial activation and dysregulate
phagocytosis [20]. CD68, a lysosomal marker, is consistently shown to be upregulated in
activated microglia [21]. Therefore, we utilized immunocytochemistry (ICC) to stain for CD68
in MMCs treated with 50nM of Aβ-42-WT, Aβ-42-N27D, and Aβ-42-Q15E for 2 hours, as this
is a time period where studies have shown increased levels of Aβ-42 uptake [22]. ICC
examination of CD68 showed that CD68 labeling is reduced in the vehicle, Aβ-42-N27D, and
Aβ-42-Q15E (**Fig. 3A**). Quantification of CD68 fluorescent intensity normalized to DAPI cell
count revealed reduced CD68 expression in Aβ-42-N27D and Aβ-42-Q15E compared to Aβ-42-
WT (**Fig. 3B**), indicating that Aβ deamidation reduces its potency of inducing microglial
activity.
169

170
171 **Figure 3. Deamidation abrogates Aβ-induced CD68 expression in mouse microglial cell**
172 **line. A**. ICC for CD68 (red) and DAPI (blue) after 2 hours of incubation with 50nM of each Aβ variant
173 (scale bar: 200 μm, representative sections from n=12-13 sample well/group). **B**. Integrated CD68
174 florescent intensity normalized by number of nuclei (n=12-13, mean±SEM, Wilcoxon rank sum tests with
175 Bonferroni adjustment for multiple comparisons).
176
Discussion
Aβ oligomers are widely recognized as drivers of microglial activation, inflammation, and
neurodegeneration [16, 23]. Thus, Aβ mutants that promote oligomerization would be expected

to induce stronger microglial activation than mutants that favor monomeric forms. Building on
our previous study where we examined the impact of Aβ-42-N27D on aggregation and
microglial activity [15], this work differs methodologically in that we freshly prepared our
samples by isolating monomers using SEC to remove any pre-seeding effects that could affect
the aggregation kinetics and behavior of the Aβ-42 peptide. This modification led to a more
consistent microglial response than observed previously.

Our SEC data indicates that Aβ-42-Q15E and Aβ-42-N27D aggregate less than Aβ-42-WT.
Therefore, the observed differences in microglial activity could be partially due to altered
oligomer formation. Similar findings have been reported for Aβ-40, where the Aβ-40-Q15E and
Aβ-40-N27D variants showed slower aggregation compared to Aβ-40-WT [14]. Thus, if
microglial activation were solely dependent on aggregation state, these findings would support
the conclusion that Aβ-42-Q15E and Aβ-42-N27D are less activating than Aβ-42-WT.

While the aggregation dynamics of deamidated mutants of Aβ-40 have been documented [14],
the aggregation dynamics and microglial response to Aβ-42-Q15E and Aβ-42-N27D remains
poorly understood. Our SEC data shows that Aβ-42-Q15E aggregates similarly to Aβ-42-N27D,
despite the Q15E switch occurring in a different region of Aβ-42 than N27D. Research has
shown that the loss of negative charges at specific residues on Aβ-42 leads to increased
aggregation, enhanced stability, and greater neurotoxicity of the peptide [24]. Given that both
N27D and Q15E introduce a negative charge to Aβ-42, this supports the idea that deamidation at
these residues could result in slower aggregation. A simple explanation is that WT is net tri-
anionic, while N27D and Q15E are net tetra-anionic. The higher negative charge is expected to
enhance water solubility and disfavor hydrophobic collapse that drives oligomer formation.
Another effect added on top is the loss of fibrillogenicity of the Q15 or N27 sidechain upon
deamidation. Future structural work may reveal additional, more subtle contributing factors.

One major finding in our study is that both Aβ-42-Q15E and Aβ-42-N27D reduce microglial
activation and diminish expression of inflammatory cytokines compared to Aβ-42-WT. As Aβ-
42-WT forms more oligomers than Aβ-42-Q15E and Aβ-42-N27D, Aβ-42-WT could be able to
bind to more microglial surface receptors than the deamidated mutants. The reduced oligomer
populations in Aβ-42-Q15E and Aβ-42-N27D could explain the decreased inflammatory
cytokine expression observed in the deamidated mutants [25, 26]. While limited differences were
found between Aβ-42-Q15E and Aβ-42-N27D in terms of cytokine expression, our data suggest
that Aβ-42-Q15E and Aβ-42-N27D may interact differently with microglia [27]. One such
analyte, IL-6, is expressed significantly higher in microglia treated with Aβ-42-Q15E than Aβ-
42-N27D. Since IL-6 production is regulated by multiple receptor-mediated signaling pathways
[28, 29], these findings suggest that Aβ-42-Q15E and Aβ-42-N27D may engage distinct
microglial surface receptors. This suggestion is plausible given that the deamidation sites differ
between the two mutants, potentially altering their conformation and resultant receptor binding.

In addition to cytokine profiling, we quantified CD68 because it is upregulated in AD pathology
and cognitive decline [17]. We found that both Aβ-42-Q15E and Aβ-42-N27D induce
significantly lower expression of CD68 compared to Aβ-42-WT. Since CD68 is a marker of
lysosomal activity, it serves as a measure for evaluating lysosomal degradation of Aβ-42 [21]. As
the expression of CD68 is lower in the deamidated variants, this may indicate that Aβ-42-Q15E

and Aβ-42-N27D are more efficiently degraded or result in reduced lysosomal activation
compared to Aβ-42-WT.
228
The N27D and Q15E mutations were designed to mimic spontaneous deamidation events and
involve substitutions that disrupt critical structural elements governing Aβ-42 aggregation and
fibril stability. Both asparagine and glutamine residues are considered fibril-stabilizing due to
their ability to form periodic tightly interdigitated bidirectional H-bonding side-chain interactions
observed in both in-vitro and brain-derived Aβ fibril structures characterized by NMR and Cryo-
EM that provide crucial stability to amyloid fibril structures [30, 31]. The N27D mutation
substitutes asparagine with aspartic acid at residue 27, a position within the turn region (residues
22-29) that forms a critical structural motif facilitating aggregation through β-hairpin formation
[20, 21]. This region contains the D23-K28 salt bridge that stabilizes β-hairpin conformations
[32], with N27 normally participating in both steric zipper formation and a hydrogen bonding
network involving D23 and neighboring residues G25, S26, and K28 [21, 33]. The β-hairpin
motif serves as an important intermediate along the aggregation pathway, with transient β-
hairpin formation promoting oligomerization [24, 34]. Similarly, the Q15E mutation introduces a
negative charge within the central hydrophobic region of Aβ-42 [14]. In this region, Q15
normally contributes to fibril stability through glutamine ladder formation and steric zipper
interactions that stabilize the fibril core [30]. The substitution of these fibril-stabilizing residues
with negatively charged aspartic and glutamic acid creates multiple disruptive effects:
electrostatic repulsion near existing salt bridges (N27D), energetically unfavorable conditions in
hydrophobic regions (Q15E), and loss of the stabilizing fibrillogenic sidechain-to-sidechain
interactions that both original residues provide. These combined structural perturbations
destabilize aggregation-prone conformations and disrupt the peptide-peptide interactions
necessary for β-sheet formation and propagation, explaining the dramatic reduction in oligomer
formation observed in our SEC profiles. Previous studies demonstrating that loss of negative
charges at specific residues leads to increased aggregation and enhanced neurotoxicity [35]
support our observation that the addition of negative charges through deamidation has the
opposite effect.
While our study provides important insights into how deamidated Aβ-42 mutants modulate
microglial activation, several limitations should be acknowledged. First, the experiments were
conducted in an immortalized microglial monoculture, which shares key behavioral
characteristics with primary microglia but does not fully capture the complexity of primary
microglial phenotypes, particularly in the context of AD. Additionally, although CD68 staining
modulation suggests deamidation altered phagocytic activity, CD68 primarily reflects lysosomal
activity and does not directly measure microglial Aβ uptake via phagocytosis, a process heavily
implicated in AD pathology [36]. Also, our cytokine profile suggests that the deamidated Aβ-42
variants may elicit slightly different microglial responses, motivating future analysis of receptor
binding kinetics or downstream signaling activation. Together, these limitations highlight the
need for in vivo studies and expanded mechanistic analyses.
266
267 Overall, our data show that deamidation of Aβ-42 reduces aggregation and dampens microglial
268 activation. Given the growing clinical significance of Aβ aggregation and immune activity in AD
269 [37], our findings therefore support further interrogation of Aβ-42 deamidation as a clinically
270 relevant approach to reducing its toxicity. In particular, the formulation of new pharmacological
271 strategies for post-translational deamidation could represent a viable strategy for AD treatment.

Methods

Aβ Preparation

Aβ-42-WT, Aβ-42-N27D, and Aβ-42-Q15E peptides were synthesized using a CEM Liberty Blue peptide synthesizer in-house following Fmoc solid-phase peptide synthesis protocols and subsequently double-purified as described in our previous work [38]. Following synthesis, crude peptides were verified for molecular weight using electrospray ionization mass spectrometry (ESI-MS) at the proteomics facility, confirming the presence of peptides with correct molecular masses corresponding to their expected sequences. See **Supplementary Figure 1**.

Monomer Isolation

For monomer isolation, lyophilized peptides were dissolved in 50 mM sodium hydroxide (NaOH) and allowed to equilibrate for 20 minutes at room temperature to ensure complete dissolution. The peptide solutions were then immediately subjected to size exclusion chromatography (SEC) to isolate monomeric fractions.

Size Exclusion Chromatography

Size exclusion chromatography (SEC) was performed using a Superdex 75 increase 10/300 GL column (GE Healthcare) connected to an ÄKTA GO chromatography system. The running buffer consisted of 20mM sodium phosphate (NaPi, pH 7.4). Peptide samples (3 mL) were injected at a flow rate of 0.5 mL/min, and elution was monitored by UV absorbance at 280 nm. Monomeric fractions were identified based on their elution volume (~14 mL) and collected accordingly. The isolated monomers were immediately diluted to a final concentration of 100 μM in the same buffer and used for aggregation studies.

For aggregation analysis, freshly isolated monomers were incubated at 37°C for 24 hours without agitation. Following incubation, samples were re-injected into the SEC column under identical conditions to analyze the aggregation state.

Cell Conditioning and Media Collection

Mouse microglial cells (MMCs) were cultured in DMEM-F12 medium supplemented with 10% fetal bovine serum and 1% Penicillin-Streptomycin until they reached 70% confluency. Cells were stored in an incubator with 95% air and 5% CO_2 atmosphere at 37 °C. Upon reaching 70% confluency, cells were seeded into 12-well plates and incubated for 24 hours. Cells were treated with either 50 nM Aβ42-WT, 50 nM Aβ42-N27D, 50nM Aβ42-Q15E or vehicle (20mM NaPi) for 24h followed by medium collection. Collected medium was centrifuged at 13,000 RPM for 10 min at 4°C, then the supernatant was collected and stored at -80°C. Multiplexed cytokine quantification was conducted using the Milliplex® MAP Mouse Cytokine/Chemokine 18-Plex kit (Eotaxin[*], GM-CSF[*], IFN-γ[*], IL-1β[*], IL-2[*], IL-3[*], IL-4[*], IL-6, IL-10, IL-12p70[*], IL-13[*], IP-10, M-CSF[*], MIP-1α, MIP-1β, RANTES, TNFα, and VEGF) (Millipore Sigma MCYTOMAG-70K). Cytokines marked with an asterisk did not fall within a linear range of bead fluorescent intensity vs. protein concentration and were not included in the analysis.

Multivariate Analysis of Cytokine Data

Data were analyzed and figures were generated via RStudio (Boston, MA) using the R programming language. The tidyverse collection of packages was used to conduct data processing. R package heatmap3 were used to generate heatmaps; R packages ggplot2, ggpubr,

ggbeeswarm, and ggprism were used to generate dotplots and barplots. Clustering was conducted using the hclust function of the stats package in R using Euclidean distance in the unweighted pair group method with arithmetic mean. For functional and protein analysis, Bonferroni adjustment for applied to correct for multiple comparisons.

Multivariate cytokine data were analyzed using Principal Component Analysis (PCA) to identify axes, called principal components (PCs), that capture the greatest variance in cytokine expression across samples. PCA was performed using the R package ropls, with all data z-scored prior to analysis. Two principal components were extracted, and 5-fold cross-validation was applied to evaluate model robustness. All cytokine data were z-scored prior to analysis. PCA loadings were used to represent the contribution of each analyte to the observed variance.

Cell Conditioning for CD68 Analysis
Mouse microglial cells were cultured in DMEM-F12 medium supplemented with 10% fetal bovine serum and 1% Penicillin-Streptomycin until they reached 70% confluency. Cells were stored in an incubator with 95% air and 5% CO_2 atmosphere at 37□°C. Upon reaching 70% confluency, cells were seeded into a 96-well plate and incubated for 24 hours. Cells were treated with either 50 nM Aβ42-WT, 50 nM Aβ42-N27D, 50nM Aβ42-Q15E or vehicle (20mM NaPi) for 2h followed by fixing with 4% PFA on ice for 10 minutes. Cells were then rinsed 3x with cold PBS and then stored at 4°C.

Immunocytochemistry
Fixed cells were permeabilized with 0.01% Triton X-100 in PBS for 10 minutes at room temperature, then rinsed twice with PBS. Cells were blocked for 1 hour at room temperature using blocking buffer containing 5% BSA and 3% goat serum in PBS. Following blocking, cells were incubated overnight at 4°C with rat anti-CD68 antibody (1:200; ThermoFisher). The following day, cells were washed five times with PBS containing 5% BSA, with each wash lasting 5 minutes. Wells were then incubated with Alexa Fluor 555 conjugated secondary antibody (1:2000; ThermoFisher) for 2 hours at room temperature. Afterward, cells were washed three times with 5% BSA in PBS for 5 minutes each wash. Nuclei were counterstained with DAPI for 20 minutes at room temperature, followed by three additional 5-minute washes with 5% BSA in PBS. Cells were then rinsed three times with PBS and stored in PBS until imaging. Samples were imaged using epifluorescent microscopy on a Zeiss Axio Observer Z.1 inverted microscope with a 20x lens and halogen bulb illumination using Zeiss filter set 49 to image DAPI and Zeiss filter set 20 to image Alexa fluor 555. Quantification of CD68 florescent intensity was analyzed by dividing the CD68 intensity by DAPI cell counts (ImageJ Macro).

ImageJ Macro
An ImageJ macro was developed to automate the quantification of CD68/IBA1 fluorescent intensity. The macro recursively scans a directory for merged images containing targeted (green for IBA1, red for CD68) and blue (DAPI) channels, while excluding single-channel files. Each image was then split into individual channels. The DAPI channel is thresholded using the Otsu method and segmented via watershed to count DAPI-positive nuclei, while measuring the green/red channel integrated fluorescent intensity. The intensity value was then normalized to the number of DAPI-positive nuclei to account for cell density. Finally, output values were printed to a log for export.

363
364 **Data Availability**
365 The data described in the manuscript are located within the manuscript and supplemental
366 information.
367
368 **Supporting Information**
369 This article contains supporting information.
370
Funding Sources
This work was funded by Georgia W. Woodruff School of Mechanical Engineering Faculty
Fellowship at Georgia Tech (L.B.W.) and in part by the National Institute of Health under grant
number R01AG075820. This work was also funded in part by the National Institute of Health
under grant number R01AG074954. This work was also supported by the National Institute on
Aging of the National Institutes of Health and the National Institute on Minority Health and
Health Disparities (award number RF1AG073434-01A1). The content is solely the responsibility
of the authors and does not necessarily represent the official views of the National Institutes of
Health.
380
381 **Conflict of Interest**
382 The authors declare that they have no conflicts of interest with the contents of this article.

References

1. *2024 Alzheimer's disease facts and figures.* Alzheimers Dement, 2024. **20**(5): p. 3708–3821.
2. Sadigh-Eteghad, S., et al., *Amyloid-beta: a crucial factor in Alzheimer's disease.* Med Princ Pract, 2015. **24**(1): p. 1–10.
3. Gouras, G.K., T.T. Olsson, and O. Hansson, *beta-Amyloid peptides and amyloid plaques in Alzheimer's disease.* Neurotherapeutics, 2015. **12**(1): p. 3–11.
4. Tiwari, S., et al., *Alzheimer's disease: pathogenesis, diagnostics, and therapeutics.* Int J Nanomedicine, 2019. **14**: p. 5541–5554.
5. Sarlus, H. and M.T. Heneka, *Microglia in Alzheimer's disease.* J Clin Invest, 2017. **127**(9): p. 3240–3249.
6. Li, Y., et al., *Mitochondrial dysfunction in microglia: a novel perspective for pathogenesis of Alzheimer's disease.* J Neuroinflammation, 2022. **19**(1): p. 248.
7. Gao, C., et al., *Microglia in neurodegenerative diseases: mechanism and potential therapeutic targets.* Signal Transduct Target Ther, 2023. **8**(1): p. 359.
8. Dani, M., et al., *Microglial activation correlates in vivo with both tau and amyloid in Alzheimer's disease.* Brain, 2018. **141**(9): p. 2740–2754.
9. Yang, I., et al., *The role of microglia in central nervous system immunity and glioma immunology.* J Clin Neurosci, 2010. **17**(1): p. 6–10.
10. Rajmohan, R. and P.H. Reddy, *Amyloid-Beta and Phosphorylated Tau Accumulations Cause Abnormalities at Synapses of Alzheimer's disease Neurons.* J Alzheimers Dis, 2017. **57**(4): p. 975–999.
11. Kinney, J.W., et al., *Inflammation as a central mechanism in Alzheimer's disease.* Alzheimers Dement (N Y), 2018. **4**: p. 575–590.
12. Zhou, Y., et al., *Human and mouse single-nucleus transcriptomics reveal TREM2-dependent and TREM2-independent cellular responses in Alzheimer's disease.* Nat Med, 2020. **26**(1): p. 131–142.
13. Kuperstein, I., et al., *Neurotoxicity of Alzheimer's disease Abeta peptides is induced by small changes in the Abeta42 to Abeta40 ratio.* EMBO J, 2010. **29**(19): p. 3408–20.
14. Sanagavarapu, K., et al., *A method of predicting the in vitro fibril formation propensity of Abeta40 mutants based on their inclusion body levels in E. coli.* Sci Rep, 2019. **9**(1): p. 3680.
15. Sajimon, M., et al., *Asparagine Deamidation Attenuates Toxicity, Aggregation, and Microglial Responses of Alzheimer's Amyloid-β.* ACS Chemical Neuroscience, 2025.
16. Iliyasu, M.O., et al., *Amyloid-beta aggregation implicates multiple pathways in Alzheimer's disease: Understanding the mechanisms.* Front Neurosci, 2023. **17**: p. 1081938.
17. Naslund, J., et al., *Correlation between elevated levels of amyloid beta-peptide in the brain and cognitive decline.* JAMA, 2000. **283**(12): p. 1571–7.
18. Raskatov, J.A., *What Is the "Relevant" Amyloid beta42 Concentration?* Chembiochem, 2019. **20**(13): p. 1725–1726.
19. Wood, L.B., et al., *Identification of neurotoxic cytokines by profiling Alzheimer's disease tissues and neuron culture viability screening.* Sci Rep, 2015. **5**: p. 16622.

20. Pan, X.D., et al., *Microglial phagocytosis induced by fibrillar beta-amyloid is attenuated by oligomeric beta-amyloid: implications for Alzheimer's disease.* Mol Neurodegener, 2011. **6**: p. 45.

21. Lier, J., W.J. Streit, and I. Bechmann, *Beyond Activation: Characterizing Microglial Functional Phenotypes.* Cells, 2021. **10**(9).

22. Thomas, A.L., et al., *Naturally-aged microglia exhibit phagocytic dysfunction accompanied by gene expression changes reflective of underlying neurologic disease.* Sci Rep, 2022. **12**(1): p. 19471.

23. Doens, D. and P.L. Fernandez, *Microglia receptors and their implications in the response to amyloid beta for Alzheimer's disease pathogenesis.* J Neuroinflammation, 2014. **11**: p. 48.

24. Cabrera, E., et al., *Abeta truncated species: Implications for brain clearance mechanisms and amyloid plaque deposition.* Biochim Biophys Acta Mol Basis Dis, 2018. **1864**(1): p. 208–225.

25. Jarosz-Griffiths, H.H., et al., *Amyloid-beta Receptors: The Good, the Bad, and the Prion Protein.* J Biol Chem, 2016. **291**(7): p. 3174–83.

26. Amin, L. and D.A. Harris, *Abeta receptors specifically recognize molecular features displayed by fibril ends and neurotoxic oligomers.* Nat Commun, 2021. **12**(1): p. 3451.

27. Sondag, C.M., G. Dhawan, and C.K. Combs, *Beta amyloid oligomers and fibrils stimulate differential activation of primary microglia.* J Neuroinflammation, 2009. **6**: p. 1.

28. Bobbo, V.C.D., et al., *Interleukin-6 Expression by Hypothalamic Microglia in Multiple Inflammatory Contexts: A Systematic Review.* Biomed Res Int, 2019. **2019**: p. 1365210.

29. Garner, K.M., et al., *Microglia priming by interleukin-6 signaling is enhanced in aged mice.* J Neuroimmunol, 2018. **324**: p. 90–99.

30. Gremer, L., et al., *Fibril structure of amyloid-beta(1-42) by cryo-electron microscopy.* Science, 2017. **358**(6359): p. 116–119.

31. Walti, M.A., et al., *Atomic-resolution structure of a disease-relevant Abeta(1-42) amyloid fibril.* Proc Natl Acad Sci U S A, 2016. **113**(34): p. E4976–84.

32. Baumketner, A., et al., *Structure of the 21-30 fragment of amyloid beta-protein.* Protein Sci, 2006. **15**(6): p. 1239–47.

33. Foley, A.R., H.W. Lee, and J.A. Raskatov, *A Focused Chiral Mutant Library of the Amyloid beta 42 Central Electrostatic Cluster as a Tool To Stabilize Aggregation Intermediates.* J Org Chem, 2020. **85**(3): p. 1385–1391.

34. Khaled, M., et al., *A Hairpin Motif in the Amyloid-beta Peptide Is Important for Formation of Disease-Related Oligomers.* J Am Chem Soc, 2023. **145**(33): p. 18340–18354.

35. Guo, M., et al., *Charge substitution shows that repulsive electrostatic interactions impede the oligomerization of Alzheimer amyloid peptides.* FEBS Lett, 2005. **579**(17): p. 3574–8.

36. Chistiakov, D.A., et al., *CD68/macrosialin: not just a histochemical marker.* Lab Invest, 2017. **97**(1): p. 4–13.

37. Azargoonjahromi, A., *The duality of amyloid-beta: its role in normal and Alzheimer's disease states.* Mol Brain, 2024. **17**(1): p. 44.

38. Dutta, S., et al., *Suppression of Oligomer Formation and Formation of Non-Toxic Fibrils upon Addition of Mirror-Image Abeta42 to the Natural l-Enantiomer.* Angew Chem Int Ed Engl, 2017. **56**(38): p. 11506–11510.