Title

CYFIP1 overexpression amplifies IL-6/STAT3 and IFN-gamma/STAT1 signaling: potential

implications for neuroinflammation and autism spectrum disorder

Authors

Emily-Rose Martin[1], Josan G. Martin[1], Mark A. Russell[1], Asami Oguro-Ando[1,2*]

[1] = University of Exeter Medical School, University of Exeter, Exeter, United Kingdom

[2] = Faculty of Pharmaceutical Sciences, Tokyo University of Science, Tokyo, Japan

*Corresponding Author: a.oguro-ando@exeter.ac.uk

Abstract

Autism spectrum disorder (ASD) encompasses a group of neurodevelopmental disorders

characterised by impaired social interaction, delayed language development, and

repetitive or restrictive behaviours. While both genetic and environmental factors

contribute to ASD, the specific molecular mechanisms underlying these interactions

remain unclear. In a recent preprint, we showed that cytokine signaling may be

dysregulated in chromosome 15q-duplication syndrome (Dup(15q)), one of the most

common syndromic forms of ASD. Specifically, Dup(15q) induced pluripotent stem cells

(iPSC)-derived neurons exhibit an amplified Signal Transducer and Activator of

Transcription-3 (STAT3) response to Interleukin-6 (IL-6), a proinflammatory cytokine often

upregulated in ASD. To identify which gene in the 15q region may be responsible for

modifying cytokine signalling responses in Dup(15q), we focus on the *Cytoplasmic*

FMRP-Interacting Protein 1 (CYFIP1) gene, as increased *CYFIP1* dosage in Dup(15q) is

associated with increased severity of neurobehavioral symptoms in individuals with

Dup(15q) and *CYFIP1* dysregulation has been linked to altered expression of various

genes involved in immunoregulatory signalling pathways. Here, we explore the effects of

CYFIP1 overexpression (*CYFIP1*-OE) on cytokine signaling, demonstrating that *CYFIP1*-

OE in HEK293 cells modifies the expression and activity of several cytokine signaling-

related transcription factors, including STAT3 and STAT1. Additionally, we use SH-SY5Y

neuroblastoma cells to assess neuron-related phenotypes, showing that *CYFIP1*-OE

alters IL-6-induced neurite outgrowth. These findings provide novel insights into how

CYFIP1 may contribute to dysregulated cytokine responses in ASD, advancing our

understanding of the molecular mechanisms that underlie neuroinflammatory

processes in this disorder.

Keywords

Background

Autism spectrum disorder (ASD) is a heterogeneous group of neurodevelopmental

conditions in which both genetic variation and environmental exposures contribute to

pathogenesis (1). Due to the highly heterogeneous nature of the disorder, both diagnosis

and treatment remain challenging. Genetic factors, including both rare and common

variants, are implicated in ASD, with over 1,000 genes identified as potentially

contributing to its onset and development (2). In addition to genetic predisposition,

environmental risk factors, particularly maternal immune activation (MIA) and

neuroinflammation, have been recognised as key contributors to ASD pathogenesis (3,4).

Therefore, it is likely that the interaction between an individual's genetic background and

specific environmental exposures plays a crucial role in triggering the pathogenesis

underlying ASD (3). Consequently, elucidating the molecular mechanisms by which ASD-

associated genes can alter neurodevelopmental processes in response to environmental

stressors is critical for advancing our understanding of the disorder.

Janus Kinase and Microtubule-Interacting Protein 1 (*JAKMIP1*) was previously identified

as a potential molecular link between syndromic forms of ASD and neuroinflammation

(5). *JAKMIP1* expression was commonly dysregulated in Fragile X syndrome (FXS) and

chromosome 15q-duplication syndrome (Dup(15q)), both genetic disorders often

comorbid with ASD (5). Furthermore, we discovered a role for JAKMIP1 in regulating

Interleukin-6 (IL-6)-induced Signal Transducer and Activator of Transcription-3 (STAT3)

signaling in neuronal cells, whereby deficiency of JAKMIP1 in SH-SY5Y neuroblastoma

cells impairs STAT3 activation and neuritogenesis in response to IL-6 stimulation (5).

These findings suggest that dysregulation of *JAKMIP1* in ASD may alter the ability of

neurons to respond to inflammatory cytokines, potentially leading to aberrations in

neuronal morphology.

To investigate whether *JAKMIP1* dysregulation in syndromic ASD could modify how cells

respond to cytokines, we previously assessed IL-6-induced STAT3 signalling in a

Dup(15q) induced pluripotent stem cell (iPSC)-derived neuronal model (6), as Dup(15q)

is one of the most common forms of syndromic ASD, accounting for around 1% of ASD

cases (7). Compared to a control iPSC line, we observed that *JAKMIP1* expression

fluctuates in Dup(15q) iPSC and iPSC-derived cortical neurons (iNeurons), being

downregulated in Dup(15q) iPSC and upregulated in iNeurons. Despite these opposing

trends, we found that *STAT3* expression was consistently reduced in both the Dup(15q)

iPSCs and iNeurons, indicating that Dup15q cells would likely display altered responses

to IL-6. Curiously, though *STAT3* expression was reduced, IL-6-induced STAT3 activity was

significantly enhanced in both Dup(15q) iPSCs and iNeurons. These findings suggest that

unlike in the SH-SY5Y cells, JAKMIP1 is likely not regulating *STAT3* expression or activity in

Dup(15q) and instead, another gene in the chromosome 15q region exerts stronger

influence on STAT3 activity.

Several genes are encoded within the 15q region of chromosome 15, however, a gene of

particular interest in relation to ASD and cytokine signalling is *Cytoplasmic FMRP-*

Interacting Protein 1 (*CYFIP1*), as studies have demonstrated that duplications (or even

deletions) encompassing *CYFIP1* increase the severity of neurobehavioural phenotypes

(8), and that *CYFIP1* expression is upregulated in the brains of individuals with Dup(15q)

and ASD (9). Although CYFIP1 is most known for its roles in repressing neuronal protein

translation as part of Fragile X Messenger Ribonucleoprotein (FMRP)-containing

ribonucleoprotein complexes (10) and regulating actin polymerisation as a core

component of the WAVE regulatory complex (WRC) (11), there is evidence to suggest that

CYFIP1 dysregulation may alter the expression of various genes involved in

immunoregulatory signalling pathways (12). For example, a recent study profiling

transcriptomic changes in Dup(15q) stem cell-derived neurons with upregulated *CYFIP1*

expression suggested that increased dosage of chromosome 15q11.2 genes could alter

the expression gene targets of SMAD Family Member 3 (SMAD3), Nuclear Factor Kappa B

(NF-κB) or Interferon-Stimulated Response Elements (ISRE) (12). Moreover, another

study identified *CYFIP1* as a potential hub gene that may coordinate Transforming Growth

Factor Beta 1 (TGF-β1)/SMAD3 signalling (13). Therefore, we hypothesised that CYFIP1

may be responsible for the observed enhanced IL-6-induced activity in Dup(15q) iPSCs

and iNeurons (6).

This paper therefore explores the effects of increased *CYFIP1* expression (modelled by

CYFIP1-overexpression (OE)) on cytokine signalling. We provide the first evidence to

directly support a novel role for CYFIP1 in modulating cytokine signalling, demonstrating

that *CYFIP1*-OE in HEK293 cells alters the expression and activity of cytokine signalling-

related transcription factors, *STAT1* and *STAT3*. Taking this further, we show how

alterations in the STAT3 signalling pathway due to *CYFIP1*-OE affects IL-6-induced neurite

outgrowth in human SH-SY5Y neuroblastoma cells, illustrating that these findings

continue to be relevant in a more neuronal-relevant model.

Methods

HEK293 and SH-SY5Y cell culture and treatment

Human embryonic kidney (HEK)-293 cells (#CRL-1573™, *American Type Culture*

Collection) and human neuroblastoma cells (#CRL-2266™, *American Type Culture*

Collection) were thawed from frozen stocks stored in 10% dimethyl sulfoxide (#D2650-

5X5ML, *Sigma-Aldrich®*) at −80°C. HEK293 cells and SH-SY5Y cells were maintained in

10-cm diameter Petri dishes (#P7612-360EA, *Sigma-Aldrich®*) with 10 mL Dulbecco's

modified eagle medium/nutrient mixture F-12 with GlutaMAX (referred to henceforth as

DMEM/F-12) (#11524436, *Fisher Scientific™*) supplemented with 10% foetal bovine

serum (FBS) (#11550356, *Fisher Scientific™*), and incubated at 37°C, 5% CO_2, 95%

humidity. Cells were passaged at 70-80% confluency using 3 mL of pre-warmed TrypLE™

(#10718463, *Fisher Scientific™*) for 5 min and TrypLE™ was inactivated by dilution in an

equal volume of DMEM/F-12 + 10% FBS.

Cells were stimulated with 20 ng/mL IL-6 (#7270IL-025, *R&D Systems*) or 20 ng/mL

Interferon-γ (IFN-γ) (#285-IF-100, *R&D Systems*) diluted in DMEM/F-12 + FBS (or in

DMEM/F-12 supplemented with Retinoic Acid (RA) and Brain-Derived Neurotrophic

Factor (BDNF) for neurite tracing experiments). For all experiments using IL-6 or IFN-γ,

control cells were used which were treated with corresponding media not containing IL-

6 or IFN-γ. Cells were treated with IL-6 or IFN-γ for different amounts of time depending

on the purpose of the experiment. To assess STAT3 activation by Western blotting for

phosphorylated STAT3 (P-STAT3^{Y705}), cells were treated with IL-6 for 30 minutes. To assess

STAT3 or STAT1 transcriptional activity by Dual-Luciferase® Reporter (DLR™) assays, cells

were treated with IL-6 or IFN-γ for 18 hours post-transfection. To measure mRNA

expression of STAT3 or STAT1 responsive genes, cells were treated with IL-6 or IFN-γ for 4

hours. Finally, to assess IL-6-induced neurite outgrowth, cells were treated with IL-6 over

a 48-hour period of differentiation.

HEK293 cell transfection

24 hours prior to transfection, 300,000 HEK293 cells were seeded into Nunc cell-culture

treated 6-well plates (#10469282, *Fisher Scientific™*) in 3 mL DMEM/F-12 + 10% FBS. The

following day, HEK293 cells were transfected with either an RFP-T2A-CYFIP1 plasmid to

induce *CYFIP1*-overexpression (these cells are referred to as '*CYFIP1*-OE cells), or the

backbone RFP-T2A plasmid as a control (these cells are referred to as 'control cells')

using Lipofectamine™ LTX Reagent (#15338100, *Fisher Scientific™*) following

manufacturer instructions (i.e., final amount of 2.5 µg of plasmid and 5 µL of

Lipofectamine™ LTX per well). Media was replaced the following day. All experiments were

performed 72 hours post-transfection.

SH-SY5Y cell transfection

2,000,000 SH-SY5Y cells were transfected by Nucleofection™ (#V4XC-2012, *Lonza*

Bioscience) following manufacturer instructions with 5 µg of either a mixture of the RFP-

T2A backbone vector and pCAβ-YFP (encoding yellow fluorescent protein (YFP)) in a 1:1

ratio as the control condition; or a mixture of RFP-T2A-CYFIP1 and pCAβ-YFP (4:1 ratio) to

assess the effects of *CYFIP1*-OE on neuritogenesis. A pCAβ-YFP plasmid was co-

transfected to enable better visualisation of the cells for neurite tracing as the RFP

encoded by the *CYFIP1*-OE plasmid photo-bleaches quickly. Media was replaced the

following day.

Protein extraction and Western blotting

72 hours post-transfection, cells were washed with ice-cold 1x phosphate buffered

saline (1xPBS) (#P4417-50TAB, Sigma-Aldrich®) then collected with a cell scraper in 300

µL of ice-cold RIPA lysis buffer (150 mM NaCl, 50 mM Tris/Cl pH = 8.0, 0.5% Sodium

Deoxycholate, 0.1% SDS, 1% Triton X-100) supplemented with 1 mM PMSF, 1%

Phosphatase Inhibitor Cocktail 2 (#P5726-1ML, Sigma Aldrich®), 1% Phosphatase

Inhibitor Cocktail 3 (#P0044-1ML, Sigma Aldrich®) and 1% Complete Protease Inhibitor

Cocktail (#P8340-1ML, Sigma Aldrich®). Cell lysates were incubated on ice for 30

minutes, cleared by centrifugation at 17,000xg for 15 minutes at 4°C, then supernatants

collected. Bicinchoninic Acid assays (#10678484, *Thermo Scientific*) were performed to

estimate protein concentrations and 50 μg of total protein was diluted with Laemmli

buffer (final concentration of 60 mM Tris/Cl pH = 6.8, 10% Glycerol, 2% SDS, 5% 2-

Mercaptoethanol, 0.02% Bromophenol Blue), then boiled for 5 minutes at 95°C. 50 μg of

total protein for each sample was loaded into 4–15% precast polyacrylamide gels

(#4561085, *Bio-Rad Laboratories*) alongside a pre-stained protein ladder (#PL00001,

Proteintech Group). Electrophoresis was performed in the Mini-PROTEAN® Tetra Vertical

Electrophoresis Cell (#1658004, *Bio-Rad Laboratories*) with 1×SDS-PAGE running buffer

(2.5 mM tris-base; 19 mM glycine (#10070150, *Fisher Chemical™*); 0.01% SDS; pH = 8.3

in ddH$_2$O). Proteins were then transferred onto methanol-activated PVDF membranes

(#IPVH304F0, *Millipore*) by a wet transfer method in the Mini-PROTEAN® Tetra Vertical

Electrophoresis Cell. PVDF membranes were then blocked for at least 1 hour at room

temperature with 5 mL of 5% skim milk in 1x tris-buffered saline (TBS; 2 mM Tris-Base; 15

mM NaCl; pH = 7.6 in ddH$_2$O) supplemented with 0.1% Tween® 20 (#10485733, *Fisher*

BioReagents™) (1xTBS-T). After blocking, membranes were sequentially incubated with 5

mL of primary antibody (gentle rocking for 2-3 hours at room temperature or overnight at

4°C), then secondary antibody (gentle rocking for 30 minutes-1 hour at room

temperature). Primary and secondary antibodies were diluted in 5% skim milk in 1x TBS-

T (or 5% BSA in 1x TBS-T for anti-P-STAT3^{Y705}). The dilution factors and specific antibodies

used are listed in **Table 1**. Three 5-minute washes with 1x TBS-T were performed between

antibody incubations. GAPDH was used as a loading control to normalise expression of

target proteins. Proteins were visualised using the Odyssey® CLx Imager (*LI-COR*

Biosciences) and the Image Studio software (Version 5.2; *LI-COR Biosciences*) was used

to perform densitometric analyses to compare the abundance of target proteins between

samples.

Catalogue Number	Supplier	Host Species	Target	Dilution	Conjugated Fluorophore
ab156016	*Abcam*	Rabbit	CYFIP1	1:1000	
sc-47724	*Santa Cruz Biotechnology*	Mouse	GAPDH	1:1000	
13846-1-AP	*Proteintech*	Rabbit	JAKMIP1	1:1000	
9139S	*Cell Signaling Technology*	Mouse	STAT3	1:1000	
9131l	*Cell Signaling Technology*	Rabbit	P-STAT3^{Y705}	1:1000	
35519	*Invitrogen*	Goat	Mouse	1:5000	DyLight™ 680
SA5-10036	*Invitrogen*	Goat	Rabbit	1:5000	DyLight™ 800

Table : Antibodies used in this study.

RNA extraction, cDNA conversion and qRT-PCR

72 hours post-transfection, total RNA was isolated and purified using the Direct-zol™ RNA

Miniprep kit following manufacturer instructions (#R2051, *Zymo Research*). 500 ng of

RNA was converted to cDNA using PrimeScript™ RT reagent kit (#RR037A, *Takara Bio*

Europe) following manufacturer instructions. qRT-PCR was then performed using HOT

FIREPol® EvaGreen® qPCR Master Mix with ROX (#01-02-00500, *Solis BioDyne*) with the

QuantStudio 12K Flex qPCR machine (*Thermo Fisher Scientific*™). For primer sequences,

see **Table 2**. mRNA expression levels were standardized against two reference genes,

Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH) and *RNA Polymerase II Subunit*

A (POLR2A) using the Pfaffl method (14) and normalised to the control cells.

Gene	Strand	Sequence
CYFIP1	Forward	TAC GAG ACG CTG CTG AAG CAG A
CYFIP1	Reverse	TCG TCC AAT CGC CAG TTC TAG G
GAPDH	Forward	TCC TCT GAC TTC AAC AGC GAC
GAPDH	Reverse	GCT GTA GCC AAA TTC GTT GTC A
NF-KB	Forward	GCA GCA CTA CTT CTT GAC CAC C
NF-KB	Reverse	TCT GCT CCT GAG CAT TGA CGT C
POLR2A	Forward	CCA TCA AGA GAG TCC AGT TCG
POLR2A	Reverse	ACC CTC CGT CAC AGA CAT TC
SOCS1	Forward	TTC GCC CTT AGC GTG AAG ATG G
SOCS1	Reverse	TAG TGC TCC AGC AGC TCG AAG A
SOCS3	Forward	CCA AGG ACG GAG ACT TCG ATT C
SOCS3	Reverse	GGG AAA CTT GCT GTG GGT GA
STAT1	Forward	ATG GCA GTC TGG CGG CTG AAT T
STAT1	Reverse	CCA AAC CAG GCT GGC ACA ATT G
STAT3	Forward	GGG AAG AAT CAC GCC TTC TAC
STAT3	Reverse	ATC TGC TGCT TCT CCG TCA C
STAT5B	Forward	GCC ACT GTT CTC TGG GAC AAT G
STAT5B	Reverse	ACA CGA GGT TCT CCT TGG TCA G

Table 2: Primers used in this study.

Dual-luciferase reporter assays

50,000 HEK293 cells were seeded into 24-well plates and incubated overnight. The

following day, cells were transfected with 2 μL of STAT3 Cignal Reporter plasmid

(#336841, GeneGlobe ID: CCS-9028L, *Qiagen*) or 2 μL of STAT1 (GAS) Cignal Reporter

plasmid (#336841, GeneGlobe ID: CCS-009L, *Qiagen*) using Lipofectamine™ LTX as

previously described. Four hours post-transfection, cells were treated with DMEM/F-12

+10% FBS with or without IL-6- or IFN-γ and incubated overnight. 18 hours later, Firefly

and Renilla luciferase activities were measured using the Dual-Luciferase® Reporter

(DLR™) Assay System (#E1960, *Promega*) following manufacturer instructions.

Luminescence was measured in white-bottomed LUMITRAC™ 96-well plates (#655074,

Greiner Bio-One) using the PHERAstar FS microplate reader (*BMG LABTECH*).

SH-SY5Y differentiation and neurite outgrowth analysis

Sterile 13-mm-diameter glass coverslips (#631-1578, *VWR®*) were placed into 24-well

plates and coated with 300 μL of 20 μg/mL poly-D-Lysine (PDL; #354210, *Corning®*) by

overnight incubation at 37°C. 24 hours post-transfection, 30,000 SH-SY5Y cells were

seeded onto PDL-coated coverslips in DMEM/F-12 + 10% FBS. The following day, media

was replaced with 20 ng/mL BDNF and 10 μM RA in DMEM/F-12 (without FBS). 48 hours

later, SH-SY5Y cells were fixed with 4% paraformaldehyde (PFA), and nuclei were

counterstained with 4',6-diamidino-2-phenylindole (DAPI) (#D9542, *Sigma-Aldrich®*) for

15 minutes. Coverslips were mounted onto glass microscope slides with ProLong™

Diamond Antifade Mountant (#15468070, *Fisher Scientific™*) and slides were imaged on

the inverted DMi8 widefield microscope (*Leica Microsystems*).

All image analysis was performed manually using the *FIJI* software (*ImageJ*; Version 2.9.0;

(15)). Prior to image analysis, a scale bar was used to set a pixel to μm ratio to each image,

after which the "Segmented Line" tool was used to measure the distance from the edge

of the nucleus to the furthest end of each neurite. The length of the longest neurite

produced by a single cell (referred to as longest neurite length (LNL); a measure of neurite

extension) and the sum length of all neurites produced by a single cell (referred to as total

neurite length (TNL); a measure of neuritogenesis) were measured.

Statistical analysis

Statistical analyses were performed using GraphPad Prism (GraphPad Prism Version

9.0.0 for Windows, GraphPad Software, San Diego, California USA, www.graphpad.com).

Unpaired Student's t-Tests were used to assess statistical differences in measurements

between two groups (e.g., *CYFIP1*-OE against control). When investigating the effects of

and interactions between two independent variables (e.g., the effect of both *CYFIP1*-OE

and IL-6 treatment on STAT3 transcriptional activity), two-way analysis of variance

(ANOVA) were performed. Appropriate post hoc tests were also performed to account for

multiple testing. Data are expressed as mean values ± standard error of the mean (SEM).

P-values < 0.05 were considered statistically significant.

257
258
Results
Modelling CYFIP1-overexpression

To model increased *CYFIP1* expression, we utilised plasmid vectors encoding an RFP-

T2A-CYFIP1 construct (to induce *CYFIP1*-OE) or RFP-T2A as a control, which we

transfected into HEK293 cells, referred to as *CYFIP1*-OE and control cells respectively

(**Figures 1A and 1B**). We then confirmed increased *CYFIP1* mRNA expression using qRT-

PCR (**Figure 1C**) and protein expression using Western blotting (**Figures 1D and E**).

Figure 1 | Modelling CYFIP1 overexpression in HEK293 cells. A) *HEK293 cells were*

transfected with either RFP-T2A plasmid as a control (referred to as Control), or RFP-T2A-

CYFIP1 to induce CYFIP1-overexpression (OE) (referred to as CYFIP1-OE). Created with

BioRender.com. ***B)*** *Control and CYFIP1-OE cells were imaged 72 hours post-transfection*

live on the Leica DMi8 widefield microscope. Scale bar = 100 μm. ***C)*** *Relative mRNA*

expression of CYFIP1 in Control and CYFIP1-OE cells 72 hours post-transfection

measured by qRT-PCR. ***D)*** *Increased CYFIP1 protein expression in CYFIP1-OE cells 72*

hours post-transfection confirmed via Western blotting. ***E)*** *Densitometric analysis of total*

CYFIP1 protein expression relative to the GAPDH loading control. All values are

normalised to the Control cells. For all experiments, values are presented as mean ± SEM

of N = 3 independent experiments, 2-3 technical replicates each. Statistical significance

*against Control was determined using Unpaired Student's T-Tests; ****P < 0.0001.*

CYFIP1-OE leads to reduced JAKMIP1 expression

It has been previously demonstrated that *JAKMIP1* expression is reduced following

CYFIP1-OE in human neuroblastoma cells (5). Therefore, to confirm whether this is the

case in HEK293 cells, we measured *JAKMIP1* expression in *CYFIP1*-OE HEK293 cells by

qRT-PCR and Western blotting. Agreeing with Nishimura et al., we found that *JAKMIP1*

mRNA (**Figure 2A**) and protein expression (**Figure 2B and C**) is significantly reduced in

CYFIP1-OE HEK293 cells.

289
Figure 2 | CYFIP1-OE leads to reduced JAKMIP1 mRNA and protein expression. A)

Relative mRNA expression of JAKMIP1 in Control and CYFIP1-OE cells measured by qRT-

PCR. B) Western blotting of JAKMIP1 expression in Control and CYFIP1-OE cells. C)

Densitometric analysis of total JAKMIP1 protein expression relative to the GAPDH loading

control. For all experiments, values are presented as mean ± SEM of N = 3 independent

experiments, 2-3 technical replicates each. Statistical significance against Control was

*determined using Unpaired Student's T-Tests; *P < 0.05; **P < 0.01.*

298 ***CYFIP1*-OE alters the expression of several cytokine signalling-related transcription**

299 **factors**

300 We have previously demonstrated that Dup(15q) iPSCs and iNeurons with increased

301 *CYFIP1* expression show reduced *STAT3* expression (6). Furthermore, we have

302 demonstrated that *JAKMIP1*-knockout SH-SY5Y cells also show reduced *STAT3*

303 expression and that JAKMIP1 deficiency may alter other cytokine signalling pathways too

304 (6). Therefore, to determine whether *CYFIP1*-OE may also lead to altered expression of

305 cytokine signalling-related transcription factors, qRT-PCR was performed to measure

306 mRNA expression of *STAT1*, *STAT3*, *STAT5B* and *NFKB1*. We observed that the expression

307 of *STAT1* (**Figure 3A**) and *STAT3* (**Figure 3B**) were reduced in *CYFIP1*-OE HEK293 cells,

308 whereas the expression of *STAT5B* (**Figure 5C**) and *NFKB1* (**Figure 5D**) were increased.

309 This suggests that CYFIP1 may also modulate the expression of cytokine signalling-

310 related transcription factors, likely via modulation of *JAKMIP1*.

311

Figure 3 | CYFIP1-OE alters the expression of other cytokine signal-regulated

transcription factors. *Relative mRNA expression of **A)** STAT1, **B)** STAT3, **C)** STAT5B or **D)***

NFKB1 in Control and CYFIP1-OE cells measured by qRT-PCR. For all experiments, values

are presented as mean ± SEM of N = 3 independent experiments, 2-3 technical replicates

each. Statistical significance against Control was determined using Unpaired Student's

T-Tests; *P < 0.05; **P <0.01; ****P < 0.0001.

***CYFIP1*-OE enhances IL-6-induced STAT3 activity**

To investigate whether reduced *STAT3* expression in *CYFIP1*-OE HEK293 cells leads to

altered cytokine-induced STAT3 responses we first performed Western blotting to assess

IL-6-induced STAT3^{Y705} phosphorylation (representative of STAT3 activation) (**Figure 4A**).

Despite the reduction in STAT3 expression (**Figure 4B**), *CYFIP1*-OE cells still show strong

responses to IL-6, demonstrating no significant difference in IL-6-stimulated STAT3^{Y705}

phosphorylation levels compared to control cells, both at the 'whole-cell level' (when

STAT3^{Y705} phosphorylation levels are normalised to GAPDH) (**Figure 4C**) and upon closer

examination of the phosphorylation kinetics of STAT3 itself (when STAT3^{Y750}

phosphorylation levels are normalised to total STAT3 protein levels) (**Figure 4D**). This

suggests that although *CYFIP1*-OE cells show reduced STAT3 expression, STAT3

activation following IL-6 stimulation is unaltered.

STAT3 activity is modulated by various post-translational modifications in addition to

phosphorylation at the Y705 residue. Therefore, despite observing no change in STAT3^{Y705}

phosphorylation following *CYFIP1*-OE, we examined whether *CYFIP1*-OE cells show

altered IL-6-induced transcriptional activity by performing a DLR™ assay. This assay

places the control of a Firefly luciferase under a STAT3 response element, meaning that

increased Firefly luminescence is indicative of increased STAT3 transcriptional activity.

Using this DLR™ assay, we found that *CYFIP1*-OE cells show increased Firefly

luminescence following IL-6 stimulation than control cells (**Figure 4E**), suggesting that

IL-6-induced STAT3 transcriptional activity is enhanced in *CYFIP1*-OE cells regardless of

the reduced STAT3 expression.

To validate the DLR™ assay results, qRT-PCR was used to measure the mRNA expression

of STAT3-responsive genes in *CYFIP1*-OE cells following IL-6 stimulation. Somewhat

contradictory to the DLR™ assay, levels of *STAT3* mRNA in IL-6-treated *CYFIP1*-OE cells

were not higher than those in untreated control cells (**Figure 4F**). However, considering

that *CYFIP1*-OE cells already show reduced *STAT3* mRNA expression at baseline, this is

unsurprising. Therefore, after normalising the levels of IL-6-stimulated *STAT3* expression

to unstimulated *STAT3* expression, *CYFIP1*-OE cells show an enhancement in IL-6-

induced *STAT3* mRNA expression compared to control cells (**Figure 4G**). The expression

of *SOCS3* shows a similar phenomenon whereby IL-6-stimulated *CYFIP1*-OE cells

appear to show lower *SOCS3* mRNA levels compared to IL-6-treated control cells (**Figure**

4H), however the fold change in *SOCS3* expression relative to unstimulated cells shows

that IL-6-induced *SOCS3* expression is magnified in *CYFIP1*-OE cells (**Figure 4I**). Taken

together, these findings complement the DLR™ assay results, suggesting that *CYFIP1*-OE

cells show enhanced STAT3 transcriptional activity following IL-6 stimulation.

Figure 4 | CYFIP1-OE enhances STAT3 transcriptional activity but not STAT3 activation. A) *Western blotting of STAT3 expression and phosphorylation (P-STAT3^{Y705}) in Control and CYFIP1-OE cells following 30-minute treatment with 0 ng/mL IL-6 or 20 ng/mL*

*IL-6. Densitometric analysis of **B)** total STAT3 protein expression relative to the GAPDH loading control, **C)** P-STAT3 relative to the GAPDH loading control or **D)** P-STAT3 relative to total STAT3 protein expression. **E)** Total transcriptional activity at a STAT3-regulated promoter in Control and CYFIP1-OE cells stimulated with or without 20 ng/mL IL-6 for 18 hours post-transfection with Cignal Reporter plasmids. Firefly luminescence intensity is expressed as a ratio to Renilla luminescence intensity to adjust for transfection efficiency. **F)** Relative mRNA expression of STAT3 in Control and CYFIP1-OE cells treated with 0 ng/mL or 20 ng/mL IL-6 for 4 hours, measured by qRT-PCR. **G)** Fold change in STAT3 mRNA expression following IL-6 treatment relative to untreated cells in part F). Values presented as mean ± SEM of N = 3 independent experiments, 3 technical replicates each. Statistical significance against Control was determined using Unpaired Student's T-Tests; ****P < 0.0001. **H)** Relative mRNA expression of SOCS3 in Control and CYFIP1-OE cells treated with 0 ng/mL or 20 ng/mL IL-6 for 4 hours, measured by qRT-PCR. **I)** Fold change in SOCS3 mRNA expression following IL-6 treatment relative to untreated cells in part H). For all experiments, values are presented as mean ± SEM of N = 3 independent experiments, 3-4 technical replicates each. Statistical significance against Control was determined using Unpaired Student's T-Tests when comparing two groups, or two-way ANOVAs with Tukey's Honest Significant Difference tests when comparing multiple groups; ns – P > 0.05; *P < 0.05; **P < 0.01; ***P < 0.001; ****P < 0.0001.*

CYFIP1-OE enhances IL-6-induced neuritogenesis

To understand how this enhanced STAT3 response to IL-6 in *CYFIP1*-OE cells observed at a molecular level affects a broader IL-6-stimulated cellular process, we differentiated *CYFIP1*-OE SH-SY5Y cells in the presence of IL-6 and measured the neurite lengths

produced (**Figure 5A**), as we have previously shown that IL-6 induces neurite outgrowth

in SH-SY5Y cells (6). As expected, control and *CYFIP1*-OE cells differentiated in the

presence of IL-6 exhibit longer LNL (**Figure 5B**) and TNL (**Figure 5C**) measurements than

cells differentiated without IL-6. *CYFIP1*-OE cells appear to have enhanced neurite

outgrowth responses to IL-6, as although *CYFIP1*-OE cells produce shorter neurites

compared to the control as baseline, the LNL (**Figure 5B**) and TNL (**Figure 5C**)

measurements are no longer significantly different between the two populations when

differentiated with IL-6. These data appear to agree with the suggestion that *CYFIP1*-OE

leads to enhanced STAT3 responses to IL-6, and that this enhancement in IL-6-induced

STAT3 activity leads to increased IL-6-stimulated neurite extension and production.

Figure 5 | CYFIP1-OE enhances IL-6-induced neuritogenesis. SH-SY5Y cells were

Nucleofected with either the RFP-T2A backbone vector and pCAβ-YFP as the control

condition; or RFP-T2A-CYFIP1 and pCAβ-YFP to assess the effects of CYFIP1-OE on IL-6-

influenced neuritogenesis. Cells were differentiated for two days with 20 ng/mL BDNF

and 10 μM RA under serum starvation with or without 20 ng/mL IL-6, fixed with 4% PFA

and stained with DAPI. **A)** *Fluorescence micrographs of differentiated SH-SY5Y cells.*

Images captured using an upright Leica DM4000 B LED microscope. Scale bar = 50 µm.

*Neurite tracing analysis was performed using the FIJI software. **B)** The length of the*

*longest neurite produced by the cell was measured for all conditions. **C)** The sum length*

of all neurites produced per cell is presented for each condition. For all experiments,

values are presented as mean ± SEM of N = 4 independent experiments, 3 technical

replicates each. Statistical significance against Control was determined using two-way

ANOVAs with Tukey's Honest Significant Difference tests; ns – P > 0.05; *P < 0.05; **P <

0.01; ****P < 0.0001.

***CYFIP1*-OE enhances IFN-γ-induced STAT1 transcriptional activity**

Having determined that *CYFIP1*-OE alters the expression of several cytokine signalling-

related transcription factors, and that *CYFIP1*-OE enhances IL-6-induced STAT3 activity,

we aimed to investigate whether *CYFIP1*-OE alters the activity of another cytokine

signalling-related transcription factor. STAT1 was chosen to follow up with because

similar to *STAT3* expression, the expression of *STAT1* was also reduced in *CYFIP1*-OE

HEK293 cells. A DLR™ assay which placed the control of a Firefly luciferase under a GAS

element (the response element STAT1 binds to initiate transcription) was performed. We

found that *CYFIP1*-OE HEK293 cells show increased Firefly luminescence following IFN-

γ stimulation compared to control cells (**Figure 6A**), suggesting that IFN-γ-induced STAT1

transcriptional activity is enhanced in *CYFIP1*-OE cells, again regardless of the reduced

STAT1 expression.

As validation of the DLR™ assay, qRT-PCR was performed to measure the expression of

STAT1-response genes following IFN-g stimulation. *CYFIP1*-OE cells show increased IFN-

γ-induced *STAT1* expression (**Figure 6B**), suggesting an enhanced transcription response

to IFN-γ stimulation. Additionally, although not significantly different at baseline, *CYFIP1*-

OE cells show increased *SOCS1* expression following IFN-γ treatment (**Figure 6C**), again

suggesting an enhanced STAT1 transcriptional response to IFN-γ stimulation. This also

suggests that the observed alterations in IL-6/STAT3 signalling are not limited to STAT3

and that CYFIP1 may play a role in modulating a larger range of cytokine signalling

pathways.

Figure 6 | CYFIP1-OE leads to enhanced STAT1 transcriptional activity following IFN-

γtreatment. A) *Total transcriptional activity at a STAT1-regulated promoter in Control and*

CYFIP1-OE cells stimulated with or without 20 ng/mL IFN-γfor 18 hours post-transfection

with Cignal Reporter plasmids. Firefly luminescence intensity is expressed as a ratio to

Renilla luminescence intensity to adjust for transfection efficiency. ***B)*** *Relative mRNA*

expression of STAT1 in Control and CYFIP1-OE cells treated with 0 ng/mL or 20 ng/mL IFN-

γ for 4 hours, measured by qRT-PCR. **C)** *Relative mRNA expression of SOCS1 in Control*

and CYFIP1-OE cells treated with 0 ng/mL or 20 ng/mL IFN-γ for 4 hours, measured by

qRT-PCR. For all experiments, values are presented as mean ± SEM of N = 4 independent

experiments, 3 technical replicates each. Statistical significance against Control was

determined using two-way ANOVAs with Tukey's Honest Significant Difference tests; ns –

P > 0.05; *P < 0.05; ***P < 0.001; ****P < 0.0001.

Discussion

A novel role for CYFIP1 in modulating IL-6/STAT3 signalling independent from

JAKMIP1

Nishimura *et al*. (5) first linked *JAKMIP1* to ASD and Dup(15q), reporting that *JAKMIP1*

expression was altered in Dup(15q) and that overexpression of the chromosome 15q

gene *CYFIP1* correspondingly resulted in *JAKMIP1* downregulation. We expanded upon

this in our previous work, identifying JAKMIP1 as a modulator of IL-6 cytokine signaling in

SH-SY5Y cells, and demonstrating that Dup(15q) iPSCs and cortical neurons with

elevated *CYFIP1* dosage and expression exhibit enhanced IL-6-induced STAT3 activity (6).

However, given that the magnified IL-6 signaling observed in these Dup(15q) cells did not

correlate with *JAKMIP1* expression levels, we hypothesised that another gene within the

15q region, likely *CYFIP1*, might be driving this increased responsiveness to IL-6.

To test this hypothesis, we induced *CYFIP1* overexpression in HEK293 cells and

discovered that this resulted in the altered expression of several cytokine signaling-

related transcription factors. In particular, *STAT1* and *STAT3* expression were reduced, whereas *STAT5B* and *NFKB1* expression were raised, suggesting that CYFIP1 may potentially exert broad cellular functions that impact the signaling networks of multiple cytokines. Taking this forward, we measured IL-6-induced activation and activity of STAT3. Consistent with the Dup(15q) iPSCs and neurons from our previous work, we observed that despite the reduced *STAT3* expression, IL-6-induced STAT3 responses were enhanced in *CYFIP1*-OE HEK293 cells. Furthermore, *CYFIP1* overexpression also potentiated IL-6-triggered neuritogenesis in SH-SY5Y cells. As such, this study provides the first evidence that CYFIP1 may play a role in regulating IL-6/STAT3 signalling in both HEK293 and SH-SY5Y cells, and its upregulation could be responsible for the enhanced responsiveness to IL-6 exhibited by Dup(15q) iPSCs and neurons.

How does CYFIP1 influence STAT1 and STAT3 activity?

The mechanisms underlying CYFIP1-mediated regulation of *STAT1* and *STAT3* expression and activity remain unclear. Regarding STAT3, reduced *STAT3* expression in *JAKMIP1*-deficient cells (6) may suggest that JAKMIP1, rather than CYFIP1, is responsible for regulating *STAT3* expression. This could occur through its interactions with various RNA-binding proteins (RBPs) or splicing factors (6). However, it is important to note that *CYFIP1* overexpression enhances IL-6-induced STAT3 transcriptional activity, whereas *JAKMIP1*-knockout impairs it (6). Thus, whilst JAKMIP1 could be responsible for controlling *STAT3* expression, when considering STAT3 activity, our results indicate that it is more likely that CYFIP1 is responsible by a mechanism independent of JAKMIP1.

Interestingly, despite *CYFIP1* overexpression leading to increased STAT3 transcriptional

activity following IL-6 treatment, we observed no significant difference in the levels

STAT3^{Y705} phosphorylation in *CYFIP1*-OE cells, suggesting that CYFIP1 does not alter

STAT3 phosphorylation kinetics, and thus STAT3 activation. One reason for this

discrepancy could be because of the influence of other post-translational modifications

of STAT3. For example, STAT3^{K685} acetylation may be necessary for IL-6-induced STAT3-

responsive gene expression (16). Rather than modulating STAT3 activation through

phosphorylation, perhaps CYFIP1 may somehow control STAT3 activation through

acetylation, however, the exact mechanism by which CYFIP1 could regulate acetylation

is unclear.

In terms of *STAT1* expression and activity, it would not be unreasonable to suggest that

these changes in STAT1 activity may be mediated by the changes in *JAKMIP1* expression.

JAKMIP1 was previously reported to inhibit Interferon-α (IFN-α) signalling (17).

Importantly, both IFN-α and IFN-γ signalling cascades involve the phosphorylation and

activation of STAT1. It is therefore likely that CYFIP1-induced downregulation of *JAKMIP1*

may reduce the inhibiting influence of JAKMIP1 on IFN signalling, thereby enhancing the

IFN response. Furthermore, STAT1 activity is also known to be altered by acetylation,

similar to STAT3 (18). If CYFIP1-mediated regulation of STAT3 is dependent on post-

translational modifications, the same mechanism(s) could be responsible for CYFIP1

regulation both STAT3 and STAT1 activity. Of course, all of these possibilities would

require further validation.

Relevance to ASD and neurodevelopment

Cytokine signalling plays crucial roles in neurodevelopment beyond immune responses,

with STAT3 and STAT1 signalling in particular regulating several key processes. For

example, STAT3 signalling has been demonstrated to control the switch between

neurogenesis and gliogenesis, promoting both neural progenitor cell self-renewal and

astrocyte differentiation (19). Similarly, STAT1 signalling has been shown to be involved

in neurogenesis, and promoting neurite outgrowth (20). Given how crucial STAT3 and

STAT1 signalling are for neurodevelopment, it is unsurprising that alterations in STAT3 and

STAT1 signalling are associated with NDDs such as ASD (21,22). MIA, where an immune

response is triggered by infection of exposure to immunogenic material during

pregnancy, is associated with increased incidence of ASD (23). Both IL-6 and IFN-γ are

pro-inflammatory cytokines that have been implicated in the pathogenesis of ASD, with

IL-6 being a key player in MIA-induced ASD-like behaviours (23,24). Considering that

CYFIP1-OE enhances STAT3 and STAT1 responses to IL-6 and IFN-γ respectively suggests

that CYFIP1 may play a pivotal role in amplifying the inflammatory signals during MIA,

potentially exacerbating the effects of immune activation on foetal brain development.

Amplification of both the IL-6/STAT3 and IFN-γ/STAT1 signalling pathways by CYFIP1

provides a potential molecular mechanism that explains the overlap between immune

activation and ASD, particularly in cases where no obvious inflammatory trigger is

identified. Elevated cytokine levels are frequently reported in individuals with ASD (25),

and our results suggest that *CYFIP1* overexpression could act as a molecular

convergence point, increasing the sensitivity of the developing brain to immune signals

even under baseline inflammatory conditions. This could help to explain cases of ASD

where immune activation is not readily apparent but inflammation-induced

neurodevelopmental disruption still occurs.

Conclusion

In summary, this study identifies a novel role for CYFIP1 in regulating the expression of

cytokine signalling-related transcription factors, as well as enhancing their activity

following cytokine stimulation. By amplifying these inflammatory pathways, CYFIP1

could play a central role in exacerbating the neuroinflammatory environment, leading to

altered neurodevelopment and increased susceptibility to ASD. Further research is

required to understand the mechanism by which CYFIP1 modulates these cytokine

signalling pathways and to investigate this in the context of ASD with MIA.

List of abbreviations

ASD – Autism spectrum disorder

Dup(15q) – Chromosome 15q-duplication syndrome

CYFIP1 – Cytoplasmic FMR1-intercating protein 1

iPSC – Induced pluripotent stem cell

IL-6 – Interleukin-6

STAT3 – Signal transducer and activator of transcription 3

OE – Overexpression

MIA – Maternal immune activation

JAKMIP1 – Janus kinase and microtubule-interacting protein 1

FXS – Fragile X syndrome

iNeurons – iPSC-derived neurons

FMRP – Fragile X messenger ribonuclear protein

WRC – WAVE regulatory complex

SMAD3 – SMAD family member 3

NF-kB – Nuclear factor of κB

ISRE – Interferon response element

TGF-β1 – Transforming growth factor β1

GAPDH – Glyceraldehyde-3-phosphate dehydrogenase

DLR™ – Dual-luciferase reporter™

SOCS3 – Suppressor of cytokine signalling 3

LNL – Longest neurite length

TNL – Total neurite length

IFN-γ - Interferon-γ

RBP – RNA-binding protein

<u>Declarations</u>

Ethics approval and consent to participate:

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

This work was supported by a Wellcome Trust funded Translational Research Exchange

@ Exeter (TREE) pump-priming fund and the Bristol Japanese Cultural Showcase

Author's contributions

E-R.M. conceived and designed the research, performed the laboratory research,

generated the data and figures, and contributed to manuscript writing and editing. J.G.M.

provided critical input and contributed to manuscript editing. M.A.R. provided critical

input and reviewed the manuscript. A.O-A. conceived and designed the research,

provided critical input, reviewed the manuscript, and secured funding. All authors

reviewed and approved the final manuscript.

Acknowledgements

The research was carried out at the National Institute for Health and Care Research

(NIHR) Exeter Biomedical Research Centre (BRC). We gratefully acknowledge the

generous support of the charitable organisations that contributed to this work, including

the Wellcome Trust Translational Research Exchange @ Exeter (TREE) pump-priming

fund and the Bristol Japanese Culture Showcase. We thank Liming Yang for valuable

discussions during lab meetings, and we are also grateful to Dr. Kaiyven Afi Leslie for his

technical advice and support.

Corresponding author

Dr. Asami Oguro-Ando (ORCID: https://orcid.org/0000-0003-4029-3250)

References

1.   American Psychiatric Publishing. . American Psychiatric Association Diagnostic

and statistical manual of mental disorders, fifth edition [DSM-5®]. American

Psychiatric Publishing. 2016;51(13).

2.   Abrahams BS, Arking DE, Campbell DB, Mefford HC, Morrow EM, Weiss LA, et al.

SFARI Gene 2.0: A community-driven knowledgebase for the autism spectrum

disorders (ASDs). Mol Autism. 2013;4(1).

3.   Modabbernia A, Velthorst E, Reichenberg A. Environmental risk factors for

autism: an evidence-based review of systematic reviews and meta-analyses. Vol.

8, Molecular Autism. 2017.

4.   Zawadzka A, Cieślik M, Adamczyk A. The role of maternal immune activation in

the pathogenesis of autism: A review of the evidence, proposed mechanisms and

implications for treatment. Vol. 22, International Journal of Molecular Sciences.

2021.

5.   Nishimura Y, Martin CL, Vazquez-Lopez A, Spence SJ, Alvarez-Retuerto AI, Sigman

M, et al. Genome-wide expression profiling of lymphoblastoid cell lines

distinguishes different forms of autism and reveals shared pathways. Hum Mol

Genet [Internet]. 2007 Jul 15 [cited 2019 Jan 20];16(14):1682–98. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/17519220

6.   Martin JG, Martin ER, Takamura N, Harlow CE, Bamford RA, Smith RG, et al.

Exploring the functions of JAKMIP1 in neuronal IL-6/STAT3 signaling and its

relevance to chromosome 15q-duplication syndrome. 2025.

7.   DiStefano C, Gulsrud A, Huberty S, Kasari C, Cook E, Reiter LT, et al. Identification

of a distinct developmental and behavioral profile in children with Dup15q

syndrome. J Neurodev Disord [Internet]. 2016 Dec 6 [cited 2019 May 30];8(1):19.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/27158270

8.   Butler MG, Bittel DC, Kibiryeva N, Talebizadeh Z, Thompson T. Behavioral

Differences among Subjects with Prader-Willi Syndrome and Type I or Type II

Deletion and Maternal Disomy. Pediatrics. 2004;113(3 I).

9.   Oguro-Ando A, Rosensweig C, Herman E, Nishimura Y, Werling D, Bill BR, et al.

Increased CYFIP1 dosage alters cellular and dendritic morphology and

dysregulates mTOR. Mol Psychiatry. 2015;20(9).

10.   Napoli I, Mercaldo V, Boyl PP, Eleuteri B, Zalfa F, De Rubeis S, et al. The Fragile X

Syndrome Protein Represses Activity-Dependent Translation through CYFIP1, a

New 4E-BP. Cell. 2008;134(6).

11.   DeRubeis S, Pasciuto E, Li KW, Fernández E, DiMarino D, Buzzi A, et al. CYFIP1

coordinates mRNA translation and cytoskeleton remodeling to ensure proper

dendritic Spine formation. Neuron. 2013;79(6).

12.   Urraca N, Hope K, Victor AK, Belgard TG, Memon R, Goorha S, et al. Significant

656   transcriptional changes in 15q duplication but not Angelman syndrome deletion

657   stem cell-derived neurons. Mol Autism. 2018;9(1).

658   13.   Erickson EK, Blednov YA, Harris RA, Mayfield RD. Glial gene networks associated

659   with alcohol dependence. Sci Rep. 2019;9(1).

660   14.   Pfaffl MW. A new mathematical model for relative quantification in real-time RT–

661   PCR. Nucleic Acids Res. 2001;29(9):e45.

662   15.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji:

663   an open-source platform for biological-image analysis. Nat Methods. 2012 Jun

664   28;9(7):676–82.

665   16.   Wang R, Cherukuri P, Luo J. Activation of Stat3 sequence-specific DNA binding

666   and transcription by p300/CREB-binding protein-mediated acetylation. Journal of

667   Biological Chemistry. 2005;280(12).

668   17.   Steindler C, Li Z, Algarté M, Alcover A, Libri V, Ragimbeau J, et al. Jamip1 (Marlin-1)

669   defines a family of proteins interacting with Janus kinases and microtubules.

670   Journal of Biological Chemistry. 2004;279(41).

671   18.   Leslie KA, Lekka C, Richardson SJ, Russell MA, Morgan NG. Regulation of STAT1

672   Signaling in Human Pancreatic β-Cells by the Lysine Deacetylase HDAC6: A New

673   Therapeutic Opportunity in Type 1 Diabetes? Diabetes [Internet]. 2024 Aug 20

674   [cited 2025 Sep 4];73(9):1473–85. Available from:

675   https://dx.doi.org/10.2337/db24-0008

676   19.   Hong S, Song MR. STAT3 but not STAT1 is required for astrocyte differentiation.

677   PLoS One. 2014;9(1).

678   20.   Warre-Cornish K, Perfect L, Nagy R, Duarte RRR, Reid MJ, Raval P, et al. Interferon-

679   γ signaling in human iPSC–derived neurons recapitulates neurodevelopmental

680   disorder phenotypes. Sci Adv. 2020;6(34).

681   21.   Garay PA, McAllister AK. Novel roles for immune molecules in neural

682   development: Implications for neurodevelopmental disorders. Frontiers in

683   Synaptic Neuroscience. 2010.

684   22.   Majerczyk D, Ayad EG, Brewton KL, Saing P, Hart PC. Systemic maternal

685   inflammation promotes ASD via IL-6 and IFN-γ. Vol. 42, Bioscience Reports. 2022.

686   23.   Choudhury Z, Lennox B. Maternal Immune Activation and Schizophrenia–

687   Evidence for an Immune Priming Disorder. Vol. 12, Frontiers in Psychiatry. 2021.

688   24.   Samuelsson AM, Jennische E, Hansson HA, Holmäng A. Prenatal exposure to

689   interleukin-6 results in inflammatory neurodegeneration in hippocampus with

690   NMDA/GABAA dysregulation and impaired spatial learning. Am J Physiol Regul

691   Integr Comp Physiol. 2006;290(5).

692   25.   Ashwood P, Krakowiak P, Hertz-Picciotto I, Hansen R, Pessah I, Van de Water J.

693   Elevated plasma cytokines in autism spectrum disorders provide evidence of

694   immune dysfunction and are associated with impaired behavioral outcome. Brain

695   Behav Immun. 2011;25(1).

Supplementary Files

Supplementary Figure 1 | Modelling CYFIP1 overexpression in HEK293 cells.

Increased CYFIP1 protein expression in CYFIP1-OE cells 72 hours post-transfection

confirmed via Western blotting. Membrane probed with: Rabbit Anti-CYFIP1

(#ab156016, Abcam), Rabbit Anti-FMRP (#13755-1-AP, Proteintech), Mouse Anti-GAPDH

(#sc-47724, Santa Cruz Biotechnology), Goat Anti-Mouse DyLight™-680 (#35519,

Invitrogen) and Goat Anti-Rabbit DyLight™-800 (#SA5-10036, Invitrogen).

Supplementary Figure 2 | CYFIP1-OE leads to reduced JAKMIP1 protein expression.

Western blotting of JAKMIP1 expression in Control and CYFIP1-OE cells. Membrane

probed with: Rabbit Anti-JAKMIP1 (#13846-1-AP, Proteintech), Mouse Anti-GAPDH (#sc-

47724, Santa Cruz Biotechnology), Goat Anti-Mouse DyLight™-680 (#35519, Invitrogen)

and Goat Anti-Rabbit DyLight™-800 (#SA5-10036, Invitrogen). Known non-specific bands

produced are indicated on the blot.

Supplementary Figure 3 | CYFIP1-OE has no effect on STAT3 activation. *Western*

blotting of STAT3 expression and phosphorylation (P-STAT3^{Y705}) in Control and CYFIP1-OE

cells following 30-minute treatment with 0 ng/mL IL-6 or 20 ng/mL IL-6. Membrane

probed with: Mouse Anti-STAT3 (#9139S, Cell Signaling Technology), Rabbit Anti-P-

STAT3^{Y705} (#9131l, Cell Signaling Technology), Mouse Anti-GAPDH (#sc-47724, Santa

Cruz Biotechnology), Goat Anti-Mouse DyLight™-680 (#35519, Invitrogen) and Goat Anti-

Rabbit DyLight™-800 (#SA5-10036, Invitrogen). Known non-specific bands are indicated

on the blot.