Cardiac synchrony remains stable across repeated intergenerational encounters but is enhanced during high stakes collaboration

Ryssa Moffat[1], Luca A. Naudszus[1], Emily S. Cross[1]

[1]Social Brain Sciences Lab, ETH Zurich

* Correspondence:
Ryssa Moffat
rmoffat@ethz.ch

Emily S. Cross
ecross@ethz.ch

Keywords: intergenerational, collaboration, art-making, physiological synchrony, relationship
formation

Abstract

Intergenerational social programs provide opportunities for people of all ages to form new
relationships. Furthermore, existing qualitative and behavioural evidence from such programs points
to health and wellbeing benefits, yet the physiological consequences of repeated intergenerational
encounters, particularly those involving collaboration, remain unknown. A deeper understanding of
how such programs shape physiological responses in dyads will illuminate the mechanisms of
relationship formation. Across a six-session creative art-making program, we tracked cardiac
synchrony (CS) within 31 intergenerational (older/younger adult) and 30 same generation (younger
adult) dyads. Each session dyads completed self-report measures, then drew together and alone,
while we recorded participants' actions with motion capture and their physiological signals (neural
and cardiac) using fNIRS. Collaborative behaviour, self-reported social closeness, and interpersonal
distance (i.e., proximity) showed group-specific patterns, whereby interpersonal distance emerged as
a promising objective measure of relationship development. CS did not covary with group, task, an
interaction thereof or any measure of behaviour or social closeness–yet there was a trending
relationship between collaboration while drawing together and CS for intergenerational dyads only.
This work has implications for understanding the mechanisms underpinning intergenerational social
interaction and for the implementation of CS as an objective measure of social connection or
collaboration.

1   Introduction

In response to worrying reports of declining social connection, many communities have begun to
employ intergenerational arts programs to improve community members' health and wellbeing
(WHO, 2019, 2025). Growing behavioural and qualitative data from such programs suggests that
intergenerational arts programs offer much needed opportunities to build social connection, reduce
stereotypes about other generations, and provide structures for people to engage in activities that add
meaning to their lives (Anderson et al., 2017; Jenkins et al., 2021; Lokon et al., 2020; Rubin et al.,
2015; WHO, 2025). Intergenerational arts programs that offer repeated contact between generations,
as opposed to one-off sessions, seem to yield the greatest improvement in social connectedness and
feelings of affiliation between generations (Lokon et al., 2020). More broadly, feelings of social

connectedness or closeness (e.g., being strangers vs. friends vs. romantic partners) are known to
modulate physiological signals (e.g., heart rate and brain activity) during collaboration (Mayo et al.,
2021). Yet, longitudinal evidence tracking the extent to which physiological signals index
relationship formation is only just beginning to emerge (Moffat et al., 2025). A deeper understanding
of how repeated intergenerational collaboration fosters relationship formation, across experiential and
physiological levels, holds potential to inform program structure and resource allocation for
intergenerational arts programs. Moreover, new insights into intergenerational collaborative
behaviour can only strengthen efforts to promote healthy ageing.
46
In this study, intergenerational and same generation dyads completed a 6-week art-making program,
in which they met as strangers and interacted with the same partner for all 6 sessions. We examined
how collaboration and physiological synchrony, specifically *cardiac synchrony (CS)*, evolved across
sessions. Specifically, we tracked how CS levels during collaboration were shaped by repeated social
encounters, the generational constellation (intergenerational vs. same generation), as well as dyads'
feelings of social closeness and interpersonal distance (i.e., physical proximity to one another).

1.1   Cardiac synchrony an index of shared arousal during collaboration

The cardiovascular system is reactive to changes in both psychological and physiological factors
(Oliveira-Silva & Gonçalves, 2011). Fluctuations in heart rate reflect changes in arousal of the
central nervous system resulting from changes in an individual's environment (Wascher, 2021).
Increases in heart rate reflect sympathetic activity, which is associated with excitement and stress.
Reductions in heart rate reflect parasympathetic activity, relating to relaxation and divided attention
(Danyluck & Page-Gould, 2019; Tschacher & Meier, 2020). When people gather in groups of two or
more, changes in their physiological signals including heart rate reflect a mixture of self-regulation
and co-regulation (Butler, 2011; Ferrer & Helm, 2013), which can result in greater temporal
alignment of heart rates among two or more people. Precise time-locking of fluctuations can arise
from external environmental factors–take for example a jumpscare (i.e., a startling event) in a horror
movie causing a spike in cinemagoers' heart rates simultaneously. Time-lagged alignment occurs
more commonly in live social interaction involving behavioural coordination, for example turn
taking in conversations (Coutinho et al., 2021). Such alignment in arousal can result from shared
attention (Golland et al., 2015), emotional contagion (Bizzego et al., 2019; Park et al., 2019), or joint
action (Noy et al., 2015; Palumbo et al., 2017). In other words, measures of temporal alignment of
heart rates between individuals, or CS, can shed light on shared fluctuations in physiological arousal
during social interactions. Accordingly, shared fluctuations in arousal, tracked using measures of CS,
are reported to underpin important aspects of collaboration, including perceptions of group cohesion,
satisfaction, commitment, subjective togetherness, trust, and perceived empathy (for review see:
Mayo et al., 2021).

Several studies have investigated the extent to which CS can predict the outcomes of collaboration
(Boukarras et al., 2025; Gordon et al., 2020; Mitkidis et al., 2015; Noy et al., 2015; Sharika et al.,
2024). Sharika et al. (2024) recorded 44 groups of 204 students completing a verbal consensus-
seeking task. The authors reported that CS between group members predicted the likelihood of
reaching consensus significantly better than individuals' heart rates, self-reported ratings of team
function, or duration of the consensus-seeking process. On this basis, Sharika et al. proposed that CS
may be useful as a biomarker of engagement during collaboration (Sharika et al., 2024). Gordon et al.
(2020) compared levels of CS during synchronised drumming, freely improvised drumming, and
resting state (i.e., no movement). They reported that CS was greater during drumming than baseline
but did not differ between drumming conditions, highlighting the role of joint action in aligning

fluctuations in arousal. Gordon et al. further reported that levels of CS during synchronised
drumming predicted levels of coordination of drumming in the subsequent improvised drumming
task. Alignment of arousals levels, as indexed by CS, may thus be a precursor for successful
collaboration (Gordon et al., 2020).

Other studies suggest that perceptual shifts during collaboration drive changes in shared arousal
(Boukarras et al., 2025; Mitkidis et al., 2015; Noy et al., 2015). Noy et al. tracked levels of perceived
and kinematically measured togetherness when dyads played a mirror game. Controlling for
differences in movement intensity, the authors found that CS increased during periods of
togetherness, more so for perceived than kinematic togetherness. Periods of togetherness were also
notably accompanied by increased heart rates, reflecting increased arousal (Noy et al., 2015).
Boukarras et al. (2025) recorded grasp timing from dyads told to grasp two bottle shaped objects
according to specific instructions. In half the trials, the instructions prompted the same or
complementary grasping behaviours, and in the other half of trials, the instructions prompted evoked
peer-to-peer or leader-follower grasp dynamics. None of these prompted grasping behaviours showed
a statistically significant relationship between grasping timing and CS. However, Boukarras et al. did
observe heightened levels of CS when dyads shifted between peer-to-peer to a leader-follower
dynamics or from same to complementary movements. This findings led the authors to suggest that
perceptual shifts during collaboration dynamics, rather than specific patterns of behavioural
coordination, may influence arousal levels, and by extension, CS (Boukarras et al., 2025). Findings
from a study by Mitkidis et al. (2015) lend support to the idea that perceptual shifts during
collaboration may alter shared arousal. Mitkidis et al. compared CS in a group that completed
interleaved collaborative Lego building tasks with public goods games to a group that completed
only Lego building tasks. They report that the public goods game led to greater CS during Lego
building. They also found that CS predicted the participants' expectations of their partners during the
public goods game. Mitkidis et al. interpret both results as being reflective of increased awareness of
the partner relating to trust-building processes. Considered together, these findings highlight that
perceptual shifts during collaboration impact arousal levels as quantified by CS. Notably, too much
shared arousal can have deleterious effects on collaboration (Dimitroff et al., 2017; Mayo & Gordon,
2020), suggesting that co-regulation of arousal levels is central to successful collaboration (Mayo et
al., 2021).

Changes in familiarity that occur within or across encounters may represent a sufficient perceptual
shifts to influence shared arousal, and by extension, CS. Fusaroli et al. (2016) invited five groups of
4-5 students to build Lego models in 5-minute blocks, alternating between building alone and
together as a group, and computed measures of heart rate (HR) coordination (i.e., the similarity and
stability of signals within groups–complementary measures to CS). The authors reported that HR
coordination within groups increased from the first block of collaborative model-building to the last
and that coordination of movement and speech within groups predicted HR coordination (Fusaroli et
al., 2016). These findings illustrate how increasing familiarity, on a short time scale, co-occurs with
increasing alignment of behaviour and arousal levels. Highlighting the relevance of understanding
changes in dyadic levels of shared arousal over repeated encounters, Gernert et al. (2025) measured
CS during psychotherapy sessions targeting affective disorders at two sessions, approximately two
weeks apart. The authors found a positive association between CS across both sessions and changes
in self-reported depressive symptom severity between the two sessions (Gernert et al., 2025).
Unfortunately, this study does not report the difference in CS between sessions, leaving the trajectory
of shared arousal an open question. While empirical attention has been paid to first encounters (e.g.,
Adel et al., 2025), little is known about the trajectory of fluctuations in shared arousal across repeated
encounters. To advance our understanding of the prerequisites for relationships formed over repeated

encounters to blossom into robust social bonds and the conditions that predict relationships fizzling
out, we need to document the trajectories of shared arousal across multiple encounters alongside
measures of relationship quality. By considering these trajectories in same generation and
intergenerational dyads, we can provide insights for optimising relationship formation in
intergenerational community programs.

1.2   Relationship quality constrains shared arousal, and by extension, cardiac synchrony

Relationship quality and shared fluctuations in arousal, as measured by CS, are known to covary,
though the effect size derived from a meta-analysis of 26 studies is quite small (Mayo et al., 2021),
and the directionality of the effect is not known. Many of the studies conducted have been dedicated
to understanding shared arousal in romantic partners and are summarised in Mayo et al.'s meta-
analysis (2021). For example, watching emotional videos together has been shown to elicit greater
CS between strangers, relative to friends and romantic partners (Bizzego et al., 2019). Bizzego et al.
propose that greater familiarity between friends and romantic couples results in their autonomic
systems being less attuned to changes in the other person's autonomic responses. This may be a
feature of the task involving non-interactive copresence with both people's attention directed to the
videos. Other tasks, involving co-presence and interaction show the opposite pattern. At a
firewalking performance, Konvalinka et al. (2011) recorded CS between firewalkers and their friends
and between firewalkers and strangers. The authors compared these groups and reported greater CS
between firewalkers and friends and family members than strangers. The authors similarly suggest
that social closeness (Konvalinka et al., 2011) is a driving factor in the coupling of autonomic
responses. Focusing on first-time encounters, Adel et al. (2025) recorded CS in dyads comprising
strangers who shared emotional stories. Dyads showed greater CS when they self-reported high
levels of mutual liking. Relationship quality seems to have a more nuanced impact on shared arousal
when collaboration is compared with competition. Danyluck and Page-Gould (2019) demonstrated
that self-reported perceived similarity and interest in friendship between dyad members can shape CS
levels differentially for collaboration and competition. Whereas CS is positively associated with
perceived similarity in both conditions, CS is positively associated with interest in friendship only
during collaboration and not during competition (Danyluck & Page-Gould, 2019). Recently,
Andersen et al. (2025) recorded CS between visitors who went through a recreational commercial
haunted house in groups, designed to induce fear in the visitors. The authors reported that CS is
closely related to individuals' ratings of subjective arousal and that CS was greater between dyads
within each visitor group who were socially close than those who were not. The findings contribute
to the evidence that social relationships shape co-regulation through shared fluctuations in arousal
(Andersen et al., 2025). Based on these studies, we would expect that in interactive contexts evoking
average physiological arousal (i.e., non-frightening; not involving haunted houses or risky behaviour
such as firewalking), CS would be highest at a first encounter and decrease as dyads who began as
strangers become acquainted and socially closer.

Self-reported feelings of social closeness to another person tends to result in individuals being
comfortable in closer physical proximity to each other (Cristani et al., 2011; Hebel & Rentzsch,
2022; Lahnakoski et al., 2020). Like social closeness, physical proximity (henceforth interpersonal
distance) is bidirectionally related with shared arousal (Andersen et al., 2025; Jackson et al., 2018).
In a 2x2 design, Jackson et al. (2018) recorded groups of people walking freely vs. in synchrony as a
group vs. manipulated arousal levels by having the leader walk above (high arousal) or below (low
arousal) average walking speed. The researchers found that the pack was the tightest (least distance
between walkers) during the condition combining synchronous and high arousal walking. The
researchers propose that their findings reflect how arousal and synchrony can be employed to

manufacture group cohesion (Jackson et al., 2018). To our knowledge, no studies to date investigate
the relationship between shared arousal measured via CS and interpersonal distance. Nonetheless,
previous research using other measures of physiological synchrony has revealed that arousal levels
are likely associated with interpersonal distance. For example, teams of paramedic trainees who show
greater synchrony in electrodermal activity work physically closer to one another and engage in more
cooperative dialogue compared to teams showing lower synchrony (Joshi et al., 2025). Interpersonal
distance appears be associated with shared arousal levels during collaboration, yet additional
evidence is needed to establish this relationship's reliability.

Notably, the relationships between shared arousal, social closeness, interpersonal distance and
collaboration have (to our knowledge) only been examined in dyads consisting of two young adults
at a single point in time (as opposed to longitudinally). In very few cases, intergenerational patterns
of shared arousal have been explored within mother-child dyads (for a review, see: DePasquale,
2020). Older adults have yet to be represented in such studies, a trend that is observed in
experimental studies more broadly (Pauly et al., 2021; Van Marum, 2020). It is critical to remedy the
underrepresentation of older adults as the proportion of older adults in the world's population is
steadily climbing (United Nations, 2024) and older adults are at elevated risk of experiencing
loneliness (WHO, 2025). Moreover, longitudinal insights extending beyond first encounters and
covering stages before friendships and romantic relationships are necessary for a deeper
understanding of the physiological underpinnings of relationship development.

1.3   Present study

This research sought to shed light on the physiological mechanisms of collaboration in budding
relationships–both same generation and intergenerational, including older adults. We preregistered
(https://osf.io/xuvy5) the three following experimental aims:

• Document the trajectory of CS over the course of six sessions of collaborative art-
making within both same generation and intergenerational dyads.
• Explore how levels of collaboration visible in drawings (evaluated by external raters)
relate to dyad members' feelings of social closeness, physical closeness, and
performance on a separate collaborative task (i.e., completing a jigsaw puzzle).
• Explore how CS is related to levels of collaboration visible in drawings. Specifically,
we wish to assess whether this relationship differs between same generation and
intergenerational dyads, and the extent to which this relationship is impacted by
dyad's feelings of social closeness, physical closeness, and performance on a separate
collaborative measure (i.e., jointly completing a jigsaw puzzle).

In reporting these findings, charting CS within intergenerational and same generation dyads across 6
sessions in an everyday context similar to a community art program, we reveal first insights into
longitudinal patterns of shared arousal, in a unique design that involves older adults. The implications
of this work are central to understanding the behavioural and physiological mechanisms governing
relationship formation and collaboration across the lifespan. Moreover, this work sheds light on
important considerations regarding the implementation of CS as an objective measure of social
connection or collaboration in everyday, real-world settings.

2   Methods

2.1   Participants

Main sample. We recruited 122 community-dwelling participants from Zurich, Switzerland (31 older adults, aged 69+ years; 91 younger adults, aged 18–35 years). Included participants were fluent in German and had no known history of neurological or psychological disorders (e.g., stroke, concussion, ADHD, autism, schizophrenia, or depression). Participants were assigned to intergenerational dyads ($n = 31$) or same generation dyads ($n = 30$) based on availability for sessions (e.g., matching people available at the same day/time for 6 consecutive weeks). All dyads started the study as strangers. Intergenerational dyads comprised one older adult (76 ± 4 years); 18 female, 13 male) and one younger adult (24 ± 4 years; 19 female, 11 male, 1 other). Same generation dyads comprised two younger adults (22.5 ± 3.6 years; 36 female, 24 male). Supplementary Table 1 details the number of individual participants' recordings and dyads' sessions that were excluded, as well as reasons for exclusion.

Our recruitment campaign involved advertising through the University of Zurich's Healthy Longevity Centre, ETH Zurich's DeSciL student pool, clubs and organisations for senior citizens (e.g., theatre groups and choirs/orchestras), and social media. All participants provided written informed consent prior to participation. Ethical approval was obtained from the Ethics Committee of the Canton Zurich (Ref: 2023-01073). The study was conducted according to the Declaration of Helsinki. We report how we determined our sample size, all data exclusions, all manipulations, and all measures in the study.

External ratings of drawings. To obtain ratings of drawings produced by the main sample (described further in 2.3.1), we recruited 148 new participants via Prolific after the main study concluded. To be included, participants had to be over18 years old, proficient English speakers, residents of the United Kingdom. To ensure response quality, we only accepted participants who had previously completed a minimum of 50 tasks on Prolific with a 100% approval rate. Of these participants, 45 were excluded for failing to respond to all attention checks correctly (i.e., by moving all sliders all the way to the right of the sliding scale when prompted). We thus use ratings from 103 raters in our analyses. We did not collect demographic information from these participants. All participants provided written informed consent. Ethical approval was obtained from the Ethics Committee of ETH Zurich (Ref: 2024-N-265). The study was conducted according to the Declaration of Helsinki.

2.2   Procedure

Each dyad completed six weekly sessions (once per week, for six weeks). At each session, participants completed questionnaires querying feelings of loneliness (De Jong Gierveld & van Tilburg, 2006) and attitudes toward their own and the other (older or younger) generation (Pittinsky et al., 2011). Participants also completed a questionnaire measuring empathy (Paulus, 2009) at the first session. Dyads drew alone once and collaboratively twice with oil pastels provided by Caran D'Ache, for a duration of five minutes (Figure 1; Supplementary Figure 1). Participants were not given any prompts regarding what to draw or how to draw (e.g., no suggestions for motifs). Participants were instructed not to talk while drawing, so that speech would not be a confound in comparisons of drawing alone (of particular relevance for neural and cardiac measures). Participants were instructed that they could talk freely during the rest of the session. After drawing, participants completed a collaborative activity, with different activities each week. In the third session, the collaborative activity was to complete a 54-piece jigsaw puzzle. Activities completed on other weeks include sorting pastels, a divergent thinking task, and a joint verbal fluency task. While participants drew and completed collaborative activities, we recorded their cortical activity using functional near-infrared spectroscopy (fNIRS) over the bilateral inferior frontal gyri (IFG) and bilateral temporo-

parietal junction (TPJ). Furthermore, we video-recorded participants' movement using a GoPro
camera to capture movements of the head and hands. At the end of each session, after we stopped the
fNIRS and video recordings, participants completed the Inclusion of Other in Self scale (Aron et al.,
1992). Analyses of fNIRS recordings, loneliness scores, and attitudes toward generations are reported
elsewhere, as are additional specifics regarding the fNIRS recording methodology (Moffat et al.,
2025).

Figure 1. A) Left: An intergenerational dyad drawing independently on separate pieces of paper during fNIRS recording.
The dyad members are separated by a grey felt divider that obscures their view of each other. Right: The same
intergenerational dyad drawing together on a single piece of paper during fNIRS recording. The colourful lines overlaid
on top of the participants show the 2D motion tracking used to capture interpersonal distance, specifically the distance
between dyad members' neck joints. B) IBI times series of two different dyads, illustrating high and low levels CS with
the produced drawing (see further examples sorted by collaboration ratings in Figure 2).

2.3 Measures

2.3.1 Drawing collaboration

Each drawing that the dyads created was judged by independent raters recruited via Prolific. Raters
($n = 103$) were instructed that they would view 122 drawings, each created by two people at once on
a single piece of paper. They were asked to rate the degree of harmony between the elements of the
drawings (on a sliding scale from 0 "entirely independent" to 100 "entirely coordinated") and the use
of space (on a sliding scale from 0 "divided" to 100 "unified"). To obtain a measure of the perceived
coherence of the drawing, which we propose reflects the level of collaboration during drawing, we

summed the ratings of harmony and use of space. Thus, raw scores of drawing collaboration could
range from 0-200. Examples of drawings with highest, average, and lowest collaboration ratings are
shown in Figure 2.

Figure 2. Examples of drawings organised by rating of collaboration. Top row = highest collaboration ratings; middle
row = average collaboration ratings; bottom row = lowest collaboration ratings. Ratings are derived by summing raters'
ratings of the harmony between the elements of the drawings (0 "entirely independent" to 100 "entirely coordinated") and
use of space (0 "divided" to 100 "unified"). The left-most drawings in the top and bottom row are also presented in
Figure 1B alongside the IBI time series recorded while the drawings were made.

2.3.2 Puzzle collaboration

After completing the drawing component of the third session, dyads completed a puzzle as the
collaborative activity for that session. Speaking was permitted. Dyads had six minutes to complete a
54-piece jigsaw puzzle of cartoon animals in jungle (pieces' dimensions: ~ 8 x 8 cm). Dyads' scores
corresponded to the number of pieces assembled in 6 minutes. In the case that participants assembled
all pieces in less than 6 minutes, the remaining number of seconds to 6 minutes was divided by 6.66
(the average number of seconds per piece if completed in 6 minutes; 360 s /54 piece = 6.66 s) and
added to a score of 54.

2.3.3 Social Closeness

We measured feelings of social closeness using the Inclusion of Other in Self (IOS) scale (Aron et al., 1992). Participants viewed seven sets of overlapping circles representing the self and the partner, ranging from no overlap to near complete overlap of the two circles, and indicate which set of circles best represents their relationship with their partner (i.e., the other dyad member). To capture cumulative social closeness within dyads and differences in feelings of social closeness within dyads, we computed summed and absolute difference IOS scores per dyad (Boukarras et al., 2025; Kenny et al., 2006). We refer to these measures as social closeness (Sum) and social closeness (Dif).

2.3.4 Interpersonal distance

We used the videos recorded at each session to track upper body movements while dyads drew together. Specifically, we used OpenPose (Cao et al., 2021) to estimate 2D pose per frame, returning coordinates for predifined body parts. To assess interpersonal distance (sometimes referred to as proximity in other work), we computed the average distance between the coordinates of each dyad members' neck joint for each 5-minute instance of drawing together. In our analyses, this measure is referred to as distance.

2.4 Data analysis

2.4.1 Extracting inter-beat intervals

As per our preregistration, we extracted the heart rate signal by identifying peaks in HbO concentrations (Scholkmann et al., 2012). We cleaned the HbO signal from neural and further physiological components such as Mayer waves by upsampling it to 100 Hz and applying a zero-phase third-order Butterworth IIR band-pass filter (Perdue et al., 2014) with cut-offs at 0.5 and 2.5 Hz (Sun et al., 2018). Then, we smoothed the signals using a Savitzky-Golay filter with window length 15 and a 5^{th}-order polynomial. We identified peaks with the AMPD algorithm (Scholkmann et al., 2012), implemented in the python *ampdlib* package (Cerina, 2023), and transformed the signal peak timestamps to an inter-beat interval (IBI) time series, smoothly interpolated at 4 Hz. We originally preregistered ARIMA cleaning (Feldman et al., 2011), but in practice it reduced the signal to near-zero fluctuations, largely removing the slow structure that may reflect the physiology of interest in this context. Given the lower signal-to-noise ratio of fNIRS-derived IBIs when compared to ECG (electrocardiography) signals, and our total number of samples (fNIRS signals recorded at ~ 5 Hz), we instead analysed IBI series on their original scale to retain physiologically meaningful variance.

2.4.2 Calculating synchrony

We calculated cross-correlation functions (CCFs) between all dyads' cleaned physiological signals in each session and activity across time lags of +/- 3 seconds (-12 to +12 lags at 4 Hz) and retained non-absolute CCF values. Examples of time series yielding high and low CS estimates are visualised in Figure 1B.

2.4.3 Group-level analysis

We completed our preregistered exploratory statistical analyses using a Bayesian approach to multilevel regression (McElreath, 2020). We fit models using the brms package (version 22.2; Bürkner, 2017) in the R language (version 4.5.1; R Core Team, 2022) within the RStudio IDE (version 2025.05.1+513; RStudio Team, 2020). We report and interpret the posterior distribution of relevant model parameters using a 95% credible interval, calculated via the 95% highest posterior

density region (HPD) method (McElreath, 2020). For readers familiar with Frequentist approaches
that include p-values, we recommend perusing Kruschke & Liddell (2018), and we share a
(simplified) heuristic for interpreting HPDs: Parameter and comparison estimates can be considered
substantial when the HPD does not contain zero and can be considered trends when the tip of an
HPD-tail overlaps with zero.

Measures of collaboration, social closeness, and interpersonal distance. To assess the extent to
which collaboration, social closeness, and interpersonal distance changed across the six sessions, as
well as relationships between measures, we fit a series of models in the following structure:
*MeasureA ~ 1 + MeasureB * Group * Session + (1|DyadID)*. We extracted point estimates for all
sessions combined and slopes across sessions, and subsequently computed contrasts between groups
(intergenerational vs. same generation).

CS for all sessions combined and across sessions. We first sought to establish if CS levels differed
substantially between real and pseudo dyads. We fit the following model: *CS ~ DyadType * Group ***
*Task * Session*. We did not include random intercepts at this stage, as the total number of real and
pseudo dyads exceeded 6000. Random intercepts, accounting for dyad-level variance, were included
in subsequent group level analyses. Contrasts between real and pseudo dyads were computed to
confirm that some component of the CS was attributable to real social interaction, as opposed to
similar experiences or homeostatic biological rhythms.

Relationships between CS and measures of collaboration, social closeness, and interpersonal
distance. First, we fit a model to the CS data, which we averaged across all lags. The model was: *CS*
*~ Group * Drawing Collaboration + Group * Puzzle Collaboration + Group * Social Closeness*
*(Sum)+ Group * Social Closeness (Dif) + Group * Interpersonal Distance + Session + (1|DyadID)*.
Subsequently, we fit a model to examine relationships between CS at separate lags and our measures.
Models were structured as follows and fit per measure: *CS ~ Lag * Group * Measure + (1|DyadID)*.
For both steps, we extracted point estimates for all sessions combined and slopes across sessions, and
subsequently computed contrasts between groups (intergenerational vs. same generation).

3   Results

3.1   Behavioural and self-report measures

3.1.1 Contrasts between groups

Drawing collaboration. Drawing collaboration differed substantially between groups, with the
intergenerational drawings showing greater collaboration than same generation drawings. Drawing
collaboration did not change substantially across sessions for either group, though there was a trend
of intergenerational dyads showing a greater increase in collaboration than same generation dyads.
Puzzle collaboration. Puzzle collaboration differed markedly between groups, with same generation
dyads scoring substantially higher than intergenerational dyads. **Social closeness (Sum).** Summed
social closeness was substantially greater for intergenerational dyads than same generation dyads.
Summed social closeness increased substantially across sessions for both groups at a similar rate.
Social closeness (Dif). Intergenerational dyads showed substantially greater differences in dyadic
social closeness than same generation dyads. Dyadic differences in social closeness decreased across
sessions substantially for same generation dyads and showed the same trend for intergenerational
dyads. **Interpersonal distance.** Interpersonal distance showed a trending group difference, with
same generation dyads showing substantially lower interpersonal distance than intergenerational
dyads. Interpersonal distance increased substantially across sessions for same generation dyads only.

Comparisons between groups showed a trending group difference, where the slope of change in interpersonal distance across sessions was more positive for same generation than intergenerational dyads.

Table 1. Standardised estimates computed per behavioural and self-report measure with HPD in square brackets. Point estimates for all six sessions combined are presented first, followed by the estimates of the slope of change across sessions in the lower section. No slope of change is available for puzzle collaboration, as dyads only completed the puzzle task once, at the third session.

	Drawing Collaboration	Puzzle Collaboration	Social Closeness (Sum)	Social Closeness (Dif)	Interpersonal Distance
Sessions combined					
Intergenerational	0.20 [0.08, 0.32]	-0.50 [-0.59, -0.42]	0.17 [0.7, 0.26]	0.13 [0.03, 0.23]	0.07 [-0.18, 0.03]
Same gen	-0.03 [-0.15, 0.08]	0.45 [0.36, 0.53]	-0.18 [-0.27, -0.08]	-0.08 [-0.18, 0.03]	-0.06 [-0.04, 0.17]
Intergen-Same gen	0.23 [0.07, 0.40]	-0.95 [-1.07, -0.84]	0.35 [0.22, 0.47]	0.20 [0.07, 0.34]	-0.13 [-0.28, 0.00]
Change across six sessions					
Intergenerational	0.04 [-0.01, 0.93]	-	0.22 [0.17, 0.27]	-0.05 [-0.10, 0.00]	0.02 [-0.02, 0.07]
Same gen	-0.01 [-0.07, 0.04]	-	0.16 [0.11, 0.21]	-0.07 [-0.13, -0.02]	0.07, [0.01, 0.12]
Intergen-Same gen	0.05 [0.00, 0.11]	-	0.09 [-0.02, 0.20]	0.02 [-0.03, 0.07]	-0.04 [-0.10, 0.00]

3.1.2 Relationships between measures

Relationships between measures of collaboration, self-reported social closeness, and interpersonal distance per group are illustrated in Figure 3. Drawing collaboration was substantially positively associated with puzzle collaboration for intergenerational group only, yielding a substantial difference between intergenerational and same generation dyads (ß = 0.26, [0.08, 0.42]). Drawing collaboration was substantially positively associated with summed social closeness for both groups, but the association was stronger for same generation dyads, relative to intergenerational dyads (ß = -0.16, HPD = [-0.30, -0.02]). Drawing collaboration was not substantially associated with differences in dyadic social closeness. Drawing collaboration was substantially negatively associated with interpersonal distance for intergenerational dyads and substantially positively associated with interpersonal distance for same generation dyads, yielding a substantial group difference (ß = -0.21, HPD = [-0.25, -0.07]). Puzzle collaboration was substantially negatively associated with summed social closeness for same generation dyads only, resulting in a substantial group difference (ß = 0.18, HPD = [0.06, 0.30]). Puzzle collaboration was also substantially negatively associated with differences in dyadic social closeness for same generation dyads only, resulting in a substantial group difference (ß = 0.21, HPD = [0.09, 0.33]). Puzzle collaboration was substantially negatively associated with interpersonal distance for same generation dyads only, and no substantial group difference was observed. Summed social closeness was substantially positively associated with dyadic differences in social closeness for both groups and did not differ substantially between groups. Interpersonal distance was substantially positively associated with summed social closeness for intergenerational dyads and substantially negatively associated with summed social closeness for same generation dyads, with a substantial difference between groups (ß = 0.22, HPD = [0.08, 0.36]). Interpersonal distance showed a trending negative association with differences in dyadic social

closeness for intergenerational dyads and a substantial positive association for same generation
dyads, yielding a substantial difference between groups (ß = -0.20, HPD = [-0.35, 0.06]).

Figure 3. Standardised estimates of relationship between behavioural and self-report measure per group. Black squares
indicate that relationships are substantial (i.e., 95% HPD does not overlap with zero). Note: the relationship between
Interpersonal Distance and Social Closeness (Dif) for same generation dyads is trending, rather than substantial, but in a
black square to facilitate interpretation of results. Standardised parameter estimates per measure shown with 95% HPD in
Supplementary Table 2.

3.2   Heart rate synchrony

3.2.1 Comparing real and pseudo dyads

All sessions combined. For all lags averaged and separate lags, we observed no substantial
difference in CS levels between real and pseudo dyads for drawing together or alone (Supplementary
Figure 2). **Across sessions.** For all lags averaged and separate lags, the rate of change in CS across
sessions (all lags averaged) did not differ substantially between real and pseudo dyads for drawing
together or alone (Figure 4; Supplementary Figure 3).

We interpret these findings as follows: Any two individuals meeting the inclusion criteria, who
experience the specific dyadic experimental environment applied here would show comparable levels
of CS. True interaction with an assigned partner does not appear to contribute uniquely to CS levels.
We present the further analyses in accordance with our preregistration.

3.2.2 Comparisons between drawing conditions and groups

All sessions combined. For all lags averaged, we observed no substantial effects of drawing task,
group or task*group interactions on CS levels. **Across sessions.** For all lags averaged, we observed
no substantial main effect (i.e., no substantial change in CS levels across sessions) and no substantial
effects of drawing task, group or task*group interactions on the slope of CS across sessions.

Figure 4. Trajectories of CS in real and pseudo dyads across sessions. Top panel shows individual data points (dots) and estimated slopes (lines). Bottom panel zooms in on the estimated slopes, with grey shading showing 95%, 89%, and 50% intervals of the posterior predictive distributions. Neither the overall level of CS for all sessions combined nor the rate of change in CS across sessions differed substantially between real and pseudo dyads. Though real dyads' CS the trajectories across sessions may appear to differ, the 95 % HPD overlaps > 10 % (threshold for a trend, Section 2.4.3) for drawing alone and together.

3.3 Relationships between CS, behaviour, and self-report measures

As per our preregistration, we first explored relationships between CS with all lags averaged, collaborative behaviour, social closeness, and distance. We found no substantial relationships between CS and either measure of collaboration, social closeness or interpersonal distance. Also in line with our preregistration, we explored relationships between CS and each measure per lag (n = 25; from -3000 ms to 3000 ms in 250 ms increments). We observed no substantial relationships for any measure and no substantial difference between groups at any lag (Figure 5). For drawing collaboration, intergenerational dyads showed a trending positive relationship between CS and drawing collaboration at all lags except -3000 ms, -2750 ms, -2500 ms, and -1750 ms. At these four lags, the degree of 95 % HPD overlap with 0 ranged from 10-14 %, (where no overlap indicates a slope substantially different than 0, < 10 % indicate a trending difference from 0). For intergenerational dyads, positive lags reflect the older dyad member's IBI signal leading or coming earlier in time, and negative lags reflect the younger member's IBI signal leading or coming earlier in time. As we observed trending positive and negative lags trending, there is no evidence that a dyad member of a certain generations fills a leader role.

Figure 5. Standardised estimates of relationship between CS and session number, as well as behavioural measures of collaboration, self-reported measures of social closeness, and interpersonal distance per lag (n = 25; from -3000 ms to 3000 ms in 250 ms increments). No substantial relationships observed. Drawing collaboration panel shows trending relationships between CS and drawing collaboration scores for intergenerational dyads at all but 4 lags. See Supplementary Figure 4 for visualisation including error bars showing 95 % HDP.

4   Discussion

In this study, we charted cardiac synchrony (CS), collaborative behaviours and social closeness in intergenerational and same generation dyads across 6 sessions in an everyday context resembling a community art program. Dyads' collaborative behaviours and measures of social closeness showed nuanced relationships, primarily characterised by greater collaboration in cases of greater social closeness. For CS–both averaged and separate lags–we found no substantial differences between real and pseudo dyads overall or across sessions, no substantial task or group differences among real dyads overall or across sessions, and no substantial relationships between CS and any measure of behaviour. We found trends of substantial relationships for intergenerational dyads between CS and drawing collaboration at all but four lags, with the four non-trending lags showing the same numerical relationship. We propose that these findings, considered together, suggest that cardiac synchrony is to a limited extent sensitive to higher stakes, open-ended, collaboration. We discuss the implications of these findings below.

4.1   Relationships between behavioural and self-report measures

Intergenerational and same generation dyads exhibited some similarities in behavioural measures alongside marked differences. For both groups, greater cumulative feelings of social closeness within dyads co-occurred with greater ratings of collaboration of the dyads' drawings. This finding is consistent with previous findings (Ford & Aberdein, 2015; Shafaei et al., 2020; Wolf et al., 2016). Notably, however, same generation dyads who reported more similar feelings of social closeness (as opposed to intergenerational dyads) completed the jigsaw puzzle more quickly. This pattern was not present in intergenerational dyads. We propose that drawing and puzzle assembly, both collaborative tasks, differ in the clarity of the goal. The goal of puzzle assembly is clear to both dyad members (i.e., correctly assemble all pieces before time runs out) while the goal of drawing may be more open-ended, based on the amount and specificity of dyads' verbal planning prior to drawing and the process of co-drawing without talking. For reasons relating to experimental control, most studies to

date have focused collaborative tasks with clear end goals (Danyluck & Page-Gould, 2019; Fusaroli et al., 2016; Mitkidis et al., 2015; Strang et al., 2014). The value of the large dataset and analyses that we present here lies in their ecological validity. That is, they reflect everyday collaboration arguably more accurately than some laboratory-based measures reported by previous studies. Everyday collaboration inherently entails free choice regarding the extent to which one engages and mutual adaptation to one's partner, which have consequences for the development of social relationships.

We did not expect groups to differ in how close they sat and/or leaned toward one another while drawing together. Yet, intergenerational dyads drew sitting closer together than same generation dyads. Further, the relationships between interpersonal distance and feelings of social closeness, measured as a cumulative score per dyad and the discrepancy between dyad members, was opposing between groups. Intergenerational dyads sat closer when they felt more socially close or had smaller discrepancies in their feelings of social closeness–showing the pattern we anticipated based on existing findings (Cristani et al., 2011; Hebel & Rentzsch, 2022; Lahnakoski et al., 2020). Same generation dyads, on the other hand, sat closer when they felt less socially close or had greater discrepancies in their feelings of social closeness. These opposite patterns suggest that reduced levels of social comfort during collaboration may amplify dyads' perceptions of the stakes of the collaboration and result in compensatory behaviours, such as sitting closer to one another. Previous works corroborates this interpretation (Lieberz et al., 2021; Saporta et al., 2021; Scheele et al., 2012; Shamay-Tsoory & Kanterman, 2024). Thus, interpersonal distance in collaboration may be a promising objective measure of social closeness and relationship development within recently acquainted dyads.

4.2   CS not sensitive to true interaction, unless stakes are increased

Our analyses of CS revealed no substantial differences between real and pseudo dyads overall or across sessions, no substantial task or group differences among real dyads overall or across sessions, and no substantial relationships between CS and any measure of behaviour. This was true for averaged and separate lags. The absence of differences between real and pseudo dyads can be interpreted as meaning that any two individuals meeting the inclusion criteria of our study (whether same generation or intergenerational), who drew following the same prompts given by the researchers, in the same context are likely to exhibit comparable levels of CS, whether or not they are in the same room at the same time or acquainted. In other words, the levels of CS observed in this study do not index *true* interaction between two people. Moreover, the levels of CS observed in this study do not index changes in collaborative behaviour or feelings of social closeness across repeated encounters.

A small number of studies have previously reported that CS does not differ between real and pseudo dyads (Flory et al., 2023; Strang et al., 2014). Flory et al. (2023) present alternative explanations. One is that CS is a spurious finding (i.e., could occur for any two people in the world). We do not think this interpretation applies here as the experimental setting, structure, and inclusion criteria provide substantial structure to the experience of each participant–that is, the description of CS as completely spurious is not founded. Another of Flory et al.'s alternative explanations is that participants operate in a common psychophysiological mode (i.e., similar mental state evoked by the experimental design), aligning with Danyluck and Page-Gould's (2019, p. 1) view that "physiological synchrony could simply reflect the mutually experienced demands of a shared environment". While this is in line with our rationale against spurious CS, we disagree on the basis that we observed trending relationships between CS and drawing collaboration for intergenerational

dyads for 21 of 25 lags. We propose that these relationships point to a noteworthy, albeit non-
substantial, influence of high stakes collaboration on CS.

We observed no task or group differences in CS measured within real dyads. Prior research has also
reported the absence of task-specific or performance-related changes in CS (Behrens et al., 2020;
Boukarras et al., 2025; Fusaroli et al., 2016; Gordon et al., 2020). More specifically, Gordon et al.
(2020) detected evidence for task differences in CS when participants drummed together at a
predetermined speed or drummed freely, improvising together. Similarly, other authors report having
found no evidence to support associations between CS and task performance. Boukarras et al. (2025)
report that reaction times in a dyadic grasping task are not associated with CS. Behrens et al. (2020)
report that cooperative success is not associated with CS when dyads played the Prisoners Dilemma
game. Strang et al. report that cooperative success is not associated with CS when dyads play Tetris
with one person controlling object rotation and the other controlling the positioning (Strang et al.,
2014). Finally, with respect to measures of social connection and closeness, HR coordination
(consisting of alternate measures to CS) during Lego building was reported not to be associated with
measures of perceived group relatedness or competence, but rather with speech and movement
dynamics (Fusaroli et al., 2016; see review for more examples: Mayo & Gordon, 2020). In summary,
despite the large number of studies dedicated to understanding CS on the grounds that CS can offer a
window into collaboration and social connection, the extent to which this is true remains a topic
worthy of debate (as well as replication).

From a practical perspective, the consistency of non-differences in CS across our preregistered
exploratory analyses suggest that CS is not an optimal measure for predicting collaborative outcomes
or relationship development in real-world environments. Given the mounting interest expressed in
using wearable devices to collect objective insights relating to social wellbeing (Fendrich et al., 2021;
Jiang et al., 2019; Moffat et al., 2024; UNICEF, 2015), we offer the following recommendation:
Organisations, policy makers, and researchers aspiring to use wearable physiological monitoring to
track changes in social behaviours and social wellbeing (i.e., social connection) should consider the
low explanatory power of CS demonstrated here and pilot extensively before dedicating funding to
large-scale roll-outs of these devices to measure program engagement or success. We feel justified in
making this recommendation based on the sample size (366 samples involving within-
participant/dyad design, which reduces inter-individual variability in our analyses). Moreover, other
work by our research group shows that inter-brain synchrony may index changes in social wellbeing
with greater sensitivity (Moffat et al., 2025), though we acknowledge the logistical and financial
barriers to larger-scale introduction of brain imaging measures.

Nonetheless, from an empirical perspective, it is promising that collaborative drawing scores show a
trending relationship with CS levels within real intergenerational dyads. This suggests that further
research could uncover which social, cognitive, and behavioural conditions need to be met for CS to
be sensitive to true dyadic interaction. Based on the behavioural data reported here, we believe these
conditions should include higher perceived stakes for successful collaboration. Our behavioural data
revealed that intergenerational dyads' drawings were rated more collaborative than same generation
dyads', and that the elevated levels of drawing collaboration emerged from the group (i.e.,
intergenerational dyads) that reported lower feelings of social closeness within dyads, more disparate
feelings of social closeness within dyads, and sat closer together while drawing together (Table 1). In
concert, these behaviours highlight how lower levels of social comfort during collaboration may raise
the stakes of the collaboration and may, in turn, contribute to shared fluctuations in arousal during
collaboration.

4.3 Limitations

In using the term "higher stakes collaboration", we are attempting to articulate the origin of the decreased social comfort, or increased social tension, as it relates to reputation management (Barclay, 2012; Cavazza et al., 2015) or the additional cognitive load stemming from interpersonal uncertainty (i.e., lack of common ground: Clark & Marshall, 1981; Speer et al., 2024). We are open to this being conceptualised in other ways. Follow up experiments could probe the extent to which individual strategies and subjective experiences shed light on the nature of such collaboration.

Measures of dyadic synchrony can influence observed synchrony levels (Helm et al., 2018). We considered this carefully before preregistering our planned pipeline to calculate IBI time series. We compared an HR synchrony approach implemented previously with fNIRS (Jones, 2024) with more common IBI approaches (Adel et al., 2025; Boukarras et al., 2025). We opted for IBI to align our work with the majority of existing studies examining CS. With respect to our comparisons of real and pseudo dyads' levels of CS, a complementary approach would be to employ segment-shuffled surrogates within each dyad. In this method, each person's cardiac time series is cut into short segments that are then randomly reordered so that the overall signal properties are preserved but true timing is disrupted (e.g., Coutinho et al., 2021). This would reveal whether true dyads display CS on top of task-driven similarity. Duration of heart rate recordings has been demonstrated to influence the likelihood of finding significant CS with dyads, particularly for smaller group sizes such as dyads (Stuldreher et al., 2023). From this perspective, our design is optimised to identify true CS, should it be present.

5 Conclusion

We charted CS, collaborative behaviour, social closeness and interpersonal distance within intergenerational and same generation dyads across six sessions involving a creative art-making task. Our analyses of collaborative behaviour, self-reported social closeness, and interpersonal distance, when considered together, suggest that social comfort drives relationships between social closeness, interpersonal distance and collaboration. Of these, interpersonal distance emerged as a promising objective measure of relationship development. We found that CS did not co-vary substantially with group, task, an interaction thereof or collaborative behaviour, social closeness or interpersonal distance. Nonetheless, we did find a trending relationship between collaboration while drawing together and CS that was specific to intergenerational dyads. These findings have implications for understanding the mechanisms underpinning social interaction within and between generations and for the implementation of CS as an objective measure of social connection or collaboration in everyday, real-world contexts.

Author contributions

RM: Conceptualisation, Equipment donation, Data curation, Formal analysis, Investigation, Visualisation, Writing – original draft, Writing – review & editing. LN: Data curation, Validation, Writing – original draft. ESC: Conceptualisation, Funding acquisition, Project administration, Writing – review & editing.

Data availability statement

These data were collected after the submission of the following preregistration: https://osf.io/xuvy5. The data collected and analysed for this study can be found in our repository entitled 'Cardiac

synchrony during collaborative drawing: Intergenerational and longitudinal perspectives' on OSF: https://osf.io/akmqw.

Acknowledgements

We thank the survey center operated by UZH Healthy Longevity Center, University of Zurich, for their assistance with recruitment of older adults from the Zurich community. We thank Tessa Portier and Medea Häuselmann for their assistance with data collection, Linda Fanconi for scheduling participants' sessions, Alistair Gadola for coding participant behaviour, and Fanny Mougel for digitising the drawings. We thank Chantal Nagel for assisting with literature review. We thank Courtney Casale and Tessa Portier for their help with proofreading.

Conflict of interest

The authors declare no potential conflict of interest.

Funding

Two Cortivision Photon Cap devices were provided to RM through the Cortivision Pathfinder Program (grant number CPP-2023/09/01). Oil pastels were provided to RM by Caran D'Ache (agreement number 712517). RM, LN, and ESC were supported by the Social Brain Sciences Lab at ETH Zurich.

References

Adel, L., Manova, V., Bain, T., Ambwani, D., Khoury, B., & Lifshitz, M. (2025). *Interpersonal cardiac synchrony during emotional sharing between strangers*. PsyArXiv.

Andersen, M. M., Rosas, F. E., Schjoedt, U., Mediano, P. A. M., Price, H., Konvalinka, I., & Clasen, M. (2025). Scared together: Heart rate synchrony and social closeness in a high-intensity horror setting. *Emotion*. https://doi.org/10.1037/emo0001556

Anderson, S., Fast, J., Keating, N., Eales, J., Chivers, S., & Barnet, D. (2017). Translating knowledge: Promoting health through intergenerational community arts programming. *Health Promotion Practice*, *18*(1), 15–25. https://doi.org/10.1177/1524839915625037

Aron, A., Aron, E. N., & Smollan, D. (1992). Inclusion of other in the self scale and the structure of interpersonal closeness. *Journal of Personality and Social Psychology*, *63*(4), 17. https://doi.org/10.1037/0022-3514.63.4.596

Barclay, P. (2012). Harnessing the power of reputation: Strengths and limits for promoting cooperative behaviors. *Evolutionary Psychology*, *10*(5), 868–883. https://doi.org/10.1177/147470491201000509

Behrens, F., Snijdewint, J. A., Moulder, R. G., Prochazkova, E., Sjak-Shie, E. E., Boker, S. M., & Kret, M. E. (2020). Physiological synchrony is associated with cooperative success in real-life interactions. *Scientific Reports*, *10*(1), 19609. https://doi.org/10.1038/s41598-020-76539-8

Bizzego, A., Azhari, A., Campostrini, N., Truzzi, A., Ng, L. Y., Gabrieli, G., Bornstein, M. H., Setoh, P., & Esposito, G. (2019). Strangers, friends, and lovers show different physiological synchrony in different emotional states. *Behavioral Sciences*, *10*(1), 11. https://doi.org/10.3390/bs10010011

Boukarras, S., Placidi, V., Rossano, F., Era, V., Aglioti, S. M., & Candidi, M. (2025). Interpersonal physiological synchrony during dyadic joint action is increased by task novelty and reduced by social anxiety. *Psychophysiology*, *62*(3), e70031. https://doi.org/10.1111/psyp.70031

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, *80*(1), 1–28. https://doi.org/10.18637/jss.v080.i01

Butler, E. A. (2011). Three views of emotion regulation and health. *Social and Personality Psychology Compass*, *5*(8), 563–577. https://doi.org/10.1111/j.1751-9004.2011.00372.x

Cao, Z., Hidalgo, G., Simon, T., Wei, S.-E., & Sheikh, Y. (2021). Openpose: Realtime multi-person 2d pose estimation using part affinity fields. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, *43*(1), 172–186. https://doi.org/10.1109/TPAMI.2019.2929257

Cavazza, N., Guidetti, M., & Pagliaro, S. (2015). Who cares for reputation? Individual differences and concern for reputation. *Current Psychology*, *34*(1), 164–176. https://doi.org/10.1007/s12144-014-9249-y

Cerina, L. (2023). *Automatic Multiscale-based Peak Detection (AMPD)* [Computer software].

Clark, H. H., & Marshall, C. R. (1981). Definite reference and mutual knowledge. In A. K. Joshi, B. L. Webber, & I. A. Sag (Eds), *Elements of Discourse Understanding* (pp. 10–63). Cambridge University Press.

Coutinho, J., Pereira, A., Oliveira-Silva, P., Meier, D., Lourenço, V., & Tschacher, W. (2021). When our hearts beat together: Cardiac synchrony as an entry point to understand dyadic co-regulation in couples. *Psychophysiology*, *58*(3), e13739. https://doi.org/10.1111/psyp.13739

Cristani, M., Paggetti, G., Vinciarelli, A., Bazzani, L., Menegaz, G., & Murino, V. (2011). Towards computational proxemics: Inferring social relations from interpersonal distances. *2011 IEEE Third International Conference on Privacy, Security, Risk and Trust and 2011 IEEE Third International Conference on Social Computing*, 290–297. https://doi.org/10.1109/PASSAT/SocialCom.2011.32

Danyluck, C., & Page-Gould, E. (2019). Social and physiological context can affect the meaning of physiological synchrony. *Scientific Reports*, *9*(1), 8222. https://doi.org/10.1038/s41598-019-44667-5

De Jong Gierveld, J., & van Tilburg, T. (2006). A 6-item scale for overall, emotional, and social loneliness: Confirmatory tests on survey data. *Research on Aging*, *28*(5), 582–598. https://doi.org/10.1177/0164027506289723

DePasquale, C. E. (2020). A systematic review of caregiver–child physiological synchrony across systems: Associations with behavior and child functioning. *Development and Psychopathology*, *32*(5), 1754–1777. https://doi.org/10.1017/S0954579420001236

Dimitroff, S. J., Kardan, O., Necka, E. A., Decety, J., Berman, M. G., & Norman, G. J. (2017). Physiological dynamics of stress contagion. *Scientific Reports*, *7*(1), 6168. https://doi.org/10.1038/s41598-017-05811-1

Feldman, R., Magori-Cohen, R., Galili, G., Singer, M., & Louzoun, Y. (2011). Mother and infant coordinate heart rhythms through episodes of interaction synchrony. *Infant Behavior and Development*, *34*(4), 569–577. https://doi.org/10.1016/j.infbeh.2011.06.008

Fendrich, S. J., Balachandran, M., & Patel, M. S. (2021). Association between behavioral phenotypes and sustained use of smartphones and wearable devices to remotely monitor physical activity. *Scientific Reports*, *11*(1), 21501. https://doi.org/10.1038/s41598-021-01021-y

Ferrer, E., & Helm, J. L. (2013). Dynamical systems modeling of physiological coregulation in dyadic interactions. *International Journal of Psychophysiology*, *88*(3), 296–308. https://doi.org/10.1016/j.ijpsycho.2012.10.013

Flory, S., Guglielmini, S., Scholkmann, F., Marcar, V. L., & Wolf, M. (2023). How our hearts beat together: A study on physiological synchronization based on a self-paced joint motor task. *Scientific Reports*, *13*(1), 11987. https://doi.org/10.1038/s41598-023-39083-9

Ford, R. M., & Aberdein, B. (2015). Exploring social influences on the joint Simon task: Empathy and friendship. *Frontiers in Psychology*, *6*. https://doi.org/10.3389/fpsyg.2015.00962

Fusaroli, R., Bjørndahl, J. S., Roepstorff, A., & Tylén, K. (2016). A heart for interaction: Shared physiological dynamics and behavioral coordination in a collective, creative construction task. *Journal of Experimental Psychology: Human Perception and Performance*, *42*(9), 1297–1310. https://doi.org/10.1037/xhp0000207

Gernert, C. C., Falkai, P., & Falter-Wagner, C. M. (2025). *Heart rate synchrony as an objective biomarker for therapy outcome in affective disorders* (p. 2024.08.29.24312787). medRxiv. https://doi.org/10.1101/2024.08.29.24312787

Golland, Y., Arzouan, Y., & Levit-Binnun, N. (2015). The mere co-presence: Synchronization of autonomic signals and emotional responses across co-present individuals not engaged in direct interaction. *PLOS ONE*, *10*(5), e0125804. https://doi.org/10.1371/journal.pone.0125804

Gordon, I., Gilboa, A., Cohen, S., Milstein, N., Haimovich, N., Pinhasi, S., & Siegman, S. (2020). Physiological and behavioral synchrony predict group cohesion and performance. *Scientific Reports*, *10*(1), 8484. https://doi.org/10.1038/s41598-020-65670-1

Hebel, V., & Rentzsch, K. (2022). One, two, three, sit next to me: Personality and physical distance. *Personality and Individual Differences*, *198*, 111798. https://doi.org/10.1016/j.paid.2022.111798

Helm, J. L., Miller, J. G., Kahle, S., Troxel, N. R., & Hastings, P. D. (2018). On measuring and modeling physiological synchrony in dyads. *Multivariate Behavioral Research*, *53*(4), 521–543. https://doi.org/10.1080/00273171.2018.1459292

Jackson, J. C., Jong, J., Bilkey, D., Whitehouse, H., Zollmann, S., McNaughton, C., & Halberstadt, J. (2018). Synchrony and physiological arousal increase cohesion and cooperation in large naturalistic groups. *Scientific Reports*, *8*(1), 127. https://doi.org/10.1038/s41598-017-18023-4

Jenkins, L. K., Farrer, R., & Aujla, I. J. (2021). Understanding the impact of an intergenerational arts and health project: A study into the psychological well-being of participants, carers and artists. *Public Health*, *194*, 121–126. https://doi.org/10.1016/j.puhe.2021.02.029

Jiang, L., Gao, B., Gu, J., Chen, Y., Gao, Z., Ma, X., Kendrick, K. M., & Woo, W. L. (2019). Wearable long-term social sensing for mental wellbeing. *IEEE Sensors Journal*, *19*(19), 8532–8542. https://doi.org/10.1109/JSEN.2018.2877427

Jones, I. (2024). *Neural, physiological and cognitive mechanisms of being watched* [Doctoral thesis]. University College London.

Joshi, V., Akiri, S., Taherzadeh, S., Williams, G., & Kleinsmith, A. (2025). Investigating differences in paramedic trainees' multimodal interaction during low and high physiological synchrony. *Proceedings of the 27th International Conference on Multimodal Interaction*, 526–534. https://doi.org/10.1145/3716553.3750796

Kenny, D. A., Kashy, D. A., & Cook, W. A. (2006). Using multilevel modeling to study dyads. In *Dyadic Data Analysis,* (pp. 78–99). Guilford Publications. https://ebookcentral.proquest.com/lib/ethz/detail.action?docID=362550

Konvalinka, I., Xygalatas, D., Bulbulia, J., Schjødt, U., Jegindø, E.-M., Wallot, S., Van Orden, G., & Roepstorff, A. (2011). Synchronized arousal between performers and related spectators in a fire-walking ritual. *Proceedings of the National Academy of Sciences*, *108*(20), 8514–8519. https://doi.org/10.1073/pnas.1016955108

Kruschke, J. K., & Liddell, T. M. (2018). The Bayesian New Statistics: Hypothesis testing, estimation, meta-analysis, and power analysis from a Bayesian perspective. *Psychonomic Bulletin & Review*, *25*(1), 178–206. https://doi.org/10.3758/s13423-016-1221-4

Lahnakoski, J. M., Forbes, P. A. G., McCall, C., & Schilbach, L. (2020). Unobtrusive tracking of interpersonal orienting and distance predicts the subjective quality of social interactions. *Royal Society Open Science*, *7*(8), 191815. https://doi.org/10.1098/rsos.191815

Lieberz, J., Shamay-Tsoory, S. G., Saporta, N., Esser, T., Kuskova, E., Stoffel-Wagner, B., Hurlemann, R., & Scheele, D. (2021). Loneliness and the social brain: How perceived social isolation impairs human interactions. *Advanced Science*, *8*(21), 2102076. https://doi.org/10.1002/advs.202102076

Lokon, E., Li, Y., & Kunkel, S. (2020). Allophilia: Increasing college students' "liking" of older adults with dementia through arts-based intergenerational experiences. *Gerontology & Geriatrics Education*, *41*(4), 494–507. https://doi.org/10.1080/02701960.2018.1515740

Mayo, O., & Gordon, I. (2020). In and out of synchrony—Behavioral and physiological dynamics of dyadic interpersonal coordination. *Psychophysiology*, *57*(6), e13574. https://doi.org/10.1111/psyp.13574

Mayo, O., Lavidor, M., & Gordon, I. (2021). Interpersonal autonomic nervous system synchrony and its association to relationship and performance – a systematic review and meta-analysis. *Physiology & Behavior*, *235*, 113391. https://doi.org/10.1016/j.physbeh.2021.113391

McElreath, R. (2020). *Statistical Rethinking; A Bayesian Course with Examples in R and Stan* (2nd Edition). Chapman and Hall/CRC.

Mitkidis, P., McGraw, J. J., Roepstorff, A., & Wallot, S. (2015). Building trust: Heart rate synchrony and arousal during joint action increased by public goods game. *Physiology & Behavior*, *149*, 101–106. https://doi.org/10.1016/j.physbeh.2015.05.033

Moffat, R., Casale, C. E., & Cross, E. S. (2024). Mobile fNIRS for exploring inter-brain synchrony across generations and time. *Frontiers in Neuroergonomics*, *4*. https://doi.org/10.3389/fnrgo.2023.1260738

Moffat, R., Dumas, G., & Cross, E. S. (2025). *Longitudinal intergenerational hyperscanning reveals indices of relationship formation and loneliness*. bioRxiv. https://doi.org/10.1101/2025.10.14.682029

Noy, L., Levit-Binun, N., & Golland, Y. (2015). Being in the zone: Physiological markers of togetherness in joint improvisation. *Frontiers in Human Neuroscience*, *9*. https://doi.org/10.3389/fnhum.2015.00187

Oliveira-Silva, P., & Gonçalves, Ó. F. (2011). Responding empathically: A question of heart, not a question of skin. *Applied Psychophysiology and Biofeedback*, *36*(3), 201–207. https://doi.org/10.1007/s10484-011-9161-2

Palumbo, R. V., Marraccini, M. E., Weyandt, L. L., Wilder-Smith, O., McGee, H. A., Liu, S., & Goodwin, M. S. (2017). Interpersonal autonomic physiology: A systematic review of the literature. *Personality and Social Psychology Review*, *21*(2), 99–141. https://doi.org/10.1177/1088868316628405

Park, S., Choi, S. J., Mun, S., & Whang, M. (2019). Measurement of emotional contagion using synchronization of heart rhythm pattern between two persons: Application to sales managers and sales force synchronization. *Physiology & Behavior*, *200*, 148–158. https://doi.org/10.1016/j.physbeh.2018.04.022

Paulus, C. (2009). *Der Saarbrücker Persönlichkeitsfragebogen SPF (IRI) zur Messung von Empathie: Psychometrische Evaluation der deutschen Version des Interpersonal Reactivity Index*. https://doi.org/10.23668/psycharchives.9249

Pauly, T., Gerstorf, D., Wahl, H.-W., & Hoppmann, C. A. (2021). A developmental–contextual model of couple synchrony across adulthood and old age. *Psychology and Aging*, *36*(8), 943–956. https://doi.org/10.1037/pag0000651

Perdue, K. L., Westerlund, A., McCormick, S. A., & Nelson, C. A. (2014). Extraction of heart rate from functional near-infrared spectroscopy in infants. *Journal of Biomedical Optics*, *19*(6), 067010. https://doi.org/10.1117/1.JBO.19.6.067010

Pittinsky, T. L., Rosenthal, S. A., & Montoya, R. M. (2011). Measuring positive attitudes toward outgroups: Development and validation of the Allophilia Scale. In L. R. Tropp & R. K. Mallett (Eds), *Moving beyond prejudice reduction: Pathways to positive intergroup relations.* (pp. 41–60). American Psychological Association. https://doi.org/10.1037/12319-002

R Core Team. (2022). *R: A language and environment for statistical computing* [Computer software]. R Foundation for Statistical Computing. http://www.r-project.org/

RStudio Team. (2020). *RStudio: Integrated Development for R* [Computer software]. RStudio, Inc. http://www.rstudio.com/

Rubin, S. E., Gendron, T. L., Wren, C. A., Ogbonna, K. C., Gonzales, E. G., & Peron, E. P. (2015). Challenging gerontophobia and ageism through a collaborative intergenerational art program. *Journal of Intergenerational Relationships*, *13*(3), 241–254. https://doi.org/10.1080/15350770.2015.1058213

Saporta, N., Scheele, D., Lieberz, J., Stuhr-Wulff, F., Hurlemann, R., & Shamay-Tsoory, S. G. (2021). Opposing association of situational and chronic loneliness with interpersonal distance. *Brain Sciences*, *11*(9), 1135. https://doi.org/10.3390/brainsci11091135

Scheele, D., Striepens, N., Güntürkün, O., Deutschländer, S., Maier, W., Kendrick, K. M., & Hurlemann, R. (2012). Oxytocin modulates social distance between males and females. *The Journal of Neuroscience*, *32*(46), 16074–16079. https://doi.org/10.1523/JNEUROSCI.2755-12.2012

Scholkmann, F., Boss, J., & Wolf, M. (2012). An efficient algorithm for automatic peak detection in noisy periodic and quasi-periodic signals. *Algorithms*, *5*(4), 588–603. https://doi.org/10.3390/a5040588

Shafaei, R., Bahmani, Z., Bahrami, B., & Vaziri-Pashkam, M. (2020). Effect of perceived interpersonal closeness on the joint Simon effect in adolescents and adults. *Scientific Reports*, *10*(1), 18107. https://doi.org/10.1038/s41598-020-74859-3

Shamay-Tsoory, S. G., & Kanterman, A. (2024). Away from the herd: Loneliness as a dysfunction of social alignment. *Social Cognitive and Affective Neuroscience*, *19*(1), nsae005. https://doi.org/10.1093/scan/nsae005

Sharika, K. M., Thaikkandi, S., Nivedita, & Platt, M. L. (2024). Interpersonal heart rate synchrony predicts effective information processing in a naturalistic group decision-making task. *Proceedings of the National Academy of Sciences*, *121*(21), e2313801121. https://doi.org/10.1073/pnas.2313801121

Speer, S. P. H., Mwilambwe-Tshilobo, L., Tsoi, L., Burns, S. M., Falk, E. B., & Tamir, D. I. (2024). Hyperscanning shows friends explore and strangers converge in conversation. *Nature Communications*, *15*(1), 7781. https://doi.org/10.1038/s41467-024-51990-7

Strang, A. J., Funke, G. J., Russell, S. M., Dukes, A. W., & Middendorf, M. S. (2014). Physio-behavioral coupling in a cooperative team task: Contributors and relations. *Journal of Experimental Psychology: Human Perception and Performance*, *40*(1), 145–158. https://doi.org/10.1037/a0033125

Stuldreher, I. V., Van Erp, J. B. F., & Brouwer, A.-M. (2023). Robustness of physiological synchrony in wearable electrodermal activity and heart rate as a measure of attentional engagement to movie clips. *Sensors*, *23*(6), 3006. https://doi.org/10.3390/s23063006

Sun, J., Rao, L., & Gao, C. (2018). Extracting heartrate from optical signal of functional near-infrared spectroscopy based on mathematical morphology. *Journal of Innovative Optical Health Sciences*, *11*(03), 1850010. https://doi.org/10.1142/S1793545818500104

Tschacher, W., & Meier, D. (2020). Physiological synchrony in psychotherapy sessions. *Psychotherapy Research*, *30*(5), 558–573. https://doi.org/10.1080/10503307.2019.1612114

UNICEF. (2015). *Innovating for Impact*. Innovating for Impact Wearable Technology Opens New Channels for Connecting the Hardest to Reach to Services. https://www.unicef.org/innovation/stories/innovating-impact

United Nations. (2024). *World population prospects 2024: Summary of results*. United Nations.

Van Marum, R. J. (2020). Underrepresentation of the elderly in clinical trials, time for action. *British Journal of Clinical Pharmacology*, *86*(10), 2014–2016. https://doi.org/10.1111/bcp.14539

Wascher, C. A. F. (2021). Heart rate as a measure of emotional arousal in evolutionary biology. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *376*(1831), 20200479. https://doi.org/10.1098/rstb.2020.0479

WHO. (2019). *What is the evidence on the role of the arts in improving health and well-being? A scoping review*. World Health Organization. https://iris.who.int/handle/10665/329834

WHO. (2025). *From loneliness to social connection*. World Health Organization. https://www.who.int/publications/i/item/978240112360

Wolf, W., Launay, J., & Dunbar, R. I. M. (2016). Joint attention, shared goals, and social bonding. *British Journal of Psychology*, *107*(2), 322–337. https://doi.org/10.1111/bjop.12144