Cardiac synchrony during collaborative drawing: A longitudinal comparison of same generation and intergenerational dyads

Ryssa Moffat[1], Luca A. Naudszus[1], Emily S. Cross[1]

[1]Social Brain Sciences Lab, ETH Zurich

* Correspondence:
Ryssa Moffat
rmoffat@ethz.ch

Emily S. Cross
ecross@ethz.ch

Keywords: intergenerational, collaboration, drawing, physiological synchrony, relationship formation

Abstract

Intergenerational social programs provide opportunities for people of all ages to form new relationships. Furthermore, existing qualitative and behavioural evidence from such programs points to health and wellbeing benefits, yet the physiological consequences of repeated intergenerational encounters remain unknown. A deeper understanding of how such programs shape dyadic physiological responses will illuminate the mechanisms of relationship formation. Across a six-session collaborative drawing program, we tracked cardiac synchrony within 31 intergenerational (older/younger adult) and 30 same generation (younger adult) dyads. Each session dyads completed self-report measures, then drew together and alone, while we recorded participants' actions with motion capture and physiological signals (neural and cardiac) using fNIRS. Collaborative behaviour, self-reported social closeness, and interpersonal distance (i.e., proximity) showed group-specific patterns, whereby interpersonal distance emerged as a promising objective measure of relationship development. Cardiac synchrony did not covary with group, task, an interaction thereof or any measure of behaviour or social closeness–yet there was a trending relationship between collaboration while drawing together and cardiac synchrony for intergenerational dyads only. In summary, cardiac synchrony pointed to marginally enhanced arousal during active collaboration between older and younger adults. Relationship development was better characterised, in this study, by behaviour and self-report measures than cardiac synchrony.

1 Introduction

In response to worrying reports of declining social connection, many communities have begun to employ intergenerational arts programs to improve community members' health and wellbeing [1,2]. Growing behavioural and qualitative data from such programs suggests that intergenerational arts programs offer much needed opportunities to build social connection, reduce stereotypes about other generations, and provide structures for people to engage in activities that add meaning to their lives [2–6]. Intergenerational arts programs that offer repeated contact older and younger generations, as opposed to one-off sessions, seem to yield the greatest improvement in social connectedness and

feelings of affiliation between generations [5]. More broadly, feelings of social connectedness or
closeness (e.g., being strangers vs. friends vs. romantic partners) are known to modulate
physiological signals (e.g., cardiac signals and brain activity) during collaboration [7]. Yet,
longitudinal evidence tracking the extent to which physiological signals index relationship formation
is only just beginning to emerge [8]. A deeper understanding of how repeated intergenerational
collaboration fosters relationship formation, across experiential and physiological levels, holds
potential to inform program structure and resource allocation for intergenerational arts programs.
Moreover, new insights into intergenerational collaborative behaviour can only strengthen efforts to
promote healthy ageing.
47
In this study, intergenerational and same generation dyads completed a 6-week collaborative drawing
program, in which they met as strangers and interacted with the same partner for all 6 sessions. We
examined how collaboration and physiological synchrony, specifically *cardiac synchrony*, evolved
across sessions. Specifically, we tracked how cardiac synchrony levels during collaboration were
shaped by repeated social encounters, the generational constellation (intergenerational vs. same
generation), as well as dyads' feelings of social closeness and interpersonal distance (i.e., physical
proximity to one another).

1.1   Cardiac synchrony an index of shared arousal during collaboration

The cardiovascular system is reactive to changes in both psychological and physiological factors [9].
Fluctuations in heart rate reflect changes in arousal of the central nervous system resulting from
changes in an individual's environment [10]. Increases in heart rate reflect sympathetic activity, which
is associated with excitement and stress. Reductions in heart rate reflect parasympathetic activity,
relating to relaxation and divided attention [11,12]. When people gather in groups of two or more,
changes in their physiological signals including heart rate reflect a mixture of self-regulation and co-
regulation [13,14], which can result in greater temporal alignment of heart rates, per se, and heart rate
variability among two or more people. Precise time-locking of fluctuations can arise from external
environmental factors–take for example a jumpscare (i.e., a startling event) in a horror movie causing
a spike in cinemagoers' heart rates simultaneously. Time-lagged alignment occurs more commonly
in live social interaction involving behavioural coordination, for example turn taking in conversations
[15]. Such alignment in arousal can result from shared attention [16], emotional contagion [17,18], or joint
action [19,20]. In other words, measures of temporal alignment of heart rates and heart rate variability
between individuals, or cardiac synchrony, can shed light on shared fluctuations in physiological
arousal during social interactions. Accordingly, shared fluctuations in arousal, tracked using
measures of cardiac synchrony, are reported to underpin important aspects of collaboration, including
perceptions of group cohesion, satisfaction, commitment, subjective togetherness, trust, and
perceived empathy (see review by Mayo and colleagues [7]).

Several studies have investigated the extent to which cardiac synchrony can predict the outcomes of
collaboration [19,21–24]. Sharika et al. [24] recorded 44 groups of 204 students completing a verbal
consensus-seeking task. The authors reported that cardiac synchrony between group members
predicted the likelihood of reaching consensus significantly better than individuals' heart rates, self-
reported ratings of team function, or duration of the consensus-seeking process. On this basis,
Sharika et al. proposed that cardiac synchrony may be useful as a biomarker of engagement during
collaboration [24]. Gordon et al. [22] compared levels of cardiac synchrony during synchronised
drumming, freely improvised drumming, and resting state (i.e., no movement). They reported that
cardiac synchrony was greater during drumming than baseline but did not differ between drumming
conditions, highlighting the role of joint action in aligning fluctuations in arousal. Gordon et al.

further reported that levels of cardiac synchrony during synchronised drumming predicted levels of
coordination of drumming in the subsequent improvised drumming task. Alignment of arousals
levels, as indexed by cardiac synchrony, may thus be a precursor for successful collaboration [22].

Other studies suggest that perceptual shifts during collaboration drive changes in shared arousal
[19,21,23]. Noy et al. tracked levels of perceived and kinematically measured togetherness when dyads
played a mirror game. Controlling for differences in movement intensity, the authors found that
cardiac synchrony increased during periods of togetherness, more so for perceived than kinematic
togetherness. Periods of togetherness were also notably accompanied by increased heart rates,
reflecting increased arousal [19]. Boukarras et al. [21] recorded grasp timing from dyads told to grasp
two bottle shaped objects according to specific instructions. In half the trials, the instructions
prompted the same or complementary grasping behaviours, and in the other half of trials, the
instructions prompted evoked peer-to-peer or leader-follower grasp dynamics. None of these
prompted grasping behaviours showed a statistically significant relationship between grasping timing
and cardiac synchrony. However, Boukarras et al. did observe heightened levels of cardiac synchrony
when dyads shifted between peer-to-peer to a leader-follower dynamics or from same to
complementary movements. These findings led the authors to suggest that perceptual shifts during
collaboration dynamics, rather than specific patterns of behavioural coordination, may influence
arousal levels, and by extension, cardiac synchrony [21]. Findings from a study by Mitkidis et al. [23]
lend support to the idea that perceptual shifts during collaboration may alter shared arousal. Mitkidis
et al. compared cardiac synchrony in a group that completed interleaved collaborative Lego building
tasks with public goods games to a group that completed only Lego building tasks. They report that
the public goods game led to greater cardiac synchrony during Lego building. They also found that
cardiac synchrony predicted the participants' expectations of their partners during the public goods
game. Mitkidis et al. interpret both results as being reflective of increased awareness of the partner
relating to trust-building processes. Considered together, these findings highlight that perceptual
shifts during collaboration impact arousal levels as quantified by cardiac synchrony. Notably, too
much shared arousal can have deleterious effects on collaboration [25,26], suggesting that co-regulation
of arousal levels is central to successful collaboration [7].

Changes in familiarity that occur within or across encounters may represent a sufficient perceptual
shift to influence shared arousal, and by extension, cardiac synchrony. Fusaroli et al. [27] invited five
groups of 4-5 students to build Lego models in 5-minute blocks, alternating between building alone
and together as a group, and computed measures of heart rate coordination (i.e., the similarity and
stability of signals within groups–complementary measures to cardiac synchrony). The authors
reported that heart rate coordination within groups increased from the first block of collaborative
model-building to the last and that coordination of movement and speech within groups predicted
heart rate coordination [27]. These findings illustrate how increasing familiarity, on a short time scale,
co-occurs with increasing alignment of behaviour and arousal levels. Highlighting the relevance of
understanding changes in dyadic levels of shared arousal over repeated encounters, Gernert et al. [28]
measured cardiac synchrony during psychotherapy sessions targeting affective disorders at two
sessions, approximately two weeks apart. The authors found a positive association between cardiac
synchrony across both sessions and changes in self-reported depressive symptom severity between
the two sessions [28]. Unfortunately, this study does not report the difference in cardiac synchrony
between sessions, leaving the trajectory of shared arousal an open question. While empirical attention
has been paid to first encounters [e.g., 29], little is known about the trajectory of fluctuations in shared
arousal across repeated encounters. To advance our understanding of the prerequisites for
relationships formed over repeated encounters to blossom into robust social bonds and the conditions
that predict relationships fizzling out, we need to document the trajectories of shared arousal across

multiple encounters alongside measures of relationship quality. By considering these trajectories in
same generation and intergenerational dyads, we can provide insights for optimising relationship
formation in intergenerational community programs.

1.2 Relationship quality constrains shared arousal, and by extension, cardiac synchrony

Relationship quality and shared fluctuations in arousal, as measured by cardiac synchrony, are known
to covary, though the effect size derived from a meta-analysis of 26 studies is quite small [7], and the
directionality of the effect is not known. Many of the studies conducted have been dedicated to
understanding shared arousal in romantic partners and are summarised in Mayo et al.'s meta-analysis
[7]. For example, watching emotional videos together has been shown to elicit greater cardiac
synchrony between strangers, relative to friends and romantic partners [17]. Bizzego et al. propose that
greater familiarity between friends and romantic couples results in their autonomic systems being
less attuned to changes in the other person's autonomic responses. This may be a feature of the task
involving non-interactive copresence with both people's attention directed to the videos. Other tasks,
involving co-presence and interaction show the opposite pattern. At a firewalking performance,
Konvalinka et al. [30] recorded cardiac synchrony between firewalkers and their friends and between
firewalkers and strangers. The authors compared these groups and reported greater cardiac synchrony
between firewalkers and friends and family members than strangers. The authors similarly suggest
that social closeness [30] is a driving factor in the coupling of autonomic responses. Focusing on first-
time encounters, Adel et al. [29] recorded cardiac synchrony in dyads comprising strangers who shared
emotional stories. Dyads showed greater cardiac synchrony when they self-reported high levels of
mutual liking. Relationship quality seems to have a more nuanced impact on shared arousal when
collaboration is compared with competition. Danyluck and Page-Gould [11] demonstrated that self-
reported perceived similarity and interest in friendship between dyad members can shape cardiac
synchrony levels differentially for collaboration and competition. Whereas cardiac synchrony is
positively associated with perceived similarity in both conditions, cardiac synchrony is positively
associated with interest in friendship only during collaboration and not during competition [11].
Recently, Andersen et al. [31] recorded cardiac synchrony between visitors who went through a
recreational commercial haunted house in groups, designed to induce fear in the visitors. The authors
reported that cardiac synchrony is closely related to individuals' ratings of subjective arousal and that
cardiac synchrony was greater between dyads within each visitor group who were socially close than
those who were not. The findings contribute to the evidence that social relationships shape co-
regulation through shared fluctuations in arousal [31]. Based on these studies, we would expect that in
interactive contexts evoking average physiological arousal (i.e., non-frightening; not involving
haunted houses or risky behaviour such as firewalking), cardiac synchrony would be highest at a first
encounter and decrease as dyads who began as strangers become acquainted and socially closer.

Self-reported feelings of social closeness to another person tends to result in individuals being
comfortable in closer physical proximity to each other [32–34]. Like social closeness, physical
proximity (henceforth interpersonal distance) is bidirectionally related with shared arousal [31,35]. In a
2x2 design, Jackson et al. [35] recorded groups of people walking freely vs. in synchrony as a group
vs. manipulated arousal levels by having the leader walk above (high arousal) or below (low arousal)
average walking speed. The researchers found that the pack was the tightest (least distance between
walkers) during the condition combining synchronous and high arousal walking. The researchers
propose that their findings reflect how arousal and synchrony can be employed to manufacture group
cohesion [35]. To our knowledge, no studies to date investigate the relationship between shared
arousal measured via cardiac synchrony and interpersonal distance. Nonetheless, previous research
using other measures of physiological synchrony has revealed that arousal levels are likely associated

with interpersonal distance. For example, teams of paramedic trainees who show greater synchrony
in electrodermal activity work physically closer to one another and engage in more cooperative
dialogue compared to teams showing lower synchrony [36]. Interpersonal distance appears to be
associated with shared arousal levels during collaboration, yet additional evidence is needed to
establish this relationship's reliability.

1.3   Cardiac synchrony in intergenerational relationships is poorly documented

Notably, the relationships between shared arousal, social closeness, interpersonal distance and
collaboration have (to our knowledge) only been examined in dyads consisting of two young adults
at a single point in time (as opposed to longitudinally). In very few cases, intergenerational patterns
of shared arousal have been explored within mother-child dyads (see review by DePasquale and
colleagues [37]). Older adults have yet to be represented in such studies, a trend that is observed in
experimental studies more broadly [38,39]. It is critical to remedy the underrepresentation of older
adults as the proportion of older adults in the world's population is steadily climbing [40] and older
adults are at elevated risk of experiencing loneliness [2]. Moreover, longitudinal insights extending
beyond first encounters and covering stages before friendships and romantic relationships are
necessary for a deeper understanding of the physiological underpinnings of relationship development.

1.4   Present study

This research sought to shed light on the physiological mechanisms of collaboration in budding
relationships–both same generation and intergenerational, including older adults. We preregistered
(https://osf.io/xuvy5) the three following experimental aims:

• Document the trajectory of cardiac synchrony over the course of six sessions of
collaborative drawing within both same generation and intergenerational dyads.
• Explore how levels of collaboration visible in drawings (evaluated by external raters)
relate to dyad members' feelings of social closeness, physical closeness, and
performance on a separate collaborative task (i.e., completing a jigsaw puzzle).
• Explore how cardiac synchrony is related to levels of collaboration visible in
drawings. Specifically, we wish to assess whether this relationship differs between
same generation and intergenerational dyads, and the extent to which this relationship
is impacted by dyad's feelings of social closeness, physical closeness, and
performance on a separate collaborative measure (i.e., jointly completing a jigsaw
puzzle).

In reporting these findings, charting cardiac synchrony within intergenerational and same generation
dyads across 6 sessions in an everyday context similar to a community art program, we reveal first
insights into longitudinal patterns of shared arousal, in a unique design that involves older adults. The
implications of this work are central to understanding the behavioural and physiological mechanisms
governing relationship formation and collaboration across the lifespan. Moreover, this work sheds
light on important considerations regarding the implementation of cardiac synchrony as an objective
measure of social connection or collaboration in everyday, real-world settings.

2   Methods

2.1   Participants

Main sample. We recruited 122 community-dwelling participants from Zurich, Switzerland (31 older adults, aged 69+ years; 91 younger adults, aged 18–35 years). Included participants were fluent in German and had no known history of neurological or psychological disorders (e.g., stroke, concussion, ADHD, autism, schizophrenia, or depression). Participants were assigned to intergenerational dyads ($n = 31$) or same generation dyads ($n = 30$) based on availability for sessions (e.g., matching people available at the same day/time for 6 consecutive weeks). All dyads started the study as strangers. Intergenerational dyads comprised one older adult (76 ± 4 years; 18 female, 13 male) and one younger adult (24 ± 4 years; 19 female, 11 male, 1 other). Same generation dyads comprised two younger adults (22.5 ± 3.6 years; 36 female, 24 male). Supplementary Table 1 details the number of individual participants' recordings and dyads' sessions that were excluded, as well as reasons for exclusion.

Our recruitment campaign involved advertising through the University of Zurich's Healthy Longevity Centre, ETH Zurich's DeSciL student pool, clubs and organisations for senior citizens (e.g., theatre groups and choirs/orchestras), and social media. All participants provided written informed consent prior to participation. Ethical approval was obtained from the Ethics Committee of the Canton Zurich (Ref: 2023-01073). The study was conducted according to the Declaration of Helsinki. We report how we determined our sample size, all data exclusions, all manipulations, and all measures in the study.

External ratings of drawings. To obtain ratings of drawings produced by the main sample (described further in 2.3.1), we recruited 148 new participants via Prolific after the main study concluded. To be included, participants had to be over18 years old, proficient English speakers, residents of the United Kingdom. To ensure response quality, we only accepted participants who had previously completed a minimum of 50 tasks on Prolific with a 100% approval rate. Of these participants, 45 were excluded for failing to respond to all attention checks correctly (i.e., by moving all sliders all the way to the right of the sliding scale when prompted). We thus use ratings from 103 raters in our analyses. We did not collect demographic information from these participants. All participants provided written informed consent. Ethical approval was obtained from the Ethics Committee of ETH Zurich (Ref: 2024-N-265). The study was conducted according to the Declaration of Helsinki.

2.2 Procedure

Each dyad completed six weekly sessions (once per week, for six weeks). At each session, participants completed questionnaires querying feelings of loneliness [41] and attitudes toward their own and the other (older or younger) generation [42]. Participants also completed a questionnaire measuring empathy [43] at the first session. Dyads drew alone once and collaboratively twice with oil pastels provided by Caran D'Ache, for a duration of five minutes (Figure 1; Supplementary Figure 1). Participants were not given any prompts regarding what to draw or how to draw (e.g., no suggestions for motifs). Participants were instructed not to talk while drawing, so that speech would not be a confound in comparisons of drawing alone (of particular relevance for neural and cardiac measures). Participants were instructed that they could talk freely during the rest of the session. After drawing, participants completed a collaborative activity, with different activities each week. In the third session, the collaborative activity was to complete a 54-piece jigsaw puzzle. Activities completed on other weeks include sorting pastels, a divergent thinking task, and a joint verbal fluency task. While participants drew and completed collaborative activities, we recorded their cortical activity using functional near-infrared spectroscopy (fNIRS) over the bilateral inferior frontal gyri (IFG) and bilateral temporo-parietal junction (TPJ). Furthermore, we video-recorded

262 participants' movement using a GoPro camera to capture movements of the head and hands. At the
263 end of each session, after we stopped the fNIRS and video recordings, participants completed the
264 Inclusion of Other in Self scale [44]. Analyses of fNIRS recordings, loneliness scores, and attitudes
265 toward generations are reported elsewhere, as are additional specifics regarding the fNIRS recording
266 methodology [8].

A

Figure 1. A) Left: An intergenerational dyad drawing independently on separate pieces of paper during fNIRS recording.
The dyad members are separated by a grey felt divider that obscures their view of each other. Right: The same
intergenerational dyad drawing together on a single piece of paper during fNIRS recording. The colourful lines overlaid
on top of the participants show the 2D motion tracking used to capture interpersonal distance, specifically the distance
between dyad members' neck joints. B) IBI times series of two different dyads, illustrating high and low levels cardiac
synchrony with the produced drawing (see further examples sorted by collaboration ratings in Figure 2).

2.3   Measures

2.3.1 Drawing collaboration

Each drawing that the dyads created was judged by independent raters recruited via Prolific. Raters
($n = 103$) were instructed that they would view 122 drawings, each created by two people at once on
a single piece of paper. They were asked to rate the degree of harmony between the elements of the
drawings (on a sliding scale from 0 "entirely independent" to 100 "entirely coordinated") and the use
of space (on a sliding scale from 0 "divided" to 100 "unified"). To obtain a measure of the perceived
coherence of the drawing, which we propose reflects the level of collaboration during drawing, we

summed the ratings of harmony and use of space. Thus, raw scores of drawing collaboration could
range from 0-200. Examples of drawings with highest, average, and lowest collaboration ratings are
shown in Figure 2.

Figure 2. Examples of drawings organised by rating of collaboration. Top row = highest collaboration ratings; middle
row = average collaboration ratings; bottom row = lowest collaboration ratings. Ratings are derived by summing raters'
ratings of the harmony between the elements of the drawings (0 "entirely independent" to 100 "entirely coordinated") and
use of space (0 "divided" to 100 "unified"). The left-most drawings in the top and bottom row are also presented in
Figure 1B alongside the IBI time series recorded while the drawings were made.

2.3.2 Puzzle collaboration

After completing the drawing component of the third session, dyads completed a puzzle as the
collaborative activity for that session. Speaking was permitted. Dyads had six minutes to complete a
54-piece jigsaw puzzle of cartoon animals in jungle (pieces' dimensions: ~ 8 x 8 cm). Dyads' scores
corresponded to the number of pieces assembled in 6 minutes. In the case that participants assembled
all pieces in less than 6 minutes, the remaining number of seconds to 6 minutes was divided by 6.66
(the average number of seconds per piece if completed in 6 minutes; 360 s /54 piece = 6.66 s) and
added to a score of 54.

2.3.3 Social Closeness

To understand how collaborative behaviour shapes feelings of social closeness, we administered the Inclusion of Other in Self (IOS) scale (Aron et al., 1992 after dyads drew together and completed the final collaborative task. Participants viewed seven sets of overlapping circles representing the self and the partner, ranging from no overlap to near complete overlap of the two circles, and indicate which set of circles best represents their relationship with their partner (i.e., the other dyad member). To capture cumulative social closeness within dyads and differences in feelings of social closeness within dyads, we computed summed and absolute difference IOS scores per dyad [21,45]. We refer to these measures as social closeness (Sum) and social closeness (Dif).

2.3.4 Interpersonal distance

We used the videos recorded at each session to track upper body movements while dyads drew together. Specifically, we used OpenPose [46] to estimate 2D pose per frame, returning coordinates for predifined body parts. To assess interpersonal distance (sometimes referred to as proximity in other work), we computed the average distance between the coordinates of each dyad members' neck joint for each 5-minute instance of drawing together. In our analyses, this measure is referred to as distance.

2.4 Data analysis

2.4.1 Extracting inter-beat intervals

As per our preregistration, we extracted the cardiac signal by identifying peaks in HbO concentrations [47]. We cleaned the HbO signal from neural and further physiological components such as Mayer waves by upsampling it to 100 Hz and applying a zero-phase third-order Butterworth IIR band-pass filter [48] with cut-offs at 0.5 and 2.5 Hz [49]. Then, we smoothed the signals using a Savitzky-Golay filter with window length 15 and a 5^{th}-order polynomial. We identified peaks with the AMPD algorithm [47], implemented in the python *ampdlib* package [50], and transformed the signal peak timestamps to an inter-beat interval (IBI) time series, smoothly interpolated at 4 Hz. We originally preregistered ARIMA cleaning [51], but in practice it reduced the signal to near-zero fluctuations, largely removing the slow structure that may reflect the physiology of interest in this context. Given the lower signal-to-noise ratio of fNIRS-derived IBIs when compared to ECG (electrocardiography) signals, and our total number of samples (fNIRS signals recorded at ~ 5 Hz), we instead analysed IBI series on their original scale to retain physiologically meaningful variance.

2.4.2 Calculating synchrony

We calculated cross-correlation functions (CCFs) between all dyads' cleaned physiological signals in each session and activity across time lags of +/- 3 seconds (-12 to +12 lags at 4 Hz) and retained non-absolute CCF values. Examples of time series yielding high and low cardiac synchrony estimates are visualised in Figure 1B.

2.4.3 Group-level analysis

We completed our preregistered exploratory statistical analyses using a Bayesian approach to multilevel regression [52]. We fit models using the brms package [version 22.2; ,53] in the R language [version 4.5.1; ,54] within the RStudio IDE [version 2025.05.1+513; ,55]. We report and interpret the posterior distribution of relevant model parameters using a 95% credible interval, calculated via the 95% highest posterior density region (HPD) method [52]. For readers familiar with Frequentist approaches that include p-values, we recommend perusing Kruschke & Liddell [56], and we share a (simplified) heuristic for

interpreting HPDs: Parameter and comparison estimates can be considered *substantial* when the HPD
does not contain zero and can be considered trends when the tip of an HPD-tail overlaps with zero.

Measures of collaboration, social closeness, and interpersonal distance. To assess the extent to
which collaboration, social closeness, and interpersonal distance changed across the six sessions, as
well as relationships between measures, we fit a series of models in the following structure:
*MeasureA ~ 1 + MeasureB * Group * Session + (1|DyadID)*. We extracted point estimates for all
sessions combined and slopes across sessions, and subsequently computed contrasts between groups
(intergenerational vs. same generation).

Cardiac synchrony for all sessions combined and across sessions. We first sought to establish if
cardiac synchrony levels differed substantially between real and pseudo dyads. We fit the following
model: *CardiacSynchrony ~ DyadType * Group * Task * Session*. We did not include random
intercepts at this stage, as the total number of real and pseudo dyads exceeded 6000. Random
intercepts, accounting for dyad-level variance, were included in subsequent group level analyses.
Contrasts between real and pseudo dyads were computed to confirm that some component of the
cardiac synchrony was attributable to real social interaction, as opposed to similar experiences or
homeostatic biological rhythms.

**Relationships between cardiac synchrony and measures of collaboration, social closeness, and
interpersonal distance.** First, we fit a model to the cardiac synchrony data, which we averaged
across all lags. The model was: *CardiacSynchrony ~ Group * Drawing Collaboration + Group *
Puzzle Collaboration + Group * Social Closeness (Sum)+ Group * Social Closeness (Dif) + Group *
Interpersonal Distance + Session + (1|DyadID)*. Subsequently, we fit a model to examine
relationships between cardiac synchrony at separate lags and our measures. Models were structured
as follows and fit per measure: *CardiacSynchrony ~ Lag * Group * Measure + (1|DyadID)*. For both
steps, we extracted point estimates for all sessions combined and slopes across sessions, and
subsequently computed contrasts between groups (intergenerational vs. same generation).

3   Results

3.1   Behavioural and self-report measures

3.1.1 Contrasts between groups
See Table 1 for estimates and 95% HPD (highest posterior density region) for each measure.
Analyses of drawing collaboration, puzzle collaboration and interpersonal distance are original to this
study. Social closeness (Sum) and (Dif) have previously been analysed by our research team [8]; we
include these analyses here for completeness. **Drawing collaboration.** Drawing collaboration
differed substantially between groups, with the intergenerational drawings showing greater
collaboration than same generation drawings. Drawing collaboration did not change substantially
across sessions for either group, though there was a trend of intergenerational dyads showing a
greater increase in collaboration than same generation dyads. **Puzzle collaboration.** Puzzle
collaboration differed markedly between groups, with same generation dyads scoring substantially
higher than intergenerational dyads. **Social closeness (Sum).** Summed social closeness was
substantially greater for intergenerational dyads than same generation dyads. Summed social
closeness increased substantially across sessions for both groups at a similar rate. **Social closeness
(Dif).** Intergenerational dyads showed substantially greater differences in dyadic social closeness
than same generation dyads. Dyadic differences in social closeness decreased across sessions
substantially for same generation dyads and showed the same trend for intergenerational dyads.

Interpersonal distance. Interpersonal distance showed a trending group difference, with same
generation dyads showing substantially lower interpersonal distance than intergenerational dyads.
Interpersonal distance increased substantially across sessions for same generation dyads only.
Comparisons between groups showed a trending group difference, where the slope of change in
interpersonal distance across sessions was more positive for same generation than intergenerational
dyads.

Table 1. Standardised estimates computed per behavioural and self-report measure with HPD in square brackets. Point
estimates for all six sessions combined are presented first, followed by the estimates of the slope of change across
sessions in the lower section. No slope of change is available for puzzle collaboration, as dyads only completed the
puzzle task once, at the third session.

	Drawing Collaboration	Puzzle Collaboration	Social Closeness (Sum)	Social Closeness (Dif)	Interpersonal Distance
Sessions combined					
Intergenerational	0.20 [0.08, 0.32]	-0.50 [-0.59, -0.42]	0.17 [0.7, 0.26]	0.13 [0.03, 0.23]	0.07 [-0.18, 0.03]
Same gen	-0.03 [-0.15, 0.08]	0.45 [0.36, 0.53]	-0.18 [-0.27, -0.08]	-0.08 [-0.18, 0.03]	-0.06 [-0.04, 0.17]
Intergen-Same gen	0.23 [0.07, 0.40]	-0.95 [-1.07, -0.84]	0.35 [0.22, 0.47]	0.20 [0.07, 0.34]	-0.13 [-0.28, 0.00]
Change across six sessions					
Intergenerational	0.04 [-0.01, 0.93]	-	0.22 [0.17, 0.27]	-0.05 [-0.10, 0.00]	0.02 [-0.02, 0.07]
Same gen	-0.01 [-0.07, 0.04]	-	0.16 [0.11, 0.21]	-0.07 [-0.13, -0.02]	0.07, [0.01, 0.12]
Intergen-Same gen	0.05 [0.00, 0.11]	-	0.09 [-0.02, 0.20]	0.02 [-0.03, 0.07]	-0.04 [-0.10, 0.00]

3.1.2 Relationships between measures

Relationships between measures of collaboration, self-reported social closeness, and interpersonal
distance per group are illustrated in Figure 3. Drawing collaboration was substantially positively
associated with puzzle collaboration for the intergenerational group only, yielding a substantial
difference between intergenerational and same generation dyads (ß = 0.26, [0.08, 0.42]). Drawing
collaboration was substantially positively associated with summed social closeness for both groups,
but the association was stronger for same generation dyads, relative to intergenerational dyads (ß = -
0.16, HPD = [-0.30, -0.02]). Drawing collaboration was not substantially associated with differences
in dyadic social closeness. Drawing collaboration was substantially negatively associated with
interpersonal distance for intergenerational dyads and substantially positively associated with
interpersonal distance for same generation dyads, yielding a substantial group difference (ß = -0.21,
HPD = [-0.25, -0.07]). Puzzle collaboration was substantially negatively associated with summed
social closeness for same generation dyads only, resulting in a substantial group difference (ß = 0.18,
HPD = [0.06, 0.30]). Puzzle collaboration was also substantially negatively associated with
differences in dyadic social closeness for same generation dyads only, resulting in a substantial group
difference (ß = 0.21, HPD = [0.09, 0.33]). Puzzle collaboration was substantially negatively
associated with interpersonal distance for same generation dyads only, and no substantial group
difference was observed. Summed social closeness was substantially positively associated with
dyadic differences in social closeness for both groups and did not differ substantially between
groups. Interpersonal distance was substantially positively associated with summed social closeness

for intergenerational dyads and substantially negatively associated with summed social closeness for
same generation dyads, with a substantial difference between groups (ß = 0.22, HPD = [0.08, 0.36]).
Interpersonal distance showed a trending negative association with differences in dyadic social
closeness for intergenerational dyads and a substantial positive association for same generation
dyads, yielding a substantial difference between groups (ß = -0.20, HPD = [-0.35, 0.06]).

Figure 3. Standardised estimates of relationship between behavioural and self-report measure per group. Black squares
indicate that relationships are substantial (i.e., 95% HPD does not overlap with zero). Note: the relationship between
Interpersonal Distance and Social Closeness (Dif) for same generation dyads is trending, rather than substantial, but in a
black square to facilitate interpretation of results. Standardised parameter estimates per measure shown with 95% HPD in
Supplementary Table 2.

In plain language, these findings can be summarised as follows: Intergenerational dyads'
collaboration was similar across different tasks. Dyads who drew more collaboratively also
assembled more pieces of the jigsaw puzzle collaboratively. For both groups, dyads' summed social
closeness predicted their level of drawing collaboration (i.e., both members feeling very close co-
occurred with higher drawing collaboration). Intergenerational dyads who sat physically closer
together made more collaborative-looking drawings, whereas the same was true for same-generation
dyads who sat further apart. Intergenerational dyads who sat closer together reported feeling socially
closer to each other, whereas same-generation dyads who sat further apart reported feeling socially
closer to each other.

3.2 Cardiac synchrony

3.2.1 Comparing real and pseudo dyads

All sessions combined. For all lags averaged and separate lags, we observed no substantial
difference in cardiac synchrony levels between real and pseudo dyads for drawing together or alone
(Supplementary Figure 2). **Across sessions.** For all lags averaged and separate lags, the rate of
change in cardiac synchrony across sessions (all lags averaged) did not differ substantially between
real and pseudo dyads for drawing together or alone (Figure 4; Supplementary Figure 3).

We interpret these findings as follows: Any two individuals meeting the inclusion criteria, who
experience the specific dyadic experimental environment applied here would show comparable levels
of cardiac synchrony. True interaction with an assigned partner does not appear to contribute

uniquely to cardiac synchrony levels. We present the further analyses in accordance with our
preregistration.

3.2.2 Comparisons between drawing conditions and groups

448 **All sessions combined.** For all lags averaged, we observed no substantial effects of drawing task,
449 group or task*group interactions on cardiac synchrony levels. **Across sessions.** For all lags averaged,
450 we observed no substantial main effect (i.e., no substantial change in cardiac synchrony levels across
451 sessions) and no substantial effects of drawing task, group or task*group interactions on the slope of
452 cardiac synchrony across sessions.

Figure 4. Trajectories of cardiac synchrony in real and pseudo dyads across sessions. Top panel shows individual data
points (dots) and estimated slopes (lines). Bottom panel zooms in on the estimated slopes, with grey shading showing
95%, 89%, and 50% intervals of the posterior predictive distributions. Neither the overall level of cardiac synchrony for
all sessions combined nor the rate of change in cardiac synchrony across sessions differed substantially between real and
pseudo dyads. Though real dyads' cardiac synchrony the trajectories across sessions may appear to differ, the 95 % HPD
overlaps > 10 % (threshold for a trend, Section 2.4.3) for drawing alone and together.

3.3   Relationships between cardiac synchrony, behaviour, and self-report measures

As per our preregistration, we first explored relationships between cardiac synchrony with all lags
averaged, collaborative behaviour, social closeness, and distance. We found no substantial
relationships between cardiac synchrony and either measure of collaboration, social closeness or
interpersonal distance. Also in line with our preregistration, we explored relationships between
cardiac synchrony and each measure per lag (n = 25; from -3000 ms to 3000 ms in 250 ms
increments). We observed no substantial relationships for any measure and no substantial difference
between groups at any lag (Figure 5). For drawing collaboration, intergenerational dyads showed a
trending positive relationship between cardiac synchrony and drawing collaboration at all lags except

-3000 ms, -2750 ms, -2500 ms, and -1750 ms. Note: 95% HPD with no overlap with 0 indicates a
slope substantially different than 0, < 10 % overlap with 0 indicates a trending difference from 0. At
the four lags that did not show trending differences, the degree of 95 % HPD overlap with 0 ranged
from 10-14 %. For intergenerational dyads, positive lags reflect the older dyad member's IBI signal
leading or coming earlier in time, and negative lags reflect the younger member's IBI signal leading
or coming earlier in time. As we observed trending positive and negative lags trending, there is no
evidence that a dyad member of a certain generations fills a leader role.

Figure 5. Standardised estimates of relationship between cardiac synchrony and session number, as well as behavioural measures of collaboration, self-reported measures of social closeness, and interpersonal distance per lag (n = 25; from -3000 ms to 3000 ms in 250 ms increments). No substantial relationships observed. Drawing collaboration panel shows trending relationships between cardiac synchrony and drawing collaboration scores for intergenerational dyads at all but 4 lags. See Supplementary Figure 4 for visualisation including error bars showing 95 % HDP.

4   Discussion

In this study, we charted cardiac synchrony, collaborative behaviours and social closeness in intergenerational and same generation dyads across 6 sessions in an everyday context resembling a community art program. Dyads' collaborative behaviours and measures of social closeness showed nuanced relationships, primarily characterised by greater collaboration in cases of greater social closeness. For cardiac synchrony–both averaged and separate lags–we found no substantial differences between real and pseudo dyads overall or across sessions, no substantial task or group differences among real dyads overall or across sessions, and no substantial relationships between cardiac synchrony and any measure of behaviour. We found trends of substantial relationships for intergenerational dyads between cardiac synchrony and drawing collaboration at all but four lags, with the four non-trending lags showing the same numerical relationship. We propose that these findings, considered together, suggest that cardiac synchrony is to a limited extent sensitive to higher stakes, open-ended, collaboration. We discuss the implications of these findings below.

4.1   Relationships between behavioural and self-report measures

Intergenerational and same generation dyads exhibited some similarities in behavioural measures alongside marked differences. For both groups, greater cumulative feelings of social closeness within dyads co-occurred with greater ratings of collaboration of the dyads' drawings. This finding is consistent with previous findings [57–59]. Notably, however, same generation dyads who reported

more similar feelings of social closeness (as opposed to intergenerational dyads) completed the jigsaw puzzle more quickly. This pattern was not present in intergenerational dyads. We propose that drawing and puzzle assembly, both collaborative tasks, differ in the clarity of the goal. The goal of puzzle assembly is clear to both dyad members (i.e., correctly assemble all pieces before time runs out) while the goal of drawing may be more open-ended, based on the amount and specificity of dyads' verbal planning prior to drawing and the process of co-drawing without talking. For reasons relating to experimental control, most studies to date have focused collaborative tasks with clear end goals [11,23,27,60]. The value of the large dataset and analyses that we present here lies in their ecological validity. That is, they reflect everyday collaboration arguably more accurately than some laboratory-based measures reported by previous studies. Everyday collaboration inherently entails free choice regarding the extent to which one engages and mutual adaptation to one's partner, which have consequences for the development of social relationships.

We did not expect groups to differ in how close they sat and/or leaned toward one another while drawing together. Yet, intergenerational dyads drew sitting closer together than same generation dyads. Further, the relationships between interpersonal distance and feelings of social closeness, measured as a cumulative score per dyad and the discrepancy between dyad members, was opposing between groups. Intergenerational dyads sat closer when they felt more socially close or had smaller discrepancies in their feelings of social closeness–showing the pattern we anticipated based on existing findings [32–34]. Same generation dyads, on the other hand, sat closer when they felt less socially close or had greater discrepancies in their feelings of social closeness. These opposite patterns suggest that reduced levels of social comfort during collaboration may amplify dyads' perceptions of the stakes of the collaboration and result in compensatory behaviours, such as sitting closer to one another. Previous works corroborates this interpretation [61–64]. Thus, interpersonal distance in collaboration may be a promising objective measure of social closeness and relationship development within recently acquainted dyads.

4.2   Cardiac synchrony not sensitive to true interaction, unless stakes are increased

Our analyses of cardiac synchrony revealed no substantial differences between real and pseudo dyads overall or across sessions, no substantial task or group differences among real dyads overall or across sessions, and no substantial relationships between cardiac synchrony and any measure of behaviour. This was true for averaged and separate lags. The absence of differences between real and pseudo dyads can be interpreted as meaning that any two individuals meeting the inclusion criteria of our study (whether same generation or intergenerational), who drew following the same prompts given by the researchers, in the same context are likely to exhibit comparable levels of cardiac synchrony, whether or not they are in the same room at the same time or acquainted. In other words, the levels of cardiac synchrony observed in this study do not index *true* interaction between two people. Moreover, the levels of cardiac synchrony observed in this study do not index changes in collaborative behaviour or feelings of social closeness across repeated encounters.

A small number of studies have previously reported that cardiac synchrony does not differ between real and pseudo dyads [60,65]. Flory et al. [65] present alternative explanations. One is that cardiac synchrony is a spurious finding (i.e., could occur for any two people in the world). We do not think this interpretation applies here as the experimental setting, structure, and inclusion criteria provide substantial structure to the experience of each participant–that is, the description of cardiac synchrony as completely spurious is not founded. Another of Flory et al.'s alternative explanations is that participants operate in a common psychophysiological mode (i.e., similar mental state evoked by the experimental design), aligning with Danyluck and Page-Gould's [11(p. 1)] view that "physiological

synchrony could simply reflect the mutually experienced demands of a shared environment". While
this is in line with our rationale against spurious cardiac synchrony, we disagree on the basis that we
observed trending relationships between cardiac synchrony and drawing collaboration for
intergenerational dyads for 21 of 25 lags. We propose that these relationships point to a noteworthy,
albeit non-substantial, influence of high stakes collaboration on cardiac synchrony.

We observed no task or group differences in cardiac synchrony measured within real dyads. Prior
research has also reported the absence of task-specific or performance-related changes in cardiac
synchrony [21,22,27,66]. More specifically, Gordon et al. [22] detected evidence for task differences in
cardiac synchrony when participants drummed together at a predetermined speed or drummed freely,
improvising together. Similarly, other authors report having found no evidence to support
associations between cardiac synchrony and task performance. Boukarras et al. [21] report that
reaction times in a dyadic grasping task are not associated with cardiac synchrony. Behrens et al. [66]
report that cooperative success is not associated with cardiac synchrony when dyads played the
Prisoners Dilemma game. Strang et al. report that cooperative success is not associated with cardiac
synchrony when dyads play Tetris with one person controlling object rotation and the other
controlling the positioning [60]. Finally, with respect to measures of social connection and closeness,
heart rate coordination (consisting of alternate measures to cardiac synchrony) during Lego building
was reported not to be associated with measures of perceived group relatedness or competence, but
rather with speech and movement dynamics [27]. Interested readers can find further examples in Mayo
and Gordon's review [26]. In summary, despite the large number of studies dedicated to understanding
cardiac synchrony on the grounds that cardiac synchrony can offer a window into collaboration and
social connection, the extent to which this is true remains a topic worthy of debate (as well as
replication).

From a practical perspective, the consistency of non-differences in cardiac synchrony across our
preregistered exploratory analyses suggest that cardiac synchrony is not an optimal measure for
predicting collaborative outcomes or relationship development in real-world environments. Given the
mounting interest expressed in using wearable devices to collect objective insights relating to social
wellbeing [67–70], we offer the following recommendation: Organisations, policy makers, and
researchers aspiring to use wearable physiological monitoring to track changes in social behaviours
and social wellbeing (i.e., social connection) should consider the low explanatory power of cardiac
synchrony demonstrated here and pilot extensively before dedicating funding to large-scale roll-outs
of these devices to measure program engagement or success. We feel justified in making this
recommendation based on the sample size (366 samples involving within-participant/dyad design,
which reduces inter-individual variability in our analyses). Moreover, other work by our research
group shows that inter-brain synchrony may index changes in social wellbeing with greater
sensitivity [8], though we acknowledge the logistical and financial barriers to larger-scale introduction
of brain imaging measures.

Nonetheless, from an empirical perspective, it is promising that collaborative drawing scores show a
trending relationship with cardiac synchrony levels within real intergenerational dyads. This suggests
that further research could uncover which social, cognitive, and behavioural conditions need to be
met for cardiac synchrony to be sensitive to true dyadic interaction. Based on the behavioural data
reported here, we believe these conditions should include higher perceived stakes for successful
collaboration. Our behavioural data revealed that intergenerational dyads' drawings were rated more
collaborative than same generation dyads', and that the elevated levels of drawing collaboration
emerged from the group (i.e., intergenerational dyads) that reported lower feelings of social closeness
within dyads, more disparate feelings of social closeness within dyads, and sat closer together while

drawing together (Table 1). In concert, these behaviours highlight how lower levels of social comfort
during collaboration may raise the stakes of the collaboration and may, in turn, contribute to shared
fluctuations in arousal during collaboration.

4.3   Limitations

In using the term "higher stakes collaboration", we are attempting to articulate the origin of the
decreased social comfort, or increased social tension, as it relates to reputation management [71,72] or
the additional cognitive load stemming from interpersonal uncertainty [i.e., lack of common ground: ,73,74]. We
are open to this being conceptualised in other ways. Follow up experiments could probe the extent to
which individual strategies and subjective experiences shed light on the nature of such collaboration.

Measures of dyadic synchrony can influence observed synchrony levels [75]. We considered this
carefully before preregistering our planned pipeline to calculate IBI time series. We compared an
cardiac synchrony approach implemented previously with fNIRS [76] with more common IBI
approaches [21,29]. We opted for IBI to align our work with existing studies examining cardiac
synchrony. Future research could use non-linear methods such as cross recurrence quantification
analysis (CQRA), which is suited for non-stationary signals such as IBI [77]. With respect to our
comparisons of real and pseudo dyads' levels of cardiac synchrony, a complementary approach
would be to employ segment-shuffled surrogates within each dyad. In this method, each person's
cardiac time series is cut into short segments that are then randomly reordered so that the overall
signal properties are preserved but true timing is disrupted [e.g., 15]. This would reveal whether true
dyads display cardiac synchrony on top of task-driven similarity. Duration of cardiac signals has
been demonstrated to influence the likelihood of finding significant cardiac synchrony, particularly
for smaller group sizes such as dyads [78]. From this perspective, our design is optimised to identify
true cardiac synchrony, should it be present. Nonetheless, future studies could use dedicated heart-
rhythm monitoring, such as electrocardiograms, to examine free-choice, unstructured collaboration in
intergenerational dyads. In suggesting this, we recognise that a complete replication of this study is
likely not feasible, and we support smaller targeted replication efforts.

5   Conclusion

We charted cardiac synchrony, collaborative behaviour, social closeness and interpersonal distance
within intergenerational and same generation dyads across six sessions involving a collaborative
drawing task. Our analyses of collaborative behaviour, self-reported social closeness, and
interpersonal distance, when considered together, suggest that social comfort drives relationships
between social closeness, interpersonal distance and collaboration. Of these, interpersonal distance
emerged as a promising objective measure of relationship development. We found that cardiac
synchrony did not co-vary substantially with group, task, an interaction thereof or collaborative
behaviour, social closeness or interpersonal distance. Nonetheless, we did find a trending relationship
between collaboration while drawing together and cardiac synchrony that was specific to
intergenerational dyads. These findings have implications for understanding the mechanisms
underpinning social interaction within and between generations and for the implementation of cardiac
synchrony as an objective measure of social connection or collaboration in everyday, real-world
contexts.

Author contributions

RM: Conceptualisation, Equipment donation, Data curation, Formal analysis, Investigation, Visualisation, Writing – original draft, Writing – review & editing. LN: Data curation, Validation, Writing – original draft. ESC: Conceptualisation, Funding acquisition, Project administration, Writing – review & editing.

Data availability statement

These data were collected after the submission of the following preregistration: https://osf.io/xuvy5. The data collected and analysed for this study can be found in our repository entitled 'Cardiac synchrony during collaborative drawing: Intergenerational and longitudinal perspectives' on OSF: https://osf.io/akmqw.

Acknowledgements

We thank the survey center operated by UZH Healthy Longevity Center, University of Zurich, for their assistance with recruitment of older adults from the Zurich community. We thank Tessa Portier and Medea Häuselmann for their assistance with data collection, Linda Fanconi for scheduling participants' sessions, Alistair Gadola for coding participant behaviour, and Fanny Mougel for digitising the drawings. We thank Chantal Nagel for assisting with literature review. We thank Courtney Casale and Tessa Portier for their help with proofreading.

Conflict of interest

The authors declare no potential conflict of interest.

Funding

Two Cortivision Photon Cap devices were provided to RM through the Cortivision Pathfinder Program (grant number CPP-2023/09/01). Oil pastels were provided to RM by Caran D'Ache (agreement number 712517). RM, LN, and ESC were supported by the Social Brain Sciences Lab at ETH Zurich.

References

1. WHO. (2019). *What is the evidence on the role of the arts in improving health and well-being? A scoping review*. World Health Organization. https://iris.who.int/handle/10665/329834

2. WHO. (2025). *From loneliness to social connection*. World Health Organization. https://www.who.int/publications/i/item/978240112360

3. Anderson, S., Fast, J., Keating, N., Eales, J., Chivers, S., & Barnet, D. (2017). Translating knowledge: Promoting health through intergenerational community arts programming. *Health Promotion Practice*, *18*(1), 15–25. https://doi.org/10.1177/1524839915625037

4. Jenkins, L. K., Farrer, R., & Aujla, I. J. (2021). Understanding the impact of an intergenerational arts and health project: A study into the psychological well-being of participants, carers and artists. *Public Health*, *194*, 121–126. https://doi.org/10.1016/j.puhe.2021.02.029

5. Lokon, E., Li, Y., & Kunkel, S. (2020). Allophilia: Increasing college students' "liking" of older adults with dementia through arts-based intergenerational experiences. *Gerontology & Geriatrics Education*, *41*(4), 494–507. https://doi.org/10.1080/02701960.2018.1515740

6.   Rubin, S. E., Gendron, T. L., Wren, C. A., Ogbonna, K. C., Gonzales, E. G., & Peron, E. P. (2015). Challenging
gerontophobia and ageism through a collaborative intergenerational art program. *Journal of Intergenerational*
Relationships, *13*(3), 241–254. https://doi.org/10.1080/15350770.2015.1058213

7.   Mayo, O., Lavidor, M., & Gordon, I. (2021). Interpersonal autonomic nervous system synchrony and its association
to relationship and performance – a systematic review and meta-analysis. *Physiology & Behavior*, *235*, 113391.
https://doi.org/10.1016/j.physbeh.2021.113391

8.   Moffat, R., Dumas, G., & Cross, E. S. (2025). *Longitudinal intergenerational hyperscanning indexes changes in*
social connection. bioRxiv. https://doi.org/10.1101/2025.10.14.682029

9.   Oliveira-Silva, P., & Gonçalves, Ó. F. (2011). Responding empathically: A question of heart, not a question of skin.
Applied Psychophysiology and Biofeedback, *36*(3), 201–207. https://doi.org/10.1007/s10484-011-9161-2

10.   Wascher, C. A. F. (2021). Heart rate as a measure of emotional arousal in evolutionary biology. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *376*(1831), 20200479.
https://doi.org/10.1098/rstb.2020.0479

11.   Danyluck, C., & Page-Gould, E. (2019). Social and physiological context can affect the meaning of physiological
synchrony. *Scientific Reports*, *9*(1), 8222. https://doi.org/10.1038/s41598-019-44667-5

12.   Tschacher, W., & Meier, D. (2020). Physiological synchrony in psychotherapy sessions. *Psychotherapy Research*,
30(5), 558–573. https://doi.org/10.1080/10503307.2019.1612114

13.   Butler, E. A. (2011). Three views of emotion regulation and health. *Social and Personality Psychology Compass*,
5(8), 563–577. https://doi.org/10.1111/j.1751-9004.2011.00372.x

14.   Ferrer, E., & Helm, J. L. (2013). Dynamical systems modeling of physiological coregulation in dyadic interactions.
International Journal of Psychophysiology, *88*(3), 296–308. https://doi.org/10.1016/j.ijpsycho.2012.10.013

15.   Coutinho, J., Pereira, A., Oliveira-Silva, P., Meier, D., Lourenço, V., & Tschacher, W. (2021). When our hearts beat
together: Cardiac synchrony as an entry point to understand dyadic co-regulation in couples. *Psychophysiology*,
58(3), e13739. https://doi.org/10.1111/psyp.13739

16.   Golland, Y., Arzouan, Y., & Levit-Binnun, N. (2015). The mere co-presence: Synchronization of autonomic signals
and emotional responses across co-present individuals not engaged in direct interaction. *PLOS ONE*, *10*(5),
e0125804. https://doi.org/10.1371/journal.pone.0125804

17.   Bizzego, A., Azhari, A., Campostrini, N., Truzzi, A., Ng, L. Y., Gabrieli, G., Bornstein, M. H., Setoh, P., &
Esposito, G. (2019). Strangers, friends, and lovers show different physiological synchrony in different emotional
states. *Behavioral Sciences*, *10*(1), 11. https://doi.org/10.3390/bs10010011

18.   Park, S., Choi, S. J., Mun, S., & Whang, M. (2019). Measurement of emotional contagion using synchronization of
heart rhythm pattern between two persons: Application to sales managers and sales force synchronization.
Physiology & Behavior, *200*, 148–158. https://doi.org/10.1016/j.physbeh.2018.04.022

19.   Noy, L., Levit-Binun, N., & Golland, Y. (2015). Being in the zone: Physiological markers of togetherness in joint
improvisation. *Frontiers in Human Neuroscience*, *9*. https://doi.org/10.3389/fnhum.2015.00187

20.   Palumbo, R. V., Marraccini, M. E., Weyandt, L. L., Wilder-Smith, O., McGee, H. A., Liu, S., & Goodwin, M. S.
(2017). Interpersonal autonomic physiology: A systematic review of the literature. *Personality and Social*
Psychology Review, *21*(2), 99–141. https://doi.org/10.1177/1088868316628405

21.   Boukarras, S., Placidi, V., Rossano, F., Era, V., Aglioti, S. M., & Candidi, M. (2025). Interpersonal physiological
synchrony during dyadic joint action is increased by task novelty and reduced by social anxiety. *Psychophysiology*,
62(3), e70031. https://doi.org/10.1111/psyp.70031

22.   Gordon, I., Gilboa, A., Cohen, S., Milstein, N., Haimovich, N., Pinhasi, S., & Siegman, S. (2020). Physiological and
behavioral synchrony predict group cohesion and performance. *Scientific Reports*, *10*(1), 8484.
https://doi.org/10.1038/s41598-020-65670-1

23.   Mitkidis, P., McGraw, J. J., Roepstorff, A., & Wallot, S. (2015). Building trust: Heart rate synchrony and arousal
during joint action increased by public goods game. *Physiology & Behavior*, *149*, 101–106.
https://doi.org/10.1016/j.physbeh.2015.05.033

24. Sharika, K. M., Thaikkandi, S., Nivedita, & Platt, M. L. (2024). Interpersonal heart rate synchrony predicts effective
information processing in a naturalistic group decision-making task. *Proceedings of the National Academy of*
Sciences, *121*(21), e2313801121. https://doi.org/10.1073/pnas.2313801121

25. Dimitroff, S. J., Kardan, O., Necka, E. A., Decety, J., Berman, M. G., & Norman, G. J. (2017). Physiological
dynamics of stress contagion. *Scientific Reports*, *7*(1), 6168. https://doi.org/10.1038/s41598-017-05811-1

26. Mayo, O., & Gordon, I. (2020). In and out of synchrony—Behavioral and physiological dynamics of dyadic
interpersonal coordination. *Psychophysiology*, *57*(6), e13574. https://doi.org/10.1111/psyp.13574

27. Fusaroli, R., Bjørndahl, J. S., Roepstorff, A., & Tylén, K. (2016). A heart for interaction: Shared physiological
dynamics and behavioral coordination in a collective, creative construction task. *Journal of Experimental*
Psychology: Human Perception and Performance, *42*(9), 1297–1310. https://doi.org/10.1037/xhp0000207

28. Gernert, C. C., Falkai, P., & Falter-Wagner, C. M. (2025). *Heart rate synchrony as an objective biomarker for*
therapy outcome in affective disorders (p. 2024.08.29.24312787). medRxiv.
https://doi.org/10.1101/2024.08.29.24312787

29. Adel, L., Manova, V., Bain, T., Ambwani, D., Khoury, B., & Lifshitz, M. (2025). *Interpersonal cardiac synchrony*
during emotional sharing between strangers. PsyArXiv.

30. Konvalinka, I., Xygalatas, D., Bulbulia, J., Schjødt, U., Jegindø, E.-M., Wallot, S., Van Orden, G., & Roepstorff, A.
(2011). Synchronized arousal between performers and related spectators in a fire-walking ritual. *Proceedings of the*
National Academy of Sciences, *108*(20), 8514–8519. https://doi.org/10.1073/pnas.1016955108

31. Andersen, M. M., Rosas, F. E., Schjoedt, U., Mediano, P. A. M., Price, H., Konvalinka, I., & Clasen, M. (2025).
Scared together: Heart rate synchrony and social closeness in a high-intensity horror setting. *Emotion*.
https://doi.org/10.1037/emo0001556

32. Cristani, M., Paggetti, G., Vinciarelli, A., Bazzani, L., Menegaz, G., & Murino, V. (2011). Towards computational
proxemics: Inferring social relations from interpersonal distances. *2011 IEEE Third International Conference on*
Privacy, Security, Risk and Trust and 2011 IEEE Third International Conference on Social Computing, 290–297.
https://doi.org/10.1109/PASSAT/SocialCom.2011.32

33. Hebel, V., & Rentzsch, K. (2022). One, two, three, sit next to me: Personality and physical distance. *Personality and*
Individual Differences, *198*, 111798. https://doi.org/10.1016/j.paid.2022.111798

34. Lahnakoski, J. M., Forbes, P. A. G., McCall, C., & Schilbach, L. (2020). Unobtrusive tracking of interpersonal
orienting and distance predicts the subjective quality of social interactions. *Royal Society Open Science*, *7*(8),
191815. https://doi.org/10.1098/rsos.191815

35. Jackson, J. C., Jong, J., Bilkey, D., Whitehouse, H., Zollmann, S., McNaughton, C., & Halberstadt, J. (2018).
Synchrony and physiological arousal increase cohesion and cooperation in large naturalistic groups. *Scientific*
Reports, *8*(1), 127. https://doi.org/10.1038/s41598-017-18023-4

36. Joshi, V., Akiri, S., Taherzadeh, S., Williams, G., & Kleinsmith, A. (2025). Investigating differences in paramedic
trainees' multimodal interaction during low and high physiological synchrony. *Proceedings of the 27th International*
Conference on Multimodal Interaction, 526–534. https://doi.org/10.1145/3716553.3750796

37. DePasquale, C. E. (2020). A systematic review of caregiver–child physiological synchrony across systems:
Associations with behavior and child functioning. *Development and Psychopathology*, *32*(5), 1754–1777.
https://doi.org/10.1017/S0954579420001236

38. Pauly, T., Gerstorf, D., Wahl, H.-W., & Hoppmann, C. A. (2021). A developmental–contextual model of couple
synchrony across adulthood and old age. *Psychology and Aging*, *36*(8), 943–956.
https://doi.org/10.1037/pag0000651

39. Van Marum, R. J. (2020). Underrepresentation of the elderly in clinical trials, time for action. *British Journal of*
Clinical Pharmacology, *86*(10), 2014–2016. https://doi.org/10.1111/bcp.14539

40. United Nations. (2024). *World population prospects 2024: Summary of results*. United Nations.

41. De Jong Gierveld, J., & van Tilburg, T. (2006). A 6-item scale for overall, emotional, and social loneliness:
Confirmatory tests on survey data. *Research on Aging*, *28*(5), 582–598. https://doi.org/10.1177/0164027506289723

42. Pittinsky, T. L., Rosenthal, S. A., & Montoya, R. M. (2011). Measuring positive attitudes toward outgroups:
Development and validation of the Allophilia Scale. In L. R. Tropp & R. K. Mallett (Eds), *Moving beyond prejudice*

reduction: Pathways to positive intergroup relations. (pp. 41–60). American Psychological Association.
https://doi.org/10.1037/12319-002

43. Paulus, C. (2009). *Der Saarbrücker Persönlichkeitsfrageboden SPF (IRI) zur Messung von Empathie:*
Psychometrische Evaluation der deutschen Version des Interpersonal Reactivity Index.
https://doi.org/10.23668/psycharchives.9249

44. Aron, A., Aron, E. N., & Smollan, D. (1992). Inclusion of other in the self scale and the structure of interpersonal
closeness. *Journal of Personality and Social Psychology*, *63*(4), 17. https://doi.org/10.1037/0022-3514.63.4.596

45. Kenny, D. A., Kashy, D. A., & Cook, W. A. (2006). Using multilevel modeling to study dyads. In *Dyadic Data*
Analysis, (pp. 78–99). Guilford Publications. https://ebookcentral.proquest.com/lib/ethz/detail.action?docID=362550

46. Cao, Z., Hidalgo, G., Simon, T., Wei, S.-E., & Sheikh, Y. (2021). Openpose: Realtime multi-person 2d pose
estimation using part affinity fields. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, *43*(1), 172–
186. https://doi.org/10.1109/TPAMI.2019.2929257

47. Scholkmann, F., Boss, J., & Wolf, M. (2012). An efficient algorithm for automatic peak detection in noisy periodic
and quasi-periodic signals. *Algorithms*, *5*(4), 588–603. https://doi.org/10.3390/a5040588

48. Perdue, K. L., Westerlund, A., McCormick, S. A., & Nelson, C. A. (2014). Extraction of heart rate from functional
near-infrared spectroscopy in infants. *Journal of Biomedical Optics*, *19*(6), 067010.
https://doi.org/10.1117/1.JBO.19.6.067010

49. Sun, J., Rao, L., & Gao, C. (2018). Extracting heartrate from optical signal of functional near-infrared spectroscopy
based on mathematical morphology. *Journal of Innovative Optical Health Sciences*, *11*(03), 1850010.
https://doi.org/10.1142/S1793545818500104

50. Cerina, L. (2023). *Automatic Multiscale-based Peak Detection (AMPD)* [Computer software].

51. Feldman, R., Magori-Cohen, R., Galili, G., Singer, M., & Louzoun, Y. (2011). Mother and infant coordinate heart
rhythms through episodes of interaction synchrony. *Infant Behavior and Development*, *34*(4), 569–577.
https://doi.org/10.1016/j.infbeh.2011.06.008

52. McElreath, R. (2020). *Statistical Rethinking; A Bayesian Course with Examples in R and Stan* (2nd Edition).
Chapman and Hall/CRC.

53. Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical*
Software, *80*(1), 1–28. https://doi.org/10.18637/jss.v080.i01

54. R Core Team. (2022). *R: A language and environment for statistical computing* [Computer software]. R Foundation
for Statistical Computing. http://www.r-project.org/

55. RStudio Team. (2020). *RStudio: Integrated Development for R* [Computer software]. RStudio, Inc.
http://www.rstudio.com/

56. Kruschke, J. K., & Liddell, T. M. (2018). The Bayesian New Statistics: Hypothesis testing, estimation, meta-
analysis, and power analysis from a Bayesian perspective. *Psychonomic Bulletin & Review*, *25*(1), 178–206.
https://doi.org/10.3758/s13423-016-1221-4

57. Ford, R. M., & Aberdein, B. (2015). Exploring social influences on the joint Simon task: Empathy and friendship.
Frontiers in Psychology, *6*. https://doi.org/10.3389/fpsyg.2015.00962

58. Shafaei, R., Bahmani, Z., Bahrami, B., & Vaziri-Pashkam, M. (2020). Effect of perceived interpersonal closeness on
the joint Simon effect in adolescents and adults. *Scientific Reports*, *10*(1), 18107. https://doi.org/10.1038/s41598-
020-74859-3

59. Wolf, W., Launay, J., & Dunbar, R. I. M. (2016). Joint attention, shared goals, and social bonding. *British Journal of*
Psychology, *107*(2), 322–337. https://doi.org/10.1111/bjop.12144

60. Strang, A. J., Funke, G. J., Russell, S. M., Dukes, A. W., & Middendorf, M. S. (2014). Physio-behavioral coupling in
a cooperative team task: Contributors and relations. *Journal of Experimental Psychology: Human Perception and*
Performance, *40*(1), 145–158. https://doi.org/10.1037/a0033125

61. Lieberz, J., Shamay-Tsoory, S. G., Saporta, N., Esser, T., Kuskova, E., Stoffel-Wagner, B., Hurlemann, R., &
Scheele, D. (2021). Loneliness and the social brain: How perceived social isolation impairs human interactions.
Advanced Science, *8*(21), 2102076. https://doi.org/10.1002/advs.202102076

62. Saporta, N., Scheele, D., Lieberz, J., Stuhr-Wulff, F., Hurlemann, R., & Shamay-Tsoory, S. G. (2021). Opposing
association of situational and chronic loneliness with interpersonal distance. *Brain Sciences*, *11*(9), 1135.
https://doi.org/10.3390/brainsci11091135

63. Scheele, D., Striepens, N., Güntürkün, O., Deutschländer, S., Maier, W., Kendrick, K. M., & Hurlemann, R. (2012).
Oxytocin modulates social distance between males and females. *The Journal of Neuroscience*, *32*(46), 16074–16079.
https://doi.org/10.1523/JNEUROSCI.2755-12.2012

64. Shamay-Tsoory, S. G., & Kanterman, A. (2024). Away from the herd: Loneliness as a dysfunction of social
alignment. *Social Cognitive and Affective Neuroscience*, *19*(1), nsae005. https://doi.org/10.1093/scan/nsae005

65. Flory, S., Guglielmini, S., Scholkmann, F., Marcar, V. L., & Wolf, M. (2023). How our hearts beat together: A study
on physiological synchronization based on a self-paced joint motor task. *Scientific Reports*, *13*(1), 11987.
https://doi.org/10.1038/s41598-023-39083-9

66. Behrens, F., Snijdewint, J. A., Moulder, R. G., Prochazkova, E., Sjak-Shie, E. E., Boker, S. M., & Kret, M. E.
(2020). Physiological synchrony is associated with cooperative success in real-life interactions. *Scientific Reports*,
10(1), 19609. https://doi.org/10.1038/s41598-020-76539-8

67. Fendrich, S. J., Balachandran, M., & Patel, M. S. (2021). Association between behavioral phenotypes and sustained
use of smartphones and wearable devices to remotely monitor physical activity. *Scientific Reports*, *11*(1), 21501.
https://doi.org/10.1038/s41598-021-01021-y

68. Jiang, L., Gao, B., Gu, J., Chen, Y., Gao, Z., Ma, X., Kendrick, K. M., & Woo, W. L. (2019). Wearable long-term
social sensing for mental wellbeing. *IEEE Sensors Journal*, *19*(19), 8532–8542.
https://doi.org/10.1109/JSEN.2018.2877427

69. Moffat, R., Casale, C. E., & Cross, E. S. (2024). Mobile fNIRS for exploring inter-brain synchrony across
generations and time. *Frontiers in Neuroergonomics*, *4*. https://doi.org/10.3389/fnrgo.2023.1260738

70. UNICEF. (2015). *Innovating for Impact*. Innovating for Impact Wearable Technology Opens New Channels for
Connecting the Hardest to Reach to Services. https://www.unicef.org/innovation/stories/innovating-impact

71. Barclay, P. (2012). Harnessing the power of reputation: Strengths and limits for promoting cooperative behaviors.
Evolutionary Psychology, *10*(5), 868–883. https://doi.org/10.1177/147470491201000509

72. Cavazza, N., Guidetti, M., & Pagliaro, S. (2015). Who cares for reputation? Individual differences and concern for
reputation. *Current Psychology*, *34*(1), 164–176. https://doi.org/10.1007/s12144-014-9249-y

73. Clark, H. H., & Marshall, C. R. (1981). Definite reference and mutual knowledge. In A. K. Joshi, B. L. Webber, & I.
A. Sag (Eds), *Elements of Discourse Understanding* (pp. 10–63). Cambridge University Press.

74. Speer, S. P. H., Mwilambwe-Tshilobo, L., Tsoi, L., Burns, S. M., Falk, E. B., & Tamir, D. I. (2024). Hyperscanning
shows friends explore and strangers converge in conversation. *Nature Communications*, *15*(1), 7781.
https://doi.org/10.1038/s41467-024-51990-7

75. Helm, J. L., Miller, J. G., Kahle, S., Troxel, N. R., & Hastings, P. D. (2018). On measuring and modeling
physiological synchrony in dyads. *Multivariate Behavioral Research*, *53*(4), 521–543.
https://doi.org/10.1080/00273171.2018.1459292

76. Jones, I. (2024). *Neural, physiological and cognitive mechanisms of being watched* [Doctoral thesis]. University
College London.

77. Young, H., & Benton, D. (2015). We should be using nonlinear indices when relating heart-rate dynamics to
cognition and mood. *Scientific Reports*, *5*(1), 16619. https://doi.org/10.1038/srep16619

78. Stuldreher, I. V., Van Erp, J. B. F., & Brouwer, A.-M. (2023). Robustness of physiological synchrony in wearable
electrodermal activity and heart rate as a measure of attentional engagement to movie clips. *Sensors*, *23*(6), 3006.
https://doi.org/10.3390/s23063006