Interrogating the structure and function of the human voltage-gated proton channel (hH$_v$1) with a fluorescent noncanonical amino acid.

Emerson M. Carmona, William N. Zagotta*, Sharona E. Gordon*.

Department of Neurobiology and Biophysics, University of Washington, Seattle, WA, U.S.

*William N. Zagotta and Sharona E. Gordon.

Email: zagotta@uw.edu, seg@uw.edu

Author Contributions: All the authors conceived the study and designed the experiments. E.M.C. performed the experiments and analyzed the data. All the authors interpreted the data. E.M.C. wrote the first version of the manuscript. All the authors critically revised the article.

Competing Interest Statement: The authors declare no competing interests.

Abstract

The human voltage-gated proton channel (hH$_v$1) is a dimer of voltage-
sensor domains (VSDs) containing highly selective proton permeation pathways
in each monomer. In addition to voltage, hH$_v$1 is regulated by other stimuli,
including pH gradients, mechanical forces, and ligands such as Zn^{2+}. Aside from
the VSDs, this membrane protein contains an N-terminal domain and a C-
terminal coiled-coil domain (CC) formed between the monomers. To address the
need for direct measurements of conformational rearrangements in hH$_v$1, we
developed a Förster resonance energy transfer (FRET) approach to measuring
the conformational rearrangements in full-length hH$_v$1 purified from *E. coli*. We
used genetic code expansion (GCE) to generate a library of 14 full-length hH$_v$1
constructs, each incorporating the fluorescent noncanonical amino acid acridon-
2-ylalanine (Acd) at a different site throughout the various structural domains.
Following the expression and purification of these hH$_v$1-Acd proteins, we found
that 12 sites yielded stable and functional proton-permeable channels. The
fluorescence properties of Acd at each site showed small site-specific
differences. Furthermore, we measured site-specific FRET efficiencies from
tryptophan (Trp) and tyrosine (Tyr) to Acd in the hH$_v$1-Acd proteins and found
results consistent with correct folding in detergent micelles. Finally, the addition
of Zn^{2+} produced reversible changes in FRET, with affected residues clustered
on the intracellular side of the channel.

Introduction

The human voltage-gated proton channel (hH$_v$1) is a membrane protein that forms a proton selective channel (1, 2). Despite its physiological importance, its molecular mechanisms are still poorly understood, including the precise structural identity of the permeation pathway and the gating mechanism of voltage (3, 4), pH gradients (3, 5), membrane stretch (6), and ligands such as the classical inhibitor Zn^{2+} (7, 8). Understanding these processes requires connecting the functional properties to the hH$_v$1 structure.

Each monomer of hH$_v$1 dimers includes an intracellular N-terminal domain of unknown function, a transmembrane voltage-sensor domain (VSD), and an intracellular coiled-coil (CC), which forms the primary intersubunit interface (9, 10). Current structural models, however, have been obtained from truncated or chimeric channel constructs (11, 12), and thus, it is not clear whether they correspond to physiological or functional conformations. Moreover, the multiple kinetic components of H$_v$1 gating currents suggest that the protein's resting state comprises an ensemble of conformations (4, 5), which is consistent with the conformational flexibility inferred from structural approaches (12–14). Because protein function arises from this distribution of conformational states (15), approaches to directly measure structural heterogeneity in the full-length hH$_v$1 are needed.

An attractive approach to interrogate the structural heterogeneity of hH$_v$1 is fluorescence spectroscopy, which offers high sensitivity at nanomolar protein concentrations and compatibility with near-physiological conditions. Fluorescence can report on local environment and accessibility changes, as well as quantify distance changes through Förster resonance energy transfer (FRET) (16). Importantly, time-resolved FRET measured using lifetimes preserves information about conformational heterogeneity on the nanosecond timescale, which is lost in steady-state measurements due to the averaging process (17). Together, these capabilities make fluorescence, especially time-resolved FRET, a powerful method to connect hH$_v$1 structure and function (18).

A few obstacles remain to directly measuring conformational

heterogeneity of hH_v1. First, a method to express and purify stable and functional

protein is needed, which is challenging for membrane proteins. Second, site-

specific fluorophore labeling is difficult at buried sites, resulting in low labeling

efficiency. Third, traditional fluorophores are bulky and have long linkers (14),

resulting in protein structure perturbation and heterogeneity dominated by the

properties of the label (19). A recent method was optimized for robust expression

and purification of full-length, functional hH_v1 in *E. coli* (20, 21). The remaining

obstacles can be overcome by labeling the protein with the fluorescent

noncanonical amino acid acridon-2-ylalanine (Acd) using genetic code expansion

(GCE) (22–24). This technology allows site-specific labeling during translation by

reassigning an amber stop codon using an orthogonal aminoacyl-tRNA

synthetase (RS)/tRNA (25, 26) (Figure 1A). Acd's small size and minimal linker

reduce perturbations and heterogeneity, and its favorable photophysics support

both steady-state and time-resolved measurements (27). Here, we incorporated

Acd across all hH_v1 structural domains and purified 12 of 14 constructs as stable,

functional proteins. We then used Acd's environmental sensitivity and FRET from

tryptophan (Trp)/tyrosine (Tyr) to Acd to validate the protein folding and report

Zn^{2+}-dependent conformational changes. These data establish a platform for

interrogating conformational heterogeneity and dynamics in the full-length hH_v1.

Results

We generated a library of hH_v1 constructs with Acd incorporated across the

different structural domains. Because available H_v1 structural models were

produced from either truncated or modified proteins (11, 12), we used an

AlphaFold model of the full-length, dimeric hH_v1 to guide Acd site selection (28,

29) (Figure 1B and Dataset S1). The model recapitulates the expected hH_v1

architecture with high confidence: S0 as a short helix parallel to the membrane,

S1-S4 transmembrane segments forming a canonical VSD, and a C-terminal

helix extending from S4 that forms the CC intersubunit interface. In addition, two

helices in the N-terminal domain (P1 and P2) were predicted (Figure 1C), albeit

with low confidence. The dimer interface in the membrane is primarily formed by

S4-S4 contacts (Figure S1A), although S1-S1 contacts have been observed

experimentally (9, 13). This model appears to represent an intermediate state,

with the first S4 positive charged residue R208 proximal to D112 in the selectivity

filter, and the third S4 positive charged residue R211 proximal to F150 in the

charge-transfer center (Figure S1B). Although the model's accuracy and

functional state are unknown, we used it as a rough starting point for selecting

positions to incorporate Acd.

To test the feasibility and specificity of Acd incorporation in hH_v1, we

initially selected two positions in the intracellular domains: A18 in the N-terminal

domain and F247 in the CC. For each position, the wild-type codon was replaced

with an amber stop codon (TAG). Bacteria co-transformed with the Acd RS/tRNA

pair and hH_v1-TAG constructs expressed fluorescent full-length protein only

when Acd was present in the culture medium (Figure 1D). We next generated a

library of 14 constructs in which Acd was incorporated into every structural

domain: the N-terminal domain, each helix of the VSD, and the CC (Figure 2A-

B). Remarkably, most of the 14 hH_v1-Acd proteins expressed at levels

comparable to the control (No TAG) (Figure 2C and S2). As expected, several

hH_v1-Acd proteins also produced TAG-truncated products (Figure 2C).

Increasing the Acd concentration in the *E. coli* growth medium did not reduce the

fraction of truncated protein (Figure S3). One of the constructs, hH_v1-C107Acd,

showed a weak in-gel fluorescence signal, although the Western blot signal was

comparable to that of other constructs.

We successfully purified stable and functional hH_v1 labeled with Acd at 12

of the 14 selected positions. We attempted to purify all expressed hH_v1-Acd

constructs by immobilized metal affinity chromatography using the detergent

Anzergent 3-12 (20, 21). For 12 of 14 constructs, purification yielded a single

fluorescent band on SDS-PAGE (Figure 3A and S4). In contrast, no purified

protein was detected for the V187Acd or Q233Acd constructs (Figure S4).

Because hH$_v$1-Q233Acd was robustly expressed (Figure 2C), its loss during

purification suggests reduced stability. The hH$_v$1-Acd yields were construct-

dependent (Figure 3B), and the TAG-truncated products did not co-purify with the

full-length protein (Figure 3A and S4).

We then evaluated the stability and homogeneity of the hH$_v$1-Acd proteins

using fluorescence-detection size exclusion chromatography (FSEC) (30).

Similar to the control (Figure S5), most hH$_v$1-Acd proteins showed a main peak

(~11.5 mL) followed by a minor peak at higher elution volumes (Figure 3C and

S6; Table S1). The proportion of these two peaks varied between constructs

(Figure S6). To determine whether the purified hH$_v$1-Acd proteins were

functional, we reconstituted the FSEC fractions containing both peaks in

asolectin liposomes and assessed their function using a liposome proton flux

assay (20, 31). All 12 reconstituted constructs produced ACMA fluorescence

quenching upon valinomycin addition, indicating that our purified proteins formed

functional proton-permeable channels (Figure 3D and S7). These results

highlight the versatility of GCE for incorporating Acd as a fluorescent probe for

structural studies of hH$_v$1.

Acd was relatively insensitive to the hydrophobicity of its local

environment. To explore how the local environment can alter Acd's spectral

properties, we measured the emission spectra of the free amino acid in different

solvents (Figure 4A and Table S2). The environmental sensitivity of Acd has

been studied previously (24, 27, 32), but we expanded this work here by using

the aprotic solvent ethyl acetate (EtAc) and a series of alcohols with varying alkyl

chain lengths. Consistent with a general solvent effect produced by polarity (16),

the emission spectrum was blue-shifted by approximately 20 nm in EtAc, the

least polar solvent measured, relative to water (Figure 4A and Table S2).

Alcohols produced a consistently smaller blue shift, independent of their alkyl-

chain length (Figure 4A). The spectrum in Buffer-H2, our standard experimental

solution containing detergent micelles, was only very slightly blue-shifted relative

to water (Figure 4A), suggesting that Acd did not substantially partition into the

hydrophobic core of the detergent micelles. We also tested whether solvent polarity affects the fluorescence lifetime of Acd using time-correlated single photon counting (TCSPC). The fluorescence lifetime of Acd showed a similar trend to the spectral shifts (Figure 4B): EtAc shortened the lifetime markedly, Acd exhibited shorter lifetimes in alcohols than in water, and Acd in Buffer-H2 had a lifetime similar to that measured in water.

The spectral blue-shift and short lifetime of Acd in EtAc suggest that Acd may be able to report on the local environment at different sites in hH_v1, especially those that are surface-exposed versus those that face the lipid core. The emission spectra of the hH_v1-Acd proteins showed small, site-specific shifts (Figure 4C and Table S3). The local environment of Acd in hH_v1 spanned the range between those observed for free Acd in Buffer-H2 and in alcohols, with Q56Acd in the N-terminal intracellular domain and C107Acd in S1 showing the most red-shifted and blue-shifted spectra, respectively (Figure 4C). Consistent with the spectra, the fluorescence lifetimes of the hH_v1-Acd proteins fell between those measured for free Acd in Buffer-H2 and in alcohols, with Q56Acd exhibiting the slowest, and C107Acd the fastest, lifetimes (Figure 4D). While small, the differences in spectral properties of Acd in hH_v1 agree with their predicted location in the structural model, with C107 being the site closest to the detergent micelles' hydrophobic core. The low sensitivity to polarity changes of Acd in different hH_v1 local environments makes it well suited for measuring FRET (27, 33).

FRET between Trp/Tyr and Acd in hH_v1 suggested that the proteins are properly folded. Trp and Tyr have fluorescence emission spectra that overlap with the Acd absorption spectrum and therefore can act as FRET donors to Acd, with relatively short R_0 values (23.5 Å for Trp; 20.9 Å for Tyr) (22) (Figure 5A). Each hH_v1 monomer contains four Trp and four Tyr residues (yellow spheres in Figure 5B). In this structural model, the Trp residues are located in the N-terminal domain (W4, W38, and W45) and midway along transmembrane segment S4 in the VSD (W207). The Tyr residues are located in the N-terminal domain (Y35

and Y42) and near the extracellular end of transmembrane segment S2 (Y134

and Y141).

To quantify Trp/Tyr to Acd FRET, we used spectral FRET analysis (34).

The total emission spectrum from Trp/Tyr and Acd was collected (Figure 5C).

The Acd emission spectrum was extracted by subtracting a scaled Trp/Tyr

spectrum collected from control wild-type protein ($F_{280,NoTAG}$) (Figure 5D and S8).

For each Acd-containing construct, the ratio of the extracted spectrum ($F_{280} -$

$F_{280,NoTAG}$) to the Acd spectrum with direct excitation (F_{370}) was calculated as

Ratio A (Figure 5D and 5E). Because *Ratio A* is not wavelength-dependent, it

reports the linearity of the detectors and the absence of significant contamination

from other sources of fluorescence. The *Ratio A* component caused by the direct

excitation of Acd (termed *Ratio A$_0$*) was measured with free Acd (35) (Figure 5E

and S9). The difference between *Ratio A* and *Ratio A$_0$* is directly proportional to

the FRET efficiency.

As expected, the *Ratio A* values for our hH$_v$1-Acd proteins were largely flat

across emission wavelengths for our hH$_v$1-Acd proteins (Figure 5E). For all Acd

sites, *Ratio A* was greater than *Ratio A$_0$* (dashed line in Figure 5E), reflecting

FRET between Trp/Tyr and Acd at every site in hH$_v$1. Most *Ratio A* values were

distributed between 0.60 and 0.80, with two clear exceptions: A18Acd (blue) and

F247Acd (magenta) (Figure 5E). A18Acd showed the highest *Ratio A*, consistent

with the multiple Trp/Tyr residues in the N-terminal domain. Conversely,

F247Acd showed the lowest *Ratio A*, consistent with its location in the CC, far

from most Trp/Tyr residues (Figure 5B).

We next compared *Ratio A – Ratio A$_0$* values with FRET efficiencies

predicted by the AlphaFold structural model. We modeled rotamer ensembles for

Trp, Tyr, and our Acd sites in the hH$_v$1 structural model using chiLife (36) and

calculated donor-acceptor distance distributions. These distributions were then

used to calculate FRET efficiencies (details in SI Text). The experimentally

determined *Ratio A – Ratio A$_0$* values showed a moderate correlation (Pearson's

r = 0.48) with the calculated efficiencies (Figure 5F).

There were two spatially clustered deviations between experiment and
model. Positions at the extracellular ends of the VSD helices (Y134 in S2, K125
in S1, and F195 in S4) showed higher experimental FRET than predicted,
indicating shorter effective donor-acceptor distances relative to the model.
Conversely, positions on the intracellular side of the VSD (K169 in S3 and I217 in
S4) showed lower experimental FRET than predicted, implying longer effective
donor-acceptor distances. Together, the spatial clustering of residues with similar
deviations indicates specific deficits in the AlphaFold model or the failure to
quantify the protein dynamics, which can guide future improvements.
231
Zinc binding changed the conformation of hH$_v$1. We next asked whether
FRET from Trp/Tyr to Acd could report conformational changes upon addition of
Zn^{2+}, the classical inhibitor of H$_v$1 proton currents in cells (7). The two
experimental structures of H$_v$1 were resolved with this cation bound to the
extracellular side of the VSD (11, 12). We observed site-specific spectral
changes in the hH$_v$1-Acd proteins in the presence of Zn^{2+} (Figure 6 and S10).
The largest changes occurred for hH$_v$1-K169Acd, where 1 mM Zn^{2+} caused a
decrease in the Trp/Tyr emission and an increase in the Acd emission (Figure
6A). This decrease in donor fluorescence, paired with increased acceptor
fluorescence, is the hallmark of increased FRET and was corroborated by
spectral analysis (Figure 6B). Additionally, a slight blue shift of the Trp/Tyr
fluorescence was observed for all hH$_v$1-Acd proteins in the presence of Zn^{2+}
(Figure S11). These changes were completely reversed by the addition of EDTA,
confirming that the changes arise from a reversible conformational change upon
Zn^{2+} binding (Figure 6A and 6B). Large changes in effective FRET efficiency
were observed with Acd incorporated at Q56 in the N-terminal domain and C107,
F159, and K169 in the VSD (Figure 6C). Notably, these positions are
intracellular, even though H$_v$1 currents are inhibited by extracellular Zn^{2+} (7, 37).
These data demonstrate that Zn^{2+} binds to hH$_v$1 and alters its conformation in
detergent micelles, consistent with a nuclear magnetic resonance study of the
VSD of hH$_v$1 (12).

Discussion

Incorporating Acd into hH$_v$1 by GCE opens new possibilities for studying

the structure and function of this channel. We systematically incorporated Acd at

multiple positions in hH$_v$1 and confirmed protein stability and function after

purification. 12 of 14 hH$_v$1-Acd proteins produced acceptable yields upon

purification and were functional, despite being a human membrane protein

expressed in *E. coli*. Two factors likely underlie the robustness of our system: (i)

an optimized purification protocol for hH$_v$1 (20, 21) that offsets the expected

reduction in yield associated with GCE (22, 25), and (ii) the high efficiency and

specificity of the evolved aminoacyl-tRNA synthetase for incorporating Acd (17,

27).

Fluorescence spectroscopy has been used to detect conformational

changes in H$_v$1. Sites at the extracellular portion of S4 (38) and S1 (39) in the

VSD of *Ciona intestinalis* H$_v$1 have been labeled with the fluorophores Alexa488

and tetramethylrhodamine (TAMRA) through cysteine-selective reagents to

capture conformational changes during activation using voltage-clamp

fluorometry. Similarly, the fluorescent noncanonical amino acid 3-(6-

acetylnaphthalen-2-ylamino)-2-aminopropanoic acid (Anap) has been

incorporated at several sites in the S4 in hH$_v$1 (40) to study conformational

changes in response to voltage and pH. In these cases, the fluorophore's

environmental sensitivity has been used as a readout of conformational changes.

By contrast, FRET provides richer structural information due to its distance

dependence. Single-molecule FRET has been measured in purified hH$_v$1 labeled

with Cy3 and Cy5 to track the VSD movements in response to voltage, pH,

cholesterol, and zinc (14, 41). The small size and minimal linker of Acd, along

with its low environmental sensitivity and high photostability, are advantageous

for measuring FRET in hH$_v$1 compared with previously used fluorophores.

The high number of FRET donors, 4 tryptophans and 4 tyrosines per

subunit, limits the utility of the Trp/Tyr-Acd FRET pair as a strategy to measure

conformational rearrangements in hH$_v$1. Our library of a dozen single Acd-

incorporating constructs, with sites distributed across the entire primary

sequence of the protein, opens the door to using Acd as a FRET donor. We have

previously used time-resolved transition metal ion FRET (tmFRET), which utilizes

a transition metal ion as a FRET acceptor (27, 33), to measure conformational

energetics in the bacterial cyclic nucleotide-gated ion channel SthK (42).

However, in that work, we incorporated Acd at a single position in the intracellular

C-terminal domain of SthK. With Acd incorporated at multiple sites in full-length

hH$_v$1, we can interrogate its gating mechanisms in unprecedented detail.

Materials and Methods

Protein expression and purification. hH$_v$1 was expressed and purified

according to the published protocol with minor modifications (20, 21). Briefly,

fresh competent BL21-Gold(DE3) cells (Agilent) were co-transformed with the

His-EK-hH$_v$1-C107A-C249A.pET15-b (hH$_v$1-Cysless, referred to as No TAG) and

MjA9Acd-RS.pDule2 (22, 23) plasmids (sequences in SI Text) using the heat

shock method. For expressing hH$_v$1-Acd proteins, the codon corresponding to

the selected position was changed to the amber stop codon (TAG) by site-

directed mutagenesis. All plasmids were sequenced before use. Pre-cultures

were grown overnight in LB medium supplemented with 0.2% glucose, 0.4

mg/mL ampicillin, and 0.1 mg/mL spectinomycin at 37 °C and 250 rpm. The next

day, cultures were diluted 1:100 in complete autoinduction medium (43)

supplemented with 0.4 mg/mL ampicillin and 0.1 mg/mL spectinomycin, and then

grown at 37 °C and 250 rpm until an OD$_{600}$ of 1.5 was reached. The protein

expression level was proportional to the Acd concentration in the culture medium

(Figure S11). We chose 0.6 mM Acd in the culture medium as optimal for hH$_v$1

expression, as higher concentrations only slightly increased the final biomass

(Figure S11B). Cultures supplemented with 0.6 mM Acd were grown overnight at

20 °C and 250 rpm. Cells were then harvested, resuspended in Buffer-H1 (50

mM Tris, 150 mM NaCl, 1 mM benzamidine, 0.17 mg/mL PMSF, pH 8.0)

supplemented with 0.5 mg/mL lysozyme, and incubated for 30 min at 4 °C before

being flash-frozen. After thawing, the extracts were supplemented with 5 mM

$MgCl_2$, fresh protease inhibitors, and 12.5 μg/mL DNase I, incubated for 1 h at 4

°C, sonicated, and centrifuged at 100,000g at 4 °C for 1 h. The membrane pellet

was resuspended in Buffer-H1 and stored at -80 °C until further use. After

thawing, the membranes were solubilized with 1.5% Anzergent 3-12 (Anatrace)

(Anz3-12) at room temperature for 1 hour. Insoluble material was removed by

centrifuging at 100,000g and 4 °C for 1 h, and the solubilized membrane extract

was incubated with His-Pur Ni-NTA (Thermo Scientific) resin equilibrated with

Buffer-H2 (50 mM Tris, 150 mM NaCl, 12 mM Anz3-12, pH 8.0) for 1 hour at

room temperature. Resin was collected in a gravity column and washed with 20

column volumes (CV) of Buffer-H2, followed by 16 CV of Buffer-H2 with 90 mM

imidazole. The hH_v1 protein was eluted with 20 CV of Buffer-H2 with 0.5 M

imidazole, was concentrated with 50kDa cut-off centrifugal filters, and imidazole

was removed by FSEC (Ex/Em = 385/450 nm) in an ENrich SEC 650 (Bio-Rad)

column with Buffer-H2. The hH_v1-containing fractions were concentrated to 0.5-

1.5 mg/mL, depending on the amount of purified protein, and then aliquoted,

flash-frozen, and stored at -80 °C until use.

Reconstitution and fluorescence proton flux assays. The hH_v1-Acd samples

were reconstituted and assayed using a liposome proton flux assay, as described

previously, with minor modifications (20). Briefly, soy polar lipid extract (Avati) in

chloroform was dried overnight at room temperature in a rotary evaporator under

vacuum and then hydrated in Buffer-K (20 mM HEPES, 150 mM KCl, and 1 mM

EDTA, pH 7.0) to a concentration of 10 mg/mL. The vesicles were sonicated in a

water bath until the solution became translucent, aliquoted, flash-frozen, and

stored at -80 °C until further use. Reconstitutions were performed at a 1:100

(protein:lipid, by mass) ratio by mixing 12.5 μg of purified hH_v1-Acd with 1.25 mg

of liposomes in a final volume of 330 μL Buffer-K containing 8 mM Anz3-12. The

mixture was incubated at room temperature for 1 h, then diluted with 5 mL of

Buffer-K and incubated for 30 min. The mixture was further incubated with three

cumulative pulses of 100 mg Bio-Beads SM2 (Bio-Rad), each for 1 h at room

temperature. After the final Bio-Beads pulse, the mixture was incubated overnight

at 4 °C. The next day, two successive 100 mg additions of Bio-Beads were

performed, with a 1-hour incubation at room temperature following each addition.

Bio-Beads were removed in a gravity column, the mixture was diluted with 10 mL

of Buffer-K, and then centrifuged at 150,000g for 2 hours at 4 °C. The

proteoliposome pellets were resuspended in 250 μL of Buffer-K, aliquoted, flash-

frozen, and stored at -80 °C until further use. On the day of the liposome proton

flux assays, samples were thawed at 37 °C and then placed on ice. The sample

(40 μL) was diluted in Buffer-Na (20 mM HEPES, 150 mM NaCl, 1 mM EDTA, pH

7.0). 9-Amino-6-chloro-2-methoxyacridine (ACMA; Sigma) was added to a final

concentration of 2 μM from a 2 mM stock solution in DMSO, and the mixture was

incubated at room temperature for 5 minutes before the fluorescence

measurement began. The baseline was recorded for 3 min (F_{max}). Subsequently,

10 nM of valinomycin (Cell Signaling) was added from a 10 μM stock solution in

DMSO, followed by 1 μM carbonyl cyanide m-chlorophenyl hydrazone (CCCP;

Sigma) from a 1 mM stock solution in DMSO to record F_{min}. The final volume was

2 mL. The fluorescence signal ($F(t)$) was normalized according to the equation

($F(t)$-F_{min})/(F_{max}-F_{min}). Fluorescence measurements were performed at 5-second

intervals with a 2-second integration time using a Fluorolog-3 spectrometer

(Horiba) configured to an excitation wavelength of 410 nm (5 nm bandpass) and

an emission wavelength of 490 nm (1 nm bandpass) at room temperature.

Absorption and fluorescence spectra recordings. The absorption and

fluorescence spectra were recorded in the indicated solvents or fresh Buffer-H2

at room temperature. Absorption spectra were recorded in a DU 800

Spectrometer (Beckman Coulter) with a wavelength interval of 0.5 nm and a scan

speed of 600 nm/min. Fluorescence emission spectra were measured using a

Fluorolog-3 spectrometer (Horiba) with an integration time of 0.1 s, 1 nm

increments, and excitation and emission slits of 5 nm. For solubilizing Acd in

different solvents, a saturated solution of the amino acid was prepared by

dissolving 0.5 mg of Acd powder in the respective solvent. Acd spectra were

corrected by subtracting a blank of the solvent. hH_v1-Acd proteins were

measured at concentrations ranging from 180 to 240 nM in Buffer-H2. Each

spectrum was corrected by subtracting a blank with the solvent or Buffer-H2

before adding the protein sample to the cuvette.

Spectral FRET analysis. We used our previously established spectral FRET

analysis to remove contamination caused by direct excitation of Acd by 280 nm

light (35). This method had the added benefit of eliminating errors arising from

the recording system's transfer function, variations in the acceptor's quantum

yield, or variations in the total concentration of fluorescent molecules. A Trp/Tyr

spectrum was collected from control protein without any TAG codons (no Acd

incorporation). This was used to subtract the Trp/Tyr emission spectra collected

at 280 nm for each Acd-incorporating protein. This yielded the extracted Acd

spectrum, F_{280}, that had two components: the component due to direct excitation

of Acd, F_{280}^{direct}, and the component due to FRET, F_{280}^{FRET}. F_{280} was normalized by

the total Acd emission with 370 nm excitation, F_{370}. The resulting ratio, termed

Ratio A, can be expressed as: $Ratio\ A = \frac{F_{280}}{F_{370}} = \frac{F_{280}^{direct}}{F_{370}} + \frac{F_{280}^{FRET}}{F_{370}}$. The direct

excitation component, $\frac{F_{280}^{direct}}{F_{370}}$, termed *Ratio A₀*, was measured using free Acd

amino acid in Buffer-H2. We then quantified the relative FRET efficiency as

$Ratio\ A - Ratio\ A_0 = \frac{F_{280}^{FRET}}{F_{370}}$. This quantity is directly proportional to FRET

efficiency. An average *Ratio A* value was calculated between 410 and 480 nm.

Time-resolved fluorescence measurements. TCSPC measurements were

performed in a FluoTime 300 spectrometer equipped with a PMA Hybrid 40

detector and LDH-P-C-375 laser diode head (PicoQuant) at room temperature.

Emitted photons were detected at 446 nm with the emission polarizer at the

magic angle. The TCSPC measurements were performed in the indicated solvent

or fresh Buffer-H2. The hH_v1-Acd samples were measured at a concentration of

1 μM and the Acd amino acid at 500 nM. The count rate was always less than

1% of the excitation repetition rate to avoid pile-up distortions of the fluorescence

decay. The IRF was measured using a dilute Ludox solution in water.

Acknowledgments

We thank the Oregon State University GCE4ALL (Center for Genetic Code

Expansion for All) for their long-standing collaboration. We also appreciate the

excellent technical support provided by Dr. James Petersson and Kyle D. Shaffer

(University of Pennsylvania) with the Acd synthesis. This work was supported by

the National Institutes of Health under award numbers R01EY037223 and

R35GM145225 to S.E.G., and R01EY010329 and R35GM148137 to W.N.Z.

E.M.C. is a Pew Latin American Fellow.

References

1. M. Sasaki, M. Takagi, Y. Okamura, A Voltage Sensor-Domain Protein Is a Voltage-
Gated Proton Channel. *Science* **312**, 589–592 (2006).

2. I. S. Ramsey, M. M. Moran, J. A. Chong, D. E. Clapham, A voltage-gated proton-
selective channel lacking the pore domain. *Nature* **440**, 1213–1216 (2006).

3. V. V. Cherny, V. S. Markin, T. E. DeCoursey, The voltage-activated hydrogen ion
conductance in rat alveolar epithelial cells is determined by the pH gradient. *The*
Journal of General Physiology **105**, 861–896 (1995).

4. E. M. Carmona, *et al.*, Gating charge displacement in a monomeric voltage-gated
proton (H_v1) channel. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 9240–9245 (2018).

5. E. M. Carmona, *et al.*, The voltage sensor is responsible for ΔpH dependence in H_v1
channels. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2025556118 (2021).

6. M. M. Pathak, *et al.*, The Hv1 proton channel responds to mechanical stimuli.
Journal of General Physiology **148**, 405–418 (2016).

7.   V. V. Cherny, T. E. DeCoursey, pH-Dependent Inhibition of Voltage-Gated H^+
Currents in Rat Alveolar Epithelial Cells by Zn^{2+} and Other Divalent Cations. *The*
Journal of General Physiology **114**, 819–838 (1999).

8.   T. E. DeCoursey, V. V. Cherny, Voltage-activated hydrogen ion currents. *J.*
Membarin Biol. **141** (1994).

9.   S.-Y. Lee, J. A. Letts, R. MacKinnon, Dimeric subunit stoichiometry of the human
voltage-dependent proton channel H_v1. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 7692–
7695 (2008).

10. H. P. Koch, *et al.*, Multimeric nature of voltage-gated proton channels. *Proc. Natl.*
Acad. Sci. U.S.A. **105**, 9111–9116 (2008).

11. K. Takeshita, *et al.*, X-ray crystal structure of voltage-gated proton channel. *Nat*
Struct Mol Biol **21**, 352–357 (2014).

12. M. Bayrhuber, *et al.*, Nuclear Magnetic Resonance Solution Structure and
Functional Behavior of the Human Proton Channel. *Biochemistry* **58**, 4017–4027
(2019).

13. Q. Li, *et al.*, Resting state of the human proton channel dimer in a lipid bilayer.
Proc. Natl. Acad. Sci. U.S.A. **112** (2015).

14. S. Han, *et al.*, Structural dynamics determine voltage and pH gating in human
voltage-gated proton channel. *eLife* **11**, e73093 (2022).

15. K. Henzler-Wildman, D. Kern, Dynamic personalities of proteins. *Nature* **450**, 964–
972 (2007).

16. J. R. Lakowicz, Ed., *Principles of Fluorescence Spectroscopy* (Springer US, 2006).

17. S. E. Gordon, M. Munari, W. N. Zagotta, Visualizing conformational dynamics of
proteins in solution and at the cell membrane. *eLife* **7**, e37248 (2018).

18. W. N. Zagotta, *et al.*, Measuring conformational equilibria in allosteric proteins
with time-resolved tmFRET. *Biophysical Journal* **123**, 2050–2062 (2024).

19. J. W. Taraska, M. C. Puljung, W. N. Zagotta, Short-distance probes for protein
backbone structure based on energy transfer between bimane and transition metal
ions. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 16227–16232 (2009).

20. E. M. Carmona, D. M. Cortes, L. G. Cuello, A novel method for expressing and
purifying large quantities of functional and stable human voltage-gated proton
channel (HH_v1). *Protein Science* **34**, e70017 (2025).

21.   E. Carmona, M. Cortes, L. Cuello, Expression and Purification of the Human
468   Voltage-Gated Proton Channel (hH$_v$1). *BIO-PROTOCOL* **15** (2025).

469   22.   L. C. Speight, *et al.*, Efficient Synthesis and In Vivo Incorporation of Acridon-2-
470   ylalanine, a Fluorescent Amino Acid for Lifetime and Förster Resonance Energy
471   Transfer/Luminescence Resonance Energy Transfer Studies. *J. Am. Chem. Soc.* **135**,
472   18806–18814 (2013).

473   23.   I. Sungwienwong, *et al.*, Improving target amino acid selectivity in a permissive
474   aminoacyl tRNA synthetase through counter-selection. *Org. Biomol. Chem.* **15**,
475   3603–3610 (2017).

476   24.   C. M. Jones, *et al.*, Genetic encoding of a highly photostable, long lifetime
477   fluorescent amino acid for imaging in mammalian cells. *Chem. Sci.* **12**, 11955–
478   11964 (2021).

479   25.   C. C. Liu, P. G. Schultz, Adding New Chemistries to the Genetic Code. *Annu. Rev.*
480   *Biochem.* **79**, 413–444 (2010).

481   26.   T. S. Young, I. Ahmad, J. A. Yin, P. G. Schultz, An Enhanced System for Unnatural
482   Amino Acid Mutagenesis in E. coli. *Journal of Molecular Biology* **395**, 361–374
483   (2010).

484   27.   W. N. Zagotta, *et al.*, An improved fluorescent noncanonical amino acid for
485   measuring conformational distributions using time-resolved transition metal ion
486   FRET. *eLife* **10**, e70236 (2021).

487   28.   J. Jumper, *et al.*, Highly accurate protein structure prediction with AlphaFold.
488   *Nature* **596**, 583–589 (2021).

489   29.   G. Kim, *et al.*, Easy and accurate protein structure prediction using ColabFold. *Nat*
490   *Protoc* **20**, 620–642 (2025).

491   30.   T. Kawate, E. Gouaux, Fluorescence-Detection Size-Exclusion Chromatography for
492   Precrystallization Screening of Integral Membrane Proteins. *Structure* **14**, 673–681
493   (2006).

494   31.   S.-Y. Lee, J. A. Letts, R. MacKinnon, Functional Reconstitution of Purified Human
495   H$_v$1 H$^+$ Channels. *Journal of Molecular Biology* **387**, 1055–1060 (2009).

496   32.   A. Szymańska, K. Wegner, L. Łankiewicz, Synthesis of
497   *N*-[(*tert*-Butoxy)carbonyl]-3-(9,10-dihydro-9-oxoacridin-2-yl)-L-alanine, a New
498   Fluorescent Amino Acid Derivative. *Helvetica Chimica Acta* **86**, 3326–3331 (2003).

499   33.   S. E. Gordon, *et al.*, Long-distance tmFRET using bipyridyl- and phenanthroline-
500   based ligands. *Biophysical Journal* **123**, 2063–2075 (2024).

501   34. R. M. Clegg, "[18] Fluorescence resonance energy transfer and nucleic acids" in
502   *Methods in Enzymology*, (Elsevier, 1992), pp. 353–388.

503   35. J. Zheng, M. D. Varnum, W. N. Zagotta, Disruption of an Intersubunit Interaction
504   Underlies Ca^{2+}-Calmodulin Modulation of Cyclic Nucleotide-Gated Channels. *J.*
505   *Neurosci.* **23**, 8167–8175 (2003).

506   36. M. H. Tessmer, S. Stoll, chiLife: An open-source Python package for in silico spin
507   labeling and integrative protein modeling. *PLoS Comput Biol* **19**, e1010834 (2023).

508   37. F. Qiu, *et al.*, Molecular mechanism of Zn^{2+} inhibition of a voltage-gated proton
509   channel. *Proc. Natl. Acad. Sci. U.S.A.* **113** (2016).

510   38. C. Gonzalez, H. P. Koch, B. M. Drum, H. P. Larsson, Strong cooperativity between
511   subunits in voltage-gated proton channels. *Nat Struct Mol Biol* **17**, 51–56 (2010).

512   39. L. Mony, T. K. Berger, E. Y. Isacoff, A specialized molecular motion opens the Hv1
513   voltage-gated proton channel. *Nat Struct Mol Biol* **22**, 283–290 (2015).

514   40. E. Suárez-Delgado, M. Orozco-Contreras, G. E. Rangel-Yescas, L. D. Islas, Activation-
515   pathway transitions in human voltage-gated proton channels revealed by a non-
516   canonical fluorescent amino acid. *eLife* **12**, e85836 (2023).

517   41. S. Han, *et al.*, Cholesterol inhibits human voltage-gated proton channel hHv1. *Proc.*
518   *Natl. Acad. Sci. U.S.A.* **119**, e2205420119 (2022).

519   42. P. Eggan, S. E. Gordon, W. N. Zagotta, Domain Coupling in Allosteric Regulation of
520   SthK Measured Using Time-Resolved Transition Metal Ion FRET. [Preprint] (2025).
521   Available at: https://elifesciences.org/reviewed-preprints/106892v2 [Accessed 2
522   August 2025].

523   43. F. W. Studier, Protein production by auto-induction in high-density shaking
524   cultures. *Protein Expression and Purification* **41**, 207–234 (2005).

525
526

527 **Figures and Tables**
528

529
Figure 1. hH$_v$1 was labeled with the fluorescent noncanonical amino acid

Acd by genetic code expansion. (A) Cartoon showing the components of the

genetic code expansion system used to incorporate Acd via amber codon

suppression in hH$_v$1. (B) AlphaFold structural model of the dimer hH$_v$1 colored by

subunit. (C) AlphaFold structural model of a single hH$_v$1 subunit colored by the

predicted Local Distance Difference Test (pLDDT). Model confidence was

represented by a gradient from red to blue, indicating low to high pLDDT. (D) In-

gel fluorescence and Western blot to evaluate Acd incorporation. An Acd

fluorescence band corresponding to the molecular weight of full-length hH$_v$1 (red

arrow) was observed in cellular extracts after separation by SDS-PAGE only

when cells contained the *Mj*A9-AcdRS/tRNA plasmid (RS) and were grown in the

presence of 1mM Acd (top). The Western blot against the His-tag present at the

N-terminus of hH$_v$1 confirmed that the band corresponded to the recombinant

protein (bottom). Note the hH$_v$1 truncated product produced when the amber stop

codon is replacing F247 in the absence of the aminoacyl-tRNA synthetase/tRNA

plasmid and Acd.

Figure 2. Acd was incorporated into 14 positions in the hH$_v$1 sequence. (A)
Cartoon showing the amino acids selected to be replaced by an amber stop
codon in the hH$_v$1 secondary structure to incorporate Acd (stars). (B) AlphaFold
dimer hH$_v$1 structural model with the amino acids selected to be replaced by Acd
as cyan spheres. (C) Acd fluorescence gel (top) and Western blot (bottom) of
cellular extracts after separation by SDS-PAGE showing the expression of the
Acd-labeled hH$_v$1 mutants. The No TAG lane contains an extract from cells
grown under identical conditions (in the presence of Acd and the aminoacyl-tRNA
synthetase/tRNA pair) and expressing hH$_v$1 without an amber stop codon.

Figure 3. Purification and functional measurements of hH$_v$1-Acd proteins.
(A) Coomassie-stained (top) and Acd fluorescence (bottom) gels after separation
by SDS-PAGE showing representative samples of the membrane extracts after
solubilization (left) and the final purified protein after immobilized metal affinity
chromatography (right). (B) Protein yield of the hH$_v$1 protein with Acd replacing
the amino acid at the indicated position. Note that V187Acd and Q233Acd
contained no detectable protein after purification. The red line indicates the
protein yield obtained from hH$_v$1 without an amber stop codon in the presence of
Acd and the aminoacyl-tRNA synthetase/tRNA pair (No TAG). (C)
Representative fluorescence-detection size-exclusion chromatograms of the
purified hH$_v$1 proteins with Acd incorporated at the indicated amino acid position.
Insets: AlphaFold dimer models, with the amino acid replaced by Acd shown as
cyan spheres. (D) Representative liposome proton flux assay of asolectin
proteoliposomes containing the indicated hH$_v$1 protein. All purified proteins
produced ACMA fluorescence quenching after the addition of valinomycin (Val).
The protonophore CCCP was added at the end of the experiment as a control.

The No TAG sample corresponds to proteoliposomes containing hH$_v$1 without an

amber stop codon expressed in the presence of Acd and the aminoacyl-tRNA

synthetase/tRNA pair.

Figure 4. Acd showed a low environmental sensitivity in hH$_v$1. (A)

Normalized fluorescence emission spectra of the Acd amino acid dissolved in the

indicated solvent (Excitation = 370 nm). EtAc: ethyl acetate, OctOH: 1-octanol,

ButOH: 1-butanol, EtOH: ethanol. (B) Normalized fluorescence decay of the Acd

amino acid dissolved in the indicated solvent. (C) Normalized fluorescence

emission spectra of the hH$_v$1-Acd proteins in Buffer-H2 (grey). The Acd amino

acid spectra in selected solvents are also shown for reference. Q56Acd and

C107Acd spectra are shown with broken lines. The emission maxima are listed in

Table S3. (D) Normalized fluorescence decay of the hH$_v$1-Acd proteins in Buffer-

H2 (grey). The Acd amino acid decays in selected solvents are also shown for

reference. Q56Acd and C107Acd decays are shown in red and black,

respectively.

Figure 5. Spectral FRET analysis using Trp and Tyr donors and Acd as acceptor. (A) Fluorescence emission spectra of the intrinsic fluorescent amino acids Trp (black) and Tyr (black, broken lines), along with the Acd absorption spectrum in Buffer-H2 (blue). (B) AlphaFold dimer hH$_v$1 structural model with the amino acids colored according to (C) and the native tryptophan and tyrosine residues shown as yellow spheres. (C) Normalized fluorescence spectra of the purified hH$_v$1-Acd proteins (Excitation = 280 nm). The spectra were normalized by the Trp/Tyr fluorescence. The colors correspond to the indicated hH$_v$1-Acd protein, and the remaining spectra are shown as grey traces in the figure. The No TAG sample (black) contains purified hH$_v$1 protein without any amber stop codon expressed in the presence of Acd and the aminoacyl-tRNA synthetase/tRNA pair. (D) Example of the spectral FRET analysis procedure. The fluorescence spectrum of hH$_v$1-Y134Acd obtained when exciting at 280 nm (black, F_{280}) minus the normalized spectrum at the same excitation wavelength of the No TAG sample (grey, $F_{280,NoTAG}$) produced the Acd emission spectrum shown in red (F_{280} − $F_{280,NoTAG}$). *Ratio A* values were calculated by dividing this spectrum by the hH$_v$1-Y134Acd fluorescence spectrum obtained by direct excitation at 370 nm (F_{370}). (E) *Ratio A* traces of the hH$_v$1-Acd proteins colored according to (C). The ratio between the spectra of free Acd amino acid excited at 280 and 370 nm

613   (*Ratio A$_0$*) is shown as broken lines. (F) Mean *Ratio A – Ratio A$_0$* values (410-480

614   nm; N=4) as a function of the predicted FRET efficiencies from the AlphaFold

615   hH$_v$1 structural model, colored according to (C). The broken line is the best fit to

616   the equation (*Ratio A – Ratio A$_0$*) = m(Predicted FRET efficiency).

Figure 6. FRET between Trp/Tyr and Acd reports a conformational rearrangement in response to Zn²⁺. (A) Normalized fluorescence emission spectra of hH$_v$1-K169Acd (Excitation = 280 nm) in Buffer-H2 in the absence of Zn²⁺ (black, Apo), in the presence of 1 mM Zn²⁺ (red), or in the presence of 1 mM Zn²⁺ and 2 mM EDTA (blue). Spectra were normalized by the maximum intensity of the Acd emission spectrum of the same sample excited at 370 nm. (B) Spectral FRET analysis of hH$_v$1-K169Acd in Buffer-H2 in the absence of Zn²⁺ (black, Apo), in the presence of 1 mM Zn²⁺ (red), or in the presence of 1 mM Zn²⁺ and 2 mM EDTA (blue). (C) Spectral FRET analysis of the hH$_v$1-Acd proteins in Buffer-H2 in the absence (black, Apo) or presence of 1 mM Zn²⁺ (red). N=4, ***p<0.005, **p<0.01.

 Supplementary Text

 Calculation of the FRET efficiency from the structural model. We calculated

 the FRET efficiency using the AlphaFold model of the full-length hH$_v$1 (Dataset

 S1). For each Acd position, the measured steady-state FRET would correspond

 to the mean FRET efficiency (E_{FRET}), which will be proportional to the *Ratio A –*

 Ratio A$_0$ value obtained from the spectral FRET analysis.

     For *n* number of donors and two acceptors in the hH$_v$1 dimer:

$$E_{FRET} = \sum_{i=1}^{n} P_{Ex,i} E_i \quad (1)$$

 where $P_{Ex,i}$ is the probability of excitation and E_i is the FRET efficiency of the *i*th

 donor. $P_{Ex,i}$ is the number of photons absorbed by the *i*th donor divided by the

 total number of photons absorbed by the total number *n* of donors, which is

 quantified by the donor's extinction coefficient. Supposing the same extinction

 coefficient for each of the Trp ($e_W = 5,501$ M^{-1} cm^{-1}) and Tyr ($e_Y = 1,209$ M^{-1} cm^{-1}

)(1), equation 1 can be written as:

$$E_{FRET} = \sum_{i=1}^{n} \left(\frac{\varepsilon_i}{\sum_{i=1}^{n} \varepsilon_i} \right) E_i = \frac{1}{n_W \varepsilon_W + n_Y \varepsilon_Y} \left(\varepsilon_W \sum_{i=1}^{n_W} E_{W,i} + \varepsilon_Y \sum_{i=1}^{n_Y} E_{Y,i} \right) \quad (2)$$

 where the sum of the total number of Trp (n_W) and Tyr (n_Y) is equal to *n*. The

 efficiency of FRET for the *i*th donor can be expressed as the sum of rates of

 energy transfer to each of the *j* = 2 Acd acceptors, k_{ij}, divided by the total rates of

 photon emission. Therefore, the right terms of equation 2 can be expressed as:

$$\sum_{i=1}^{n_W} E_{W,i} = \sum_{i=1}^{n_W} \left(\frac{\sum_{j=1}^{2} k_{ij}}{\sum_{j=1}^{2} k_{ij} + (1/\tau_W)} \right) \quad \text{and} \quad \sum_{i=1}^{n_Y} E_{Y,i} = \sum_{i=1}^{n_Y} \left(\frac{\sum_{j=1}^{2} k_{ij}}{\sum_{j=1}^{2} k_{ij} + (1/\tau_Y)} \right) \quad (3)$$

 where the Trp lifetime ($t_W = 3.1$ ns) and tyrosine lifetime ($t_Y = 3.6$ ns) in the

 absence of donors are considered constant(1). Finally, each energy transfer rate

 k_{ij} is a function of the distance r_{ij} between the *i*th donor and the *j*th acceptor:

$$k_{ij}(r) = \frac{1}{\tau_D} \left(\frac{R_0}{r_{ij}} \right)^6 \quad (4)$$

 where the donor lifetime t$_D$ and R_0 would be t_W and 23.5 Å or t_Y and 20.9 Å for

 Trp or Tyr residues, respectively. Calculating E_{FRET} using the above equations

 considers a fixed distance between donors and acceptors. A better approach is

 to consider the distance distributions $P_{ij}(r)$ produced by the different rotameric

states of the FRET pair, which can be modeled easily using chiLife(2). Then,

equation 4 will be replaced by equation 5:

$$k_{ij}(r) = \int_0^\infty P_{ij}(r) \left[\frac{1}{\tau_D} \left(\frac{R_0}{r} \right)^6 \right] dr \quad (5).$$

DNA plasmid sequences in FASTA format.

>His-EK-hHv1-C107A-C249A.pET15-b

TTGAGATCCTTTTTTTCTGCGCGTAATCTGCTGCTTGCAAACAAAAAACCAC

CGCTACCAGCGGTGGTTTGTTTGCCGGATCAAGAGCTACCAACTCTTTTTCC

GAAGGTAACTGGCTTCAGCAGAGCGCAGATACCAAATACTGTCCTTCTAGT

GTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACCGCCTACATA

CCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGGCGATAAGTC

GTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGGATAAGGCGCAGCG

GTCGGGCTGAACGGGGGGTTCGTGCACACAGCCCAGCTTGGAGCGAACGA

CCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAGCGCCACGC

TTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAGGGTCGGA

ACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTATCTTTA

TAGTCCTGTCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTTGTGATGC

TCGTCAGGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGCGGCCTTTTT

ACGGTTCCTGGCCTTTTGCTGGCCTTTTGCTCACATGTTCTTTCCTGCGTTA

TCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGCTGATACCG

CTCGCCGCAGCCGAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGC

GGAAGAGCGCCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTCA

CACCGCATATATGGTGCACTCTCAGTACAATCTGCTCTGATGCCGCATAGTT

AAGCCAGTATACACTCCGCTATCGCTACGTGACTGGGTCATGGCTGCGCCC

CGACACCCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGTCTGCTCCC

GGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATGTGTCA

GAGGTTTTCACCGTCATCACCGAAACGCGCGAGGCAGCTGCGGTAAAGCT

CATCAGCGTGGTCGTGAAGCGATTCACAGATGTCTGCCTGTTCATCCGCGT

CCAGCTCGTTGAGTTTCTCCAGAAGCGTTAATGTCTGGCTTCTGATAAAGCG

GGCCATGTTAAGGGCGGTTTTTTCCTGTTTGGTCACTGATGCCTCCGTGTAA

GGGGGATTTCTGTTCATGGGGGTAATGATACCGATGAAACGAGAGAGGATG

CTCACGATACGGGTTACTGATGATGAACATGCCCGGTTACTGGAACGTTGT

GAGGGTAAACAACTGGCGGTATGGATGCGGCGGGACCAGAGAAAATCAC

TCAGGGTCAATGCCAGCGCTTCGTTAATACAGATGTAGGTGTTCCACAGGG

TAGCCAGCAGCATCCTGCGATGCAGATCCGGAACATAATGGTGCAGGGCG

CTGACTTCCGCGTTTCCAGACTTTACGAAACACGGAAACCGAAGACCATTCA

TGTTGTTGCTCAGGTCGCAGACGTTTTGCAGCAGCAGTCGCTTCACGTTCG

CTCGCGTATCGGTGATTCATTCTGCTAACCAGTAAGGCAACCCCGCCAGCC

TAGCCGGGTCCTCAACGACAGGAGCACGATCATGCGCACCCGTGGCCAGG

ACCCAACGCTGCCCGAGATGCGCCGCGTGCGGCTGCTGGAGATGGCGGA

CGCGATGGATATGTTCTGCCAAGGGTTGGTTTGCGCATTCACAGTTCTCCG

CAAGAATTGATTGGCTCCAATTCTTGGAGTGGTGAATCCGTTAGCGAGGTG

CCGCCGGCTTCCATTCAGGTCGAGGTGGCCCGGCTCCATGCACCGCGACG

CAACGCGGGGAGGCAGACAAGGTATAGGGCGGCGCCTACAATCCATGCCA

ACCCGTTCCATGTGCTCGCCGAGGCGGCATAAATCGCCGTGACGATCAGC

GGTCCAGTGATCGAAGTTAGGCTGGTAAGAGCCGCGAGCGATCCTTGAAG

CTGTCCCTGATGGTCGTCATCTACCTGCCTGGACAGCATGGCCTGCAACGC

GGGCATCCCGATGCCGCCGGAAGCGAGAAGAATCATAATGGGGAAGGCCA

TCCAGCCTCGCGTCGCGAACGCCAGCAAGACGTAGCCCAGCGCGTCGGCC

GCCATGCCGGCGATAATGGCCTGCTTCTCGCCGAAACGTTTGGTGGCGGG

ACCAGTGACGAAGGCTTGAGCGAGGGCGTGCAAGATTCCGAATACCGCAA

GCGACAGGCCGATCATCGTCGCGCTCCAGCGAAAGCGGTCCTCGCCGAAA

ATGACCCAGAGCGCTGCCGGCACCTGTCCTACGAGTTGCATGATAAAGAAG

ACAGTCATAAGTGCGGCGACGATAGTCATGCCCCGCGCCCACCGGAAGGA

GCTGACTGGGTTGAAGGCTCTCAAGGGCATCGGTCGAGATCCCGGTGCCT

AATGAGTGAGCTAACTTACATTAATTGCGTTGCGCTCACTGCCCGCTTTCCA

GTCGGGAAACCTGTCGTGCCAGCTGCATTAATGAATCGGCCAACGCGCGG

GGAGAGGCGGTTTGCGTATTGGGCGCCAGGGTGGTTTTTCTTTTCACCAGT

GAGACGGGCAACAGCTGATTGCCCTTCACCGCCTGGCCCTGAGAGAGTTG

CAGCAAGCGGTCCACGCTGGTTTGCCCCAGCAGGCGAAAATCCTGTTTGAT

GGTGGTTAACGGCGGGATATAACATGAGCTGTCTTCGGTATCGTCGTATCC

CACTACCGAGATATCCGCACCAACGCGCAGCCCGGACTCGGTAATGGCGC

GCATTGCGCCCAGCGCCATCTGATCGTTGGCAACCAGCATCGCAGTGGGA

ACGATGCCCTCATTCAGCATTTGCATGGTTTGTTGAAAACCGGACATGGCAC

TCCAGTCGCCTTCCCGTTCCGCTATCGGCTGAATTTGATTGCGAGTGAGATA

TTTATGCCAGCCAGCCAGACGCAGACGCGCCGAGACAGAACTTAATGGGC

CCGCTAACAGCGCGATTTGCTGGTGACCCAATGCGACCAGATGCTCCACGC

CCAGTCGCGTACCGTCTTCATGGGAGAAATAATACTGTTGATGGGTGTCT

GGTCAGAGACATCAAGAAATAACGCCGGAACATTAGTGCAGGCAGCTTCCA

CAGCAATGGCATCCTGGTCATCCAGCGGATAGTTAATGATCAGCCCACTGA

CGCGTTGCGCGAGAAGATTGTGCACCGCCGCTTTACAGGCTTCGACGCCG

CTTCGTTCTACCATCGACACCACCACGCTGGCACCCAGTTGATCGGCGCGA

GATTTAATCGCCGCGACAATTTGCGACGGCGCGTGCAGGGCCAGACTGGA

GGTGGCAACGCCAATCAGCAACGACTGTTTGCCCGCCAGTTGTTGTGCCAC

GCGGTTGGGAATGTAATTCAGCTCCGCCATCGCCGCTTCCACTTTTTCCCG

CGTTTTCGCAGAAACGTGGCTGGCCTGGTTCACCACGCGGGAAACGGTCT

GATAAGAGACACCGGCATACTCTGCGACATCGTATAACGTTACTGGTTTCAC

ATTCACCACCCTGAATTGACTCTCTTCCGGGCGCTATCATGCCATACCGCGA

AAGGTTTTGCGCCATTCGATGGTGTCCGGGATCTCGACGCTCTCCCTTATG

CGACTCCTGCATTAGGAAGCAGCCCAGTAGTAGGTTGAGGCCGTTGAGCAC

CGCCGCCGCAAGGAATGGTGCATGCAAGGAGATGGCGCCCAACAGTCCCC

CGGCCACGGGGCCTGCCACCATACCCACGCCGAAACAAGCGCTCATGAGC

CCGAAGTGGCGAGCCCGATCTTCCCCATCGGTGATGTCGGCGATATAGGC

GCCAGCAACCGCACCTGTGGCGCCGGTGATGCCGGCCACGATGCGTCCG

GCGTAGAGGATCGAGATCTCGATCCCGCGAAATTAATACGACTCACTATAG

GGGAATTGTGAGCGGATAACAATTCCCCTCTAGAAATAATTTTGTTTAACTTT

AAGAAGGAGATATACCATGGGCAGCAGCCATCATCATCATCATCACAGCAG

CGGCGATGATGATGATAAAATGGCGACCTGGGACGAAAAGGCGGTGACCC

GTCGTGCGAAAGTTGCGCCGGCGGAGCGTATGAGCAAGTTCCTGCGTCAC

TTTACCGTGGTTGGTGACGATTACCACGCGTGGAACATCAACTATAAGAAAT

GGGAGAACGAGGAAGAGGAAGAGGAAGAGGAACAGCCGCCGCCGACCCC

GGTTAGCGGCGAGGAAGGCCGTGCGGCGGCGCCGGATGTGGCGCCGGC

GCCGGGTCCGGCGCCGCGTGCGCCGCTGGATTTCCGTGGCATGCTGCGTA

AACTGTTCAGCAGCCACCGTTTTCAAGTTATCATTATCGCGCTGGTGGTTCT

GGATGCGCTGCTGGTGCTGGCGGAGCTGATCCTGGACCTGAAGATTATCCA

GCCGGATAAAACAACTACGCGGCGATGGTTTTCCACTATATGAGCATCAC

CATTCTGGTGTTCTTTATGATGGAAATCATCTTCAAGCTGTTCGTTTTCCGTC

TGGAGTTCTTTCACCACAAATTCGAAATCCTGGACGCGGTGGTTGTGGTTGT

GAGCTTTATCCTGGATATTGTGCTGCTGTTCCAGGAGCACCAATTTGAAGCG

CTGGGTCTGCTGATTCTGCTGCGTCTGTGGCGTGTTGCGCGTATTATCAAC

GGCATTATCATTAGCGTGAAGACCCGTAGCGAGCGTCAGCTGCTGCGTCTG

AAGCAGATGAACGTTCAACTGGCGGCGAAAATCCAGCACCTGGAATTTAGC

GCGAGCGAGAAGGAACAAGAGATTGAACGTCTGAACAAACTGCTGCGTCAG

CACGGTCTGCTGGGCGAAGTGAACTAAGGATCCGGCTGCTAACAAAGCCC

GAAAGGAAGCTGAGTTGGCTGCTGCCACCGCTGAGCAATAACTAGCATAAC

CCCCTTGGGGCCTCTAAACGGGTCTTGAGGGGTTTTTTGCTGAAAGGAGGA

ACTATATCCGGATATCCCGCAAGAGGCCCGGCAGTACCGGCATAACCAAGC

CTATGCCTACAGCATCCAGGGTGACGGTGCCGAGGATGACGATGAGCGCA

TTGTTAGATTTCATACACGGTGCCTGACTGCGTTAGCAATTTAACTGTGATAA

ACTACCGCATTAAAGCTTATCGATGATAAGCTGTCAAACATGAGAATTCTTG

AAGACGAAAGGGCCTCGTGATACGCCTATTTTTATAGGTTAATGTCATGATA

ATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGA

ACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAGA

CAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAGTAT

TCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTGCCTTCCTG

TTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGAAGATCAGTT

GGGTGCACGAGTGGGTTACATCGAACTGGATCTCAACAGCGGTAAGATCCT

TGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCACTTTTAAAGTT

CTGCTATGTGGCGCGGTATTATCCCGTGTTGACGCCGGGCAAGAGCAACTC

GGTCGCCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTCACCAGTCA

CAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTG

CCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCTGACAACGATCG

GAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGATCATGTAA

CTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACCAAACGACG

AGCGTGACACCACGATGCCTGCAGCAATGGCAACAACGTTGCGCAAACTAT

TAACTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAATAGACTGGAT

GGAGGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCTTCCGGCTG

GCTGGTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGGGTCTCGCGGTA

TCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGTAGTTATCT

ACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTG

AGATAGGTGCCTCACTGATTAAGCATTGGTAACTGTCAGACCAAGTTTACTC

ATATATACTTTAGATTGATTTAAAACTTCATTTTTAATTTAAAAGGATCTAGGT

GAAGATCCTTTTTGATAATCTCATGACCAAAATCCCTTAACGTGAGTTTTCGT

TCCACTGAGCGTCAGACCCCGTAGAAAGATCAAGGATCTTC

>MjA9Acd-RS.pDule2

TTGAGATCGTTTTGGTCTGCGCGTAATCTCTTGCTCTGAAAACGAAAAAACC

GCCTTGCAGGGCGGTTTTTCGAAGGTTCTCTGAGCTACCAACTCTTTGAACC

GAGGTAACTGGCTTGGAGGAGCGCAGTCACCAAAACTTGTCCTTTCAGTTT

AGCCTTAACCGGCGCATGACTTCAAGACTAACTCCTCTAAATCAATTACCAG

TGGCTGCTGCCAGTGGTGCTTTTGCATGTCTTTCCGGGTTGGACTCAAGAC

GATAGTTACCGGATAAGGCGCAGCGGTCGGACTGAACGGGGGGTTCGTGC

ATACAGTCCAGCTTGGAGCGAACTGCCTACCCGGAACTGAGTGTCAGGCGT

GGAATGAGACAAACGCGGCCATAACAGCGGAATGACACCGGTAAACCGAA

AGGCAGGAACAGGAGAGCGCACGAGGGAGCCGCCAGGGGGAAACGCCTG

GTATCTTTATAGTCCTGTCGGGTTTCGCCACCACTGATTTGAGCGTCAGATT

TCGTGATGCTTGTCAGGGGGGCGGAGCCTATGGAAAAACGGCTTTGCCGC

GGCCCTCTCACTTCCCTGTTAAGTATCTTCCTGGCATCTTCCAGGAAATCTC

CGCCCCGTTCGTAAGCCATTTCCGCTCGCCGCAGTCGAACGACCGAGCGT

AGCGAGTCAGTGAGCGAGGAAGCGGAATATATCCTGTATCACATATTCTGC

TGACGCACCGGTGCAGCCTTTTTTCTCCTGCCACATGAAGCACTTCACTGAC

ACCCTCATCAGTGCCAACATAGTAAGCCAGTATACACTCCGCTAGCGCTGA

TGTCCGGCGGTGCTTTTGCCGTTACGCACCACCCCGTCAGTAGCTGAACAG

GAGGGACAGCTCCCGGCGGATTTGTCCTACTCAGGAGAGCGTTCACCGAC

```
813   AAACAACAGATAAAACGAAAGGCCCAGTCTTTCGACTGAGCCTTTCGTTTTA
814   TTTGATGCCTGGCAGTTCCCTACTCTCGCATGGGGAGACCCCACACTACCA
815   TCGGCGCTACGGCGTTTCACTTCTGAGTTCGGCATGGGGTCAGGTGGGAC
816   CACCGCGCTACTGCCGCCAGGCAAATTCTGTTTTATCAGACCGCTTCTGCG
817   TTCTGATTTAATCTGTATCAGGCTGAAAATCTTCTCTCATCCGCCAAAACAGC
818   CAAGCTGGAGACCGTTTAAACTCAATGATGATGATGATGATGGTCGACGGC
819   GCTATTCAGATCCTCTTCTGAGATGAGTTTTTGTTCGGGCCCAAGCTTCGAA
820   TTCCCATATGGTACCCGTTTGAAACTGCAGTTATAATCTCTTTCTAATTGGCT
821   CTAAATCTTTATAAGTTCTTCAGCTACAGCATTTTTTAAATCCATTGGATGC
822   AATTCCTTATTTTTAAATAAACTCTCTAACTCCTCATAGCTATTAACTGTCAAA
823   TCTCCACCAAATTTTTCTGGCCTTTTTATGGTTAAAGGATATTCAAGGAAGTA
824   TTTAGCTATCTCCATTATTGGATTTCCTTCAACAACTCCAGCTGGGCAGTATG
825   CTTTCTTTATCTTAGCCCTAATCTCTTCTGGAGAGTCATCAACAGCTATAAAA
826   TTCCCTTTTGAAGAACTCATCTTTCCTTCTCCATCCAAACCCGTTAAGACAGG
827   GTTGTGAATACAAACAACCTTTTTTGGTAAAAGCTCCCTTGCTAACATGTTTA
828   TTTTTCTCTGCTCCATCCCTCCAACTGCAACATCAACGCCAGTATAATGAATA
829   GAATTAACCTGCATTATTGGATAGATAACTTCAGCAACCTTTGGATTTTCATC
830   CTCTCTTGCTATAAGTTCCATACTCCTTCTTGCTCTTTTTAAGGTAGTTTTTAA
831   AGCCAATCTATAGACATTCAGTGTATAATCCTTATCAAGCTCCAGTTCACTTC
832   CATAAACATATTTTGCCTTTAACCCCATTGCTTCAAAAACTTTTTTGTTATAAT
833   CTCCTATTTTTCTAATCTCATCCAACTCTCCTTTCTGGTTTAAATAGGCGTGT
834   AAATCAGCCAAATCTATAATTATATCAAATCCAGCATTTTGTAAATCAATCAT
835   CTTTTTTATTTGGAGATAATGCCCTAAATGTATTTTACCACTTGGTTCAAAAC
836   CTATCGCAGCAGATTTTTCATCTTTTTTTAAAACCTCTCTTAACTCTTCCTCG
837   CTGATAATTTCAGATGTGTTTCTCTTTATCATTTCAAATTCGTCCATGGGGGA
838   TTCCTCAAAGCGTAAACTCAGCGTTACAAGTATTACACAAAGTTTTTTATGTT
839   GAGAATATTTTTTTGATGGGGCGCCACTTATTTTTGATCGTTCGCTCAAAGAA
840   GCGGCGCCAGGGTTGTTTTTCTTTTCACCGGTGAGACGGGCAACAGAACGC
841   CATGAGCGGCCTCATTTCTTATTCTGAGTTACAACAGTCCGCACCGCTGTCC
842   GTATATATGAGTAAACTTGGTCCCGGGTTACCGGTTTGGTTAGCGAGAAGA
843   GCCAGTAAAAGACGCAGTGACGGCAATGTCTGATGCAATATGGACAATTGG
```

TTTCTTCTCTGAATGGCGGGAGTATGAAAAGTATGGCTGAAGCGCAAAATGA

TCCCCTGCTGCCGGGATACTCGTTTAATGCCCATCTGGTGGCGGGTTTAAC

GCCGATTGAGGCCAACGGTTATCTCGATTTTTTTATCGACCGACCGCTGGG

AATGAAAGGTTATATTCTCAATCTCACCATTCGCGGTCAGGGGGTGGTGAAA

AATCAGGGACGAGAATTTGTTTGCCGACCGGGTGATATTTTGCTGTTCCCG

CCAGGAGAGATTCATCACTACGGTCGTCATCCGGAGGCTCGCGAATGGTAT

CACCAGTGGGTTTACTTTCGTCCGCGCGCCTACTGGCATGAATGGCTTAAC

TGGCCGTCAATATTTGCCAATACGGGGTTCTTTCGCCCGGATGAAGCGCAC

CAGCCGCATTTCAGCGACCTGTTTGGGCAAATCATTAACGCCGGGCAAGGG

GAAGGGCGCTATTCGGAGCTGCTGGCGATAAATCTGCTTGAGCAATTGTTA

CTGCGGCGCATGGAAGCGATTAACGAGTCGCTCCATCCACCGATGGATAAT

CGGGTACGCGAGGCTTGTCAGTACATCAGCGATCACCTGGCAGACAGCAAT

TTTGATATCGCCAGCGTCGCACAGCATGTTTGCTTGTCGCCGTCGCGTCTG

TCACATCTTTTCCGCCAGCAGTTAGGGATTAGCGTCTTAAGCTGGCGCGAG

GACCAACGTATCAGCCAGGCGAAGCTGCTTTTGAGCACCACCCGGATGCCT

ATCGCCACCGTCGGTCGCAATGTTGGTTTTGACGATCAACTCTATTTCTCGC

GGGTATTTAAAAAATGCACCGGGGCCAGCCCGAGCGAGTTCCGTGCCGGT

TGTGAAGAAAAGTGAATGATGTAGCCGTCAAGTTGTCATAATTGGTAACGA

ATCAGACAATTGACGGCTTGACGGAGTAGCATAGGGTTTGCAGAATCCCTG

CTTCGTCCATTTGACAGGCACATTATGCATGCCGCTTCGCCTTCGCGCGCG

AATTGATCTGCTGCCTCGCGCGTTTCGGTGATGACGGTGAAAACCTCTGAC

ACATGCAGCTCCCGGAGACGGTCACAGCTTGTCTGTAAGCGGATGCCGGG

AGCAGACAAGCCCGTCAGGGCGCGTCAGCGGGTGTTGGCGGGTGTCGGG

GCGCAGCCATGACCCAGTCAACTGCGATGAGTGGCAGGGCGGGGCGTAAT

TTTTTTAAGGCAGTTATTGGTGCCCTTAAACGCCTGGTTGCTACGCCTGAAT

AAGTGATAATAAGCGGATGAATGGCAGAAATTCGAAAGCAAATTCGACCCG

GTCGTCGGTTCAGGGCAGGGTCGTTAAATAGCCGCTTATGTCTATTGCTGG

TTTACCGGTTTATTGACTACCGGAAGCAGTGTGACCGTGTGCTTCTCAAATG

CCTGAGGCCAGTTTGCTCAGGCTCTCCCCGTGGAGGTAATAATTGACGATA

TGATCATTTATTCTGCCTCCCAGAGCATGATAAAAACGGTTAGCGCTTCGTT

AATACAGATGTAGGTGTTCCACAGGGTAGCCAGCAGCATCCTGCGATGCAG

```
875   ATCCGGAACATAATGGTGCAGGGCGCTTGTTTCGGCGTGGGTATGGTGGCA
876   GGCCCCGTGGCCGGGGGACTGTTGGGCGCTGCCGGCACCTGTCCTACGA
877   GTTGCATGATAAAGAAGACAGTCATAAGTGCGGCGACGATAGTCATGCCCC
878   GCGCCCACCGGAAGGAGCTACCGGCAGCGGTGCGGACTGTTGTAACTCAG
879   AATAAGAAATGAGGCCGCTCATGGCGTTCTGTTGCCCGTCTCACTGGTGAA
880   AAGAAAACAACCCTGGCGCCGCTTCTTTGAGCGAACGATCAAAAATAAGT
881   GGCGCCCCATCAAAAAAATATTCTCAACATAAAAAACTTTGTGTAATACTTGT
882   AACGCTGAATTCCCGGCGGTAGTTCAGCAGGGCAGAACGGCGGACTCTAA
883   ATCCGCATGGCGCTGGTTCAAATCCGGCCCGCCGGACCACTGCAGATCCTT
884   AGCGAAAGCTAAGGATTTTTTTTAAGCTTGGCACTGGCCGTCGTTTTACAAC
885   GTCGTGACTGGGAAAACCCTGGCGTTACCCAACTTAATCGCCTTGCAGCAC
886   ATCCCCCTTTCGCCAGACGCTCTCCCTTATGCGACTCCTGCATTAGGAAGC
887   AGCCCAGTAGTAGGTTGAGGCCGTTGAGCACCGCCGCCGCAAGGAATGGT
888   GCATGCAAGGAGCCCGAGATGCGCCGCGTGCGGCTGCTGGAGATGGCGG
889   ACGCGATGGATATGTTCTGCCAAGGGTTGGTTTGCGCATTCACAGTTCTCC
890   GCAAGAATTGATTGGCTCCAATTCTTGGAGTGGTGAATCCGTTAGCGAGGT
891   GCCGCCGGCTTCCATTCAGGTCGAGGTGGCCCGGCTCCATGCACCGCGAC
892   GCAACGCGGGGAGGCAGACAAGGTATAGGGCGGCGCCTACAATCCATGCC
893   AACCCGTTCCATGTGCTCGCCGAGGCGGCATAAATCGCCGTGACGATCAG
894   CGGTCCAATGATCGAAGTTAGGCTGGTAAGAGCCGCGAGCGATCCTTGAAG
895   CTGTCCCTGATGGTCGTCATCTACCTGCCTGGACAGCATGGCCTGCAACGC
896   GGGCATCCCGATGCCGCCGGAAGCGAGAAGAATCATAATGGGGAAGGCCA
897   TCCAGCCTCGCGTCGCGAACGCCAGCAAGACGTAGCCCAGCGCGTCGGCC
898   GCCATGCCGGCGATAATGGCCTGCTTCTCGCCGAAACGTTTGGTGGCGGG
899   ACCAGTGACGAAGGCTTGAGCGAGGGCGTGCAAGATTCCGAATACCGCAA
900   GCGACAGGCCGATCATCGTCGCGCTCCAGCGAAAGCGGTCCTCGCCGAAA
901   ATGACCCAGAGCGCTGCCGGCACCTGTCCTACGAGTTGCATGATAAAGAAG
902   ACAGTCATAAGTGCGGCGACGATAGTCATGCCCCGCGCCCACCGGAAGGA
903   GCTGACTGGGTTGAAGGCTCTCAAGGGCATCGGTCGACGCTCTCCCTTATG
904   CGACTCCTGCATTAGGAAGCAGCCCAGTAGTAGGTTGAGGCCGTTGAGCAC
905   CGCCGCCGCAAGGAATGGTGCATGCAAGGAGATGGCGCCCAACAGTCCCC
```

CGGCCACGGGGCCTGCCACCATACCCACGCCGAAACAAGCGCTCATGAGC

CCGAAGTGGCGAGCCCGATCTTCCCCATCGGTGATGTCGGCGATATAGGC

GCCAGCAACCGCACCTGTGGCGCCGGTGATGCCGGCCACGATGCGTCCG

GCGTAGAGGATCCCGGTAAACCAGCAATAGACATAAGCGGCTATTTAACGA

CCCTGCCCTGAACCGACGACCGGGTCATCGTGGCCGGATCTTGCGGCCCC

TCGGCTTGAACGAATTGTTAGACATTATTTGCCGACTACCTTGGTGATCTCG

CCTTTCACGTAGTGGACAAATTCTTCCAACTGATCTGCGCGCGAGGCCAAG

CGATCTTCTTCTTGTCCAAGATAAGCCTGTCTAGCTTCAAGTATGACGGGCT

GATACTGGGCCGGCAGGCGCTCCATTGCCCAGTCGGCAGCGACATCCTTC

GGCGCGATTTTGCCGGTTACTGCGCTGTACCAAATGCGGGACAACGTAAGC

ACTACATTTCGCTCATCGCCAGCCCAGTCGGGCGGCGAGTTCCATAGCGTT

AAGGTTTCATTTAGCGCCTCAAATAGATCCTGTTCAGGAACCGGATCAAAGA

GTTCCTCCGCCGCTGGACCTACCAAGGCAACGCTATGTTCTCTTGCTTTTGT

CAGCAAGATAGCCAGATCAATGTCGATCGTGGCTGGCTCGAAGATACCTGC

AAGAATGTCATTGCGCTGCCATTCTCCAAATTGCAGTTCGCGCTTAGCTGGA

TAACGCCACGGAATGATGTCGTCGTGCACAACAATGGTGACTTCTACAGCG

CGGAGAATCTCGCTCTCTCCAGGGGAAGCCGAAGTTTCCAAAAGGTCGTTG

ATCAAAGCTCGCCGCGTTGTTTCATCAAGCCTTACGGTCACCGTAACCAGC

AAATCAATATCACTGTGTGGCTTCAGGCCGCCATCCACTGCGGAGCCGTAC

AAATGTACGGCCAGCAACGTCGGTTCGAGATGGCGCTCGATGACGCCAACT

ACCTCTGATAGTTGAGTCGATACTTCGGCGATCACCGCTTCCCTCATACTCT

TCCTTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGA

TACATATTTGAATGTATTTAGAAAAATAAACAAATAGCTAGCTCACTCGGTCG

CATCGATGATAAGCTGTCAAACATGAGAATTACAACTTATATCGTATGGGGC

TGACTTCAGGTGCTACATTTGAAGAGATAAATTGCACTGAAATCTAGAAATAT

TTTATCTGATTAATAAGATGATCTTC

Supplementary Figures

Figure S1. AlphaFold dimer hH$_v$1 structural model. (A) Extracellular view of

the model, showing each subunit in a different color. (B) One subunit of the dimer

showing the position of the arginine gating charges in S4 (blue), the selectivity

filter (red), and the charge transfer center (yellow).

Figure S2. The protein yield of hH$_v$1-Acd depended on the position of Acd incorporation. (A) Optical density at 600 nm (OD$_{600}$) of the final cultures expressing the hH$_v$1 with Acd replacing the indicated amino acid. (B) Relative protein yield calculated by multiplying the OD$_{600}$ and Western blot band intensity of cellular extracts from the final cultures expressing the hH$_v$1 with Acd replacing the indicated amino acid. Values of normalized by the No TAG condition, which are cells expressing the hH$_v$1 without any amber stop codon in the presence of Acd and the aminoacyl-tRNA synthetase/tRNA pair (red broken line).

950
951
Figure S3. The truncation of hH$_v$1-K125Acd was not decreased at higher

concentrations of Acd in the culture medium. (A) Cellular extracts of cells co-

transformed with the MjA9-AcdRS/tRNA pair plasmid and hH$_v$1-K125TAG and

grown with the indicated concentration of Acd in the culture medium were

separated by SDS-PAGE to visualize the hH$_v$1 band (red arrow) by Acd

fluorescence (top) and Western blot against the N-terminus His-tag (bottom). (B)

Optical density at 600 nm (OD$_{600}$) of the cultures at the end of the expression. (C)

Relative yield obtained by multiplying the Western blot band intensity by the

OD600, normalized by the value obtained at 1 mM Acd.

Figure S4. SDS-PAGE of the hH$_v$1 protein samples after purification.

Coomassie-stained (top) and Acd fluorescence (bottom) gels of the samples from the elution of the nickel resin column, separated by SDS-PAGE. Note that V187Acd and Q233Acd did not contain appreciable amounts of protein. The No TAG condition consisted of samples purified from cells expressing the hH$_v$1 without any amber stop codon in the presence of Acd and the aminoacyl-tRNA synthetase/tRNA pair.

972 **Figure S5. Size exclusion chromatogram of the No TAG sample.** The No

973 TAG sample was purified from cells expressing the hH$_v$1 without any amber stop

974 codon in the presence of Acd and the aminoacyl-tRNA synthetase/tRNA pair.

975 The sample eluted from the immobilized affinity chromatography column was

976 concentrated and injected into the size exclusion chromatography column.

Figure S6. Fluorescence-detection size exclusion chromatography of hH$_v$1-Acd proteins. Chromatograms of the purified hH$_v$1 proteins with Acd incorporated at the indicated amino acid position. The AlphaFold dimer model with the amino acid replaced by Acd as cyan spheres is included.

Figure S7. The purified hH$_v$1-Acd proteins were functional proton channels.
Liposome proton flux assays of asolectin proteoliposomes containing the
indicated hH$_v$1-Acd protein. The proteoliposome samples produced ACMA
fluorescence quenching after the addition of valinomycin (180 s) in the presence
of a potassium gradient. The protonophore CCCP was added at the end of the
experiment (600 s). The No TAG sample (red trace) contains proteoliposomes
containing hH$_v$1 without any amber stop codon expressed in the presence of Acd

992   and the aminoacyl-tRNA synthetase/tRNA pair. Empty asolectin liposomes (black

993   traces) showed slow ACMA fluorescence quenching.

Fig. S8. Fluorescence emission spectra of hH$_v$1-Acd. Spectra of the indicated hH$_v$1-Acd protein sample obtained when exciting at 280 nm (black) and 370 nm (blue). The isolated spectrum of Acd (red) was obtained by subtracting the black

999   trace from the normalized spectrum of the No TAG sample excited at 280 nm

1000   (grey).

1001

1002

1003 **Figure S9. Fluorescence emission spectra of the Acd amino acid in**

1004 **solution.** Spectra of Acd in Buffer-H2 using an excitation wavelength of 370 nm

1005 (blue) or 280 nm (red).

1006

1007

Fig. S10. Changes in the fluorescence emission spectra of hH$_v$1-Acd in the presence of zinc. Spectra of the indicated hH$_v$1-Acd protein sample obtained when exciting at 280 nm in the Apo state (black) and in the presence of 1 mM

1011 Zn^{2+} (red). Spectra were normalized by the maximum intensity of the Acd

1012   emission spectrum of the same sample excited at 370 nm.

1013

1014

Figure S11. The protein yield of hH$_v$1-A18Acd was proportional to the

concentration of Acd in the culture medium. (A) Cellular extracts of cells co-

transformed with the MjA9-AcdRS/tRNA pair plasmid and hH$_v$1-A18TAG and

grown with the indicated concentration of Acd in the culture medium were

separated by SDS-PAGE to visualize the hH$_v$1 band (red arrow) by Acd

fluorescence (top) and Western blot against the N-terminus His-tag (bottom). (B)

Optical density at 600 nm (OD$_{600}$) of the cultures at the end of the expression. (C)

Relative yield obtained by multiplying the Western blot band intensity by the

OD$_{600}$, normalized by the value obtained at 1 mM Acd.

1024

1025   **Supplementary Tables**

1026   **Table S1. Size-exclusion chromatography elution volumes of the hH$_v$1**

1027   **proteins.** Elution volumes were measured as the location of the maximum value

1028   of absorbance at 280 nm in the chromatograms.

hH$_v$1 sample	Elution volume (mL)
No TAG	11.5
A18Acd	11.6
Q56Acd	11.7
M91Acd	11.9
V103Acd	11.5
C107Acd	12.5
K125Acd	11.7
Y134Acd	11.5
F159Acd	11.5
K169Acd	11.5
F195 Acd	11.5
I217Acd	12.4
F247Acd	13.1

1029

1030

1031 **Table S2. Emission fluorescence spectrum maximum of Acd in different**

1032 **solvents.** The wavelength corresponds to the location of the maximum intensity

1033 of the emission fluorescence spectrum of the free amino acid dissolved in the

1034 corresponding solvent. EtAc: ethyl acetate, OctOH: 1-octanol, ButOH: 1-butanol,

1035 EtOH: ethanol.

Solvent	Emission Maximum (nm)
EtAc	402
OctOH	414
ButOH	415
EtOH	414
Water	421
Buffer-H2	420

1036

1037

1038 **Table S3. The emission fluorescence spectrum maximum of the hH$_v$1-Acd**

1039 **proteins.** The wavelength corresponds to the location of the maximum intensity

1040 of the emission fluorescence spectrum of the samples.

hH$_v$1 sample	Emission Maximum (nm)
A18Acd	418
Q56Acd	420
M91Acd	419
V103Acd	417
C107Acd	415
K125Acd	418
Y134Acd	416
F159Acd	418
K169Acd	415
F195 Acd	416
I217Acd	415
F247Acd	418

1041

1042

Supplementary Dataset S1 (separate file). Coordinates of the AlphaFold predicted model of the hH$_v$1 dimer (in PDB format).

Supplementary References

1. J. R. Lakowicz, Ed., *Principles of Fluorescence Spectroscopy* (Springer US, 2006).

2. M. H. Tessmer, S. Stoll, chiLife: An open-source Python package for in silico spin labeling and integrative protein modeling. *PLoS Comput Biol* **19**, e1010834 (2023).