Interrogating the structure and function of the human voltage-gated proton channel (hH$_v$1) with a fluorescence noncanonical amino acid.

Emerson M. Carmona, William N. Zagotta*, Sharona E. Gordon*.

Department of Neurobiology and Biophysics, University of Washington, Seattle, WA, U.S.

Corresponding Authors: William N. Zagotta and Sharona E. Gordon (zagotta@uw.edu, seg@uw.edu).

Author Contributions: All the authors conceived the study and designed the experiments. E.M.C. performed the experiments and analyzed the data. All the authors interpreted the data. E.M.C. wrote the first version of the manuscript. All the authors critically revised the article.

Competing Interest Statement: The authors declare no competing interests.

Abstract

The human voltage-gated proton channel (hH$_v$1) is a dimer of voltage-sensor domains (VSDs) containing highly selective proton permeation pathways in each monomer. In addition to voltage, hH$_v$1 is regulated by other stimuli, including pH gradients, mechanical forces, and ligands such as Zn^{2+}. Aside from the VSDs, this membrane protein contains an N-terminal domain and a C-terminal coiled-coil domain (CC) formed between the monomers. To address the need for direct measurements of conformational rearrangements in hH$_v$1, we developed a Förster resonance energy transfer (FRET) approach to measuring the conformational rearrangements in full-length hH$_v$1 purified from *E. coli*. We used genetic code expansion (GCE) to generate a library of 14 full-length hH$_v$1 constructs, each incorporating the fluorescent noncanonical amino acid acridon-2-ylalanine (Acd) at a different site throughout the various structural domains. Following the expression and purification of these hH$_v$1-Acd proteins, we found that 12 sites yielded stable and functional proton-permeable channels. The fluorescence properties of Acd at each site showed small site-specific differences. Furthermore, we measured site-specific FRET efficiencies from tryptophan (Trp) and tyrosine (Tyr) to Acd in the hH$_v$1-Acd proteins and found results consistent with correct folding in detergent micelles. Notably, the addition of Zn^{2+} produced reversible changes in FRET, with affected residues clustered on the intracellular side of the channel.

Introduction

The human voltage-gated proton channel (hH$_v$1) is a membrane protein that forms a highly selective proton channel (1, 2). Despite its physiological importance, its molecular mechanisms are still poorly understood, including the precise structural identity of the permeation pathway and the gating mechanism of voltage (3, 4), pH gradients (3, 5), membrane stretch (6), and ligands such as the classical inhibitor Zn^{2+} (7, 8). Understanding these processes requires connecting the functional properties to the hH$_v$1 structure.

Each monomer of hH$_v$1 dimers includes an intracellular N-terminal domain of unknown function, a transmembrane voltage-sensor domain (VSD), and an intracellular coiled-coil (CC), which forms the primary intersubunit interface (9, 10). Current structural models, however, have been obtained from truncated or chimeric channel constructs (11, 12), and thus, it is not clear whether they correspond to physiological or functional conformations. Moreover, the multiple kinetic components of H$_v$1 gating currents suggest that the protein's resting state comprises an ensemble of conformations (4, 5), which is consistent with the conformational flexibility inferred from structural approaches (12–14). Because protein function arises from this distribution of conformational states (15), approaches to directly measure structural heterogeneity in the full-length hH$_v$1 are needed.

An attractive approach to interrogate the structural heterogeneity of hH$_v$1 is fluorescence spectroscopy, which offers high sensitivity at nanomolar protein concentrations and compatibility with near-physiological conditions. Fluorescence can report on local environment and accessibility changes, as well as quantify distance changes through Förster resonance energy transfer (FRET) (16). Importantly, time-resolved FRET measured using lifetimes preserves information about conformational heterogeneity on the nanosecond timescale, which is lost in steady-state measurements due to the averaging process (17). Together, these capabilities make fluorescence, especially time-resolved FRET, a powerful method to connect hH$_v$1 structure and function (18).

A few obstacles remain to directly measuring conformational heterogeneity of hH$_v$1. First, a method to express and purify stable and functional protein is needed, which is challenging for membrane proteins. Second, site-specific fluorophore labeling is difficult at buried sites, resulting in low labeling efficiency. Third, traditional fluorophores are bulky and have long linkers (14), resulting in protein structure perturbation and heterogeneity dominated by the properties of the label (19). A recent method was optimized for robust expression and purification of full-length, functional hH$_v$1 in *E. coli* (20, 21). The remaining obstacles can be overcome by labeling the protein with the fluorescent noncanonical amino acid acridon-2-ylalanine (Acd) using genetic code expansion (GCE) (22–24). This technology allows site-specific labeling during translation by reassigning an amber stop codon to an orthogonal aminoacyl-tRNA synthetase (RS)/tRNA (25, 26) (Figure 1A). Acd's small size and minimal linker reduce perturbations and heterogeneity, and its favorable photophysics support both steady-state and time-resolved measurements (27). Here, we incorporated Acd across all hH$_v$1 structural domains and purified 12 of 14 constructs as stable, functional proteins. We then used Acd's environmental sensitivity and FRET from tryptophan (Trp)/tyrosine (Tyr) to Acd to validate the protein folding and report Zn^{2+}-dependent conformational changes. These data establish a platform for interrogating conformational heterogeneity and dynamics in the full-length hH$_v$1.

Results

We generated a library of hH$_v$1 constructs with Acd incorporated across the different structural domains. Because available H$_v$1 structural models were produced from either truncated or modified proteins (11, 12), we used an AlphaFold model of the full-length, dimeric hH$_v$1 to guide Acd site selection (28, 29) (Figure 1B and Dataset S1). The model recapitulates the expected hH$_v$1 architecture with high confidence: S0 as a short helix parallel to the membrane, S1-S4 transmembrane segments forming a canonical VSD, and a C-terminal

helix extending from S4 that forms the CC intersubunit interface. In addition, two helices in the N-terminal domain (P1 and P2) were predicted (Figure 1C), albeit with low confidence. The dimer interface in the membrane is primarily formed by S4-S4 contacts (Figure S1A), although S1-S1 contacts have been observed experimentally (9, 13). This model appears to represent an intermediate state, with the first S4 positive charged residue R208 proximal to D112 in the selectivity filter, and the third S4 positive charged residue R211 proximal to F150 in the charge-transfer center (Figure S1B). Although the model's accuracy and functional state are unknown, we used it as a rough starting point for selecting positions to incorporate Acd.

To test the feasibility and specificity of Acd incorporation in hH$_v$1, we initially selected two positions in the intracellular domains: A18 in the N-terminal domain and F247 in the CC. For each position, the wild-type codon was replaced with an amber stop codon (TAG). Bacteria co-transformed with the Acd RS/tRNA pair and hH$_v$1-TAG constructs expressed fluorescent full-length protein only when Acd was present in the culture medium (Figure 1D). We next generated a library of 14 constructs in which Acd was incorporated into every structural domain: the N-terminal domain, each helix of the VSD, and the CC (Figure 2A-B). Remarkably, most of the 14 hH$_v$1-Acd proteins expressed at levels comparable to the control (No TAG) (Figure 2C and S2). As expected, several hH$_v$1-Acd proteins also produced TAG-truncated products (Figure 2C). Increasing the Acd concentration in the *E. coli* growth medium did not reduce the fraction of truncated protein (Figure S3). One of the constructs, hH$_v$1-C107Acd, showed a weak in-gel fluorescence signal, although the Western blot signal was comparable to that of other constructs.

We successfully purified stable and functional hH$_v$1 labeled with Acd at 12 of the 14 selected positions. We attempted to purify all expressed hH$_v$1-Acd constructs by immobilized metal affinity chromatography using the detergent Anzergent 3-12 (20, 21). For 12 of 14 constructs, purification yielded a single fluorescent band on SDS-PAGE (Figure 3A and S4). In contrast, no purified

protein was detected for the V187Acd or Q233Acd constructs (Figure S4). Because hH$_v$1-Q233Acd was robustly expressed (Figure 2C), its loss during purification suggests reduced stability. The hH$_v$1-Acd yields were construct-dependent (Figure 3B), and the TAG-truncated products did not co-purify with the full-length protein (Figure 3A and S4).

We then evaluated the stability and homogeneity of the hH$_v$1-Acd proteins using fluorescence-detection size exclusion chromatography (FSEC) (30). Similar to the control (Figure S5), most hH$_v$1-Acd proteins showed a main peak (~11.5 mL) followed by a minor peak at higher elution volumes (Figure 3C and S6; Table S1). The proportion of these two peaks varied between constructs (Figure S6). To determine whether the purified hH$_v$1-Acd proteins were functional, we reconstituted the FSEC fractions containing both peaks in asolectin liposomes and assessed their function using the liposome proton flux assay (20, 31). All 12 reconstituted constructs produced ACMA fluorescence quenching upon valinomycin addition, indicating that our purified proteins formed functional proton-permeable channels (Figure 3D and S7). These results highlight the versatility of GCE for incorporating Acd as a fluorescent probe for structural studies of hH$_v$1.

Acd was relatively insensitive to the hydrophobicity of its local environment. To explore how the local environment can alter Acd's spectral properties, we measured the emission spectra of the free amino acid in different solvents (Figure 4A and Table S2). The environmental sensitivity of Acd has been studied previously (24, 27, 32), but we expanded this work here by using the aprotic solvent ethyl acetate (EtAc) and a series of alcohols with varying alkyl chain lengths. Consistent with a general solvent effect produced by polarity (16), the emission spectrum was blue-shifted by approximately 20 nm in EtAc, the least polar solvent measured, relative to water (Figure 4A and Table S2). Alcohols produced a consistently smaller blue shift, independent of their alkyl-chain length (Figure 4A). The spectrum in Buffer-H2, our standard experimental solution containing detergent micelles, was only very slightly blue-shifted relative

to water (Figure 4A), suggesting that Acd did not substantially partition into the hydrophobic core of the detergent micelles. We also tested whether solvent polarity affects the fluorescence lifetime of Acd using time-correlated single photon counting (TCSPC). The fluorescence lifetime of Acd showed a similar trend to the spectral shifts (Figure 4B): EtAc shortened the lifetime markedly, Acd exhibited shorter lifetimes in alcohols than in water, and Acd in Buffer-H2 had a lifetime similar to that measured in water.

The spectral blue-shift and short lifetime of Acd in EtAc suggest that Acd may be able to report on the local environment at different sites in hH_v1, especially those that are surface-exposed versus those that face the lipid core. The emission spectra of the hH_v1-Acd proteins showed small, site-specific shifts (Figure 4C and Table S3). The local environment of Acd in hH_v1 spanned the range between those observed for free Acd in Buffer-H2 and in alcohols, with Q56Acd in the N-terminal intracellular domain and C107Acd in S1 showing the most red-shifted and blue-shifted spectra, respectively (Figure 4C). Consistent with the spectra, the fluorescence lifetimes of the hH_v1-Acd proteins fell between those measured for free Acd in Buffer-H2 and in alcohols, with Q56Acd exhibiting the slowest, and C107Acd the fastest, lifetimes (Figure 4D). The spectral properties of Acd in hH_v1 agree with its location in the structural model, with C107 being the site closest to the detergent micelles' hydrophobic core. The low sensitivity to polarity changes of Acd in different hH_v1 local environments makes it well suited for measuring FRET (27, 33).

FRET between Trp/Tyr and Acd in hH_v1 suggested that the proteins are properly folded. Trp and Tyr have fluorescence emission spectra that overlap with the Acd absorption spectrum and therefore can act as FRET donors to Acd, with relatively short R_0 values (23.5 Å for Trp; 20.9 Å for Tyr) (22) (Figure 5A). Each hH_v1 monomer contains four Trp and four Tyr residues (yellow spheres in Figure 5B). In this structural model, the Trp residues are located in the N-terminal domain (W4, W38, and W45) and midway along transmembrane segment S4 in the VSD (W207). The Tyr residues are located in the N-terminal domain (Y35

and Y42) and near the extracellular end of transmembrane segment S2 (Y134 and Y141).

To quantify Trp/Tyr to Acd FRET, we used spectral FRET analysis (34). The total emission spectrum from Trp/Tyr and Acd was collected (Figure 5C). The Acd emission spectrum was extracted by subtracting a scaled Trp/Tyr spectrum collected from control wild-type protein ($F_{280,NoTAG}$). For each Acd-containing construct, the ratio of the extracted spectrum (F_{280}) to the Acd spectrum with direct excitation (F_{370}) was calculated as *Ratio A* (Figure 5D and S8). Because *Ratio A* is not wavelength-dependent, it reports the linearity of the detectors and the presence of significant contamination from other sources of fluorescence. The *Ratio A* component caused by the direct excitation of Acd (termed *Ratio A_0*) was measured with free Acd (35) (Figure S9). The difference between *Ratio A* and *Ratio A_0* is directly proportional to the FRET efficiency.

As expected, the *Ratio A* values were largely flat across emission wavelengths for our hH$_v$1-Acd proteins (Figure 5E). For all Acd sites, *Ratio A > Ratio A_0* (dashed line in Figure 5E), reflecting FRET between Trp/Tyr and Acd at every site in hH$_v$1. Most *Ratio A* values were distributed between 0.60 and 0.80, with two clear exceptions: A18Acd (blue) and F247Acd (magenta) (Figure 5E). A18Acd showed the highest *Ratio A*, consistent with the multiple Trp/Tyr residues in the N-terminal domain. Conversely, F247Acd showed the lowest *Ratio A*, consistent with its location in the CC, far from most Trp/Tyr residues (Figure 5B).

We next compared *Ratio A – Ratio A_0* values with FRET efficiencies predicted by the AlphaFold structural model. We modeled rotamer ensembles for Trp, Tyr, and our Acd sites in the hH$_v$1 structural model using chiLife (36) and calculated donor-acceptor distance distributions. These distributions were then used to calculate FRET efficiencies (details in SI Text). The experimentally determined *Ratio A – Ratio A_0* values showed a moderate correlation (Pearson's r = 0.54) with the calculated efficiencies (Figure 5F).

There were two spatially clustered deviations between experiment and model. Positions at the extracellular ends of the VSD helices (Y134 in S2, K125

in S1, and F195 in S4) showed higher experimental FRET than predicted, indicating shorter effective donor-acceptor distances relative to the model. Conversely, positions on the intracellular side of the VSD (K169 in S3 and I217 in S4) showed lower experimental FRET than predicted, implying longer effective donor-acceptor distances. Together, the spatial clustering of residues with similar deviations indicates specific deficits in the AlphaFold model and can guide future improvements.

Zinc binding changed the conformation of hH$_v$1. We next asked whether FRET from Trp/Tyr to Acd could report conformational changes upon addition of Zn^{2+}, the classical inhibitor of H$_v$1 proton currents in cells (7). The two experimental structures of H$_v$1 were resolved with this cation bound to the extracellular side of the VSD (11, 12). We observed site-specific spectral changes in the hH$_v$1-Acd proteins in the presence of Zn^{2+} (Figure 6 and S10). The largest changes occurred for hH$_v$1-K169Acd, where 1 mM Zn^{2+} caused a decrease in the Trp/Tyr emission and an increase in the Acd emission (Figure 6A). This decrease in donor fluorescence, paired with increased acceptor fluorescence, is the hallmark of increased FRET and was corroborated by spectral analysis (Figure 6B). Additionally, a slight blue shift of the Trp/Tyr fluorescence was observed for all hH$_v$1-Acd proteins in the presence of Zn^{2+} (Figure S11). These changes were completely reversed by the addition of EDTA, confirming that the changes arise from a reversible conformational change upon Zn^{2+} binding (Figure 6A and 6B). Large changes in effective FRET efficiency were observed with Acd incorporated at Q56 in the N-terminal domain and C107, F159, and K169 in the VSD (Figure 6C). Notably, these positions are intracellular, even though H$_v$1 currents are inhibited by extracellular Zn^{2+} (7, 37). These data demonstrate that Zn^{2+} binds to hH$_v$1 and alters its conformation in

detergent micelles, consistent with a nuclear magnetic resonance study of the VSD of hH$_v$1 (12).

Discussion

Incorporating Acd into hH$_v$1 by GCE opens new possibilities for studying the structure and function of this channel. We systematically incorporated Acd at multiple positions in hH$_v$1 and confirmed protein stability and function after purification. 12 of 14 functional hH$_v$1-Acd proteins produced acceptable yields upon purification, despite the additional challenges associated with recombinant expression of a human membrane protein in *E. coli*. Two factors likely underlie the robustness of our system: (i) an optimized purification protocol for hH$_v$1 (20, 21) that offsets the expected reduction in yield associated with GCE (22, 25), and (ii) the high efficiency of the evolved aminoacyl-tRNA synthetase for incorporating Acd (17, 27).

Fluorescence spectroscopy has been used to detect conformational changes in H$_v$1. Sites at the extracellular portion of S4 (38) and S1 (39) in the VSD of *Ciona intestinalis* H$_v$1 have been labeled with the fluorophores Alexa488 and tetramethylrhodamine (TAMRA) through cysteine-selective reagents to capture conformational changes during activation using voltage-clamp fluorometry. Similarly, the fluorescent noncanonical amino acid 3-(6-acetylnaphthalen-2-ylamino)-2-aminopropanoic acid (Anap) has been incorporated at several sites in the S4 in hH$_v$1 (40) to study conformational changes in response to voltage and pH. In these cases, the fluorophore's environmental sensitivity has been used as a readout of conformational changes. By contrast, FRET provides richer structural information due to its distance dependence. Single-molecule FRET has been measured in purified hH$_v$1 labeled with Cy3 and Cy5 to track the VSD movements in response to voltage, pH, cholesterol, and zinc (14, 41). The small size and minimal linker of Acd, along

with its low environmental sensitivity and high photostability, are advantageous for measuring FRET in hH$_v$1 compared with previously used fluorophores.

The high number of FRET donors, 4 tryptophans and 4 tyrosines per subunit, limits the utility of the Trp/Tyr-Acd FRET pair as a strategy to measure conformational rearrangements in hH$_v$1. Our library of a dozen single Acd-incorporating constructs, with sites distributed across the entire primary sequence of the protein, opens the door to using Acd as a FRET donor. We have previously used time-resolved transition metal ion FRET (tmFRET), which utilizes a transition metal ion as a FRET acceptor (27, 33), to measure conformational energetics in the bacterial cyclic nucleotide-gated ion channel SthK (42). However, in that work, we incorporated Acd at a single position in the intracellular C-terminal domain of SthK. With Acd incorporated at multiple sites in full-length hH$_v$1, we can interrogate its gating mechanisms in unprecedented detail.

Materials and Methods

Protein expression and purification. hH$_v$1 was expressed and purified according to the published protocol with minor modifications (20, 21). Briefly, fresh competent BL21-Gold(DE3) cells (Agilent) were co-transformed with the His-EK-hH$_v$1-C107A-C249A.pET15-b (hH$_v$1-Cysless, referred to as No TAG) and MjA9Acd-RS.pDule2 (22, 23) plasmids (sequences in SI Text) using the heat shock method. For expressing hH$_v$1-Acd proteins, the codon corresponding to the selected position was changed to the amber stop codon (TAG) by site-directed mutagenesis. All plasmids were sequenced before use. Pre-cultures were grown overnight in LB medium supplemented with 0.2% glucose, 0.4 mg/mL ampicillin, and 0.1 mg/mL spectinomycin at 37 °C and 250 rpm. The next day, cultures were diluted 1:100 in complete autoinduction medium (43) supplemented with 0.4 mg/mL ampicillin and 0.1 mg/mL spectinomycin, and then grown at 37 °C and 250 rpm until an OD$_{600}$ of 1.5 was reached. The protein expression level was proportional to the Acd concentration in the culture medium (Figure S11). We chose 0.6 mM Acd in the culture medium as optimal for hH$_v$1

expression, as higher concentrations only slightly increased the final biomass (Figure S11B). Cultures supplemented with 0.6 mM Acd were grown overnight at 20 °C and 250 rpm. Cells were then harvested, resuspended in Buffer-H1 (50 mM Tris, 150 mM NaCl, 1 mM benzamidine, 0.17 mg/mL PMSF, pH 8.0) supplemented with 0.5 mg/mL lysozyme, and incubated for 30 min at 4 °C before being flash-frozen. After thawing, the extracts were supplemented with 5 mM $MgCl_2$, fresh protease inhibitors, and 12.5 μg/mL DNase I, incubated for 1 h at 4 °C, sonicated, and centrifuged at 100,000g at 4 °C for 1 h. The membrane pellet was resuspended in Buffer-H1 and stored at -80 °C until further use. After thawing, the membranes were solubilized with 1.5% Anzergent 3-12 (Anatrace) (Anz3-12) at room temperature for 1 hour. Insoluble material was removed by centrifuging at 100,000g and 4 °C for 1 h, and the solubilized membrane extract was incubated with His-Pur Ni-NTA (Thermo Scientific) resin equilibrated with Buffer-H2 (50 mM Tris, 150 mM NaCl, 12 mM Anz3-12, pH 8.0) for 1 hour at room temperature. Resin was collected in a gravity column and washed with 20 column volumes (CV) of Buffer-H2, followed by 16 CV of Buffer-H2 with 90 mM imidazole. The hH_v1 protein was eluted with 20 CV of Buffer-H2 with 0.5 M imidazole, was concentrated with 50kDa cut-off centrifugal filters, and imidazole was removed by FSEC (Ex/Em = 385/450 nm) in an ENrich SEC 650 (Bio-Rad) column with Buffer-H2. The hH_v1-containing fractions were concentrated to 0.5-1.5 mg/mL, depending on the amount of purified protein, and then aliquoted, flash-frozen, and stored at -80 °C until use.

Reconstitution and fluorescence proton flux assays. The hH_v1-Acd samples were reconstituted and assayed using a liposome proton flux assay, as described previously, with minor modifications (20). Briefly, soy polar lipid extract (Avati) in chloroform was dried overnight at room temperature in a rotary evaporator under vacuum and then hydrated in Buffer-K (20 mM HEPES, 150 mM KCl, and 1 mM EDTA, pH 7.0) to a concentration of 10 mg/mL. The vesicles were sonicated in a water bath until the solution became translucent, aliquoted, flash-frozen, and stored at -80 °C until further use. Reconstitutions were performed at a 1:100

(protein:lipid, by mass) ratio by mixing 12.5 μg of purified hH$_v$1-Acd with 1.25 mg of liposomes in a final volume of 330 μL Buffer-K containing 8 mM Anz3-12. The mixture was incubated at room temperature for 1 h, then diluted with 5 mL of Buffer-K and incubated for 30 min. The mixture was further incubated with three cumulative pulses of 100 mg Bio-Beads SM2 (Bio-Rad), each for 1 h at room temperature. After the final Bio-Beads pulse, the mixture was incubated overnight at 4 °C. The next day, two successive 100 mg additions of Bio-Beads were performed, with a 1-hour incubation at room temperature following each addition. Bio-Beads were removed in a gravity column, the mixture was diluted with 10 mL of Buffer-K, and then centrifuged at 150,000g for 2 hours at 4 °C. The proteoliposome pellets were resuspended in 250 μL of Buffer-K, aliquoted, flash-frozen, and stored at -80 °C until further use. On the day of the liposome proton flux assays, samples were thawed at 37 °C and then placed on ice. The sample (40 μL) was diluted in Buffer-Na (20 mM HEPES, 150 mM NaCl, 1 mM EDTA, pH 7.0). 9-Amino-6-chloro-2-methoxyacridine (ACMA; Sigma) was added to a final concentration of 2 μM from a 2 mM stock solution in DMSO, and the mixture was incubated at room temperature for 5 minutes before the fluorescence measurement began. The baseline was recorded for 3 min (F_{max}). Subsequently, 10 nM of valinomycin (Cell Signaling) was added from a 10 μM stock solution in DMSO, followed by 1 μM carbonyl cyanide m-chlorophenyl hydrazone (CCCP; Sigma) from a 1 mM stock solution in DMSO to record F_{min}. The final volume was 2 mL. The fluorescence signal ($F(t)$) was normalized according to the equation ($F(t)$-F_{min})/(F_{max}-F_{min}). Fluorescence measurements were performed at 5-second intervals with a 2-second integration time using a Fluorolog-3 spectrometer (Horiba) configured to an excitation wavelength of 410 nm (5 nm bandpass) and an emission wavelength of 490 nm (1 nm bandpass) at room temperature.

Absorption and fluorescence spectra recordings. The absorption and fluorescence spectra were recorded in the indicated solvents or fresh Buffer-H2 at room temperature. Absorption spectra were recorded in a DU 800 Spectrometer (Beckman Coulter) with a wavelength interval of 0.5 nm and a scan

speed of 600 nm/min. Fluorescence emission spectra were measured using a Fluorolog-3 spectrometer (Horiba) with an integration time of 0.1 s, 1 nm increments, and excitation and emission slits of 5 nm. For solubilizing Acd in different solvents, a saturated solution of the amino acid was prepared by dissolving 0.5 mg of Acd powder in the respective solvent. Acd spectra were corrected by subtracting a blank of the solvent. hH$_v$1-Acd proteins were measured at concentrations ranging from 180 to 240 nM in Buffer-H2. Each spectrum was corrected by subtracting a blank with the solvent or Buffer-H2 before adding the protein sample to the cuvette.

Spectral FRET analysis. We used our previously established spectral FRET analysis to remove contamination caused by direct excitation of Acd by 280 nm light (35). This method had the added benefit of eliminating errors arising from the recording system's transfer function, variations in the acceptor's quantum yield, or variations in the total concentration of fluorescent molecules. A Trp/Tyr spectrum was collected from control protein without any TAG codons (no Acd incorporation). This was used to subtract the Trp/Tyr emission spectra collected at 280 nm for each Acd-incorporating protein. This yielded the extracted Acd spectrum, F_{280}, that had two components: the component due to direct excitation of Acd, F_{280}^{direct}, and the component due to FRET, F_{280}^{FRET}. F_{280} was normalized by the total Acd emission with 370 nm excitation, F_{370}. The resulting ratio, termed *Ratio A*, can be expressed as: $Ratio\ A = \frac{F_{280}}{F_{370}} = \frac{F_{280}^{direct}}{F_{370}} + \frac{F_{280}^{FRET}}{F_{370}}$. The direct excitation component, $\frac{F_{280}^{direct}}{F_{370}}$, termed *Ratio A$_0$*, was measured using free Acd amino acid in Buffer-H2. We then quantified the relative FRET efficiency as $Ratio\ A - Ratio\ A_0 = \frac{F_{280}^{FRET}}{F_{370}}$. This quantity is directly proportional to FRET efficiency. An average *Ratio A* value was calculated between 410 and 480 nm.

Time-resolved fluorescence measurements. TCSPC measurements were performed in a FluoTime 300 spectrometer equipped with a PMA Hybrid 40

detector and LDH-P-C-375 laser diode head (PicoQuant) at room temperature. Emitted photons were detected at 446 nm with the emission polarizer at the magic angle. The TCSPC measurements were performed in the indicated solvent or fresh Buffer-H2. The hH$_v$1-Acd samples were measured at a concentration of 1 μM and the Acd amino acid at 500 nM. The count rate was always less than 1% of the excitation repetition rate to avoid pile-up distortions of the fluorescence decay. The IRF was measured using a dilute Ludox solution in water.

Acknowledgments

We thank the Oregon State University GCE4ALL (Center for Genetic Code Expansion for All) for their long-standing collaboration, Dr. James Petersson, and Kyle D. Shaffer (University of Pennsylvania) for excellent technical support with the Acd synthesis. This work was supported by the National Institutes of Health under award numbers R01EY037223 and R35GM145225 to S.E.G., and R01EY010329 and R35GM148137 to W.N.Z. E.M.C. is a Pew Latin American Fellow.

References

1. M. Sasaki, M. Takagi, Y. Okamura, A Voltage Sensor-Domain Protein Is a Voltage-Gated Proton Channel. *Science* **312**, 589–592 (2006).

2. I. S. Ramsey, M. M. Moran, J. A. Chong, D. E. Clapham, A voltage-gated proton-selective channel lacking the pore domain. *Nature* **440**, 1213–1216 (2006).

3. V. V. Cherny, V. S. Markin, T. E. DeCoursey, The voltage-activated hydrogen ion conductance in rat alveolar epithelial cells is determined by the pH gradient. *The Journal of General Physiology* **105**, 861–896 (1995).

4. E. M. Carmona, *et al.*, Gating charge displacement in a monomeric voltage-gated proton (H$_v$1) channel. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 9240–9245 (2018).

5. E. M. Carmona, *et al.*, The voltage sensor is responsible for ΔpH dependence in H_v1 channels. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2025556118 (2021).

6. M. M. Pathak, *et al.*, The Hv1 proton channel responds to mechanical stimuli. *Journal of General Physiology* **148**, 405–418 (2016).

7. V. V. Cherny, T. E. DeCoursey, pH-Dependent Inhibition of Voltage-Gated H^+ Currents in Rat Alveolar Epithelial Cells by Zn^{2+} and Other Divalent Cations. *The Journal of General Physiology* **114**, 819–838 (1999).

8. T. E. DeCoursey, V. V. Cherny, Voltage-activated hydrogen ion currents. *J. Membarin Biol.* **141** (1994).

9. S.-Y. Lee, J. A. Letts, R. MacKinnon, Dimeric subunit stoichiometry of the human voltage-dependent proton channel H_v1. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 7692–7695 (2008).

10. H. P. Koch, *et al.*, Multimeric nature of voltage-gated proton channels. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 9111–9116 (2008).

11. K. Takeshita, *et al.*, X-ray crystal structure of voltage-gated proton channel. *Nat Struct Mol Biol* **21**, 352–357 (2014).

12. M. Bayrhuber, *et al.*, Nuclear Magnetic Resonance Solution Structure and Functional Behavior of the Human Proton Channel. *Biochemistry* **58**, 4017–4027 (2019).

13. Q. Li, *et al.*, Resting state of the human proton channel dimer in a lipid bilayer. *Proc. Natl. Acad. Sci. U.S.A.* **112** (2015).

14. S. Han, *et al.*, Structural dynamics determine voltage and pH gating in human voltage-gated proton channel. *eLife* **11**, e73093 (2022).

15. K. Henzler-Wildman, D. Kern, Dynamic personalities of proteins. *Nature* **450**, 964–972 (2007).

16. J. R. Lakowicz, Ed., *Principles of Fluorescence Spectroscopy* (Springer US, 2006).

17. S. E. Gordon, M. Munari, W. N. Zagotta, Visualizing conformational dynamics of proteins in solution and at the cell membrane. *eLife* **7**, e37248 (2018).

18. W. N. Zagotta, *et al.*, Measuring conformational equilibria in allosteric proteins with time-resolved tmFRET. *Biophysical Journal* **123**, 2050–2062 (2024).

19. J. W. Taraska, M. C. Puljung, W. N. Zagotta, Short-distance probes for protein backbone structure based on energy transfer between bimane and transition metal ions. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 16227–16232 (2009).

20. E. M. Carmona, D. M. Cortes, L. G. Cuello, A novel method for expressing and purifying large quantities of functional and stable human voltage-gated proton channel (нHᵥ1). *Protein Science* **34**, e70017 (2025).

21. E. Carmona, M. Cortes, L. Cuello, Expression and Purification of the Human Voltage-Gated Proton Channel (hHᵥ1). *BIO-PROTOCOL* **15** (2025).

22. L. C. Speight, *et al.*, Efficient Synthesis and In Vivo Incorporation of Acridon-2-ylalanine, a Fluorescent Amino Acid for Lifetime and Förster Resonance Energy Transfer/Luminescence Resonance Energy Transfer Studies. *J. Am. Chem. Soc.* **135**, 18806–18814 (2013).

23. I. Sungwienwong, *et al.*, Improving target amino acid selectivity in a permissive aminoacyl tRNA synthetase through counter-selection. *Org. Biomol. Chem.* **15**, 3603–3610 (2017).

24. C. M. Jones, *et al.*, Genetic encoding of a highly photostable, long lifetime fluorescent amino acid for imaging in mammalian cells. *Chem. Sci.* **12**, 11955–11964 (2021).

25. C. C. Liu, P. G. Schultz, Adding New Chemistries to the Genetic Code. *Annu. Rev. Biochem.* **79**, 413–444 (2010).

26. T. S. Young, I. Ahmad, J. A. Yin, P. G. Schultz, An Enhanced System for Unnatural Amino Acid Mutagenesis in E. coli. *Journal of Molecular Biology* **395**, 361–374 (2010).

27. W. N. Zagotta, *et al.*, An improved fluorescent noncanonical amino acid for measuring conformational distributions using time-resolved transition metal ion FRET. *eLife* **10**, e70236 (2021).

28. J. Jumper, *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

29. G. Kim, *et al.*, Easy and accurate protein structure prediction using ColabFold. *Nat Protoc* **20**, 620–642 (2025).

30. T. Kawate, E. Gouaux, Fluorescence-Detection Size-Exclusion Chromatography for Precrystallization Screening of Integral Membrane Proteins. *Structure* **14**, 673–681 (2006).

31. S.-Y. Lee, J. A. Letts, R. MacKinnon, Functional Reconstitution of Purified Human H_v1 H^+ Channels. *Journal of Molecular Biology* **387**, 1055–1060 (2009).

32. A. Szymańska, K. Wegner, L. Łankiewicz, Synthesis of *N*-[(*tert*-Butoxy)carbonyl]-3-(9,10-dihydro-9-oxoacridin-2-yl)-L-alanine, a New Fluorescent Amino Acid Derivative. *Helvetica Chimica Acta* **86**, 3326–3331 (2003).

33. S. E. Gordon, *et al.*, Long-distance tmFRET using bipyridyl- and phenanthroline-based ligands. *Biophysical Journal* **123**, 2063–2075 (2024).

34. R. M. Clegg, "[18] Fluorescence resonance energy transfer and nucleic acids" in *Methods in Enzymology*, (Elsevier, 1992), pp. 353–388.

35. J. Zheng, M. D. Varnum, W. N. Zagotta, Disruption of an Intersubunit Interaction Underlies Ca^{2+}-Calmodulin Modulation of Cyclic Nucleotide-Gated Channels. *J. Neurosci.* **23**, 8167–8175 (2003).

36. M. H. Tessmer, S. Stoll, chiLife: An open-source Python package for in silico spin labeling and integrative protein modeling. *PLoS Comput Biol* **19**, e1010834 (2023).

37. F. Qiu, *et al.*, Molecular mechanism of Zn^{2+} inhibition of a voltage-gated proton channel. *Proc. Natl. Acad. Sci. U.S.A.* **113** (2016).

38. C. Gonzalez, H. P. Koch, B. M. Drum, H. P. Larsson, Strong cooperativity between subunits in voltage-gated proton channels. *Nat Struct Mol Biol* **17**, 51–56 (2010).

39. L. Mony, T. K. Berger, E. Y. Isacoff, A specialized molecular motion opens the Hv1 voltage-gated proton channel. *Nat Struct Mol Biol* **22**, 283–290 (2015).

40. E. Suárez-Delgado, M. Orozco-Contreras, G. E. Rangel-Yescas, L. D. Islas, Activation-pathway transitions in human voltage-gated proton channels revealed by a non-canonical fluorescent amino acid. *eLife* **12**, e85836 (2023).

41. S. Han, *et al.*, Cholesterol inhibits human voltage-gated proton channel hHv1. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2205420119 (2022).

42. P. Eggan, S. E. Gordon, W. N. Zagotta, Domain Coupling in Allosteric Regulation of SthK Measured Using Time-Resolved Transition Metal Ion FRET. [Preprint] (2025). Available at: https://elifesciences.org/reviewed-preprints/106892v2 [Accessed 2 August 2025].

43. F. W. Studier, Protein production by auto-induction in high-density shaking cultures. *Protein Expression and Purification* **41**, 207–234 (2005).

Figures

Figure 1. hH$_v$1 was labeled with the fluorescent noncanonical amino acid Acd by genetic code expansion. (A) Cartoon showing the components of the genetic code expansion system used to incorporate Acd via amber codon suppression in hH$_v$1. (B) AlphaFold structural model of the dimer hH$_v$1 colored by subunit. (C) AlphaFold structural model of a single hH$_v$1 subunit colored by the predicted Local Distance Difference Test (pLDDT). Model confidence was represented by a gradient from red to blue, indicating low to high pLDDT. (D) In-gel fluorescence and Western blot to evaluate Acd incorporation. An Acd fluorescence band corresponding to the molecular weight of full-length hH$_v$1 (red arrow) was observed in cellular extracts after separation by SDS-PAGE only when cells contained the *Mj*A9-AcdRS/tRNA plasmid (RS) and were grown in the presence of 1mM Acd (top). The Western blot against the His-tag present at the N-terminus of hH$_v$1 confirmed that the band corresponded to the recombinant protein (bottom). Note the hH$_v$1 truncated product produced when the amber stop

codon is replacing F247 in the absence of the aminoacyl-tRNA synthetase/tRNA plasmid and Acd.

Figure 2. Acd was incorporated into 14 positions in the hH$_v$1 sequence. (A) Cartoon showing the amino acids selected to be replaced by an amber stop codon in the hH$_v$1 secondary structure to incorporate Acd (stars). (B) AlphaFold dimer hH$_v$1 structural model with the amino acids selected to be replaced by Acd as cyan spheres. (C) Acd fluorescence gel (top) and Western blot (bottom) of cellular extracts after separation by SDS-PAGE showing the expression of the Acd-labeled hH$_v$1 mutants. The No TAG lane contains an extract from cells grown under identical conditions (in the presence of Acd and the aminoacyl-tRNA synthetase/tRNA pair) and expressing hH$_v$1 without an amber stop codon.

Figure 3. Purification and functional measurements of hH$_v$1-Acd proteins.
(A) Coomassie-stained (top) and Acd fluorescence (bottom) gels after separation by SDS-PAGE showing representative samples of the membrane extracts after solubilization (left) and the final purified protein after immobilized metal affinity chromatography (right). (B) Protein yield of the hH$_v$1 protein with Acd replacing the amino acid at the indicated position. Note that V187Acd and Q233Acd contained no detectable protein after purification. The red line indicates the protein yield obtained from hH$_v$1 without an amber stop codon in the presence of Acd and the aminoacyl-tRNA synthetase/tRNA pair (No TAG). (C) Representative fluorescence-detection size-exclusion chromatograms of the purified hH$_v$1 proteins with Acd incorporated at the indicated amino acid position. Insets: AlphaFold dimer models, with the amino acid replaced by Acd shown as cyan spheres. (D) Representative liposome proton flux assay of asolectin proteoliposomes containing the indicated hH$_v$1 protein. All purified proteins produced ACMA fluorescence quenching after the addition of valinomycin (Val).

The protonophore CCCP was added at the end of the experiment as a control. The No TAG sample corresponds to proteoliposomes containing hH$_v$1 without an amber stop codon expressed in the presence of Acd and the aminoacyl-tRNA synthetase/tRNA pair.

Figure 4. Acd showed a low environmental sensitivity in hH$_v$1. (A) Normalized fluorescence emission spectra of the Acd amino acid dissolved in the indicated solvent (Excitation = 370 nm). EtAc: ethyl acetate, OctOH: 1-octanol, ButOH: 1-butanol, EtOH: ethanol. (B) Normalized fluorescence decay of the Acd amino acid dissolved in the indicated solvent. (C) Normalized fluorescence emission spectra of the hH$_v$1-Acd proteins in Buffer-H2 (grey). The Acd amino acid spectra in selected solvents are also shown for reference. Q56Acd and C107Acd spectra are shown with broken lines. The emission maxima are listed in Table S3. (D) Normalized fluorescence decay of the hH$_v$1-Acd proteins in Buffer-H2 (grey). The Acd amino acid decays in selected solvents are also shown for reference. Q56Acd and C107Acd decays are shown in red and black, respectively.

Figure 5. Spectral FRET analysis using Trp and Tyr donors and Acd as acceptor. (A) Fluorescence emission spectra of the intrinsic fluorescent amino acids Trp (black) and Tyr (black, broken lines), along with the Acd absorption spectrum in Buffer-H2 (blue). (B) AlphaFold dimer hH_v1 structural model with the amino acids colored according to (C) and the native tryptophan and tyrosine residues shown as yellow spheres. (C) Normalized fluorescence spectra of the purified hH_v1-Acd proteins (Excitation = 280 nm). The spectra were normalized by the Trp/Tyr fluorescence. The colors correspond to the indicated hH_v1-Acd protein, and the remaining spectra are shown as grey traces in the figure. The No TAG sample (black) contains purified hH_v1 protein without any amber stop codon expressed in the presence of Acd and the aminoacyl-tRNA synthetase/tRNA pair. (D) Example of the spectral FRET analysis procedure. The fluorescence spectrum of hH_v1-Y134Acd obtained when exciting at 280 nm (black, F_{280}) minus the normalized spectrum at the same excitation wavelength of the No TAG sample (grey, $F_{280,NoTAG}$) produced the Acd emission spectrum shown in red ($F_{280} - F_{280,NoTAG}$). *Ratio A* values were calculated by dividing this spectrum by the hH_v1-Y134Acd fluorescence spectrum obtained by direct excitation at 370 nm (F_{370}). (E) *Ratio A* traces of the hH_v1-Acd proteins colored according to (C). The ratio between the spectra of free Acd amino acid excited at 280 and 370 nm

(*Ratio A$_0$*) is shown as broken lines. (F) Average *Ratio A – Ratio A$_0$* values (410-480 nm) as a function of the predicted FRET efficiencies from the AlphaFold hH$_v$1 structural model, colored according to (C). The broken line is the best fit to the equation (*Ratio A – Ratio A$_0$*) = m(Predicted FRET efficiency).

Figure 6. FRET between Trp/Tyr and Acd reports a conformational rearrangement in response to Zn^{2+}. (A) Normalized fluorescence emission spectra of hH$_v$1-K169Acd (Excitation = 280 nm) in Buffer-H2 in the absence of Zn^{2+} (black, Apo), in the presence of 1 mM Zn^{2+} (red), or in the presence of 1 mM Zn^{2+} and 2 mM EDTA (blue). Spectra were normalized by the maximum intensity of the Acd emission spectrum of the same sample excited at 370 nm. (B) Spectral FRET analysis of hH$_v$1-K169Acd in Buffer-H2 in the absence of Zn^{2+} (black, Apo), in the presence of 1 mM Zn^{2+} (red), or in the presence of 1 mM Zn^{2+} and 2 mM EDTA (blue). (C) Spectral FRET analysis of the hH$_v$1-Acd proteins in Buffer-H2 in the absence (black, Apo) or presence of 1 mM Zn^{2+} (red).