Global shifts in vegetation compositional resilience over the past 8,000 years

Mengna Liao[1,*], Kai Li[1,*], Chenzhi Li[2,3], Ulrike Herzschuh[3], Jian Ni[1]

[1] College of Life Sciences, Zhejiang Normal University, Jinhua 321004, China
[2] State Key Laboratory of Lake and Watershed Science for Water Security, Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, Nanjing 211135, China
[3] Polar Terrestrial Environmental Systems, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Potsdam 14473, Germany

*Corresponding authors. mnliao@zjnu.edu.cn (ML), likai@zjnu.edu.cn (KL)

Abstract

Vegetation resilience is critical for sustaining biodiversity and ecosystem services, yet its

long-term dynamics and underlying drivers remain poorly resolved. Here, we assess changes

in continental-scale vegetation compositional resilience over the past 8,000 years by applying

established leading indicators (including autocorrelation, standard deviation, skewness, and

kurtosis) to 482 fossil pollen records from all major landmasses except Antarctica. Using a

machine learning approach, we evaluate the relative contributions of anthropogenic land

cover change (ALCC) and climatic parameters (annual mean temperature, temperature

seasonality, annual precipitation, and precipitation seasonality) in driving resilience shifts

across continents. We further employ structural equation modeling (SEM) to disentangle how

abiotic factors (ALCC and climatic variables) and biotic factors (species richness, evenness,

synchrony, and temporal β diversity) jointly shape long-term trends in vegetation resilience at

the continental scale. Our results reveal a persistent, millennia-scale decline in resilience

during the past 1.6 to 4.4 thousand years before present, with intensified anthropogenic

land-use change identified as the dominant global driver. This finding suggests that the global

decline in vegetation resilience observed over recent decades likely represents a continuation

of the long-term trend established during the preceding Holocene. Notably, North America

exhibited a resurgence in resilience over the past 1,200 years, primarily attributed to enhanced

resilience in tundra and savanna ecosystems. SEM analyses uncover complex pathways

through which abiotic and biotic predictors interact to shape millennial-scale resilience

changes across continents. Despite the complexity of these underlying mechanisms, the

analyses demonstrate that biotic factors collectively exert substantial, and often predominant,

influences on long-term resilience dynamics. This underscores the indispensable role of biotic

attributes in modulating long-term vegetation resilience, highlighting their potential as critical

targets for ecosystem conservation and restoration.

Keywords: pollen analysis, early warning signals, vegetation resilience, anthropogenic

land-use, climate change, diversity, synchrony, multi-millennial timescale

Introduction

Ongoing global changes have profoundly impacted Earth's ecosystems, forcing abruptly and irreversibly regime shifts in some ecosystems (Cavanaugh et al., 2019; Smol et al., 2005; Wernberg et al., 2016). At the planetary scale, evidence suggests that the biosphere as a whole may be approaching a catastrophic bifurcation point (a 'tipping point') due to sustained loss of resilience (Barnosky et al., 2012). Given that current rates of change in key Earth System features already match or exceed those of abrupt geophysical events in the past (Steffen et al., 2018), assessing the global-scale resilience of Earth's ecosystems has become an urgent priority for developing adaptive management strategies.

Resilience was defined as the capacity of a system to resist disturbance, recover to its original state after a disturbance, and sustain fundamental functions at pre-disturbance levels (Folke et al., 2004). It represents two stability paradigms, namely local stability and non-local stability (Dakos & Kéfi, 2022). The local stability, also termed engineering resilience, typically refers to an ecosystem's capacity to return to its original state following a small disturbance when near a stable equilibrium state (Pimm, 1984). The non-local stability, widely known as ecological resilience, can evaluate the magnitude of disturbance that can be absorbed before the system changes state (Holling, 1973). Fundamentally, engineering resilience focuses on preserving the recovery rate of a structure and/or function, while ecological resilience emphasizes maintaining the existence of the structure and/or function (Holling, 1996). Over the past two decades, ecologists proposed generic early warning signals (or leading indicators) to bridging the gap between the intuitive conceptual clarity of ecosystem resilience and its operational, quantifiable metrics (Carpenter & Brock, 2006; Dakos & Kéfi, 2022; Guttal & Jayaprakash, 2008; Scheffer et al., 2009; Wang et al., 2012). Resilience reduction, which is often associated with phenomena recognized in multi-stable ecosystems as "critical slowing down" or "flicking" can be quantitatively characterized by increases in leading indicators such as variance, autocorrelation, skewness, and kurtosis of system state variables (Dakos et al., 2008; Dakos et al., 2013; Scheffer et al., 2009; Wang et al., 2012). These phenomena can theoretically be assessed within any temporally and spatially bounded system (Barnosky et al., 2012). As different individual metric may have different predictive ability, the combination of several leading indicators into a single composite metric could enable more reliable inference of resilience changes (Drake & Griffen, 2010).

Vegetation unquestionably forms the most crucial component of terrestrial ecosystems, which provides essential ecosystem services while also maintaining the balance of the Earth System (Duveiller et al., 2018). Recent studies, employing either individual leading indicators or composite resilience indices, have demonstrated a widespread decline in vegetation resilience across many regions over the past decades (Boulton et al., 2022; Feng et al., 2021; Forzieri et al., 2022; Smith et al., 2022), based on satellite-derived vegetation metrics such as Vegetation Optical Depth (VOD) and Normalized Difference Vegetation Index (NDVI).

While these metrics serve as indicators of gross primary production (Liu et al., 2015) and vegetation greenness or canopy coverage (Myneni et al., 1997), they poorly reflect vegetation composition, which is a critical dimension for understanding shifts in ecosystem structure and function disturbed by climate change and human activities. Moreover, satellite-derived vegetation metrics are limited to a temporal coverage of only several decades, which is exceedingly brief when compared to the extensive history of human-induced landscape alterations and the long-term process of vegetation succession. In addition, vegetation's response to climate change manifests across a range of timescales from hundreds to tens of thousands of years, potentially showing non-linear characteristics or demonstrating temporal lags (Fastovich et al., 2025). Therefore, the limitations in data observation post significant challenges for empirical data in detecting critical shifts of vegetation composition resilience and attributing these shifts.

Fossil pollen records offer long-term perspectives for inferring past plant community properties and their dynamics across local, regional, and global scales spanning the last few centuries to millions of years (Birks, 2019; Jackson & Blois, 2015). By compiling pollen data, recent studies have revealed that vegetation composition has been transformed globally by climate changes and increasing human disturbances since the last glacial maximum (Gordon et al., 2024; Nolan et al., 2018). Moreover, the rates of vegetation compositional change accelerated unprecedently during the late Holocene in both magnitude and extent (Mottl et al., 2021). Given the profound transformations of vegetation ecosystems caused by climate changes and, particularly, human disturbances, large-scale vegetation resilience could have undergone significantly shifts during the Holocene. However, as of now, there is no evidence to support this hypothesis, nor has it been confirmed whether human disturbances play the crucial role in driving continental or global changes in vegetation compositional resilience.

Ecosystems not only make direct responses to external forcings but also possess inherent ecological feedbacks that enable them to mediate responses to exogenous stress, thereby sustaining the resilience of ecosystems (Blois et al., 2013; Folke et al., 2004). The stabilizing effect of biodiversity (generally measured by species richness) has received the most extensive attention and discussion (Ives & Carpenter, 2007; Loreau & De Mazancourt, 2013). Most biodiversity–stability relationships involve some forms of compensatory dynamics or insurance effect, which require either asynchrony of species responses to environmental fluctuation or the presence of functionally redundant species with differing sensitivities to specific perturbations (de Bello et al., 2021; Loreau & De Mazancourt, 2013; Oliver et al., 2015). While this certainly appears to be the case at the community level in some ecosystems over relatively short ecological timescales spanning years to no longer than decades (Hatton et al., 2024; Hisano et al., 2018; Isbell et al., 2015), these patterns may not generalize to larger spatial scales or longer temporal scales (Peterson et al., 1998; Wang & Loreau, 2014). Other biological attributes, such as species evenness and spatial self-organization, regulate ecosystem resilience as well (Kéfi et al., 2024; Valencia et al., 2020). While, the applicability

of these stability mechanisms to explaining long-term resilience trajectories in vegetation

composition remains uncertain. This knowledge gap underscores an urgent need to extend

current knowledge to broader spatial extents and longer temporal baselines, thereby aligning

research scales with the operational dimensions of global biodiversity management and

conservation strategies.

Here, we assessed continental-scale vegetation resilience over the past 8,000 years using

fossil pollen sequences from the recently published global late Quaternary pollen dataset (Li

et al., 2022). A total of 482 pollen records (Figure 1) were selected from all continents, with

the exception of Antarctica. We calculated a composite resilience index (CRI) for each record

by integrating four leading indicators (*i.e.*, autocorrelation, standard deviation, skewness, and

kurtosis). The estimated CRI sequences were aggregated by continent, and each continental

CRI dataset was fitted using generalized additive models (GAMs) to reconstruct long-term

trajectories of vegetation compositional resilience. We then evaluated the relative importance

of abiotic (including climate variables and land-use change) and biotic factors (including

diversity and synchrony) to vegetation resilience, as well as their direct and indirect effects,

employing a machine learning approach and piecewise structural equation model (SEM).

With this study, we aim to provide a long-term perspective on trends in vegetation resilience

across continental to global scales and provide new insight into the mechanisms underlying

long-term vegetation resilience dynamics.

Figure 1. Spatial distribution of the fossil pollen sequences analyzed in this study, with

sequence counts by continent and biome. (A) Spatial distribution of pollen sequences. (B) The

number of pollen sequences per continent [symbols match those in (A)]. (C) The number of

pollen sequences per biome [colors match those in (A)].

Methods

Fossil pollen data source

Fossil pollen data were obtained from the LegacyPollen 2.0 dataset, a globally harmonized

compilation of 3680 palynological records (Li et al., 2024, 2025). To ensure analytical

consistency across diverse records, the dataset underwent rigorous taxonomic

standardization—woody and dominant herbaceous taxa were harmonized at the genus level,

while other herbaceous taxa were standardized at the family level (Herzschuh et al., 2022).

Temporal standardization was achieved by constructing robust Bayesian age–depth models

(Blaauw & Christen, 2011; Li et al., 2022). Each record contains both percentage and count

data for identified taxa, along with calibrated age estimates corresponding to sediment depth.

Data filtering

We applied several criteria for data filtering, including archive type, the youngest and oldest

depositional ages of records, the mean temporal resolution of samples, the total numbers of

chronological control points, the presence or absence of sedimentary hiatus, and the temporal

gap between adjacent samples. Firstly, we excluded records collected from marine, ice, and

animal midden, as these archives rarely allow clear defining of the geographical source of

fossil pollen. Regarding the depositional ages, we retained records with a minimal age no

older than 2 cal ka BP and a basal age no younger than 10 cal ka BP. The adoption of this

criterion ensures that the timeseries of the calculated vegetation resilience (see below for how

CRI was calculated) spans at least from the mid- to the late Holocene. We used a mean

temporal resolution of 200 years as another data filtering criterion, based on a comprehensive

assessment of both data availability and the capacity of pollen data to well resolve long-term

vegetation dynamics over multi-millennial timescales. To minimize uncertainties arising from

potential bias in estimated depositional ages, we excluded records containing fewer than three

independent chronological control points from the analysis. The presence of pronounced

sedimentary hiatus or large chronological gaps between adjacent samples in pollen records

may result in either misinterpretation or failure to reconstruct vegetation dynamics accurately.

Given this, we calculated the chronological gaps between each pair of adjacent samples in all

records that met the aforementioned criteria. If a record contains a chronological gap

exceeding 2,000 years, only its upper section is retained for subsequent analysis, contingent

upon this section meeting the aforementioned criteria. Ultimately, we identified 482 pollen

sequences that met all the filtering criteria, including a total of 56,939 pollen samples. Among

these sequences, 60.5% contain samples that predate the onset of the Holocene (11.7 cal ka

BP), 19.7% extend back to 15 cal ka BP, and only 3.1% reach as far back as 20 cal ka BP

(Figure S1).

Compositional resilience index (CRI)

CRI can be computed as the sum of the standardized differences of each employed resilience

metric (Drake & Griffen, 2010). In this study, we applied four leading indicators of ecological

resilience namely, autocorrelation at-lag-1 (AR1), standard deviation (SD), skewness (SK),

and kurtosis (KURT), due to their positive performance (Dakos et al., 2024) to compute CRI

for each selected pollen record (Eqs. 1–9) (Dakos et al., 2024; Dakos et al., 2012; Drake &

Griffen, 2010).

$AR1 = \frac{\sum_{t=1}^{n-1}(z_t - \mu)(z_{t+1} - \mu)}{\sigma_z^2}$ (1)

$SD = \frac{1}{n-1}\sum_{t=1}^{n}(z_t - \mu)^2$ (2)

$SK = \frac{\frac{1}{n}\sum_{t=1}^{n}(z_t - \mu)^3}{\sqrt{\frac{1}{N}\sum_{t=1}^{n}(z_t - \mu)^2}}$ (3)

$KURT = \frac{\frac{1}{n}\sum_{t=1}^{n}(z_t - \mu)^4}{\left(\sqrt{\frac{1}{N}\sum_{t=1}^{n}(z_t - \mu)^2}\right)^2}$ (4)

$\widehat{AR1}_t = \frac{AR1_t - \overline{AR1}_{1:t}}{sd(AR1_{1:t})}$ (5)

$\widehat{SD}_t = \frac{SD_t - \overline{SD}_{1:t}}{sd(SD_{1:t})}$ (6)

$\widehat{SK}_t = \frac{SK_t - \overline{SK}_{1:t}}{sd(SK_{1:t})}$ (7)

$\widehat{KURT}_t = \frac{KURT_t - \overline{KURT}_{1:t}}{sd(KURT_{1:t})}$ (8)

$CRI_t = -(\widehat{AR1}_t + \widehat{SD}_t + \widehat{SK}_t + \widehat{KURT}_t)$ (9)

where z_t is the subset of time-detrended ecological state series (*i.e.*, scores of DCA1 in this

study) within a sliding window; z_{t+1} is the first-order lag series; μ and σ_z^2 are the mean and

variance of z_t, respectively. n is the size of the sliding window. $AR1_t$, SD_t, SK_t, $KURT_t$, and

CRI_t are the *AR1*, *SD*, *SK*, *KURT*, and *CRI* at time *t*, respectively. $AR1_{1:t}$, $SD_{1:t}$, $SK_{1:t}$, and

$KURT_{1:t}$ denote the timeseries of *AR1*, *SD*, *SK*, and *KURT* from times 1 to *t*, respectively.

$\overline{AR1}_{1:t}$, $\overline{SD}_{1:t}$, $\overline{SK}_{1:t}$, and $\overline{KURT}_{1:t}$ are the mean values of *AR1*, *SD*, *SK*, *KURT*, and *CRI*

from times 1 to *t*, respectively. $\widehat{AR1}_t$, $\widehat{SD}_t$, $\widehat{SK}_t$, and $\widehat{KURT}_t$ indicate the normalized *AR1*,

SD, *SK*, and *KURT* at time *t*. Given that an increase in these leading indicators theoretically

signifies a decline in system resilience, a rising in the composite index (calculated as the sum

of standardized differences across all employed leading indicator) would similarly indicate

reduced resilience (Drake & Griffen, 2010). To enable the composite index more intuitively

reflect the direction of resilience change, we defined the CRI_t as the negative value of the

summed standardized differences. Specifically, an increasing CRI trend indicates enhanced

vegetation resilience, whereas a decreasing CRI trend signifies declined vegetation resilience.

Given that these leading indicators are applicable exclusively to one-dimensional

timeseries (Dakos & Kéfi, 2022), we performed detrended correspondence analysis (DCA)

using 'decorana' function from the 'vegan' package (Oksanen et al., 2022) to generate an

aggregate measure reflecting the compositional state of multi-dimensional pollen assemblages.

The first axis of DCA (hereafter DCA1), which generally captures the dominant gradient of

compositional variation, was used as the aggregate measure for calculating the leading

indicators. Additionally, an underlying assumption for applying these leading indicators is that

the observations or timeseries must be equidistant. Therefore, we linearly interpolated the

DCA1 scores at 200-year intervals (corresponding to the maximum mean temporal resolution
of the samples) to generate equivalent timeseries (Dakos et al., 2008). Subsequently, we
detrended the interpolated DCA1 scores using a locally weighted regression smoother
(LOESS) to remove long-term trends. The residuals between interpolated and fitted values of
DCA scores were then used to compute the leading indicators (Wang et al., 2012).

The choice of the rolling window size is a trade-off between the time-resolution of the
available data and reliability of the estimate for the indicators (Lenton et al., 2012). To
determine the optimal window size, we estimated AR1, SD, SK, and KURT using rolling
window sizes ranging from 10% to 80% of the timeseries length, incremented in steps of five
data points. Trends in the leading indicators across different window sizes were quantified
using the Kendall tau rank correlation coefficient. For each indicator, the optimal rolling
window size was independently selected for individual DCA1 residual timeseries based on
the maximum Kendall tau value. Given that the optimal rolling window sizes generally vary
across the four leading indicators for each residual timeseries, the temporal coverage of the
composite index is determined by the common overlapping period among all four indicators.
Eventually, continental CRI sequences for North America, Europe, and Asia span the past
8,000 years, whereas those for South America, Africa, and Oceania cover the past 7,200 or
7,000 years (Figure S1).

Leading indicator calculations and trend estimations of statistical moments with different
rolling window sizes were performed using functions 'genetic_ews' and 'sensitively_ews',
respectively, in 'earlywarnings' package (Dakos et al., 2012).

Richness and evenness

Richness was quantified as the effective number of taxa (*Hill* number of order $q = 0$) in each
pollen assemblage, while evenness was measured by assessing the equality in the abundances
of individual pollen types within the assemblage (Birks et al., 2016; Hill, 1973). Given that
empirical estimates of *Hill* numbers tend to be an increasing function of sampling effort or
sample completeness (Chao et al., 2014), richness was determined by calculating the expected
number of taxa in samples rarefied to a fixed size of 300 pollen grains. Evenness was
estimated by using the modified *Hill*'s ratio (Eq. 10) (Alatalo, 1981):
$E = N2/N1$ (10)
where E is evenness; *N1* is the *Hill* number of order $q = 1$ (equivalent to the exponential of
Shannon entropy or diversity measure), serving as an estimate of the number of abundant taxa
in an assemblage; *N2* is the *Hill* numbers of order $q = 2$ (equivalent to the inverse of
Simpson's measure index of concentration), providing an estimate of the number of very
abundant taxa in an assemblage. *Hill* numbers ($q = 0, 1, 2$) were calculated using 'iNEXT'
function from the 'iNEXT' package (Chao et al., 2014; Hsieh et al., 2022).

Temporal *β* diversity

Temporal *β* diversity between temporally adjacent sample pairs within each pollen record was quantified using the Sørensen pair-wase dissimilarity, calculated via the 'beta.pair' function in the 'betapart' package (Baselga et al., 2018). To guarantee the accuracy of *β* diversity estimations along a temporal gradient, it is essential to maintain a uniform time interval between adjacent samples. For this, pollen data from each record were linearly interpolated to generate a dataset with evenly spaced interval prior to calculating temporal *β* diversity. Dataset interpolation was performed using the function 'interp.dataset' from the package 'rioja' (Juggins, 2022), with an interpolated interval set to 200 years to match the temporal resolution used for computing the CRI. Given that the algorithm for calculating the Sørensen pair-wase dissimilarity is exclusively applicable to present/absent data, all non-zero values in the interpolated pollen dataset were converted to 1.

Synchrony

We defined synchrony between two species or taxa as the extent of correlation in their abundances over a specified time period, with values ranging from +1 (perfect synchrony) to 0 (maximum asynchrony). For a multispecies community, community-wide synchrony can commonly be measured as the mean correlation between the abundance of each species versus the rest of the community (Gross et al., 2014) or the arithmetic mean correlation coefficient across all species pairs (Loreau & de Mazancourt, 2008; Thibaut & Connolly, 2013). Given that pollen data are presented as either the percentage abundance of each taxon in samples or the grain count of each taxon within rarefied samples with equal size, it is inappropriate to apply Gross's synchrony metric (Gross et al., 2014), as the correlation between the percentage abundance or grain count of each taxon and the rest of the assemblage would exhibit perfect correlation. Therefore, we considered community-wide synchrony metric based on correlation coefficient across all species pairs to be a feasible approach for reflecting the overall synchrony in pollen assemblages. In this study, we developed a new measure of overall synchrony for pollen assemblages as

$$\eta_w = \sum_i w_{ij} \cdot \left| r(A_i, A_j) \right| \qquad (11)$$

where η_w denotes the weighted synchrony index; A_i and A_j represent the timeseries of percentage abundance for taxon *i* and taxon *j* ($i \neq j$), respectively; *r* is the correlation coefficient between A_i and A_j, with a significant level < 0.05; w_{ij} is the sum of the mean percentage abundances of taxa *i* and *j* within an assemblage over the targeted period, namely the sum of the averages of A_i and A_j. We used Kendall's tau coefficient to quantified the correlations between paired taxa, and the significance of these correlations was evaluated using a two-tailed *t*-test. To obtain temporal trajectory of synchrony change, η_w was calculated within a five-point rolling window (1000 years) of the interpolated pollen dataset (200-year resolution).

Resilience trends at continental and biome scales

We modeled continental and biome CRIs against time with GAMs. GAMs are nonparametric data-driven regression models, which are effective in assessing non-linear relationships between response and predictor variables without any restrictive assumptions (Hastie & Tibshirani, 1987). To facilitate model convergence, we standardized each CRI timeseries using z-score normalization, which involves subtracting the mean from each data point and then dividing by the standard deviation. The standardized CRI timeseries were subsequently aggregated into continental subsets and biome subsets according to their geological locations. The information on the continents where the pollen records were obtained is from the LegacyPollen2.0 dataset. The ISLSCP II Potential Natural Vegetation dataset, sourced from the Earthdata platform (https://www.earthdata.nasa.gov/) and comprising fifteen vegetation types, was applied as the reference for defining biome subsets. For each subset, we fitted a GAM to the pooled standardized CRI values (hereafter, CRI_st) using the 'gam' function in the 'mgcv' package (Wood, 2011). In this process, we estimated a smooth function representing the relationship between CRI_st and time, assuming a Gaussian distribution. When formulating GAM equation, we designated CRI_st as the response variable and used the age corresponding to each CRI_st value as the predictor variable. We incorporated a thin plate regression spline smoother into the formula, setting the smoothing basis dimension to five for representing the smooth term. Given the heterogeneity (*e.g.*, variations in environmental conditions, local ecological differences) across sampling sites from which pollen records were derived, we included a random effect of sampling sites into the model to provide a more realistic representation of the underlying data structure and yield more robust results. The significance of the GAMs was evaluated using the *F* statistic.

Investigating key drivers for continental vegetating resilience shifts

We used a random forest (RF) model to identify the primary driver(s) responsible for the transition of continental resilience from an increasing to a declining trend during the late Holocene. RF analysis is a flexible, non-parametric regression tool which can handle a large number of predictors even in small datasets with a low number of observations, and is suited to analyzing non-linear relationships and complex interactions (Feld et al., 2016). To capture the states of resilience trends, we created a dummy variable based on the fitted continental-scale trends of CRI_st prior to conducting the analysis. Specifically, we assigned a value of 1 to periods showing an increasing trend and 0 to those showing a decreasing trend. It is worth noting that, given our focus on the increase-to-decline transition during the late Holocene, we excluded data from the dummy variable for two specific periods: (1) post-1.4 cal ka BP in North America, where trajectories deviated from the post-transition decline, and (2) pre-5.8 cal ka BP in South America, where trajectories deviated from the pre-transition increase (Figure 2). This approach allowed us to eliminate potential bias in assessing the relative importance of predictors that may arise from the existence of other transitions in the

CRI_st trends. Additionally, RF analysis was not applied to Africa due to the absence of a

discernible transition in its CRI_st trend (Figure 2).

In RF models, we used the dummy variable representing the resilience state as the

response variable. The predictor variables included annual mean precipitation (AMT),

temperature seasonality (TS), annual precipitation (AP), and precipitation seasonality (PS),

which were derived from CHELSA-TraCE21k dataset (Karger et al, 2023), as well as

anthropogenic land use, which was obtained from the KK10 dataset scenario of

Anthropogenic Land Cover Change (ALCC) (Kaplan et al., 2011). Based on the geographical

coordinates of the pollen records, we extracted timeseries data of the predictor variables for

all pollen record from the two datasets using the function 'extract' from the package 'terra'

(Hijmans, 2023). The time slices for the extracted data were selected to match those of the

CRI_st timeseries. To generate continental-scale timeseries for each predictor variable, we

firstly standardized each predictor's timeseries using z-score transformation and then

partitioned the standardized timeseries into six continental subsets based on their geographical

locations (Figure S2). For each predictor, continental-scale timeseries were generated by

computing the median values across all time slices within each continental subset.

Variable importance was quantified by the increase in out-of-bag error when the values

of the variable are randomly permuted (Breiman, 2001). We estimated the importance of

predictor variables using the percentage increase in the mean squared error ($\%InMSE$), where

higher $\%InMSE$ values indicate greater variable importance (Breiman, 2001). The

significance of each predictor was assessed with 1,000 permutations of the response variable.

A predictor was considered statistically significant if the permutation-induced increase in

MSE exceeded the 95[th] percentile of the null distribution generated from these permutations.

RF analyses and the permutation were conducted using the functions 'rfPermute' and

'importance' from the package 'rfPermute' (Archer, 2023).

Attributing long-term changes of continental resilience

We performed piecewise structural equation models (SEMs) based on a maximum likelihood

method to test direct and indirect pathways of biotic and abiotic factors on continental-scale

resilience of vegetation composition. SEM provides a way to test postulated relationships

between causal and confounding variables and to disentangle their effects (Shipley, 2016). We

constructed an initial SEM including climatic variables (AMT, TS, AP, PS) and ALCC as

observed variables, biotic attributes (richness, evenness, temporal β diversity, synchrony) as

latent variables, and the CRI as the dependent variable. The initial structure contained all

possible biotic and abiotic relationships with CRI, but did not define structural paths among

latent variables. Standardized path coefficients were used to measure the direct and indirect

effects of predictors (Grace & Bollen, 2005). We refined the model structure by adding

potentially significant pathway(s) and/or removing those with extremely weak standardized

path coefficient to improve the fit. The minimal and best model was selected based on the

Akaike Information Criterion (AIC). Fisher's C test was conducted to assess the global

goodness-of-fit of the models ($P \geq 0.05$ indicated reliable models). We conducted SEM

analyses using the function 'psem' from the package 'piecewiseSEM' (Lefcheck, 2016),

performing separately for each continent to examine continental-specific causal relationships

among variables. Following a similar data processing protocol to that used for RF analysis,

we generated continental-scale timeseries for each variable through multi-step procedure

involving standardization of data using z-score normalization, partitioning of standardized

data into continental subsets, and calculating median values for each time slices within each

subset (Figure S3).

In SEM, direct effects were measured as the standardized path coefficients between

variables, while indirect effects were calculated as the product of direct path coefficients

along a specific causal chain. We calculated the direct and indirect effects for each observed

and laten variable, independent of the statistical significance of the path coefficient(s). The net

effect of each variable was estimated by summing its direct and indirect effects. In addition,

we classified the observed and laten variables into three types: climate, ALCC, and ecological

attributes, to evaluate the relative contributions of different factor types on resilience. The net

effects of each factor type were quantified as the sum of all direct and indirect effects of the

variables grouped within that type. Then, we evaluated the relative effect sizes of distinct

factor types for each continental-scale SEM. Effect size was defined as the absolute value of

the net effect. The relative contributions of each factor type were quantified as the proportion

(%) of its effect size relative to the total effect size across all factor types.

We performed all data manipulations and analyses using R (Team, 2022). All packages

mentioned in this study are software extensions to the R environment.

Results

Continental-scale variations in vegetation resilience

All continents except Africa show significantly reversed CRI trends over the past 8,000 years

(Figure 2). Notably, these reversed trends predominantly comprise a pattern of increase

during the Mid-Holocene [8.0 to 4.2 thousand calibrated years before radiocarbon present (cal

ka BP, where 0 ka BP is 1950 CE)], followed by a decline in the Late Holocene (4.2 to 0 cal

ka BP), although the onset of these declines varied across continents (Figure 2). The earliest

decline signal was detected in North America (4.4 cal ka BP), followed by Europe where the

CRI began to decrease at 3.6 cal ka BP. For South America, Asia, and Oceania, the declines in

CRI trends occurred at 2.8, 2.4, and 1.6 cal ka BP, respectively. Interestingly, in North

America, the CRI trajectory over the past 1,200 years deviates from the "increase–decrease"

pattern, showing an increasing trend instead.

Figure 2. Continental CRI trends and relative importance of abiotic drivers. Solid lines

denote significant fits ($P < 0.05$) of generalized additive model (GAM), while dashed lines

indicate non-significant relationship ($P > 0.05$). The shaded areas represent the 95%

confidential intervals of the GAM fits. Histograms show the relative importance of each

driver, quantified as percentage increase in the mean squared error (%*InMSE*), with

significant drivers marked with asterisks ($P < 0.05$ and ** $P < 0.01$). ALCC, anthropogenic

land-cover change; AMT, annual mean temperature; TS, temperature seasonality; AP, annual

precipitation; PS, precipitation seasonality.

The main abiotic predictors driving the transition of continental-scale CRI trends from

increasing to declining during the late Holocene were identified through RF analysis (Figure

2). ALCC was the most important variable for predicting the transition across all continents,

except Oceania where AMT surpassed ALCC as the top-ranked driver. In addition to ALCC,

the trend transition of CRIs in North America and Europe was significantly influenced by TS,

PS, and AMT, whereas in Asia it was significantly affected by TS, AMT, and AP. In Oceania,

TS was another significant predictor that had considerable impact on the CRI shift.

Pathways and effects of abiotic and biotic attributes on continental-scale vegetation resilience

Pathways and effects of abiotic factors (AM, TS, AP, PS, ALCC) and ecological attributes (taxonomic richness, evenness, temporal β diversity, and synchrony) on CRIs have been estimated by SEMs (Figure 3). In North America, the most prominent path is the direct link from ALCC to CRI. Evenness plays a significant role in influencing CRI, serving both as a significant direct driver to CRI variations and as a mediator that relays the effects of other factors to CRI. TS and richness have significant impacts on evenness, subsequently further affecting CRI. Moreover, ALCC exerts an indirect influence on CRI through richness. In Europe, temporal β diversity is the only factor that displays a significant direct path influencing the CRI. Richness shows a significant indirect path to CRI through temporal β diversity. ALCC exerts a substantial influence on richness, which subsequently further impacts CRI. In Asia, TS displays a strong positive direct path to CRI, while PS shows a strong negative direct path to CRI. Furthermore, synchrony exhibits a significant negative direct link to CRI, through which ALCC exerts negative effects on CRI. Similar to Europe, there are no significant direct paths from biotic factors to CRI, and temporal β diversity is the only factor that shows a significant direct path influencing CRI in South America. In this SEM, abiotic factors contribute to CRI via several significant indirect paths, including those from AMT and evenness via temporal β diversity, from ALCC, PS, and richness via evenness, and from ALCC via richness. The SEM for Africa reveals only one significant direct path originating from AP. Similar to Africa, the significant direct paths originate solely from abiotic factors in Oceania; however, the abiotic factors exerting significant impacts are ALCC and AMT, rather than AP.

Figure 3. SEMs showing the direct and indirect effects of multiple abiotic and biotic drivers on CRI across six continents. Solid lines denote statistically significant paths ($P < 0.05$), while dashed lines indicate non-significant paths. Red and blue lines represent positive and negative effects, respectively. R^2 values indicate model-explained variance for each response variable. Standardized path coefficients are reported only for significant paths with significant level marked (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). Overall fit of SEM was evaluated using Shipley's test of d-separation: Fisher's C statistic (if $P > 0.05$, no paths are missing and the model is a good fit) and Akaike information criterion (AIC). ALCC, anthropogenic land-cover change; AMT, annual mean temperature; TS, temperature seasonality; AP, annual precipitation; PS, precipitation seasonality.

The factors incorporated into the SEMs show varying effect sizes across different continents (Figure 4). In North America, evenness contributes most to CRI variations, with net effect size (*n.e.s.*, thereafter) of 0.7057. Although richness demonstrates both substantial positive and indirect negative effects on CRI, its net effect is weak. ALCC exerts a relatively strong effect on CRI in North America (*n.e.s.* = 0.1579), while other abiotic factors contribute comparatively minimal influence to CRI variation. In Europe, temporal *β* diversity (*n.e.s.* = 0.4274) emerges as the strongest driver of CRI variation, followed by evenness (*n.e.s.* = 0.3028). In Europe, the *n.e.s.* of richness on CRI is weak and ALCC is identified as the strongest abiotic driver of long-term CRI variation (*n.e.s.* = -.2745). In Asia, PS (*n.e.s.* = 0.5886) and TS (*n.e.s.* = 0.54) show the notably stronger effects on CRI variation than all other evaluated factors. Synchrony also contributes a substantial effect on CRI (*n.e.s.* = -0.511). In South America, richness (*n.e.s.* = -0.5452) and temporal *β* diversity (*n.e.s* = -0.5142) are two strongest predictors of CRI variations, and AP (*n.e.s.* = 0.2169), ALCC (*n.e.s.* = 0.2023) and PS (*n.e.s.* = 0.1973) are the most important abiotic drivers. In Africa, richness (*n.e.s.* = 0.4255) and synchrony (*n.e.s.* = 0.3274) show obvious larger effect sizes on CRI variations than all other evaluated factors. Among abiotic factors, ALCC (*n.e.s.* = 0.1781) is the strongest predictor of CRI variation in Africa, following by AP (*n.e.s.* = -0.1563) and TS (*n.e.s.* = 0.1811). AMT (*n.e.s.* = 0.4119) and ALCC (*n.e.s.* = 0.3216) are the two most important factors influencing CRI in Oceania. Among biotic factors, the effect of evenness (*n.e.s.* = 0.3692) is stronger than that of richness (*n.e.s.* = 0.2069).

Figure 4. Effect sizes of abiotic and biotic drivers on continental CRI trajectories. Donut charts illustrate the relative net effect of each grouped variable to continental CRI changes. ALCC, anthropogenic land-cover change; AMT, annual mean temperature; TS, temperature seasonality; AP, annual precipitation; PS, precipitation seasonality.

When quantifying the total net effects, ecological attributes account for a predominant proportion of the total *n.e.s.*, ranging from *ca* 51% to *ca* 80%, across all continents except Europe (Figure 4). In Europe, the three contributors of ALCC, climate (*i.e.*, AMT, TS, AP, and PS), and ecological exhibit relatively comparable *n.e.s.*, with ALCC emerging as the most important factor. Although evenness and temporal *β* diversity have the largest *n.e.s.* among all evaluated factors in the SEM for Europe, their opposing effects diminish the overall impact of ecological attributes on CRI. A similar situation emerged in Asia, where PS and TS are the two most influential variables affecting CRI, yet their counteracting effects result in climate contributing only a minor portion of the total net effect on CRI. When comparing abiotic factors, climate exerts a notable stronger influence than ALCC on CRI variations in South America and Asia. Conversely, ALCC plays a more important role than climate in North America and Africa, while the *n.e.s.* of these two contributors are comparable in Europe and Oceania.

Discussion

Pollen archives have recorded the global shifts of vegetation compositional resilience during the Holocene, with the onsets of resilience declines occurred approximately between 4.4 and 1.6 cal ka BP across different continents except Africa (Figure 2). This aligns with that the

Holocene Earth experienced significant vegetation changes, manifesting in multiple
dimensions such as shifts in the compositional and structural in terrestrial vegetation (Nolan
et al., 2018), accelerated rates of compositional change (Mottl et al., 2021), and alterations in
floristic diversity (Gordon et al., 2024). However, the continental-scale trends in resilience
differ from those observed in other ecosystem properties, including composition (Figure S4),
turnover rates (Mottl et al., 2021), and biodiversity (Gordon et al., 2024). In addition,
transitions in these properties often occur asynchronously across continents and/or biomes,
reflecting the inherent non-linearity and heterogeneity of ecosystem responses to Holocene
environmental changes. Interestingly, the global resilience decline reversed in North America
at *ca* 1.2 cal ka BP (Figure 2). This phenomenon can be collectively attributed to the shifts in
the resilience trajectories of steppe, tundra, and savanna biomes during the past 1,500 and
2,000 years, as well as the concurrent slowdown in the resilience loss in forest and shrubland
in North America (Figure S5). On a global scale, compositional resilience in tundra and
savanna ecosystems diverged from their preceding declining trajectories and initiated a
recovery at around 3–2 cal ka BP (Figure S6). Thus, the continental-to-global scale shifts in
vegetation resilience should be predominantly shaped by the integrated effects of different
ecosystem responses to external stress, with such differential responses likely contingent upon
the intrinsic properties of these ecosystems (Pennekamp et al., 2018).

ALCC was primary factor that triggered the reversed trends of continental vegetation
resilience during the late Holocene (Figure 2). This was particularly evident in South America
where ALCC emerged as the sole statistically significant predictor driving reverse in the CRI
trend. Although human modification on Earth's landscapes can be traced back to the
Paleolithic era, its adverse effects on vegetation compositional resilience did not occur until
the late Holocene. Among the diverse array of resource procurement and land management,
intensive agriculture's high demand for resources and its continuous modification of land
makes it the strongest force in changing the Earth's landscapes. Intensive agriculture was
prevalent only in a few regions with suitable climatic conditions until 4–3 cal ka BP, after
which it began to spread more widely across the globe (Stephens et al., 2019). The rapid
expansion of intensive agriculture has largely coincided with the decline in the compositional
resilience of global vegetation, suggesting the existence of a land-use threshold inducing
shifts in the long-term trajectories of continental vegetation resilience. In addition to ALCC,
climatic variables contributed substantially to the shifts in the long-term trends of vegetation
resilience (Figure 2), suggesting that the observed resilience decline was commonly triggered
by the synergistic effects of human disturbance and climate change.

Both theoretical and empirical researches have delved into how various features of
ecosystems, such as complexity of ecosystems (Pimm, 1984), biodiversity (Liang et al., 2025;
Pennekamp et al., 2018; Rodrigues et al., 2025; S. Wang & Loreau, 2016), species synchrony
(Schnabel et al., 2021; Valencia et al., 2020), species evenness (Isbell et al., 2009; Valencia et
al., 2020), spatial self-organization (de Paoli et al., 2017; Kéfi et al., 2024), affect ecosystem

stability/resilience. The relationship between biodiversity and stability has long been a

contentious issue, primarily focused on the question of whether greater diversity (number of

species) leads to less variability in communities. Our findings reveals that species richness

can exert positive, negative, or negligible effects on vegetation resilience, at least in terms of

vegetation composition, when assessed across continental scale and millennial timescales

(Figure 3). This conclusion supports that a general conclusion about the diversity-stability

cannot be anticipated (Ives & Carpenter, 2007), since external pressures often affect stability

and diversity simultaneously. The stabilizing effects of biodiversity arise from species'

asynchronous responses to environmental fluctuations, a phenomenon commonly referred as

insurance effects (Loreau et al., 2021; Yachi & Loreau, 1999). However, species synchrony

was identified as an important structural component of the SEMs only for Asia and Africa

(Figure 3). In Asia, the SEM demonstrates a negative link between synchrony and vegetation

compositional resilience, which appears to align with the insurance effect. In contrast, the

SEM of Africa reveals a positive pathway between synchrony and resilience. Furthermore, we

observe a positive richness-synchrony relationship in Africa, which appears to contradict the

prevailing hypothesis that higher species richness enhances the likelihood of a community

containing species with differing responses to abiotic drivers or competition, thereby reducing

synchrony (McCann, 2000). However, the positive correlation between species richness and

synchrony observed in this study aligns with findings from an empirical global assessment

that examined how species richness and synchrony influence community stability (Valencia et

al., 2020). These findings collectively suggest that the effect of richness on stability/resilience

is not necessarily mediated through its negative influence on synchrony, either at the local

scale or at the continental scale.

Our study identifies evenness as another important direct or indirect predictor of

long-term changes in vegetation compositional resilience across North and South America,

Europe, and Oceania (Figure 3). Theoretically, declines in evenness may decrease temporal

stability/resilience by weakening the portfolio effect or statistical averaging (Doak et al., 1998;

Hillebrand et al., 2008). While, the SEMs predominantly reveal a negative association

between evenness and resilience (Figure 3). This is likely because, analogous to the

mass-ratio hypothesis (Grime, 1998), dominant species exert stronger effect on vegetation

dynamics than common and rare taxa do. When a vegetation ecosystem operates under a

background where the dominant taxa progressively lose their ecological dominance (*i.e.*,

evenness increase), a gradual decline of composition stability or resilience will arise from the

reduced involvement of the dominants in shaping ecosystem structure and dynamics.

Temporal β diversity, defined as temporal variation in community composition or species

assemblages at a specific location (Magurran et al., 2019), comprises two distinct components:

species turnover (resulting from species replacement) and nestedness (resulting from species

loss or gain) (Baselga, 2010). In this study, temporal β diversity is quantified using adjacent

samples from time series with fixed temporal intervals, thereby serving as a metric for the rate

of compositional change in an assemblage. It was expected that the variations in

compositional change rates were expected to influence vegetation resilience, and this

hypothesis is verified by the SEMs for Europe and South America (Figures 3 & 4). However,

similar to species richness and evenness, the direction of temporal β diversity's impact on

resilience is not fixed.

Despite these complexities present above, there are some general underlying mechanisms

that regulate changes in vegetation compositional resilience. Our study reveals that the

assessed abiotic factors exert certain impacts on the biotic factors, and that some or all of

these abiotic factors have direct influences on resilience (Figure 3). These findings emphasize

that biotic factors unlikely change in isolation from abiotic drivers influencing stability, and in

fact these abiotic drivers are typically the main causes of changes in ecological attributes and

may serve as the original driving force for variations in ecosystem stability or resilience (Ives

& Carpenter, 2007). As another generality, both biotic and abiotic factors may have opposing

effects on resilience through their direct and direct pathways (Figure 4). In some cases, a

certain factor may appear to have a negligible impact on resilience because it simultaneously

generates positive and negative effects of comparable magnitude. However, this does not

necessarily imply that the positive and negative effects produced by this factor are weak *per*

se. Regarding the relative importance of different factor types, biotic factors as a whole

generally exert a substantially greater influence on continental-scale changes in compositional

resilience than abiotic factors do, although Europe represents an exception to this pattern

(Figure 4). Furthermore, while ALCC has been identified as the primary driver for the shift in

the long-term trend of vegetation compositional resilience during the late Holocene, its

cumulative effect on resilience trajectories over the past 8,000 years has not exceeded the

cumulative influence of climate change across all continents.

In conclusion, our results suggest that human-induced modifications to Earth's

landscapes have led to a persistent, global-scale decline in vegetation resilience that has lasted

for several millennia. However, we have also found that over the last 1,000–2,000 years, in

some regions constrained by moisture and/or temperature conditions, such as tundra and

savanna ecosystems, vegetation resilience has continuously enhanced. Regarding the

mechanisms driving millennia-scale changes in continental vegetation resilience, this study

reveals a complex picture when incorporating factors representing human disturbance, climate

changes, and ecological attributes. Despite this, the results demonstrate that biological

attributes generally contribute more than other factors in regulating the long-term changes in

vegetation resilience. There are two limitations in our data that could potentially affect the

findings presented here. First, the sparse pollen records in Africa and Australia may lead to a

lack of precision when analyzing the long-term trends of vegetation resilience in these two

continents. In addition, since pollen data generally lack high taxonomic resolution, in cases

where the changes in vegetation composition primarily manifest as alternations in plant taxa

within the same genus or family, the variations in vegetation composition are likely to be
underestimated. This may subsequently bias the estimated resilience metrics based on
taxonomic compositional changes, thereby affecting accurate assessments of resilience
dynamics.

Acknowledgements
We acknowledge the contributions of all data providers who supported this study. We
appreciate Professor Rong Wang (Nanjing Institute of Geography & Limnology, Chinese
Academy of Sciences) for his insightful discussions on methodological approaches. This work
was supported by the National Natural Science Foundation of China (No. 42477484, No.
42577519) and the State Key Laboratory for Vegetation Structure, Function and Construction
(No. VegLabOF2025006).

Author Contributions
Mengna Liao: Conceptualization (lead); data curation (equal); formal analysis (lead);
funding acquisition (equal); methodology (lead); software (lead); supervision (equal);
validation (lead); visualization (lead); writing – original draft (lead); writing – review and
editing (equal). **Kai Li:** Conceptualization (supporting); funding acquisition (equal); writing –
original draft (supporting); writing – review and editing (equal). **Chenzhi Li:** Data Curation
(equal); investigation (equal); writing – review and editing (equal). **Ulrike Herzschuh:** Data
Curation (equal); investigation (equal); writing – review and editing (equal). **Jian Ni:**
Conceptualization (supporting); investigation (equal); supervision (equal); writing – review
and editing (equal).

Competing Interests statement
The authors declare no competing interests.

Data Availability and Code Availability
The LegacyPollen2.0 pollen dataset is openly available at
https://doi.pangaea.de/10.1594/PANGAEA.965907. The CHELSA-TraCE21k climate data is
openly available at https://www.chelsa-climate.org/datasets/chelsa-trace21k-centennial. The
KK10 Anthropogenic Land Cover data is openly available at
https://doi.pangaea.de/10.1594/PANGAEA.871369. The R scripts are available on GitHub at
https://github.com/mnliao/Global-shifts-vegetation-resilience.git.

Supplementary Materials was provided as a separated PDF file.

References

Alatalo, R. V. (1981). Problems in the measurement of evenness in ecology. *Oikos*, 199-204.

Archer, E. (2023). rfPermute: Estimate Permutation p-Values for Random Forest Importance Metrics. R package (Version 2.5.2): RCRAN. Retrieved from http:cran.r-project.org/package=rfPermute

Barnosky, A. D., Hadly, E. A., Bascompte, J., Berlow, E. L., Brown, J. H., Fortelius, M., . . . Marquet, P. A. (2012). Approaching a state shift in Earth's biosphere. *Nature, 486*(7401), 52-58.

Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global ecology and Biogeography, 19*(1), 134-143.

Baselga, A., Orme, D., Villeger, S., De Bortoli, J., & Leprieur, F. (2018). betapart: Partitioning Beta Diversity into Turnover and Nestedness Components. R package (Version 1.5.1). Retrieved from https://CRAN.R-project.org/package=betapart

Birks, H. J. B. (2019). Contributions of Quaternary botany to modern ecology and biogeography. *Plant Ecology & Diversity, 12*(3-4), 189-385.

Birks, H. J. B., Felde, V. A., Bjune, A. E., Grytnes, J.-A., Seppä, H., & Giesecke, T. (2016). Does pollen-assemblage richness reflect floristic richness? A review of recent developments and future challenges. *Review of Palaeobotany and Palynology, 228*, 1-25.

Blaauw, M., & Christen, J. A. (2011). Flexible paleoclimate age-depth models using an autoregressive gamma process. *Bayesian Analysis, 6*(3), 457-474.

Blois, J. L., Zarnetske, P. L., Fitzpatrick, M. C., & Finnegan, S. (2013). Climate change and the past, present, and future of biotic interactions. S*cience, 341*(6145), 499-504.

Boulton, C. A., Lenton, T. M., & Boers, N. (2022). Pronounced loss of Amazon rainforest resilience since the early 2000s. *Nature Climate Change, 12*(3), 271-278.

Breiman, L. (2001). Random forests. *Machine Learning, 45*(1), 5-32.

Carpenter, S. R., & Brock, W. A. (2006). Rising variance: a leading indicator of ecological transition. *Ecology Letters, 9*(3), 311-318.

Cavanaugh, K. C., Dangremond, E. M., Doughty, C. L., Williams, A. P., Parker, J. D., Hayes, M. A., . . . Feller, I. C. (2019). Climate-driven regime shifts in a mangrove–salt marsh ecotone over the past 250 years. *Proceedings of the National Academy of Sciences, 116*(43), 21602-21608.

Chao, A., Gotelli, N. J., Hsieh, T., Sander, E. L., Ma, K., Colwell, R. K., & Ellison, A. M. (2014). Rarefaction and extrapolation with Hill numbers: a framework for sampling and estimation in species diversity studies. *Ecological Monographs, 84*(1), 45-67.

Dakos, V., Boulton, C. A., Buxton, J. E., Abrams, J. F., Arellano-Nava, B., Armstrong McKay, D. I., . . . Dylewsky, D. (2024). Tipping point detection and early warnings in climate, ecological, and human systems. *Earth System Dynamics, 15*(4), 1117-1135.

Dakos, V., Carpenter, S. R., Brock, W. A., Ellison, A. M., Guttal, V., Ives, A. R., . . . van Nes,
E. H. (2012). Methods for detecting early warnings of critical transitions in time
series illustrated using simulated ecological data. *Plos ONE, 7*(7), e41010.

Dakos, V., & Kéfi, S. (2022). Ecological resilience: What to measure and how. *Environmental*
Research Letters, 17(4), 043003.

Dakos, V., Scheffer, M., Van Nes, E. H., Brovkin, V., Petoukhov, V., & Held, H. (2008).
Slowing down as an early warning signal for abrupt climate change. *Proceedings of*
the National Academy of Sciences, 105(38), 14308-14312.

Dakos, V., van Nes, E. H., & Scheffer, M. (2013). Flickering as an early warning signal.
Theoretical Ecology, 6(3), 309-317.

de Bello, F., Lavorel, S., Hallett, L. M., Valencia, E., Garnier, E., Roscher, C., . . . Májeková,
M. (2021). Functional trait effects on ecosystem stability: assembling the jigsaw
puzzle. *Trends in Ecology & Evolution, 36*(9), 822-836.

De Paoli, H., Van Der Heide, T., Van Den Berg, A., Silliman, B. R., Herman, P. M., & Van De
Koppel, J. (2017). Behavioral self-organization underlies the resilience of a coastal
ecosystem. *Proceedings of the National Academy of Sciences, 114*(30), 8035-8040.

Doak, D. F., Bigger, D., Harding, E., Marvier, M., O'malley, R., & Thomson, D. (1998). The
statistical inevitability of stability-diversity relationships in community ecology. *The*
American Naturalist, 151(3), 264-276.

Drake, J. M., & Griffen, B. D. (2010). Early warning signals of extinction in deteriorating
environments. *Nature, 467*(7314), 456-459.

Duveiller, G., Hooker, J., & Cescatti, A. (2018). The mark of vegetation change on Earth's
surface energy balance. *Nature Communications, 9*(1), 679.

Fastovich, D., Meyers, S. R., Saupe, E. E., Williams, J. W., Dornelas, M., Dowding, E. M., . . .
Kiessling, W. (2025). Coupled, decoupled, and abrupt responses of vegetation to
climate across timescales. *Science, 389*(6755), 64-68.

Feld, C. K., Segurado, P., & Gutierrez-Canovas, C. (2016). Analysing the impact of multiple
stressors in aquatic biomonitoring data: A 'cookbook' with applications in R. *Science*
of the Total Environment, 573, 1320-1339.

Feng, Y., Su, H., Tang, Z., Wang, S., Zhao, X., Zhang, H., . . . Fang, J. (2021). Reduced
resilience of terrestrial ecosystems locally is not reflected on a global scale.
Communications Earth & Environment, 2(1), 88.

Folke, C., Carpenter, S., Walker, B., Scheffer, M., Elmqvist, T., Gunderson, L., & Holling, C.
S. (2004). Regime shifts, resilience, and biodiversity in ecosystem management.
Annual Review of Ecology, Evolution, and Systematics, 35(1), 557-581.

Forzieri, G., Dakos, V., McDowell, N. G., Ramdane, A., & Cescatti, A. (2022). Emerging
signals of declining forest resilience under climate change. *Nature, 608*(7923),
534-539.

Gordon, J. D., Fagan, B., Milner, N., & Thomas, C. D. (2024). Floristic diversity and its

relationships with human land use varied regionally during the Holocene. *Nature ecology & evolution*, 1-13. doi:https://doi.org/10.1038/s41559-024-02457-x

Grace, J. B., & Bollen, K. A. (2005). Interpreting the results from multiple regression and structural equation models. *Bulletin of the Ecological Society of America, 86*(4), 283-295.

Grime, J. (1998). Benefits of plant diversity to ecosystems: immediate, filter and founder effects. *Journal of Ecology, 86*(6), 902-910.

Gross, K., Cardinale, B. J., Fox, J. W., Gonzalez, A., Loreau, M., Wayne Polley, H., . . . van Ruijven, J. (2014). Species richness and the temporal stability of biomass production: a new analysis of recent biodiversity experiments. *The American Naturalist, 183*(1), 1-12.

Guttal, V., & Jayaprakash, C. (2008). Changing skewness: an early warning signal of regime shifts in ecosystems. *Ecology Letters, 11*(5), 450-460.

Hastie, T., & Tibshirani, R. (1987). Generalized additive models: some applications. *Journal of the American Statistical Association, 82*(398), 371-386. doi:https://doi.org/10.1080/01621459.1987.10478440

Hatton, I. A., Mazzarisi, O., Altieri, A., & Smerlak, M. (2024). Diversity begets stability: Sublinear growth and competitive coexistence across ecosystems. *Science, 383*(6688), eadg8488.

Herzschuh, U., Li, C., Böhmer, T., Postl, A. K., Heim, B., Andreev, A. A., . . . Ni, J. (2022). LegacyPollen 1.0: a taxonomically harmonized global late Quaternary pollen dataset of 2831 records with standardized chronologies. *Earth System Science Data, 14*(7), 3213-3227.

Hijmans, R. J. (2023). terra: Spatial Data Analysis. R package version 1.7-29 (Version 1.7-29): RCRAN. Retrieved from http:cran.r-project.org/package=terra

Hill, M. O. (1973). Diversity and Evenness: A Unifying Notation and Its Consequences. *Ecology, 54*(2), 427-432. doi:doi:10.2307/1934352

Hillebrand, H., Bennett, D. M., & Cadotte, M. W. (2008). Consequences of dominance: a review of evenness effects on local and regional ecosystem processes. *Ecology, 89*(6), 1510-1520.

Hisano, M., Searle, E. B., & Chen, H. Y. (2018). Biodiversity as a solution to mitigate climate change impacts on the functioning of forest ecosystems. *Biological Reviews, 93*(1), 439-456.

Holling, C. S. (1973). Resilience and stability of ecological systems. *Annual Review of Ecology and Systematics, 4*, 1-23.

Holling, C. S. (1996). Engineering Resilience Versus Ecological Resilience. In P. E. Schulze (Ed.), *Engineering within Ecological Constraints* (pp. 31-43). Washington DC: National Academy Press.

Hsieh, T. C., Ma, K. H., & Chao, A. (2022). iNEXT: iNterpolation and Extrapolation for

species diversity (Version 3.0.0). Retrieved from

http://chao.stat.nthu.edu.tw/wordpress/software_download/

Isbell, F., Craven, D., Connolly, J., Loreau, M., Schmid, B., Beierkuhnlein, C., . . . De Luca, E.

(2015). Biodiversity increases the resistance of ecosystem productivity to climate

extremes. *Nature, 526*(7574), 574-577.

Isbell, F. I., Polley, H. W., & Wilsey, B. J. (2009). Biodiversity, productivity and the temporal

stability of productivity: patterns and processes. *Ecology Letters, 12*(5), 443-451.

Ives, A. R., & Carpenter, S. R. (2007). Stability and diversity of ecosystems. *Science,*

317(5834), 58-62.

Jackson, S. T., & Blois, J. L. (2015). Community ecology in a changing environment:

Perspectives from the Quaternary. *Proceedings of the National Academy of Sciences,*

112(16), 4915-4921.

Juggins, S. (2022). rioja: Analysis of Quanternary Science Data, R package (Version 1.0-5):

RCRAN. Retrieved from http:cran.r-project.org/package=rioja

Kéfi, S., Génin, A., Garcia-Mayor, A., Guirado, E., Cabral, J. S., Berdugo, M., . . . Maestre, F.

T. (2024). Self-organization as a mechanism of resilience in dryland ecosystems.

Proceedings of the National Academy of Sciences, 121(6), e2305153121.

Kaplan, J. O., Krumhardt, K. M., Ellis, E. C., Ruddiman, W. F., Lemmen, C., & Goldewijk, K.

K. (2011). Holocene carbon emissions as a result of anthropogenic land cover change.

The Holocene, 21(5), 775-791.

Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H., & Zimmermann, N. E. (2023).

CHELSA-TraCE21k–high-resolution (1 km) downscaled transient temperature and

precipitation data since the Last Glacial Maximum. *Climate of the Past, 19*(2),

439-456.

Lefcheck, J. S. (2016). piecewiseSEM: Piecewise structural equation modelling in r for

ecology, evolution, and systematics. *Methods in Ecology and Evolution, 7*(5),

573-579.

Lenton, T. M., Livina, V., Dakos, V., van Nes, E. H., & Scheffer, M. (2012). Early warning of

climate tipping points from critical slowing down: comparing methods to improve

robustness. *Philosophical Transactions of the Royal Society A: Mathematical,*

Physical and Engineering Sciences, 370(1962), 1185-1204.

Li, C., Dallmeyer, A., Ni, J., Chevalier, M., Willeit, M., Andreev, A. A., . . . Wieczorek, M.

(2025). Global biome changes over the last 21 000 years inferred from model–data

comparisons. *Climate of the Past, 21*(6), 1001-1024.

Li, C., Ni, J., Böhmer, T., Cao, X., Zhou, B., Liao, M., . . . Heim, B. (2024). LegacyPollen2. 0:

an updated global taxonomically and temporally standardized fossil pollen dataset of

3680 palynological records, PANGAEA [data set bundled publication].

PANGAEA–Data Publisher for Earth & Environmental Science.

Li, C., Postl, A. K., Böhmer, T., Cao, X., Dolman, A. M., & Herzschuh, U. (2022).

795 Harmonized chronologies of a global late Quaternary pollen dataset (LegacyAge 1.0).

796 *Earth System Science Data, 14*(3), 1331-1343.

797 Liang, M., Yang, Q., Chase, J. M., Isbell, F., Loreau, M., Schmid, B., . . . Wang, S. (2025).

798 Unifying spatial scaling laws of biodiversity and ecosystem stability. *Science,*

799 *387*(6740), eadl2373.

800 Liu, Y. Y., Van Dijk, A. I., De Jeu, R. A., Canadell, J. G., McCabe, M. F., Evans, J. P., & Wang,

801 G. (2015). Recent reversal in loss of global terrestrial biomass. *Nature Climate*

802 *Change, 5*(5), 470-474.

803 Loreau, M., Barbier, M., Filotas, E., Gravel, D., Isbell, F., Miller, S. J., . . . Germain, R.

804 (2021). Biodiversity as insurance: from concept to measurement and application.

805 *Biological Reviews, 96*(5), 2333-2354.

806 Loreau, M., & de Mazancourt, C. (2008). Species synchrony and its drivers: neutral and

807 nonneutral community dynamics in fluctuating environments. *The American*

808 *Naturalist, 172*(2), E48-E66.

809 Loreau, M., & De Mazancourt, C. (2013). Biodiversity and ecosystem stability: a synthesis of

810 underlying mechanisms. *Ecology Letters, 16*, 106-115.

811 Magurran, A. E., Dornelas, M., Moyes, F., & Henderson, P. A. (2019). Temporal *β*

812 diversity—A macroecological perspective. *Global Ecology and Biogeography, 28*(12),

813 1949-1960.

814 McCann, K. S. (2000). The diversity–stability debate. *Nature, 405*(6783), 228-233.

815 Mottl, O., Flantua, S. G. A., Bhatta, K. P., Felde, V. A., Giesecke, T., Goring, S., . . . Ivory, S.

816 (2021). Global acceleration in rates of vegetation change over the past 18,000 years.

817 *Science, 372*(6544), 860-864.

818 Myneni, R. B., Keeling, C., Tucker, C. J., Asrar, G., & Nemani, R. R. (1997). Increased plant

819 growth in the northern high latitudes from 1981 to 1991. *Nature, 386*(6626), 698-702.

820 Nolan, C., Overpeck, J. T., Allen, J. R., Anderson, P. M., Betancourt, J. L., Binney, H. A., . . .

821 Cheddadi, R. (2018). Past and future global transformation of terrestrial ecosystems

822 under climate change. *Science, 361*(6405), 920-923.

823 Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., . . . Weedon, J.

824 (2022). vegan: Community Ecology Package. R package (Version 2.6-4). Retrieved

825 from https://CRAN.R-project.org/package=vegan

826 Oliver, T. H., Heard, M. S., Isaac, N. J., Roy, D. B., Procter, D., Eigenbrod, F., . . . Petchey, O.

827 L. (2015). Biodiversity and resilience of ecosystem functions. *Trends in Ecology &*

828 *Evolution, 30*(11), 673-684.

829 Pennekamp, F., Pontarp, M., Tabi, A., Altermatt, F., Alther, R., Choffat, Y., . . . Griffiths, J. I.

830 (2018). Biodiversity increases and decreases ecosystem stability. *Nature, 563*(7729),

831 109-112.

832 Peterson, G., Allen, C. R., & Holling, C. S. (1998). Ecological resilience, biodiversity, and

833 scale. *Ecosystems, 1*, 6-18.

834   Pimm, S. L. (1984). The complexity and stability of ecosystems. *Nature, 307*(5949), 321-326.

835   Rodrigues, A. V., Rissanen, T., Jones, M. M., Huikkonen, I. M., Huitu, O., Korpimäki, E., . . .
836   Pietiäinen, H. (2025). Cross‐Taxa Analysis of Long‐Term Data Reveals a Positive
837   Biodiversity‐Stability Relationship With Taxon‐Specific Mechanistic Underpinning.
838   *Ecology Letters, 28*(4), e70003.

839   Scheffer, M., Bascompte, J., Brock, W. A., Brovkin, V., Carpenter, S. R., Dakos, V., . . .
840   Sugihara, G. (2009). Early-warning signals for critical transitions. *Nature, 461*(7260),
841   53-59.

842   Schnabel, F., Liu, X., Kunz, M., Barry, K. E., Bongers, F. J., Bruelheide, H., . . . Pfaff, C.-T.
843   (2021). Species richness stabilizes productivity via asynchrony and drought-tolerance
844   diversity in a large-scale tree biodiversity experiment. *Science Advances, 7*(51),
845   eabk1643.

846   Shipley, B. (2016). *Cause and correlation in biology: A user's guide to path analysis,*
847   *structural equations and causal inference with R (2nd edn)*. Cambridge: Cambridge
848   University Press.

849   Smith, T., Traxl, D., & Boers, N. (2022). Empirical evidence for recent global shifts in
850   vegetation resilience. *Nature Climate Change, 12*(5), 477-484.

851   Smol, J. P., Wolfe, A. P., Birks, H. J. B., Douglas, M. S., Jones, V. J., Korhola, A., . . .
852   Antoniades, D. (2005). Climate-driven regime shifts in the biological communities of
853   arctic lakes. *Proceedings of the National Academy of Sciences, 102*(12), 4397-4402.

854   Steffen, W., Rockström, J., Richardson, K., Lenton, T. M., Folke, C., Liverman, D., . . .
855   Crucifix, M. (2018). Trajectories of the Earth System in the Anthropocene.
856   *Proceedings of the National Academy of Sciences, 115*(33), 8252-8259.

857   Stephens, L., Fuller, D., Boivin, N., Rick, T., Gauthier, N., Kay, A., . . . Denham, T. (2019).
858   Archaeological assessment reveals Earth's early transformation through land use.
859   *Science, 365*(6456), 897-902.

860   Team, R. C. (2022). R: A language and environment for statistical computing. R Foundation
861   for Statistical Computing. Vienna, Austria.

862   Thibaut, L. M., & Connolly, S. R. (2013). Understanding diversity–stability relationships:
863   towards a unified model of portfolio effects. *Ecology Letters, 16*(2), 140-150.

864   Valencia, E., De Bello, F., Galland, T., Adler, P. B., Lepš, J., E-Vojtkó, A., . . . Dengler, J.
865   (2020). Synchrony matters more than species richness in plant community stability at
866   a global scale. *Proceedings of the National Academy of Sciences, 117*(39),
867   24345-24351.

868   Wang, R., Dearing, J. A., Langdon, P. G., Zhang, E., Yang, X., Dakos, V., & Scheffer, M.
869   (2012). Flickering gives early warning signals of a critical transition to a eutrophic
870   lake state. *Nature, 492*(7429), 419-422.

871   Wang, S., & Loreau, M. (2014). Ecosystem stability in space: α, β and γ variability. *Ecology*
872   *Letters, 17*(8), 891-901.

Wang, S., & Loreau, M. (2016). Biodiversity and ecosystem stability across scales in
metacommunities. *Ecology Letters, 19*(5), 510-518.
Wernberg, T., Bennett, S., Babcock, R. C., De Bettignies, T., Cure, K., Depczynski, M., . . .
Hovey, R. K. (2016). Climate-driven regime shift of a temperate marine ecosystem.
Science, 353(6295), 169-172.
Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood
estimation of semiparametric generalized linear models. *Journal of the Royal*
Statistical Society Series B: Statistical Methodology, 73(1), 3-36.
Yachi, S., & Loreau, M. (1999). Biodiversity and ecosystem productivity in a fluctuating
environment: the insurance hypothesis. *Proceedings of the National Academy of*
Sciences, 96(4), 1463-1468.