Direct and diffuse cross-kingdom interactions in plant microbiome assembly

Rachel A. Hammer[1], Marissa R. Lee[1], Jeffrey Kimbrel[2], Rhona Stuart[2], Christine V. Hawkes[2]*

[1]Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC 27607, USA
[2]Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

*Corresponding author: chawkes@ncsu.edu

Abstract

Studies of plant-associated microbial communities consistently indicate a role for classic assembly mechanisms, such as environmental and host filters, but often leave substantial unexplained variation. We propose that biotic interactions within microbial communities may help to fill this gap, specifically cross-kingdom interactions between fungi and bacteria, as these are increasingly found to be important to both assembly and function. We hypothesized that direct interactions between bacterial and fungal taxa are an important driver of composition in low-diversity leaf habitats, where pairwise reciprocal interactions are more likely. In high-diversity root habitats, however, we expected diffuse, indirect interactions to be more relevant to composition. To test these hypotheses, we used 16S and ITS rRNA sequence analysis to characterize bacterial and fungal communities of switchgrass (*Panicum virgatum* L.) leaves and roots within and across 14 stands spanning mountain to coastal ecoregions of North Carolina, USA. We analyzed putative direct and diffuse interactions using ecological network inference and examined the variance explained in microbial community composition using variance partitioning analysis with spatial, environmental, and biotic interactions as explanatory factors. Variance partitioning showed that cross-kingdom biotic interactions contribute to microbial community structure. The largest improvements to variance explained (5-11%) were from including direct interactions, except for root fungal communities where the variance explained by diffuse interactions (7.5%) was more than double that of direct interactions (2.8%). These contributions were comparable with those from environmental and spatial factors. The joint effects of putative biotic interactions and environmental conditions also contributed to the explained variation, highlighting the importance of the environmental tracking in microbes. These findings provide insight into using network inference for identifying putative cross-kingdom biotic interactions in plant-associated microbiomes and their contributions to microbiome assembly, which could guide the direction of future research on biotic interactions.

Keywords
Bacteria, fungi, foliar, root, switchgrass, *Panicum virgatum*

INTRODUCTION

Studies of plant-associated microbial communities consistently indicate a role for classic assembly mechanisms, such as environmental and host filters, but often leave substantial unexplained variation (Waring and Hawkes 2018; Wang et al. 2023; Vieira et al. 2020; Mészárošová et al. 2024; Giauque and Hawkes 2016; Lee and Hawkes 2021a). Here, we propose that this knowledge gap may be partly filled by considering biotic interactions, particularly between fungi and bacteria. Advances in models (e.g., network, joint-species distribution, trait-based (Layeghifard et al. 2017; Pollock et al. 2014; Aguilar-Trigueros et al. 2014) and mechanistic approaches (Jacoby and Kopriva 2019) are driving progress in assessing the role of biotic interactions in community assembly to analyze larger datasets and multiple data matrices (Pichler and Hartig 2021; Jiang et al. 2019). These advances have facilitated our ability to consider biotic interactions and uncover key microbes that respond both to abiotic factors and each other (e.g., Agler et al. 2016).

To date, the vast majority of experimental microbiome studies address biotic interactions pairwise, which assumes that interactions are direct and independent of other taxa in the community. Yet emergent properties arising from indirect interactions, which are unpredictable from pairwise effects, are common in both plant and animal communities (Lai et al. 2022; Bever et al. 2015; Werner and Peacor 2003) and their occurrence in simple microbial communities is supported by both empirical evidence and predictive models (Gallardo-Navarro and Santillán 2019; Mickalide and Kuehn 2019; Sanchez-Gorostiaga et al. 2019; Morin et al. 2022). Indirect effects occur when the interaction between two species is mediated via a third species by, for example, interaction chains, modification of environmental conditions, or changes in resource availability (Wootton 1994). Indirect interactions in multispecies communities may be diffuse if the effect of one species is spread over many other species (Moen 1989). In microbial communities, such diffuse effects might occur through extracellular production of volatiles, enzymes, signaling molecules, and other metabolites (Weiland-Bräuer 2021; Lobanov et al. 2023; Rizaludin et al. 2023). In plant-associated microbial communities, diffuse effects might also include modifications to the host environment (Seybold et al. 2020).

Cross-kingdom interactions may be as important as within-kingdom interactions in microbial communities (Lee et al. 2022; Rossmann et al. 2020), including both direct and indirect effects. Bacteria and fungi often live in close physical associations that can facilitate interactions, such as in mixed biofilms within soil micro-niches and the rhizosphere (Pandit et al. 2020; Miquel Guennoc et al. 2018), where both competition and facilitation can occur (Wang and Kuzyakov 2024; Deveau et al. 2018). For instance, mycorrhizal hyphae are known to harbor bacteria internally, on the hyphal surface, and in the hyphosphere (Emmett et al. 2021; Frey-Klett et al. 2011; Warmink et al. 2011), where recent studies point to carbon-nutrient exchange (Nuccio et al. 2022; Duan et al. 2023) and facilitation of stress tolerance (Hestrin et al. 2022). Indirect interactions are less studied, but a clear example was reported by Sharma et al. (2021), in which a nitrogen-fixing bacterium modified the interaction between a fungus and a fungus-feeding bacterium. Despite evidence supporting cross-kingdom interactions (Yuan et al. 2021), these

are rarely incorporated into assessments of community assembly and our understanding of microbial interaction effects on assembly therefore remain limited (Dastogeer et al. 2020).

To examine the role of both direct and indirect (diffuse) cross-kingdom interactions in microbiome community assembly, we characterized bacterial and fungal communities associated with switchgrass (*Panicum virgatum* L.) at 14 stands spanning mountain to coastal ecoregions of North Carolina, USA. We identified putative cross-kingdom direct and diffuse interactions using ecological network inference. Using variance partitioning analysis, we examined how direct and diffuse cross-kingdom interactions obtained from the networks contributed to community assembly relative to spatial and environmental factors. We hypothesized that direct interactions between bacterial and fungal taxa would be an important driver of leaf microbiome composition, because these are oligotrophic, low-diversity habitats and microbes tend to aggregate in more favorable microsites (Vorholt 2012; Andrews et al. 2002; Monier and Lindow 2004; Lodge et al. 1996), making pairwise interactions more likely. In roots, however, we expected diffuse, indirect interactions to be more relevant to composition, because these are exudate-rich habitats with high microbial diversity, density, and activity (Grayston et al. 1997; Herman et al. 2006).

METHODS

Experimental design and study sites
Switchgrass (*Panicum virgatum* L.) was sampled at 14 sites across a 462-km range in North Carolina, USA, in September and October 2018, as described in Lee and Hawkes (Lee and Hawkes 2021a). Briefly, we sampled 8 plants per site in a stratified random design. At each plant, we collected 1) five healthy, undamaged leaves spaced throughout the plant canopy, and 2) roots and soils to 15-cm depth immediately adjacent to the north and south side of the plant. Samples were stored on ice for transport to the lab, where fine roots < 2 mm were separated from soils and washed. Subsamples of leaves, roots, and soils were used for biogeochemical analyses or frozen at -80 °C until DNA extraction.

Plant and soil properties
Plant and soil field measurements included plant height, basal width and length, and GPS location. In the lab, we measured soil pH, microbial biomass carbon (MBC), moisture and soil texture as described in Lee and Hawkes (Lee and Hawkes 2021b). Total soil C and N were measured by combustion at the North Carolina State University Environmental and Agricultural Testing Service. Other soil nutrients (P, K) were measured at the NC Department of Agriculture and Consumer Services. Climate variables included mean annual precipitation (MAP) and mean annual temperature (MAT) based on 30-year normal ("PRISM Climate Group" 2014) .

Microbial community composition
We extracted DNA from 40-50 mg of leaves or roots using the Synergy 2.0 Plant DNA Extraction Kit (OPS Diagnostics, Lebanon, New Jersey, USA). Two leaf samples were excluded because they were senesced (brown) at collection. Amplicons were generated for bacteria in the 16S rRNA

region using the primers 515F and 806R (Caporaso et al. 2011) and for fungi in the 5.8S and ITS2 rRNA region with the primers 5.8S_FUN and ITS4_FUN (Taylor et al. 2016), modified with Illumina adapters. PCR reactions included 1× KAPA Taq Ready Mix (Roche, Pleasanton, CA, USA), 0.2 µM of each primer, 10 ng µL^{-1} of DNA template, and water up to 25 µL. For 16S, we also added 0.25 µM each of universal peptide nucleic acid (PNA) blockers for host mitochondria and plastids (Lundberg et al. 2013). For ITS, we added 0.5 mg mL-1 bovine serum albumin and 15 µM of a custom PNA designed for *P. virgatum* (Lee and Hawkes 2021a). Amplicon fragments <50 bp in size were removed using Agencourt AMpure XP beads in a ratio of 1:1.8 (Beckman Coulter, Indianapolis, IN, USA). Negative controls included one pooled blank from DNA extractions and two PCR blanks. Libraries were prepared using the Nextera XT Index Kit (Illumina, San Diego, CA, USA) and submitted for paired-end (2 x 250) sequencing on Illumina MiSeq v2 in the Genomic Sciences Laboratory at North Carolina State University (Raleigh, NC, USA). Sequences are available on the NCBI SRA under BioProject PRJNA648664.

Primers were removed with *cutadapt* v3.3 (Martin 2011). Amplicon sequence variants were identified using *dada2* v1.10.1, (Callahan et al. 2016) with maxEE=c(2,5) and otherwise default settings. Seven leaf samples and one root sample were omitted from the dataset due to insufficient 16S or ITS reads (< 150). The resulting ASVs were curated using *lulu* v0.1.0 (Frøslev et al. 2017) with default settings. Fourteen contaminants were removed from the 16S dataset based on analysis of negative controls with *decontam* v1.10.0 (Davis et al. 2018), which led to the removal of 9 samples with < 150 reads. Initial taxonomic identification of 16S ASVs relied on SILVA v132 (Glöckner et al. 2017) and ITS were based on UNITE (Nilsson et al. 2019). We then removed singleton ASVs and any ASVs that were not identified as bacteria or fungi in the 16S and ITS data, respectively. To improve ASV identification and generate phylogenies, we used T-BAS v2.3 to place both 16S and ITS sequences on reference trees (Carbone et al. 2019). The resulting tables included 16,296 bacterial ASVs from 323 samples and 932 fungal ASVs from 332 samples (Figure S1).

Statistics
To determine whether to assess the microbial communities from leaf and root compartments together or separately, we first evaluated the similarity of the microbial communities using redundancy analysis on the centered log-ratio transformed data for both bacterial and fungal communities. We tested how plant compartment affected fungal and bacterial community composition using residual randomized permutation procedures (RRPP v.1.2.1; Collyer and Adams 2018). Consistent with previous analyses (Lee and Hawkes 2021a), we found that the leaf and root microbiota had minimal overlap (P = 0.001, Table S1; Figures S2). Therefore, we chose to investigate potential drivers of community assembly for the leaf and root compartments separately.

In order to incorporate biotic interactions in community analyses, we first inferred cross-kingdom ecological association networks from which direct and diffuse interactions could be defined for the leaf and root communities (*SpiecEasi* v1.1.2 and *NetCoMi* v1.1.0; Kurtz et al. 2015; Peschel et al. 2021). Sparse inverse covariance estimation for ecological association inference (SPIEC-EASI) improves upon correlation methods, such as co-occurrence or joint

species distribution models, without relying on pairwise interactions. Instead, SPIEC-EASI acknowledges the conditional states of all nodes such that the resulting inverse covariance matrix contains only values for conditionally dependent (direct) interactions, while the conditionally independent (indirect) interactions are near zero (Kurtz et al. 2015). Prior to network construction, we filtered samples to only those present in both fungal and bacterial datasets and removed rare and low abundance taxa. Specifically, we 1) filtered leaf ASVs to include only those present in ≥10% of samples, 2) filtered root ASVs to only those present in either ≥10% of samples for fungi or ≥25% for bacteria, and 3) retained only those ASVs with >1% relative abundance. Filtering by prevalence and abundance resulted in 235 out of 1,002 bacterial ASVs and 114 out of 215 fungal ASVs in leaf communities and 388 out of 11,167 bacterial ASVs and 75 out of 730 fungal ASVs in root communities. The filtering did not remove any sites from the analysis, and overall microbial abundance patterns across sites were comparable before and after filtering. Networks were created for each site and filtered to include only high-confidence edges, where the edge-wise variability estimate was >0.8, indicating that the edge appeared in 80% of subsampled datasets. We performed a union of the site-level ASV networks and their putative interactions and inferred final networks for roots and leaves (Figure 1, Table S2; *Cytoscape* v3.10.1, Shannon et al. 2003).

We defined two categories of cross-kingdom interactions: direct and diffuse. First, directly interacting taxa were defined by network edges connecting bacterial and fungal nodes. Second, diffuse interactions were represented in two ways, 1) by keystone hub taxa and 2) by alpha diversity. Keystone hub taxa—defined as network nodes in the 75th percentile of degree and eigenvector centrality within each kingdom—represent those that are most influential to the network and may be necessary to community assembly without directly interacting with the other kingdom (Berry and Widder 2014). As an alternative to the keystone model, alpha diversity—defined as the richness of either bacteria or fungi in each sample—may better represent the set of all possible interactions between the kingdoms (Lee et al. 2019; Fiore-Donno et al. 2024). However, because alpha diversity never explained more than 1.5% of the variation in community composition, these results are only presented in the supplement (Tables S11 and S14, Figure S3). To better understand the bacterial and fungal taxa identified as direct or diffuse interactors, we tested for phylogenetic conservation of interaction type with Pagel's λ (*geiger* v2.0.10, *ape* v5.6-2, *phytools* v1.0-1; Pennell et al. 2014; Paradis and Schliep 2019; Revell 2012). We also conducted literature searches to identify putative ecological roles of all direct and diffuse ASVs (Tables S3-6).

To address our hypotheses, we examined whether 1) interactions with bacteria (direct or diffuse) improved our ability to explain fungal community assembly and 2) whether interactions with fungi (direct or diffuse) improved our ability to explain bacterial community assembly. To do this, we used variance partitioning analysis (*vegan* v2.6-4; Oksanen et al. 2001) with either bacterial or fungal community composition as the dependent variable, represented as the Hellinger-transformed ASV count matrix (Peres-Neto et al. 2006; Legendre and Gallagher 2001). Independent predictors were matrices of environmental variables, spatial variables, and a set of biotic interactions (either direct, diffuse keystone hubs, or diffuse alpha diversity). Each variance partitioning analysis included the same environmental and spatial variables (Table S7).

Environmental variables included climate (MAP, MAT), plant size (height), microbial biomass, soil macronutrients (C, N, P, K), and soil properties (pH, texture, moisture). Spatial variables were stand area and sampling location (latitude, longitude). All environmental and spatial variables were standardized by scaling to zero mean and unit variance using the *decostand* function from the *vegan* R package (v2.6-4; Oksanen et al. 2001).

Each variance partitioning analysis was run four times: 1) without biotic interactions, then with biotic interactions represented by a secondary ASV matrix from the opposite kingdom including either 2) the directly interacting ASVs, 3) the diffusely interacting ASVs represented by keystone taxa, or 4) the diffusely interacting ASVs represented by alpha diversity. The sensitivity of these results to different network definitions is provided in supplemental Tables S9-14. All testable fractions from the variance partitioning analyses were analyzed for significance with redundancy analysis (RDA) models and *P*-values were calculated based on a distribution of R^2 values from permuted data (*vegan* v2.6-4; Oksanen et al. 2001). Shared variation fractions between environmental factors and biotic interactions were further investigated using permanova (*RRPP* v1.2.1; Collyer and Adams 2018) to determine environmental variables significantly associated ($P < 0.05$) with the ASVs. A subset of the significant environmental variables with the highest R^2 values were further examined using gradient analysis (*TITAN2* v2.4.3; Baker and King 2010) to characterize associated changes in ASV abundance.

RESULTS

Identification of direct and diffuse interactions
Direct interactions.—Putative cross-kingdom direct interactions identified by ecological association networks included 69 taxa in leaves (37 bacteria and 32 fungi) and 68 taxa in roots (36 bacteria and 32 fungi), representing 22% and 15% of all taxa included in the networks, respectively (Figure 1). These directly interacting taxa were phylogenetically diverse, covering 26 different bacterial orders and 19 different fungal orders (Figure 1). In both the leaf and root networks, a similar number of direct edges were found with positive association estimates indicating co-occurrence between fungal and bacterial taxa (leaf: 24, root: 18) and negative associations indicating inverse relationships between taxa (leaf: 19, root: 20, respectively). The distribution of ASV abundances across sites indicated that both bacteria and fungi with direct cross-kingdom interactions ranged from rare (2-3 sites) to prevalent (8-14 sites) (Figure S4). Most taxa involved in cross-kingdom direct interactions were only connected to one other ASV (78%), but 21% had multiple direct links (Table S16). For example, in the leaf network, an Alpha-Proteobacteria ASV in the genus *Bosea*, which is known for plant growth promotion (Table S3), shared edges with ASVs in three different fungal genera: two basidiomycetous yeasts (*Fibulobasidium*, *Bulleribasidium*) and one agaric fungus (*Chrysomycena*). Similarly, in the root network, a bacterial ASV in the genus *Streptomyces*, also known for promoting plant growth, shared edges with ASVs an ascomycetous yeast (*Citeromyces* sp.) and a widespread basidiomycetous fungus (*Hydnum* sp.) (Table S4).

Diffuse interactions.—Keystone taxa representing putative diffuse interactions included 25 bacteria and 9 fungi identified for the leaf community and 69 bacteria and 3 fungi for the root community (Figure 1). Keystone ASVs were generally prevalent, with bacterial keystone taxa found in 7 to 14 sites and fungal keystones found in 4 to 14 sites (Figure S4). At the genus-level, the most frequently occurring keystone bacteria in leaves were Actinomycetes represented by three *Nocardiodes* ASVs that are potentially associated with plant growth promotion (Table S16). The most common keystone bacteria in the root were ASVs in the Alpha-Proteobacteria (*Filomicrobium* sp.)*,* Acidobacteria (*Koribacter* sp.)*,* and Actinomycetes (*Mycobacterium* sp.; a putative plant growth promoter) (Table S16). Of the 9 keystone fungal ASVs in the leaf network, 7 were Ascomycota (5 Dothideomycetes, 2 Eurotiomycetes) and 2 were Basidiomycota yeasts (Tremellomycetes, Spiculogloeomycetes). In roots, 2 of the 3 keystone fungi were Glomeromycetes—arbuscular mycorrhizas in the genus *Paraglomus*—and the third ASV was a basidiomycetous pathogen from the genus *Shivasia* (Ustilaginomycetes) (Table S16).

Diversity and phylogenetic relationships.—As described above, both direct and diffuse interactors *within* each kingdom included phylogenetically diverse taxa with a range of putative ecological roles (Figure 1, Table S3-6). No phylogenetic signals were detected for interaction type (Table S15), so further analyses did not account for phylogeny. This is supported in the observed composition patterns, where neither bacteria nor fungi had much taxonomic overlap below the level of order. For example, Dothideomycetes fungal ASVs were found across three out of four conditions (direct leaf, diffuse leaf, and direct root), but included 13 different genera with the dominant ones being *Sphaerellopsis* and *Cumuliphoma* (both potentially pathogenic) in leaves and *Poaceascoma* (potentially endophytic) in roots based on relative abundance.

Contributions of cross-kingdom biotic interactions to community assembly
In the absence of biotic interactions, environmental variables explained 27.5% and 18.6% of fungal and 22.2% and 21.9% of bacterial leaf and root communities, respectively (Figure 2, Table S8). When biotic interactions were considered, standalone environmental contributions to community composition declined by ~15-20% in leaves and ~8-10% in roots (Figure 2, Table S8). In all analyses with and without biotic interactions, spatial variables and their intersections explained less than ~10% of the variation.

Putative direct cross-kingdom interactions were better predictors than diffuse interactions in most cases, explaining 2 to 6 times more variation in community composition of foliar fungi (10.4% direct vs. 2.6% diffuse), foliar bacteria (11.4% direct vs. 5.6% diffuse), and root bacteria (4.6% direct vs. 0.7% diffuse; Table S8, Figure 2). The opposite pattern was observed only for root fungi, where diffuse interactions with bacteria explained more than twice as much of the variation in fungal community composition (8.4%) compared to direct interactions (4.0%).

The joint effects of environment and biotic interactions also helped to explain community patterns. These joint effects were dominant in leaves when considering direct interactions for both fungi (15.2%) and bacteria (14.4%) and in root fungal communities when considering either direct (8.7%) or diffuse (10.7%) cross-kingdom interactions (Table S8). In all other cases,

the joint effects of environment and cross-kingdom interactions explained less than environment alone (Table S8).

The shared effects of environment and biotic interactions might be due to correlations between the interacting ASVs and environmental variation, as indicated by the PERMANOVA results (Table S17). Based on environmental gradient analysis, many foliar bacteria and fungi identified as direct and diffuse interactors were associated with changes in soil texture (% sand) (Figure 3, Tables S18). Additionally, foliar bacteria identified as direct interactors were associated with soil pH and plant basal area, whereas foliar fungal direct interactors were associated with MAT (Tables S18, Figure S6). In roots, bacterial and fungal ASVs involved in putative interactions tracked different soil properties–soil pH and soil %C or soil %N, respectively (Figure 3). Fungal taxa involved in direct interactions also tracked MAP (Figure 3) and one fungal diffuse (keystone) taxon tracked plant height (Table S19, Figure S6).

DISCUSSION

Cross-kingdom interactions between bacteria and fungi represented almost a quarter of all links in their networks and were an important driver of microbial community composition similar in importance to environmental and spatial predictors. This importance increased even further when considering the joint effects of environment and biotic interactions, which highlights the complexity of factors influencing microbial community composition and the need to expand the classic approach to describing community assembly. Our findings add to the growing pool of research studies that have found biotic interactions to be important in explaining community structure (Lee et al. 2019; Fiore-Donno et al. 2024; García-Girón et al. 2020), while offering a method for better defining those biotic interactions in ecological terms.

Our original hypothesis, that cross-kingdom direct interactions would be more important than diffuse interactions in lower diversity leaf environments, was largely supported. Microorganisms tend to cluster in microhabitats on leaves, enabling direct competition for space and resources or other antagonistic or mutualistic interactions (Monier and Lindow 2004; Chaudhry et al. 2020). For example, several studies have shown the importance of bacteria-bacteria interactions in *Arabidopsis*—these interactions tend to be negative, supported by identification of antimicrobial compound production and shifts in community composition that can lead to damaged plant tissue (Chen et al. 2020; Schäfer et al. 2022). Other studies have identified cross-kingdom interactions in plants where bacteria may have an antagonistic effect on fungal pathogens. In one case, Kamensky et al. (Kamensky et al. 2003) demonstrated that a gram-negative bacterial strain was able to protect the plant host from multiple necrotrophic fungi and identified production of several anti-fungal compounds as the most likely mechanism. In the current study, the reported ecological roles of the taxa involved in putative direct interactions suggest the potential for either competition (negative edges) or facilitation (positive edges) between bacteria and plant pathogenic fungi. Putative direct interactions identified from ecological association networks may point to a key subset of ecologically important taxa that can be prioritized for further study.

In roots, we originally expected cross-kingdom diffuse interactions to be better predictors than direct interactions given the typically high diversity of root microbial communities. However, this was only the case for root fungi and their diffuse interactions with bacteria. This outcome aligns with their observed alpha diversities—bacterial richness in roots was six times higher than in leaves, but fungal richness was comparatively lower and did not change between the plant compartments. The higher bacterial richness we found in roots may increase the potential for diffuse interactions or modifications to the environment. Supporting this, evidence indicates that selective reductions in soil microbial diversity result in expansions of previously rare taxa (Romdhane et al. 2022) and modifications of specific biogeochemical processes, potentially altering local environmental conditions (Peter et al. 2011; Trivedi et al. 2019). Alternatively, the larger number of bacteria may have allowed for more robust identification of keystone taxa compared to the fungi.

Multiple studies have reported that keystone taxa in microbial networks tend to be important for community structure, ecosystem functioning, and interactions with abiotic and biotic factors (Wu et al. 2023; Yang et al. 2021; Hu et al. 2024; Agler et al. 2016). This approach to defining keystones has additional support from Berry and Widder (Berry and Widder 2014), who demonstrated that high degree, closeness centrality, and transitivity corresponded to "keystoneness." We found keystone taxa to be a good representation of diffuse interactions, likely because defining keystone taxa as ASVs that were highly connected (degree) and had a strong influence (eigenvector centrality) partly captured their role in determining overall community structure (Cottee-Jones and Whittaker 2012; Jones et al. 1994). In contrast, we found that alpha diversity had poor explanatory power for community assembly, perhaps because using overall species richness to represent diffuse interactions can dilute the effects of keystone taxa. This differs from other studies that reported modest effects of biotic interactions defined as species richness on other taxonomic group richness (Lee et al. 2019), though Fiore-Donno et. al. (Fiore-Donno et al. 2024) did find seasonal effects on the relative importance of species richness on community composition which may be a reflection of the important intersection between environmental and biotic factors that we saw.

Up to 15% of the variation in microbial community structure was present in the joint effects of environment and biotic interactions, likely caused by both direct and diffuse ASVs that tracked specific environmental conditions. Although there were no universal patterns across taxa, there were clear trends that suggest potential ecological processes. For example, 70-75% of leaf ASVs were either positively (direct interactors) or negatively (diffuse interactors) associated with soil sand content, suggesting that soil texture may influence source pools, local dispersal, or recruitment to the plant host. Sand content affects multiple soil properties, including nutrient and water availability, for which it may serve as a proxy, but we also found some taxa that were correlated with soil nutrients or rainfall. In roots, out of the five directly interacting fungi associated with both MAP and nitrogen, three were associated with either high MAP and lower nitrogen or lower MAP and high nitrogen, suggesting that these are key ecological filters. In most cases, environmental factors are likely driving ASV distributions, which can affect the structure of microbial co-occurrence networks by increasing or reducing potential interactions

(Lekberg et al. 2021). Alternatively, biotic interactions can lead to changes in environmental factors, such as altering available nutrients via the formation and degradation of soil organic matter, that are important to both the plant host and microbial community (Sokol et al. 2022). Understanding the relationships between abiotic factors and microbial species distributions can offer insight into what drives the correlations underlying joint effects.

Inferring biotic interactions from ecological association networks based on amplicon sequencing of rDNA necessarily means that these interactions are putative. Although this approach is not purely correlational, it is still dependent on distributional co-occurrence patterns captured at a single time point rather than explicit tests of interaction. Similarly, we identified putative ecological roles of taxa from trait databases and cases reported in the literature, which we note are particularly problematic at the genus level when both conspecifics and congeners can exhibit multiple or context-dependent trophic strategies. The identification of putative interactions and ecological roles might be improved by using methods that characterize active bacteria and fungi instead of the entire rDNA community (Emerson et al. 2017), by measuring the community over time to identify how consistent these putative interactions might be (Fiore-Donno et al. 2024; Argiroff et al. 2024; Agler et al. 2016), or by employing multi-omics networks that model metabolic interactions (Muller et al. 2018). Alternatively, stable isotope probing could identify active taxa that use the same substrates and likely compete (Dumont and Murrell 2005; Prosser et al. 2006), taxa turnover using labeled oxygen (Schwartz et al. 2016), or taxa that are part of a trophic web (Lueders et al. 2004).

Nevertheless, our results were robust to the changes in network parameters and keystone threshold. Variance partitioning results at the different edge-wise variability estimates were all within a 10% range of total variation explained. The largest difference observed with increasing stringency of the edge-wise variability estimate was in network connectedness, evidenced by the dramatic decrease in the number of edges especially those that were cross-kingdom. In the current study, the majority of keystone taxa were retained across different edge-wise variability estimates, though there were unique keystones identified in each due to the change in nodes and edges maintained at increasing stringency. The minimal sensitivity of overall outcomes to these changes suggests that putative direct and diffuse interactions identified from networks are largely internally consistent and can improve our assessment of community drivers.

The incorporation of biotic interactions into investigating microbial community assembly is a relatively recent and growing trend (Dastogeer et al. 2020) that is supported by our results. We took this a step further by defining different types of interactions—direct and diffuse—that proved to be important sources of variation in microbial community composition aligned with expectations from ecological theory. However, it is important to note that there was still 54-74% residual variation that was unexplained by our models. These efforts are hampered by reliance on putative interactions and difficulty in identifying true interactions in complex microbial communities. To partly overcome this, others have successfully used both taxonomic and functional characteristics of taxa to better capture interactions in well-characterized systems (Li et al. 2023). Moving forward, community assembly models will be improved by a

greater understanding of how we can better estimate and represent cross-kingdom biotic interactions.

ACKNOWLEDGEMENTS
For assistance with laboratory and field work, we thank N. Yang, B. Whitaker, I. McBryde, J. Czel, E. Hardy, and M. Alvarez. Site access was generously provided by NCSU, North Carolina Department of Agriculture and Consumer Services, BASF Corp., Weyerhaeuser Co., C. Wilson, and E. Deal. The manuscript was improved by comments from M. Cubeta. This research was supported by a DOE Genomic Sciences Program SFA (SCW1039) subaward from Lawrence Livermore National Lab and startup funding from North Carolina State University to CVH.

AUTHOR CONTRIBUTIONS
The experiment was designed by CVH. RAH analyzed the data. RAH and CVH wrote the manuscript. MRL led field and lab work and carried out the bioinformatics analysis of amplicon sequencing data. All authors edited the manuscript.

DATA and CODE AVAILABILITY
Raw sequence data are available on the NCBI Short Read Archive under BioProject PRJNA648664. Site and sample data, as well as R code, are available on GitHub.

REFERENCES

Agler, Matthew T., Jonas Ruhe, Samuel Kroll, et al. 2016. "Microbial Hub Taxa Link Host and Abiotic Factors to Plant Microbiome Variation." *PLOS Biology* 14 (1): e1002352. https://doi.org/10.1371/journal.pbio.1002352.

Aguilar-Trigueros, Carlos A., Jeff R. Powell, Ian C. Anderson, Janis Antonovics, and Matthias C. Rillig. 2014. "Ecological Understanding of Root-Infecting Fungi Using Trait-Based Approaches." *Trends in Plant Science* 19 (7): 432–38. https://doi.org/10.1016/j.tplants.2014.02.006.

Andrews, John H, Russell N Spear, and Erik V Nordheim. 2002. "Population Biology of Aureobasidium Pullulans on Apple Leaf Surfaces." *Canadian Journal of Microbiology* 48 (6): 500–513. https://doi.org/10.1139/w02-044.

Argiroff, William A., Alyssa A. Carrell, Dawn M. Klingeman, et al. 2024. "Seasonality and Longer-Term Development Generate Temporal Dynamics in the Populus Microbiome." *mSystems* 9 (3): e0088623. https://doi.org/10.1128/msystems.00886-23.

Baker, Matthew E., and Ryan S. King. 2010. "A New Method for Detecting and Interpreting Biodiversity and Ecological Community Thresholds." *Methods in Ecology and Evolution* 1 (1): 25–37. https://doi.org/10.1111/j.2041-210X.2009.00007.x.

Berry, David, and Stefanie Widder. 2014. "Deciphering Microbial Interactions and Detecting Keystone Species with Co-Occurrence Networks." *Frontiers in Microbiology* 5. https://doi.org/10.3389/fmicb.2014.00219.

Bever, James D., Scott A. Mangan, and Helen M. Alexander. 2015. "Maintenance of Plant Species Diversity by Pathogens." *Annual Review of Ecology, Evolution, and Systematics* 46 (Volume 46, 2015): 305–25. https://doi.org/10.1146/annurev-ecolsys-112414-054306.

Callahan, Benjamin J., Paul J. McMurdie, Michael J. Rosen, Andrew W. Han, Amy Jo A. Johnson, and Susan P. Holmes. 2016. "DADA2: High-Resolution Sample Inference from Illumina Amplicon Data." *Nature Methods* 13 (7): 581–83. https://doi.org/10.1038/nmeth.3869.

Caporaso, J. Gregory, Christian L. Lauber, William A. Walters, et al. 2011. "Global Patterns of 16S rRNA Diversity at a Depth of Millions of Sequences per Sample." *Proceedings of the National Academy of Sciences* 108 (supplement_1): 4516–22. https://doi.org/10.1073/pnas.1000080107.

Carbone, Ignazio, James B. White, Jolanta Miadlikowska, et al. 2019. "T-BAS Version 2.1: Tree-Based Alignment Selector Toolkit for Evolutionary Placement of DNA Sequences and Viewing Alignments and Specimen Metadata on Curated and Custom Trees." *Microbiology Resource Announcements* 8 (29): 10.1128/mra.00328-19. https://doi.org/10.1128/mra.00328-19.

Chaudhry, Vasvi, Paul Runge, Priyamedha Sengupta, Gunther Doehlemann, Jane E Parker, and Eric Kemen. 2020. "Shaping the Leaf Microbiota: Plant–Microbe–Microbe Interactions." *Journal of Experimental Botany* 72 (1): 36–56. https://doi.org/10.1093/jxb/eraa417.

Chen, Tao, Kinya Nomura, Xiaolin Wang, et al. 2020. "A Plant Genetic Network for Preventing Dysbiosis in the Phyllosphere." *Nature* 580 (7805): 653–57. https://doi.org/10.1038/s41586-020-2185-0.

Collyer, Michael L., and Dean C. Adams. 2018. "RRPP: An r Package for Fitting Linear Models to High-Dimensional Data Using Residual Randomization." *Methods in Ecology and Evolution* 9 (7): 1772–79. https://doi.org/10.1111/2041-210X.13029.

Cottee-Jones, Henry Eden W., and Robert J. Whittaker. 2012. "Perspective: The Keystone Species Concept: A Critical Appraisal." *Frontiers of Biogeography* 4 (3). https://doi.org/10.21425/F5FBG12533.

Dastogeer, Khondoker M. G., Farzana Haque Tumpa, Afruja Sultana, Mst Arjina Akter, and Anindita Chakraborty. 2020. "Plant Microbiome–an Account of the Factors That Shape Community

Composition and Diversity." *Current Plant Biology*, Plant Interactions with Microbes and Environment, vol. 23 (September): 100161. https://doi.org/10.1016/j.cpb.2020.100161.

Davis, Nicole M., Diana M. Proctor, Susan P. Holmes, David A. Relman, and Benjamin J. Callahan. 2018. "Simple Statistical Identification and Removal of Contaminant Sequences in Marker-Gene and Metagenomics Data." *Microbiome* 6 (1): 226. https://doi.org/10.1186/s40168-018-0605-2.

Deveau, Aurélie, Gregory Bonito, Jessie Uehling, et al. 2018. "Bacterial–Fungal Interactions: Ecology, Mechanisms and Challenges." *FEMS Microbiology Reviews* 42 (3): 335–52. https://doi.org/10.1093/femsre/fuy008.

Duan, Shilong, Stéphane Declerck, Gu Feng, and Lin Zhang. 2023. "Hyphosphere Interactions between and Promote Carbon–Phosphorus Exchange at the Peri-Arbuscular Space In." *Environmental Microbiology* 25 (4): 867–79. https://doi.org/10.1111/1462-2920.16333.

Dumont, Marc G., and J. Colin Murrell. 2005. "Stable Isotope Probing — Linking Microbial Identity to Function." *Nature Reviews Microbiology* 3 (6): 499–504. https://doi.org/10.1038/nrmicro1162.

Emerson, Joanne B., Rachel I. Adams, Clarisse M. Betancourt Román, et al. 2017. "Schrödinger's Microbes: Tools for Distinguishing the Living from the Dead in Microbial Ecosystems." *Microbiome* 5 (1): 86. https://doi.org/10.1186/s40168-017-0285-3.

Emmett, Bryan D, Véronique Lévesque-Tremblay, and Maria J Harrison. 2021. "Conserved and Reproducible Bacterial Communities Associate with Extraradical Hyphae of Arbuscular Mycorrhizal Fungi." *The ISME Journal* 15 (8): 2276–88. https://doi.org/10.1038/s41396-021-00920-2.

Fiore-Donno, Anna Maria, Jule Freudenthal, Mathilde Borg Dahl, Christian Rixen, Tim Urich, and Michael Bonkowski. 2024. "Biotic Interactions Explain Seasonal Dynamics of the Alpine Soil Microbiome." *ISME Communications* 4 (1): ycae028. https://doi.org/10.1093/ismeco/ycae028.

Frey-Klett, P., P. Burlinson, A. Deveau, M. Barret, M. Tarkka, and A. Sarniguet. 2011. "Bacterial-Fungal Interactions: Hyphens between Agricultural, Clinical, Environmental, and Food Microbiologists." *Microbiology and Molecular Biology Reviews* 75 (4): 583–609. https://doi.org/10.1128/mmbr.00020-11.

Frøslev, Tobias Guldberg, Rasmus Kjøller, Hans Henrik Bruun, et al. 2017. "Algorithm for Post-Clustering Curation of DNA Amplicon Data Yields Reliable Biodiversity Estimates." *Nature Communications* 8 (1): 1188. https://doi.org/10.1038/s41467-017-01312-x.

Gallardo-Navarro, Óscar Adrián, and Moisés Santillán. 2019. "Three-Way Interactions in an Artificial Community of Bacterial Strains Directly Isolated From the Environment and Their Effect on the System Population Dynamics." *Frontiers in Microbiology* 10 (November). https://doi.org/10.3389/fmicb.2019.02555.

García-Girón, Jorge, Jani Heino, Francisco García-Criado, Camino Fernández-Aláez, and Janne Alahuhta. 2020. "Biotic Interactions Hold the Key to Understanding Metacommunity Organisation." *Ecography* 43 (8): 1180–90. https://doi.org/10.1111/ecog.05032.

Giauque, Hannah, and Christine V. Hawkes. 2016. "Historical and Current Climate Drive Spatial and Temporal Patterns in Fungal Endophyte Diversity." *Fungal Ecology* 20 (April): 108–14. https://doi.org/10.1016/j.funeco.2015.12.005.

Glöckner, Frank Oliver, Pelin Yilmaz, Christian Quast, et al. 2017. "25 Years of Serving the Community with Ribosomal RNA Gene Reference Databases and Tools." *Journal of Biotechnology*, Bioinformatics Solutions for Big Data Analysis in Life Sciences presented by the German Network for Bioinformatics Infrastructure, vol. 261 (November): 169–76. https://doi.org/10.1016/j.jbiotec.2017.06.1198.

Grayston, S. J., D. Vaughan, and D. Jones. 1997. "Rhizosphere Carbon Flow in Trees, in Comparison with Annual Plants: The Importance of Root Exudation and Its Impact on Microbial Activity and

Nutrient Availability." *Applied Soil Ecology* 5 (1): 29–56. https://doi.org/10.1016/S0929-1393(96)00126-6.

Herman, D. J., K. K. Johnson, C. H. Jaeger, E. Schwartz, and M. K. Firestone. 2006. "Root Influence on Nitrogen Mineralization and Nitrification in Avena Barbata Rhizosphere Soil." *Soil Science Society of America Journal* 70 (5): 1504–11. https://doi.org/10.2136/sssaj2005.0113.

Hestrin, Rachel, Megan Kan, Marissa Lafler, et al. 2022. "Plant-Associated Fungi Support Bacterial Resilience Following Water Limitation." *The ISME Journal* 16 (12): 2752–62. https://doi.org/10.1038/s41396-022-01308-6.

Hu, Minjie, Jordi Sardans, Dongyao Sun, et al. 2024. "Microbial Diversity and Keystone Species Drive Soil Nutrient Cycling and Multifunctionality Following Mangrove Restoration." *Environmental Research* 251 (June): 118715. https://doi.org/10.1016/j.envres.2024.118715.

Jacoby, Richard P, and Stanislav Kopriva. 2019. "Metabolic Niches in the Rhizosphere Microbiome: New Tools and Approaches to Analyse Metabolic Mechanisms of Plant–Microbe Nutrient Exchange." *Journal of Experimental Botany* 70 (4): 1087–94. https://doi.org/10.1093/jxb/ery438.

Jiang, Duo, Courtney R. Armour, Chenxiao Hu, et al. 2019. "Microbiome Multi-Omics Network Analysis: Statistical Considerations, Limitations, and Opportunities." *Frontiers in Genetics* 10 (November). https://doi.org/10.3389/fgene.2019.00995.

Jones, Clive G., John H. Lawton, and Moshe Shachak. 1994. "Organisms as Ecosystem Engineers." *Oikos* 69 (3): 373–86. https://doi.org/10.2307/3545850.

Kamensky, Merav, Marianna Ovadis, Ilan Chet, and Leonid Chernin. 2003. "Soil-Borne Strain IC14 of *Serratia Plymuthica* with Multiple Mechanisms of Antifungal Activity Provides Biocontrol of *Botrytis Cinerea* and *Sclerotinia Sclerotiorum* Diseases." *Soil Biology and Biochemistry* 35 (2): 323–31. https://doi.org/10.1016/S0038-0717(02)00283-3.

Kurtz, Zachary D., Christian L. Müller, Emily R. Miraldi, Dan R. Littman, Martin J. Blaser, and Richard A. Bonneau. 2015. "Sparse and Compositionally Robust Inference of Microbial Ecological Networks." *PLOS Computational Biology* 11 (5): e1004226. https://doi.org/10.1371/journal.pcbi.1004226.

Lai, Hao Ran, Kwek Yan Chong, Alex Thiam Koon Yee, Margaret M. Mayfield, and Daniel B. Stouffer. 2022. "Non-Additive Biotic Interactions Improve Predictions of Tropical Tree Growth and Impact Community Size Structure." *Ecology* 103 (2): e03588. https://doi.org/10.1002/ecy.3588.

Layeghifard, Mehdi, David M. Hwang, and David S. Guttman. 2017. "Disentangling Interactions in the Microbiome: A Network Perspective." *Trends in Microbiology* 25 (3): 217–28. https://doi.org/10.1016/j.tim.2016.11.008.

Lee, Charles K., Daniel C. Laughlin, Eric M. Bottos, et al. 2019. "Biotic Interactions Are an Unexpected yet Critical Control on the Complexity of an Abiotically Driven Polar Ecosystem." *Communications Biology* 2 (1): 1–10. https://doi.org/10.1038/s42003-018-0274-5.

Lee, Kiseok Keith, Hyun Kim, and Yong-Hwan Lee. 2022. "Cross-Kingdom Co-Occurrence Networks in the Plant Microbiome: Importance and Ecological Interpretations." *Frontiers in Microbiology* 13 (July): 953300. https://doi.org/10.3389/fmicb.2022.953300.

Lee, Marissa R., and Christine V. Hawkes. 2021a. "Plant and Soil Drivers of Whole-Plant Microbiomes: Variation in Switchgrass Fungi from Coastal to Mountain Sites." *Phytobiomes Journal* 5 (1): 69–79. https://doi.org/10.1094/pbiomes-07-20-0056-fi.

Lee, Marissa R., and Christine V. Hawkes. 2021b. "Widespread Co-Occurrence of Sebacinales and Arbuscular Mycorrhizal Fungi in Switchgrass Roots and Soils Has Limited Dependence on Soil Carbon or Nutrients." *Plants People Planet* 3 (5). https://doi.org/10.1002/ppp3.10181.

Legendre, Pierre, and Eugene D. Gallagher. 2001. "Ecologically Meaningful Transformations for Ordination of Species Data." *Oecologia* 129 (2): 271–80. https://doi.org/10.1007/s004420100716.

Lekberg, Ylva, Carlos A. Arnillas, Elizabeth T. Borer, et al. 2021. "Nitrogen and Phosphorus Fertilization Consistently Favor Pathogenic over Mutualistic Fungi in Grassland Soils." *Nature Communications* 12 (1): 3484. https://doi.org/10.1038/s41467-021-23605-y.

Lobanov, Alexander, Samantha Dyckman, Helen Kurkjian, and Babak Momeni. 2023. "Spatial Structure Favors Microbial Coexistence except When Slower Mediator Diffusion Weakens Interactions." *eLife* 12 (June): e82504. https://doi.org/10.7554/eLife.82504.

Lodge, D. Jean, P. J. Fisher, and B. C. Sutton. 1996. "Endophytic Fungi of Manilkara Bidentata Leaves in Puerto Rico." *Mycologia* 88 (5): 733–38. https://doi.org/10.2307/3760967.

Lueders, Tillmann, Bianca Wagner, Peter Claus, and Michael W. Friedrich. 2004. "Stable Isotope Probing of rRNA and DNA Reveals a Dynamic Methylotroph Community and Trophic Interactions with Fungi and Protozoa in Oxic Rice Field Soil." *Environmental Microbiology* 6 (1): 60–72. https://doi.org/10.1046/j.1462-2920.2003.00535.x.

Lundberg, Derek S., Scott Yourstone, Piotr Mieczkowski, Corbin D. Jones, and Jeffery L. Dangl. 2013. "Practical Innovations for High-Throughput Amplicon Sequencing." *Nature Methods* 10 (10): 999–1002. https://doi.org/10.1038/nmeth.2634.

Martin, Marcel. 2011. "Cutadapt Removes Adapter Sequences from High-Throughput Sequencing Reads." *EMBnet.Journal* 17 (1): 1. https://doi.org/10.14806/ej.17.1.200.

Mészárošová, Lenka, Eliška Kuťáková, Petr Kohout, Zuzana Münzbergová, and Petr Baldrian. 2024. "Plant Effects on Microbiome Composition Are Constrained by Environmental Conditions in a Successional Grassland." *Environmental Microbiome* 19 (1): 8. https://doi.org/10.1186/s40793-024-00550-z.

Mickalide, Harry, and Seppe Kuehn. 2019. "Higher-Order Interaction between Species Inhibits Bacterial Invasion of a Phototroph-Predator Microbial Community." *Cell Systems* 9 (6): 521-533.e10. https://doi.org/10.1016/j.cels.2019.11.004.

Miquel Guennoc, Cora, Christophe Rose, Jessy Labbé, and Aurélie Deveau. 2018. "Bacterial Biofilm Formation on the Hyphae of Ectomycorrhizal Fungi: A Widespread Ability under Controls?" *FEMS Microbiology Ecology* 94 (7): fiy093. https://doi.org/10.1093/femsec/fiy093.

Moen, Jon. 1989. "Diffuse Competition: A Diffuse Concept." *Oikos* 54 (2): 260–63. https://doi.org/10.2307/3565280.

Monier, J.-M., and S. E. Lindow. 2004. "Frequency, Size, and Localization of Bacterial Aggregates on Bean Leaf Surfaces." *Applied and Environmental Microbiology* 70 (1): 346–55. https://doi.org/10.1128/AEM.70.1.346-355.2004.

Morin, Manon A., Anneliese J. Morrison, Michael J. Harms, and Rachel J. Dutton. 2022. "Higher-Order Interactions Shape Microbial Interactions as Microbial Community Complexity Increases." *Scientific Reports* 12 (1): 22640. https://doi.org/10.1038/s41598-022-25303-1.

Muller, Emilie E. L., Karoline Faust, Stefanie Widder, Malte Herold, Susana Martínez Arbas, and Paul Wilmes. 2018. "Using Metabolic Networks to Resolve Ecological Properties of Microbiomes." *Current Opinion in Systems Biology*, • Regulatory and metabolic networks • Special Section: Single cell and noise, vol. 8 (April): 73–80. https://doi.org/10.1016/j.coisb.2017.12.004.

Nilsson, Rolf Henrik, Karl-Henrik Larsson, Andy F. S. Taylor, et al. 2019. "The UNITE Database for Molecular Identification of Fungi: Handling Dark Taxa and Parallel Taxonomic Classifications." *Nucleic Acids Research* 47 (D1): D259–64. https://doi.org/10.1093/nar/gky1022.

Nuccio, Erin E., Steven J. Blazewicz, Marissa Lafler, et al. 2022. "HT-SIP: A Semi-Automated Stable Isotope Probing Pipeline Identifies Cross-Kingdom Interactions in the Hyphosphere of Arbuscular Mycorrhizal Fungi." *Microbiome* 10 (1): 199. https://doi.org/10.1186/s40168-022-01391-z.

Oksanen, Jari, Gavin L. Simpson, F. Guillaume Blanchet, et al. 2001. "Vegan: Community Ecology Package." September 6. https://doi.org/10.32614/CRAN.package.vegan.

Pandit, A., A. Adholeya, D. Cahill, L. Brau, and M. Kochar. 2020. "Microbial Biofilms in Nature: Unlocking Their Potential for Agricultural Applications." *Journal of Applied Microbiology* 129 (2): 199–211. https://doi.org/10.1111/jam.14609.

Paradis, Emmanuel, and Klaus Schliep. 2019. "Ape 5.0: An Environment for Modern Phylogenetics and Evolutionary Analyses in R." *Bioinformatics (Oxford, England)* 35 (3): 526–28. https://doi.org/10.1093/bioinformatics/bty633.

Pennell, Matthew W., Jonathan M. Eastman, Graham J. Slater, et al. 2014. "Geiger v2.0: An Expanded Suite of Methods for Fitting Macroevolutionary Models to Phylogenetic Trees." *Bioinformatics (Oxford, England)* 30 (15): 2216–18. https://doi.org/10.1093/bioinformatics/btu181.

Peres-Neto, Pedro R., Pierre Legendre, Stéphane Dray, and Daniel Borcard. 2006. "Variation Partitioning of Species Data Matrices: Estimation and Comparison of Fractions." *Ecology* 87 (10): 2614–25. https://doi.org/10.1890/0012-9658(2006)87%255B2614:VPOSDM%255D2.0.CO;2.

Peschel, Stefanie, Christian L Müller, Erika von Mutius, Anne-Laure Boulesteix, and Martin Depner. 2021. "NetCoMi: Network Construction and Comparison for Microbiome Data in R." *Briefings in Bioinformatics* 22 (4): bbaa290. https://doi.org/10.1093/bib/bbaa290.

Peter, Hannes, Sara Beier, Stefan Bertilsson, Eva S Lindström, Silke Langenheder, and Lars J Tranvik. 2011. "Function-Specific Response to Depletion of Microbial Diversity." *The ISME Journal* 5 (2): 351–61. https://doi.org/10.1038/ismej.2010.119.

Pichler, Maximilian, and Florian Hartig. 2021. "A New Joint Species Distribution Model for Faster and More Accurate Inference of Species Associations from Big Community Data." *Methods in Ecology and Evolution* 12 (11): 2159–73. https://doi.org/10.1111/2041-210X.13687.

Pollock, Laura J., Reid Tingley, William K. Morris, et al. 2014. "Understanding Co-Occurrence by Modelling Species Simultaneously with a Joint Species Distribution Model (JSDM)." *Methods in Ecology and Evolution* 5 (5): 397–406. https://doi.org/10.1111/2041-210X.12180.

"PRISM Climate Group." 2014. February 4. https://prism.oregonstate.edu/.

Prosser, Jim I, J Ignacio Rangel-Castro, and Ken Killham. 2006. "Studying Plant–Microbe Interactions Using Stable Isotope Technologies." *Current Opinion in Biotechnology*, Analytical biotechnology, vol. 17 (1): 98–102. https://doi.org/10.1016/j.copbio.2006.01.001.

Revell, Liam J. 2012. "Phytools: An R Package for Phylogenetic Comparative Biology (and Other Things)." *Methods in Ecology and Evolution* 3 (2): 217–23. https://doi.org/10.1111/j.2041-210X.2011.00169.x.

Rizaludin, Muhammad Syamsu, Paolina Garbeva, Mark Zwart, and Jie Hu. 2023. "Microbial Volatiles Mediate Bacterial Evolutionary Dynamics." *The ISME Journal* 17 (12): 2144–46. https://doi.org/10.1038/s41396-023-01530-w.

Romdhane, Sana, Aymé Spor, Julie Aubert, et al. 2022. "Unraveling Negative Biotic Interactions Determining Soil Microbial Community Assembly and Functioning." *The ISME Journal* 16 (1): 296–306. https://doi.org/10.1038/s41396-021-01076-9.

Rossmann, Maike, Juan E Pérez-Jaramillo, Vanessa N Kavamura, et al. 2020. "Multitrophic Interactions in the Rhizosphere Microbiome of Wheat: From Bacteria and Fungi to Protists." *FEMS Microbiology Ecology* 96 (4): fiaa032. https://doi.org/10.1093/femsec/fiaa032.

Sanchez-Gorostiaga, Alicia, Djordje Bajić, Melisa L. Osborne, Juan F. Poyatos, and Alvaro Sanchez. 2019. "High-Order Interactions Distort the Functional Landscape of Microbial Consortia." *PLOS Biology* 17 (12): e3000550. https://doi.org/10.1371/journal.pbio.3000550.

Schäfer, Martin, Christine M. Vogel, Miriam Bortfeld-Miller, Maximilian Mittelviefhaus, and Julia A. Vorholt. 2022. "Mapping Phyllosphere Microbiota Interactions in Planta to Establish Genotype–Phenotype Relationships." *Nature Microbiology* 7 (6): 856–67. https://doi.org/10.1038/s41564-022-01132-w.

Schwartz, Egbert, Michaela Hayer, Bruce A Hungate, et al. 2016. "Stable Isotope Probing with 18O-Water to Investigate Microbial Growth and Death in Environmental Samples." *Current Opinion in Biotechnology*, Analytical biotechnology, vol. 41 (October): 14–18. https://doi.org/10.1016/j.copbio.2016.03.003.

Seybold, Heike, Tobias J. Demetrowitsch, M. Amine Hassani, et al. 2020. "A Fungal Pathogen Induces Systemic Susceptibility and Systemic Shifts in Wheat Metabolome and Microbiome Composition." *Nature Communications* 11 (1): 1910. https://doi.org/10.1038/s41467-020-15633-x.

Shannon, Paul, Andrew Markiel, Owen Ozier, et al. 2003. "Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks." *Genome Research* 13 (11): 2498–504. https://doi.org/10.1101/gr.1239303.

Sharma, Shubhangi, Stéphane Compant, Philipp Franken, Silke Ruppel, and Max-Bernhard Ballhausen. 2021. "It Takes Two to Tango: A Bacterial Biofilm Provides Protection against a Fungus-Feeding Bacterial Predator." *Microorganisms* 9 (8): 8. https://doi.org/10.3390/microorganisms9081566.

Sokol, Noah W., Eric Slessarev, Gianna L. Marschmann, et al. 2022. "Life and Death in the Soil Microbiome: How Ecological Processes Influence Biogeochemistry." *Nature Reviews Microbiology* 20 (7): 415–30. https://doi.org/10.1038/s41579-022-00695-z.

Taylor, D. Lee, William A. Walters, Niall J. Lennon, et al. 2016. "Accurate Estimation of Fungal Diversity and Abundance through Improved Lineage-Specific Primers Optimized for Illumina Amplicon Sequencing." *Applied and Environmental Microbiology* 82 (24): 7217–26. https://doi.org/10.1128/AEM.02576-16.

Trivedi, Chanda, Manuel Delgado-Baquerizo, Kelly Hamonts, Kaitao Lai, Peter B. Reich, and Brajesh K. Singh. 2019. "Losses in Microbial Functional Diversity Reduce the Rate of Key Soil Processes." *Soil Biology and Biochemistry* 135 (August): 267–74. https://doi.org/10.1016/j.soilbio.2019.05.008.

Vieira, Selma, Johannes Sikorski, Sophie Dietz, et al. 2020. "Drivers of the Composition of Active Rhizosphere Bacterial Communities in Temperate Grasslands." *The ISME Journal* 14 (2): 463–75. https://doi.org/10.1038/s41396-019-0543-4.

Vorholt, Julia A. 2012. "Microbial Life in the Phyllosphere." *Nature Reviews Microbiology* 10 (12): 828–40. https://doi.org/10.1038/nrmicro2910.

Wang, Chaoqun, and Yakov Kuzyakov. 2024. "Mechanisms and Implications of Bacterial–Fungal Competition for Soil Resources." *The ISME Journal* 18 (1): wrae073. https://doi.org/10.1093/ismejo/wrae073.

Wang, Xing, Zuoqiang Yuan, Arshad Ali, et al. 2023. "Leaf Traits and Temperature Shape the Elevational Patterns of Phyllosphere Microbiome." *Journal of Biogeography* 50 (12): 2135–47. https://doi.org/10.1111/jbi.14719.

Waring, Bonnie, and Christine V Hawkes. 2018. "Ecological Mechanisms Underlying Soil Bacterial Responses to Rainfall along a Steep Natural Precipitation Gradient." *FEMS Microbiology Ecology* 94 (2): fiy001. https://doi.org/10.1093/femsec/fiy001.

Warmink, J. A., R. Nazir, B. Corten, and J. D. van Elsas. 2011. "Hitchhikers on the Fungal Highway: The Helper Effect for Bacterial Migration via Fungal Hyphae." *Soil Biology and Biochemistry*, Knowledge gaps in soil C and N interactions, vol. 43 (4): 760–65. https://doi.org/10.1016/j.soilbio.2010.12.009.

Weiland-Bräuer, Nancy. 2021. "Friends or Foes—Microbial Interactions in Nature." *Biology* 10 (6): 496. https://doi.org/10.3390/biology10060496.

Werner, Earl E., and Scott D. Peacor. 2003. "A Review of Trait-Mediated Indirect Interactions in Ecological Communities." *Ecology* 84 (5): 1083–100. https://doi.org/10.1890/0012-9658(2003)084%255B1083:AROTII%255D2.0.CO;2.

Wootton, J. Timothy. 1994. "The Nature and Consequences of Indirect Effects in Ecological Communities." *Annual Review of Ecology and Systematics* 25: 443–66.

Wu, Chunxiao, Benshuai Yan, Furong Wei, et al. 2023. "Long-Term Application of Nitrogen and Phosphorus Fertilizers Changes the Process of Community Construction by Affecting Keystone Species of Crop Rhizosphere Microorganisms." *Science of The Total Environment* 897 (November): 165239. https://doi.org/10.1016/j.scitotenv.2023.165239.

Yang, Fan, Qiong Chen, Qian Zhang, Chunyan Long, Wei Jia, and Xiaoli Cheng. 2021. "Keystone Species Affect the Relationship between Soil Microbial Diversity and Ecosystem Function under Land Use Change in Subtropical China." *Functional Ecology* 35 (5): 1159–70. https://doi.org/10.1111/1365-2435.13769.

Yuan, Mengting Maggie, Anne Kakouridis, Evan Starr, et al. 2021. "Fungal-Bacterial Cooccurrence Patterns Differ between Arbuscular Mycorrhizal Fungi and Nonmycorrhizal Fungi across Soil Niches." *mBio* 12 (2): 10.1128/mbio.03509-20. https://doi.org/10.1128/mbio.03509-20.

Figure 1. Ecological association networks for (A) leaves and (D) roots used to define cross-kingdom interactions. Direct interactors are red nodes, keystone hub taxa are blue nodes, and taxa with only within-kingdom interactions are white nodes. Edge line type indicates the direction of association (negative are dashed and positive are solid). Bar plots of the relative abundance for each interaction type in (B) leaves and (D) roots, with the total number of taxa at the top of each bar and color representing taxonomic order.

Figure 2. Variance partitioning of microbial community composition when including only environmental and spatial (E+S) factors, or additionally including direct or diffuse biotic interactions. Rows are the bacterial and fungal communities in either the (A, B) leaf or (B, D) root, with each column indicating the subset of taxa used to represent the cross-kingdom biotic interactions. Residuals are reported adjacent to each Venn diagram. Only values > 3% are displayed.

Figure 3. Selected results of TITAN2 gradient analysis of ASVs identified as having direct or diffuse cross-kingdom interactions in (A-D) leaf and (E-H) root communities. Only ASVs that significantly changed in abundance across key environmental factors are shown (for others see Supplemental Fig. S6). Triangle symbols indicate the median change point, with symbol size relative to the median z-score, symbol direction indicating increases (up triangle) or decreases (down triangle) along the gradient, and color representing the taxonomic order as in Fig. 1. The horizontal lines around each symbol show the 5-95% quartile of bootstrapped change point distribution.