Spatial Regulation of Lck Activation at the Immune Synapse Revealed by a FRET-Based Biosensor

Clara Meana[1,3], Gonzalo San-José[1], María Beatriz Durán-Alonso[1,2], Javier Casas[1,2]

[1]Unidad de Excelencia Instituto de Biomedicina y Genética Molecular (IBGM), Consejo Superior de Investigaciones Científicas (CSIC) - Universidad de Valladolid (UVa), Spain

[2]Departamento de Bioquímica y Biología Molecular y Fisiología, Facultad de Medicina, Universidad de Valladolid, 47003 Valladolid, Spain

[3]Departamento de Pediatría e Inmunología, Universidad de Valladolid, Valladolid 47005, Spain

* Correspondence:
Corresponding Author
javier.casas@uva.es

Keywords: TCR, LCK, Immune Synapse, FRET, Spatial Regulation.

Abstract

T cell receptor (TCR) signaling is critically dependent on the Src-family kinase Lck, whose activation is tightly regulated both spatially and conformationally during antigen recognition. Here, we employ a third-generation FRET-based biosensor (TqLck-V2.3) to visualize Lck conformational dynamics in live T cells with high spatial resolution. Upon TCR engagement, we observe a paradoxical increase in whole-cell FRET signal, which immunolabeling reveals to be due to selective internalization of inactive Lck (pY505), while active Lck (pY394) remains membrane-associated and enriched at the immune synapse (IS). Using CD8α mutants that disrupt Lck binding, we demonstrate that free Lck undergoes more pronounced conformational activation than coreceptor-bound Lck. Furthermore, we show that C-terminal Src kinase (Csk) preferentially phosphorylates free Lck at Y505, while CD45 suppresses its activation via dephosphorylation of Y394, suggesting a dual regulatory mechanism that maintains free Lck in an inactive state under resting conditions. High-resolution imaging confirms sustained activation of Lck at the IS and transient inactivation at the periphery, revealing a spatially confined signaling architecture. These findings uncover a novel regulatory mechanism involving selective internalization and spatial segregation of Lck conformations and establish TqLck-V2.3 as a powerful tool for dissecting TCR signaling dynamics.

Introduction

T cell activation is a highly regulated process initiated by the engagement of the T cell receptor (TCR) with peptide–major histocompatibility complex (pMHC) molecules on antigen-presenting cells (APCs). This interaction triggers a cascade of intracellular signaling events, many of which are orchestrated by the Src-family kinase Lck. Lck plays a pivotal role in phosphorylating immunoreceptor tyrosine-based activation motifs (ITAMs) within the CD3 complex, thereby initiating downstream signaling pathways essential for T cell function and fate decisions.

Despite its central role, the mechanisms governing Lck activation and spatial regulation during TCR engagement remain incompletely understood. Lck exists in multiple conformational states, regulated by phosphorylation at two key tyrosine residues: Y394 (activating) and Y505 (inhibitory). The dynamic balance between these states is influenced by interactions with coreceptors (CD4/CD8), phosphatases, and kinases, as well as by the spatial organization of signaling complexes at the immune synapse.

Our previous work has proposed that free Lck, not bound to coreceptors, may be primarily responsible for initiating TCR signaling, while coreceptor-bound Lck contributes to signal amplification or stabilization. However, direct evidence supporting this model has been limited by the lack of tools capable of resolving Lck conformational changes in live cells with high spatial and temporal resolution. Current research indicates that in unstimulated T cells, the proportion of active Lck varies significantly, with studies reporting ranges from 2% to approximately 50% of the total Lck pool(1–3). This variation has inspired considerable debate in the field regarding the precise regulation and activation mechanisms of Lck(4).

Traditional approaches to studying Lck activation have relied heavily on biochemical methods such as Western blot and immunoprecipitation, which, while informative, cannot capture the spatial and temporal dynamics of Lck activation in living cells(2,3,5). Furthermore, technical challenges in sample preparation and antibody specificity have contributed to conflicting results in the literature(1,6). These limitations have highlighted the need for more sophisticated approaches to studying Lck regulation in situ.

Recent advances in fluorescence-based imaging techniques, particularly the development of FRET-based biosensors, have opened new possibilities for investigating the spatial and temporal aspects of Lck activation(7–9). These approaches allow for real-time monitoring of Lck conformational changes and activation states within the complex environment of the immune synapse, providing unprecedented insight into the spatial regulation of TCR signaling machinery.

We used a third-generation FRET-based biosensor, TqLck-V2.3, optimized for monitoring Lck conformational dynamics in T cells. Ratiometric FRET enables real-time monitoring of dynamic molecular interactions with high temporal resolution, which is critical for capturing the fast kinetics of early T cell signaling events. Unlike Fluorescence Lifetime Imaging Microscopy (FLIM), which requires longer acquisition times and complex data fitting to resolve fluorescence lifetimes, ratiometric FRET can be implemented with standard widefield or confocal microscopy, allowing for faster image

acquisition and analysis. This speed is essential for studying transient signaling intermediates and spatially localized activation patterns at the immunological synapse. TqLck-V2.3 enables the visualization of Lck conformational states in live T cells and provides insights into the spatial compartmentalization of active and inactive Lck during immune synapse formation. In this study, we combine live-cell imaging, phospho-specific labeling, and mutant T cell models to dissect the conformational regulation of Lck during antigen recognition. Our findings reveal a novel mechanism of Lck regulation involving the selective internalization of its inactive form and demonstrate that free Lck undergoes more substantial conformational changes than coreceptor-bound Lck upon TCR engagement

Material & Methods

2.1 Constructs and cells.

CHO cells expressing the tetracycline repressor (Thermo TRex) under blasticidin selection (10 μg/ml) were grown in Ham's F12 media with 10% (vol/vol) FCS, 100 U/ml of penicillin, 10 mg/ml of streptomycin, hygromycin 0.3 mg/ml and G418 0.8 mg/ml. Hybridomas expressing the OT-I TCR with or without CD8α and CD8β were made by retroviral transfection of TCR-deficient 58α-β cells (10). The plasmid pEF-TqLckV2.3 was provided by Dr. Burkhart Schraven from Magdeburg University. TqLckV2.3 was amplified by PCR adding 5′Hind-III and Not-I 3′specific restriction sites and inserted into retroviral vector pBMN-Z (S. Kinoshita and G. Nolan, www.stanford.edu/group/nolan). All the mutants were made using the Quickchange Site Mutagenesis Kit (Stratagene). Supernatants of these cells were used to transduce OT-I hybridomas. OT-I hybridoma cells were maintained in IMDM medium containing 10% fetal bovine serum, 2 mM L-glutamine, 100 U/ml penicillin/streptomycin, 50 μM β-mercaptoethanol,500 μg/ml G418 (selecting for TCRα) and 3 μg/ml puromycin (selecting for TCRβ).

pHR-Csk-P2A-Cbp and pHCM-mCherry-CD45cytCD43ex were kindly provided by Dr. John James from Oxford University.

2.2 Flow cytometry.

All flow cytometry was conducted on Cytek Aurora cytometer at the Institute of Biomedicine and Molecular Genetics flow cytometry facility. Flow cytometry data were analyzed in FlowJo (Tree Star) and exported where necessary.

2.3 Lentiviral and retroviral Transduction

VSV-G–pseudotyped lentiviral particles were produced by co-transfection of 293FT cells with transfer constructs (CD45cyt-CD43ex and Csk-P2A-Cbp) and the compatible packaging plasmids pMD2.G and psPAX2 in the presence of PEI 25K. Ecotropic retroviral particles were produced by co-transfection of Plat-E cells with transfer constructs (pBMN-TqLckV2.3, pBMN-CD8α-CxCP or pBMN-CD8β) and the compatible packaging plasmid pCL-Eco in the presence of PEI 25K. Viruses were harvested at 48 and 72 hours after transfection. Viral transduction of OT-I T cell hybridomas were carried out using concentrated viral particles in the presence of 8 μg/ml Polybrene (Sigma-

113 Aldrich), and infected cells were selected by FACS sorting or antibiotic resistance and subjected to
114 imaging experiments.

2.4 Live Cell FRET Imaging

A dual superfast external excitation/emission wheels specifically designed for FRET were used for
imaging, allowing almost simultaneous acquisition of donor and acceptor emission during donor
excitation (20 ms delay). This consisted in a Leica K5 sCMOS camera attached to a Leica Dmi8
inverted microscope equipped with aforementioned ultrafast external filter wheels and a CoolLED 3
illumination system. Cells expressing Lck activity FRET-based sensor (TqLckV2.3) were cultured in
glass-bottomed culture dishes (Mat-Tek). When improved adhesion was required, the glass was coated
with poly-L-lysine (Sigma). The culture medium was replaced with HBSS supplemented with 10 mM
HEPES, pH 7.4, 1.3 mM $CaCl_2$, and 1.3 mM $MgCl_2$. Cells were visualized at 37°C using a Leica Dmi8
system with an HC PL FLUOTAR 63X/1.3 NA oil immersion lens and an infrared focus maintenance
system. Proper localization of the sensor was confirmed by obtaining YFP images. For ratiometric
imaging, Turquoise (Tq) fluorescence was obtained using an excitation filter at 430/24 nm and an
emission filter at 470/24 nm, Venus using an excitation filter at 500/20 nm and an emission filter at
535/30 nm, and FRET, without correction, using the Tq excitation filter at 430/24 nm and the Venus
emission filter. The acquisition of the three channels was virtually simultaneous. We recorded a 2 min
baseline before adding T cells acquiring images each 5 s. ImageJ software with custom plugins was
used to correct both the movement and the background of the raw images and to create pixel-by-pixel
FRET/Tq ratio images.

2.5 Immunocytochemistry Analyses

For phospho-specific staining of Lck, cells were fixed with 4% paraformaldehyde for 20 min in
suspension and then permeabilized with 0.1% Triton X-100 for 20 min at room temperature. Cells were
treated with 5% goat serum for 30 min and incubated with antibodies against phospho Y505, phospho
Y394, total Lck (Cell Signaling) or phospho-Tyr (4G10 clone, Merck) for 1h. After washing, the cells
were incubated with spectrally appropriate Alexa Fluor-conjugated Fab fragments against primary
species antibodies (Thermo) for 1h. After intracellular staining, nuclei were counterstained with the
DNA binding dye Hoechst 33342 (Thermo). All images were captured with a Leica confocal system
TCS SP5X inverted microscope with a HCS Plan Apo CS 63X/1.4 NA oil immersion lens. Leica
Application Suite Advanced Fluorescence software was used for the capture, and ImageJ was used for
deconvolution (Parallel Iterative 3D plugin), image presentation and quantification.

Results

3.1 Functionality of a third-generation FRET-based Lck biosensor

To monitor Lck conformational dynamics in live cells, we employed an optimizedand based on second-
generation FRET-based biosensor, TqLck-V2.3. In this construct, the original CFP and YFP
fluorophores were replaced with monomeric Turquoise2 (Tq) and monomeric Venus (V),
respectively(7) . Additionally, the entire insert was codon-optimized to minimize recombination events
during retroviral cloning. Importantly, the donor fluorophore (Tq) was repositioned between the SH3

and SH2 domains of Lck, based on prior findings by Paster et al. demonstrating that this configuration
(as in CLck-Y2) more accurately recapitulates TCR-mediated signaling compared to other biosensor
designs (Fig. 1A)(11).

We first confirmed the biosensor's localization by confocal microscopy, validating its predominant
presence at the plasma membrane and exclusion from the cytoplasm (Fig. 1B). Upon treatment with
the Lck inhibitor PP2, we observed an increase in the FRET/Tq ratio, consistent with a closed (inactive)
Lck conformation. Conversely, stimulation with pervanadate, a phosphatase inhibitor that mimics TCR
activation, led to a decrease in the FRET/Tq ratio, indicative of an open (active) conformation (Fig.
1C). Notably, unstimulated cells exhibited a broad distribution of FRET/Tq ratios, likely reflecting the
presence of distinct Lck pools in different conformational states.

3.2 Quantitative Analysis of Lck Regulation via TCR–pMHC Conjugate Formation in antigen
presenting surfaces

To investigate Lck regulation in the context of antigen presentation, we developed an improved assay
system tailored to measure T-cell receptor (TCR) engagement with peptide–MHC (pMHC) complexes.
The OT-I TCR recognizes the OVA peptide (SIINFEKL) with high affinity when presented by H-2Kb,
but does not respond to the VSV peptide (RGYVYQGL) in the same context. Specifically, we analyzed
conjugate formation between OT-I cells expressing CD8α and CD8β (hereafter referred to as OT-
I.WT) and a surrogate antigen-presenting cell line. We used CHO cells engineered to express
doxycycline (DOX) inducible single-chain H-2Kb molecules constitutively loaded with either the
agonist peptide OVA (scH-2Kb–OVA) or the non-stimulatory peptide VSV (scH-2Kb–VSV), as
previously described (12,13) (Fig. 2A, B). To generate stable and functional single-chain pMHCI (sc-
pMHCI) molecules, we employed recombinant constructs in which the MHC-binding peptide was
fused to the N-terminus of β2-microglobulin, followed by linkage to the N-terminus of the MHCI
heavy chain, as previously described(14–16). When expressed on the cell surface, these constructs
exhibited high stability and effectively triggered T cell activation.

Quantification of conjugate formation revealed a time-dependent increase in the percentage of OT-
I.WT cells forming conjugates with CHO cells upon induction of pMHC expression. This increase was
significantly greater when CHO cells presented the antigenic OVA peptide compared to the non-
antigenic VSV peptide. In the absence of doxycycline (repressed state), there was some "leaky"
expression of sc-KbOVA, which was sufficient to trigger partial CD69 upregulation in CD8 OT-I T
cells, however, this level of expression did not lead to significant TCR downregulation (12,13) . Flow
cytometry using a KbOVA-specific antibody revealed that sc-KbOVA expression under these
conditions was nearly undetectable.

To assess early T-cell activation, we measured global tyrosine phosphorylation levels after 15 minutes
of co-culture using phospho-tyrosine staining (Fig. 2C). OT-I.WT cells exposed to CHO cells
presenting Kb:OVA exhibited elevated phospho-tyrosine levels relative to those exposed to Kb:VSV.
Pre-treatment of OT-I.WT cells with either a TCR-blocking antibody or the Src-family kinase inhibitor
PP2 abrogated this phosphorylation response, reducing it to levels comparable to those observed with
the non-stimulatory VSV peptide (Fig. 2C). Together, these results demonstrate that OT-I.WT T cells
selectively form stable conjugates and initiate early activation signaling in response to antigenic pMHC
complexes, highlighting the specificity and functional sensitivity of TCR–pMHC interactions in this
system.

3.3 Validation of a third-generation FRET-based Lck biosensor for live-cell imaging

Having validated the TqLck-V2.3 biosensor for monitoring Lck conformational changes, we next
employed confocal microscopy to assess Lck dynamics in a more spatially resolved context. OT-I T
cells expressing the biosensor were seeded onto monolayers of CHO cells inducibly expressing pMHC
complexes. Z-projection imaging was used to capture the full three-dimensional interaction interface
between T cells and the CHO monolayer as shown in Fig. 2D.

Unexpectedly, when CHO cells presented the antigenic OVA peptide, we observed a significant
increase in the FRET/Tq ratio across the entire T cell (Fig. 2E), suggesting a shift toward a closed
(inactive) Lck conformation. To investigate this paradoxical finding, we performed immunolabeling
using phospho-specific antibodies targeting Lck phosphorylated at Y394 (active form) and Y505
(inactive form).

Immunofluorescence analysis revealed that following TCR engagement, active Lck (pY394) remained
localized at the plasma membrane, whereas the inactive form (pY505) was predominantly internalized
(Fig 3A). These results support the FRET-based observation of increased closed conformation when
analyzing the whole cell and suggest that internalization of inactive Lck contributes to the overall
FRET signal, while active Lck remains membrane-associated. The selective internalization of Y505-
phosphorylated (inactive) Lck following TCR engagement suggests a previously unrecognized
regulatory mechanism that may spatially restrict Lck activity by compartmentalizing its inactive pool
away from the site of TCR stimulation, normally the immune synapse. Altogether, these findings
establish TqLck-V2.3 as a powerful biosensor for dissecting Lck conformational dynamics in live cells
and uncover a novel mechanism of spatial regulation whereby inactive Lck is selectively internalized
following TCR engagement, potentially fine-tuning signaling at the immune synapse.

3.4 Lck biosensor reveals conformational activation at the immune synapse upon TCR
216
engagement and distinguishes coreceptor independent Lck pools
217

To investigate the spatial and conformational dynamics of Lck activation during T-cell receptor (TCR)
engagement we used fast FRET live cell microscopy. Stimulation with crosslinking biotinylated
αCD3/αCD8 antibodies induced a 5 minute decrease in the FRET/Tq ratio as shown in Fig. 4A. This
conformational change was abolished by PP2 pre-treatment, confirming that the observed FRET signal
corresponds to Lck activation. To assess the biosensor's performance in a more physiological context,
we co-cultured TqLck-V2.3-expressing OT-I T cells with CHO cells presenting either scH-2Kb–OVA
or scH-2Kb–VSV and monitored FRET changes at the immune synapse. A significant decrease in the
FRET/Tq ratio was observed only in response to OVA presentation, indicating Lck activation in the
immune synapse upon antigen recognition (Fig. 4B).

Previous results indicated that free Lck is initially responsible for TCR triggering (17). To further
dissect the contribution of coreceptor-bound versus free Lck, we expressed the TqLck-V2.3 biosensor
in OT-I T cells carrying a mutation in the intracellular CxCP motif of CD8α (C227K/C229P mutated
to S227K/S229P), which disrupts Lck binding (17). This is hereafter referred to as OT-I.CxCP.
Interestingly, these mutant cells exhibited a more pronounced FRET/Tq ratio decrease upon OVA
recognition compared to wild-type cells, suggesting that free Lck undergoes more substantial
conformational changes than coreceptor-bound Lck during TCR engagement (Fig. 4C). These results
align with previous findings that initial TCR triggering is mediated predominantly by free Lck rather
than coreceptor-bound Lck (17).To further investigate the spatial dynamics of free Lck during antigen
presentation, we manually quantified FRET efficiency of the Tq-LckV2.3 biosensor at the IS and
across the remaining plasma membrane (Fig. 4D). During the initial phase of T cell–APC contact (0–
60 s), we observed a marked decrease in FRET signal at the IS, indicative of sensor opening and most

likely a higher Lck activity. In contrast, an increase in FRET was detected in the rest of the membrane,
suggesting Lck inactivation in those regions. In the subsequent 60 seconds, the sensor at the rest of the
membrane exhibited a closure, consistent with increased Lck activity, which remained stable
throughout the rest of the interaction. Notably, the previously opened sensor in IS regions maintained
its conformation (Fig. 4D), indicating a sustained spatial segregation of Lck activity during the T cell–
APC engagement.

3.5 High-Resolution Imaging Reveals Spatial Restriction of Active Lck at the Immune Synapse

Since free Lck seems to be responsible for initial TCR triggering, to enhance spatial resolution and
better characterize Lck conformational dynamics at the immune synapse (IS), we performed
deconvolution confocal microscopy on fixed cells. Fig. 4E shows a representative conjugate between
an OT-I CxCP mutant T cell expressing the TqLck-V2.3 biosensor, where Lck can´t bind to coreceptor,
and a Cy5-labeled CHO antigen-presenting cell expressing Kb-OVA (Fig. 4E) or Kb-VSV (Fig. 4F).

FRET/Tq ratio imaging revealed a distinct decrease in the ratio at the IS upon recognition of Kb-OVA
(but not Kb-VSV), indicating the presence of Lck in an open (active) conformation upon cognate
antigen recognition. Notably, a subtle but consistent increase in the FRET/Tq ratio was observed at the
periphery of the IS. This is consistent with the elevated FRET/Tq levels previously detected in the non-
synaptic membrane regions by live-cell fluorescence microscopy (Fig. 4D). This spatial pattern
suggests that only the open conformation of Lck is selectively enriched within the synapse, while the
closed (inactive) form may be excluded or retained at the borders.

3.6 Csk preferentially phosphorylates Lck not bound to coreceptors

To establish a latent system where we can distinguish the role of spatial conformation of free Lck and
coreceptor bound Lck, we next sought to restrain the kinase activity of Lck by overexpressing C-
terminal Src kinase (Csk). Csk is a key negative regulator of Lck, phosphorylating its inhibitory
tyrosine residue (Y505) to promote a closed, inactive conformation. This phosphorylation event
counterbalances the activating phosphorylation at Y394, thereby modulating Lck catalytic activity.
Importantly, Csk lacks intrinsic membrane-targeting domains and relies on adaptor proteins such as
PAG/Cbp to localize to the plasma membrane, where it can access Lck. To investigate whether Csk
preferentially phosphorylates coreceptor-bound or free Lck, we expressed the FRET-based Lck activity
sensor Tq-LckV2.3, along with Csk and Cbp, in OT-I T cells co-expressing either wild-type CD8α or
a CxCP mutant that disrupts Lck–coreceptor interaction. In cells expressing the CxCP mutant, we
observed a marked increase in the FRET/Tq ratio of the Lck sensor when Csk and Cbp were co-
expressed, indicative of a closed Lck conformation and consistent with enhanced phosphorylation at
the inhibitory Y505 site (Fig. 5). These results suggest that Csk preferentially targets coreceptor-free
Lck for inhibitory phosphorylation.

3.7 CD45 Suppresses Free Lck Activation via Tyr394 Dephosphorylation to Prevent Spontaneous
TCR Signaling

CD45 is a tyrosine phosphatase that modulates T cell signaling in a complex manner by
dephosphorylating both the inhibitory Tyr505 and the activating Tyr394 of Lck, thereby exerting both
activating and suppressive control depending on the context and localization of Lck. To investigate the
regulatory role of CD45 on distinct pools of Lck in resting T cells, we examined the conformational
changes in free (non coreceptor-bound) Lck. We made use of the construct CD45cytCD43ex-mCherry
(18,19) where the cytoplasmic phosphatase domains of CD45 were fused to the large extracellular and
transmembrane domains of CD43. Although CD43 ectodomain shares no amino acid similarity with
CD45, it is as long and as heavily glycosylated as that of CD45 and should therefore mimic its basic
physicochemical properties. This construct has previously been shown to functionally replicate
CD45(18). We used the TqLckV-2.3 biosensor in the presence or absence of CD45cytCD43ex-
mCherry in resting cells, We observed a consistent FRET increase in the free Lck pool when
CD45cytCD43ex-mCherry was expressed in CD8α-CxCP mutant expressing cells (Fig. 5). This
suggests that CD45 preferentially targets Tyr394 in free Lck, in contrast to its well-characterized role
in dephosphorylating the inhibitory Tyr505 site as shown when Lck can be also bound to CD8
coreceptor (Fig. 5). These findings imply that CD45 may act as a negative regulator of free Lck by
maintaining it in an inactive state through dephosphorylation of the activating site. This observation
suggests that CD45 actively suppresses the activation of free Lck by targeting its activating
phosphorylation site. It further suggests that CD45 functions as a gatekeeper, maintaining free Lck in
an inactive state under resting conditions. This mechanism may serve to prevent spontaneous or ligand-
independent TCR triggering, ensuring that T cell activation is tightly regulated and spatially restricted
to appropriate signaling contexts.

299 4. Discussion

T cell activation is a tightly regulated process that relies on the precise spatial and temporal
coordination of signaling molecules at the immune synapse. Among these, the Src-family kinase Lck
plays a pivotal role in initiating TCR signaling by phosphorylating ITAMs within the CD3 complex.
While the importance of Lck is well established, the mechanisms that regulate its activation and
localization during antigen recognition remain incompletely understood. In this study, we employed a
third-generation FRET-based biosensor, TqLck-V2.3, to visualize Lck conformational dynamics in
live T cells and dissect the contributions of coreceptor-bound versus free Lck during TCR engagement.

TqLck-V2.3 biosensor reports on the conformational state of Lck but does not differentiate between
the primed (open, non-phosphorylated) and double-phosphorylated (open, phosphorylated at both
Tyr394 and Tyr505) forms, as both adopt a constitutively open conformation that is insensitive to TCR
stimulation(7,8) . Latest studies show that the proportion of active Lck in resting T cells might be as
low as 2% of the total Lck population (1) therefore, any conformational opening changes detected by
the biosensor upon TCR engagement are likely indicative of Lck activation events.

Our results demonstrate that Lck undergoes distinct conformational changes upon TCR stimulation,
with a rapid decrease in the FRET/Tq ratio indicating a transition to an open, active conformation. This
activation was abrogated by the Src-family kinase inhibitor PP2, confirming the specificity of the
biosensor for Lck activity. Importantly, we observed that antigen-specific stimulation with CHO cells
presenting OVA-loaded pMHC induced a robust FRET response, whereas non-stimulatory VSV-
loaded pMHC did not, validating the biosensor's sensitivity to physiologically relevant TCR signals.

Unexpectedly, whole-cell imaging revealed an increase in the FRET/Tq ratio upon antigen recognition,
suggesting a paradoxical shift toward a closed Lck conformation. Immunofluorescence analysis
resolved this discrepancy by showing that inactive Lck (pY505) is selectively internalized following

TCR engagement, while active Lck (pY394) remains at the plasma membrane. This spatial segregation implies a novel regulatory mechanism in which the internalization of inactive Lck may serve to sharpen the signaling interface by restricting kinase activity to the immune synapse.

Further, our use of a CD8α mutant that disrupts Lck binding revealed that free Lck undergoes more substantial conformational activation than coreceptor-bound Lck. This finding supports previous models suggesting that free Lck is primarily responsible for initiating TCR signaling, while coreceptor-bound Lck may play a more modulatory role(2,17). High-resolution imaging confirmed that active Lck is enriched within the immune synapse, whereas the inactive form is excluded to the periphery, reinforcing the idea that spatial compartmentalization of Lck activity is critical for signal fidelity.

In mice, the presence of LCK bound to CD8 is not essential for the antiviral and antitumor functions of cytotoxic T cells. However, it plays a kinase-dependent role in enhancing CD8+ T cell responses to weaker antigens. Conversely, LCK associated with CD4 is crucial for the proper development and activity of helper T cells, primarily through a kinase-independent mechanism that stabilizes CD4 on the cell surface(20). As free Lck is the primary initiator of TCR signaling(17), then the Csk preference for phosphorylating free Lck suggests a built-in regulatory checkpoint. This ensures that only the right spatial and temporal context allows TCR triggering, preventing spontaneous or inappropriate activation. The system may keep free Lck in a controlled but restrained state (most likely in a closed configuration), where it can rapidly respond to TCR engagement but is otherwise kept inactive by Csk. This would allow rapid signal initiation while maintaining immune tolerance. Coreceptor-bound Lck might serve more as a stabilized, less dynamic pool, possibly involved in signal amplification rather than initiation. This division of labor could explain why coreceptor-bound Lck is less accessible to Csk.

During antigen recognition at the IS, Lck remains in a sustained open conformation, supporting continuous TCR signaling. This localized activation ensures that signaling is tightly focused at the site of contact between the T cell and the antigen-presenting cell. In contrast, Lck outside the IS undergoes only transient inactivation, suggesting a dynamic but restrained regulation across the rest of the plasma membrane. This spatial compartmentalization of Lck activity likely serves to prevent unintended signaling while maintaining the cell's readiness to respond to new stimuli, highlighting a finely tuned mechanism of T cell activation.

Our data refine the current understanding of Lck regulation in resting T cells by demonstrating that Csk preferentially phosphorylates free, non-coreceptor-bound Lck, while CD45 dephosphorylates not only the inhibitory Y505 but also the activating Y394 residue. These findings challenge the notion of a static, constitutively active Lck pool and suggest a more dynamic regulatory landscape in which Lck activity is tightly controlled through spatial segregation and substrate specificity. In this context, CD45 emerges as a dual-function phosphatase capable of both priming and attenuating Lck activity depending on its localization and substrate accessibility. CD45 is a critical regulator that maintains T cells in a quiescent state and sets the threshold for TCR activation(21) . This nuanced model aligns with and extends the kinetic segregation framework, emphasizing the importance of membrane microdomain organization and the differential regulation of coreceptor-associated versus free Lck in shaping the threshold and timing of TCR signaling.

Together, these findings provide new insights into the spatial and conformational regulation of Lck during T cell activation. The selective internalization of inactive Lck and the preferential activation of free Lck at the immune synapse represent previously unrecognized layers of regulation that may fine-

365 tune TCR signaling. The TqLck-V2.3 biosensor offers a powerful tool for further dissecting these
366 dynamics in physiological and pathological contexts, including autoimmunity and immunotherapy.

Conflict of Interest

370 The author declare that the research was conducted in the absence of any commercial or financial
371 relationships that could be construed as a potential conflict of interest.

Author Contributions

373 Conceptualization, J.C.; methodology, J.C. and B.D.;.; validation, J.C., B.D.; formal analysis, J.C. and
374 G.S.J.; investigation, J.C., M.B. and G.S.J; resources, J.C.; data curation, J.C., B.D. and G.S.J.;
375 writing—original draft preparation, J.C.; writing—review and editing, J.C., M.B., and G.S.J.;
376 visualization, J.C.; supervision, J.C..; project administration J.C.; funding acquisition, J.C. All authors
377 have read and agreed to the published version of the manuscript.

Funding

380 This research was funded by several intramural funds from the Institute of Biomedicine and Molecular
381 Genetics (IBGM) through the Regional Government of Castile and Leon.

Acknowledgments

I would like to specifically thank Nicholas Gascoigne for all the materials and advice provided through
the years.

Bibliography

1. Ondrej B, Valecka J, Manning J, Filipp D. The pool of preactivated Lck in the initiation of T-cell
signaling: a critical re-evaluation of the Lck standby model. (2014)1–12. doi: 10.1038/icb.2014.100

2. Wei Q, Brzostek J, Sankaran S, Casas J, Hew LS-Q, Yap J, Zhao X, Wojciech L, Gascoigne NRJ.
Lck bound to coreceptor is less active than free Lck. *P Natl Acad Sci Usa* (2020)201913334. doi:
10.1073/pnas.1913334117

3. Nika K, Nika K, Soldani C, Soldani C, Salek M, Salek M, Paster W, Paster W, Gray A, Gray A, et
al. Constitutively Active Lck Kinase in T Cells Drives Antigen Receptor Signal Transduction.
Immunity (2010) 32:766–777. doi: 10.1016/j.immuni.2010.05.011

4. Simeoni L. Lck activation: puzzling the pieces together. *Oncotarget* (2017) 8:102761–102762. doi:
10.18632/oncotarget.22309

5. Moogk D, Zhong S, Yu Z, Liadi I, Rittase W, Fang V, Dougherty J, Perez-Garcia A, Osman I, Zhu
C, et al. Constitutive Lck Activity Drives Sensitivity Differences between CD8+ Memory T Cell
Subsets. *The Journal of Immunology* (2016)1600178. doi: 10.4049/jimmunol.1600178

6. Ballek O, Broučková A, Manning J, Filipp D. A specific type of membrane microdomains is
involved in the maintenance and translocation of kinase active Lck to lipid rafts. *Immunol Lett* (2012)
142:64–74. doi: 10.1016/j.imlet.2012.01.001

7. Philipsen L, Reddycherla AV, Hartig R, Gumz J, Kästle M, Kritikos A, Poltorak MP, Prokazov Y,
Turbin E, Weber A, et al. De novo phosphorylation and conformational opening of the tyrosine kinase
Lck act in concert to initiate T cell receptor signaling. *Science signaling* (2017) 10:eaaf4736. doi:
10.1126/scisignal.aaf4736

8. Stirnweiss A, Hartig R, Gieseler S, Lindquist JA, Reichardt P, et al. T Cell Activation Results in
Conformational Changes in the Src Family Kinase Lck to Induce Its Activation. *Science signaling*
(2013) 6:ra13–ra13. doi: 10.1126/scisignal.2003607

9. Kästle M, Merten C, Hartig R, Kaehne T, Liaunardy-Jopeace A, Woessner NM, Schamel WW,
James J, Minguet S, Simeoni L, et al. Tyrosine 192 within the SH2 domain of the Src-protein tyrosine
kinase p56Lck regulates T-cell activation independently of Lck/CD45 interactions. *Cell Commun*
Signal (2020) 18:183. doi: 10.1186/s12964-020-00673-z

10. Letourneur F, Malissen B. Derivation of a T cell hybridoma variant deprived of functional T cell
receptor α and β chain transcripts reveals a nonfunctional α-mRNA of BW5147 origin. *Eur J Immunol*
(1989) 19:2269–2274. doi: 10.1002/eji.1830191214

11. Paster W, Paar C, Eckerstorfer P, Jakober A, Drbal K, Schütz GJ, Sonnleitner A, Stockinger H.
Genetically encoded Förster resonance energy transfer sensors for the conformation of the Src family
kinase Lck. (2009) 182:2160–2167. doi: 10.4049/jimmunol.0802639

12. Hoerter JAH, Brzostek J, Artyomov MN, Abel SM, Casas J, Rybakin V, Ampudia J, Lotz C,
Connolly JM, Chakraborty AK, et al. Coreceptor affinity for MHC defines peptide specificity
requirements for TCR interaction with coagonist peptide-MHC. *J Exp Med* (2013) 210:1807–1821.
doi: 10.1084/jem.20122528

13. Zhao X, Wu L-Z, Ng EKY, Leow KWS, Wei Q, Gascoigne NRJ, Brzostek J. Non-Stimulatory
pMHC Enhance CD8 T Cell Effector Functions by Recruiting Coreceptor-Bound Lck. *Front Immunol*
(2021) 12:721722. doi: 10.3389/fimmu.2021.721722

14. Yu YYL, Netuschil N, Lybarger L, Connolly JM, Hansen TH. Cutting Edge: Single-Chain Trimers
of MHC Class I Molecules Form Stable Structures That Potently Stimulate Antigen-Specific T Cells
and B Cells. *J Immunol* (2002) 168:3145–3149. doi: 10.4049/jimmunol.168.7.3145

15. Choudhuri K, Wiseman D, Brown MH, Gould K, Merwe PA van der. T-cell receptor triggering is
critically dependent on the dimensions of its peptide-MHC ligand. *Nature* (2005) 436:578–582. doi:
10.1038/nature03843

16. Palmowski MJ, Parker M, Choudhuri K, Chiu C, Callan MFC, Merwe PA van der, Cerundolo V,
Gould KG. A Single-Chain H-2Db Molecule Presenting an Influenza Virus Nucleoprotein Epitope
Shows Enhanced Ability at Stimulating CD8+ T Cell Responses In Vivo. *J Immunol* (2009) 182:4565–
4571. doi: 10.4049/jimmunol.0803893

17. Casas J, Brzostek J, Zarnitsyna VI, Hong J, Wei Q, Hoerter JAH, Fu G, Ampudia J, Zamoyska R,
Zhu C, et al. Ligand-engaged TCR is triggered by Lck not associated with CD8 coreceptor. *Nat
Commun* (2014) 5:5624. doi: 10.1038/ncomms6624

18. Irles C, Symons A, Michel F, Bakker TR, Merwe PA van der, Acuto O. CD45 ectodomain controls
interaction with GEMs and Lck activity for optimal TCR signaling. *Nat Immunol* (2003) 4:189–197.
doi: 10.1038/ni877

19. James JR, Vale RD. Biophysical mechanism of T-cell receptor triggering in a reconstituted system.
Nature (2012) 487:64–69. doi: 10.1038/nature11220

20. Horkova V, Drobek A, Paprckova D, Niederlova V, Prasai A, Uleri V, Glatzova D, Kraller M,
Cesnekova M, Janusova S, et al. Unique roles of co-receptor-bound LCK in helper and cytotoxic T
cells. *Nat Immunol* (2023) 24:174–185. doi: 10.1038/s41590-022-01366-0

21. Hui E, Vale RD. In vitro membrane reconstitution of the T-cell receptor proximal signaling
network. *Nature Structural & Molecular Biology* (2014) 21:133–142. doi: 10.1038/nsmb.2762

471   **Figures**

Figure 1. Schematic representation of the conformational changes that result in FRET changes in the TqLckV2.3 biosensor. A) The model of the open conformation (left pane) and closed conformation (right pane), modified from Prakaash et al. Sci Rep (2022). B) OT-I T cell hybridoma expressing TqLckV2.3 biosensor were analyzed by confocal microscopy. Scale bar, 5 μm. C) Quantitation of FRET/Tq ratio from TqLckV2.3 biosensor in OT-I T cell hybridoma treated with vehicle (blue, n= 62) 10 μM PP2 Lck inhibitor (purple, n=60) or 2 min stimulation with 1 mM Pervanadate (green, n=65). ***, p < 0.001 by Student's t test.

Figure 2. Expression of inducible sc-MHCI agonist molecules KbOVA inCHO cells and activation of corresponding
OT-I CD8 T cells. (A) Trex CHO cells expressing DOX-inducible sc-MHC were stained with anti-Kb:OVA and analyzed
by flow cytometry. Green shading: Repressed (no DOX) Trex CHO cells. Purple: Induced (+ DOX) sc-KbOVA. (B) OT-I
hybridomas were stimulated for the indicated times with Trex CHO cells expressing DOX-inducible sc-MHC and
percentage of conjugates were represented. (C) OT-I hybridomas were stimulated as in (B) for 15 min and stained with
anti-phospho-Tyr. Images were taken using confocal microscopy. Mean fluorescence of Control n = 23, KbOVA n = 20,
KbOVA + αTCR n= 18, KbOVA +PP2 n= 18 and KbVSV n = 17 ± s.e.m. is shown. (D) OT-I T cell hybridoma expressing
TqLckV2.3 biosensor cells were plated onto a Cy5 labeled monolayer of Trex CHO cells expressing DOX-inducible sc-
MHC and analyzed by confocal microscopy. A representative Z-max projection image is presented, along with two distinct
3D reconstructions illustrating T cell–APC interactions. E) Individual FRET/Tq ratio (representative images shown) of OT-
I T cell hybridoma expressing TqLckV2.3 cells. Control n = 98, KbOVA n = 65 and KbVSV n = 43.). #, p < 0.1; ***, p <
0.001 by Student's t test. Scale bar 5 μm.

Figure 3. Spatial segregation of different phospho-Lck species. A) OT-I T cells, unstimulated and 15 min after
stimulation with αCD3/CD8 crosslinking and analyzed by epifluorescecnce deconvolution microscopy B) Differential
phospho-Lck antibody staining (green Y394P and red Y505P) distribution analysis from A. Bars represent % of
cytoplasmic phospho-Lck ± s.e.m. >40 cells/sample. ***, $p < 0.001$ by Student's t test. C) OT-I T cells 15 min after
stimulation with αCD3/CD8 crosslinking and analyzed by confocal deconvolution microscopy. Nuclei were counterstained
with Hoechst 33342. Scale bar 5 μm.

Figure 4. Lck dynamics at the Immune Synapse A) OT-I T cells expressing TqLckV2.3, untreated or pretreated with 10
mM PP2 were stimulated with αCD3/CD8 crosslinking and analyzed by live cell FRET microscopyMean FRET/Tq ratio ±
s.e.m was represented. Multiple t test was applied. ***, p < 0.001. B) OT-I hybridomas expressing TqLckV2.3 were
stimulated with Trex CHO cells expressing DOX-inducible KbOVA (purple circles) or KbVSV (blue triangles) and
FRET/Tq ratio was represented since the initial contact between T cell and APC. C) Same as in B but with OT-I expressing
CD8α-CxCP mutant (green squares). D) Same as in C (green circles) and rest of the plasma membrane FRET/Tq ratio
analysis (red circles). The results are representative of at least three independent experiments. E) Representative image of
OT-I. TqLckV2.3 expressing cells forming the immune synapse (Tq fluorescence in cyan, mVenus in yellow, CHO APC
membrane labelled with Cy5 in purple and transmitted light image is shown as well) Bottom-right panel shows FRET/Tq
ratio of the T cell. F) Same as in E but with unlabeled CHO. Bottom right panel shows magnification of the selected region.
The results are representative of at least three independent experiments.

530 **Figure 5. Lck regulation in resting cells.** FRET/Tq ratio from TqLckV2.3 biosensor in unstimulated OT-I.WT (purple)
531 or OT-I.CxCP (cyan) cell hybridomas expressing Csk/Cbp or CD45cytCD43ex. WT n=233, CxCP n=139; Csk-Cbp
532 expressing WT n=262 CxCP Csk-Cbp CxCP n=149; CD45 expressing WT n=119 CD45 expressing CxCP n =52.