Carbapenem-resistance *oprD* mutations reshape *Pseudomonas*

***aeruginosa* host-pathogen interactions during infection**

Pablo Laborda[1,*], Claudia Antonella Colque[1], Ruggero La Rosa[1,2], Søren Molin[3], Helle

Krogh Johansen[1,2,4].

[1] Department of Clinical Microbiology 9301, Rigshospitalet, 2100 Copenhagen, Denmark

[2] Department of Health Technology, Technical University of Denmark, 2800 Kgs. Lyngby,

Denmark

[3] The Novo Nordisk Foundation Center for Biosustainability, Technical University of

Denmark, 2800 Kgs. Lyngby, Denmark

[4] Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of

Copenhagen, 2200 Copenhagen, Denmark

*Corresponding author: Pablo Laborda (palama@biosustain.dtu.dk)

Running title: **OprD mutations remodel *P. aeruginosa* infection**

Abstract

Antibiotic resistance is a major global health threat. While its role in reducing drug
susceptibility is well established, the broader consequences of resistance mutations on
bacterial physiology and behavior during infection remain poorly understood.
Carbapenem-resistant *Pseudomonas aeruginosa* is considered among the highest-
priority bacterial threats, with resistance commonly driven by loss-of-function mutations
in the carbapenem uptake porin OprD. Here we show that such mutations can arise in
clinical isolates even without prior carbapenem treatment, indicating that their effects
during infection extend beyond antibiotic resistance. Consistent with this, we found that
oprD mutants exhibit enhanced early attachment to and translocation across airway
epithelial barriers in infection models, an effect maintained across strains with distinct
clinical genomic backgrounds and infection dynamics. Our findings indicate that loss of
OprD alters the bacterial outer membrane charge and reduces mucus entrapment,
thereby facilitating epithelial barrier colonization. Overall, these results illustrate how
antibiotic resistance mutations can directly shape infection dynamics, extending their
impact well beyond antimicrobial susceptibility.

Introduction

Antibiotic resistance is one of the most important medical challenges of the 21st century, undermining the effectiveness of current treatments and driving higher morbidity and mortality associated with infections worldwide [1]. Gram-negative pathogens are of particular concern due to their intrinsic resistance mechanisms, low outer membrane permeability to antimicrobial agents, and ability to acquire additional resistance traits [2]. Among them, *Pseudomonas aeruginosa* stands out as a highly adaptable opportunistic pathogen capable of causing a broad spectrum of infections, especially in immunocompromised patients and individuals with chronic lung diseases such as cystic fibrosis (CF), Primary Ciliary Dyskinesia (PCD) and Chronic Obstructive Pulmonary Disease (COPD) [3–5].

In chronic infections, *P. aeruginosa* often acquires mutations that reduce the efficacy of antimicrobial agents [6,7], a phenomenon typically attributed to the selective pressure of antibiotic exposure. Resistance to carbapenems, a last resort antibiotics, is of particular concern because it severely restricts therapeutic options [8,9], placing carbapenem-resistant *P. aeruginosa* among the highest-priority bacterial threats [10]. A central determinant of carbapenem susceptibility is the outer membrane porin OprD, which mediates drug uptake of carbapenem antibiotics. Loss or alteration of OprD is among the most common mechanisms of carbapenem resistance in *P. aeruginosa* [4,11,12], and it is frequently observed in clinical isolates [6,7].

While the genetic basis of antibiotic resistance is generally well-characterized, its broader physiological consequences remain largely undiscovered. Resistance mutations can exert pleiotropic effects on bacterial envelope organization, metabolism, and virulence [13–17]. These changes may occur directly, when the altered resistance determinant is involved in core physiological processes, or indirectly through perturbation of complex regulatory networks. Notably, many of these networks are intertwined with virulence and pathogenicity pathways [18,19], suggesting that antimicrobial resistance can reshape infection dynamics in ways that extend beyond drug susceptibility.

Here, we show that loss-of-function mutations in *oprD* appear in clinical isolates in the

absence of prior carbapenem exposure, suggesting that they can provide infection-

related advantages beyond antibiotic resistance. Consistent with this, we observed that

these mutations increase *P. aeruginosa* attachment to and crossing of airway epithelial

barriers. Our data suggest that loss of OprD leads to a reorganization of the bacterial

surface and reduces mucus entrapment, thereby facilitating earlier access to the

epithelial surface and more rapid penetration. Together, these findings highlight how

examining antimicrobial resistance and bacterial physiology as interconnected traits can

reveal important links with direct consequences for infection outcomes.

Results

Mutations in oprD *appear in clinical isolates without prior carbapenem exposure but*

often disappear

We first investigated the clinical prevalence of *oprD* mutations in chronically infected

patients. For this, we examined a collection of 474 longitudinally sampled clinical

isolates of *P. aeruginosa* obtained from 34 chronically infected individuals with CF at the

Copenhagen CF Center [6]. Most mutations in *oprD* acquired during infection were

frameshifts, likely preventing porin production or assembly into the membrane (Figure

1A, Supplementary Table 1). These mutations were observed at multiple timepoints

throughout chronic infection. When comparing these data with carbapenem treatment

periods, we found that the appearance of *oprD*-mutated strains did not always coincide

with antibiotic exposure (Figure 1B). Importantly, carbapenem administration in these

patients was exclusively intravenous, as recorded in hospital treatment records, ensuring

that treatment histories are not confounded by undocumented oral use at different

timepoints. Consistent with this, meropenem Minimal Inhibitory Concentration (MIC)

testing showed that a substantial fraction of *oprD*-mutated isolates (26%) remained

susceptible according to EUCAST clinical breakpoints [20] (MIC ≤ 2; Supplementary Table

1), with susceptible isolates recovered both after carbapenem therapy and from patients

without prior exposure. Together, these observations indicate that the emergence of *oprD*

mutations is not solely driven by carbapenem pressure and may reflect infection-related

advantages beyond antibiotic resistance.

Despite their recurrent acquisition, *oprD* mutations were almost never maintained in the

infecting populations of this cohort (Figure 1B). The single case in which a mutation (3

nucleotide insertion) persisted throughout chronic infection (DK36 clone type) was also

the only one that did not cause a frameshift, and thus likely had a lower impact on protein

functionality. This suggests that while OprD loss may provide a selective advantage

during infection, the trait appears to be disadvantageous for long-term persistence in

chronic infection.

A

B

Figure 1. Characterization of *oprD* mutations in *P. aeruginosa* clinical isolates. (A) Non-

synonymous mutations identified in the *oprD* gene of *P. aeruginosa* clinical isolates from

chronically infected individuals with CF, belonging to different clone types [6]. Information

about the specific position of each mutation can be found in Supplementary Table 1. (B)

Temporal distribution of *oprD*-mutated strains during chronic infection, shown together

with carbapenem treatment periods of the corresponding patients. Isolates without

mutations are indicated by blue circles, isolates carrying *oprD* mutations by red circles,

and carbapenem treatment periods by orange rectangles.

Mutations in oprD *increase early attachment and translocation across airway epithelia*

The presence of OprD mutations independently of carbapenem treatment suggested that

OprD loss may confer infection-related advantages beyond antibiotic resistance. To

directly test this hypothesis, we engineered two PAO1 derivative *oprD* mutant strains

harboring previously identified mutations [6], to avoid the confounding effects of additional

mutations present in clinical genomes. One mutant carried a 13-nucleotide deletion

(positions 856–868) originally found in a strain with DK41 clone type (*oprD41*) that was

selected right after a carbapenem treatment period. The other reconstructed mutant

harbored a two-nucleotide (AC) deletion at positions 243–244 previously detected in a

clone type DK09 strain (*oprD09*), that appeared without carbapenem treatment (Figure

1).

We then compared the infection dynamics of these mutants with the wild-type PAO1

strain in a fully differentiated airway epithelium cultured at the air–liquid interface (ALI)

which exhibits barrier function and mucus production as a human airway epithelium *in*

vivo [18,21]. Following apical inoculation, bacterial growth, attachment to the epithelial cells

and translocation to the basolateral compartment were monitored from 1.5 to 14 hours

post-infection.

Both *oprD* mutants displayed consistently higher initial attachment and a markedly

increased capacity to cross the epithelial barrier compared with PAO1. CFU counts of

bacteria firmly attached to the epithelium (not removed by apical PBS washing) were

significantly higher for the mutants than for PAO1 as early as 1.5 hours post-inoculation

(Figure 2A, one-way ANOVA with Tukey's multiple comparisons test, $p < 0.05$ for *oprD*41

and $p < 0.001$ for *oprD*09). Mutants were also detected in the basolateral compartment by

10 hours post-infection, whereas wild-type PAO1 bacteria were absent at this time point,

and their numbers remained higher than PAO1 at 14 hours (Figure 2A, one-way ANOVA

with Tukey's multiple comparisons test, $p < 0.0001$). Confocal microscopy analysis

confirmed these findings: at 10 hours, mutant bacteria were observed to be already

penetrating the epithelium while PAO1 cells remained largely superficial, and by 14

hours, mutants showed substantially deeper and more abundant epithelial penetration

(Figure 2B). To assess whether these infection dynamics translated into detectable

disruption of the epithelial barrier, we measured transepithelial electrical resistance

(TEER) as a general indicator of tight junction integrity and thus overall epithelial damage.

TEER progressively declined over the course of all infections, consistent with the large

epithelial barrier disruption. However, no significant differences were detected between

strains at the monitored timepoints (Figure 2C, two-way ANOVA with Tukey's multiple

comparisons test, $p>0.05$).

These results indicate that loss of OprD enhances *P. aeruginosa* early adhesion to and

translocation across the airway epithelium, although this does not affect initial overall

tissue disruption.

P. aeruginosa (GFP)   F-actin (Phaloidin)   Nuclei (To-pro)

Figure 2. Effects of *oprD* mutations on airway infection dynamics of *P. aeruginosa*. (A)

Colony-forming units (CFUs) of PAO1, *oprD*41, and *oprD*09 in the apical compartment

(non-firmly attached cells, in yellow), attached to the epithelium (pink), and in the

basolateral compartment (blue), measured from 1.5 to 14 hours post-infection in fully

differentiated BCi-NS1.1 ALI airway epithelial cultures. Data represent mean ± SEM from

three biological replicates, each with three technical replicates. Statistical significance

was determined by compartment and timepoint by one-way ANOVA with Tukey's multiple

comparisons test and is indicated as *p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001 and

****p ≤ 0.0001. (B) Confocal microscopy images of BCi-NS1.1 ALI cultures at 10- and 14

hours post-infection with PAO1, *oprD*41, and *oprD*09. Bacteria are shown in green (GFP),

epithelial F-actin in red (Phalloidin-AF555), and nuclei in blue (TO-PRO-3). Nuclei signal

was omitted from maximum intensity projections for image clarity. Scale bar, 100 μm.

Images are representative of three independent experiments. (C) Fold change in

transepithelial electrical resistance (TEER) of fully differentiated BCi-NS1.1 ALI airway

epithelial cultures left uninfected (Mock, in grey) or infected with PAO1 (blue), *oprD*41

(dark red), or *oprD*09 (light red), calculated at the end of the experiment (1.5 to 14 hours)

relative to baseline values at the beginning. Data represent mean ± SEM from three

biological replicates, each with three technical replicates. Statistical significance was

assessed by two-way ANOVA with Tukey's multiple comparisons test; connecting letters

indicate significant differences between groups (*p*<0.05).

Mutations in oprD *are associated with changes in membrane organization and adherence*

to mucin

To investigate the mechanisms underlying the enhanced epithelial crossing of the *oprD*

mutants, we performed both phenotypic screenings and transcriptomic analyses on the

bacterial monocultures grown in laboratory media Luria–Bertani (LB). In *in vitro* virulence

analyses such as pyocyanin production, protease secretion, biofilm formation, and

swarming motility, the mutant strains showed no significant differences from the wild-

type PAO1 (Supplementary Figure 1A, 1B, 1C and 1D, one-way ANOVA with Tukey's

multiple comparisons test, p>0.05). Likewise, RNA-seq analysis revealed no significant

transcriptional changes between strains (Supplementary Figure 2A and 2B), with no

genes meeting differential expression thresholds ($|\log_2$ fold change$| \geq 1$, adjusted $p <$

0.05). These results indicate that the absence of OprD does not involve a global

transcriptional reorganization.

Because type III secretion (T3SS) has been described as a key determinant of epithelial

colonization and penetration [22], and its expression is strongly host-induced [23], we also

quantified the transcript levels of five T3SS genes representing different operons by RT-

qPCR from total RNA extracted from infected ALI cultures after 14 hours of infection.

Consistent with the RNA-seq data obtained in LB, no significant differences were

detected between PAO1 and the *oprD* mutants (Supplementary Figure 2C), supporting

that the enhanced epithelial phenotype is not due to altered T3SS expression.

We next assessed bacterial fitness in LB and synthetic cystic fibrosis sputum medium

(SCFM) which mimics the nutritional composition of the CF airways [24]. Growth of the

mutants was indistinguishable from PAO1 (Supplementary Figure 1E, one-way ANOVA

with Tukey's multiple comparisons test, $p>0.05$), ruling out a contribution of general

fitness. Antibiotic susceptibility profiling confirmed the expected increase in

carbapenem resistance and additionally revealed a modest decrease in colistin

susceptibility (Figure 3A, Supplementary Figure 3). Because colistin is an antimicrobial

peptide that interacts with the bacterial surface, we hypothesized that OprD loss altered

membrane properties.

Consistent with this hypothesis, we quantified alterations in outer membrane

organization. The *oprD* mutants exhibited a significantly more negative net surface

charge than PAO1 (Figure 3B, one-way ANOVA with Tukey's multiple comparisons test,

$p<0.05$ for *oprD*41 and $p<0.01$ for *oprD*09), as determined by zeta potential

measurements, which report on the electrostatic properties of the bacterial surface. To

further assess membrane organization, we used the fluorescent probe Laurdan, which is

sensitive to lipid packing and allows discrimination between changes in membrane

fluidity (generalized polarization, GP) and overall probe accessibility (fluorescence

intensity). Laurdan GP values remained unchanged ($GP_{PAO1} = 0.51 \pm 0.004$, $GP_{oprD41} = 0.52$

± 0.002, $GP_{oprD09} = 0.50 \pm 0.002$, one-way ANOVA with Tukey's multiple comparisons test,

$p>0.05$), indicating that lipid order and membrane fluidity are preserved. However, total

Laurdan fluorescence increased in the mutants (Figure 3C one-way ANOVA with Tukey's

multiple comparisons test, p<0.05), suggesting greater accessibility of outer membrane

components to external molecules, consistent with a structural reorganization of the

envelope.

We hypothesized that such alterations in surface properties could directly influence how

bacteria interact with the host environment. Given that mucus is a highly negatively

charged barrier that bacteria must overcome to establish infection [25,26], we next examined

mucin binding after incubating bacteria with commercially sourced mucin pre-coated

onto the wells of a microplate. Non-adherent cells were removed by washing, and the

number of bound bacteria was quantified by CFU counting. Notably, *oprD* mutants

showed impaired attachment compared to PAO1 (Figure 3D, one-way ANOVA with

Tukey's multiple comparisons test, p<0.0001), which may reduce their entrapment within

mucus and thereby facilitate earlier access to the epithelial surface and more rapid

penetration through the tissue.

These findings indicate that OprD loss primarily affects membrane properties and mucus

interaction, rather than global virulence regulation or bacterial fitness, providing a

plausible mechanistic basis for the enhanced epithelial penetration observed.

Figure 3. Phenotypic characterization of changes caused by *oprD* mutations in *P. aeruginosa*. (A) Antibiotic susceptibility of *oprD*41 and *oprD*09 relative to the parental PAO1 strain against tobramycin (TOB), amikacin (AMK), colistin (COL), ciprofloxacin (CIP), ceftazidime (CAZ), piperacillin/tazobactam (PIP/TAZ), aztreonam (AZT), and meropenem (MER). Bacterial growth data (OD$_{600}$ values at the end of the assay across antibiotic concentrations) are provided in Supplementary Figure 2. (B) Surface charge and (C) Laurdan fluorescence intensity for PAO1 (blue), *oprD*41 (dark red), and *oprD*09 (light red). Surface charge values are shown as mean ± SEM from three biological replicates with twenty technical replicates each, whereas Laurdan fluorescence total intensity is presented as mean ± SEM from three biological replicates with three technical replicates each. Statistical significance was determined by one-way ANOVA with Tukey's multiple comparisons test and is indicated as *p ≤ 0.05 and **p ≤ 0.01. (D)

Mucin adherence of PAO1 (blue), *oprD*41 (dark red), and *oprD*09 (light red), expressed as

the ratio of attached to non-attached bacteria after incubation with mucin. Data

represent mean ± SEM from three biological replicates with three technical replicates

each. Statistical significance was determined by one-way ANOVA with Tukey's multiple

comparisons test and is indicated as ****$p \leq 0.0001$.

Effects of oprD *mutations on airway infection are preserved in clinical genomic*

backgrounds

Finally, to validate the relevance of these findings in patient-derived isolates, we tested

whether the infection phenotype was preserved in clinical genomic backgrounds. For

that, we analyzed epithelial penetration of pairs of clinical strains with or without *oprD*

mutation, isolated from the same sputum sample and phylogenetically related,

minimizing the number of confounding genetic differences (Supplementary Table 2). The

pairs, previously whole-genome sequenced [6], belonged to clone types DK06 (strains 305

and 306) and DK44 (strains 330 and 333). In DK06, strain 306 carried a deletion of a C at

position 927 of *oprD* and had emerged after carbapenem therapy (Figure 1), whereas

strain 305 retained the wild-type allele. In DK44, strain 330 harbored a two-nucleotide

insertion (AC) at position 184, while strain 333 carried an intact *oprD* gene; these isolates

were obtained from a patient who had not received prior carbapenem treatment (Figure

1).

Infections were performed in the airway epithelium ALI model, with bacterial growth,

attachment and translocation to the basolateral side assessed at 14 and 22 hours post-

inoculation. Consistent with the phenotype observed in the PAO1 background, the *oprD*

mutant clinical strains (306 and 330) were detected in the basolateral compartment as

early as 14 hours, whereas their *oprD* wild-type counterparts (305 and 333) were absent

at this time point, and the *oprD* mutant strains were attached to the epithelium in higher

numbers (Figure 4A, left panel; t-test with Holm–Šídák multiple comparisons test,

$p<0.05$). By 22 hours, bacteria from all strains were detectable in the basolateral

compartment, but *oprD* mutants remained significantly more abundant than their *oprD*

wild-type counterparts (Figure 4A, right panel; t-test with Holm–Šídák multiple

comparisons test, p<0.05). Confocal microscopy images at 22 hours post-infection
visually confirmed these quantitative results, showing deeper epithelial penetration of
the *oprD*-mutated strains across distinct colonization patterns (Figure 4B).

TEER values, measured as an indicator of epithelial barrier integrity, decreased
progressively over time in all strains (Figure 4C; two-way ANOVA with Tukey's multiple
comparisons test, p>0.05 between strains). However, no significant TEER reduction
relative to uninfected controls was observed at 14 hours post-infection for any strain
(Figure 4C; two-way ANOVA with Tukey's multiple comparisons test, p>0.05), indicating
that these clinical isolates are less damaging to the epithelium than the laboratory strain
PAO1 (Figure 2C).

Consistent with the observations in the PAO1 background, the *oprD* mutant clinical
strains also showed reduced adhesion to mucin compared with their *oprD* wild-type
counterparts (Figure 4D, two-way ANOVA with Tukey's multiple comparisons test,
p ≤ 0.0001), suggesting that loss of OprD similarly decreases mucus entrapment in
clinical isolates and may contribute to their enhanced epithelial translocation.

Together, these findings demonstrate that the enhanced epithelial translocation
conferred by *oprD* mutations is preserved in clinical isolates with distinct infection
dynamics and virulence, underscoring the relevance of this phenotype in patient
infection contexts.

Figure 4. Effects of *oprD* mutations on airway infection dynamics of *P. aeruginosa* clinical

strains. (A) Colony-forming units (CFUs) of clinical strains 305 (dark purple), 306 (light

purple), 333 (dark green), and 330 (light green) recovered from the apical compartment

(non-attached cells), firmly attached to the epithelium, and from the basolateral

compartment at 14 (left panel) and 22 hours (right panel) post-infection in fully

differentiated BCi-NS1.1 ALI airway epithelial cultures. Data represent mean ± SEM from

three biological replicates, each performed in technical triplicates. Statistical

significance between strains of the same clone type was assessed by t-test with Holm–

Šídák multiple comparisons test and is indicated as *p ≤ 0.05, **p ≤ 0.01 and

***p ≤ 0.001. (B) Confocal microscopy images of BCi-NS1.1 ALI cultures at 22 hours

post-infection with clinical strains 305, 306, 333, and 330. Bacteria are shown in green

(GFP), epithelial F-actin in red (Phalloidin-AF555), and nuclei in blue (TO-PRO-3). Nuclei

signal was omitted from maximum intensity projections for image clarity. Scale bar,

100 μm. Images are representative of three independent experiments. (C) Fold change in

transepithelial electrical resistance (TEER) of fully differentiated BCi-NS1.1 ALI airway

epithelial cultures left uninfected (Mock, in grey) or infected with clinical strains 305 (dark

purple), 306 (light purple), 333 (dark green) or 330 (light green), calculated at the end of

the experiment (14 or 22 hours) relative to the values at the beginning. Data represent

mean ± SEM from three biological replicates, each with three technical replicates.

Statistical significance was assessed by two-way ANOVA with Tukey's multiple

comparisons test; connecting letters indicate significant differences between groups

($p<0.05$). (C) Mucin adherence of clinical strains 305 (dark purple), 306 (light purple), 333

(dark green) and 330 (light green), expressed as the ratio of attached to non-attached

bacteria after incubation with mucin. Data represent mean ± SEM from three biological

replicates with three technical replicates each. Statistical significance was determined

by two-way ANOVA with Tukey's multiple comparisons test and is indicated as

***p ≤ 0.0005 and ****p ≤ 0.0001.

Discussion

Antibiotic resistance is typically only studied in the narrow context of drug susceptibility,

yet its broader effects on bacterial physiology and host interactions remain

underexplored. Resistance mutations can reconfigure fundamental aspects of bacterial

biology, thereby shaping infection outcomes in ways not directly related to antimicrobial

therapy. Here, we provide a clear example: loss-of-function mutations in the carbapenem

porin OprD in *P. aeruginosa* alter infection dynamics independently of resistance.

Specifically, we show that *oprD* mutants exhibit reduced mucin adhesion and enhanced

epithelial translocation (Figure 5).

This phenotype appears to arise from structural alterations of the outer membrane rather

than transcriptional reprogramming. Although direct genetic reversion of only the

membrane changes caused by OprD loss is not feasible, and chemical modulation of

membrane charge would also affect the host environment, the convergence of our data

strongly supports membrane reorganization as the most plausible driver of the observed

effects. This interpretation is supported by parallels with *Streptococcus pneumoniae*,

whose polysaccharide capsule, particularly when highly negatively charged, reduces

mucus-mediated clearance, facilitating transit to and colonization of the epithelial

surface [27]. Notably, *P. aeruginosa* has been shown to breach the airway epithelium via

goblet cell invasion [22], which requires overcoming entrapment within the mucin secreted

by these cells. This suggests that surface modifications reducing mucus entrapment may

represent a conserved strategy among respiratory pathogens.

Interestingly, we found that clinical *oprD* mutations can arise both with and without prior

carbapenem treatment. Notably, the clinical mutations analyzed in this study, both

acquired following carbapenem therapy and in patients without exposure, conferred

similar infection-relevant phenotypes. This observation underscores that selective

forces acting during infection are not limited to antibiotic exposure. Indeed, previous

reports have also described *oprD* mutations emerging in patients without therapy [28],

further supporting the idea that the selective advantage of *oprD* loss extends beyond drug

resistance. One possible explanation is that reduced mucin binding facilitates evasion of

mucociliary clearance, providing a selective advantage during airway colonization even

in the absence of antibiotic pressure. Importantly, we found that these infection-related

phenotypes were maintained across clinical strains with distinct genomic backgrounds,

virulence and infection dynamics, indicating that the effect of *oprD* loss is robust and not

restricted to a particular strain context. Since the enhanced epithelial penetration

observed in *oprD* mutants likely stems from an altered interaction with mucus that

promotes earlier contact with epithelial cells, rather than from changes in their

colonization strategy, the phenotype can be maintained across strains with different

infection dynamics. Nevertheless, we also observed that *oprD* mutations were rarely

maintained over time in chronic infections. This transient presence suggests an

evolutionary trade-off, as OprD loss facilitates short-term invasion of the airway

epithelium but may incur fitness costs, possibly immunological, that limit long-term

persistence during chronic colonization. In this sense, *oprD* mutations exemplify the

classic evolutionary trade-off in host-pathogen interactions: what benefits acute

invasion may ultimately hinder long-term persistence [29]. Still, it is unclear which other

selective pressures may select for *oprD* mutations.

Despite their limited fixation, *oprD* mutations can have clinically relevant consequences.

For example, it was reported that, in one CF patient, the acquisition of an *oprD* mutation

after carbapenem therapy coincided with a rare increase in virulence during chronic

infection [30]. Although concomitant mutations in the type III secretion system likely

contributed substantially to this phenotype, the simultaneous emergence of an *oprD*

mutation suggests a possible combined effect. Consistent with this, our data show that

the enhanced epithelial penetration associated with *oprD* loss is maintained across

clinical isolates with different baseline levels of virulence, as reflected by TEER

measurements of the performed infections, indicating that this phenotype is largely

independent of strain-specific virulence traits. More broadly, a recent large-scale

analysis found that *oprD* mutations were the strongest genomic predictors of septic

shock in *P. aeruginosa* infections, outperforming even classical virulence genes [31].

Together, these findings indicate that *oprD* mutations may shape clinical trajectories in

ways not captured by traditional views of resistance.

In addition to their impact on patient outcomes, the enhanced translocation ability of

oprD mutants may also influence bacterial dissemination. In a murine model of gut

infection, *oprD* mutants displayed higher fitness when bacterial recovery from the spleen

was used as a readout [32], consistent with our mechanistic observations of improved

epithelial penetration. Furthermore, in critically ill patients, a well-described gut-to-lung

translocation event of *P. aeruginosa* coincided with carbapenem treatment and the

emergence of *oprD* mutations [33]. While these mutations were originally interpreted

merely as genetic markers to track strain identity, our results suggest they may actively

promote epithelial traversal, facilitating within-host dissemination.

Taken together, our findings reveal that *oprD* loss-of-function mutations promote airway

barrier penetration, likely through alterations in outer membrane properties and mucus

interaction. However, their lack of stable persistence in chronic infection highlights a

trade-off between short-term invasiveness and long-term adaptation. More broadly, our

study illustrates that resistance mutations should not be viewed solely through the lens

of antibiotic susceptibility. Instead, their interconnection with bacterial physiology can

have significant implications for infection dynamics and clinical outcomes beyond the

reduction of drug susceptibility. Recognizing these dual roles may ultimately improve

how we interpret bacterial genomes in clinical microbiology and refine our predictions of

infection progression, with direct implications for patient management.

Figure 5. Schematic representation of the effects of *oprD* mutations in *P. aeruginosa*

infection. Loss-of-function mutations in *oprD* alter outer membrane organization and

increase surface negative charge. These changes likely reduce bacterial binding to

mucus, thereby facilitating closer contact with the epithelial surface at early timepoints.

This enhanced epithelial interaction promotes more efficient colonization and

subsequent translocation across the airway barrier.

Materials and methods

Bacterial strains growth and antibiotic susceptibility testing

The *P. aeruginosa* laboratory strain PAO1 was employed as the reference background for constructing recombinant strains [34]. *P. aeruginosa* clinical strains used in this study were previously described and whole-genome sequenced [6]. The local ethics committee at the Capital Region of Denmark (Region Hovedstaden) approved the use of the stored *P. aeruginosa* isolates (registration number H-21078844). Bacterial strains were cultured routinely in LB medium at 37 °C under aerobic conditions.

Bacterial growth was monitored in a BioTek ELx808 Absorbance Microtiter Reader (BioTek Instruments) at 37 °C with shaking at 250 rpm. Optical density (OD) at 600 nm was recorded at 20-minutes intervals over a 24-hours period. Cultures were initiated at an OD_{600} of 0.01, obtained by diluting overnight LB precultures into fresh LB or SCFM. SCFM was prepared following the protocol described previously [35], excluding DNA and mucins to reduce viscosity.

The minimum inhibitory concentrations (MICs) of ciprofloxacin, ceftazidime, tobramycin, amikacin, colistin, piperacillin/tazobactam, aztreonam, and meropenem were determined for multiple *P. aeruginosa* strains using the broth microdilution method in Mueller–Hinton (MH) medium, following the guidelines established by the European Committee on Antimicrobial Susceptibility Testing (EUCAST) [36].

Construction of bacterial strains

Genetic modifications were carried out using a CRISPR-Cas9-based recombineering system with a protocol previously described [37], that is briefly described below.

Electrocompetent *P. aeruginosa* PAO1 cells were prepared by washing twice with 300 mM sucrose, followed by electroporation with the pSH624-ssr plasmid. Transformants were selected on LB-agar plates supplemented with gentamicin (50 μg/mL). Spacer sequences targeting the *oprD41* mutation (Fw: 5'–GCGCGGGATGCGCACACTTTCACCT–3'; Rv: 5'–AAACAGGTGAAAGTGTGCGCATCCC–3') and *oprD09* mutation (Fw: 5'–GCGCGTATGAATCCGGCTTCACCCA–3'; Rv: 5'–AAACTGGGTGAAGCCGGATTCATAC–3')

were phosphorylated using T4 Polynucleotide Kinase (New England Biolabs) and cloned

into Eco31I (FastDigest, Thermo Fisher Scientific)-digested pS548-CsR vector via T4 DNA

ligase (Thermo Fisher Scientific), following the manufacturers' protocols.

To induce ssr expression, an overnight culture of PAO1 pSH624-ssr was incubated with 1

mM IPTG for 3 hours. Subsequently, the constructed pS548-CsR-*oprD41* and pS548-

CsR-*oprD09* plasmids were introduced by electroporation together with the

recombineering oligonucleotides containing the desired genetic modification flaked by

50 nucleotides upstream and downstream of its genetic context. Transformants were

selected on LB agar containing tetracycline (100 µg/mL), and Cas9 activity was induced

by adding 5 mM 3-methylbenzoate (3-mBz) into the LB-agar plate. Successful genome

editing was confirmed by PCR screening and Sanger sequencing of the targeted genomic

region.

GFP-tagging of bacterial strains

To enable visualization by confocal microscopy, *P. aeruginosa* strains were genetically

modified to express GFP. Tagging was achieved by conjugative transfer of the plasmid

pUC18T-mini-Tn7T-Gm-Tp-gfp as previously described [18,38]. The process involved a four-

parental mating, in which equal volumes of cultures of *Escherichia coli* DH5α/pUC18T-

mini-Tn7T-Gm-Tp-gfp, *E. coli* SM10(λpir)/pTNS1, *E. coli* HB101/pRK2013, and the target *P.*

aeruginosa strain were applied onto a 3-mm cellulose acetate filter placed on non-

selective LB agar. Plates were incubated overnight at 37 °C. Transformants were selected

on Pseudomonas Isolation Agar (PIA) supplemented with gentamycin (50 µg/mL).

Subsequently, the antibiotic resistance cassette was removed by Flp-mediated excision

using the plasmid pFLP2. For this step, electrocompetent *P. aeruginosa* cells from the

previous conjugation were prepared by two washes with 300 mM sucrose, followed by

electroporation with pFLP2. Transformants were selected on LB agar containing

carbenicillin (200 µg/mL). Plasmid curing was performed by streaking onto LB agar with

5% sucrose. GFP expression in the final strains was confirmed by detection of green

fluorescence.

Air Liquid Interface infection model preparation

BCi-NS1.1 cells [21] were employed to establish ALI-based infection systems as previously

described [17,18]. Cells were first expanded in standard culture flasks under humidified

conditions at 37 °C and 5% CO_2, using PneumaCult™-Ex Plus medium (STEMCELL

Technologies) according to the manufacturer's guidelines. For ALI setup, $1×10^5$ cells were

seeded onto 1 μm pore polyester membrane inserts (Falcon) pre-coated with type I

collagen (Gibco). Once full confluence was achieved, the apical medium was removed,

and the basolateral compartment was replenished with PneumaCult™-ALI maintenance

medium (STEMCELL Technologies) supplemented with 4 μg/mL heparin, 480 ng/mL

hydrocortisone, PneumaCult™-ALI 10× supplement, and PneumaCult™-ALI maintenance

supplement. Cultures were maintained for 28 days at 37 °C and 5% CO_2, with media

replacement every 3–4 days. Epithelial polarization was monitored by regular

measurements of transepithelial electrical resistance (TEER) using an STX2 chopstick

electrode (World Precision Instruments). From day 15 onward, accumulated mucus was

removed weekly by rinsing the apical surface with PBS.

Infection procedure and characterization

Overnight cultures of *P. aeruginosa* were diluted in LB medium to an initial OD_{600} of 0.01

and incubated until reaching mid-exponential growth. One milliliter of culture was then

pelleted by centrifugation (4500 × g, 5 minutes), washed with PBS, and resuspended in

PBS to a final concentration of $1×10^5$ CFU/mL. For infection, 10 μL of this suspension

(≈$1×10^3$ CFU) were applied to the apical compartment of fully differentiated BCi-NS1.1

ALI cultures, with 600 μL of supplemented PneumaCult™-ALI maintenance medium in the

basolateral chamber. Control wells received the same volume of sterile PBS. Cultures

were incubated at 37 °C and 5% CO_2 for the desired infection period, after which 200 μL

of PBS were added to the apical chamber.

Apical washes (200 μL) and basolateral media (600 μL) were collected for CFU counting

by serial dilution and plating on LB agar. To quantify adherent bacteria, the apical surface

was rinsed with 200 μL PBS, epithelial cells were detached using a cell scraper, and

resulting suspensions were serially diluted and plated.

For confocal microscopy, infected ALI cultures were rinsed with PBS and fixed with 4%

paraformaldehyde applied to both apical and basolateral chambers for 20 minutes at 4

°C. After three PBS washes, samples were permeabilized and blocked with PBS

containing 3% BSA, 1% saponin, and 1% Triton X-100 for 1 hour at room temperature.

Staining was performed by adding 100 µL of Phalloidin-AF555 (1:400, Invitrogen) and TO-

PRO-3 (1:1000, BioLegend) diluted in 3% BSA/1% saponin-PBS to the apical chamber for

2 hours at room temperature. Inserts were then excised, mounted on glass slides with

VECTASHIELD® Antifade Mounting Medium (VWR), and imaged using a Leica Stellaris 8

confocal microscope (40× oil objective, NA 1.3). Images were processed with LAS X

software v1.4.4.26810 (Leica).

RNA preparation and RNA-sequencing

Bacteria were inoculated into LB medium at an initial OD_{600} of 0.01 and cultured at 37 °C

with shaking at 200 rpm until reaching mid-exponential phase ($OD_{600} \approx 0.6$). Three

independent biological cultures were assessed for PAO1, *oprD*41, and *oprD*09. Prior to

RNA extraction, transcription was blocked by adding RNA Protect Bacteria solution

(Qiagen) following the manufacturer's protocol. Total RNA was then isolated using the

RNeasy Mini Kit (Qiagen) according to the supplier's instructions. RNA quality and

integrity were assessed using a Fragment Analyzer 2100 (Agilent Technologies). Samples

were run in a NovaSeq Illumina sequencer with 2x150bp configuration yielding 10 million

reads per sample.

Reads were trimmed, and low-quality reads and potential contamination from adapters

were removed using Trimmomatic (v0.35) tool [39]. Reads were further processed using

SortMeRNA tool (v2.1) [40] to remove reads generated from residual rRNA transcripts.

Reads were mapped against PAO1 genome (NCBI: NC_002516.2) using BWA-MEM

algorithm, and duplicated reads were marked using Picard tools. Reads mapping on each

annotated coding sequence were counted using htseq-count v0.7.2 [41] imported and

processed in RStudio.

Counts were normalized using log2-negative binomial transformation performed using

the rld transformation function contained in the R package DESeq2 [42] with the option

blind set as "true." Normalized counts were used to evaluate whole transcriptome

similarities using hierarchical clustering analysis (HCA), principal component analysis

(PCA), and k-mean clustering. DEG analysis between transcriptomes was performed

using the R package DESeq2, considering statistically significant genes with a Log2(fold

change) $\geq |1|$ and an adjusted P $\leq$ 0.01.

RNA extraction and quantitative RT-PCR from infected cultures

Fully differentiated BCi-NS1.1 ALI cultures were infected with approximately 10^3 CFU of

each bacterial strain as described above. For each strain, three biological replicates were

performed, each consisting of two cell culture inserts pooled to increase bacterial RNA

yield. After 14 hours incubation at 37 °C and 5 % CO_2, transcription was stopped by

adding 300 µL RNAprotect Bacteria Reagent (Qiagen) per replicate. Total RNA was

extracted using the RNeasy Mini Kit (Qiagen) according to the manufacturer's

instructions and treated with TURBO DNase (Invitrogen) to remove residual genomic

DNA. cDNA synthesis was performed from 10 µg of RNA with the GoScript Reverse

Transcription Mix (Promega). Quantitative PCR was run on a QuantStudio 5 Real-Time

PCR System (Applied Biosystems) using 96-well plates and GoTaq qPCR Master Mix

(Promega), with 100 ng cDNA per reaction. Cycling conditions were 95 °C for 10 min

followed by 40 cycles of 95 °C for 15 s and 60 °C for 1 min. Primers (400 nM) are listed in

Supplementary Table 3. Relative expression was calculated by the $2^{-\Delta\Delta CT}$ method [43] using

rplU as housekeeping reference. Data represents the mean of three biological replicates

with three technical replicates each.

Protease secretion and pyocyanin production

Protease secretion and pyocyanin production were measured as previously performed

[17].Overnight cultures of *P. aeruginosa* were adjusted to an OD_{600} of 0.01 in 10 mL of LB

medium and incubated for 24 hours at 37 °C in 50 mL flasks under constant 250 rpm

agitation. Each experiment was performed in three independent biological replicates.

One milliliter of each resulting culture was centrifuged (7000 × g, 5 minutes), and the

supernatant was sterilized by filtration through a 0.22 µm membrane. Two microliters of

the sterile supernatant were spotted onto 1% skim milk agar plates (prepared with 1%

agar), and proteolytic activity was evaluated by measuring clearing zones after 24 hours

of incubation at 37 °C. The same supernatant was also used for semi-quantitative

assessment of pyocyanin production by measuring absorbance at 690 nm in 100 µL

aliquots using a BioTek ELx808 Absorbance Microtiter Reader (BioTek Instruments).

Biofilm production

Biofilm production was assessed as previously described [44]. *P. aeruginosa* strains were

grown in Nunc™ MicroWell™ 96-Well Flat-Bottom Microplates (Thermo Fisher Scientific)

containing 150 µL of LB medium per well, with Nunc™ Immuno TSP Lids (Thermo Fisher

Scientific) replacing the standard lids. Bacteria were added at an initial OD_{600} of 0.05.

Plates were incubated at 37 °C for 20 hours to allow biofilm formation on the pegs. After

incubation, the peg lids were rinsed by immersion in a microtiter plate filled with 180 µL

PBS to remove non-adherent cells. Subsequently, the lids were transferred to a plate

containing 160 µL of 0.01% crystal violet solution for 15 minutes to stain the biofilms.

Following staining, the lids underwent three washes in separate PBS-filled microtiter

plates (180 µL per well) to eliminate excess dye. Finally, pegs were transferred to a fresh

plate with 180 µL of 99% ethanol to solubilize the bound crystal violet. The ethanol

solution was then used to quantify biofilm biomass by measuring absorbance at 590 nm

in a BioTek ELx808 Absorbance Microtiter Reader (BioTek Instruments).

Motility assay

Swarming motility was assessed using Petri dishes containing 25 mL of a defined medium

composed of 0.5% casamino acids, 0.5% Bacto agar, 0.5% filtered glucose, 3.3 mM

K_2HPO_4, and 3 mM $MgSO_4$, as previously performed [45]. Four microliters of *P. aeruginosa*

cultures adjusted to an OD_{600} of 1.0 were spotted onto the center of each plate. Plates

were incubated at 37 °C for 17 hours. Experiments included three technical replicates per

strain. Swarming areas were quantified by analyzing photographic images with ImageJ

software, measuring the surface area (cm^2) covered by bacterial spread.

Membrane properties determination

Zeta potential measurements of *P. aeruginosa* strains were performed using a Zetasizer

NANO-ZS (Malvern Instruments) equipped with a universal dip cell (ZEN1002). Overnight

cultures were washed once with ultrapure water to remove residual medium and salts,

then adjusted to an OD_{600} of 1. Measurements were obtained from three independent

biological replicates, each comprising 30 readings. Electrophoretic mobility values were

converted to zeta potentials using the Smoluchowski equation.

Membrane fluidity was assessed via Laurdan staining. Washed bacterial suspensions at

OD_{600} = 1 in PBS were incubated with 2.5 µM Laurdan dye for 60 minutes at room

temperature in the dark. 150 µL of samples and blanks were aliquoted into Nunc™ 96-

Well Optical-Bottom black microplates (Thermo Fisher Scientific) with three technical

replicates each. Fluorescence was measured at emission wavelengths of 440 and 490

nm upon excitation at 350 nm in a BioTek ELx808 Absorbance Microtiter Reader (BioTek

Instruments). Laurdan GP values were calculated using the formula:

$GP=(I_{440}-I_{490})/(I_{440}+I_{490})$, where I_{440} and I_{490} represent fluorescence intensities at 440 nm and

490 nm, respectively.

Bacterial mucin adhesion assay

Adhesion assays were conducted as previously performed [46] with minor modifications.

Wells of Nunc™ MicroWell™ 96-Well Flat-Bottom Microplate (Thermo Fisher Scientific)

were coated with 200 µL of porcine mucin (type II, Sigma-Aldrich, M2378) solution (10

µg/mL in PBS), while control wells received 200 µL of PBS. Plates were incubated at 37 °C

for 18 hours prior to the assay. Overnight bacterial cultures were washed and diluted in

PBS to a final concentration of $5×10^7$ CFU/mL. PBS was aspirated from mucin-coated and

control wells, which were then inoculated in triplicate with 100 µL of the bacterial

suspension (≈$5×10^6$ CFU). Plates were incubated for 30 minutes at 37 °C to allow

bacterial adhesion. After incubation, the supernatant was removed, and wells were

washed five times with 200 μL PBS to eliminate non-attached cells. The wash solutions

were plated in serial dilutions to quantify non-attached bacteria. To detach adherent

bacteria along with mucin, 200 μL of 1% (v/v) Triton X-100 in PBS was added to each well

and incubated at room temperature for 30 minutes. Serial dilutions of the resulting

suspensions were plated to determine CFUs of adhered bacteria. Adhesion efficiency

was expressed as the ratio of adhered to non-adhered bacteria. All assays were

performed in three biological replicates.

Statistical analysis

All statistical analyses were performed with GraphPad Prism version 10.3.1 (GraphPad

Software). Depending on the experimental design, differences among groups were

assessed using ordinary one-way ANOVA or two-way ANOVA followed by Tukey's post

hoc test for multiple comparisons, or by unpaired *t*-tests with Holm–Šídák correction for

multiple comparisons. A p-value <0.05 was considered indicative of statistical

significance.

Data availability

All data supporting the findings of this study are available within the paper, its

Supplementary Materials, and Supplementary Data files. Raw sequencing data have

been deposited in the NCBI Sequence Read Archive (SRA) under BioProject accession

number PRJNA1346407. Source data are provided with this paper.

Acknowledgements

We thank the Infection Microbiology group for their insightful comments and discussion. The Basal Cell Immortalized Non-Smoker 1.1 (BCi-NS1.1) cell line was a gift from Professor Ronald G. Cristal (Weil Cornell Medical College, New York, USA). This research was funded by the "Novo Nordisk Foundation Center for Biosustainability (CfB)" grant number NNF10CC1016517 and by the Challenge Grant NNF19OC0056411 to H.K.J. from The Novo Nordisk Foundation. P.L. is the recipient of an ERS/EU RESPIRE4 Marie Skłodowska-Curie Postdoctoral Research Fellowship (Ref. nr: R4202305-01047; this project has received funding from the European Respiratory Society and the European Union's H2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 847462).

Author contributions

PL designed the study and performed the experimental work. CAC performed the bioinformatic analysis of transcriptomic data. CAC, RLR, HKJ and SM participated in the design of the study. All authors participated in writing the manuscript and approved the submitted version.

Conflict of interest

The authors declare to have no conflict of interest.

References

1. Murray, C. J. *et al.* Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet* **399**, 629–655 (2022).

2. Macesic, N., Uhlemann, A. C. & Peleg, A. Y. Multidrug-resistant Gram-negative bacterial infections. *The Lancet* **405**, 257–272 (2025).

3. Laborda, P., Sanz-García, F., Hernando-Amado, S. & Martínez, J. L. Pseudomonas aeruginosa: an antibiotic resilient pathogen with environmental origin. *Curr Opin Microbiol* **64**, 125–132 (2021).

4. Laborda, P., Hernando-Amado, S., Martínez, J. L. & Sanz-García, F. Antibiotic Resistance in Pseudomonas. *Adv Exp Med Biol* **1386**, 117–143 (2022).

5. Qin, S. *et al.* Pseudomonas aeruginosa: pathogenesis, virulence factors, antibiotic resistance, interaction with host, technology advances and emerging therapeutics. *Signal Transduct Target Ther* **7**, 1–27 (2022).

6. Marvig, R. L., Sommer, L. M., Molin, S. & Johansen, H. K. Convergent evolution and adaptation of Pseudomonas aeruginosa within patients with cystic fibrosis. *Nat Genet* **47**, 57–64 (2015).

7. Weimann, A. *et al.* Evolution and host-specific adaptation of Pseudomonas aeruginosa. *Science (1979)* **385**, eadi0908 (2024).

8. World Health Organization. *WHO Bacterial Priority Pathogens List, 2024: Bacterial Pathogens of Public Health Importance, to Guide Research, Development, and Strategies to Prevent and Control Antimicrobial Resistance*. (2024).

9. Canton, R., Doi, Y. & Simner, P. J. Treatment of carbapenem-resistant Pseudomonas aeruginosa infections: a case for cefiderocol. *Expert Rev Anti Infect Ther* **20**, 1077–1094 (2022).

10. Sati, H. *et al.* The WHO Bacterial Priority Pathogens List 2024: a prioritisation study to guide research, development, and public health strategies against antimicrobial resistance. *Lancet Infect Dis* **25**, 1033–1043 (2025).

11. Darby, E. M. *et al.* Molecular mechanisms of antibiotic resistance revisited. *Nat Rev Microbiol* **21**, 280–295 (2022).

12. Wu, W., Huang, J. & Xu, Z. Antibiotic influx and efflux in Pseudomonas aeruginosa: Regulation and therapeutic implications. *Microb Biotechnol* **17**, e14487 (2024).

13. Blanco, P. *et al.* Bacterial Multidrug Efflux Pumps: Much More Than Antibiotic Resistance Determinants. *Microorganisms* **4**, 14 (2016).

14. Laborda, P., Gil-Gil, T., Martínez, J. L. & Hernando-Amado, S. Preserving the
efficacy of antibiotics to tackle antibiotic resistance. *Microb Biotechnol* **17**,
e14528 (2024).

15. Sanz-García, F. *et al.* Translating eco-evolutionary biology into therapy to tackle
antibiotic resistance. *Nat Rev Microbiol* **21**, 671–685 (2023).

16. Kong, K. F. *et al.* Pseudomonas aeruginosa AmpR is a global transcriptional factor
that regulates expression of AmpC and PoxB β-lactamases, proteases, quorum
sensing, and other virulence factors. *Antimicrob Agents Chemother* **49**, 4567–
4575 (2005).

17. Goltermann, L. *et al.* Macrolide resistance through uL4 and uL22 ribosomal
mutations in Pseudomonas aeruginosa. *Nat Commun* **15**, 8906 (2024).

18. Laborda, P. *et al.* Mutations in the efflux pump regulator MexZ shift tissue
colonization by Pseudomonas aeruginosa to a state of antibiotic tolerance. *Nat
Commun* **15**, 2584 (2024).

19. Laborda, P., Molin, S., Johansen, H. K., Martínez, J. L. & Hernando-Amado, S. Role
of bacterial multidrug efflux pumps during infection. *World J Microbiol Biotechnol*
40, 226 (2024).

20. European Committee on Antimicrobial Susceptibility Testing (EUCAST).
Breakpoint tables for interpretation of MICs and zone diameters. Version 15.0.
(2025).

21. Walters, M. S. *et al.* Generation of a human airway epithelium derived basal cell
line with multipotent differentiation capacity. *Respir Res* **14**, 135 (2013).

22. Leoni Swart, A. *et al.* Pseudomonas aeruginosa breaches respiratory epithelia
through goblet cell invasion in a microtissue model. *Nat Microbiol* **9**, 1725–1737
(2024).

23. Dasgupta, N., Ashare, A., Hunninghake, G. W. & Yahr, T. L. Transcriptional
Induction of the Pseudomonas aeruginosa Type III Secretion System by Low Ca2+
and Host Cell Contact Proceeds through Two Distinct Signaling Pathways. *Infect
Immun* **74**, 3334 (2006).

24. Turner, K. H., Wessel, A. K., Palmer, G. C., Murray, J. L. & Whiteley, M. Essential
genome of Pseudomonas aeruginosa in cystic fibrosis sputum. *Proc Natl Acad
Sci U S A* **112**, 4110–4115 (2015).

25. Sheng, Y. H. & Hasnain, S. Z. Mucus and Mucins: The Underappreciated Host
Defence System. *Front Cell Infect Microbiol* **12**, 856962 (2022).

26. Linden, S. K., Sutton, P., Karlsson, N. G., Korolik, V. & McGuckin, M. A. Mucins in
the mucosal barrier to infection. *Mucosal Immunol* **1**, 183–197 (2008).

27. Nelson, A. L. *et al.* Capsule Enhances Pneumococcal Colonization by Limiting
Mucus-Mediated Clearance. *Infect Immun* **75**, 83 (2006).

28. Wolter, D. J. *et al.* Emergence of carbapenem resistance in Pseudomonas
aeruginosa isolates from a patient with cystic fibrosis in the absence of
carbapenem therapy. *Clinical Infectious Diseases* **46**, e137-41 (2008).

29. La Rosa, R., Molin, S. & Johansen, H. K. Pseudomonas aeruginosa: persistence
beyond antibiotic resistance. *Trends Microbiol* **33**, 1076–1084 (2025).

30. Jorth, P. *et al.* Cystic fibrosis lung function decline after within-host evolution
increases virulence of infecting pseudomonas aeruginosa. *Am J Respir Crit Care*
Med **203**, 637–640 (2021).

31. Valik, J. K. *et al.* Genomic virulence markers are associated with severe outcomes
in patients with Pseudomonas aeruginosa bloodstream infection.
Communications Medicine **4**, 264 (2024).

32. Skurnik, D. *et al.* Enhanced in vivo fitness of carbapenem-resistant oprD mutants
of Pseudomonas aeruginosa revealed through high-throughput sequencing. *Proc*
Natl Acad Sci U S A **110**, 20747–20752 (2013).

33. Wheatley, R. M. *et al.* Gut to lung translocation and antibiotic mediated selection
shape the dynamics of Pseudomonas aeruginosa in an ICU patient. *Nat Commun*
13, 6523 (2022).

34. Stover, C. K. *et al.* Complete genome sequence of Pseudomonas aeruginosa
PAO1, an opportunistic pathogen. *Nature* **406**, 959–964 (2000).

35. Turner, K. H., Wessel, A. K., Palmer, G. C., Murray, J. L. & Whiteley, M. Essential
genome of Pseudomonas aeruginosa in cystic fibrosis sputum. *Proc Natl Acad*
Sci U S A **112**, 4110–4115 (2015).

36. European Committee on Antimicrobial Susceptibility Testing (EUCAST). *EUCAST*
Reading Guide for Broth Microdilution - Version 5.0. (2024).

37. Pankratz, D. *et al.* An expanded CRISPR–Cas9-assisted recombineering toolkit for
engineering genetically intractable Pseudomonas aeruginosa isolates. *Nat Protoc*
18, 3253–3288 (2023).

38. Choi, K. H. & Schweizer, H. P. mini-Tn7 insertion in bacteria with single attTn7
sites: Example Pseudomonas aeruginosa. *Nat Protoc* **1**, 153–161 (2006).

39. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

40. Kopylova, E., Noé, L. & Touzet, H. SortMeRNA: Fast and accurate filtering of
ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**, 3211–3217
(2012).

41. Anders, S. *et al.* Count-based differential expression analysis of RNA sequencing
data using R and Bioconductor. *Nat Protoc* **8**, 1765–1786 (2013).

42. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

43. Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT-
PCR. *Nucleic Acids Res* **29**, e45 (2001).

44. Bartell, J. A. *et al.* Evolutionary highways to persistent bacterial infection. *Nat
Commun* **10**, 629 (2019).

45. Muñoz-Cazalla, A., Martínez, J. L. & Laborda, P. Crosstalk between Pseudomonas
aeruginosa antibiotic resistance and virulence mediated by phenylethylamine.
Microb Biotechnol **16**, 1492–1504 (2023).

46. Herrmann, C., Lingner, M., Herrmann, S., Brockhausen, I. & Tümmler, B. Mucin
adhesion of serial cystic fibrosis airways Pseudomonas aeruginosa isolates.
Front Cell Infect Microbiol **14**, 1448104 (2024).

772