**Genetic dissection of transcription start site selection by RNA Polymerase II in**

***Saccharomyces cerevisiae***

Payal Arora[1], Kieran F. Brennan[1], Marcie H. Warner[1], Craig D. Kaplan[1]*

[1]Department of Biological Sciences, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

*Address for correspondence: Department of Biological Sciences, University of Pittsburgh,

Pittsburgh, PA 15260, Life Science Annex 202, Fifth and Ruskin Avenues, Pittsburgh, PA

15260. Email: craig.kaplan@pitt.edu

**ABSTRACT**

Transcription initiation by RNA Pol II is driven by a preinitiation complex (PIC) comprising Pol II

and general transcription factors (GTFs): TFIIA, TFIIB, TFIID, TFIIE, TFIIF and TFIIH. In

*Saccharomyces cerevisiae*, transcription start site (TSS) selection proceeds by a promoter

scanning mechanism where the PIC scans downstream for TSSs. To determine which factors

participate in TSS selection by promoter scanning, we designed and implemented forward

genetic selections using initiation-sensitive genetic reporters. These reporters were of two types,

either being sensitive to upstream or downstream TSS shifts, allowing detection of alleles that

affect promoter scanning in different directions. From >1000 candidates, we identified three

primary classes of mutants: existing and novel mutants within known PIC components, including

novel alleles of multiple TFIIH subunits; mutants that alter promoter scanning indirectly through

cellular conditions (GTP levels and $Mn^{2+}$ levels); and mutants in chromatin and transcription

elongation-related factors. Genome-wide analysis using TSS mapping shows that tested PIC

mutants alter TSS usage globally across promoters while tested chromatin and transcription

elongation mutants had much more limited effects. These studies expand the spectrum of

mutants able to perturb initiation by promoter scanning and suggest potential plasticity in the

TFIIH structure that may be part of the evolution of this initiation mechanism.

## INTRODUCTION

Transcription initiation in eukaryotes requires assembly of a large pre-initiation complex (PIC)

comprising RNA Polymerase II (Pol II) and general transcription factors (GTF) (CONAWAY AND

CONAWAY 1993; ORPHANIDES *et al.* 1996; ROEDER 1996; KORNBERG 2007). Together these

components, in conjunction with transcriptional activators and coactivators effect promoter DNA

opening, identification of the transcription start site (TSS), and de novo initiation of RNA

synthesis using DNA-templated NTP substrates (KAPLAN 2013; ROEDER 2019). Additionally, a

large number of other factors are recruited to promoters, including chromatin remodelers and

factors functioning in the transition from initiation to elongation (RANDO AND WINSTON 2012).

During PIC assembly, TBP and TFIIB bind to the core promoter transiently with TFIIA stabilizing

TBP-DNA interactions. Pol II is then recruited to core promoter along with GTFs TFIIF and TFIIE

stabilizing the TFIIB-TBP-DNA interactions. Thereafter, TFIIH is recruited into the PIC to drive

open complex formation (promoter melting) and subsequent transcription initiation (BAEK *et al.*

2021).

In eukaryotes, the majority of promoters specifies multiple TSSs. In most eukaryotes, a

TSS is stereotypically positioned approximately 30-31 basepairs downstream from a site of TBP

binding (a canonical TATA-box or otherwise AT-rich sequence) (LIFTON *et al.* 1978;

BREATHNACH AND CHAMBON 1981; DUDNYK *et al.* 2024). It is proposed that each TSS is

supported by an associated PIC assembly point upstream (KADONAGA 1990; SMALE AND

KADONAGA 2003; SANDELIN *et al.* 2007; KADONAGA 2012; VO NGOC *et al.* 2017a; VO NGOC *et al.*

2017b; LUSE *et al.* 2020; DUDNYK *et al.* 2024). In contrast, transcription initiation in the budding

yeast *Saccharomyces cerevisiae* occurs through a promoter scanning mechanism (GIARDINA

AND LIS 1993; KUEHNER AND BROW 2006; FAZAL *et al.* 2015; QIU *et al.* 2020). The driving factors

for this novel initiation mechanism can only be speculated upon but given the loss of most

splicing in budding yeast and its highly compact genome, scanning may allow for 5′ UTR

diversification that cannot be accomplished by alternative splicing coupled to alternative

promoter usage. Intriguingly, PIC components and their assembled structure are highly

conserved between yeast and humans (BURATOWSKI *et al.* 1988; HE *et al.* 2016; AIBARA *et al.*

2021; SCHILBACH *et al.* 2021; TOMKO *et al.* 2021). While DNA melting by the yeast PIC occurs

proximal to its site of binding as in other eukaryotes (GIARDINA AND LIS 1993), the yeast PIC

scans downstream for TSSs, which are generally located 40-120 bp or further downstream of

the TATA box, when present (STRUHL 1987; PARK *et al.* 2014; LU AND LIN 2019; LU AND LIN

2021). TFIIH, required for Pol II promoter DNA opening in all eukaryotes, is proposed to use its

DNA translocase activity to continue pumping downstream DNA into the Pol II active site to

enable promoter scanning and usage of downstream TSSs (HAMPSEY 2006; MILLER AND HAHN

2006; FAZAL *et al.* 2015; FISHBURN *et al.* 2015).

Promoter scanning identifies TSSs based on their inherent sequence characteristics

together with the contributions of initiation factor and Pol II activities. Yeast share a universal

eukaryotic preference for $Y_{-1}R_{+1}$ as TSSs initiating at a purine (R, representing A or G), have a

pyrimidine (Y, representing C or T) just upstream of the TSS, and with an additional species-

specific contribution of the -8 position and of additional bases (HUROWITZ AND BROWN 2003;

ZHANG AND DIETRICH 2005; MIURA *et al.* 2006; NAGALAKSHMI *et al.* 2008; ARRIBERE AND GILBERT

2013; PELECHANO *et al.* 2013; MALABAT *et al.* 2015; LU AND LIN 2019; QIU *et al.* 2020; LU AND LIN

2021; ZHU *et al.* 2024). Scanning in yeast was conceptualized by Kuehner and Brow (KUEHNER

AND BROW 2006) where TSSs have innate efficiencies of usage, and the alteration of initiation

factor activity can alter individual TSS usage and the distribution of TSS usage at promoters

(KUEHNER AND BROW 2006; QIU *et al.* 2020; ZHAO *et al.* 2021). Genetic and molecular

experiments have shown that TSS selection in yeast is sensitive to mutation in multiple PIC

components (*e.g.* Pol II and GTFs (PINTO *et al.* 1992a; PINTO *et al.* 1994; FAITAR *et al.* 2001;

GHAZY *et al.* 2004; FREIRE-PICOS *et al.* 2005; KHAPERSKYY *et al.* 2008; JIN AND KAPLAN 2014;

QIU *et al.* 2020; ZHAO *et al.* 2021) and PIC-associated components (*e.g.* the conserved factor

Sub1, which is the yeast homolog of PC4 (KNAUS *et al.* 1996; WU *et al.* 1999; SIKORSKI *et al.*

2011; BRABERG *et al.* 2013). Mutations in the active site of Pol II that increase Pol II catalytic

activity in vitro increase efficiency of initiation at each available TSS (KAPLAN *et al.* 2012; JIN

AND KAPLAN 2014; QIU *et al.* 2020). With each TSS being more efficient in promoting initiation,

the flux of scanning Pol II can be exhausted earlier than WT, leading to an observed upstream

shift in observed TSS usage (KUEHNER AND BROW 2006; QIU *et al.* 2020). Similarly, decrease in

Pol II catalytic activity causes reduced efficiency of individual TSSs, resulting in increases in the

flux of scanning Pol II reaching downstream positions, and an observed downstream shift in

TSS usage (KUEHNER AND BROW 2006; QIU *et al.* 2020).

In contrast to Pol II, because TFIIH is hypothesized to drive scanning through its Ssl2

translocase subunit (FAZAL *et al.* 2015; FISHBURN *et al.* 2015; TOMKO *et al.* 2017; TOMKO *et al.*

2021; ZHAO *et al.* 2021), the effects of mutations in Ssl2 on TSS distributions have been

interpreted as being due to altered TFIIH processivity (ZHAO *et al.* 2021). *ssl2* alleles that confer

downstream shifts in TSS distributions have been interpreted as increasing scanning

processivity due to a higher probability of scanning reaching downstream positions. In contrast,

*ssl2* alleles that confer upstream shifts in TSS distributions appear to do so by limiting scanning

processivity, meaning that downstream positions are reached at a lower probability (KUEHNER

AND BROW 2006; ZHAO *et al.* 2021). While TFIIH and Pol II are the primary activities driving TSS

selection, how GTFs contribute to either efficiency or processivity pathways or through

additional mechanisms is not known (JIN AND KAPLAN 2014; QIU *et al.* 2020).

Mutations in other GTFs like TFIIB (PINTO *et al.* 1992a; PINTO *et al.* 1994; BANGUR *et al.*

1997; PARDEE *et al.* 1998; FAITAR *et al.* 2001; JIN AND KAPLAN 2014) and TFIIF (SUN AND

HAMPSEY 1996; GHAZY *et al.* 2004; JIN AND KAPLAN 2014) also affect TSS distributions, and

among those tested, to do so genome wide  (QIU *et al.* 2020; BASNET *et al.* 2026), as do tested

Pol II and *ssl2* alleles of TFIIH (QIU *et al.* 2020; ZHAO *et al.* 2021). Mutations in TFIIB cause

downstream shifts to TSS usage in ways that appear consistent with both TFIIB and the Pol II

active site determining initiation efficiency at all TSSs (WU *et al.* 1999; FAITAR *et al.* 2001;

KOSTREWA *et al.* 2009; JIN AND KAPLAN 2014; QIU *et al.* 2020). Additionally, published mutations

in TFIIF, the GTF recruiting Pol II to the core promoter, have also been shown to shift TSSs

upstream (SUN AND HAMPSEY 1995; GHAZY *et al.* 2004; FREIRE-PICOS *et al.* 2005; JIN AND

KAPLAN 2014; QIU *et al.* 2020). Because specific alleles of TFIIF and TFIIB have been found to

mutually suppress each other (GHAZY *et al.* 2004; FREIRE-PICOS *et al.* 2005) and also additively

interact with Pol II alleles (JIN AND KAPLAN 2014; QIU *et al.* 2020), we have proposed that at

least some TFIIF mutants affect TSS selection by functioning alongside the Pol II active site in

promoting initiation efficiency (GHAZY *et al.* 2004; FREIRE-PICOS *et al.* 2005; JIN AND KAPLAN

2014). The interface between Pol II and TFIIH is also important for scanning and has been

proposed to be a key factor in the evolution of the scanning mechanism (YANG *et al.* 2022;

YANG *et al.* 2024; BASNET *et al.* 2026). TFIIH is tethered to the PIC via the RING finger domain

of Tfb3 (TFIIK kinase subdomain of TFIIH) through interactions with TFIIE and the Rpb7 subunit

of Pol II (SCHILBACH *et al.* 2017; YANG *et al.* 2022; YANG *et al.* 2024). In yeast, mutations in both

Rpb7 (BRABERG *et al.* 2013) and Tfb3 (YANG *et al.* 2024) can confer upstream shifts to TSS

usage *in vivo*, consistent with a defective scanning process (BASNET *et al.* 2026). Factors

outside of PIC such as chromatin and elongation factors have also been implicated to affect

promoter choice and potentially TSS selection.

PIC factors (Pol II and GTFs) and PIC-associated factors like Sub1 have been

genetically identified as contributing to TSS-selection during promoter scanning. What is not

known is how many additional factors might also contribute to the process of TSS selection. Are

there additional factors that cooperate or contribute to TSS selection but have not been

structurally located in the PIC akin to Sub1? What are the surfaces and networks within the PIC

that contribute to TSS selection by promoter scanning? Finally, promoter chromatin structure

can easily be imagined to affect PIC function and TSS distribution profiles because most yeast

promoters have TSSs that map within boundaries of the +1 nucleosome (RANDO AND WINSTON

2012). We have aimed to take an unbiased approach to identify any additional factors that

participate in TSS selection during initiation by promoter scanning using forward genetic

selections (BROW 2019; WANG *et al.* 2025). Through the use of existing (BERROTERAN *et al.*

1994) and novel genetic reporters, we have identified ~300 mutations in proteins within and

outside the PIC affecting TSS selection, including a number of alleles of Pol II

elongation/chromatin genes, *eg. SPT2, SPT4, SPT5, SPT6, SPT16, SPN1, BUR2*, and

histones. We then examined a subset of the novel alleles in genetic interaction studies to probe

relationships between mutant alleles. Finally, we examined the genome-wide effects of selected

PIC and chromatin/elongation-class alleles for effects on TSS-selection. We have found that

tested GTF alleles alter TSS usage genome wide as predicted from effects on promoter

scanning in general. In contrast, chromatin or Pol II elongation-class alleles did not show similar

widespread effects. We interpret this class of mutant as acting on TSSs in a promoter-selective

fashion.

**MATERIALS AND METHODS**

**Yeast strains and oligos**

Yeast strains and oligos used in this study are listed in **Table S2**.

**Yeast media**

Yeast media used in the current studies have been prepared as previously described in

(AMBERG *et al.* 2005; KAPLAN *et al.* 2012; JIN AND KAPLAN 2014; MALIK *et al.* 2017; ZHAO *et al.*

2021). Briefly, all solid media contain 2% w/v agar (BD); YPD is 1% w/v yeast extract (Gibco™

Bacto™), 2% w/v peptone (Gibco™ Bacto™) and 2% w/v glucose (VWR) supplemented with

0.15 mM adenine (Sigma) and 0.4 mM tryptophan (Sigma) for solid medium (not supplemented

when liquid); YPD+3% formamide contains 3% v/v formamide (Ambion); YP Raffinose (YPRaf)

and YP Raffinose Galactose (YPRafGal) contain 2% w/v raffinose (Amresco) (YPRaf) or 1% w/v

galactose (Sigma) and 2% w/v raffinose (YPRafGal). Both YPRaf and YPRafGal are additionally

supplemented with 1 µg/ml of (Sigma) Antimycin A to prevent aerobic respiration. Synthetic

minimal media (SC-) were what is referred to as "Hopkins mix" with minor alterations as

previously noted (AMBERG *et al.* 2005; KAPLAN *et al.* 2012) (amino acid dropout mix, 0.17% w/v

yeast nitrogen base w/o amino acids and 0.5% w/v ammonium sulphate (BD), 2% glucose), with

the appropriate dropouts. Mycophenolic acid (MPA) (ThermoScientific chemicals) containing SC

media were supplemented at 20 µg/ml. Other drugs in the respective media were: G418

(GoldBio) and clonNAT (GoldBio) at 200 µg/ml, hygromycin (GoldBio) at 300 µg/ml and

canavanine (Sigma) at 60 µg/ml. Synthetic minimal media containing G418, clonNAT or

hygromycin are prepared as above with 0.1% w/v of monosodium glutamate (MSG) replacing

ammonium sulphate. Drug 3-aminotriazole (3AT) (Sigma) a competitive inhibitor of His3p was

used in some SC-His media at varying concentrations (later noted), for setting the threshold for

a visible His$^+$ phenotype.

**Genetic selections**

Each of the genetic selection strains possessing initiation reporters was grown overnight in

liquid YPD. Cultures were then washed twice in sterile distilled water to remove residual amino

acids before plating on respective selection media. Selection media for the reporters were SC-

His + 2mM 3AT for *imd2Δ::HIS3* and SC-His + 0.1mM 3AT for *cyc1-1019::HIS3*, *adhΔ50::HIS3*

and *hmo1Δ38::HIS3*. For dual selection of His$^+$ Lys$^+$ candidates for downstream reporters,

conditions used were SC-His-Lys +5mM 3AT for *imd2Δ::HIS3* SC-His-Lys +0.1mM 3AT, Cells

were spread using sterile 3mm glass beads at $10^8$ cells per plate. Multiple colonies of each

parent strain were inoculated for genetic selection to increase chances of independent

candidates. Plates were then kept in a lightly sealed plastic bag to prevent drying out and

incubated at 30˚C for ~7 days until suppressor colonies appeared.

**Single spot phenotyping**

Suppressor candidates were picked from selection plates and spotted onto solid YPD petri plate

in 8x6 grid (48-well format) containing 2 parents and 46 candidates apiece for the purpose of

candidate rescreening and additional phenotyping. Briefly, after growth on YPD, cells were

transferred into 100 µl of sterile distilled water in a 96-well tray using a 48-pin replicator,

followed by a second dilution of 6 µl of this cell suspension into 150µl of water. A sterilized 48-

pin replicator was then used to spot from the 150µl cell suspension onto phenotyping media.

Phenotype scoring timing was day 2 for YPD 30˚C and YPD 37˚C, day 3 for YPD+3%

formamide and synthetic complete medium (SC), day 4 for YPRaf, day 5 for SC-Lys and YPD at

16˚C and day 6 for YPRafGal. Reporter phenotypes on SC-His +/- 3AT were scored on days 3

through 5 depending on strength and SC-His-Lys +/- 3AT on day 5. **Scoring:** Scoring was

performed as in (JIN AND KAPLAN 2014). Single phenotype spots were given scores from 0 to 5,

where 0 means no growth and 5 is WT parent growth on complete medium. For conditional

phenotypes where growth is indicative of mutant phenotype (i.e. Spt$^-$ phenotype or His$^+$

phenotypes in reporter strains), scoring was 0 for no growth (WT) with positive numbers

indicating increasing growth reflecting strength of mutant phenotype. Relative scores of mutants

were first calculated per condition by Score$_{mutant}$-Score$_{WT}$ (Relative score$_{mutant: condition}$). These

relative scores were then corrected to respective control growth condition for estimating mutant

score corrected to control condition (e.g.: Relative score$_{mutant: YPRafGal}$ - Relative score$_{mutant: YPRaf}$).

These would assign positive values to mutants where growth is better than WT, negative to

those where growth is worse in an any condition and a 0 to WT.

**Mutation validation by re-introduction:**

Mutations were reintroduced into a clean parental strain possessing relevant initiation reporter

to validate effects of candidate mutations on reporter expression. Mutations were amplified from

suppressor strains and fused with a drug resistance marker by PCR sewing to generate a

*mutation::drug resistance marker* cassette and reintroduced into parent strains (JANKE *et al.*

2004). The marker used was *kanMX6* containing an insulated resistance cassette (pFA6a-

KanMX6-ins2, Addgene 195039) to minimize potential effects of marker expression on adjacent

loci (POWERS *et al.* 2022). Multiple independent transformants were phenotyped and genotyped

to assess correlation of reporter suppression and introduction of the mutant genotype along with

the resistance marker. The marker was also used to tag mutation loci to enable synthetic

genetic array analysis and selection of mutant loci through genetic crosses. CRISPR based

mutagenesis was performed to generate *rpb10* and *tfg1* mutations in the genome as described

in (LAUGHERY *et al.* 2015).

**Diploid generation**

Complementation assays and genetic interaction studies were performed with diploid

generation. Diploids were generated by mating suppressor candidates against parent or mutant

strains of opposite mating type, followed by double selection of individual haploid markers for

selecting diploids. Briefly, candidate suppressors, pre-integrated with auxotrophic markers

(*LEU2* for *MATalpha* and *TRP1* for *MATa*) were mated with the opposite mating type of the

required strains with a second auxotrophic selection marker; or appropriate drug resistance

markers were used for diploid selection for genetic interactions. Mating was performed by

transferring solid YPD-grown candidates to pre-aliquoted 100 µl-per-well of saturated culture of

test strain per well in a 96-well plate using a 48-pin (8x6) replicator and spotting onto solid YPD

using the same replicator. Post incubation for 2 days at 30˚C on plates, cells were transferred

into 100µl of pre-aliquoted sterile distilled water with the 48-pin replicator and resuspended by

shaking of replicator followed by transfer onto a double-selection media. These were incubated

at 30˚C to select for diploids for 2 days.

**Double mutant generation by synthetic genetic array**

Double mutants were generated from diploids of "tagged" single mutants mated in high

throughput following protocol from the Boone lab

(https://boonelab.ccbr.utoronto.ca/pdf/SGA_protocol_final.pdf). Briefly, a panel of *MATα single*

*mutation 1::drug resistance marker 1* strains was mated to a query *mutation 2::drug resistance*

*marker 2 MATa* strain possessing a *MATa* marker *can1Δ::STE2pr-URA3,* and diploids were

selected by double drug resistance as mentioned. After two days of incubation on double drug

diploids were transferred from solid medium directly onto solid enriched sporulation media (1%

potassium acetate enriched with auxotrophic amino acids in 2% agar) for 5 days at 27˚C. After

sporulation, sporulated spots were transferred into 50 μl sterile distilled water in 96 well plate

using a pinner, resuspended and spotted onto haploid selection media (SC-Ura-

Arg+MSG+canavanine), single mutant selection media (SC-Ura-Arg+MSG+canavanine + drug

1 or drug 2) and double mutant (SC-Ura-Arg+MSG+canavanine+drug 1+drug 2) selection media

in parallel instead of the traditional approach where mutants (single or double) are selected after

*MATa* meiotic progeny selection in tandem. We have observed that sicker haploids are lost if

selections are performed in tandem.

**Genomic DNA sequencing**

Candidate mutations were identified via AmpliSeq-based custom amplicon sequencing or whole

genome sequencing (WANG *et al.* 2025). Genomic DNA was isolated from candidates using

PureLink™ Pro 96 Genomic DNA Purification Kit and YeaStar genomic DNA kit from Zymo

research for AmpliSeq based sequencing. The Plasmid miniprep kit from Zymo research,

customized with bead lysis was used for genomic DNA isolation for whole genome sequencing.

AmpliSeq libraries were prepared and sequenced by the Texas A&M TIGGS genomics core

facility. AmpliSeq libraries were prepared following AmpliSeq for Illumina protocol for a custom

oligo pool comprising 1917 encompassing 310 genes. These genes comprise ORFs of

transcription machinery and transcription-associated genes. Libraries were sequenced in

batches of 96 samples using full Illumina MiSeq of 15 million reads by 250bpx2 run to aim for an

average coverage of ~80X per amplicon. Whole genome sequencing of samples was performed

by Seqcoast or Seqcenter. Each genomic DNA sample was sequenced at a depth of 400Mbp

yielding an average coverage of ~30X.

**AmpliSeq data processing**

Paired-end sequencing fastq files were first trimmed for adaptors at the 3′ end, followed by 3′

and 5′ oligo trimming (amplification oligos do not contribute towards mutation detection) by

cutadapt v2.1 (MARTIN 2011). These were then processed by MutantHuntWGS v1.1 (ELLISON *et*

*al.* 2020), a container-based package deploying predefined tools for alignments and variant

calling. Briefly, oligo and adaptor-free paired-end fastq files were aligned to *Saccharomyces*

*cerevisiae* S288C reference genome R64-2-1 by bowtie2 v2.2.9 (LANGMEAD AND SALZBERG

2012) resulting in bam alignment files followed by bam sorting and indexing by SAMtools v1.3.1

(LI *et al.* 2009; DANECEK *et al.* 2021). Genotype likelihoods were calculated followed by variant

calling by BCFtools v1.3.1 (DANECEK *et al.* 2021) using haploid ploidy file provided by

MutantHuntWGS v1.1 (ELLISON *et al.* 2020) (ELLISON *et al.* 2020). This resulted in vcf files for all

sequenced candidates. Mutant variants in mutant vcf files were differentially called relative to

parent vcf files by VCFtools v 0.1.14 (DANECEK *et al.* 2011). Resulting differential mutants were

annotated using SNPeff v4.3p (CINGOLANI *et al.* 2012) and SIFT4G (VASER *et al.* 2016), which

uses EF4.74 *Saccharomyces cerevisiae* library for annotating severity of amino acid change.

For WGS, fastq reads were analyzed by Seqcenter and Seqcoast for mutation detection by

breseq (DEATHERAGE AND BARRICK 2014). *Saccharomyces cerevisiae* S288C reference genome

R64-1-1 was used for annotation which generated a list of variants. Variants specific to

suppressor candidate were curated by a custom shell script comparing mutant variants to

parent variants.

**STRIPE-seq**

Total RNA was isolated from 50 mL of mid-log yeast culture with density of $1\times10^7$ cells/mL as

described previously (SCHMITT *et al.* 1990) and resuspended in 90 µL nuclease free water. RNA

samples were in-column DNase I treated using the RNeasy mini kit from Qiagen with 100 µg

RNA input per column. DNased RNAs (10 µg per sample) were depleted of rRNA using

bioltinylated riboPOOL kit for *Saccharomyces cerevisiae* from siTOOLS Biotech, purified by

Dynabeads MyOne Streptavidin C1 and eluted in 9 µL nuclease free water. STRIPE-seq

libraries were then generated from rRNA depleted RNAs using 200 ng RNA per sample as

previously published (POLICASTRO *et al.* 2020) with our modifications. Briefly, 200 ng of each

RNA sample, split into two 100 ng reactions were terminator exonuclease (TEX) treated to

remove uncapped RNA. Pooled, TEXed reactions were then reverse transcribed using a

random 10-mer with partial homology to P7 adaptor RTO

(GCCTTGGCACCCGAGAATTCCANNNNNNNNNN) instead of a complete P7:oligo-random

pentamer (modification; P7 adaptor added during final library amplification). Non-templated

cytosines at the 3′ end of cDNA then allow annealing of a template switching oligo, containing

partial P5 adaptor, 9-base UMI (instead of 8-base UMI) and a linker (TATAGGG) (Biotin-

GTTCAGAGTTCTACAGTCCGACGATCNNNNNNNNNNTATArGrGrG). After template switching,

libraries were amplified by dual indexing primers with complete P5 and P7 adpaters. Libraries

(>200-800 bp) were extracted from 3% polyacrylamide TBE gel to remove oligos and then

sequenced by Illumina NextSeq 2000 system with 1X50 bp sequencing cycle. Sequencing

reads were delivered demultiplexed.

**STRIPE-seq analysis**

STRIPE-seq fastq files were processed as TSS-seq reads we had previously published (ZHAO

*et al.* 2021) with STAR alignment and additional bam processing followed previously published

of goSTRIPES and TSRexploreR packages (POLICASTRO *et al.* 2020; POLICASTRO *et al.* 2021;

POLICASTRO AND ZENTNER 2022). Fastq files were filtered for quality with fastx-toolkit v 0.0.14 (-

q 20 -p 75) (http://hannonlab.cshl.edu/fastx_toolkit/), followed by trimming of adaptors by

trimmomatic v 0.38 (BOLGER *et al.* 2014) and then 9 base UMI extraction by umi-tools v 1.0.0

(SMITH *et al.* 2017). Reads were then trimmed of linker TATAGGG and then trimmed to 25

bases by cutadapt v 2.10 (MARTIN 2011) before local alignment by STAR v 2.7.9a (DOBIN *et al.*

2013) generating bam files. Bam files with uniquely mapping reads were then filtered of

unfavorably flagged reads by samtools v 1.9 (samtools view –F 3844) (LI *et al.* 2009; DANECEK

*et al.* 2021) followed by deduplication by umi-tools v 1.0.0 (SMITH *et al.* 2017). Deduplicated and

indexed bam files were then processed by the protocol followed by the Zentner lab (POLICASTRO

*et al.* 2020; POLICASTRO *et al.* 2021; POLICASTRO AND ZENTNER 2022). Briefly, reads with more

than 3 soft-clipped 5′ bases are listed and filtered out by picard v 2.18.12 (Picard Toolkit 2019)

following GoSTRIPES (Robert Policastro). Indexed filtered bams were then processed by

TSRexploreR (POLICASTRO *et al.* 2021) for G correction to generate bedgraphs for 5′ ends of

reads (TSSs) that we processed to split into + and – strand bedgraphs and downstream

analyses following our TSS-seq pipeline (ZHAO *et al.* 2021). Pearson correlation matrices were

generated for bedgraphs of all libraries using our in-house Rscript. Promoter matrices

representing TSSs were generated comprising 401 base window (250 bases upstream and 150

bases downstream from median TSS position including the median) across 5979 annotated

promoters (RHEE AND PUGH 2012). These matrices were utilized for estimating median and

spread (base distance between $10^{th}$ and $90^{th}$ percentile) of TSS distributions across the 5979

promoter and eventually Δmedian and Δspread of mutants relative to WT for differences (ZHAO

*et al.* 2021). Promoter matrices were row normalized before generating average matrices across

replicates and estimating average median and average spread values for calculating average

Δmedian and Δspread values relative to average WT. Top 2829 promoters with expression

cutoff of 10 were selected based on library normalized average of all WTs sorted by expression:

sum of TSSs per 401 base promoter window.

**RESULTS**

**TSS-shifting mutants identified by transcription initiation genetic reporters**

To understand the genetic requirements for initiation by promoter scanning, we established

genetic selection systems for the isolation of transcription start site-shifting mutants. We

deployed four initiation reporter alleles designed or chosen because expression from the

reporter is sensitive to altered TSS selection: *imd2Δ::HIS3, cyc1-1019::HIS3, adhΔ50::HIS3,*

and *hmo1Δ38::HIS3* (**Fig. 1**). These reporters comprise native (*imd2Δ::HIS3*) (MALIK *et al.* 2017)

or engineered (*cyc1-1019::HIS3*, *adhΔ50::HIS3* and *hmo1Δ38::HIS3*) (FURTER-GRAVES AND

HALL 1990; PINTO *et al.* 1992a; PINTO *et al.* 1992b) promoters that have each been fused to the

*HIS3* ORF. In wild type cells, they generally do not express sufficient levels of *HIS3* gene

product to confer growth on medium lacking histidine (the His⁻ phenotype). In the presence of a

TSS selection mutant conferring an upstream or downstream shift appropriate to the reporter

configuration, a shift in TSS usage leads to expression of a functional *HIS3* mRNA, allowing

cells to grow on medium lacking histidine (the His⁺ phenotype) (**Fig. 1a**). We tuned our

phenotypic selections through the use of the drug 3-aminotriazole (3AT), a competitive inhibitor

of His3p (KLOPOTOWSKI AND WIATER 1965; STRUHL AND DAVIS 1977), allowing a shift in the

threshold of reporter activity needed to confer the His⁺ phenotype and beneficial for reporters

that allowed leaky *HIS3* expression in WT cells (**Fig. 1b**). We reasoned that using two different

promoters for each class of TSS-shift reporter would allow us to prioritize mutants more likely to

affect promoter scanning globally as opposed to reporter-specific alleles. We previously

developed and characterized the *imd2Δ::HIS3* reporter where *HIS3* replaces the *IMD2* ORF and

is under control of the *IMD2* promoter (MALIK *et al.* 2017). Downstream shifts to TSS usage

bypass a terminator element in the *IMD2* promoter and allow expression of the downstream

ORF (KOPCEWICZ *et al.* 2007; JENKS *et al.* 2008; KUEHNER AND BROW 2008). We found this

reporter exquisitely sensitive to defects in Pol II and GTFs leading to downstream TSS shifts

(MALIK *et al.* 2017) and also that we could identify alleles of *SSL2* (ZHAO *et al.* 2021) or *TFA1*,

*TFB3*, or *RPB7* (BASNET *et al.* 2026) able to suppress this reporter through genetic screens or

site-directed mutagenesis. Therefore, this reporter appears sensitive to mutations in multiple

PIC components that might each alter promoter scanning in distinct ways, *i.e.* controlling Pol II

initiation efficiency or controlling scanning processivity.

Our other downstream shift-detecting reporter *cyc1-1019::HIS3* functions analogously to

*imd2Δ::HIS3* (NA *et al.* 1992; PINTO *et al.* 1992a; PINTO *et al.* 1992b). Here, downstream shifts in

TSS usage allow bypass of an introduced upstream ATG (the *cyc1-1019* allele). The introduced

ATG allows translation initiation out of frame with *HIS3* and confers a His⁻ phenotype. Shift to

use of downstream TSSs allows translation to begin at the normal *HIS3* ATG. A variant of this

reporter configuration (*cyc1-5000*) was used to identify the first *SUA7* (TFIIB) alleles (PINTO *et*

*al.* 1992a; PINTO *et al.* 1992b) as well as alleles of *RPB1* (*SUA8*) (BERROTERAN *et al.* 1994).

This reporter is less leaky than *imd2Δ::HIS3* and thus more tightly His⁻ even in the absence of 3-

AT (**Fig. 1b**). We demonstrated that strong downstream shifting mutants *rpb1* H1085Y and

*sub1Δ* were His⁺ using this reporter (**Fig. 1b**).

Our upstream shifting reporters function analogously to each other. In each case, a

reduced TSS to ATG distance likely hampers translation initiation of *HIS3* because the *HIS3*

ATG is too close to the 5′ end of the mRNA (FURTER-GRAVES AND HALL 1990). 5′ extension of

the mRNA through usage of an upstream TSS allows more efficient *HIS3* translation initiation.

We employed *adh50Δ::HIS3*, previously shown to detect *rpb9* alleles as upstream shifters

(FURTER-GRAVES AND HALL 1990; FURTER-GRAVES *et al.* 1991), and a second, novel upstream

reporter. To create an additional upstream shift-detecting reporter, we analyzed our previous

genome wide TSS data (QIU *et al.* 2020) for promoters that showed especially large shifts in

TSS usage in response to upstream TSS mutants, settling on *HMO1*. We then fused different

lengths of the *HMO1* promoter, truncated by 38, 45 and 53 basepairs to *HIS3* to ask if any of

these configurations could cleanly detect upstream TSS-shifting mutants (**Fig. S1a**). First, we

tested if these variants could function effectively as potential initiation reporters by assessing if

they showed suppressibility when exposed to UV mutagenesis (**Fig. S1a**), an approach

previously tested for reporter suppressibility (HAMPSEY 1991). After observing reporter

suppressibility, we tested them in conjunction with TSS-shifting Pol II and *rpb9* mutants (**Fig.**

**1b**).

**Identification of novel upstream and downstream TSS-shifting mutants**

For our selections, we integrated our engineered reporters (*cyc1-1019::HIS3*, *adhΔ50::HIS3* and

*hmo1Δ38::HIS3*) at a gene-free locus on chromosome I (**Fig. S1b**) (GRUBER *et al.* 2012) with

*imd2Δ::HIS3* present at the native *IMD2* locus. We isolated suppressors of initiation reporters by

genetic selection of His⁺ candidates and did so in two batches with two versions of genetic

selection strains. In an initial pilot genetic selection, ~700 candidates were isolated across the

four first generation strains carrying the reporters described above, with the majority isolated

from the *IMD2* reporter. Candidates were re-phenotyped for their His$^+$ phenotypes and for

secondary conditional or transcription-related phenotypes. These included growth on YPD at

30 °C, 16 °C, 37 °C and with 3% formamide, SC supplemented with MPA to detect MPA

sensitivity commonly observed for known upstream shifting TSS alleles (ZHAO *et al.* 2021), SC-

Lys medium to test for suppression of the Spt$^-$ reporter allele *lys2-128∂* (SIMCHEN *et al.* 1984;

CUI *et al.* 2016), and YPRafGal with YPRaf as a control for suppression of *gal10Δ56* (GREGER

AND PROUDFOOT 1998; GREGER *et al.* 2000; KAPLAN *et al.* 2005), a reporter allele able to detect

different kinds of transcription-associated phenotypes. We utilized profiles of phenotypes to

potentially cluster mutants by similarity of their additional phenotypes and to choose candidates

from distinct phenotypic groups for further study. Additionally, known phenotypes of existing

mutants allowed provisional attribution of candidates to possible candidate genes. We chose

168 of these candidates for sequencing using a custom Illumina AmpliSeq amplicon pool (WANG

*et al.* 2025). Of 168 candidates sequenced using Illumina AmpliSeq, we identified a potential

causative mutation in 72 of them. For genes already linked to TSS selection, we identified

alleles of *SUA7* (2 unique alleles), *RPB1* (8 unique alleles), *RPB2* (8 alleles from 11

candidates), *RPB9* (2 unique alleles), and *SUB1* (3 alleles from 4 candidates). Because of

inconsistent coverage with AmpliSeq, we piloted whole genome sequencing for 10 of the initial

candidates where AmpliSeq failed to identify a candidate mutation. Whole genome sequencing

(WGS) identified potential causative mutations for 7/10 candidates sequenced and provided

information allowing refinement of our selection procedures as described below.

This preliminary analysis allowed us to refine our selections for a second round where we

identified additional both upstream- and downstream-shifting candidates (**Figs. 2, 3, 4, 5**). We

identified transcription-related mutants (subset displayed in **Figs. 2, 3, 4, S2, S3**) but also

identified classes of mutant that appear reporter specific (*e.g.* guanine metabolism, **Fig. 5a**) and

a likely indirect modulator of TSS selection (e.g. *PMR1*, **Fig. 5b**), classes that AmpliSeq would

have failed to identify. Candidates mutations identified by sequencing are compiled in **Table S1.**

Previously identified TSS shifting alleles are compiled in **Table S3**. We then incorporated

secondary screening to filter these latter classes out as discussed in the next section. Among

upstream-shifting candidates, we filtered out recessive loss of function mutants in *RPB9* by a

complementation assay. In a subsequent third round genetic selections we isolated ~1750 more

candidates comprising 447 suppressors of *imd2Δ::HIS3*, 672 of *cyc1-1019::HIS3* (**Fig. 2a**), 293

of *adhΔ50::HIS3,* and 350 of *hmo1Δ38::HIS3* (**Fig. 3a**) and after phenotyping-based filtering we

prioritized ~250 candidates for whole genome sequencing. In a separate selection of

*imd2Δ::HIS3* suppressors performed by the students in the BIOSC 0068 Foundations of Biology

Laboratory 2 course at the University of Pittsburgh, 48 His$^+$ candidates were prioritized by

phenotypic and genotypic screens for whole genome sequencing, yielding 46 candidates. These

were combined with the above 447 *imd2Δ::HIS3* suppressors amounting to 495 *imd2Δ::HIS3*

suppressors (**Fig. 2a**). Prioritization across all reporter classes resulted in a total of ~300

candidates sequenced by whole genome sequencing. Out of the ~300 candidates sequenced,

we identified reporter transcription-associated mutations in ~240 of these and further ~160

unique mutants.

Of all transcription-related mutants isolated in this study, we observed three primary

groups: PIC mutants, indirect effectors of TSS selection (*e.g. PMR1*) or promoter specific non-

transcription factors *e.g.* guanine mutants), and chromatin/Pol II elongation factors. We

validated 8 candidates through reconstruction of mutants by generating alleles in a clean

genetic background and all 8 recapitulated candidate suppression (**Fig. 2b**). In addition to direct

validation by reconstruction of a subset of mutants, several lines of evidence support the

mutations identified by WGS are causative for phenotypes. First, we selected spontaneous

suppressors, so the expectation was for most strains to have essentially a single causative

mutation. Second, we identified previously described alleles of *SUA7, RPB1, RPB2, TFB3,*

*SPT6*, and *TFA1* while not observing changes elsewhere in their genomes. Third, we identified

multiple inactivating alleles of *SUB1* and *RPB9* as expected for non-essential genes with known

effects on TSS selection, while not observing changes elsewhere in their genomes. Fourth,

there were five cases where we identified the identical allele independently in separate

selections. Finally, nearly all candidate sequencing data showed only a single change in a gene

that could rationally be linked to transcription initiation. Because previous genetic interaction

studies have helped annotate functions to mutations (JIN AND KAPLAN 2014; ZHAO *et al.* 2021),

we tested inter-subunit interface mutants and genetic interactions across a subset of our

mutants before assessing TSS shifts caused by our novel mutants genome wide, which we

discuss later.

Among mutants in PIC components, we identified alleles of Pol II (*RPO21/RPB1*, *RPB2*,

*RPB9*, *RPB10*) and of GTFs (TFIIB, *TFA1*/TFIIE, *TFG2*/TFIIF, TFIIH: *SSL2*, *SSL1*, *TFB2*, *TFB3*,

*TFB4* and *RAD3*) (**Fig. 4**). We first observed a high prevalence of mutations in the downstream

shifting class in *RPB1* and *RPB2* lining the cleft leading to and surrounding the active site.

Because *RPB1* and *RPB2* are larger genes, these mutants predominated in our selections,

though many of these mutations were novel (**Figs. 4a, b**). Because downstream shifting is

expected from reduced Pol II activity (KAPLAN *et al.* 2012; MALIK *et al.* 2017), catalytic or

structural defects that lead to reduced catalytic activity or PIC function are expected to be

isolated as downstream shifters. However, not all *rpb* mutants clustered around the active site

(**Figs. 4a, b**). We observed two types of mutations amongst those surrounding the activity cleft.

We found that in both Rpo21/Rpb1 and Rpb2, mutations within structural domains surrounding

the cleft displayed formamide sensitivity (**Fig. 4d**), a phenotype that could reflect protein folding

or stability defects. Mutations facing or on the surface of the cleft generally were formamide

resistant. We also isolated mutations at intra-subunit interfaces including those linked to control

Pol II catalysis. For example, we isolated an upstream-shifting mutation G707V in the Rpb1

anchor loop region (**Fig. 4e**). This loop lies between α20 and α21 helices of Rpb1 (KASTER *et al.*

2016) and a poly-alanine substitution at Rpb1 707-709 in the anchor loop was previously shown

to confer MPA sensitivity, consistent with an upstream shift in TSS usage (KASTER *et al.* 2016).

We additionally observed mutations in the Rpb1 α21 helix, A725T and E734D, both resulting in

downstream shifts. A prior mutation in Rpb1 α21, G730D, was identified as a reduced catalytic

activity mutant and appeared to function through incompatibility with Rpb9 in that slow growth of

G730D was suppressed by *rpb9Δ* (KASTER *et al.* 2016). It is possible that these substitutions

work similarly though they are predicted to be more subtle. Alternatively, E734 is positioned to

potentially interact with a network of residues that connects to an Rpb1-Rpb8 interaction but is

also adjacent to TFIIS when it is inserted into the active site. However, no role for TFIIS in TSS

selection has yet been described. Regardless, perturbation to E734 could result in reduced Pol

II catalysis, leading to a downstream shift in TSS usage.

Of *rpb2* mutations, beyond the active cleft, we observed mutations in Pol II-lobe-jaw

domain along the path of DNA entry near the Rpb9-TFIIF interface (HEMMING *et al.* 2000) (**Fig.**

**4f**). Because previous studies have shown that perturbation of both TFIIF and Rpb9 functions

results in upstream shifts (FURTER-GRAVES *et al.* 1991; HULL *et al.* 1995; SUN AND HAMPSEY

1995; SUN *et al.* 1996; KHAPERSKYY *et al.* 2008; EICHNER *et al.* 2010), the upstream-sensitive

reporter suppressor Rpb2 C317R could be consistent with disruption of this interface. Here,

Rpb2 C317 is buried, and substitution of an arginine might be predicted to perturb structure and

alter positioning of either Rpb9 or TFIIF or both. Disruption of Tfg1 at this interface

encompassing residue R422 (homologous to R417 in the studied *S. mikate* Tfg1) (EICHNER *et*

*al.* 2010) and deletion of *RPB9* (WOYCHIK *et al.* 1991) each result in a cold sensitive phenotype,

a phenotype shared by Rpb2 C317R (**Fig. 3b**). In contrast to Rpb2 C317R, a nearby

substitution E319D results in the opposite phenotype – suppression of a downstream-shift

sensitive reporter. While Rpb2 E319 is close to Tfg1, it appears to make direct interactions with

Rpb9 Y15 and R30 and perturbing the length of the 319 sidechain in E319D may alter this

interaction with Rpb9 (**Fig. 4e**).

We also isolated Rpb2 G369C and G369R mutants as upstream shifters (**Fig. 4f**).

Substitutions in Rpb2 G369 were previously isolated as Spt⁻ (HEKMATPANAH AND YOUNG 1991),

as a suppressor of a TFIIB downstream-shifting allele (CHEN AND HAMPSEY 2004), and to be

MPA$^S$ with altered termination at a genetic reporter allele involving *ADH2* (KUBICEK *et al.* 2013).

Bulky substitutions for glycine at this residue would be in position to interfere with the Rpb1-Tfg1

interface (CHEN *et al.* 2007; SCHILBACH *et al.* 2021). Previous studies show that mutation in the

TFIIF dimerization domains within both Tfg1 and Tfg2 subunits also resulted in upstream TSS

shifts (SUN AND HAMPSEY 1995; GHAZY *et al.* 2004; EICHNER *et al.* 2010). Intriguingly, we also

isolated a downstream shifting mutation in the Pol II subunit Rpb10, a subunit shared amongst

the three eukaryotic nuclear RNA polymerases. This allele is discussed below where we probe

how this substitution may be working (**Fig. 6**).

Among GTFs, we identified novel alleles of *TFG2* (TFIIF), *SUA7* (TFIIB)*, RAD3, SSL1,*

*SSL2, TFB2*, *TFB4* and *TFB3* (TFIIH) and *TFA1* (TFIIE). For Tfg2, we identified three

suppressors of a downstream-shift sensitive reporter: V128G, C222Y (**Figs. 2b**, **4g**) and T366P

(**Table S1**) (with C222Y isolated twice). This is the first occurrence of *tfg2* alleles potentially

conferring downstream shifts, because previously identified mutants of Tfg2 that affected TSSs

resulted in upstream shifts (SUN AND HAMPSEY 1995; GHAZY *et al.* 2004; EICHNER *et al.* 2010).

Tfg2 V128 and C222 are close to each other in space but there is not an obvious interpretation

for their structural perturbations. Interestingly, while Tfg2 substitutions V128G and C222Y lie

within the Tfg2 dimerization domain (but not at the dimerization interface), T366P lies in the

winged helix domain (WH) and is not visualized in the yeast PIC. The WH domain has been

proposed to interact with upstream DNA (EICHNER *et al.* 2010).

In *SUA7,* we isolated novel suppressors of downstream-shift sensitive reporters in the B-

finger/B-reader domain such as R60W and G80R alleles as well as those in previously studied

residues R64K (BANGUR *et al.* 1997; PARDEE *et al.* 1998) and R78C (PINTO *et al.* 1994) (**Fig. 4h,**

**Table S1**). We note that G80R is cold sensitive just as are classical *sua7* alleles (PINTO *et al.*

1992b). Besides substitutions in the B-reader domain, we also identified R182C in the

Sua7/TFIIB B-core N-terminal cyclin fold (**Fig. S2**), the domain that interacts with the Pol II wall,

involved in directing the DNA-RNA hybrid (CRAMER *et al.* 2001; KOSTREWA *et al.* 2009). The

TFIIB B-core domain binds to the Pol II complex at the Rpb2 "wall" domain, which undergoes

reorganization as a consequence of the TFIIB-Pol II interaction. Previously isolated substitutions

in C149 at the TFIIB/Rpb2 interface resulted in a downstream TSS shift (WU *et al.* 1999). Our

new substitution, Sua7 R182C is in a residue that packs with Sua7 C149, which is adjacent to

the Rpb2 wall domain residue M868 (KOSTREWA *et al.* 2009; SAINSBURY *et al.* 2013) (**Fig. 4h**).

We also isolated Rpb2 substitutions in the wall near the N-terminal B-cyclin fold domain

proximal to M868 (Rpb2 G867S, Y931C, L961S). Unbiased isolation of TSS mutants at this

interface highlights the ability of our selections to identify interfaces involved in TSS selection.

Within TFIIH, we isolated alleles across the complex, consistent with a network of

interactions throughout its subunits able to modulate the scanning process. Within Ssl1, *ssl1*

I122T and *ssl1* R230G were isolated and are the first *ssl1* alleles directly connected to TSS

selection. *SSL1* was initially genetically isolated through a suppressor of a stem loop placed at

the 5′ UTR of *HIS4* (YOON *et al.* 1992) and potentially regulates Rad3 helicase activity involved

in NER (COIN *et al.* 1998; KIM *et al.* 2015) via its VWA family protein domain. Rad3 activity is not

implicated in TFIIH initiation functions and the location of our *ssl1* alleles suggests that they

might instead alter internal Ssl1 interactions or the organization of residues that interact with

Tfb1 or Tfb4 or both. Ssl1 I122 is directly adjacent to a loop of Tfb4 that inserts into Ssl1 while

Ssl1 R230 is positioned internally to organize an Ssl1 loop that connects to both Tfb1 and Tfb4

interfaces (**Fig. 4i**).

Further, we isolated Tfb2 F189S that lies in the HEAT repeats domain of Tfb2 as a

suppressor of a downstream shift-sensitive reporter. Tfb2 F189 contributes to hydrophobic

packing within Tfb2 that might limit dynamics or breathing of this critical connector protein for

bridging Ssl2 to the rest of TFIIH. Relaxation of this domain through loss of the bulky

phenylalanine sidechain potentially could confer flexibility on the connection of Ssl2 to the rest

of the complex (**Fig. 4k**). Similarly to how Tfb2 connects Ssl2 to the rest of TFIIH, Tfb4 is the

connector of Tfb2 to the rest of TFIIH. We isolated *tfb4* N71K as a suppressor of a downstream

shift-sensitive reporter. Tfb4 N71 lies in the VWA domain and insertion of a bulkier and charged

lysine into an internal Tfb4 pocket might also loosen the Tfb4 structure, potentially adding

flexibility to the linkages across components (**Fig. 4k**).

Tfb3 is critical connecting TFIIH by its RING domain to Pol II (through Rpb7) and the rest

of the PIC (through TFIIE). Previously tested mutations that disrupt this interface cause

upstream shifts in TSS selection (YANG *et al.* 2022; YANG *et al.* 2024; BASNET *et al.* 2026), while

removal of the Tfb3 RING domain or the TFIIK kinase module (part of TFIIH) result in usage of

TSSs ~30 bp downstream of the TATA box *in vitro*, similarly to non-scanning eukaryotes

(MURAKAMI *et al.* 2015; YANG *et al.* 2022). We have interpreted these mutants as conferring a

defect in scanning, either a loss of scanning processivity or reduction in scanning rate. Here, we

have isolated *tfb3* S23F as suppressing a downstream shift-sensitive reporter (**Fig. 4l**). Tfb3

residue S23 lies within the Tfb3 RING finger domain and is present in a loop adjacent to a

hydrophobic interface with TFIIE subunits Tfa1 and Tfa2, formed by Tfb3 L22, Tfa1 I183, Tfa1

V102, Tfa1 H103, and Tfa2 L283 (**Fig. 4l**). Intriguingly, a potentially more disruptive substitution

at the very same residue, *tfb3* S23P, was isolated as an upstream shifter along with other

disruptive substitutions at the Tfb3-TFIIE interface (YANG *et al.* 2022; YANG *et al.* 2024; BASNET

*et al.* 2026). Because it has the opposite phenotype of Tfb3 S23P, we interpret Tfb3 S23F as

stabilizing the Tfb3-TFIIE interaction (see Discussion). Supporting our interpretations of

mutations identified in targeted *TFB3* and *TFA1* genetic screens or site-directed mutational

studies (YANG *et al.* 2022; YANG *et al.* 2024; BASNET *et al.* 2026), we isolated mutations R64T

and P51L in *TFB*3 that result in suppression of upstream shift-sensitive reporters, namely Tfb3

R64T and P51L (**Fig. 3, 4I**). Tfb3 R64 had previously been identified as making a key interaction

with Rpb7 and each side of this interaction appears to function in promoting scanning as

mutations are MPA$^S$ and/or alter TSS selection (BRABERG *et al.* 2013; YANG *et al.* 2022; BASNET

*et al.* 2026). Contemporaneously with our identification of these residues in unbiased genetic

selections, P51 mutations were directly tested based on evolutionary analysis of *TFB3*, with P51

and R64 conferring strong MPA-sensitivity. Because P51 and R64 substitutions were epistatic

with each other, this result was consistent with these residues working together, *e.g.* P51

working by enabling positioning of R64 (YANG *et al.* 2024)**.**

For TFIIH, physical interactions between Rad3 and Ssl2 may also contribute to TFIIH

function in scanning (**Fig. 4j**). Rad3 K499E resulted in suppression of a downstream-shift

sensitive reporter and is directly proximal to Ssl2 (**Figs. 2b**, **4j**). We had previously identified

mutations in Ssl2 at K357R and K357E (ZHAO *et al.* 2021) near this Rad3 interface sharing the

same phenotype as Rad3 K499E, raising the possibility that could be functional interface, with

the caveat that Ssl2 K357 may act through other interactions. Because we interpret downstream

shifts for Ssl2 mutants as a gain of processivity, there are two possibilities for Rad3 K499E in

this framework. First, the interaction between Rad3 and Ssl2 might limit Ssl2 processivity, and

therefore we would interpret K499E as a loss of function for the interaction. Alternatively, the

interaction between Rad3 and Ssl2 might promote processivity, and K499E confers phenotypes

by strengthening the interaction.

We further wanted to isolate special "*ssl2*-like" alleles that showed both a His$^+$

phenotype on the downstream shift-sensitive *imd2∆::HIS3* reporter and a Lys$^+$ phenotype due to

suppression of the *lys2-128∂* Spt$^-$ phenotype reporter, a phenotype combination exhibited by a

specific subset of downstream-shifting *ssl2* mutants (ZHAO *et al.* 2021). These previously

identified alleles were highly localized to the DRD domain of Ssl2 (ZHAO *et al.* 2021). Before our

Ssl2 work, the Spt⁻ phenotype had only been observed in conjunction with upstream-shifting

phenotypes for alleles of Pol II and the TFIIF subunit Tfg2 (KAPLAN *et al.* 2012; JIN AND KAPLAN

2014). Because the Spt⁻ *ssl2* mutants appeared novel, we wanted to ask if additional GTF

mutants with similar phenotypes could be identified, therefore we performed selections with the

*imd2Δ::HIS3* reporter selecting on SC-His-Lys medium. While we identified two new alleles of

*SSL2*, each with a mutation in Helicase domain 1 (HD1, *ssl2* R378L and *ssl2* W482G), we did

not identify additional mutations of this class in other GTFs, suggesting this class of allele may

be unique to *SSL2*. Finally, we isolated a TFIIE mutation *tfa1* E374* as a putative downstream

shifting allele (**Fig. S2**). Tfa1 has been shown to have inhibitory effect on Tfa2-activated Ssl2

translocase activity (LIN AND GRALLA 2005). This suggests *tfa1* E374* could function by relieving

restriction on Ssl2 translocase activity, resulting in downstream TSS shift due to increased

processivity.

**Non-PIC mutants conferring suppression of TSS-shift sensitive reporters**

Besides PIC factors, our study also isolated non-PIC mutants, comprising putative reporter-

specific factors like guanine pathway mutants, indirect effectors like *PMR1*, and

chromatin/transcription elongation factors. We found almost fifty percent of initial *IMD2* reporter

suppressors had mutations in guanine metabolism genes (**Fig. 5a**). *IMD2* is activated under low

GTP conditions, specifically because low GTP induces a TSS switch that bypasses the

upstream terminator in the *IMD2* promoter (**Fig. 1a**). Therefore, defects in guanine metabolism

create inducing conditions for *IMD2.* We predicted this class of mutant might be differentially

sensitive to supplementation by guanine. We found that guanine supplementation suppressed

the His⁺ phenotype of guanine pathway mutants specifically, but not a subset of tested PIC

mutants (**Fig. 5a**), and therefore incorporated guanine supplementation into subsequent

phenotyping conditions as indicated above. Our WGS also identified an allele of *PMR1* as a

suppressor of the *adhΔ50::HIS3* reporter (**Table 1S**). *PMR1* encodes an ER/golgi $Ca^{2+}/Mn^{2+}$

transporter, and loss of its activity raises cytoplasmic $Mn^{2+}$ levels, conferring $Mn^{2+}$

hypersensitivity (RUDOLPH *et al.* 1989; DURR *et al.* 1998; MANDAL *et al.* 2000) (**Fig. 5b**). Our lab

had previously found that *pmr1Δ* genetically interacts with increased activity Pol II mutants

(BRABERG *et al.* 2013), that increased activity Pol II mutants are specifically sensitive to

increased $Mn^{2+}$ (CABART *et al.* 2014), and that *pmr1Δ* itself resulted in an upstream shift to TSS

usage at the *ADH1* promoter (QIU *et al.* 2016), due to potential increased Pol II activity in

response to heightened cytoplasmic (and nuclear) $Mn^{2+}$ levels. Isolating *pmr1* alleles as

suppressors of the *hmo1Δ38::HIS3* reporter would be consistent with potential global effects of

$Mn^{2+}$ levels on TSS selection. While $Mn^{2+}$ has many effects on Pol II transcription including

perturbing fidelity, it consistently stimulates RNA polymerase catalysis *in vitro e.g.* (NIYOGI *et al.*

1981; QIU *et al.* 2024). Therefore, upstream shifts at multiple promoters *in vivo* are consistent

with expectations for increased Pol II catalytic activity during initiation caused by increase in

cellular $Mn^{2+}$ levels.

Intriguingly, we isolated chromatin-related Pol II associated elongation factors as

suppressors of both downstream reporters *imd2Δ::HIS3* and *cyc1-1019::HIS3* (**Fig. S3**). Many

of these are canonical "Spt" genes and fittingly, some of these were isolated in our double

selections for His$^+$ Lys$^+$ (Spt$^-$) suppressors, but a number were isolated from His$^+$ selections. We

isolated alleles of *spt2*, *spt5*, *spt16*, *bur2, spn1* and histones from both downstream-sensitive

reporters and *spt4* and *spt6* only from the *imd2Δ::HIS3* reporter (**Fig. S3**). This reporter bias

may potentially exist because our selections were not saturating. A role for Spt6 in *IMD2* TSS

selection has been described for specific *spt6* alleles defective for interaction with a

phosphorylated linker in Rpo21/Rpb1 (CONNELL *et al.* 2022). The chromatin/elongation factors

we isolated are all known to suppress initiation from cryptic promoters within genes (WINSTON *et*

*al.* 1984; CLARK-ADAMS *et al.* 1988; KAPLAN *et al.* 2003; CHEUNG *et al.* 2008; DORIS *et al.* 2018).

Similarly, their Spt⁻ phenotypes at *lys2-128∂* are consistent with a chromatinized-promoter within

a transcription unit (in this case, a Ty1 *∂* element within the *LYS2* gene) being activated upon

elongation-mediated chromatin disruption in specific mutants (SIMCHEN *et al.* 1984; WINSTON *et*

*al.* 1984; CLARK-ADAMS *et al.* 1988; SWANSON *et al.* 1991; PRELICH AND WINSTON 1993;

FISCHBECK *et al.* 2002; KAPLAN *et al.* 2003; CHEUNG *et al.* 2008; CUI *et al.* 2016; DORIS *et al.*

2018). This would be a reasonable model to explain suppression of *imd2Δ::HIS3* because the

*IMD2* promoter (driving His⁺) is transcribed as part of an upstream CUT transcript (KOPCEWICZ

*et al.* 2007; JENKS *et al.* 2008; KUEHNER AND BROW 2008), potentially sensitizing the

downstream TSS to chromatin defects driven by Pol II elongation. Conversely, this model does

not easily explain why we also identified chromatin/elongation gene alleles as suppressors of

*cyc1-1019::HIS3* (see Discussion) and raises the possibility that chromatin factors might globally

alter TSS selection, which we test directly below.

**Functional characterization of TSS mutants through genetic interaction analyses**

After our genetic selections, we wanted to functionally characterize mutants using genetic

interactions. In our prior studies, both additive/synergistic and epistatic interactions have been

observed for TSS-shifting mutants leading to a framework where mutants are interpreted as

affecting initiation efficiency (i.e. Pol II catalysis) or scanning processivity (TFIIH/PIC functions).

Previous examination of double mutants between Pol II catalytic mutants and TFIIB or TFIIF

alleles showed additivity across combinations (enhancement when mutants shift TSSs the same

direction and suppression when they shift in opposite directions). However, in other

combinations, we observed epistasis where double mutants exhibited the TSS phenotype of

one of the single mutants and not the other. This was especially clear when upstream shifting

Pol II mutants were combined with downstream shifting *ssl2* alleles, leading to the hypothesis

that the tested *ssl2* alleles increased scanning processivity, but this didn't lead to downstream

TSS shifts in mutants that increased initiation upstream. Regardless of mechanistic models for

how mutants are genetically interacting, double mutant analysis can reveal when mutations are

dependent on one another (epistasis), which is indicative of their defects being at the same

step, or when mutations function redundantly in an essential step (synergy or synthetic lethality),

or when mutations are perturbing a system independently, whereby double mutant phenotypes

will be the additive consequences of the single mutants combined. To therefore functionally

dissect our mutants, we tested interactions at inter-subunit interfaces and further analyzed

functional interactions across the PIC (**Figs. 6b, 6c; 7; S4**).

We first tested a potential Rpb10-Tfg1 interface in TSS selection by genetic interaction

studies and site-directed mutagenesis. We isolated *rpb10* L25S as a downstream-shifting

suppressor of the *imd2Δ::HIS3* reporter and showed this allele could also suppress the *cyc1-*

*1019::HIS3* reporter (**Fig. 2b**). Because Rpb10 is a common subunit shared amongst three RNA

polymerases, it was interesting to find a Pol II-specific phenotype. Examination of Pol PIC

structures for Pol I, II, and III PICs suggested that an interaction between Rpb10 in the vicinity of

the Rpb10 helix-loop-helix L25 residue might only be present in the Pol II PIC, where Rpb10

appears to contact a loop in the Tfg1 N-terminal domain (**Fig. 6b**). This potential interaction

suggested that Rpb10 might affect Pol II TSS distributions through an interaction with TFIIF

subunit Tfg1 that may be perturbed in the *rpb10* L25S allele. To examine this, we targeted

Rpb10 E29 and Tfg2 R26 residues that display potential interactions based on structural studies

(SCHILBACH *et al.* 2021) (**Fig. 6c**). We generated a series of genomic mutations by CRISPR

including charge reversals at both residues and assessed growth and TSS-shift reporter

phenotypes. On analysis of mutants, we did not observe overall growth defects in *tfg*1 single or

*tfg1 rpb10* double mutants (**Fig. 6c**, rows 6 and 7). Charge reversal *rpb10* mutation of E29R

resulted in suppression of the downstream shift reporter while E29A did not display effects (**Fig.**

**6c**, rows 3 and 4). However, when *rpb10* E29R was combined with the *tfg1* R26A or R26E

mutations, apparent suppression of the downstream shift phenotype was observed (**Fig. 6c**,

rows 10 and 11). Moreover, we found that *tfg1* mutants also suppressed the leaky His$^+$

phenotype of *imd2Δ::HIS3*, with Tfg1 R26E being more robust than R26A (**Fig. 6c**, rows 6 and

7, SC-His panel). Suppression of the leaky His$^+$ phenotype of *imd2Δ::HIS3* might result from an

upstream TSS-shifting phenotype, therefore we tested these *tfg1* mutants against the upstream

reporter *adhΔ50::HIS3* (**Fig. 6c**, lower panel) but *tfg1* mutants did not suppress *adhΔ50::HIS3*.

These results show that an additional *rpb10* allele can alter TSS selection (**Fig. 6c**) and that its

phenotype can be modulated by proximal *tfg1* alleles. However, we could not prove that Rpb10-

Tfg1 interactions underlie the *rpb10* L25S mutant phenotype.

We tested additional functional interactions by tetrad dissection analysis of diploids

followed by fitness analysis of each spore by pixel quantification using EBImage (PAU *et al.*

2010; NAMJILSUREN AND ARNDT 2025). We then calculated "expected" double mutant growth

phenotype from single mutant phenotypes by a multiplicative model for independent effects on

fitness of single mutants in a double mutant combination (MANI *et al.* 2008) and compared this

expected double mutant growth phenotype to the observed double mutant growth phenotype.

Genetic interactions were established based on comparisons of these double mutant metrics.

We examined upstream shifting *rpb9Δ* genetic interactions with a panel of mutants. First, we

examined *rpb9Δ* in conjunction with the upstream-shifting *rpb2 G369C* as the proximity of Rpb9

to Rpb2 and TFIIF raises the possibility that these mutants work in the same pathway, as it has

previously been proposed that *rpb2 G369C* works through effects on TFIIF. Second, we tested

*rpb9Δ* against a subset of our novel downstream shifting alleles to ask if they were additive or

epistatic. The GTF TFIIF binds at the interface between Rpb9 and Rpb2 in the protrusion-lobe

domain with proximity to Rpb2 G369 (**Fig. 4, 7b**)(CHEN *et al.* 2007). Because *rpb2* G369C and

*rpb9Δ* are both upstream shifting mutations, it is expected that the double mutant in case of non-

interaction would be a product of individual fitnesses. Conversely, if they functioned together in

the same process, we might observe less than multiplicative effects (sometimes referred to as a

specific form of epistasis). Instead, we observed growth fitness of *rpb2* G369C *rpb9Δ* double

mutants to be worse than predicted from the multiplicative model, indicating a synthetic defect

(**Fig. 7d**).

When combining upstream shifting *rpb9Δ* with a panel of downstream shifting mutants,

*tfb2* F189S, *tfb4* N71K, *ssl1* I122T, *rad3* K499E, *rpb10* L25S and *tfg2* C222Y by tetrad

dissection analysis (**Fig. S4**). We found *rpb9Δ rpb10* L25S double mutant fitness to be similar to

the *rpb9Δ* single mutant growth defect, with resultant fitness around the predicted fitness (**Fig.**

**S4**, bottom, middle). In addition, we found *rpb9Δ tfg2* C222Y double mutant fitness to be slightly

worse than expected (**Fig. S4**, bottom, right). Because these mutants have opposite effects on

TSS shifts, combination might be predicted to result in suppression of growth defects, if the

effects of each mutant on TSS selection are additive *and* the growth defects observed

correspond to the TSS selection phenotype. This kind of interaction between *rpb9Δ* and *rpb10*

L25S to be epistatic where the defect of one mutation (upstream) cannot be compensated by a

second mutation working in an opposite direction (downstream). Conversely, for interaction

between *tfg2* C22Y and *rpb9Δ*, if either of the *tfg2* C222Y or *rpb9Δ* effects required a WT allele

for their specific phenotypes, then instead of suppression of phenotypes by two opposite acting

alleles, we might observe exacerbation due to conversion of a downstream or upstream shifting

allele to the other class in the double mutant. Assuming the change in growth fitness reflects an

equivalent change in TSS selection, this kind of interaction could represent sign epistasis.

In contrast to the *rpb9Δ-tfg2* C222Y negative interaction, we observed clear double mutant

suppression between *rpb9Δ tfb2* F189S, *rpb9Δ tfb4* N71K, *rpb9Δ ssl1* I122T and *rpb9Δ rad3*

K499E, where each TFIIH allele suppressed growth defects associated with *rpb9Δ* (**Fig. S4**, top

panel and bottom, left). This indicates that the growth defect induced by upstream mutant *rpb9Δ*

was compensated by all tested TFIIH downstream mutants and suggesting that the growth

defect of *rpb9Δ* could be initiation related. This supposition is because TFIIH's gene expression

functions are limited to the PIC. This result was unexpected based on a simple model where the

*rpb9Δ* upstream-shifting phenotype would be caused by loss of its restraint of the key Pol II

catalytic domain, the trigger loop (TL) (KASTER *et al.* 2016). This model was based on *rpb9Δ*

conferring similar phenotypes to hyperactive mutants in the Rpb1 TL (upstream TSS shifts and

increased elongation rate *in vitro*). Upstream-shifting TL mutants are epistatic to downstream-

shifting *ssl2* TFIIH alleles, and therefore similar interactions between *rpb9Δ* and downstream-

shift reporter specific-TFIIH alleles was what was previously expected (see Discussion).

**Novel PIC alleles shift TSSs genome wide while chromatin/elongation alleles have muted**

**effects**

To gain insight into the function of a subset of our TSS shifting mutants, we quantified TSSs

across a curated set of promoters genome wide by STRIPE-seq in WT and mutant strains

(POLICASTRO *et al.* 2020; POLICASTRO *et al.* 2021; POLICASTRO AND ZENTNER 2022) (**Fig. 8a**).

We chose 8 downstream-shifting mutants, six from the PIC class and two from the

chromatin/elongation class isolated in our studies namely *tfb4* N71K, *rad3* K499E, *ssl1* I122T,

*tfb2* F189S, *rpb1* A480S, and *rpb10* L25S (PIC mutants) and *spn1* V199G and *spt5* G814*

(chromatin/elongation mutants) together with their corresponding WT parents (**Figs. 8**, **9**, **S5**,

**S6**). To assess TSS shifts in mutants relative to WT parents, we quantified TSSs as in our

previous studies (QIU *et al.* 2020; ZHAO *et al.* 2021) along with STRIPE-seq processing as in

(POLICASTRO *et al.* 2021). We focused our TSS analyses to 5979 Pol II promoter regions (RHEE

AND PUGH 2012; QIU *et al.* 2020; ZHAO *et al.* 2021) filtered to include promoters passing a cutoff

of a read count threshold in the promoter window analyzed resulting in analysis of n=2829

promoters. TSS usage values were then compared to those in WT by two TSS shift metrics as

in (QIU *et al.* 2020; ZHAO *et al.* 2021): shift in median TSS (Δmedian), which represents the

difference in median position of the TSS distribution at a promoter between a mutant and WT

strains (**Fig. 8b**) and change in spread (Δspread) that represents the change in width of

promoter TSS distributions relative to WT (**Fig. 9a**).

Individual replicates were generally reproducible for Δmedian TSS (**Fig. S5**). Principal

component analysis (PCA) of the Δmedian values of combined replicates showed three clusters

representing functional differences across the primary axis of variance (**Fig. 8c**): cluster 1

contained *rpb1* A480S, *rpb10* L25S, *ssl1* I122T and *tfb4* N71K, cluster 2 contained *rad3* K499E

and *tfb2* F189S, and cluster 3 contained *spt5* G814* and *spn1* V199G. When examining PCA of

individual replicates each compared to the average WT median position (**Fig. S5b**)**,** we found

chromatin alleles to cluster near the WT, suggesting they only have mild effects on TSS

distributions globally, but with greater effects for *spt5* G814* than for *spn1* V199G (**Fig. 8f**).

While both chromatin/elongation alleles did have effects beyond just an individual reporter, their

effects were quantitatively and qualitatively different from PIC alleles, consistent with potentially

relating to the chromatin context of individual promoters rather than innately affecting TSS

selection and promoter scanning at all promoters. All PIC alleles showed a large bias towards

downstream shifts in TSS distribution across the majority of promoters, with *rad3* K499E and

*tfb2* F189S showing the strongest effects (**Fig. 8f**).

PCA of Δspread values also displayed three groups and examination of PCA for

individual replicates of all strains compared to the average WT behavior shows that

chromatin/elongation mutants cluster with WT, most PIC mutants cluster together while *rad3*

K499E and *tfb2* F189S, which showed the strongest effects on Δmedian TSS, formed a

separate cluster based on Δspread (**Figs. 9, S6**). Examination of directional bias in spread

suggests that while spread was perturbed for most mutants, *rad3* K499E and *tfb2* F189S

showed the strongest increase in spread, consistent with the downstream shift in TSS usage

also expanding TSS spread to downstream positions (**Figs. 9e, S6d**). These results again

differentiated two TFIIH alleles, *rad3* K499E and *tfb2* F189S, from two others, *ssl1* I122T and

*tfb4* N71K.

**DISCUSSION**

Our studies reveal the wider landscape of genes and specific mutations that can alter the TSS

selection in yeast. In addition to independently identifying some already discovered alleles,

validating our novel reporters, we also identified an array of novel alleles suppressing our

initiation reporters. Notably, we did not find additional factors associated with promoters but not

directly connected to the PIC, such as chromatin remodelers or transcription coactivators, as

able to suppress our initiation reporters. This could simply be due to these remodelers not

strongly functioning in TSS accessibility at our reporters, or at least not in a way that might allow

reporter suppression upon their mutation. We find that mutations in many PIC components

including Rpb10 and additional TFIIH subunits can perturb initiation by promoter scanning when

mutated. Importantly, we find that the novel *rpb10* allele and tested TFIIH alleles alter TSS

selection genome wide. Our prior studies, new alleles identified here, and our parallel studies on

the Pol II/PIC interface reinforce that upstream shifting of TSSs results from compromise of

TFIIH-PIC interactions through the Tfb3-TFIIE-Rpb7 interface (BRABERG *et al.* 2013; YANG *et al.*

2022; YANG *et al.* 2024; BASNET *et al.* 2026). We extend evidence that the Rpb2-TFIIB

interaction is important for promoter scanning while also finding that perturbation to Rpo21/Rpb1

and Rpb2 broadly leads to suppression of downstream-shift sensitive reporters. This latter result

further supports the idea that defective Pol II catalysis/initiation activity results in phenotypes

consistent with downstream shifts in TSS usage. In contrast, we also have identified novel TFIIH

downstream-shifting alleles that are generally not located at inter-subunit interfaces, save one.

The downstream-shifting phenotype of these novel TFIIH alleles would be consistent

with a gain of function in TFIIH scanning processivity or rate. One interpretation of our alleles in

Rad3, Tfb2, Tfb4, and Ssl1 based on their locations is that they alter plasticity or dynamics

within the TFIIH structure. When this plasticity is altered, perturbations may propagate either to

the Ssl2 active site or to Ssl2 DNA binding properties, which in turn allow scanning to be

extended to downstream positions through increased processivity or increased scanning rate.

The idea that TFIIH processivity can be sensitive to small structural perturbations provides a

conceivable evolutionary path for TFIIH to gain ability to reach downstream TSSs during the

advent of the promoter scanning mechanism.

Potential communication between Ssl2 and the rest of the PIC can take a few paths.

First, there is a connection from Ssl2 to Tfb3 and Pol II/TFIIE through the following proteins:

Ssl2 to Tfb2 to Tfb4 to Ssl1 to Rad3 to Tfb3. Second, Tfb1 drapes over multiple TFIIH

components and connects to Tfa1 from TFIIE. Finally, Rad3 has a small number of residues

that may close the gap between it and Ssl2 and contact Ssl2 residues. We have identified

alleles along components that form the path from Ssl2 to Tfb3 (Tfb2, Tfb4, Ssl1, Tfb3) and

between Rad3 and Ssl2. If the protein connection between Ssl2 to Tfb3 has a certain amount of

flexibility or dynamics (using an analogy of an accordion's bellows or a pantograph that might

tolerate or allow extension or contraction, depending on rigidity), then alterations to this flexibility

might have consequences for Ssl2 rate or processivity as it translocates downstream on

promoter DNA. The locations of putative gain of function Tfb2, Tfb4, and Ssl1 alleles are all

consistent with perturbing internal intra-subunit interactions and we would propose they

increase flexibility of the "bellows" or "pantograph" formed by these components that connect

Ssl2 to Tfb3 and the PIC. This increased flexibility may then allow Ssl2 to translocate longer or

faster, resulting in a shift in TSS usage to downstream positions. Future biochemical and

biophysical experiments, potentially coupled with molecular dynamics simulations, might provide

tests of a model that, first, novel TFIIH alleles do indeed confer increased TFIIH processivity,

and second, that these alleles might alter structural properties of TFIIH in similar ways.

Chromatin/elongation factors can be envisioned to function in controlling initiation in two

ways. First, native chromatin architecture may reduce transcription initiation from nucleosome-

occluded TSSs (most yeast promoters have TSSs located within the boundaries of +1

nucleosomes) (RANDO AND WINSTON 2012). If there are downstream shifts in nucleosomes

dependent on elongation factors, then this could feed back to TSS usage and relax nucleosomal

inhibition of nucleosome-sensitive TSSs for promoters with appropriately positioned +1

nucleosomes. Second, if a promoter's TSSs are transcribed themselves from an upstream

promoter, then promoter-associated nucleosomes that are transcribed dependent on an

upstream TSS could become dependent on factors that restore transcribed nucleosomes in the

wake of elongating Pol II. This situation might reflect allele-selective effects on TSS usage for

transcribed promoters and less so global effects. This arrangement of an upstream promoter

driving transcription of a downstream promoter and thereby sensitizing the chromatin over the

downstream promoter is similar to that observed at the *SRG1-SER3* locus (MARTENS *et al.*

2004) and also the structure of the *lys2-128∂* allele (SIMCHEN *et al.* 1984). Each of these loci are

especially sensitive to chromatin and transcription elongation defects, and the latter of which

was used in the original identification of a number of the Spt chromatin/elongation factors –

factors that we isolated with our reporters here (SIMCHEN *et al.* 1984; CLARK-ADAMS AND

WINSTON 1987; CLARK-ADAMS *et al.* 1988; MALONE *et al.* 1991; SWANSON *et al.* 1991; CUI *et al.*

2016). In these cases, chromatin maintenance suppresses transcription initiation from within the

transcribed regions. Altered chromatin architecture in chromatin mutants can result in

widespread initiation from cryptic promoters (PRELICH AND WINSTON 1993; KAPLAN *et al.* 2003;

CHU *et al.* 2007; CHEUNG *et al.* 2008; DORIS *et al.* 2018) but perhaps also from chromatin-

sensitive TSSs adjacent to existing promoters. In either case, altered chromatin can result in

increased transcription from "suppressed" sites.

We rationalize the identification of chromatin elongation factors from the *imd2Δ::HIS3*

reporter because of its idiosyncratic transcription from upstream TSSs (generating a cryptic

unstable transcript (CUT) (DAVIS AND ARES 2006; KOPCEWICZ *et al.* 2007; JENKS *et al.* 2008;

KUEHNER AND BROW 2008)) that likely contributes to or controls nucleosome positioning over the

downstream, low-guanine inducible TSS. It was already known that certain *spt6* alleles could

increase usage of downstream TSSs at *IMD2* in the absence of known inducing conditions

(CONNELL *et al.* 2022). Here, we extend identification of chromatin and elongation factors that

can act at *IMD2* through suppression of *imd2Δ::HIS3* to *spt2*, *spt4*, *spt5*, *spt16*, *spn1*, *bur2, hta1,*

and *htb1* alleles. These results suggest that *imd2Δ::HIS3* may be similarly sensitive to the

chromatin/Spt⁻ class of mutant as is the classical *lys2-128∂* allele. However, we also isolated

chromatin mutants suppressing *cyc1-1019::HIS3* in addition to *imd2Δ::HIS3.* This *cyc1* allele

was inserted into a non-native locus that is devoid of genes that has previously been used to

house reporter genes (GRUBER *et al.* 2012). Our initial presumption was that the *CYC1* promoter

would not be itself transcribed, but examination of data from the Steinmetz lab (XU *et al.* 2009)

suggests that our *CYC1* promoter fragment also contains *SUT215*, a stable unannotated

transcript, and that integration of *cyc1-1019::HIS3* at chrI:199270-199271 causes it to be

flanked by *SUT004* upstream and *CUT442* downstream. Taken together, these provide support

that *cyc1-1019::HIS3* sensitivity to chromatin and elongation factors could be through

transcription over its promoter altering promoter nucleosomes, and we speculate

chromatin/elongation effects at other promoters may relate especially to those promoters that

have overlapping transcription units.

**Limitations of the study**

Our genetic interaction analysis indicated some interesting relationships between reporter

suppressor alleles, specifically the suppression observed between a subset of downstream-

shifting TFIIH alleles and *rpb9Δ*. Our previous genetic interaction studies showed that upstream

shifting Pol II catalytic alleles were epistatic to downstream-shifting *ssl2* alleles for both growth

phenotypes and for TSS shifts at *ADH1* (ZHAO *et al.* 2021). Our previous understanding for

*rpb9Δ* was that it was functioning similarly to upstream-shifting Pol II catalytic alleles (KASTER *et*

*al.* 2016). This assumption, coupled with the assumption that novel downstream-shifting TFIIH

alleles should function like *ssl2* downstream-shifting alleles, led to the expectation that *rpb9Δ*

would be epistatic to TFIIH alleles for TSS shift phenotypes. Additionally, we had the

expectation that *rpb9Δ* growth phenotypes related to its elongation defects and would not

necessarily be suppressible by initiation factor mutants. Contrary to these expectations, we find

the possibility of complexity in genetic interactions that might indicate different initiation alleles

are working in more complex ways than we had previously thought. We imagine these results

should be developed and extended in a few ways. For example, double mutant combinations of

*rpb9∆* and existing *ssl2* alleles and of Pol II alleles with novel TFIIH alleles would also allow

understanding of potential allele specificity. Additionally, examination of double mutants for

phenotypes on transcription reporters would allow to assess if initiation specific phenotypes

reflect overall growth phenotypes. Finally, quantitative primer extension analysis of double

mutants for *ADH1* TSS selection would also determine if growth phenotypes correlate with

specific changes to TSS distributions at promoters, where *ADH1* promoter has extensively

served as an excellent target for TSS distribution change assessment.

**Data availability**

Sequencing data generated in this study are available in the NCBI BioProject under accession

number PRJNA1337182. Strains used in this study, outlined in **Tables S1** and **S2**, are available

from the corresponding author upon request.

**Acknowledgements**

We acknowledge Dr. David Brow at the University of Wisconsin for guidance on genetic

selections. We are especially grateful to the students of BIOSC 0068 Foundations of Biology

Laboratory 2 course at the University of Pittsburgh for genetic selection of candidates leading to

identification of a subset of alleles described here. We acknowledge Gabe Zentner, Robert

Policastro, and Chhabi Govind for advice on STRIPE-seq. We would like to dedicate this paper

to the memory of Gabe Zentner.

**Funding**

Supported by funding from National Institutes of Health grants R01GM120450 and

R35GM144116 to C.D.K. This research was supported in part by the University of Pittsburgh

Center for Research Computing and Data, RRID:SCR_022735, through the resources provided.

Specifically, this work used the HTC cluster, which is supported by NIH award number

S10OD028483.

**Conflict of interest**

The authors declare no conflicts of interest.

## REFERENCES

Picard Toolkit 2019. Broad Institute, Broad Institute, GitHub Repository.
    https://broadinstitute.github.io/picard/.

Aibara, S., S. Schilbach and P. Cramer, 2021 Structures of mammalian RNA polymerase II pre-initiation complexes. Nature 594**:** 124-128.

Amberg, D. C., D. J. Burke and J. N. Strathern, 2005 Methods in Yeast Genetics: A Cold Spring Harbor Laboratory Course Manual. Cold Spring Harbor, NY: Cold Spring Harbor PressCold Spring Harbor, NY: Cold Spring Harbor Press.

Arribere, J. A., and W. V. Gilbert, 2013 Roles for transcript leaders in translation and mRNA decay revealed by transcript leader sequencing. Genome Res 23**:** 977-987.

Baek, I., L. J. Friedman, J. Gelles and S. Buratowski, 2021 Single-molecule studies reveal branched pathways for activator-dependent assembly of RNA polymerase II pre-initiation complexes. Mol Cell 81**:** 3576-3588 e3576.

Bangur, C. S., T. S. Pardee and A. S. Ponticelli, 1997 Mutational analysis of the D1/E1 core helices and the conserved N-terminal region of yeast transcription factor IIB (TFIIB): identification of an N-terminal mutant that stabilizes TATA-binding protein-TFIIB-DNA complexes. Mol Cell Biol 17**:** 6784-6793.

Basnet, P., Y. Zhu, I. O. Vvedenskaya, P. Arora, S. Hammer *et al.*, 2026 Important role in transcription start site selection for the RNA polymerase II-TFIIE-TFIIH interface in Saccharomyces cerevisiae. Genetics.

Berroteran, R. W., D. E. Ware and M. Hampsey, 1994 The sua8 suppressors of Saccharomyces cerevisiae encode replacements of conserved residues within the largest subunit of RNA polymerase II and affect transcription start site selection similarly to sua7 (TFIIB) mutations. Mol Cell Biol 14**:** 226-237.

Bolger, A. M., M. Lohse and B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30**:** 2114-2120.

Braberg, H., H. Jin, E. A. Moehle, Y. A. Chan, S. Wang *et al.*, 2013 From structure to systems: high-resolution, quantitative genetic analysis of RNA polymerase II. Cell 154**:** 775-788.

Breathnach, R., and P. Chambon, 1981 Organization and expression of eucaryotic split genes coding for proteins. Annu Rev Biochem 50**:** 349-383.

Brow, D. A., 2019 An Allosteric Network for Spliceosome Activation Revealed by High-Throughput Suppressor Analysis in Saccharomyces cerevisiae. Genetics 212**:** 111-124.

Buratowski, S., S. Hahn, P. A. Sharp and L. Guarente, 1988 Function of a yeast TATA element-binding protein in a mammalian transcription system. Nature 334**:** 37-42.

Cabart, P., H. Jin, L. Li and C. D. Kaplan, 2014 Activation and reactivation of the RNA polymerase II trigger loop for intrinsic RNA cleavage and catalysis. Transcription 5**:** e28869.

Chen, B. S., and M. Hampsey, 2004 Functional interaction between TFIIB and the Rpb2 subunit of RNA polymerase II: implications for the mechanism of transcription initiation. Mol Cell Biol 24**:** 3983-3991.

Chen, H. T., L. Warfield and S. Hahn, 2007 The positions of TFIIF and TFIIE in the RNA polymerase II transcription preinitiation complex. Nat Struct Mol Biol 14**:** 696-703.

Cheung, V., G. Chua, N. N. Batada, C. R. Landry, S. W. Michnick *et al.*, 2008 Chromatin- and transcription-related factors repress transcription from within coding regions throughout the Saccharomyces cerevisiae genome. PLoS Biol 6**:** e277.

Chu, Y., R. Simic, M. H. Warner, K. M. Arndt and G. Prelich, 2007 Regulation of histone modification and cryptic transcription by the Bur1 and Paf1 complexes. EMBO J 26**:** 4646-4656.

Cingolani, P., A. Platts, L. Wang le, M. Coon, T. Nguyen *et al.*, 2012 A program for annotating
and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the
genome of Drosophila melanogaster strain w1118; iso-2; iso-3. Fly (Austin) 6**:** 80-92.
Clark-Adams, C. D., D. Norris, M. A. Osley, J. S. Fassler and F. Winston, 1988 Changes in
histone gene dosage alter transcription in yeast. Genes Dev 2**:** 150-159.
Clark-Adams, C. D., and F. Winston, 1987 The SPT6 gene is essential for growth and is
required for delta-mediated transcription in Saccharomyces cerevisiae. Mol Cell Biol 7**:**
679-686.
Coin, F., J. C. Marinoni, C. Rodolfo, S. Fribourg, A. M. Pedrini *et al.*, 1998 Mutations in the XPD
helicase gene result in XP and TTD phenotypes, preventing interaction between XPD
and the p44 subunit of TFIIH. Nat Genet 20**:** 184-188.
Conaway, R. C., and J. W. Conaway, 1993 General initiation factors for RNA polymerase II.
Annu Rev Biochem 62**:** 161-190.
Connell, Z., T. J. Parnell, L. L. McCullough, C. P. Hill and T. Formosa, 2022 The interaction
between the Spt6-tSH2 domain and Rpb1 affects multiple functions of RNA Polymerase
II. Nucleic Acids Res 50**:** 784-802.
Cramer, P., D. A. Bushnell and R. D. Kornberg, 2001 Structural basis of transcription: RNA
polymerase II at 2.8 angstrom resolution. Science 292**:** 1863-1876.
Cui, P., H. Jin, M. R. Vutukuru and C. D. Kaplan, 2016 Relationships Between RNA Polymerase
II Activity and Spt Elongation Factors to Spt- Phenotype and Growth in Saccharomyces
cerevisiae. G3 (Bethesda) 6**:** 2489-2504.
Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks *et al.*, 2011 The variant call format
and VCFtools. Bioinformatics 27**:** 2156-2158.
Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan *et al.*, 2021 Twelve years of
SAMtools and BCFtools. Gigascience 10.
Davis, C. A., and M. Ares, Jr., 2006 Accumulation of unstable promoter-associated transcripts
upon loss of the nuclear exosome subunit Rrp6p in Saccharomyces cerevisiae. Proc
Natl Acad Sci U S A 103**:** 3262-3267.
Deatherage, D. E., and J. E. Barrick, 2014 Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using breseq. Methods Mol Biol 1151**:**
165-188.
Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski *et al.*, 2013 STAR: ultrafast
universal RNA-seq aligner. Bioinformatics 29**:** 15-21.
Doris, S. M., J. Chuang, O. Viktorovskaya, M. Murawska, D. Spatt *et al.*, 2018 Spt6 Is Required
for the Fidelity of Promoter Selection. Mol Cell 72**:** 687-699 e686.
Dudnyk, K., D. Cai, C. Shi, J. Xu and J. Zhou, 2024 Sequence basis of transcription initiation in
the human genome. Science 384**:** eadj0116.
Durr, G., J. Strayle, R. Plemper, S. Elbs, S. K. Klee *et al.*, 1998 The medial-Golgi ion pump
Pmr1 supplies the yeast secretory pathway with Ca2+ and Mn2+ required for
glycosylation, sorting, and endoplasmic reticulum-associated protein degradation. Mol
Biol Cell 9**:** 1149-1162.
Eichner, J., H. T. Chen, L. Warfield and S. Hahn, 2010 Position of the general transcription
factor TFIIF within the RNA polymerase II transcription preinitiation complex. EMBO J
29**:** 706-716.
Ellison, M. A., J. L. Walker, P. J. Ropp, J. D. Durrant and K. M. Arndt, 2020 MutantHuntWGS: A
Pipeline for Identifying Saccharomyces cerevisiae Mutations. G3 (Bethesda) 10**:** 3009-
3014.
Faitar, S. L., S. A. Brodie and A. S. Ponticelli, 2001 Promoter-specific shifts in transcription
initiation conferred by yeast TFIIB mutations are determined by the sequence in the
immediate vicinity of the start sites. Mol Cell Biol 21**:** 4427-4440.

Fazal, F. M., C. A. Meng, K. Murakami, R. D. Kornberg and S. M. Block, 2015 Real-time
observation of the initiation of RNA polymerase II transcription. Nature 525**:** 274-277.
Fischbeck, J. A., S. M. Kraemer and L. A. Stargell, 2002 SPN1, a conserved gene identified by
suppression of a postrecruitment-defective yeast TATA-binding protein mutant. Genetics
162**:** 1605-1616.
Fishburn, J., E. Tomko, E. Galburt and S. Hahn, 2015 Double-stranded DNA translocase activity
of transcription factor TFIIH and the mechanism of RNA polymerase II open complex
formation. Proc Natl Acad Sci U S A 112**:** 3961-3966.
Freire-Picos, M. A., S. Krishnamurthy, Z. W. Sun and M. Hampsey, 2005 Evidence that the
Tfg1/Tfg2 dimer interface of TFIIF lies near the active center of the RNA polymerase II
initiation complex. Nucleic Acids Res 33**:** 5045-5052.
Furter-Graves, E. M., R. Furter and B. D. Hall, 1991 SHI, a new yeast gene affecting the
spacing between TATA and transcription initiation sites. Mol Cell Biol 11**:** 4121-4127.
Furter-Graves, E. M., and B. D. Hall, 1990 DNA sequence elements required for transcription
initiation of the Schizosaccharomyces pombe ADH gene in Saccharomyces cerevisiae.
Mol Gen Genet 223**:** 407-416.
Ghazy, M. A., S. A. Brodie, M. L. Ammerman, L. M. Ziegler and A. S. Ponticelli, 2004 Amino
acid substitutions in yeast TFIIF confer upstream shifts in transcription initiation and
altered interaction with RNA polymerase II. Mol Cell Biol 24**:** 10975-10985.
Giardina, C., and J. T. Lis, 1993 DNA melting on yeast RNA polymerase II promoters. Science
261**:** 759-762.
Greger, I. H., A. Aranda and N. Proudfoot, 2000 Balancing transcriptional interference and
initiation on the GAL7 promoter of Saccharomyces cerevisiae. Proc Natl Acad Sci U S A
97**:** 8415-8420.
Greger, I. H., and N. J. Proudfoot, 1998 Poly(A) signals control both transcriptional termination
and initiation between the tandem GAL10 and GAL7 genes of Saccharomyces
cerevisiae. EMBO J 17**:** 4771-4779.
Gruber, J. D., K. Vogel, G. Kalay and P. J. Wittkopp, 2012 Contrasting properties of gene-
specific regulatory, coding, and copy number mutations in Saccharomyces cerevisiae:
frequency, effects, and dominance. PLoS Genet 8**:** e1002497.
Hampsey, M., 1991 A tester system for detecting each of the six base-pair substitutions in
Saccharomyces cerevisiae by selecting for an essential cysteine in iso-1-cytochrome c.
Genetics 128**:** 59-67.
Hampsey, M., 2006 The Pol II initiation complex: finding a place to start. Nat Struct Mol Biol 13**:**
564-566.
He, Y., C. Yan, J. Fang, C. Inouye, R. Tjian *et al.*, 2016 Near-atomic resolution visualization of
human transcription promoter opening. Nature 533**:** 359-365.
Hekmatpanah, D. S., and R. A. Young, 1991 Mutations in a conserved region of RNA
polymerase II influence the accuracy of mRNA start site selection. Mol Cell Biol 11**:**
5781-5791.
Hemming, S. A., D. B. Jansma, P. F. Macgregor, A. Goryachev, J. D. Friesen *et al.*, 2000 RNA
polymerase II subunit Rpb9 regulates transcription elongation in vivo. J Biol Chem 275**:**
35506-35511.
Hull, M. W., K. McKune and N. A. Woychik, 1995 RNA polymerase II subunit RPB9 is required
for accurate start site selection. Genes Dev 9**:** 481-490.
Hurowitz, E. H., and P. O. Brown, 2003 Genome-wide analysis of mRNA lengths in
Saccharomyces cerevisiae. Genome Biol 5**:** R2.
Janke, C., M. M. Magiera, N. Rathfelder, C. Taxis, S. Reber *et al.*, 2004 A versatile toolbox for
PCR-based tagging of yeast genes: new fluorescent proteins, more markers and
promoter substitution cassettes. Yeast 21**:** 947-962.

Jenks, M. H., T. W. O'Rourke and D. Reines, 2008 Properties of an intergenic terminator and
start site switch that regulate IMD2 transcription in yeast. Mol Cell Biol 28**:** 3883-3893.
Jin, H., and C. D. Kaplan, 2014 Relationships of RNA polymerase II genetic interactors to
transcription start site usage defects and growth in Saccharomyces cerevisiae. G3
(Bethesda) 5**:** 21-33.
Kadonaga, J. T., 1990 Assembly and disassembly of the Drosophila RNA polymerase II
complex during transcription. J Biol Chem 265**:** 2624-2631.
Kadonaga, J. T., 2012 Perspectives on the RNA polymerase II core promoter. Wiley Interdiscip
Rev Dev Biol 1**:** 40-51.
Kaplan, C. D., 2013 Basic mechanisms of RNA polymerase II activity and alteration of gene
expression in Saccharomyces cerevisiae. Biochim Biophys Acta 1829**:** 39-54.
Kaplan, C. D., M. J. Holland and F. Winston, 2005 Interaction between transcription elongation
factors and mRNA 3'-end formation at the Saccharomyces cerevisiae GAL10-GAL7
locus. J Biol Chem 280**:** 913-922.
Kaplan, C. D., H. Jin, I. L. Zhang and A. Belyanin, 2012 Dissection of Pol II trigger loop function
and Pol II activity-dependent control of start site selection in vivo. PLoS Genet 8**:**
e1002627.
Kaplan, C. D., L. Laprade and F. Winston, 2003 Transcription elongation factors repress
transcription initiation from cryptic sites. Science 301**:** 1096-1099.
Kaster, B. C., K. C. Knippa, C. D. Kaplan and D. O. Peterson, 2016 RNA Polymerase II Trigger
Loop Mobility: INDIRECT EFFECTS OF Rpb9. J Biol Chem 291**:** 14883-14895.
Khaperskyy, D. A., M. L. Ammerman, R. C. Majovski and A. S. Ponticelli, 2008 Functions of
Saccharomyces cerevisiae TFIIF during transcription start site utilization. Mol Cell Biol
28**:** 3757-3766.
Kim, J. S., C. Saint-Andre, H. S. Lim, C. S. Hwang, J. M. Egly *et al.*, 2015 Crystal structure of
the Rad3/XPD regulatory domain of Ssl1/p44. J Biol Chem 290**:** 8321-8330.
Klopotowski, T., and A. Wiater, 1965 Synergism of aminotriazole and phosphate on the
inhibition of yeast imidazole glycerol phosphate dehydratase. Arch Biochem Biophys
112**:** 562-566.
Knaus, R., R. Pollock and L. Guarente, 1996 Yeast SUB1 is a suppressor of TFIIB mutations
and has homology to the human co-activator PC4. EMBO J 15**:** 1933-1940.
Kopcewicz, K. A., T. W. O'Rourke and D. Reines, 2007 Metabolic regulation of IMD2
transcription and an unusual DNA element that generates short transcripts. Mol Cell Biol
27**:** 2821-2829.
Kornberg, R. D., 2007 The molecular basis of eukaryotic transcription. Proc Natl Acad Sci U S A
104**:** 12955-12961.
Kostrewa, D., M. E. Zeller, K. J. Armache, M. Seizl, K. Leike *et al.*, 2009 RNA polymerase II-
TFIIB structure and mechanism of transcription initiation. Nature 462**:** 323-330.
Kubicek, C. E., R. D. Chisholm, S. Takayama and D. K. Hawley, 2013 RNA polymerase II
mutations conferring defects in poly(A) site cleavage and termination in Saccharomyces
cerevisiae. G3 (Bethesda) 3**:** 167-180.
Kuehner, J. N., and D. A. Brow, 2006 Quantitative analysis of in vivo initiator selection by yeast
RNA polymerase II supports a scanning model. J Biol Chem 281**:** 14119-14128.
Kuehner, J. N., and D. A. Brow, 2008 Regulation of a eukaryotic gene by GTP-dependent start
site selection and transcription attenuation. Mol Cell 31**:** 201-211.
Langmead, B., and S. L. Salzberg, 2012 Fast gapped-read alignment with Bowtie 2. Nat
Methods 9**:** 357-359.
Laughery, M. F., T. Hunter, A. Brown, J. Hoopes, T. Ostbye *et al.*, 2015 New vectors for simple
and streamlined CRISPR-Cas9 genome editing in Saccharomyces cerevisiae. Yeast 32**:**
711-720.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25**:** 2078-2079.
Lifton, R. P., M. L. Goldberg, R. W. Karp and D. S. Hogness, 1978 The organization of the
histone genes in Drosophila melanogaster: functional and evolutionary implications. Cold
Spring Harb Symp Quant Biol 42 Pt 2**:** 1047-1051.
Lin, Y. C., and J. D. Gralla, 2005 Stimulation of the XPB ATP-dependent helicase by the beta
subunit of TFIIE. Nucleic Acids Res 33**:** 3072-3081.
Lu, Z., and Z. Lin, 2019 Pervasive and dynamic transcription initiation in Saccharomyces
cerevisiae. Genome Res 29**:** 1198-1210.
Lu, Z., and Z. Lin, 2021 The origin and evolution of a distinct mechanism of transcription
initiation in yeasts. Genome Res 31**:** 51-63.
Luse, D. S., M. Parida, B. M. Spector, K. A. Nilson and D. H. Price, 2020 A unified view of the
sequence and functional organization of the human RNA polymerase II promoter.
Nucleic Acids Res 48**:** 7767-7785.
Malabat, C., F. Feuerbach, L. Ma, C. Saveanu and A. Jacquier, 2015 Quality control of
transcription start site selection by nonsense-mediated-mRNA decay. Elife 4.
Malik, I., C. Qiu, T. Snavely and C. D. Kaplan, 2017 Wide-ranging and unexpected
consequences of altered Pol II catalytic activity in vivo. Nucleic Acids Res 45**:** 4431-
4451.
Malone, E. A., C. D. Clark, A. Chiang and F. Winston, 1991 Mutations in SPT16/CDC68
suppress cis- and trans-acting mutations that affect promoter function in Saccharomyces
cerevisiae. Mol Cell Biol 11**:** 5710-5717.
Mandal, D., T. B. Woolf and R. Rao, 2000 Manganese selectivity of pmr1, the yeast secretory
pathway ion pump, is defined by residue gln783 in transmembrane segment 6. Residue
Asp778 is essential for cation transport. J Biol Chem 275**:** 23933-23938.
Mani, R., R. P. St Onge, J. L. t. Hartman, G. Giaever and F. P. Roth, 2008 Defining genetic
interaction. Proc Natl Acad Sci U S A 105**:** 3461-3466.
Martens, J. A., L. Laprade and F. Winston, 2004 Intergenic transcription is required to repress
the Saccharomyces cerevisiae SER3 gene. Nature 429**:** 571-574.
Martin, M., 2011 Cutadapt removes adapter sequences from high-throughput sequencing reads.
2011 17**:** 3.
Miller, G., and S. Hahn, 2006 A DNA-tethered cleavage probe reveals the path for promoter
DNA in the yeast preinitiation complex. Nat Struct Mol Biol 13**:** 603-610.
Miura, F., N. Kawaguchi, J. Sese, A. Toyoda, M. Hattori *et al.*, 2006 A large-scale full-length
cDNA analysis to explore the budding yeast transcriptome. Proc Natl Acad Sci U S A
103**:** 17846-17851.
Murakami, K., P. J. Mattei, R. E. Davis, H. Jin, C. D. Kaplan *et al.*, 2015 Uncoupling Promoter
Opening from Start-Site Scanning. Mol Cell 59**:** 133-138.
Na, J. G., I. Pinto and M. Hampsey, 1992 Isolation and characterization of SUA5, a novel gene
required for normal growth in Saccharomyces cerevisiae. Genetics 131**:** 791-801.
Nagalakshmi, U., Z. Wang, K. Waern, C. Shou, D. Raha *et al.*, 2008 The transcriptional
landscape of the yeast genome defined by RNA sequencing. Science 320**:** 1344-1349.
Namjilsuren, S., and K. M. Arndt, 2025 Glc7/PP1 triggers Paf1 complex dissociation from RNA
polymerase II to enable transcription termination. bioRxiv.
Niyogi, S. K., R. P. Feldman and D. J. Hoffman, 1981 Selective effects of metal ions on RNA
synthesis rates. Toxicology 22**:** 9-21.
Orphanides, G., T. Lagrange and D. Reinberg, 1996 The general transcription factors of RNA
polymerase II. Genes Dev 10**:** 2657-2683.
Pardee, T. S., C. S. Bangur and A. S. Ponticelli, 1998 The N-terminal region of yeast TFIIB
contains two adjacent functional domains involved in stable RNA polymerase II binding
and transcription start site selection. J Biol Chem 273**:** 17859-17864.

Park, D., A. R. Morris, A. Battenhouse and V. R. Iyer, 2014 Simultaneous mapping of transcript ends at single-nucleotide resolution and identification of widespread promoter-associated non-coding RNA governed by TATA elements. Nucleic Acids Res 42: 3736-3749.

Pau, G., F. Fuchs, O. Sklyar, M. Boutros and W. Huber, 2010 EBImage--an R package for image processing with applications to cellular phenotypes. Bioinformatics 26: 979-981.

Pelechano, V., W. Wei and L. M. Steinmetz, 2013 Extensive transcriptional heterogeneity revealed by isoform profiling. Nature 497: 127-131.

Pinto, I., J. G. Na, F. Sherman and M. Hampsey, 1992a cis- and trans-acting suppressors of a translation initiation defect at the cyc1 locus of Saccharomyces cerevisiae. Genetics 132: 97-112.

Pinto, I., D. E. Ware and M. Hampsey, 1992b The yeast SUA7 gene encodes a homolog of human transcription factor TFIIB and is required for normal start site selection in vivo. Cell 68: 977-988.

Pinto, I., W. H. Wu, J. G. Na and M. Hampsey, 1994 Characterization of sua7 mutations defines a domain of TFIIB involved in transcription start site selection in yeast. J Biol Chem 269: 30569-30573.

Policastro, R. A., D. J. McDonald, V. P. Brendel and G. E. Zentner, 2021 Flexible analysis of TSS mapping data and detection of TSS shifts with TSRexploreR. NAR Genom Bioinform 3: lqab051.

Policastro, R. A., R. T. Raborn, V. P. Brendel and G. E. Zentner, 2020 Simple and efficient profiling of transcription initiation and transcript levels with STRIPE-seq. Genome Res 30: 910-923.

Policastro, R. A., and G. E. Zentner, 2022 Genome-Wide Profiling of Transcription Initiation with STRIPE-seq. Methods Mol Biol 2477: 21-34.

Powers, E. N., C. Chan, E. Doron-Mandel, L. Llacsahuanga Allcca, J. Kim Kim et al., 2022 Bidirectional promoter activity from expression cassettes can drive off-target repression of neighboring gene translation. Elife 11.

Prelich, G., and F. Winston, 1993 Mutations that suppress the deletion of an upstream activating sequence in yeast: involvement of a protein kinase and histone H3 in repressing transcription in vivo. Genetics 135: 665-676.

Qiu, C., P. Arora, I. Malik, A. J. Laperuta, E. M. Pavlovic et al., 2024 Thiolutin has complex effects in vivo but is a direct inhibitor of RNA polymerase II in vitro. Nucleic Acids Res 52: 2546-2564.

Qiu, C., O. C. Erinne, J. M. Dave, P. Cui, H. Jin et al., 2016 High-Resolution Phenotypic Landscape of the RNA Polymerase II Trigger Loop. PLoS Genet 12: e1006321.

Qiu, C., H. Jin, I. Vvedenskaya, J. A. Llenas, T. Zhao et al., 2020 Universal promoter scanning by Pol II during transcription initiation in Saccharomyces cerevisiae. Genome Biol 21: 132.

Rando, O. J., and F. Winston, 2012 Chromatin and transcription in yeast. Genetics 190: 351-387.

Rhee, H. S., and B. F. Pugh, 2012 Genome-wide structure and organization of eukaryotic pre-initiation complexes. Nature 483: 295-301.

Roeder, R. G., 1996 The role of general initiation factors in transcription by RNA polymerase II. Trends Biochem Sci 21: 327-335.

Roeder, R. G., 2019 50+ years of eukaryotic transcription: an expanding universe of factors and mechanisms. Nat Struct Mol Biol 26: 783-791.

Rudolph, H. K., A. Antebi, G. R. Fink, C. M. Buckley, T. E. Dorman et al., 1989 The yeast secretory pathway is perturbed by mutations in PMR1, a member of a Ca2+ ATPase family. Cell 58: 133-145.

Sainsbury, S., J. Niesser and P. Cramer, 2013 Structure and function of the initially transcribing RNA polymerase II-TFIIB complex. Nature 493**:** 437-440.

Sandelin, A., P. Carninci, B. Lenhard, J. Ponjavic, Y. Hayashizaki *et al.*, 2007 Mammalian RNA polymerase II core promoters: insights from genome-wide studies. Nat Rev Genet 8**:** 424-436.

Schilbach, S., S. Aibara, C. Dienemann, F. Grabbe and P. Cramer, 2021 Structure of RNA polymerase II pre-initiation complex at 2.9 A defines initial DNA opening. Cell 184**:** 4064-4072 e4028.

Schilbach, S., M. Hantsche, D. Tegunov, C. Dienemann, C. Wigge *et al.*, 2017 Structures of transcription pre-initiation complex with TFIIH and Mediator. Nature 551**:** 204-209.

Schmitt, M. E., T. A. Brown and B. L. Trumpower, 1990 A rapid and simple method for preparation of RNA from Saccharomyces cerevisiae. Nucleic Acids Res 18**:** 3091-3092.

Sikorski, T. W., S. B. Ficarro, J. Holik, T. Kim, O. J. Rando *et al.*, 2011 Sub1 and RPA associate with RNA polymerase II at different stages of transcription. Mol Cell 44**:** 397-409.

Simchen, G., F. Winston, C. A. Styles and G. R. Fink, 1984 Ty-mediated gene expression of the LYS2 and HIS4 genes of Saccharomyces cerevisiae is controlled by the same SPT genes. Proc Natl Acad Sci U S A 81**:** 2431-2434.

Smale, S. T., and J. T. Kadonaga, 2003 The RNA polymerase II core promoter. Annu Rev Biochem 72**:** 449-479.

Smith, T., A. Heger and I. Sudbery, 2017 UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. Genome Res 27**:** 491-499.

Struhl, K., 1987 Promoters, activator proteins, and the mechanism of transcriptional initiation in yeast. Cell 49**:** 295-297.

Struhl, K., and R. W. Davis, 1977 Production of a functional eukaryotic enzyme in Escherichia coli: cloning and expression of the yeast structural gene for imidazole-glycerolphosphate dehydratase (his3). Proc Natl Acad Sci U S A 74**:** 5255-5259.

Sun, Z. W., and M. Hampsey, 1995 Identification of the gene (SSU71/TFG1) encoding the largest subunit of transcription factor TFIIF as a suppressor of a TFIIB mutation in Saccharomyces cerevisiae. Proc Natl Acad Sci U S A 92**:** 3127-3131.

Sun, Z. W., and M. Hampsey, 1996 Synthetic enhancement of a TFIIB defect by a mutation in SSU72, an essential yeast gene encoding a novel protein that affects transcription start site selection in vivo. Mol Cell Biol 16**:** 1557-1566.

Sun, Z. W., A. Tessmer and M. Hampsey, 1996 Functional interaction between TFIIB and the Rpb9 (Ssu73) subunit of RNA polymerase II in Saccharomyces cerevisiae. Nucleic Acids Res 24**:** 2560-2566.

Swanson, M. S., E. A. Malone and F. Winston, 1991 SPT5, an essential gene important for normal transcription in Saccharomyces cerevisiae, encodes an acidic nuclear protein with a carboxy-terminal repeat. Mol Cell Biol 11**:** 3009-3019.

Tomko, E. J., J. Fishburn, S. Hahn and E. A. Galburt, 2017 TFIIH generates a six-base-pair open complex during RNAP II transcription initiation and start-site scanning. Nat Struct Mol Biol 24**:** 1139-1145.

Tomko, E. J., O. Luyties, J. K. Rimel, C. L. Tsai, J. O. Fuss *et al.*, 2021 The Role of XPB/Ssl2 dsDNA Translocase Processivity in Transcription Start-site Scanning. J Mol Biol 433**:** 166813.

Vaser, R., S. Adusumalli, S. N. Leng, M. Sikic and P. C. Ng, 2016 SIFT missense predictions for genomes. Nat Protoc 11**:** 1-9.

Vo Ngoc, L., C. J. Cassidy, C. Y. Huang, S. H. Duttke and J. T. Kadonaga, 2017a The human initiator is a distinct and abundant element that is precisely positioned in focused core promoters. Genes Dev 31**:** 6-11.

Vo Ngoc, L., Y. L. Wang, G. A. Kassavetis and J. T. Kadonaga, 2017b The punctilious RNA polymerase II core promoter. Genes Dev 31**:** 1289-1301.

Wang, M., P. Arora, C. D. Kaplan and D. A. Brow, 2025 Substitutions in RNA-binding protein
Hrp1 map a potential interaction surface with the yeast RNA polymerase II elongation
complex. bioRxiv.
Winston, F., D. T. Chaleff, B. Valent and G. R. Fink, 1984 Mutations affecting Ty-mediated
expression of the HIS4 gene of Saccharomyces cerevisiae. Genetics 107**:** 179-197.
Woychik, N. A., W. S. Lane and R. A. Young, 1991 Yeast RNA polymerase II subunit RPB9 is
essential for growth at temperature extremes. J Biol Chem 266**:** 19053-19055.
Wu, W. H., I. Pinto, B. S. Chen and M. Hampsey, 1999 Mutational analysis of yeast TFIIB. A
functional relationship between Ssu72 and Sub1/Tsp1 defined by allele-specific
interactions with TFIIB. Genetics 153**:** 643-652.
Xie, Y., H. Li, X. Luo, H. Li, Q. Gao *et al.*, 2022 IBS 2.0: an upgraded illustrator for the
visualization of biological sequences. Nucleic Acids Res 50**:** W420-W426.
Xu, Z., W. Wei, J. Gagneur, F. Perocchi, S. Clauder-Munster *et al.*, 2009 Bidirectional promoters
generate pervasive transcription in yeast. Nature 457**:** 1033-1037.
Yang, C., P. Basnet, S. Sharmin, H. Shen, C. D. Kaplan *et al.*, 2024 Transcription start site
scanning requires the fungi-specific hydrophobic loop of Tfb3. Nucleic Acids Res 52**:**
11602-11611.
Yang, C., R. Fujiwara, H. J. Kim, P. Basnet, Y. Zhu *et al.*, 2022 Structural visualization of de
novo transcription initiation by Saccharomyces cerevisiae RNA polymerase II. Mol Cell
82**:** 660-676 e669.
Yoon, H., S. P. Miller, E. K. Pabich and T. F. Donahue, 1992 SSL1, a suppressor of a HIS4 5'-
UTR stem-loop mutation, is essential for translation initiation and affects UV resistance
in yeast. Genes Dev 6**:** 2463-2477.
Zhang, Z., and F. S. Dietrich, 2005 Mapping of transcription start sites in Saccharomyces
cerevisiae using 5' SAGE. Nucleic Acids Res 33**:** 2838-2851.
Zhao, T., I. O. Vvedenskaya, W. K. Lai, S. Basu, B. F. Pugh *et al.*, 2021 Ssl2/TFIIH function in
transcription start site scanning by RNA polymerase II in Saccharomyces cerevisiae.
Elife 10.
Zhu, Y., I. O. Vvedenskaya, S. H. Sze, B. E. Nickels and C. D. Kaplan, 2024 Quantitative
analysis of transcription start site selection reveals control by DNA sequence, RNA
polymerase II activity and NTP levels. Nat Struct Mol Biol 31**:** 190-202.

1317

1318

1319

1320

**Figure 1. TSS shift-sensitive reporters detect TSS shifts by *HIS3* expression. a. Left.**
Schematic of initiation reporters *imd2Δ::HIS3* and *cyc1-1019::HIS3* sensitive to downstream
shifts in TSS usage. Downstream-shifting TSS mutants constitutively express *IMD2* transcripts
from a downstream TSS that allows functional *HIS3* expression when *HIS3* replaces the *IMD2*
ORF, conferring a His⁺ phenotype (top). Downstream TSS shifts at *cyc1-1019::HIS3* allow an
upstream AUG to be bypassed, resulting in usage of the appropriate in-frame AUG, allowing
functional His3p translation (bottom). **Right.** Schematic of the initiation reporters *adhΔ50::HIS3*
(top) and *hmo1Δ38::HIS3* (bottom) that are sensitive to upstream shifts in TSS usage.
Truncation places the native TSSs for these promoters too close to the native AUG rendering
ineffective translation initiation. Upstream shifts in TSS selection lengthen the 5′ UTRs for each
reporter, allowing His3p to be translated from the lengthened *HIS3* mRNA. **b.** Validation of
reporter phenotypes with known TSS-shifting mutants: 10-fold serially diluted yeast strains or a
single concentration of cells (marked with asterisk – "single spot phenotyping") are plated on
media to illustrate reporter function. Here, the *cyc1-5000* allele-based reporter was deployed for
isolating *sua* alleles, where the *cyc1-5000* allele resembles *cyc1-1019* allele (1a, left),
possessing additional C→A and C→G mutations at -23 and -17 positions respectively, relative

to +1 A of the native ATG. This makes the context flanking the upstream AUG start codon more
preferred, making the *cyc1-5000* downstream shift detection more stringent than the *cyc1-1019*
reporter. *rpb1* alleles (upstream or downstream shifts*)*, *sub1Δ* (downstream shift), and *rpb9Δ*
(upstream shift) mutants were tested against the upstream and downstream shift sensitive
reporters to illustrate allele-specific reporter activity. Note that *cyc1-1019::HIS3* can be
expressed by strong downstream shifting *sub1Δ* and *rpb1* H1085Y of all downstream *rpb1*
mutations tested.

**a**

**b**

1344

**Figure 2. Isolation of spontaneous suppressors of downstream TSS-shift sensitive**
**initiation reporters. a**. Hierarchically clustered growth phenotypes for a subset of candidate
downstream-shifting suppressors of *cyc1-1019::HIS3* (left) or *imd2Δ::HIS3* (right) reporters with
a subset of identified mutations annotated; right bottom panel displays *imd2Δ::HIS3*
suppressors isolated by selection on SC-His-Lys+5mM 3AT. Phenotypes of candidate
suppressors displayed are relative phenotypes of candidates that have been normalized to
control conditions, as assessed from single-spot phenotyping assays. Phenotypes were scored
from spot assay, corrected to WT and then corrected to the same on control condition (see
Methods). Phenotypic quantification therefore assigns a value of zero for WT-like growth, a
negative value if phenotype is worse than WT and positive, if growth is better than WT and
value of phenotype represents strength of defect. **b.** Validation of a subset of candidates by
knock-in of TSS mutants into *imd2Δ::HIS3* and *cyc1-1019::HIS3* reporter backgrounds and re-
phenotyping. Spot assay phenotyping of candidates isolated from genetic selection (candidate)
and reincorporated mutants with a *kanMX6* tag (*corresponding mutation::kanMX6*) displaying
replication of candidate phenotypes in reincorporated mutants. Mutations were amplified out of
isolated candidates, PCR sewed with a *kanMX6* drug marker cassette and re-introduced into
each of the reporter strains by integrative transformation. The *rpb10* candidate mutation was
introduced using CRISPR/Cas9.

1363
**Figure 3. Isolation of spontaneous suppressors of upstream TSS-shift sensitive initiation**
**reporters. a.** Hierarchically clustered single-spot phenotypes of a subset of upstream-shifting

suppressors of *adhΔ50::HIS3* (left) and *hmo1Δ38::HIS3* (right) initiation reporters with identified
mutations annotated. Phenotypes displayed have been scored and processed as in Fig. 2a.
Briefly, scored phenotypes have been corrected to WT on the same condition, followed by
normalization to the same on control conditions. Here, a positive value represents growth better
than WT and a negative value represents worse than WT (zero). **b.** Single spot phenotyping of
upstream isolated mutants. **c.** Cryo-EM structure of Pol II PIC annotated with upstream shifting
mutation residues identified (PDB: 7O4J).

**Figure 4. Mutations identified across PIC**. Lollipop diagram of mutations isolated in Rpb1 (**a**)
and Rpb2 (**b**). Domain diagram figure created using IBS 2.0 (XIE *et al.* 2022). **c**. PIC structure
(PDB: 7O4J) displaying mutations isolated in our study. **d.** Structural mutations surrounding
active cleft that are formamide sensitive marked in magenta. **e.** Mutations in α21 helix and
anchor loop isolated with proximity to Rpb9 and Rpb8. **f.** Residues at Rpb2-TFIIF-Rpb9
interface found to alter TSS selection. **g.** Initiation mutations near Tfg1-Tfg2 interface causing
downstream shifts. **h.** Mutations altering interactions between Sua7 B-core and Rpb2 isolated in
this study in the vicinity of previously isolated Sua7 C149 initiation mutant. **i.** Potential
interactions between Ssl1 and Tfb4 loop or Tfb1 in driving TSS selection. **j.** Rad3 K499E
mutation potentially driving TSS selection by altering interactions with Ssl2 across Rad3-Ssl2
interface with K357R and K357E previously isolated in our study altering TSS selection. **k.**
Mutations in Tfb2 F189 and Tfb4 N71 lying within the respective structural domains potentially
promoting structural "breathing". **l.** Mutation S23F in the RING domain of Tfb3 increasing
interaction between TFIIH and TFIIE by potentially promoting allele-specific hydrophobic
packing.

**Figure 5. Phenotyping of subset of non-PIC mutants identified as suppressors of initiation reporters. a.** Guanine metabolism mutants and controls plated as single concentrations of cells on phenotyping media. Guanine metabolism gene mutants confer the His⁺ phenotype to cells with the *imd2Δ::HIS3* reporter, likely due to the *IMD2* promoter's sensitivity to guanine levels. Guanine supplementation suppresses the His⁺ phenotype of guanine pathway mutants (SC-His 2 mM 3AT Gua/HCl: left panel) whereas PIC mutants remain His⁺ under the same conditions (SC-His 2 mM 3AT Gua/HCl: right panel). Guanine is solubilized in 10 mM NaOH at a final concentration of 500 µM and media must be neutralized by equivalent HCl as NaOH alone affects solubility of MPA. **b.** 10-fold serial dilutions of *PMR1* or *pmr1Δ* strains plated on phenotyping media. *pmr1Δ* suppresses both upstream reporters and confers sensitivity to Mn²⁺.

**Figure 6. Analysis of potential Rpb10-Tfg1 interactions in proper TSS selection. a.** Cryo-EM structures of PICs of Pol I, Pol II, and Pol II displaying Rpb10, a shared subunit between them. **b.** Cryo-EM structure of Rpb10 from Pol I-III PICs (Pol I PIC: 6TPS, Pol II PIC: 7O4J, Pol III PIC: 6EU0) overlayed displaying Tfg1 from the Pol II PIC, with Tfg1 R26 in the proximity of Rpb10 E29 suggesting a potential electrostatic interaction. While Rpa135 in Pol I lies adjacent to Rpb10, a specific physical interaction with Rpb10 E29 may not exist. **c.** Serially diluted spot assay phenotyping of single and double mutants of *tfg1* R26 and *rpb10* E29 resides displaying charge reversals, mutations to alanine and silent PAM control mutations (Rpb10 L30 AAC→AAT) as mutants were generated by CRISPR/Cas9.

1412

**Figure 7. Synthetic genetic interaction between *rpb9Δ* and *rpb2* G369C. a.** Cryo-EM
structure of Pol II PIC displaying Rpb2 G369 and Rpb9. Rpb2 G369 lies in the Pol II lobe
domain which interacts with Rpb9 though this region between Rpb2 and Rpb9 and has been
shown to interact with TFIIF. **b.** Tetrad dissection of *rpb9Δ/+ rpb2* G369C/+ diploid to examine
genetic interactions between them. Spores are annotated by genotype. **c.** Observed and
expected fitness of single and double mutants plotted for assessing genetic interaction.
"Expected" fitness for a double mutant is the product of the two single mutant fitnesses (MANI *et*
*al.* 2008). Observed fitness of single and double mutants estimated by quantification of spore
colony sizes. Observed fitness is worse than expected fitness indicating a synthetic genetic
interaction.

**Figure 8. STRIPE-seq analysis of novel mutants identifies three functional groups of**
**initiation mutants based on shift of median of TSS distribution. a.** Schematic of STRIPE-
seq library generation. **b.** Schematic of ∆median metric for TSS analysis. **c.** Principal
component analysis (PCA) of ∆median values of the top 2829 promoters based on an
expression cutoff of ≥10 reads within the analyzed 401 nucleotide promoter windows. PCA of
∆median values across mutants with combined biological replicates plotted along the first two
principal components display three clusters, ① chromatin mutants *spt5* G814*, *spn1* V199G, ②
*rpb1* A480S, *rpb10* L25S, *ssl1* I122T, *tfb4* N71K and ③ *rad3* K499E and *tfb2* F189S. **d.**
Hierarchical clustering of Spearman correlation of ∆median values across mutants. **e.**
Hierarchically clustered heatmap of ∆median values across top 2829 promoters of mutants with
replicates combined. **f.** Box-plot analysis of ∆median values across mutants presented by Tukey
plot.

**Figure 9. STRIPE-seq analysis of novel mutants identifies three functional groups of**
**initiation mutants based on shift of spread of TSS distribution. a.** Schematic of ∆spread
metric of analysis **b.** Principal component analysis of ∆spread values of top 2829 genes across
mutants plotted along the first two principal components displays three clusters for ∆median, ①
chromatin mutants *spt5* G814*, *spn1* V199G, ② *rpb1* A480S, *rpb10* L25S, *ssl1* I122T, *tfb4*
N71K and ③ *rad3* K499E and *tfb2* F189S. **c.** Hierarchical clustering of Spearman correlation of
∆spread values across mutants. **d.** Hierarchically clustered heatmap of ∆spread values across
top 2829 promoters of mutants. **e.** Box-plot analysis of ∆spread values across mutants
presented by Tukey plot.

1446
**Figure S1. Isolation of *HMO1* promoter variant that detects upstream TSS shifts. a.**
Schematic illustrating an *HMO1* promoter variant deletion series (left) with behavior of reporters
in cells in absence or presence of UV mutagenesis (right) to assess potential for His+ mutant
generation on the His- background, indicative of reporter suppression. **b.** Schematic of
functional *hmo1Δ38::HIS3* reporter. *HMO1* promoter sequence between –405 and –39 was
amplified and appended to the *HIS3* ORF followed by the *CYC1* terminator and integrated into a
gene free region of chromosome I (chrI:199270-199271).

1454

1455 **Figure S2. Initiation and secondary phenotypes of mutants isolated in genetic selections.**
1456 Single spot phenotyping of a subset of downstream-shifting mutants isolated in our study.

1457

1458 **Figure S3. Phenotypes of chromatin mutants isolated in the study.** Chromatin/Pol II
1459 elongation factor mutants identified for both downstream-shift sensitive reporters (subset
1460 shown).

1461

**Figure S4. Genetic interactions between upstream-shifting *rpb9*Δ and novel downstream-shifting alleles prioritized in this study.** Genetic interaction by tetrad analysis between *rpb9*Δ and downstream shifting novel alleles *tfb2* F189S, *tfb4* N71K, *ssl1* I122T across the top (left-right) and *rad3* K499E, *rpb10* L25S and *tfg2* C222Y across the bottom (left-right). Growth phenotypes quantified through spore colony size as in Figure 6. Suppression of *rpb9*Δ fitness defect by *tfb2* F189S, *tfb4* N71K, *ssl1* I122T and *rad3* K499E observed, while interactions with *rpb10* L25S and *tfg2* C222Y appear epistatic. Asterisks (*) indicate incomplete tetrads or failed dissections not considered.

**Figure S5. STRIPE-seq analysis of ∆median values across individual mutant replicates. a.**
Pearson correlation of individual TSS positions between samples across genome with cutoff per
position set at ≥ 3. **b.** Principal component analysis of ∆median values across top 2829 genes
qualifying expression cutoff of ≥10 total expression within 401 base analyzed promoter window
for individual sample replicates. **c.** Spearman correlation of ∆median of top 2829 genes across
all samples and replicates. **d.** Hierarchically clustered heatmap of ∆median values across all
promoters in individually sequenced samples. **e.** Box-plot analysis of these ∆median values
amongst all sequenced samples analyzed by Tukey's method.

**Figure S6. STRIPE-seq analysis of Δspread values across individual mutant replicates. a.** Principal component analysis of Δspread values across top 2829 genes qualifying expression cutoff of ≥10 within 401 base analyzed promoter window for individual sample replicates. **b.**

1484    Pearson correlation of $\Delta$spread of top 2829 genes across all samples and replicates. **c.**
1485    Hierarchically clustered heatmap of $\Delta$spread values across all promoters in individually
1486    sequenced samples **d.** Box-plot analysis of these $\Delta$spread values amongst all sequenced
1487    samples analyzed by Tukey's method.