Identifying the therapeutic potential of Niclosamide in overcoming IFN-gamma dependent cancer immune evasion in the Tumor Microenvironment

Yue Zhang[1], Sophia Peng[1,2], Hamid Akhbariyoon[1], Suzi Kim[1], Ellie Lai[1], En Cai[1*]

[1]Department of Biological Sciences, Carnegie Mellon University, 5000 Forbes Ave, Pittsburgh, PA 15213

[2]Department of Immunobiology, Yale University, New Haven, CT 06520

*Correspondence to: En Cai, Ph.D.

E-mail: encai@andrew.cmu.edu

Phone: 412-383-7976

Abstract

Background: Tumor cells frequently develop immune resistance through interferon-γ (IFN-γ)–induced PD-L1 expression, acquisition of cancer stem cell (CSC)–like features, and adaptation to hypoxia within the tumor microenvironment (TME). Although IFN-γ activates both STAT1 and STAT3, how these pathways interact to regulate immune evasion under hypoxia remains unclear.

Methods: Using the MC38 murine colorectal cancer model and T cell–tumor spheroid co-culture assays, we examined how IFN-γ signaling through STAT1 and STAT3 influences PD-L1 expression, CSC plasticity, and cytotoxic T cell function under normoxic and hypoxic conditions. Pharmacologic inhibitors and siRNA knockdown were used to dissect pathway function, and Niclosamide, an FDA-approved anthelmintic, was evaluated as a dual STAT1/STAT3 inhibitor.

Results: We found that IFN-γ primarily induced PD-L1 through STAT1 activation, while CSC plasticity was associated with STAT3 signaling. STAT1 and STAT3 displayed reciprocal regulation—blocking one enhanced activation of the other. Niclosamide effectively inhibited phosphorylation of both STAT1 and STAT3, which led to suppressed PD-L1 upregulation and reduced CSC enrichment. In addition, it also partially inhibited hypoxia-induced HIF-1α expression. In co-culture assays, Niclosamide improved T cell

infiltration and reduced exhaustion under hypoxic conditions, resulting in improved T cell killing.

Conclusions: Our findings identified Niclosamide as a potent dual STAT1/3 inhibitor capable of reversing IFN-γ and hypoxia-driven immune evasion. Repurposing Niclosamide may represent a promising strategy to enhance the efficacy of immune checkpoint blockade in solid tumors.

Keywords:

Tumor microenvironment, IFN-γ, PD-L1, STAT1, STAT3, Cancer Stem Cells, Niclosamide, tumor spheroids, Immune Evasion, Cancer Immunotherapy

key messages

Interferon-γ (IFN-γ) enhances cytotoxic T cell function but also promotes tumor immune evasion by upregulating PD-L1 and inducing cancer stem cell–like properties. Our study identifies a reciprocal regulatory mechanism between STAT1 and STAT3 in IFN-γ-treated tumor cells that shapes immune evasion outcomes. We demonstrate that Niclosamide, an FDA-approved anthelmintic, acts as a dual STAT1/STAT3 inhibitor, effectively suppressing PD-L1 induction, limiting cancer stemness, and reducing HIF-1α expression under hypoxia. Niclosamide also restores T cell infiltration and decreases exhaustion in a 3D tumor spheroid model. By repurposing Niclosamide, this work provides a feasible approach to enhance the efficacy of immune checkpoint blockade and guide future translational and clinical studies in immunotherapies against solid tumors.

Background

Over the past decade, cancer immunotherapy, which aims to restore immune function within the tumor microenvironment (TME), has achieved remarkable success in cancer treatment[1,2]. Therapies like immune checkpoint blockade (ICB) have demonstrated durable responses in some patients with advanced malignancies. Nevertheless, only a subset of patients (around 30%) with certain solid tumors benefits from ICBs, while most patients exhibit minimal response to these treatments[3–5]. Understanding the mechanisms of immune evasion and identifying new therapeutic targets remain critical for improving treatment outcomes.

Tumor cells employ multiple strategies to escape immune surveillance. One major mechanism is programmed cell death ligand 1 (PD-L1) upregulation, which inhibits cytotoxic T cell (CTL) function [6,7]. Another mechanism is the development of cancer stem cell (CSC)–like properties, which confer resistance to therapy and drive recurrence[8,9]. In addition, hypoxia is a hallmark of the TME that profoundly reshapes cellular metabolism, promotes PD-L1 expression via hypoxia-inducible factors (HIFs), and suppresses CTL and NK cell functions [10].

During immune responses, T cells release type II interferon (IFN-γ), which is a glycosylated protein that facilitates tumor rejection by modulating systemic anti-tumor immunity[11,12]. However, studies reveal a dichotomous nature of

IFN-γ: while it enhances immune functions, it also helps cancer cells evade immune attacks. Specifically, IFN-γ stimulation has been shown to increase cytoplasmic expression of PD-L1[13,14] and elevate cancer cell stemness[15,16]. Both the PD1/PD-L1 axis and CSCs play crucial roles in enabling tumor cells to escape anti-tumor immunity across various cancers[17–19]. IFN-γ activates downstream Janus kinas/signal transducer and activator of transcription (JAK/STAT) signaling, primarily STAT1 and STAT3, which regulate overlapping gene networks involved in immune evasion [20–22]. Therefore, there is an urgent need for therapeutic strategies that inhibit IFN-γ's effects on cancer immune evasion while preserving its role in facilitating CTL and natural killer (NK) cells in eliminating tumors.

Here, we examined the fine-tuning of the IFN/JAK/STAT pathway under normal and hypoxia condition, and identified the specific STAT pathway that drive PD-L1 upregulation and CSC transformation, respectively. Using a mouse colon cancer model, our study reveals a reciprocal relationship between STAT1 and STAT3 activities in cancer cells (MC38). Specifically, inhibition of STAT1 signaling leads to a significant enhancement in STAT3 activation, and conversely, blocking STAT3 activity results in increased STAT1 signaling. Both transcription factors, STAT1 and STAT3, regulate overlapping sets of target genes, many of which play crucial roles in immune evasion mechanisms. This dynamic interplay suggests that the balance between STAT1 and STAT3 could be a critical factor in shaping the immune

microenvironment within tumors, potentially influencing cancer progression and resistance to immune-mediated therapies. Furthermore, in evaluating pharmacological methods to block STAT1/STAT3 activities, we identified an anthelmintic drug, Niclosamide, which inhibits the activity of both STAT1 and STAT3, and partially blocks the upregulation of hypoxia-inducible factor 1α (HIF-1α) protein induced by hypoxia. The co-culture study of primary murine T cells and 3D tumor spheroids model further confirmed that Niclosamide could enhance T cell infiltration and reduce T cell exhaustion under a hypoxic TME. Together, our study reveals Niclosamide as a promising candidate for repurposing in cancer immunotherapy to overcome IFN-γ and hypoxia-induced immune evasion.

Methods

Cell Culture and Compounds: MC38 and MC38-OVA cell lines were gifts from Robert Eil lab at OHSU. Both MC38 and MC38-OVA cell line were maintained in DMEM Medium (Gibco, Gaithersburg, MD, USA), supplemented with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin. Primary OT1 mouse T cells were prepared from lymph nodes and spleens of OT-I TCR transgenic mice (6-10 weeks old), and were maintained in RPMI-1640 Medium (Gibco, Gaithersburg, MD, USA), supplemented with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin and 50 μ M β - mercaptoethanol (complete RPMI). Splenocytes were incubated in complete

RPMI with 100 ng/mL SIINFEKL peptide for 30 minutes at 37°C and then washed three times. Lymphocytes and splenocytes were then mixed at 1:1 ratio at 2 million cells/mL in complete RPMI and maintained at 37°C 2% CO_2. Interleukin-2 (HIL2-RO, Roche) were added to the cell culture on day 2 at a final concentration of 10 U/mL. Cells were replenished with fresh media and IL-2 every 2 days and used from day 4 to day 7. Mouse IFN gamma Recombinant Protein were purchased from MilliporeSigma (Sigma-Aldrich, USA). STAT1 inhibitor Fludarabine (F-ara-A, NSC 118218) (MedChemExpress, Cat. No.: HY-B0069), STAT3 inhibitor Stattic (MedChemExpress, Cat. No.: HY-13818), and Niclosamide (BAY2353, MedChemExpress, Cat. No.: HY-B0497) were purchased from MedChemExpress (Monmouth Junction, NJ, USA).

For cell cultured under hypoxia condition, cells were maintained in the Heracell™ VIOS 160i Tri-Gas CO2 Incubator from Thermo Fisher Scientific (Waltham, MA), and nitrogen gas were connected to the incubator to make the oxygen concentration down to 1.5% inside the chamber. For normoxia condition, the oxygen concentration is 20%, as detected by oxygen sensor.

Mice: C57BL/6-Tg (TcraTcrb)1100Mjb/J (OT1) mice were purchased from Jackson Laboratories (Bar Harbor, ME, USA) and were housed and bred at the Carnegie Mellon University, according to Laboratory Animal Resource Center guidelines. The protocol was approved by the Institutional Animal Care and Use Committee (IACUC) at the Carnegie Mellon University.

Formation of co-culture tumor spheroids: MC38OVA (or MC38) tumor spheroids were generated by seeding $1x10^4$ cells per well on Costar ultra-low attachment (Corning) round bottom 96 wells plates in the 3D Tumorsphere Medium XF (PromoCell). 7 days later, $1x10^5$ total or CD8 sorted mouse primary T cells were labeled with ViaFluor® 650 SE Cell Proliferation Dye, then added with the spheroids to perform co-culture. Spheroids were gently resuspended and left to sediment to the bottom of the Eppendorf tube. These steps were repeated 2 times with PBS in order to wash the spheroids from the non-infiltrating immune cells. Spheroids were then break down mechanically to obtain a single cell suspension for further analyzed by flow cytometry.

Quantitative real-time PCR: Total RNAs were isolated from cell lysate or mouse tissue with TRIzol reagent (Invitrogen Life Technologies, Carlsbad, CA, USA). The expression levels of multiple mRNAs were detected by quantitative real-time PCR using the High-Capacity cDNA Reverse Transcription Kit (Thermo Fisher Scientific, Waltham, MA, US) and the LightCycler® 480

System (Roche, Indianapolis, IN, USA). Primers used for Quantitative RT-PCR were shown in **Supplemental Table 1**.

Western Blot: Total proteins were extracted from cultured cells with radio-immunoprecipitation assay (RIPA) buffer (Thermo Fisher Scientific, Waltham, MA, USA) plus fresh protease and phosphatase inhibitors (Roche, Indianapolis, IN, USA). Proteins (10-20µg) were loaded to each lane and separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis. Proteins were then transferred to a 0.45 µm pore size PDVF membrane (Bio-Rad, Hercules, CA, US), followed by immunodetection of target proteins using specific antibodies with chemiluminescent detection. The developed films were quantified with ImageJ software. The antibodies used in Western blot were listed in **Supplemental Table 2**.

Flow study: Cell suspensions of MC38 or primary T cells were incubated with the different combinations of antibodies (for MC38: PD-L1, CD133, CD44, B7H4, For T cells: CD45, CD8, PD1, Tim3) on ice for 30 minutes. Then the samples were washed and fixed after cell surface staining according to the manufacturer's description in fixative buffer(eBioscience). All antibodies were purchased from BD Biosciences and flow data were collected on a BD Accuri C6 Plus Flow Cytometer (BD Biosciences). The antibodies used in flow study were also listed in **Supplemental Table 2**.

The data were analyzed using the FlowJo software (Tree Star Inc.).

Immunofluorescence: For immunofluorescence, cultured cells or tumor spheres were rinsed with 0.1% TX-100 in PBS (PBST) three times, blocked with 1% goat serum and 2% BSA in PBS at room temperature for 1 h and then incubated with the primary antibody conjugated with fluorescence. Then the slides were washed 3 times with PBST, and mounted with Fluoromount-G™ Mounting Medium (Invitrogen™). Fluorescent images were captured with All-in-One Fluorescence Microscope BZ-X800 (Keyence, Ōsaka, Japan). The primary antibodies used in the immunofluorescence were shown in **Supplemental Table 2**.

RNA sequencing (RNA-seq)

RNA-seq of IFN-γ treated MC38 cells was performed by Innomics Inc. (One Broadway, 14th FL, MA) using DNBSEQ Eukaryotic Strand-specific Transcriptome Resequencing. Alignment of raw sequencing reads was performed using Hisat2 software. The DNBSEQ package was used to find differentially expressed genes, and differentially expressed genes with $P <$ 0.05 were used to perform cluster analysis and enrichment analysis with ClusterProfiler package in the R program.

Statistics: Data were presented as mean ± SEM. Statistical analysis was performed with GraphPad Prism 10.0 software (GraphPad Software, San Diego, CA). Comparison between experimental variables was performed

using one-way or two-way ANOVA followed by a Tukey post hoc test.

Significance levels were denoted as *p <0.05, **p < 0.01, or ***p <0.001.

Results

IFN-γ induced PD-L1 expression through STAT1 pathway

In this study, we used the mouse colon cancer cell line MC38 as a model to study the effects of IFN-γ treatment. Consistent with previous literature reports, we found a significantly increase in PD-L1 expression on MC38 cells following treatment with IFN-γ, as shown by both flow cytometry (Fig.1A) and immunofluorescent staining images (Fig.1B-C). The up-regulation of cell surface PD-L1 demonstrated a dose and time-dependent manner (Supplemental Figure 1). In addition, PD-L1 upregulation was also observed in induced MC38 tumor spheres, which exhibited more CSC features (increased population with CD44+CD133+ expression) compared to adherent MC38 cells (Supplemental Figure 2A-B).

IFN-γ signals through STAT1 and STAT3 [23,24]. However, the primary signal transducer responsible for IFN-γ dependent PD-L1 upregulation varies between different tumor cell lines[25]. To evaluate the potential roles of STAT1 and/or STAT3 activation in mediating PD-L1 protein expression in MC38, we used small interfering RNA (siRNA) to knock down STAT1 and STAT3 and examined their distinct contributions to IFN-γ-induced PD-L1 expression. Flow cytometry assays confirmed that IFN-γ-induced PD-L1 expression was diminished when STAT1 was knocked down, while STAT3

knocking-down have an opposite effect (Fig.1D). A similar effect was observed in MC38 cells that were pre-transfected with siRNAs and then induced into tumor spheres (Supplemental Figure 2C).

We performed real-time PCR to assess siRNA knock down level and target gene expression level. The results indicate an efficient knock down of STAT1 (Fig.1E) and STAT3 (Fig.1F) by siRNA. PD-L1, and classical STAT1 transcriptional targets including CXCL10, IL-15 was upregulated in response to IFN-γ treatment, and the up-regulation was diminished with the siRNA of STAT1, but not siSTAT3 (Fig.1G-I). In comparison, the mRNA expression level of myeloid cell leukemia-1 MCL-1, which is a transcriptional target of STAT3, was down-regulated by IFN-γ treatment, and increased while treated with siRNA of STAT1 (Fig.1J). The western blot data suggest that the phosphorylated STAT1 (pSTAT1), which is the activate form, and phosphorylated STAT3 (pSTAT3) were both enhanced by IFN-γ, and the induced pSTAT1 is much stronger compared to pSTAT3 at higher IFN-γ concentrations. Also, IFN-γ mediated upregulation of PD-L1 in a dose-related manner (Fig. 1K). Knocking-down of STAT1 with siRNA (siSTAT1) partially blocks the up-regulation of PD-L1 induced by IFN-γ. Notably, this also increases pSTAT3 induced by IFN-γ compared to that in the scramble siRNA control (siCtrl) sample (Fig. 1L). Conversely, the siRNA of STAT3 (siSTAT3) further enhances the increase in PD-L1 expression and phosphorylation of STAT1 when combined with IFN-γ (Fig.1L). These results suggest competitive

interactions between pSTAT1 and pSTAT3, since these two transcriptional factors compete for similar kinases and promotor sites[25].

In a T cell–tumor cell co-culture system, we found treatment with 5.0 ng/mL IFN-γ enhanced the survival of MC38 cells when co-cultured with primary mouse T cells at a 1:1 ratio, and this effect was even more pronounced in tumor spheroids, which exhibited increased resistance to T cell-mediated killing (Supplemental Fig. 2D). Furthermore, MC38 cells pre-treated with 5.0 ng/mL IFN-γ showed markedly increased resistance to T cell cytotoxicity during short-term co-culture (30 min to 4 h) (Supplemental Fig. 3A). Consistently, colon cancer cells exposed to low concentrations of IFN-γ (0.5–5 ng/mL) developed enhanced resistance to T cell killing (at a 1:3 T cell-to-tumor ratio), whereas higher IFN-γ concentrations (10–100 ng/mL) led to increased susceptibility (Supplemental Fig. 3B). All these data suggest that the tumor killing function of T cells is impaired through the overexpression of PD-L1 on MC38 cells.

IFN-γ regulate cancer cell stemness indirectly by manipulating pSTAT1/pSTAT3 competitive activity

Parallel to our study on IFN-γ-dependent PD-L1 expression in MC38 cells, we seek to investigate the impact of IFN-γ on mitosis and cell cycle. A recent study using mouse models of breast cancer showed that IFN-γ produced by activated T cells can directly convert non-CSCs to CSCs[16]. However, it remains unclear whether this effect is specific to breast cancer models or

could be observed across multiple cancer types. Based on the theory that type I interferons (IFNs-I) promote an adaptive yet reversible transcriptional rewiring of cancer cells towards stemness and immune escape[15], we hypothesize that IFN-γ may also induce stemness in MC38 cancer cells.

To test this hypothesis, we generated colon cancer spheroids from MC38 cells. We first used ki67 to label the active proliferating cells in mix cell population. To our surprise, we found that IFN-γ treatment reduced the number of ki67 positive cells in a MC38 tumor spheroids model (Fig. 2A-C). We also found the sphere-forming unit (SFU) was significantly repressed by IFN-γ pre-treatment (Fig. 2D-E). This reduction of SFU can be neutralized by STAT1 siRNA but not by STAT3 siRNA (Fig. 2F). We then tested cancer stem cell surface markers, including CD44 and CD133, and observed a downregulation of both $CD44^{hi}CD133^{+}$ and $CD44^{hi}CD133^{-}$ populations in MC38 cells treated with IFN-γ. Both populations, previously reported to represent colorectal cancer stem cells [26–29], were significantly reduced following IFN-γ treatment for control siRNA (Fig.2G). However, when the cells were first treated with siSTAT1 to knock down STAT1, this reduction was abolished. In contrast, knocking down STAT3 did not produce a similar effect (Fig.2G). The results from real-time PCR demonstrated that IFN-γ can induce CSC markers like CD44, CD133, OCT4, and SOX2 only when tumor cells were transfected with siSTAT1 (Fig.2H-K). This suggests that when STAT1 were inhibited, the mRNA of stemness-related genes was up regulated.

Western blot results showed that the protein levels of CSC markers, including SOX2 and Nanog, were reduced by IFN-γ treatment under normal condition, but remain the same or higher level with STAT1 knocking down (Fig.2L), which is consistent with our flow cytometry results. This suggests that STAT1 may have suppression role for these CSC-related genes. When MC38 cells were transfected with siSTAT3, both the mRNA and protein of the CSC markers were significantly reduced (Fig.2H-L), suggesting STAT3 is playing an important role for self-renewal of tumor cells. Given that these CSC markers are all strongly regulated by STAT3[30,31], and our data show that knocking down STAT1 increases expression of these genes with IFN-treatment, our data suggests a potential competitive antagonism between pSTAT1 and pSTAT3 as signal transducers and transcriptional factors. We conclude that the decrease of CSCs triggered by IFN-γ is primarily due to the dominant activation of JAK/STAT1 pathway, which indirectly suppress STAT3 signaling. Notably, when STAT1 activity is inhibited, IFN-γ could facilitate the transformation to CSCs through the compensate activation of STAT3 signaling. Therefore, single inhibition of individual STAT1 or STAT3 signaling pathway activated by IFN-γ could not prevent immune evasion. We need to find a more effective inhibition of both pathways, to prevent the over-expression of PD-L1 on tumor cells while also interfering with the development of CSCs.

Niclosamide inhibits both STAT1 and STAT3 activity, thus reducing PD-L1 up-regulation induced by IFN-γ

To further investigate the role of IFN-γ in TME, we use small molecule STAT1 inhibitor Fludarabine (F-ara-A, NSC 118218) (MedChemExpress, Cat. No.: HY-B0069) and STAT3 inhibitor Niclosamide (BAY2353, MedChemExpress, Cat. No.: HY-B0497). Fludarabine works by inhibiting the cytokine-induced activation of STAT1 and STAT1-dependent gene transcription in normal resting or activated lymphocytes [32,33]. Niclosamide is reported as a STAT3 inhibitor with an IC50 of 0.25 μM in HeLa cells [34]. Both reagents were used in our experiments to block STAT1 and STAT3 transcriptional activity respectively.

The western blot result indicates that 5μM Fludarabine did not have a sufficient inhibition effect to STAT1 phosphorylation in MC38 cells. In comparison, 0.5μM Niclosamide could disrupt STAT1 protein phosphorylation while also inhibit STAT3 protein expression and activity (Fig.3A). In flow-cytometer assays, Niclosamide partially blocked the up-regulation of cell surface PD-L1 induced by IFN-γ (Fig.3B), considering Niclosamide is an STAT3 inhibitor, this effect is opposite compared to STAT3 siRNA (Fig.1D). When low dose of IFN-γ (5ng/ml) were used to increase tumor cell viability in a T cell killing assay, Niclosamide could neutralize this effect while Fludarabine could not (Fig.3C). Meanwhile, Niclosamide did not affect the effectiveness of IFN-γ in reducing SFU (Fig.3D) (Supplemental Figure 4 A-B),

and reducing the population of CD44^{hi}CD133^{+}/ CD44^{hi}CD133^{-} cell population in MC38 induced tumor spheroids (Fig. 3E). Niclosamide also reduce PD-L1 expression on the cell surface of these tumor spheroids (Supplemental Figure 4C). In western blot assay, Niclosamide did not affect the downregulation of Nanog, SOX2 proteins by IFN-γ (Fig. 3F). These data suggested Niclosamide could disrupt both STAT1 and STAT3 phosphorylation in IFN-γ signaling, and could become a potential method to prevent cancer cell's immune evasion in combination use with immune-checkpoint blockers.

Delineate the signal network in inflammatory TME under hypoxia condition.

Hypoxia condition in TME activated a complex signaling network mediated by HIFs, which are closely associated with the JAK/STAT signaling pathways. The hypoxic environment in the solid tumor is well known to mediates resistance to chemotherapy, radiotherapy, and immunotherapy through complex mechanisms[17]. To delineate the changes in multiple signaling pathways during IFN-γ treatment under hypoxia conditions, we sequenced total RNA isolated from the following 4 samples: *Ctrl_1_1*: MC38 cells under normal condition, without IFN-γ; *IFN_1_2*: MC38 cells treated with IFN-γ, under normal condition; *Ctrl_2_1*: MC38 cells under hypoxia condition, without IFN-γ; *IFN_2_2*: MC38 cells treated with IFN-γ, under hypoxia condition. The total RNA-sequencing analysis provided insights into the transcriptional landscape of cancer cells that has been modulated by the IFN-

γ and hypoxia stimulation (Fig. 4A-B). Among all the differentially expressed genes (DEGs), we validate a serious of genes related to cancer progression and immune response with real-time PCR. One of these genes is CD274 (PD-L1), which is significantly up-regulated by IFN-γ alone and further enhanced under hypoxia condition (Fig. 4D). STAT1 and its target genes were all up-regulated by IFN-γ in both conditions (Fig. 4C-F). Notably, the induced CXCL10 (also known as Interferon gamma-induced protein 10, IP-10) level is reduced under hypoxia condition compared to normal condition (Fig. 4E), consistent with literature reports that in certain cancer cells, such as colorectal, breast, ovarian, and lung cancer cells, prolonged hypoxia reduces CXCL10 expression, which is linked to the activity of the HIF-1α protein[35]. According to quantitate PCR results, STAT3 mRNA is also up-regulated by IFN-γ, but its target genes including HIF1α, and some stem cell marker such as SOX2 and OCT-4 mRNA were down-regulated under hypoxia condition (Fig. 4G-N). In general, the RNA-seq results further confirms the role of JAK-STAT signaling in regulating potential genes that mediate cancer immune evasion under hypoxia condition, such as immune-check point blockers (CD274) and chemo attractors (CXCL10).

Hypoxia enhances IFN-γ induced PD-L1 upregulation, which is also blocked by Niclosamide

To further investigate the contribution of hypoxia to immune evasion, we performed western blot analysis of varies protein under both conditions. We

found that both PD-L1 and pSTAT1 were up-regulated more significantly in hypoxia condition compared to normal condition (Fig.5A). More importantly, Niclosamide significantly neutralized the induction of HIF1α under hypoxia condition, while preserve its ability to block both STAT1 and STAT3 phosphorylation and protein levels (Fig.5B). The MC38 cells survive better with higher IFN-γ concentration in hypoxia condition compared to normal condition in a T cell-tumor cell co-culture assay (Fig.5C). In a dose-dependent cell viability assay, we found Niclosamide show more toxicity to tumor cells under hypoxia condition (Fig.5D). These findings further suggests that Niclosamide could be used as an efficient reagent to eliminate IFN-γ-induced immune resistance.

IFN-γ induce T cell exhaustion with hypoxia and enhance T cell infiltration in the TME.

Given that STATs are also key molecular determinants of T-cell fate and effector function[36] , we want to further study the recruitment and exhaustion progress of T cells in an immunosuppressive tumor microenvironment. We use tumor spheroids induced by MC38-ova cells (transgenic MC38 cell line with stably expressing **Ovalbumin**) and a T cell-tumor cell co-culture system, to mimic a simplified TME[37] (Fig. 6A). Using the established protocol of *in vitro* T cell exhaustion[38], we were able to induce mouse primary CD8+ T cells to an early or late exhausted state (Supplemental Figure 5A). With the exhausted T marker like PD-1, Tim-3, we found that both early state

exhaustion (PD-1+/Tim-3-) and late state exhaustion (PD-1+/Tim-3+) T cells were significantly increased when co-cultured with MC38ova cells that pre-treated with IFN-γ, and under hypoxia condition (Fig. 6B). Meantime, the MC38ova cells also show more induced PD-L1 expression under hypoxia condition (Fig. 6C). We found IFN-γ could promote CD8+ T cells infiltration into the tumor spheres significantly, as shown by CD8+ T cells prelabeled with ViaFluor 650 (VF650) (Fig. 6D-E). We hypothesize this effect is due to CXCL10, a powerful chemokine that acts by binding to its specific receptor, CXCR3, to attract immune cells such as T cells and macrophages to the sites of inflammation. Under hypoxia condition, the T cell infiltration with IFN-γ stimulation is reduced moderately (Fig. 6D-E), which is consistent to our finding that CXCL10 expression is also less induced under hypoxia condition (Fig. 4E). In addition, ViaFluor® 650 SE dye (**Biotium**, Fremont, CA) is a membrane-permeant compound that is converted to fluorescent dye by intracellular esterases and will covalently reacts with amine groups on intracellular protein at the same time, forming fluorescent conjugates that are retained in the cell. With each cell division, daughter cells inherit roughly half of the fluorescent label, allowing the number of cell divisions that occur after labeling to be detected by the appearance of successively dimmer fluorescent peaks on a flow cytometry histogram. In our experiment, the T cell labeled with viaFluor650 show two peaks in histogram under normal condition, but

only one peak under hypoxia condition (Fig. 6F), indicating the T cell proliferation is generally reduced.

In another co-culture assay, when the MC38ova cells were pre-treated with both IFN-γ and Niclosamide, the late state exhausted T cell (PD1+Tim3+) population was less compared to IFN-γ treated only condition (Fig.7A). In the T cell-tumor spheres co-culture model, although Niclosamide did not change IFN-γ induced CD8+ T cell infiltration under normal condition (Fig.7B-C), but when IFN-γ's effect was reduced under hypoxia condition, Niclosamide could partially rescue T cells' infiltration (Fig.7D). Also, compared to free CD8+ T cells labeled with ViaFluor650, which divided multiple times in 24hours (D0-D3), the CD8+ T cells get infiltrated into tumor spheroids show only one time division, and IFN-γ increase the subpopulation of divided T cells (D1).Niclosamide did not impair the population of total infiltrated CD8+ T cells (CD8+ Infil), nor the population of proliferated T cells that got infiltrated (CD8+ proli). (Supplemental Figure 5B-C) Adding together, the inhibition of both STAT1/STAT3 pathway with Niclosamide will not only reduce PD-L1 overexpression induced by IFN-γ, but also rescue the hypoxia induced T cell's exhaustion and less infiltration related to T cell function and metabolism changes. We anticipate that the combination use of IFN-γ plus Niclosamide in colorectal cancer treatment could reach a better therapeutical effect in slowing down the growth of tumors.

Discussion

Both hypoxia and consistent inflammation are considered hallmarks of solid tumor development[1,39]. Both conditions drive profound changes in the tumor microenvironment (TME), including metabolic reprogramming, immune suppression, and resistance to therapy [17,18]. Hypoxia impairs immune cell infiltration, promotes regulatory T cells (Tregs) and myeloid-derived suppressor cells (MDSCs), and upregulates PD-L1 expression in tumor and stromal cells [17,18]. These adaptations converge on central signaling hubs such as HIF-1α, STATs, NF-κB, and PI3K-mTOR, which collectively regulate tumor survival and immune evasion [18,40]. Meanwhile, inflammatory cells like tumor-associated macrophages (TAMs) contribute to abnormal, leaky vasculature, exacerbating the hypoxic conditions in tumors[17,41,42]. These findings triggered our great interest since IFN-γ and other cytokines secreted by tumor infiltrated lymphocytes (TILs) also activate the STAT-JAK signaling pathway, and these molecules may have synergistic effect combined with hypoxia condition[21,40].

In this study, we identified a critical interplay between IFN-γ signaling and hypoxia in shaping immune evasion mechanisms. Specifically, we demonstrated that STAT1 mediates IFN-γ-induced PD-L1 expression, while STAT3 promotes cancer stemness, with reciprocal regulation between the two pathways (Fig.1-2). Although IFN-γ could significantly induce PD-L1 up-regulation in the tumor cells, disturbance on the IFN-γ signaling could also

result in multiple mechanisms of compensation. Additionally, tumor cells use other strategies to evade immune surveillance and T cells attacking. Importantly, selective inhibition of STAT1 or STAT3 alone enhanced compensatory activation of the other pathway, suggesting that therapeutic strategies targeting only one branch may be insufficient.

We then demonstrated that Niclosamide, an FDA-approved anthelmintic, acts as a dual inhibitor of STAT1 and STAT3. Unlike selective inhibitors, Niclosamide simultaneously suppressed PD-L1 upregulation, reduced CSC enrichment, and partially blocked HIF-1α induction under hypoxia (Fig.3, Fig.5). In functional assays, Niclosamide restored T cell cytotoxicity, enhanced infiltration, and reduced exhaustion within a hypoxic TME (Fig.8). Niclosamide is orally bioavailable, inexpensive, and has a well-established safety profile from decades of clinical use against parasitic infections. These properties make it an attractive candidate for rapid translation as an adjuvant to immune checkpoint blockade (ICB). By simultaneously targeting STAT1/3 and hypoxia-driven signaling, Niclosamide may broaden the efficacy of ICB in patient populations that currently exhibit poor response rates. These findings position Niclosamide as a promising repurposed drug candidate for overcoming immune resistance in solid tumors.

Our work also clarifies the paradoxical role of IFN-γ in tumor immunity. While IFN-γ enhances T cell infiltration, it concurrently promotes PD-L1 expression

and T cell exhaustion, particularly under hypoxia. This duality highlights the complexity of cytokine signaling in the TME and explains why modulating IFN-γ signaling must be approached with caution. However, our study primarily used *in vitro* and *ex vivo* systems, including MC38 spheroids and T cell co-cultures. While these models capture key features of the TME, *in vivo* validation will be essential to confirm therapeutic efficacy and safety. Moreover, the exact molecular mechanism by which Niclosamide inhibits STAT1 and STAT3 remains to be fully defined and warrants further investigation.

Conclusion

In general, our study evaluates the effectiveness of targeting STAT1 and STAT3 molecules involved in the IFN-γ pathways to determine which pathway modulator can improve the efficacy of cancer treatments, particularly immunotherapy. Especially, our findings establish Niclosamide as a dual STAT1/3 inhibitor that disrupts IFN-γ- and hypoxia-mediated immune evasion. By restoring T cell function and limiting tumor plasticity, Niclosamide may represent a versatile repurposed agent for combination immunotherapy. This work provides a rationale for preclinical in vivo studies and future clinical trials to evaluate Niclosamide as an adjuvant strategy in refractory solid tumors.

Acknowledgments: This work was supported by institutional start-up funds

provided to Dr. En Cai by the Department of Biological Science, Carnegie

Mellon University.

We thank Dr. Eil and Robert Eil lab at School of Medicine, Ohio State

University for kindly providing the MC38 and MC38-ova cell line.

Author Contributions:

Study conception and design: YZ, EC; data collection: YZ, SP, HA, SK, EL;
analysis and interpretation of results: YZ, EC; draft manuscript preparation:
YZ, EC.

Disclosure

The authors declare no competing financial interests.

List of Abbreviations

PD-L1   Programmed Cell Death 1 Ligand 1

CSCs   Cancer Stem Cells

IFN-γ   Interferon-gamma

STAT1 Signal Transducer Activator of Transcription 1

STAT3 Signal Transducer Activator of Transcription 3

JAK Janus kinase

TME Tumor Micro-Environment

ICB Immune Checkpoint Blockade

CTL Cytotoxic T cells

PD1 Programmed Cell Death 1

HIFs Hypoxia-inducible Factors

NK Natural Killer Cells

TAMs Tumor-associated Macrophages

TILs Tumor Infiltrated Lymphocytes

References

1. Munn DH, Mellor AL, Regents G, Place F. IDO in the Tumor Microenvironment: Inflammation, Counter-regulation and Tolerance. Trends immunology. 2017;37:193–207. https://doi.org/10.1016/j.it.2016.01.002.IDO

2. Chen Z, Han F, Du Y, Shi H, Zhou W. Hypoxic microenvironment in cancer: molecular mechanisms and therapeutic interventions. Signal Transduct Target Ther. Springer Nature; 2023. https://doi.org/10.1038/s41392-023-01332-8

3. Liu Z, Xiang Y, Zheng Y, Kang X. Advancing immune checkpoint blockade in colorectal cancer therapy with nanotechnology. Front Immunol. Frontiers Media S.A.; 2022. https://doi.org/10.3389/fimmu.2022.1027124

4. Byun JK, Park M, Lee S, Yun JW, Lee J, Kim JS, et al. Inhibition of Glutamine Utilization Synergizes with Immune Checkpoint Inhibitor to Promote Antitumor Immunity. Mol Cell [Internet]. Elsevier Inc.; 2020;80:592-606.e8. https://doi.org/10.1016/j.molcel.2020.10.015

5. Peng Q, Qiu X, Zhang Z, Zhang S, Zhang Y, Liang Y, et al. PD-L1 on dendritic cells attenuates T cell activation and regulates response to immune checkpoint blockade. Nat Commun [Internet]. Springer US; 2020;11:1–8. https://doi.org/10.1038/s41467-020-18570-x

6. Lau J, Cheung J, Navarro A, Lianoglou S, Haley B, Totpal K, et al. Tumour and host cell PD-L1 is required to mediate suppression of anti-tumour immunity in mice. Nat Commun. Nature Publishing Group; 2017;8. https://doi.org/10.1038/ncomms14572

7. Juneja VR, McGuire KA, Manguso RT, LaFleur MW, Collins N, Nicholas Haining W, et al. PD-L1 on tumor cells is sufficient for immune evasion in immunogenic tumors and inhibits CD8 T cell cytotoxicity. Journal of Experimental Medicine. 2017;214:895–904. https://doi.org/10.1084/jem.20160801

8. Sancho P, Burgos-Ramos E, Tavera A, Bou Kheir T, Jagust P, Schoenhals M, et al. MYC/PGC-1α balance determines the metabolic phenotype and plasticity of pancreatic cancer stem cells. Cell Metab. 2015;22:590–605. https://doi.org/10.1016/j.cmet.2015.08.015

9. Kim JH, Lee KJ, Seo Y, Kwon JH, Yoon JP, Kang JY, et al. Effects of metformin on colorectal cancer stem cells depend on alterations in glutamine metabolism. Sci Rep. Springer US; 2018;8:1–13. https://doi.org/10.1038/s41598-017-18762-4

10. Li Y, Zhao L, Li XF. Hypoxia and the Tumor Microenvironment. Technol Cancer Res Treat. SAGE Publications Inc.; 2021. https://doi.org/10.1177/15330338211036304

11. Qian J, Wang C, Wang B, Yang J, Wang Y, Luo F, et al. The IFN-γ/PD-L1 axis between T cells and tumor microenvironment: Hints for glioma anti-PD-1/PD-L1 therapy. J Neuroinflammation. Journal of Neuroinflammation; 2018;15:1–13. https://doi.org/10.1186/s12974-018-1330-2

12. Mazet JM, Mahale JN, Tong O, Watson RA, Lechuga☐Vieco AV, Pirgova G, et al. IFNγ signaling in cytotoxic T cells restricts anti-tumor responses by inhibiting the maintenance and diversity of intra-tumoral stem-like T cells. Nat Commun. Springer US; 2023;14:1–21. https://doi.org/10.1038/s41467-023-35948-9

13. Imai Y, Chiba T, Kondo T, Kanzaki H, Kanayama K, Ao J, et al. Interferon-γ induced PD-L1 expression and soluble PD-L1 production in gastric cancer. Oncol Lett. 2020;20:2161–8. https://doi.org/10.3892/ol.2020.11757

14. Chen S, Crabill GA, Pritchard TS, McMiller TL, Wei P, Pardoll DM, et al. Mechanisms regulating PD-L1 expression on tumor and immune cells. J Immunother Cancer. Journal for ImmunoTherapy of Cancer; 2019;7:1–12. https://doi.org/10.1186/s40425-019-0770-2

15. Musella M, Guarracino A, Manduca N, Galassi C, Ruggiero E, Potenza A, et al. Type I IFNs promote cancer cell stemness by triggering the epigenetic regulator KDM1B. Nat Immunol. Springer US; 2022;23:1379–92. https://doi.org/10.1038/s41590-022-01290-3

16. Beziaud L, Young CM, Alonso AM, Norkin M, Minafra AR, Huelsken J. IFNγ-induced stem-like state of cancer cells as a driver of metastatic progression following immunotherapy. Cell Stem Cell. The Authors; 2023;30:818-831.e6. https://doi.org/10.1016/j.stem.2023.05.007

17. Vito A, El-Sayes N, Mossman K. Hypoxia-Driven Immune Escape in the Tumor Microenvironment. Cells. 2020;9:1–20. https://doi.org/10.3390/cells9040992

18. Wen Q, Han T, Wang Z, Jiang S. Role and mechanism of programmed death-ligand 1 in hypoxia-induced liver cancer immune escape (Review). Oncol Lett. 2020;19:2595–601. https://doi.org/10.3892/ol.2020.11369

19. Taube JM, Anders RA, Young GD, Xu H, Sharma R, McMiller TL, et al. Colocalization of inflammatory response with B7-H1 expression in human melanocytic lesions supports an adaptive resistance mechanism of immune escape. Sci Transl Med. 2012;4. https://doi.org/10.1126/scitranslmed.3003689

20. Tang et al. 2005, Piomelli Daniele 2013. STAT3 and HIF1α cooperatively activate HIF1 target genes in MDA-MB-231 and RCC4 cells. Bone. 2008;23:1–7. https://doi.org/10.1038/onc.2013.115.STAT3

21. Dinarello A, Betto RM, Diamante L, Tesoriere A, Ghirardo R, Cioccarelli C, et al. STAT3 and HIF1α cooperatively mediate the transcriptional and physiological responses to hypoxia. Cell Death Discov. Springer US; 2023;9:1–12. https://doi.org/10.1038/s41420-023-01507-w

22. Moon EJ, Mello SS, Li CG, Chi JT, Thakkar K, Kirkland JG, et al. The HIF target MAFF promotes tumor invasion and metastasis through IL11 and STAT3 signaling. Nat Commun [Internet]. Springer US; 2021;12. https://doi.org/10.1038/s41467-021-24631-6

23. Jahangiri A, Dadmanesh M, Ghorban K. STAT3 inhibition reduced PD-L1 expression and enhanced antitumor immune responses. J Cell Physiol. 2020;235:9457–63. https://doi.org/10.1002/jcp.29750

24. Zerdes I, Wallerius M, Sifakis EG, Wallmann T, Betts S, Bartish M, et al. STAT3 activity promotes programmed-death ligand 1 expression and suppresses immune responses in breast cancer. Cancers (Basel). 2019;11:1–20. https://doi.org/10.3390/cancers11101479

25. Qing Y, Stark GR. Alternative activation of STAT1 and STAT3 in response to interferon-γ. Journal of Biological Chemistry. © 2004 ASBMB. Currently published by Elsevier Inc; originally published by American Society for Biochemistry and Molecular Biology.; 2004;279:41679–85. https://doi.org/10.1074/jbc.M406413200

26. Senbanjo LT, Aljohani H, Majumdar S, Chellaiah MA. Characterization of CD44 intracellular domain interaction with RUNX2 in PC3 human prostate cancer cells. Cell Communication and Signaling. Cell Communication and Signaling; 2019;6:1–13.

27. Li J, Chen JN, Zeng TT, He F, Chen SP, Ma S, et al. CD133+ liver cancer stem cells resist interferon-gamma-induced autophagy. BMC Cancer [Internet]. BMC Cancer; 2016;16:1–11. https://doi.org/10.1186/s12885-016-2050-6

28. Park J, Kim SK, Hallis SP, Choi BH, Kwak MK. Role of CD133/NRF2 Axis in the Development of Colon Cancer Stem Cell-Like Properties. Front Oncol. 2022;11:1–14. https://doi.org/10.3389/fonc.2021.808300

29. Fakiruddin KS, Lim MN, Nordin N, Rosli R, Zakaria Z, Abdullah S. Targeting of CD133+ cancer stem cells by mesenchymal stem cell expressing TRAIL reveals a prospective role of apoptotic gene regulation in non-small cell lung cancer. Cancers (Basel). 2019;11:1–29. https://doi.org/10.3390/cancers11091261

30. Frank DA, Mahajan S, Ritz J. Fludarabine-induced immunosuppression is associated with inhibition of STAT1 signaling [Internet]. NATURE MEDICINE • VOLUME. 1999. http://medicine.nature.com

31. Tournilhac O, Cazin B, Leprètre S, Diviné M, Maloum K, Delmer A, et al. Impact of frontline fludarabine and cyclophosphamide combined treatment on peripheral blood stem cell mobilization in B-cell chronic lymphocytic leukemia. Blood. American Society of Hematology; 2004;103:363–5. https://doi.org/10.1182/blood-2003-05-1449

32. Ren X, Duan L, He Q, Zhang Z, Zhou Y, Wu D, et al. Identification of niclosamide as a new small-molecule inhibitor of the STAT3 signaling pathway. ACS Med Chem Lett. 2010;1:454–9. https://doi.org/10.1021/ml100146z

33. Liu M, Guo S, Hibbert JM, Jain V, Singh N, Wilson NO, et al. CXCL10/IP-10 in infectious diseases pathogenesis and potential therapeutic implications. Cytokine Growth Factor Rev. 2011. p. 121–30. https://doi.org/10.1016/j.cytogfr.2011.06.001

34. Cui Y, Li YY, Li J, Zhang HY, Wang F, Bai X, et al. STAT3 regulates hypoxia-induced epithelial mesenchymal transition in oesophageal squamous cell cancer. Oncol Rep. 2016;36:108–16. https://doi.org/10.3892/or.2016.4822

35. Courau T, Bonnereau J, Chicoteau J, Bottois H, Remark R, Assante Miranda L, et al. Cocultures of human colorectal tumor spheroids with immune cells reveal the therapeutic potential of MICA/B and NKG2A targeting for cancer treatment. J Immunother Cancer. BioMed Central Ltd.; 2019;7. https://doi.org/10.1186/s40425-019-0553-9

36. Scharping NE, Rivadeneira DB, Menk A V., Vignali PDA, Ford BR, Rittenhouse NL, et al. Mitochondrial stress induced by continuous stimulation under hypoxia rapidly drives T cell exhaustion. Nat Immunol [Internet]. Springer US; 2021;22:205–15. https://doi.org/10.1038/s41590-020-00834-9

37. Wu Y, Zhou BP. Inflammation: a driving force speeds cancer metastasis. Cell Cycle [Internet]. 2009;8:3267–73. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3624763/pdf/nihms412728.pdf

38. Ma S, Zhao Y, Lee WC, Ong LT, Lee PL, Jiang Z, et al. Hypoxia induces HIF1α-dependent epigenetic vulnerability in triple negative breast cancer to confer immune effector dysfunction and resistance to anti-PD-1 immunotherapy. Nat Commun. Springer US; 2022;13. https://doi.org/10.1038/s41467-022-31764-9

39. Cui Y, Li YY, Li J, Zhang HY, Wang F, Bai X, et al. STAT3 regulates hypoxia-induced epithelial mesenchymal transition in oesophageal squamous cell cancer. Oncol Rep. 2016;36:108–16. https://doi.org/10.3892/or.2016.4822

40. Liu L, Feng D, Chen G, Chen M, Zheng Q, Song P, et al. Mitochondrial outer-membrane protein FUNDC1 mediates hypoxia-induced mitophagy in mammalian cells. Nat Cell Biol. 2012;14:177–85. https://doi.org/10.1038/ncb2422

Figure Legends

Figure 1. **IFNγ up-regulate surface PD-L1 expression in MC38 cells through IFN-STAT1 signaling.** PD-L1 expression on MC38 cell surface was significantly increased in the presence of IFN-γ analyzed by flow cytometry (**A**) and by immune

fluorescent staining (**B-C**) at 24 hours. (**D**) When treated with siRNA of STAT1, the up-regulation of PD-L1 was significantly reduced, while siRNA of STAT3 has an opposite effect. (**E-F**) The small interfering RNA reaches a sufficient knock down of STAT1 and STAT3, as shown by real-time PCR assay. (**G-I**) The mRNA of STAT1 target genes such as CXCL10 and IL-15 shown similar trend of regulation by IFN-γ as PD-L1. (**J**) The mRNA of STAT3 target gene, MCL-1, shown totally different regulation under normal condition and under treatment of siRNAs. (**K-L**) The expression of STAT1/STAT3 normal and phosphorylated proteins, and PD-L1 proteins in MC38 cell lines were measured by Western blotting with different dose of IFN-γ stimulation **(K)** or after knocking down of STAT1 (siSTAT1) or STAT3 (siSTAT3) **(L)**. The number on the image indicates the relative abundance of PD-L1 protein (fold of control). The results are expressed as the mean ± SEM of triplicate measurements in each group. *p<0.05, **p<0.01, ***p<0.001.

Figure 2. IFN𝛾 indirectly modulate cancer cell stemness in MC38 cells through IFN-STAT3 pathway. (**A-C**) Ki67 expression in the MC38 tumor spheres was significantly reduced in the presence of IFN-γ analyzed by flow cytometry and immune fluorescence. (**D-E**) The tumor sphere forming unit induced in the MC38 cells were also reduced in the presence of IFN-γ (Arrowhead: tumor spheroid. Scale bar: 200μm). **(F)** When MC38 cells were pre-treated with siRNA of STAT1/STAT3, the SFU show different changes. **(G)** When MC38 cells were pre-treated with siRNA of STAT1/STAT3, the CD44+CD133+ population show different trend of regulation by

IFN-γ. **(H-K)** The mRNA of cancer stem cell markers was measured by real-time PCR. **(L)** The expression of cancer stem cell marker proteins in cell nucleus was measured by Western blotting. The results are expressed as the mean ± SEM of triplicate measurements in each group. *p<0.05, **p<0.01, ***p<0.001.

Figure 3. Niclosamide has inhibition effect on both STAT1 and STAT3, which blocks IFNγ-induced PD-L1 up-regulation in MC38 cells while also reduces CSCs formation. (A) The expression of STAT1/STAT3 signaling pathway and PD-L1 proteins in MC38 cell lines with Fludarabine and Niclosamide treatment was measured by Western blotting. **(B)** The surface PDL1 expression level was measured by FACS with IFN-γ and Niclosamide. (**C**) The cell viability of MC38 when co-cultured with T cells and pre-treated with Fludarabine and Niclosamide were measured by CCK8 assay. (**D**) The sphere forming units induced in MC38 cells was measured with or without Niclosamide. (**E**) The cell population of CD44+/CD133+ in MC38 treated with Niclosamide was measured by FACS. (**F**) The expression of cancer stem cell marker proteins in cell nucleus was measured by Western blot. The results are expressed as the mean ± SEM of triplicate measurements in each group. *p<0.05, **p<0.01, ***p<0.001.

Figure 4. RNA-sequencing analysis from total RNA sample of MC38 cells treated with IFNγ. (A) The volcano map of DEGs from RNA-sequencing analysis comparing IFNγ-treated group to control group**. (B)** The volcano map of DEGs from

RNA-sequencing analysis comparing hypoxia group to normoxia group. **(C-N)** The

real-time PCR validation of varies genes show changes in the RNA-sequencing. The

results are expressed as the mean ± SEM of triplicate measurements in each group.

*p<0.05, **p<0.01, ***p<0.001.

Figure 5. Hypoxia enhances PD-L1 upregulation by IFNγ, while Niclosamide

down-regulates Hif1α under hypoxia condition. (A-B) The expression level of

STAT1/STAT3 signaling pathway and PD-L1 proteins in MC38 cell lines with siRNA

(**A**) or Fludarabine and Niclosamide treatment (**B**) was measured by Western blot. (**C**)

Cell viability of MC38 cells pre-treated with different dose of IFN-γ and co-cultured

with primary T cells were measured by CCK8 assay, under normoxia or hypoxia

condition. (**D**) The Cell viability of MC38 cells treated with different dose of

Fludarabine or Niclosamide were measured by CCK8 assay, under normoxia or

hypoxia condition. The results are expressed as the mean ± SEM of measurements

from 4 ∼ 6 animals per group, and each dot represents a measurement from one

animal. *p<0.05, **p<0.01, ***p<0.001.

Figure 6. IFNγ and Hypoxia condition has synergistic effect on inducing T cell

exhaustion, and IFNγ is essential for T cell infiltration in a tumor-T cells co-

culture system. (**A**) Scheme of the protocol used to create co-culture model, using

MC38ova adherent cells or tumor-derived spheroids with mouse primary T cells. (**B**)

The exhaustion state of CD8+T cells co-cultured with MC38ova cells measured by

FACS. (**C**) The expression level of PDL1 on MC38ova cells co-cultured with primary

T cells measured by FACS. (**D-E**) The confocal image of tumor-derived spheroids

and infiltrating T cells labeled with ViaFluor650 (VF650), with quantification of

infiltrating T cell numbers. (**F**) The histogram of ViaFluor650-labeled T cells in the co-

culture system measured by flow analysis. The results are expressed as the mean ±

SEM of measurements 6 mice per group, and each dot represents a measurement

from one animal. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Figure 7. Niclosamide reduces tumor cell immune evasion while preserving

IFNγ's effect in facilitating T cell infiltration in a tumor spheroids-T cells co-

culture system under hypoxia condition. (A) The exhaustion state of CD8+T cells

co-cultured with MC38ova cells measured by FACS, pre-treated with or without

Niclosamide. **(B)** The confocal image of tumor-derived spheroids and infiltrating T

cells labeled with viaFluor650**. (C-D)** The flow analysis of infiltrating T cells labeled

with ViaFluor650 under normal or hypoxia condition. The results are expressed as

the mean ± SEM of measurements from 4 animals per group, and each dot

represents a measurement from one animal. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Supplemental Figure 1. PD-L1 upregulation was observed on the surface of

MC38 tumor cells with IFNγ treatment of different dosages and time periods.

Confocal image of immune fluorescent staining of surface PDL1 on MC38 cells.

Supplemental Figure 2. PD-L1 upregulation by IFN*γ* was observed in induced MC38 tumor spheres, which exhibited more CSC features.: (**A**) PD-L1 expression in MC38 tumor spheres level was measured by FACS. (**B**) Flow analysis revealed that the stem cell-like population was much more in tumor spheres compared to adherent cells. (**C**)The PD-L1 expression in MC38 cells treated with siRNA of scramble control, STAT1 or STAT3. (**D**) Tumor spheres show more resistant to T cells compared to adherent MC38 cells when co-cultured with T cells. The results are expressed as the mean ± SEM of triplicate measurements in each group. *p<0.05, **p<0.01, ***p<0.001).

Supplemental Figure 3. IFN*γ* up-regulate PD-L1 expression affect tumor cells' viability while co-cultured with primary T cells. (**A**) Live cells number of MC38 cells pre-treated with or without IFN-γ and then co-cultured with primary T cells of different time period were measured by Trypan Blue staining assay. (**B)** Cell viability of MC38 cells pre-treated with different dose of IFN-γ and co-cultured with primary T cells were measured by CCK8 assay. The results are expressed as the mean ± SEM of triplicate measurements in each group. *p<0.05, **p<0.01, ***p<0.001.

Supplemental Figure 4. Niclosamide reduces tumor spheres formation with or without IFN*γ*. (**A-B**) The tumor spheres forming unit in the MC38 cells were significantly reduced when treated with Niclosamide, but not with fludarabine (F-ara-A). (**C**) Niclosamide shows partially blocking of the IFN*γ* induced up-regulation of

PDL1 in MC38 tumor spheres. The results are expressed as the mean ± SEM of triplicate measurements in each group, ****p<0.0001).

Supplemental Figure 5. Mouse primary CD8+ T cells co-cultured with MC38 tumor cells were induced to massive exhaustion under hypoxia condition, and co-cultured with tumor spheroids reduced T cell proliferation. (A) The flow analysis of two different protocol of inducing T cell exhaustion *in vitro*. Upper panel: the primary T cells were co-cultured with CD3/CD28 beads for 7 days. Lower panel: the primary T cells were co-coculture with MC38-OVA cells for 7 days. **(B)** Flow analysis of control MC38ova cells (without staining), free CD8+T cells labeled with ViaFluor650, and CD8+ T cells get infiltrated into tumor spheroids, with or without IFNγ. The gating cell populations are CD8+ T cell get infiltrated into tumor spheres **(CD8+ Infil**), and CD8+ T cells get proliferated after infiltration **(CD8+ proli). (C)** ViaFluor650 dyes are membrane-permeant compounds that are converted to fluorescent dyes by intracellular esterases and will covalently react with amine groups on intracellular proteins at the same time, forming fluorescent conjugates that are retained in the cell. With each cell division, daughter cells inherit roughly half of the fluorescent label, allowing the number of cell divisions that occur after labeling to be detected by the appearance of successively dimmer fluorescent peaks on a flow cytometry histogram. Thus, CD8+ T cell divisions can be tracked after labeling with the ViaFluor650.

Fig. 1 IFNγ up-regulate surface PD-L1 expression in MC38 cells through IFN-STAT1 signaling.

Fig. 2 IFNγ indirectly modulate cancer cell stemness in MC38 cells through IFN-STAT3 pathway.

Fig. 3 Niclosamide have inhibition effect on both STAT1 and STAT3, which blocks IFNγ-induced PD-L1 up-regulation in MC38 cells while also reduces CSCs formation.

Fig. 4 RNA-sequencing analysis from total RNA sample of MC38 cells treated with IFNγ and Hypoxia.

Fig. 5 Hypoxia enhances PD-L1 upregulation by IFNγ, while Niclosamide down-regulates Hif1α under hypoxia condition.

Fig. 6 IFNγ and Hypoxia condition has synergistic effect on inducing T cell exhaustion, and IFNγ is essential for T cell infiltration in a tumor-T cells co-culture system.

Fig. 7 Niclosamide reduce tumor cell immune evasion while preserving IFNγ's effect in facilitating T cell infiltration in a tumor spheroids-T cells co-culture system under hypoxia condition.