1 **Causes and consequences of sex-chromosome turnovers in Diptera**

2 Lorena Layana1, Melissa A.Toups2@, Beatriz Vicoso1@
3
4 1 Institute of Science and Technology Austria
5 2 University of Louisiana Lafayette, Department of Biology
6 @ Correspondence: melissa.toups@louisiana.edu, bvicoso@ist.ac.at
7

8 **ORCID**
9 Beatriz Vicoso: https://orcid.org/0000-0002-4579-8306
10 Melissa Toups: https://orcid.org/0000-0002-9752-7380
11 Lorena Layana: https://orcid.org/0000-0002-1253-6297
12

Abstract
Sex-chromosome systems are highly variable across animals, but how they
transition from one to another is not well understood. Diptera have undergone multiple
sex-chromosome turnovers and expansions while maintaining their general
chromosomal content, which makes them an ideal clade to study such transitions. We
analysed more than 100 dipteran whole-genome assemblies and identified 4 new
lineages that underwent sex-chromosome turnover (in addition to the 5 previously
reported). We find the majority of turnovers happened in the group Schizophora, which
tend to have fewer genes on the F element (the chromosome homologous to the
ancestral insect X chromosome) than lower Dipterans. Most derived X chromosomes
have higher GC content than autosomes, which may support the prevalence of male-
achiasmy in Diptera. In addition, an excess of gene movement out of the X is detected
for most of these new X chromosomes, suggesting this process shapes the long-term
gene content of sex chromosomes.
Keywords: Schizophora, sex chromosomes, sex-chromosome turnover, Diptera,
genomic features, out-of-X movement.
29

Introduction
31
In many organisms, sex is determined by a pair of sex chromosomes (X and Y in
the case of male heterogamety, or Z and W in female heterogamety). As sex and
separate sexes are ancestral in animals, but sex chromosomes are not, turnover (i.e.
the reversal of sex chromosomes to autosomes, and their replacement by a novel sex-
chromosome pair) must have occurred repeatedly. Turnover is expected to be
particularly common when sex chromosomes are young and/or undifferentiated, as at
that point they still maintain most of the features of an autosomal pair (Vicoso 2019).
The accumulation of deleterious mutations on a young but already degenerating Y
chromosome should favor turnover, as this would yield a new XY pair free of such
mutation load (the so-called "hot potato" model of turnover (Blaser et al. 2014)). Indeed,
recurrent turnover has been observed in many clades with undifferentiated sex
chromosomes, such as several groups of fishes (Ansai et al. 2022; Behrens et al. 2024;
El Taher et al.) and frogs (Jeffries et al. 2018; Ma and Veltsos 2021; Evans et al. 2024).
On the other hand, once sex chromosomes are fully differentiated, they are expected to
become more stable, as reverting a highly specialized X or Y chromosome to an

autosome is likely to incur harmful fitness effects (POKORNÁ and KRATOCHVÍL 2009).
Y chromosome gene loss should prevent this chromosome from becoming fixed as an
autosome. However, fixing the X as an autosome may cause fitness effects in males as
a result of misregulation due to relictual dosage compensatory mechanisms or through
missing key male-specific genes. An example of the potential stability of well-
differentiated X chromosomes is provided by mammals, where the same XY pair has
been maintained over 190 million years (Graves 2016). Recent studies have uncovered
even older pairs of sex chromosomes, such as those of liverworts (Iwasaki et al. 2021)
and cephalopods (Coffing et al. 2025), again supporting the idea that differentiated sex
chromosomes may function as an "evolutionary trap".
Diptera (flies and mosquitoes) is an ideal order in which to study chromosomal
evolution, as in many species these are formed by different combinations of 6
conserved chromosomal arms. These so-called "Muller elements" A to F were first
described in *Drosophila*, where their homology was inferred from similar banding
patterns of salivary gland chromosomal spreads and similar segregation patterns in
their linkage maps (Schaeffer 2018). In all *Drosophila* species, element A is the X
chromosome, although additional elements have become sex-linked in many species
through fusions of autosomes to the X and/or to the Y (Ellison and Bachtrog 2019). The
sequencing of the genomes of both *Drosophila melanogaster* and *Anopheles gambiae*
showed that the conservation of gene content of Muller elements extended beyond
Drosophila, as homology could be detected between the chromosomal arms of the two
species based on their gene content (Zdobnov et al. 2002). Recent analyses of Dipteran
genomes show that the Muller elements of higher Diptera have undergone several
rearrangements compared to the ancestral dipteran karyotype (Gries, Jaron and
colleagues, submitted), but are well conserved through much of the Brachycera branch
(and in particular in the Schizophora section, i.e. muscoid flies and their relatives). Since
most of the sex chromosome changes studied previously and in this work occurred in
the Schizophora, we describe them in terms of homology to Muller elements.
A chromosome homologous to Element A was also inferred to be the X in *A.*
gambiae (Zdobnov et al. 2002), which initially suggested that the X chromosome of
dipteran insects might have been another example of a well conserved X chromosome.
However, further sampling of the clade showed that element F was instead the
ancestral X of the clade, and had been replaced by element A in both *Drosophila* and
Anopheles independently (Vicoso and Bachtrog 2015). In *Drosophila*, Muller element F
is a highly unusual autosome, which likely reflects its history as a former X
chromosome. It carries fewer than 100 genes and is dot-shaped in chromosomal
spreads (in *D. melanogaster* is also known as the "dot chromosome", or as the "fourth
chromosome") (Riddle and Elgin 2018). Element F is almost fully constitutively
heterochromatic, and is globally regulated by the protein Painting-of-Fourth (Riddle and
Elgin 2018). Similar turnover events were inferred in 6 other lineages, with another 5
having transitioned to undifferentiated sex chromosomes (Vicoso and Bachtrog 2015).
Whether the former X chromosome simply reverted to an autosome, or underwent more
complex rearrangements, could not be established in the absence of chromosome-level
genome assemblies.
This high frequency of turnover of an old and well-differentiated X chromosome is
at odds with the evolutionary trap hypothesis, but also with other insects, which typically

share an ancient X chromosome homologous to element F (Meisel et al. 2019;
Chauhan et al. 2021; Toups and Vicoso 2023; Li et al. 2024). It is currently unclear why
turnover has occurred in Diptera but not other insect orders, although one possibility is
that the shrinking of gene content on the X that occurred in the ancestor of this group
facilitated its reversal to an autosome (Vicoso and Bachtrog 2015; Lasne et al. 2023;
Toups and Vicoso 2023). However, turnover was studied using highly fragmented
genome assemblies, and a detailed and systematic genomic characterization of derived
X chromosomes has yet to be performed. Here, we take advantage of many fly and
mosquito chromosome-level genome assemblies that have recently been made
available by the Darwin Tree of Life and other projects (Richards et al. 2005; Konganti
et al. 2018; Mahajan et al. 2018; Bracewell et al. 2019; Hill et al. 2019; Renschler et al.
2019; Hemmer et al. 2020; Mai et al. 2020; Massey et al. 2020; Small et al. 2020;
Torosin et al. 2020; Chakraborty et al. 2021; Durkin et al. 2021; Urban et al. 2021;
Zamyatin et al. 2021; Lukyanchikova et al. 2022; Romine et al. 2022; The Darwin Tree
of Life Project Consortium 2022; Wei et al. 2022; Alencar et al. 2023; Leung et al. 2023;
Meisel et al. 2023; Reinhardt et al. 2023; Tandonnet et al. 2023; Trinca et al. 2023;
Congrains et al. 2024) to confirm several putative sex chromosome transitions, as well
as identify several additional ones. We also investigate some genomic and evolutionary
features of these independently derived X chromosomes that could not be studied
previously, such as the fate of reverted X chromosomes and the role of gene movement
in shaping the gene content of the X and autosomes.
114
115 **Results & discussion**
116
1 Additional sex-chromosome turnover events and X-to-autosome fusions
Earlier work identified sex-chromosome turnover (with the reversal of element F
to an autosome) using highly-fragmented genome assemblies; scaffolds were assigned
to the X or autosomes based on coverage, but their location on different autosomes was
unknown (Vicoso and Bachtrog 2015). There are now over 100 dipteran species for
which chromosome-level assemblies with an identified X are available. We mapped the
coding sequences of *D. melanogaster* to each chromosome of 149 dipteran genomes
(listed in Supplementary Table 1) to identify their homology to Muller elements. While
fewer homologs were detected in species that are more distant from *Drosophila*, more
than 6000 were identified in each, giving us sufficient power to identify chromosomal
homology. We calculated the expected number of shared genes between each pair of
D. melanogaster / outgroup chromosomes, and inferred homology when the ratio of the
observed to expected number was above 1.5 (following (Toups and Vicoso 2023)), and
this excess was significant with a Fisher's exact test. We could then compare the X of
different species in a phylogenetic framework (based on the phylogeny of (Sperling and
Glover 2023), modified from (Wiegmann et al. 2011)); when several species in a family
had the same chromosomal complement, only one was used for further analyses. As
expected of an ancestral X, element F corresponded to the sole X in 26 out of 41
families (Figure 1 and Supplementary Table 1). We recovered several transitions that
were inferred previously, including the turnover events of Drosophilidae and Diopsidae
(in which the ancestral X reverted to an autosome, e.g. the clades of Figure 1 where
element F is no longer part of the X). Interestingly, while the X of *A. gambiae* was

139
140
141
Figure 1: Multiple transitions in sex chromosomes in dipteran insects (in red asterisks).
Branches highlighted in bold/black correspond to the Schizophora group. The bars at
each edge of the phylogeny represent the different X chromosome complements that
were detected in this study (family name shaded) or previously (Vicoso and Bachtrog
2015; Ellison and Bachtrog 2019). In Polleniidae (not included in the reference tree
used here) the X is homologous to element F. The barplot summarizes the number of
families in which each Muller element has been co-opted as the X chromosome.
Numbers in parentheses next to the family names correspond to the number of species
sampled.
151

previously inferred to derive from a turnover event to element A (Vicoso and Bachtrog
2015), all but two of the *Anopheles* species sampled here (*A. rivulorum* and *A. coluzzii*)
show a significant enrichment of both element A and element F genes on the X,
suggesting an X to autosome fusion instead (Supplementary Dataset 1). A 1.4-fold
enrichment of element F genes is also present on the X of *A. rivulorum* and *A. coluzzii*
(Supplementary Dataset 1), suggesting that the difference reflects a loss of power in
these species rather than biological variation. It should be noted that some lineages,
such as Drosophilidae, have been sampled at much higher depth than others, and this
may account for the apparent excess of neo-sex chromosome origination events.
We also uncovered four additional turnover events in Clusiidae, Conopidae,
Hippoboscidae and Sciaridae, and the presence of several additional neo-X arms
through fusions to the ancestral X (in Conopidae, Bombyliidae, and Asilidae). In total, 9
full turnover events have now been uncovered in this clade, of which 6 happened in
groups with chromosome-level assemblies including an annotated X are available (this
number does not include turnovers to undifferentiated sex chromosomes (Vicoso and
Bachtrog 2015), which we do not consider here). In 3 lineages (Conopidae, Diopsidae
and Sciaridae), turnover co-occurred with the fusion of element F to an autosome, while
in two (Drosophilidae and Hippoboscidae) element F reverted to an independent
autosomal element (Supplementary Dataset 1,2 and Supplementary Figure 1). In
Clusiidae, element F genes are almost exclusively found in scaffolds that were not
assembled into chromosomes (Supplementary Dataset 3), suggesting that a "dot"
chromosome may also be present in this lineage. Given that only a small subset of
lineages have fusions involving element F, the fact that at least 3 out of 6 lineages that
underwent turnover do so suggests that such fusions may favor turnover, or stabilize
the existence of the reverted ancestral X after turnover. It should be noted that although
the X chromosome of Sciaridae corresponds to both elements A and E (Figure 1), this
represents a simple turnover event, as a homologous autosome containing parts of
elements A and E is present in the sister group Bibionidae (chromosome 1 in *Bibio*
marce, Supplementary Figure 2). This is in line with (Gries, Jaron and colleagues,
submitted), who recovered similar sex chromosome transitions when using ancestrally
reconstructed chromosomes of Diptera as their reference instead.
Turnover events and X to autosome fusions may reflect general dynamics of
rearrangements in dipteran insects, or alternatively, may have been favored by
selection if they allowed for the sex-linkage of alleles or genes with sex-specific
benefits. In particular, sexually-antagonistic alleles, i.e. alleles that benefit one sex at
the expense of the other, are theoretically expected to promote the fusion of the
autosome they are on to either the X or Y chromosomes (Charlesworth and
Charlesworth 1980; Matsumoto and Kitano 2016), and can in principle also favor
turnover (van Doorn and Kirkpatrick 2007). If such selective processes are at play,
fusions involving sex chromosomes may be more common than expected by chance
(Anderson et al. 2020). Clear 1:1 homology between Muller elements is only found in
the section Schiphozora (which includes *Drosophila* and other muscoid flies), and
therefore we limited our analysis of fusions between X-linked and autosomal elements
to this group (Supplementary Figure 3). We did not consider fusions that occurred in
Drosophila, as these have been considered elsewhere, and found to be rarer than
expected by chance (Anderson et al. 2020). We find only 3 fusions involving the sex

chromosome, and 5 autosome-autosome fusions, compared to an expected 2.66 and
5.33 respectively (Blackmon and Adams 2015; Anderson et al. 2020), small numbers
that at least do not provide evidence of selection. Similarly, it has been hypothesized
that some chromosomes may be more prone to accumulating sexually-antagonistic
mutations (if they contain genes that are under strong sexual conflict), and may
therefore be co-opted as sex chromosomes more often than expected by chance
(Charlesworth and Charlesworth 1980; Marshall Graves and Peichel 2010; Jeffries et al.
2018; Toups et al. 2019; Anderson et al. 2020; Mora et al. 2024). In our dataset
(including turnover and autosome-sex chromosome fusions), element D was co-opted
as the X in at least 8 families, element A and B in 5, and elements C and E in 4 or fewer
families. This distribution is not different from random sampling (p=0.23 of one Muller
element being represented at least 8 times with binomial testing). While it is possible
that this just reflects a lack of power due to the small number of events, we currently
have no evidence of selection favoring one chromosome over the others.
While Diptera lineages generally have highly dynamic sex-chromosome
evolution, one family stood out: in 4 species of bee grabbers (Conopidae) for which an
annotated X chromosome was available, 4 different X chromosome complements were
found (Supplementary Figure 4). *Conops quadrifasciatus* maintains the ancestral X,
while fusions to this chromosome happened in *Myopa testacea* (F+D) and *Sicus*
ferrugineus (F+A, although it is likely that this represents a Y-autosome fusion, Layana
et al., under preparation). Finally, a turnover event occurred in *Thecophora atra*, where
element B replaced the original element F as the X. The time for the divergence of
these Conopidae species is only 44 million years, highlighting how dynamic sex
chromosomes are in this group.
222
2 The section Schizophora has reduced gene content on element F and increased
turnover rates
One hypothesis for why Dipterans undergo recurrent sex-chromosome turnover
is the shrinking of the X that occurred in their ancestor (from >1000 to ~100-200 genes)
(Vicoso and Bachtrog 2015; Toups and Vicoso 2023). However, even within Diptera, the
proportion of *Drosophila* genes that map to the chromosome homologous to the
element F can vary by a factor of ~5, from 0.66% of all mapped genes in Calliphoridae
to >3% in Ptychopteridae (Figure 2). It is therefore possible that, even within this order,
turnover may be shaped by this difference in gene content. In particular, 8 out of 9 of
the turnover events (with reversal of element F to an autosome) happened in the
section Schizophora, despite a small difference in the total number of families that were
sampled inside and outside of the section (21 vs 18). To test if the rate of turnover is
higher in Schizophora than in other dipteran insects, we produced a rooted tree with
branch lengths using available gene annotations, as well as BUSCO orthologs for
lineages for which an annotation was not available. We then applied the approach of
(Revell et al. 2024) to compare the log likelihood of models of turnover with
homogeneous rates throughout the phylogeny to the log likelihood of equivalent models
that allow for variation between Schizophora and other branches (Supplementary Figure
5). Both types of models confirm the intuition that high turnover rates are a feature of
Schizophora (p=0.004 for "ER" models, which allows for equal rates of turnover and
turnover reversal, and p=0.007 for "ARD" models, which assumes different rates of

Figure 2: The proportion of genes located on the ancestral X (element F) is variable
across Diptera (outer circle). Shaded edge family labels denote sex chromosome
turnover (with reversal of the F to an autosome, either fused or unfused to another
Muller element). Branches shaded in black correspond to the Schizophora section,
where 8 out of 9 turnovers occurred.

251
252
turnover and reversal, a more realistic model in our case; both p-values were obtained
with likelihood ratio tests).
All Schizophora families still using element F as their X carry fewer than 1.25% of
mapped genes on this chromosome, suggesting that this small proportion may be
ancestral to the clade. Outside of Schizophora, on the other hand, between 1.45 and
3.19% of mapped *Drosophila* genes are on element F. We used the Phytools function
fastAnc to perform ancestral reconstruction of the percentage of genes on F throughout
the phylogeny (Sup. Fig. 5). This analysis confirms that shrinkage of the F occurred in
the ancestor of Schizophora (1.12% of genes on F, 95% confidence interval: 0.89-
1.50%), with all nodes within Schizophora having smaller percentages than those
outside of Schizophora (p=7.4e-07). These results therefore support the concurrent

shrinking of the gene content of element F and an increase in sex chromosome
turnover. While this association may be due to chance or sampling biases, better
sampling outside of Schizophora may uncover more fine-grained heterogeneity that
could contribute to turnover elsewhere. Finally, while entirely anecdotal, it is intriguing
that the smallest chromosome homologous to element F is found in Hippoboscidae
(0.6% of mapped genes), where it reverted to an independent dot-like autosome, again
in line with the idea that small chromosomes can undergo turnover.
271
3 Muscidae: the mysterious case of the disappearing element F
In most Dipteran lineages, element F was conserved as the sex chromosome,
reverted to an autosome, or was incorporated into an autosome through a fusion. In
Polietes domitor (Muscidae), on the other hand, no chromosome showed strong
evidence of homology to element F (Figure 3C). The putative X chromosome of this
species carries too few genes (1) to determine homology to a Muller element. To
investigate what might have happened to element F in this clade , we first searched for
element F genes in the 10 species of the family Muscidae for which genomes were
available, even if they did not have an assigned X chromosome in the assembly. In two
species of the genus *Hydrotaea* (Falk and Grzywacz 2024; Falk and Grzywacz 2024),
small scaffolds (assigned as B1 and B2 in *H. diabolus*, and as chromosome 6 in *H.*
cyrtoneurina) were enriched for element F genes (Figure 3B). However, these scaffolds
were much smaller than chromosomes homologous to element F in other families, with
a total of only 17 and 20 genes mapped in the two species (of which 7 were common to
both).
Karyotypic data in other *Hydrotaea* suggests that in addition to having 5 large
euchromatic chromosomes, species are often polymorphic for small heterochromatic
chromosomes, with their number varying from 1 to 7 across individuals and species
(Loeschcke et al. 1994). The two small scaffolds of *H. diabolus* had reduced coverage
in the sequenced male individual, and were in light of this karyotypic data inferred to be
supernumerary (B) chromosomes (Falk and Grzywacz 2024). However, such a
coverage pattern is equally likely to reflect a small X chromosome (or two small X
chromosomes), a simpler explanation to account for both reduced coverage and
homology to the ancestral X. It is further possible that such a small X could vary in
number across cells, as a similar pattern has been found for the dot chromosome in
Drosophila melanogaster (Mohr 1932). We therefore hypothesize that the ancestral X
underwent dramatic shrinking in the ancestor of Muscidae to form the minute X of
Hydrotaea. In other Muscidae species, no such small chromosomes with homology to
element F were reported.
To further investigate the evolution of element F in Muscidae, we selected genes
that are on F in *D. melanogaster* and on its homologous X in *Sarcophaga caerulescens*
(Figure 3A), and which therefore must have been on the ancestral X of the lineage
leading up to Muscidae (Calyptratae). Of these 55 genes, only 10 (18%) were found on
the minute X chromosomes of either *Hydrotaea* species. 18 genes (32%) cannot be
found in any of the 10 Muscidae species, suggesting that the shrinking of Muller
element F in the ancestor of this group was accompanied by substantial gene loss. A
similar pattern is observed when all *D. melanogaster* genes from element F are mapped
to each of the Muscidae species, with a relatively large proportion missing

310
311
Figure 3: Extreme shrinking of Muller element F in Muscidae. **A.** Genomic location of *D.*
melanogaster genes mapped to the genomes of *Sarcophaga caerulescens*
(Sarcophagidae), colored according to their Muller element in *D. melanogaster,* to show
an example of a chromosome homologous to element F (the X chromosome); **B.** the
same for *Hydrotaea diabolus* and **C.** for *Polietes domitor* (Muscidae). **D.** Model for the
evolution of Muller element F in Muscidae.
318
319

(Supplementary Figure 6: 49-62% of element F genes recovered, compared to >80% in comparably distant lineages). Finally, 27 of the 55 ancestrally X-linked genes (49%) are detected in at least one Muscidae species, but are not found on the *Hydrotaea* minute X, suggesting that extensive gene movement from the X to the autosomes also contributed to shrinking element F.

The 10 minute X genes had different fates in the various Muscidae that did not have an independent Muller element F (Figure 3D). In *Eudasyphora cyanicolor* and *Muscina levida*, the majority of them were found on Muller element A (8/8 and 6/9, respectively), suggesting that a minute X - A fusion occurred at least once. In the stable fly *Stomoxys calcitrans*, all 8 out of 8 minute X genes were instead found on Muller element D, supporting an independent fusion of the minute X to this chromosome. These observations are in line with (Meisel et al. 2020), who inferred that the X chromosomes of the horn fly and the stable fly correspond to A-F and D-F fusions respectively. Interestingly, this work (which was based on partly assembled genomes) already noted the small number of genes putatively assigned to element F, as well as the lack of an observed homologue of element F in cytological studies. Finally, in *Polietes domitor,* only 2 out of 10 minute X genes were recovered (one on element D, and one on an unmapped scaffold). While it is possible that a minute F is found in this species but failed to assemble, it is also possible that this small chromosome was lost in this lineage (the BUSCO score for the assembly is >99%, (Falk et al. 2024)). Muscidae lineages are very labile in their sex chromosome complements (Meisel et al. 2020). The house fly *Musca domestica* is also noted for its highly polymorphic sex chromosomes, with all of the 5 large chromosomes having been found to carry the male-determining factor (Sharma et al. 2017). Muscidae may therefore perfectly exemplify how an extreme reduction in gene content of the X can be accompanied by highly variable sex determination systems, although more work to identify the X chromosome in its various lineages is still needed to fully characterize turnover dynamics in this clade.

4. High GC content on derived X chromosomes supports widespread male achiasmy

Most *Drosophila* species are male-achiasmatic, i.e., meiotic recombination typically does not occur in males. Cytological observations in a few other dipteran species suggests that a lack of crossovers in male meiosis may be widespread in this group (Satomura et al. 2019). Male achiasmy dramatically influences sex chromosome evolution, as full sex linkage of a chromosome is expected to initially favor turnover (by linking male-favorable mutations to males), but should also promote rapid degeneration of whole Y chromosomes once turnover has occurred (Satomura et al. 2019). Since measuring recombination rates in non-model species remains challenging, how widespread male achiasmy is remains an open question, which we address here indirectly. In particular, the GC content throughout the genome is typically correlated with the local recombination rate (Charlesworth et al. 2020). In species where recombination is similar in males and females, the X has lower recombination rates than the autosomes because it does not recombine in males (where it is paired with the Y). However, in male-achiasmatic species, the X is expected to have higher rates of recombination than the autosomes, because it spends less time in nonrecombining

males. We therefore predict that if male achiasmy is widespread, novel X chromosomes should overall have acquired a higher GC content than autosomes.

We compared the GC content of the X and autosomes of the 11 lineages that acquired one of the large Muller elements as the X (either by itself or as a fusion to the ancestral X, as the small size of element F means the signal will likely be driven by the neo-X chromosome). For lineages for which several assemblies were available, only one is represented in Figure 4 (the rest are shown in Supplementary Figure 7). In 7 out

Figure 4: GC content proportion in 100,000 bp windows of the X and autosomes in Diptera species representative of lineages with derived large X chromosomes (p-value annotation legend according to a Mann-Whitney statistical test: **** for p <= 0.0001, *** for p <= 0.001, ** for p <= 0.01, * for p <= 0.05).

of 11 lineages, the X had significantly higher GC content than the autosomes,
consistent with lack of recombination in males being common in Diptera. In *D.*
melanogaster and in *Sicus ferrugineus* (Conopidae) the GC content of the X was similar
to that of the autosomes, while in only two species it had significantly lower GC content
(Sciaridae: *Bradysia coprophila*, Bombyliidae: *Bombylius discolor*). At least one
Sciaridae species appears to be male-achiasmatic (Amabis et al. 1979). However,
Sciaridae males do not transmit the paternal copy of their genome (neither the X nor the
autosomes) (Baird et al. 2025), and sex-averaged rates of recombination should
therefore be equal for all chromosomes in this group. Bombyliidae, on the other hand,
may represent a true case of loss of male heterochiasmy, as low GC content on the X is
consistent in the 4 available species. The fact that *D. melanogaster*, the best known
example of male achiasmy, does not have higher GC content on the X also shows that
other factors play a role (such as differences in female recombination rates between the
X and autosomes, or sex-biased mutation rates). This is in line with previous data that
recombination rate and GC content are negatively correlated on the X in *D.*
melanogaster, though why this occurs is not fully understood (Singh et al. 2005).
Despite these limitations, our data is therefore at least consistent with a prevalent role of
male achiasmy in shaping Dipteran sex chromosome evolution.
399
5. Excessive movement out of most but not all X chromosomes
X chromosomes often have an unusual gene content. In many insects, they are
enriched for female-biased genes, but have a deficit of male-biased genes (Sturgill et al.
2007). How this evolves is unclear, but one potential contributor is the excessive gene
movement from the X to the autosomes that has been observed in various clades
(Betrán et al. 2002; Emerson et al. 2004; Baker and Wilkinson 2010; Toups and Hahn
2010; Miller et al. 2022). Such runaway genes often have testis- or male-biased
expression (Potrzebowski et al. 2008; Vibranovski et al. 2009), and their movement
could therefore contribute to the demasculinization of the X. Most of these analyses
were done within clades that share the same X chromosome, and relied on duplicates
to infer movement, such that the original copy of the genes still remained on the X. We
therefore took advantage of the many X chromosomes of Dipterans to extend these
analyses to see if an excess of genes fully move from the X to the autosomes over
longer evolutionary times.
We selected from the phylogeny trios of species where (1) a focal species has a
newly derived X that corresponds to one of Muller elements A to E; and (2) two
independent outgroups carry element F as their X. To avoid biases due to different
qualities of annotations, we mapped the *D. melanogaster* gene sequences and used
their best mapping location in each of the other genomes as a proxy for the location of
their ortholog. We used the two outgroups to infer the ancestral location of the genes (if
it was found on homologous chromosomes in the two, otherwise an ancestral location
could not be inferred), and then inferred gene movement if in the focal species the gene
was on a different chromosome. Since the correspondence of Muller elements and
chromosomes becomes more difficult to assess with increasing distance to *Drosophila*,
we focused exclusively on new X chromosomes found in the section Schizophora (in
Diopsidae, Clusiidae, Hippoboscidae, Conopidae and Drosophilidae). We selected *D.*
virilis as representative species of the Drosophilids because each chromosome

427
Figure 5: Out of X gene movement in Schizophora species which have co-opted a large Muller
element as their X chromosome after turnover. **A.** Bar plots show the fraction of genes inferred
to have moved out of each chromosome in the focal species, using the trios shown in the
phylogenetic trees above each (focal species are in bold). Since smaller chromosomes have a
higher chance of experiencing gene loss (because their target space on other chromosomes is
larger), we normalize the fractions by dividing them by the fraction of genes found on other
chromosomes. The bars representing the X chromosome are in green. Asterisks denote a
significant excess of movement out of the X chromosome (*** for p<0.001, ** for p<0.01, * for
p<0.1) without correcting for multiple comparisons. All p-values remain significant after
Bonferroni correction. **B.** StringDB functional enrichment. The barplot shows the gene ontology
(grey) and phenotype (yellow) terms that were enriched among genes that moved out of the X in
at least 2 species, and the number of species that had evidence of enrichment.

comprises a single Muller element, making the analysis more straightforward. In 4 out of
6 species, a clear out-of-X pattern could be seen (Figure 5A), with the X having lost a
greater fraction of genes than the other large elements even after controlling for
chromosome size, which should affect the target space of gene movement. In the other
two, however, the X did not stand out from other chromosomes. While this could reflect
true biological variability, it is also possible that these patterns are influenced by the age
of these sex chromosomes, as gene loss is likely to be slow. In particular, the strongest
effect is seen in *Drosophila*, whose X chromosome has existed for >60 millions of years
(estimate of divergence time between *Drosophila* and *Phortica*, with whom the X is
shared). On the other hand, all new Conopidae X chromosomes arose since the split of
the genera *Conops* and *Myopa*, estimated at 44 million years ago (Kumar et al. 2017),
and no estimated age is available for Hippoboscidae. In the two lineages where element
F reverted to an autosome without fusing (Hippoboscidae and Drosophilidae), we detect
a very strong out-of-F effect, likely due to its history as an X during part of the history of
the focal lineages.
We also performed a functional enrichment of genes that had moved out of the X
using stringDB with the *D. melanogaster* genome as background (Szklarczyk et al.
2023). In the two lineages where no excess of gene movement could be detected, no
enrichment was found either. In the others, between 11 and 40 functional terms were
enriched (FDR<0.05, see Supplementary Tables 4-8), suggesting that the set of genes
that move out of the X is not random. Interestingly, several of these functional terms
were shared by at least two species (Figure 5B). No such overlap was found when the
same analysis was performed with genes that moved between autosomes. The gene
ontology term enriched in most species was "protein-containing complex". However, this
term was also recovered in Drosophilidae for genes that moved between autosomes
(Supplementary Table 9), suggesting that it may reflect a general property of moved
genes, independent of their original genomic location. On the other hand, several GO
categories related to metabolism and two phenotypic terms related to lethality were
enriched among genes that moved out of the X in 3 out of 4 species, but never when
genes that moved between autosomes were used as a control. This suggests that over
long evolutionary time periods, X chromosomes may become depleted of essential
genes. Such genes are the most likely to incur fitness costs when they are lost from the
Y, and selection may favor their relocation to an autosome. On the other hand, no terms
related to spermatogenesis were enriched, suggesting that either demasculinization
through gene movement is not a global pattern, or that genes involved in
spermatogenesis evolve too fast for meaningful analyses to be performed on such
broad time scales. Taken together, these results show that the out-of-X pattern is not
unique to the lineages previously sampled, and is likely to shape X chromosome gene
content over long evolutionary time periods.
479
Conclusions
We analyzed more than 100 different chromosome-level genome assemblies of
Dipterans, and identified 4 new sex-chromosome transitions from the ancestral insect X
in Diptera (in addition to the 5 previously identified). Similar to previous analyses of
Drosophila sex chromosomes, transitions to new sex-chromosome systems did not
preferentially involve a single Muller element, suggesting there was no single element

predisposed to sex-linkage. Interestingly, most turnovers occurred within the
Schizophora, which also has fewer genes on Muller element F than lower Dipterans,
suggesting that even on this scale reverting the ancestral X chromosome to an
autosome may be less deleterious with fewer genes. We indirectly tested for male
achiasmy in the species with novel X chromosomes. We found that most of these X
have higher GC content than the autosomes, supporting how widespread male
achiasmy is in Diptera. Finally, excess gene movement off the X is common and over
time, the X may become depleted for essential genes.

Methods
495
Genome assemblies
We downloaded from NCBI (https://www.ncbi.nlm.nih.gov/assembly) the set of
available Diptera genome assemblies, and their respective list of chromosomes. We
selected all species for which "X" was found in a chromosome description, and manually
checked that an X chromosome had been annotated. While this search was first
performed in February 2023, further species were added to the dataset as they became
available. The full list of analysed genomes, as well as their respective publication when
available, is provided in Supplementary Table 1 (including scaffolded genomes and
genomes with no annotated X in Muscidae and Conopidae).
505
Phylogenetic tree
A phylogenetic tree of Diptera comprising all but one of the families for which a
genome was analysed was adapted from (Sperling and Glover 2023), and visualized
with iTOL, version 7.1 (Letunic and Bork 2024). iTOL was also used to edit the
phylogenetic trees shown in Figure 2 and Supplementary Figures 3 and 4. The
published phylogeny did not include branch lengths. In order to be able to perform the
phylogenetic inferences described in section "**2 The section Schizophora has**
reduced gene content on element F and increased turnover rates", we downloaded
protein annotations for one representative of each family from Ensembl (Darwin Tree of
Life) https://projects.ensembl.org/darwin-tree-of-life/ or other sources (Supplementary
Table 10) and extract the longest sequence per protein using a custom Perl script
(GetLongestCDS_V2.pl). When no annotations were available (in the case of *Glossina*
fuscipes (Glossinidae), *Liriomyza trifolii* (Agromyzidae), *Cirrula hians* (Ephydridae
species), *Sitodiplosis mosellanafa*, (Cecidomyiidae), *Coboldia fuscipes*, (Scatopsidae),
Trichoceridae species, *Coelopa pilipes* (Coelopidae), *Dryomyza anilis* (Dryomyzidae),
Ornithomya fringillina (Hippoboscidae) and *Rhagio lineola* (Rhagionidae), we ran
BUSCO on the genome assembly with the Diptera gene set, and used the resulting
conserved genes instead. In the case of *C. hians*, we used an improved assembly that
was obtained by running Megahit (Li et al. 2015) with default parameters on all available
published Illumina genomic reads (NCBI Bioproject accession: PRJNA268392). We also
included the outgroup *Panorpa germanica* (protein set from Ensembl, see
Supplementary Table 10). All protein sets were used as input for Orthofinder (with
standard parameters), which produced a rooted species tree.
529
Mapping of *Drosophila* genes to outgroup genomes

D. melanogaster genes were retrieved from Ensembl
(https://nov2020.archive.ensembl.org/Drosophila_melanogaster/Info/Index). We filtered
for the longest sequence of each CDS using a custom Perl script
(GetLongestCDS_V2.pl). We then used Standalone BLAT, version 36x2 (Kent 2002) to
map the resulting gene sequences to the dipteran genomes using default parameters
but with translated DNA for the query and database. We kept the best hit for each gene
and kept only those genes that do not overlap or overlap by <20 bp using two Perl
scripts (1-besthitblat.pl and 2-redremov_blat_V2.pl). *D. melanogaster* genes were
mapped to all the dipteran genomes specified in Supplementary Table 1. These genes
were also mapped to the assembled Fannidae transcriptome. The set of *D.*
melanogaster genes and their assigned chromosomal location in each outgroup
genome is provided as Supplementary Dataset 4, and the graphical representation of
the location of *D. melanogaster* genes colored according to their original Muller element
is provided in Supplementary Dataset 2.
545

546   **Detection of X-linked Muller elements in Dipterans**
547   We used the results from Blat (described above) to infer X-linked Muller elements
548   in each species. In the absence of homology, the expected number of genes that should
549   be shared by a pair of chromosomes of two species is:

$$N_{expected} = P_1 * P_2 * N_{total}$$

550
551   Where P1 is the proportion of genes found on the chromosome of species 1, P2 is the
552   proportion of genes found on the chromosome of species 2, and N_total is the total
553   number of genes found in both species. Chromosomes of outgroup species were
554   inferred to be homologous to *Drosophila* Muller elements when the number of genes
555   they shared with the Muller element was greater than 1.5-fold the expected number.
556

Detection of X-autosome and autosome-autosome fusions
Within the Schizophora, each Muller element typically corresponds to only 1
chromosome (one exception is found in *Sicus ferrugineus*, where element C is found on
two chromosomes; since it is difficult to separate a fission from a misassembly, we
consider the two to be a single chromosome in our analyses). On the other hand,
Diptera chromosomes can be enriched (>1.5x) for more than 1 Muller element, showing
that fusions have occurred. Most lineages only had independent fusions that could be
counted directly. Where fusions were present in more than one species in a single
family, parsimony was used to infer the number of times they arose.
566

Conopidae tree
The Conopidae phylogenetic tree was inferred with Orthofinder, version 2.5.5
(Emms and Kelly 2019), using the multiple sequence alignment option (MSA). This
option infers a maximum likelihood tree, which uses MAFFT, version 7.505 (Katoh et al.
2002) to align the sequences and FastTree (Price et al. 2010) to infer the tree. We used
protein sequences from the six Conopidae species and a Lauxaniidae (*Calliopum*
simillimum) which was used as the outgroup. The fasta files were downloaded from
https://projects.ensembl.org/darwin-tree-of-life/ and we filtered for the longest sequence
of each protein with the Perl script GetLongestCDS_V2.pl.
576

Phylogenetic analyses

Comparison of turnover rates within and outside Schizophora The Phytools (Revell 2012) functions fitMk and fitmultiMk were used to compare the log-likelihood of models that assume a single turnover rate for the whole tree (fitMk) to models that assume two different turnover rates inside and outside of Schizophora (fitmultiMk). Two types of models were used: ER assumes turnover and turnover reversal happen at equal rates, while ARD allows for asymmetric turnover and reversal rates. P-values were obtained with likelihood ratio tests.

Ancestral reconstruction of % of mapped genes on element F For each species in our analysis, we detected the chromosome that was enriched for element F genes (>1.5 fold relative to the expected number, see description of the approach for detecting homology to X). If this chromosome was also enriched for genes from another Muller element, a fusion was inferred, and the species was excluded from this analysis. For each species where no fusion was found, we extracted the total number of genes mapped to the chromosome homologous to F, and divided it by the total number of mapped genes to obtain a fraction of genes on F. We then used the Phytools function FastAnc to perform ancestral reconstruction of the percentage of mapped genes located on element F throughout the dipteran phylogeny. Families for which only scaffold level assemblies were available, or where element F was fused to one of the large chromosomes, were excluded from this analysis (as it was not possible to infer the % of genes on F).

Fannidae transcriptome assembly

We used a Perl script (QuickAssembly.pl) to download paired-end RNA-seq reads of a male *Fannia canicularis* individual (with accession number SRR8236841 from NCBI) and run Trimmomatic (Bolger et al. 2014). We then used Trinity (Grabherr et al. 2011) to assemble the transcriptome, followed by running the script Trinity_gene_splice_modeler.py (also from the Trinity software package) to obtain super transcripts. Supertranscripts were filtered by size with faFilter (UCSC source code collection, http://genome.ucsc.edu/), keeping only those with more than 500 bp sequence length.

Estimation of GC content and repeats

GC and repeat content were estimated in the same way described in (Lasne et al. 2023). In short, for the GC content we used the Python script GCcalc.py (https://github.com/WenchaoLin/GCcalc) in 100,000 bp window size and 100,000 bp step size along the genome of each species. RepeatModeler (Flynn et al. 2020) and RepeatMasker [https://www.repeatmasker.org/] were used to obtain repeat content in windows size of 10,000 bp.

Gene movement inference and statistical testing

We focused on the section Schizophora to examine movement of genes between chromosomes, as within this section homologous chromosomes are generally well conserved, making inferences of gene movement reliable. For each species that coopted one of the large Muller elements as an X chromosome, we manually selected two independent outgroups that still had element F as the ancestral X. We first identified

homologous chromosomes between the outgroup and focal species. To do so, we
compared the mapping location of *D. melanogaster* genes on the chromosomes of each
focal-outgroup pair of species, and classified two chromosomes as homologous when
they had a more than 1.5-fold excess of observed / expected shared genes (where the
expectation is simply: proportion of mapped genes on the chromosome of species 1 *
proportion of mapped genes on the chromosome of species 2 * total number of mapped
genes). When the location was not in agreement in the two outgroup species, the gene
was not considered further; when it was, this location was assigned as its putative
ancestral location. Gene movement was then inferred when the location in the focal
species was different from the inferred ancestral one.
In order to determine if there was a significant excess of movement out of the X
for each species, we computed the expected number of movements between
chromosomes using the following formula (taken from (Toups et al. 2011), originally
proposed by (Betrán et al. 2002)):
637

$$\frac{\sum N_i L_j f_{ij}}{\sum_i \sum_{j \neq i} N_i L_j f_{ij}}$$

638
639
This formula accounts for the number of genes on the chromosome of the parent copy,
the size of the chromosome with the daughter copy, and whether movement occurred to
an X chromosome or an autosome. The indices i and j are for the chromosomes with
the parent copy and with the daughter copy, respectively, N_i is the number of genes on
the chromosome with the parent copy, L_j is the length of the chromosome of the
daughter copy, and f_{ij} is 1 when j is an autosome and 0.75 when j is an X chromosome.
We computed the expected gene movements for each chromosome, and then binned
the movements as (1) autosome to autosome (2) autosome to X and (3) X to autosome.
We then determined significance using a Chi-square goodness of fit test.
649

Functional enrichment on StringDB

The Flybase accessions of *D. melanogaster* homologs of genes that moved out
of the X in each of the focal species (previous section) were used as input for StringDB
(Szklarczyk et al. 2023) with multiple proteins, and "*Drosophila melanogaster*" as the
reference species. The enriched functional categories obtained with the standard
StringDB analysis with default parameters (as of May 2025) were then downloaded
(Supplementary Tables 3 to 6). A similar analysis was performed with sets of genes
inferred to have moved between autosomes (Supplementary Tables 7 and 8 for
Diopsidae and Drosophilidae; no enrichment was found in other lineages).
659

Data availability

661 Supplementary datasets and Supplementary Table 1 are available at:
662 https://seafile.ist.ac.at/d/036324e37c474706a91a/ [*a permanent URL will be provided*
663 *upon acceptance*]. Pipelines [link will be provided upon publication]
664

Author contributions

666 MAT and BV designed the original study. LL performed most analyses with support from
667 MAT and BV. LL, MAT and BV wrote the manuscript.

Funding

This work was supported by a grant from the Austrian Science Fund (FWF, grant number PAT 8748323) to BV.

Conflict of interest statement

The authors declare no conflict of interest.

Acknowledgements

We thank the Vicoso group for their feedback on an early version of the manuscript. We are grateful to Kamil Jaron and Julia Gries for helpful discussions and for sharing their unpublished work. Computational resources and support were provided by the Scientific Computing Unit at ISTA.

References

Alencar RM et al. 2023. Unravelling the genome of the brackish water malaria vector Anopheles aquasalis. Sci Rep. 13(1):20472 [accessed 2025 Jun 4]. https://www.nature.com/articles/s41598-023-47830-1. https://doi.org/10.1038/s41598-023-47830-1

Amabis JM, Reinach FC, Andrews N. 1979. Spermatogenesis in Trichosia pubescens (Diptera:Sciaridae). J Cell Sci. 36:199–213. https://doi.org/10.1242/jcs.36.1.199

Anderson NW, Hjelmen CE, Blackmon H. 2020. The probability of fusions joining sex chromosomes and autosomes. Biol Lett. 16(11):20200648. https://doi.org/10.1098/rsbl.2020.0648

Ansai S et al. 2022. Diversity of sex chromosomes in Sulawesian medaka fishes. J Evol Biol. 35(12):1751–1764. https://doi.org/10.1111/jeb.14076

Baird RB et al. 2025. Faster adaptation but slower divergence of X chromosomes under paternal genome elimination. Nat Commun. 16(1):5288 [accessed 2025 Oct 23]. https://www.nature.com/articles/s41467-025-60114-8. https://doi.org/10.1038/s41467-025-60114-8

Baker RH, Wilkinson GS. 2010. Comparative Genomic Hybridization (CGH) reveals a neo-X chromosome and biased gene movement in stalk-eyed flies (genus Teleopsis). PLoS Genet. 6(9):e1001121. https://doi.org/10.1371/journal.pgen.1001121

Behrens KA et al. 2024. Turnover of sex chromosomes in the Lake Tanganyika cichlid tribe Tropheini (Teleostei: Cichlidae). Sci Rep. 14(1):2471. https://doi.org/10.1038/s41598-024-53021-3

Betrán E, Thornton K, Long M. 2002. Retroposed New Genes Out of the X in Drosophila. Genome Res. 12(12):1854–1859 [accessed 2025 Apr 26]. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC187566/. https://doi.org/10.1101/gr.604902

Blackmon H, Adams R. 2015. EvobiR: tools for comparative analyses and teaching
evolutionary biology. https://github.com/coleoguy/evobir

Blaser O, Neuenschwander S, Perrin N. 2014. Sex-chromosome turnovers: the hot-
potato model. Am Nat. 183(1):140–146. https://doi.org/10.1086/674026

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 30(15):2114–2120 [accessed 2022 Mar 3].
https://doi.org/10.1093/bioinformatics/btu170.
https://doi.org/10.1093/bioinformatics/btu170

Bracewell R, Chatla K, Nalley MJ, Bachtrog D. 2019. Dynamic turnover of centromeres
drives karyotype evolution in Drosophila Henikoff S, Tautz D, Henikoff S, editors. eLife.
8:e49002 [accessed 2025 Jun 4]. https://doi.org/10.7554/eLife.49002.
https://doi.org/10.7554/eLife.49002

Chakraborty M et al. 2021. Evolution of genome structure in the Drosophila simulans
species complex. Genome Res. 31(3):380–396. https://doi.org/10.1101/gr.263442.120

Charlesworth D et al. 2020. Using GC Content to Compare Recombination Patterns on
the Sex Chromosomes and Autosomes of the Guppy, Poecilia reticulata, and Its Close
Outgroup Species. Mol Biol Evol. 37(12):3550–3562.
https://doi.org/10.1093/molbev/msaa187

Charlesworth D, Charlesworth B. 1980. Sex differences in fitness and selection for
centric fusions between sex-chromosomes and autosomes. Genet Res. 35(2):205–214.
https://doi.org/10.1017/s0016672300014051

Chauhan P et al. 2021. Genome assembly, sex-biased gene expression and dosage
compensation in the damselfly Ischnura elegans. Genomics. 113(4):1828–1837.
https://doi.org/10.1016/j.ygeno.2021.04.003

Coffing GC et al. 2025. Cephalopod sex determination and its ancient evolutionary
origin. Curr Biol. 35(4):931-939.e4 [accessed 2025 Apr 26].
https://www.cell.com/current-biology/abstract/S0960-9822(25)00005-3.
https://doi.org/10.1016/j.cub.2025.01.005

Congrains C et al. 2024. Chromosome-scale genome of the polyphagous pest
Anastrepha ludens (Diptera: Tephritidae) provides insights on sex chromosome
evolution in Anastrepha. G3 GenesGenomesGenetics. 14(12):jkae239 [accessed 2025
Jun 4]. https://doi.org/10.1093/g3journal/jkae239.
https://doi.org/10.1093/g3journal/jkae239

van Doorn GS, Kirkpatrick M. 2007. Turnover of sex chromosomes induced by sexual
conflict. Nature. 449(7164):909–912. https://doi.org/10.1038/nature06178

Durkin SM et al. 2021. Behavioral and Genomic Sensory Adaptations Underlying the
Pest Activity of Drosophila suzukii. Mol Biol Evol. 38(6):2532–2546 [accessed 2025 Jun
4]. https://doi.org/10.1093/molbev/msab048. https://doi.org/10.1093/molbev/msab048

El Taher A et al. Dynamics of sex chromosome evolution in a rapid radiation of cichlid
fishes. Sci Adv. 7(36):eabe8215 [accessed 2025 Apr 26].
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8442896/.
https://doi.org/10.1126/sciadv.abe8215

Ellison C, Bachtrog D. 2019. Recurrent gene co-amplification on Drosophila X and Y
chromosomes. PLoS Genet. 15(7):e1008251.
https://doi.org/10.1371/journal.pgen.1008251

Emerson JJ, Kaessmann H, Betrán E, Long M. 2004. Extensive gene traffic on the
mammalian X chromosome. Science. 303(5657):537–540.
https://doi.org/10.1126/science.1090042

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for
comparative genomics. Genome Biol. 20(1):238 [accessed 2021 Apr 18].
https://doi.org/10.1186/s13059-019-1832-y. https://doi.org/10.1186/s13059-019-1832-y

Evans BJ et al. 2024. Rapid Sex Chromosome Turnover in African Clawed Frogs
(Xenopus) and the Origins of New Sex Chromosomes. Mol Biol Evol. 41(12):msae234.
https://doi.org/10.1093/molbev/msae234

Falk S, Grzywacz A. 2024. The genome sequence of a muscid fly, Hydrotaea diabolus
(Harris, [1780]). Wellcome Open Res. [published online ahead of print] [accessed 2025
Oct 23]. https://doi.org/10.12688/wellcomeopenres.21211.1

Falk S, Sivell D, Webb J, Grzywacz A. 2024. The genome sequence of a muscid fly,
Polietes domitor (Harris, 1780). Wellcome Open Res. 9:58 [accessed 2025 Oct 23].
https://pmc.ncbi.nlm.nih.gov/articles/PMC11411267/.
https://doi.org/10.12688/wellcomeopenres.20648.1

Flynn JM et al. 2020. RepeatModeler2 for automated genomic discovery of
transposable element families. Proc Natl Acad Sci U S A. 117(17):9451–9457.
https://doi.org/10.1073/pnas.1921046117

Grabherr MG et al. 2011. Trinity: reconstructing a full-length transcriptome without a
genome from RNA-Seq data. Nat Biotechnol. 29(7):644–652 [accessed 2022 Mar 3].
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3571712/.
https://doi.org/10.1038/nbt.1883

Graves JAM. 2016. Evolution of vertebrate sex chromosomes and dosage
compensation. Nat Rev Genet. 17(1):33–46. https://doi.org/10.1038/nrg.2015.2

Hemmer LW et al. 2020. Hybrid dysgenesis in Drosophila virilis results in clusters of
mitotic recombination and loss-of-heterozygosity but leaves meiotic recombination

unaltered. Mob DNA. 11(1):10 [accessed 2025 Jun 4]. https://doi.org/10.1186/s13100-
020-0205-0. https://doi.org/10.1186/s13100-020-0205-0

Hill T, Koseva BS, Unckless RL. 2019. The Genome of Drosophila innubila Reveals
Lineage-Specific Patterns of Selection in Immune Genes. Mol Biol Evol. 36(7):1405–
1417 [accessed 2025 Jun 4]. https://doi.org/10.1093/molbev/msz059.
https://doi.org/10.1093/molbev/msz059

Iwasaki M et al. 2021. Identification of the sex-determining factor in the liverwort
Marchantia polymorpha reveals unique evolution of sex chromosomes in a haploid
system. Curr Biol CB. 31(24):5522-5532.e7. https://doi.org/10.1016/j.cub.2021.10.023

Jeffries DL et al. 2018. A rapid rate of sex-chromosome turnover and non-random
transitions in true frogs. Nat Commun. 9(1):4088. https://doi.org/10.1038/s41467-018-
06517-2

Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30(14):3059–
3066 [accessed 2022 Apr 3]. https://doi.org/10.1093/nar/gkf436.
https://doi.org/10.1093/nar/gkf436

Kent WJ. 2002. BLAT--the BLAST-like alignment tool. Genome Res. 12(4):656–664.
https://doi.org/10.1101/gr.229202

Konganti K et al. 2018. A Whole Genome Assembly of the Horn Fly, Haematobia
irritans, and Prediction of Genes with Roles in Metabolism and Sex Determination. G3
GenesGenomesGenetics. 8(5):1675–1686 [accessed 2025 Jun 4].
https://doi.org/10.1534/g3.118.200154. https://doi.org/10.1534/g3.118.200154

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A Resource for
Timelines, Timetrees, and Divergence Times. Mol Biol Evol. 34(7):1812–1819.
https://doi.org/10.1093/molbev/msx116

Lasne C et al. 2023. The Scorpionfly (Panorpa cognata) Genome Highlights Conserved
and Derived Features of the Peculiar Dipteran X Chromosome. Mol Biol Evol.
40(12):msad245. https://doi.org/10.1093/molbev/msad245

Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the
phylogenetic tree display and annotation tool. Nucleic Acids Res. 52(W1):W78–W82.
https://doi.org/10.1093/nar/gkae268

Leung W et al. 2023. Long-read genome assemblies for the study of chromosome
expansion: Drosophila kikkawai, Drosophila takahashii, Drosophila bipectinata, and
Drosophila ananassae. G3 GenesGenomesGenetics. 13(10):jkad191 [accessed 2025
Jun 4]. https://doi.org/10.1093/g3journal/jkad191.
https://doi.org/10.1093/g3journal/jkad191

Li D et al. 2015. MEGAHIT: an ultra-fast single-node solution for large and complex
metagenomics assembly via succinct de Bruijn graph. Bioinformatics. 31(10):1674–
1676 [accessed 2022 Apr 3]. https://doi.org/10.1093/bioinformatics/btv033.
https://doi.org/10.1093/bioinformatics/btv033

Li X, Mank JE, Ban L. 2024. The grasshopper genome reveals long-term gene content
conservation of the X Chromosome and temporal variation in X Chromosome evolution.
Genome Res. 34(7):997–1007. https://doi.org/10.1101/gr.278794.123

Loeschcke V, Nielsen BO, Andersen D. 1994. Relationship Among Hydrotaea Species
Based on Allozymes, Karyotype and Morphology (Diptera: Muscidae). Hereditas.
121(1):103–111 [accessed 2025 Oct 23].
https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1601-5223.1994.00103.x.
https://doi.org/10.1111/j.1601-5223.1994.00103.x

Lukyanchikova V et al. 2022. Anopheles mosquitoes reveal new principles of 3D
genome organization in insects. Nat Commun. 13(1):1960 [accessed 2025 Jun 4].
https://www.nature.com/articles/s41467-022-29599-5. https://doi.org/10.1038/s41467-
022-29599-5

Ma W-J, Veltsos P. 2021. The Diversity and Evolution of Sex Chromosomes in Frogs.
Genes. 12(4):483. https://doi.org/10.3390/genes12040483

Mahajan S et al. 2018. De novo assembly of a young Drosophila Y chromosome using
single-molecule sequencing and chromatin conformation capture. PLOS Biol.
16(7):e2006348 [accessed 2025 Jun 4].
https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.2006348.
https://doi.org/10.1371/journal.pbio.2006348

Mai D, Nalley MJ, Bachtrog D. 2020. Patterns of Genomic Differentiation in the
Drosophila nasuta Species Complex. Mol Biol Evol. 37(1):208–220 [accessed 2025 Jun
4]. https://doi.org/10.1093/molbev/msz215. https://doi.org/10.1093/molbev/msz215

Marshall Graves JA, Peichel CL. 2010. Are homologies in vertebrate sex determination
due to shared ancestry or to limited options? Genome Biol. 11(4):205.
https://doi.org/10.1186/gb-2010-11-4-205

Massey JH et al. 2020. Co-evolving wing spots and mating displays are genetically
separable traits in Drosophila. Evolution. 74(6):1098–1111 [accessed 2025 Jun 4].
https://onlinelibrary.wiley.com/doi/abs/10.1111/evo.13990.
https://doi.org/10.1111/evo.13990

Matsumoto T, Kitano J. 2016. The intricate relationship between sexually antagonistic
selection and the evolution of sex chromosome fusions. J Theor Biol. 404:97–108.
https://doi.org/10.1016/j.jtbi.2016.05.036

Meisel RP et al. 2020. Sex Chromosome Evolution in Muscid Flies. G3
GenesGenomesGenetics. 10(4):1341–1352 [accessed 2025 Apr 26].

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7144080/.
https://doi.org/10.1534/g3.119.400923

Meisel RP et al. 2023. New insights into immune genes and other expanded gene
families of the house fly, Musca domestica, from an improved whole genome sequence.
Arch Insect Biochem Physiol. 114(3):e22049 [accessed 2025 Jun 4].
https://onlinelibrary.wiley.com/doi/abs/10.1002/arch.22049.
https://doi.org/10.1002/arch.22049

Meisel RP, Delclos PJ, Wexler JR. 2019. The X chromosome of the German cockroach,
Blattella germanica, is homologous to a fly X chromosome despite 400 million years
divergence. BMC Biol. 17(1):100. https://doi.org/10.1186/s12915-019-0721-x

Miller D et al. 2022. Retrogene Duplication and Expression Patterns Shaped by the
Evolution of Sex Chromosomes in Malaria Mosquitoes. Genes. 13(6):968.
https://doi.org/10.3390/genes13060968

Mohr OL. 1932. GENETICAL AND CYTOLOGICAL PROOF OF SOMATIC
ELIMINATION OF THE FOURTH CHROMOSOME IN DROSOPHILA
MELANOGASTER. Genetics. 17(1):60–80 [accessed 2025 Oct 23].
https://doi.org/10.1093/genetics/17.1.60. https://doi.org/10.1093/genetics/17.1.60

Mora P et al. 2024. Sex-biased gene content is associated with sex chromosome
turnover in Danaini butterflies. Mol Ecol. 33(24):e17256.
https://doi.org/10.1111/mec.17256

POKORNÁ M, KRATOCHVÍL L. 2009. Phylogeny of sex-determining mechanisms in
squamate reptiles: are sex chromosomes an evolutionary trap? Zool J Linn Soc.
156(1):168–183 [accessed 2025 Apr 26]. https://doi.org/10.1111/j.1096-
3642.2008.00481.x. https://doi.org/10.1111/j.1096-3642.2008.00481.x

Potrzebowski L et al. 2008. Chromosomal Gene Movements Reflect the Recent Origin
and Biology of Therian Sex Chromosomes. PLoS Biol. 6(4):e80 [accessed 2025 Apr
26]. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2276528/.
https://doi.org/10.1371/journal.pbio.0060080

Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood
trees for large alignments. PloS One. 5(3):e9490.
https://doi.org/10.1371/journal.pone.0009490

Reinhardt JA et al. 2023. Impacts of Sex Ratio Meiotic Drive on Genome Structure and
Function in a Stalk-Eyed Fly. Genome Biol Evol. 15(7):evad118 [accessed 2025 Jun 4].
https://doi.org/10.1093/gbe/evad118. https://doi.org/10.1093/gbe/evad118

Renschler G et al. 2019. Hi-C guided assemblies reveal conserved regulatory
topologies on X and autosomes despite extensive genome shuffling. Genes Dev.
33(21–22):1591–1612. https://doi.org/10.1101/gad.328971.119

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and
other things). Methods Ecol Evol. 3(2):217–223 [accessed 2025 Oct 23].
https://onlinelibrary.wiley.com/doi/abs/10.1111/j.2041-210X.2011.00169.x.
https://doi.org/10.1111/j.2041-210X.2011.00169.x

Revell LJ, Schliep KP, Mahler DL, Ingram T. 2024. Testing for heterogeneous rates of
discrete character evolution on phylogenies. J Evol Biol. 37(12):1591–1602 [accessed
2025 Oct 23]. https://doi.org/10.1093/jeb/voae119. https://doi.org/10.1093/jeb/voae119

Richards S et al. 2005. Comparative genome sequencing of Drosophila pseudoobscura:
chromosomal, gene, and cis-element evolution. Genome Res. 15(1):1–18.
https://doi.org/10.1101/gr.3059305

Riddle NC, Elgin SCR. 2018. The Drosophila Dot Chromosome: Where Genes Flourish
Amidst Repeats. Genetics. 210(3):757–772 [accessed 2025 Jun 4].
https://doi.org/10.1534/genetics.118.301146.
https://doi.org/10.1534/genetics.118.301146

Romine MG et al. 2022. The genome sequence of the avian vampire fly (Philornis
downsi), an invasive nest parasite of Darwin's finches in Galápagos. G3
GenesGenomesGenetics. 12(2):jkab414 [accessed 2025 Jun 4].
https://doi.org/10.1093/g3journal/jkab414. https://doi.org/10.1093/g3journal/jkab414

Satomura K, Osada N, Endo T. 2019. Achiasmy and sex chromosome evolution. Ecol
Genet Genomics. 13:100046 [accessed 2025 Oct 23].
https://www.sciencedirect.com/science/article/pii/S2405985418300351.
https://doi.org/10.1016/j.egg.2019.100046

Schaeffer SW. 2018. Muller "Elements" in Drosophila: How the Search for the Genetic
Basis for Speciation Led to the Birth of Comparative Genomics. Genetics. 210(1):3–13
[accessed 2025 Apr 26]. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6116959/.
https://doi.org/10.1534/genetics.118.301084

Sharma A et al. 2017. Male sex in houseflies is determined by Mdmd, a paralog of the
generic splice factor gene CWC22. Science. 356(6338):642–645.
https://doi.org/10.1126/science.aam5498

Singh ND, Davis JC, Petrov DA. 2005. Codon bias and noncoding GC content correlate
negatively with recombination rate on the Drosophila X chromosome. J Mol Evol.
61(3):315–324. https://doi.org/10.1007/s00239-004-0287-1

Small ST et al. 2020. Radiation with reticulation marks the origin of a major malaria
vector. Proc Natl Acad Sci. 117(50):31583–31590 [accessed 2025 Jun 4].
https://www.pnas.org/doi/10.1073/pnas.2018142117.
https://doi.org/10.1073/pnas.2018142117

Sperling AL, Glover DM. 2023. Parthenogenesis in dipterans: a genetic perspective.
Proc Biol Sci. 290(1995):20230261. https://doi.org/10.1098/rspb.2023.0261

Sturgill D, Zhang Y, Parisi M, Oliver B. 2007. Demasculinization of X chromosomes in
the Drosophila genus. Nature. 450(7167):238–241. https://doi.org/10.1038/nature06330

Szklarczyk D et al. 2023. The STRING database in 2023: protein-protein association
networks and functional enrichment analyses for any sequenced genome of interest.
Nucleic Acids Res. 51(D1):D638–D646. https://doi.org/10.1093/nar/gkac1000

Tandonnet S et al. 2023. A chromosomal-scale reference genome of the New World
Screwworm, Cochliomyia hominivorax. DNA Res. 30(1):dsac042 [accessed 2025 Jun
4]. https://doi.org/10.1093/dnares/dsac042. https://doi.org/10.1093/dnares/dsac042

The Darwin Tree of Life Project Consortium. 2022. Sequence locally, think globally: The
Darwin Tree of Life Project. Proc Natl Acad Sci. 119(4):e2115642118 [accessed 2025
Jun 4]. https://www.pnas.org/doi/full/10.1073/pnas.2115642118.
https://doi.org/10.1073/pnas.2115642118

Torosin NS et al. 2020. 3D genome evolution and reorganization in the Drosophila
melanogaster species group. PLOS Genet. 16(12):e1009229 [accessed 2025 Jun 4].
https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1009229.
https://doi.org/10.1371/journal.pgen.1009229

Toups MA, Hahn MW. 2010. Retrogenes reveal the direction of sex-chromosome
evolution in mosquitoes. Genetics. 186(2):763–766.
https://doi.org/10.1534/genetics.110.118794

Toups MA, Pease JB, Hahn MW. 2011. No excess gene movement is detected off the
avian or lepidopteran Z chromosome. Genome Biol Evol. 3:1381–1390.
https://doi.org/10.1093/gbe/evr109

Toups MA, Rodrigues N, Perrin N, Kirkpatrick M. 2019. A reciprocal translocation
radically reshapes sex-linked inheritance in the common frog. Mol Ecol. 28(8):1877–
1889 [accessed 2025 Apr 29]. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7558804/.
https://doi.org/10.1111/mec.14990

Toups MA, Vicoso B. 2023. The X chromosome of insects likely predates the origin of
class Insecta. Evol Int J Org Evol. 77(11):2504–2511.
https://doi.org/10.1093/evolut/qpad169

Trinca V et al. 2023. Biocatalytic potential of Pseudolycoriella CAZymes (Sciaroidea,
Diptera) in degrading plant and fungal cell wall polysaccharides. iScience.
26(4):106449. https://doi.org/10.1016/j.isci.2023.106449

Urban JM et al. 2021. High contiguity de novo genome assembly and DNA modification
analyses for the fungus fly, Sciara coprophila, using single-molecule sequencing. BMC
Genomics. 22(1):643 [accessed 2025 Jun 4]. https://doi.org/10.1186/s12864-021-
07926-2. https://doi.org/10.1186/s12864-021-07926-2

Vibranovski MD, Lopes HF, Karr TL, Long M. 2009. Stage-Specific Expression Profiling
of Drosophila Spermatogenesis Suggests that Meiotic Sex Chromosome Inactivation
Drives Genomic Relocation of Testis-Expressed Genes. PLOS Genet. 5(11):e1000731
[accessed 2022 Mar 28].
https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1000731.
https://doi.org/10.1371/journal.pgen.1000731

Vicoso B. 2019. Molecular and evolutionary dynamics of animal sex-chromosome
turnover. Nat Ecol Evol. 3(12):1632–1641 [accessed 2022 Mar 28].
https://www.nature.com/articles/s41559-019-1050-8. https://doi.org/10.1038/s41559-
019-1050-8

Vicoso B, Bachtrog D. 2015. Numerous transitions of sex chromosomes in Diptera.
PLoS Biol. 13(4):e1002078. https://doi.org/10.1371/journal.pbio.1002078

Wei KH-C, Mai D, Chatla K, Bachtrog D. 2022. Dynamics and Impacts of Transposable
Element Proliferation in the Drosophila nasuta Species Group Radiation. Mol Biol Evol.
39(5):msac080 [accessed 2025 Jun 4]. https://doi.org/10.1093/molbev/msac080.
https://doi.org/10.1093/molbev/msac080

Wiegmann BM et al. 2011. Episodic radiations in the fly tree of life. Proc Natl Acad Sci
U S A. 108(14):5690–5695. https://doi.org/10.1073/pnas.1012675108

Zamyatin A et al. 2021. Chromosome-level genome assemblies of the malaria vectors
Anopheles coluzzii and Anopheles arabiensis. GigaScience. 10(3):giab017 [accessed
2025 Jun 4]. https://doi.org/10.1093/gigascience/giab017.
https://doi.org/10.1093/gigascience/giab017

Zdobnov EM et al. 2002. Comparative genome and proteome analysis of Anopheles
gambiae and Drosophila melanogaster. Science. 298(5591):149–159.
https://doi.org/10.1126/science.1077061