Causes and consequences of sex-chromosome turnovers in Diptera

Lorena Layana1, Melissa A.Toups2@, Beatriz Vicoso1@

1 Institute of Science and Technology Austria

2 University of Louisiana Lafayette, Department of Biology

@ Correspondence: melissa.toups@louisiana.edu, bvicoso@ist.ac.at

ORCID
Beatriz Vicoso: https://orcid.org/0000-0002-4579-8306
Melissa Toups: https://orcid.org/0000-0002-9752-7380
Lorena Layana: https://orcid.org/0000-0002-1253-6297

12
13

Abstract

Sex-chromosome systems are highly variable across animals, but how they transition from one

to another is not well understood. Diptera have undergone multiple sex-chromosome turnovers

and expansions while maintaining their general chromosomal content, which makes them an

ideal clade to study such transitions. We analysed more than 100 dipteran whole-genome

assemblies and identified 4 new lineages that underwent sex-chromosome turnover (in addition

to the 5 previously reported). We find the majority of turnovers happened in the group

Schizophora, which tend to have fewer genes on the F element (the chromosome homologous

to the ancestral insect X chromosome) than lower dipterans, a factor previously hypothesized to

facilitate turnover. Most derived X chromosomes have higher GC content than autosomes,

consistent with a high prevalence of male-achiasmy in Diptera. In addition, an excess of gene

movement out of the X is detected for most of these new X chromosomes, and many of these

moved genes have high testis expression in Drosophila, suggesting that out-of-X gene

movement contributes to the long-term demasculinization of X chromosomes.

Keywords: Schizophora, sex chromosomes, sex-chromosome turnover, Diptera, genomic

features, out-of-X movement.

Introduction

In many organisms, sex is determined by a pair of sex chromosomes (X and Y in the case of

male heterogamety, or Z and W in female heterogamety). As sex and separate sexes are

ancestral in animals, but sex chromosomes are not, turnover (i.e. the reversal of sex

chromosomes to autosomes, and their replacement by a novel sex-chromosome pair) must

have occurred repeatedly. Turnover is expected to be particularly common when sex

chromosomes are young and/or undifferentiated, as at that point they still maintain most of the

features of an autosomal pair (Vicoso 2019). The accumulation of deleterious mutations on a

young but already degenerating Y chromosome should favor turnover, as this would yield a new

XY pair free of such mutation load (the so-called "hot potato" model of turnover (Blaser et al.

2014)). Indeed, recurrent turnover has been observed in many clades with undifferentiated sex

chromosomes, such as several groups of fishes (Ansai et al. 2022; Behrens et al. 2024; El

Taher et al.) and frogs (Jeffries et al. 2018; Ma and Veltsos 2021; Evans et al. 2024). On the

other hand, once sex chromosomes are fully differentiated, they are hypothesized to become

more stable, as reverting a highly specialized X or Y chromosome to an autosome is likely to

incur harmful fitness effects (POKORNÁ and KRATOCHVÍL 2009). Y chromosome gene loss

should prevent this chromosome from becoming fixed as an autosome. However, fixing the X as

an autosome may cause fitness effects in males as a result of misregulation due to relictual

dosage compensatory mechanisms or through missing key male-specific genes. An example of

the potential stability of well-differentiated X chromosomes is provided by mammals, where the

same XY pair has been maintained over 190 million years (Graves 2016). Recent studies have

uncovered even older pairs of sex chromosomes, such as those of liverworts (Iwasaki et al.

2021) and cephalopods (Coffing et al. 2025), again supporting the idea that differentiated sex

chromosomes may function as an "evolutionary trap". Alternatively, the long term conservation

of some sex chromosomes (by other, currently unknown, factors) may lead to an apparent

association with differentiation.

Diptera (flies and mosquitoes) is an ideal order in which to study chromosomal evolution, as

in many species these are formed by different combinations of 6 conserved chromosomal arms.

These so-called "Muller elements" A to F were first described in *Drosophila*, where their

homology was inferred from similar banding patterns of salivary gland chromosomal spreads

and similar segregation patterns in their linkage maps (Schaeffer 2018). In all *Drosophila*

species, element A is the X chromosome, although additional elements have become sex-linked

in many species through fusions of autosomes to the X and/or to the Y (Ellison and Bachtrog

2019). The sequencing of the genomes of both *Drosophila melanogaster* and *Anopheles*

gambiae showed that the conservation of gene content of Muller elements extended beyond

Drosophila, as homology could be detected between the chromosomal arms of the two species

based on their gene content (Zdobnov et al. 2002). Recent analyses of dipteran genomes show

that the Muller elements of higher Diptera have undergone several rearrangements compared to

the ancestral dipteran karyotype (Gries, Jaron and colleagues, submitted), but are well

conserved through much of the Brachycera branch (and in particular in the Schizophora section,

i.e. muscoid flies and their relatives). Since most of the sex chromosome changes studied

previously and in this work occurred in the Schizophora, we describe them in terms of homology

to Muller elements.

A chromosome homologous to Element A was also inferred to be the X in *A. gambiae*

(Zdobnov et al. 2002), which initially suggested that the X chromosome of dipteran insects might

have been another example of a well conserved X chromosome. However, further sampling of

the clade showed that element F was instead the ancestral X of the clade, and had been

replaced by element A in both *Drosophila* and Anopheles independently (Vicoso and Bachtrog

2015). In *Drosophila*, Muller element F is a highly unusual autosome, which likely reflects its

history as a former X chromosome. It carries fewer than 100 genes and is dot-shaped in

chromosomal spreads (in *D. melanogaster* is also known as the "dot chromosome", or as the

"fourth chromosome") (Riddle and Elgin 2018). Element F is almost fully constitutively

heterochromatic, and is globally regulated by the protein Painting-of-Fourth (Riddle and Elgin

2018). Similar turnover events were inferred in 6 other lineages, with another 5 having

transitioned to undifferentiated sex chromosomes (Vicoso and Bachtrog 2015). Whether the

former X chromosome simply reverted to an autosome, or underwent more complex

rearrangements, could not be established in the absence of chromosome-level genome

assemblies.

This high frequency of turnover of an old and well-differentiated X chromosome is at odds

with the evolutionary trap hypothesis, but also with other insects, which typically share an

ancient X chromosome homologous to element F (Meisel et al. 2019; Chauhan et al. 2021;

Toups and Vicoso 2023; Li et al. 2024). It is currently unclear why turnover has occurred in

Diptera but not other insect orders, although one possibility is that the shrinking of gene content

on the X that occurred in the ancestor of this group facilitated its reversal to an autosome

(Vicoso and Bachtrog 2015; Lasne et al. 2023; Toups and Vicoso 2023). However, turnover was

studied using highly fragmented genome assemblies, and a detailed and systematic genomic

characterization of derived X chromosomes has yet to be performed. Here, we take advantage

of many fly and mosquito chromosome-level genome assemblies that have recently been made

available by the Darwin Tree of Life and other projects (Richards et al. 2005; Konganti et al.

2018; Mahajan et al. 2018; Bracewell et al. 2019; Hill et al. 2019; Renschler et al. 2019;

Hemmer et al. 2020; Mai et al. 2020; Massey et al. 2020; Small et al. 2020; Torosin et al. 2020;

Chakraborty et al. 2021; Durkin et al. 2021; Urban et al. 2021; Zamyatin et al. 2021;

Lukyanchikova et al. 2022; Romine et al. 2022; The Darwin Tree of Life Project Consortium

2022; Wei et al. 2022; Alencar et al. 2023; Leung et al. 2023; Meisel et al. 2023; Reinhardt et al.

2023; Tandonnet et al. 2023; Trinca et al. 2023; Congrains et al. 2024) to confirm several

putative sex chromosome transitions, as well as identify several additional ones. We also

investigate some genomic and evolutionary features of these independently derived X

chromosomes that could not be studied previously, such as the fate of reverted X chromosomes

and the role of gene movement in shaping the gene content of the X and autosomes.

Results & discussion

1 Additional sex-chromosome turnover events and X-to-autosome fusions

Earlier work identified sex-chromosome turnover (with the reversal of element F to an

autosome) using highly-fragmented genome assemblies; scaffolds were assigned to the X or

autosomes based on coverage, but their location on different autosomes was unknown (Vicoso

and Bachtrog 2015). There are now over 100 dipteran species for which chromosome-level

assemblies with an identified X are available. We mapped the coding sequences of *D.*

melanogaster to each chromosome of 149 dipteran genomes (listed in Table S1) to identify their

homology to Muller elements (see Figure S1 for the graphical representation of these genes in

the chromosomes of one species per family). While fewer homologs were detected in species

that are more distant from *Drosophila*, more than 6000 were identified in each, giving us

sufficient power to identify chromosomal homology. We calculated the expected number of

shared genes between each pair of *D. melanogaster* / outgroup chromosomes, and inferred

homology when the ratio of the observed to expected number was above 1.5 (following (Toups

and Vicoso 2023)), and this excess was significant with a Fisher's exact test (Table S2). We

could then compare the X of different species in a phylogenetic framework (based on the

phylogeny of (Sperling and Glover 2023), modified from (Wiegmann et al. 2011)); when several

species in a family had the same chromosomal complement, only one was used for further

analyses. As expected of an ancestral X, element F corresponded to the sole X in 26 out of 41

families (Figure 1A and Table S1). We recovered several transitions that were inferred

previously, including the turnover events of Drosophilidae and Diopsidae (in which the ancestral

X reverted to an autosome, e.g. the clades of Figure 1A where element F is no longer part of the

X). Interestingly, while the X of *A. gambiae* was previously inferred to derive from a turnover

event to element A (Vicoso and Bachtrog 2015), all but two of the *Anopheles* species sampled

here (*A. rivulorum* and *A. coluzzii*) show a significant enrichment of both element A and element

F genes on the X, suggesting an X to autosome fusion instead (Table S2). A 2-fold enrichment
of element F genes is present on the X of *A. coluzzii* (Supplementary Dataset 1), and a large
proportion of putative X-linked genes is on unmapped scaffolds in *A. rivulorum* (Supplementary
Dataset 2)*, suggesting that the difference reflects a loss of power in these species rather than
biological variation. It should be noted that some lineages, such as Drosophilidae, have been
sampled at much higher depth than others, and this may account for the apparent excess of
neo-sex chromosome origination events.
We also uncovered four additional turnover events in Clusiidae, Conopidae, Hippoboscidae
and Sciaridae, and the presence of several additional neo-X arms through fusions to the
ancestral X (in Conopidae, Bombyliidae, and Asilidae). It should be noted that although the X
chromosome of Sciaridae corresponds to both elements A and E (Figure 1), this represents a
simple turnover event, as a homologous autosome containing parts of elements A and E is
present in the sister group Bibionidae (chromosome 1 in *Bibio marce*, Figure S2). This is in line
with Gries, Jaron and colleagues (*submitted*), who recovered similar sex chromosome
transitions when using ancestrally reconstructed chromosomes of Diptera as their reference
instead. In total, 9 full turnover events have now been uncovered in this clade, of which 6
happened in groups with chromosome-level assemblies including an annotated X are available
(this number does not include turnovers to undifferentiated sex chromosomes (Vicoso and
Bachtrog 2015), which we do not consider here). In 3 lineages (Conopidae, Diopsidae and
Sciaridae), turnover co-occurred with the fusion of element F to an autosome, while in two
(Drosophilidae and Hippoboscidae) element F reverted to an independent autosomal element
(Figure 1B and Supplementary Dataset 2). In Clusiidae, element F genes are almost exclusively
found in scaffolds that were not assembled into chromosomes (Supplementary Dataset 3),
suggesting that a "dot" chromosome may also be present in this lineage. Given that only a small
subset of lineages have fusions involving element F (6 out of 30 families where no turnover
occurred, Fig 1A), the fact that at least 3 out of 6 lineages that underwent turnover do so
suggests that such fusions may favor turnover, or stabilize the existence of the reverted
ancestral X after turnover. This is in stark contrast to other clades, where X:autosome fusions
consistently lead to neo-X formation rather than turnover (Ellison and Bachtrog 2019; Sigeman
et al. 2021; Bracewell et al. 2024; Jayaprasad et al. 2024; Wright et al. 2024).
168

A

Bibionidae (2)
Sciaridae (2)
Cecidomyiidae
Scatopsidae
Culicidae (22)
Ptychopteridae (2)
Tipulidae (2)
Trichoceridae
Rhagionidae (1)
Acroceridae (1)
Stratiomyidae (5)
Bombyliidae (4)
Asilidae (5)
Therevidae (1)
Dolichopodidae (2)
Sciomyzidae (3)
Conopidae (4)
Dryomyzidae (1)
Coelopidae (1)
Lauxaniidae (1)
Heleomyzidae (1)
Pallopteridae (1)
Clusiidae (1)
Diopsidae (1)
Tephritidae (1)
Sepsidae
Calliphoridae (4)
Tachinidae (10)
Rhinophoridae (2)
Sarcophagidae (4)
Anthomyiidae (1)
Muscidae
Glossinidae
Hippoboscidae (2)
Drosophilidae (21)
Ephydridae
Agromyzidae
Pipunculidae (1)
Syrphidae (33)
Hybotidae (1)
Empididae (2)

F
8
5 5 4 2
A B C D E
Muller elements

B

Conopidae	Hippoboscidae	Drosophilidae	Diopsidae	Clusiidae	Sciaridae
Thecophora atra	*Ornithomya fringillina*	*Drosophila virilis*	*Teleopsis dalmani*	*Clusia tigrina*	*Bradysia coprophila*

Conopidae
50 Mbp
X Chr 1 Chr 2 Chr 3

Hippoboscidae
50 Mbp
X Chr 1 Chr 2 Chr 3 Chr 4 **Chr 5**

Drosophilidae
35 Mbp
X Chr 2 Chr 3 Chr 4 Chr 5 **Chr 6**

Diopsidae
180 Mbp
X Chr 1 **Chr 2**

Clusiidae
400 Mbp
X Chr 1 Chr 2 Chr 3 **Unplaced**

Sciaridae
100 Mbp
X Chr 2 **Chr 3** Chr 4

169
170

(legend on next page)

Figure 1: Multiple transitions in sex chromosomes in dipteran insects. A. Phylogeny of the families
sampled here, with their corresponding chromosome complement(s). The bars at each edge of the
phylogeny represent the Muller elements that are homologous to the X chromosome(s) of that family,
either inferred in this study (shaded family name) or previously reported (Vicoso and Bachtrog 2015;
Ellison and Bachtrog 2019). Branches highlighted in bold/black correspond to the Schizophora group, and
turnover events are marked with red asterisks. The barplot summarizes the number of families in which
each Muller element has been co-opted as an X chromosome. Numbers in parentheses next to family
names correspond to the number of species analysed in this study. In Polleniidae (which was missing in
the reference tree used here, and is therefore not shown) the X is homologous to element F. B. Graphical
representation of *D. melanogaster genes* in chromosomes of lineages where a turnover event has
occurred (only one species per lineage is represented). Autosomes homologous to F are highlighted.
182
Turnover events and X to autosome fusions may reflect general dynamics of

rearrangements in dipteran insects, or alternatively, may have been favored by selection if they

allowed for the sex-linkage of alleles or genes with sex-specific benefits. In particular, sexually-

antagonistic alleles, i.e. alleles that benefit one sex at the expense of the other, are theoretically

expected to promote the fusion of the autosome they are on to either the X or Y chromosomes

(Charlesworth and Charlesworth 1980; Matsumoto and Kitano 2016), and can in principle also

favor turnover (van Doorn and Kirkpatrick 2007). If such selective processes are at play, fusions

involving sex chromosomes may be more common than expected by chance (Anderson et al.

2020). Clear 1:1 homology between Muller elements is only found in the section Schiphozora

(which includes *Drosophila* and other muscoid flies), and therefore we limited our analysis of

fusions between X-linked and autosomal elements to this group (Figure S3). We did not

consider fusions that occurred in *Drosophila*, as these have been considered elsewhere, and

found to be rarer than expected by chance (Anderson et al. 2020). We find only 3 fusions

involving the sex chromosome, and 5 autosome-autosome fusions, compared to an expected

2.66 and 5.33 respectively (Blackmon and Adams 2015; Anderson et al. 2020), small numbers

that at least do not provide evidence of selection. Similarly, it has been hypothesized that some

chromosomes may be more prone to accumulating sexually-antagonistic mutations (if they

contain genes that are under strong sexual conflict), and may therefore be co-opted as sex

chromosomes more often than expected by chance (Charlesworth and Charlesworth 1980;

Marshall Graves and Peichel 2010; Jeffries et al. 2018; Toups et al. 2019; Anderson et al. 2020;

Mora et al. 2024). In our dataset (including turnover and autosome-sex chromosome fusions),

element D was co-opted as the X in at least 8 families, element A and B in 5, and elements C

and E in 4 or fewer families. This distribution is not different from random sampling (p=0.23 of

one Muller element being represented at least 8 times with binomial testing). While it is possible

that this just reflects a lack of power due to the small number of events, we currently have no

evidence of selection favoring one chromosome over the others.

While Diptera lineages generally have highly dynamic sex-chromosome evolution, one
family stood out: in 4 species of bee grabbers (Conopidae) for which an annotated X
chromosome was available, 4 different X chromosome complements were found (Figure S4).
Conops quadrifasciatus maintains the ancestral X, while fusions to this chromosome happened
in *Myopa testacea* (F+D) and *Sicus ferrugineus* (F+A, although it is likely that this represents a
Y-autosome fusion, Layana et al., under preparation). Finally, a turnover event occurred in
Thecophora atra, where element B replaced the original element F as the X. The time for the
divergence of these Conopidae species is only 44 million years, highlighting how dynamic sex
chromosomes are in this group.
218
2 The section Schizophora has reduced gene content on element F and increased
turnover rates
One hypothesis for why dipterans undergo recurrent sex-chromosome turnover is the shrinking
of the X that occurred in their ancestor (from >1000 to ~100-200 genes) (Vicoso and Bachtrog
2015; Toups and Vicoso 2023). However, even within Diptera, the proportion of *Drosophila*
genes that map to the chromosome homologous to the element F can vary by a factor of ~5
(Table S3), from 0.66% of all mapped genes in Calliphoridae to >3% in Ptychopteridae (Figure
2). It is therefore possible that, even within this order, turnover may be shaped by this difference
in gene content. In particular, 8 out of 9 of the turnover events (with reversal of element F to an
autosome) happened in the section Schizophora, despite a small difference in the total number
of families that were sampled inside and outside of the section (21 vs 18). To test if the rate of
turnover is higher in Schizophora than in other dipteran insects, we produced a rooted tree with
branch lengths using available gene annotations, as well as BUSCO orthologs for lineages for
which an annotation was not available (Table S4). We then applied the approach of (Revell et
al. 2024) to compare the log likelihood of models of turnover with homogeneous rates
throughout the phylogeny to the log likelihood of equivalent models that allow for variation
between Schizophora and other branches (Figure S5A). Both types of models confirm the
intuition that high turnover rates are a feature of Schizophora (p=0.004 for "ER" models, which
allows for equal rates of turnover and turnover reversal, and p=0.007 for "ARD" models, which
assumes different rates of turnover and reversal, a more realistic model in our case; both p-
values were obtained with likelihood ratio tests).
240

Figure 2: **The proportion of genes located on the ancestral X (element F) is variable across Diptera (outer circle).** Shaded edge family labels denote sex chromosome turnover (with reversal of the F to an autosome, either fused or unfused to another Muller element). Branches shaded in black correspond to the Schizophora section, where 8 out of 9 turnovers occurred.

All Schizophora families still using element F as their X carry fewer than 1.25% of mapped genes on this chromosome, suggesting that this small proportion may be ancestral to the clade. Outside of Schizophora, on the other hand, between 1.45 and 3.19% of mapped *Drosophila* genes are on element F. We used the Phytools function fastAnc to perform ancestral reconstruction of the percentage of genes on F throughout the phylogeny (Figure S5B and Supplementary Dataset 4). This analysis confirms that shrinkage of the F occurred in the ancestor of Schizophora (1.12% of genes on F, 95% confidence interval: 0.89-1.50%), with all nodes within Schizophora having smaller percentages than those outside of Schizophora (p=7.4e-07). These results therefore support the concurrent shrinking of the gene content of element F and an increase in sex chromosome turnover. While this association may be due to

257 chance or sampling biases, better sampling outside of Schizophora may uncover more fine-
258 grained heterogeneity that could contribute to turnover elsewhere. Finally, while entirely
259 anecdotal, it is intriguing that the smallest chromosome homologous to element F is found in
260 Hippoboscidae (0.6% of mapped genes), where it reverted to an independent dot-like
261 autosome, again in line with the idea that small chromosomes can undergo turnover.
262
3 Muscidae: the mysterious case of the disappearing element F
In most dipteran lineages, element F was conserved as the sex chromosome, reverted to an
autosome, or was incorporated into an autosome through a fusion. In *Polietes domitor*
(Muscidae), on the other hand, no chromosome showed strong evidence of homology to
element F (Figure 3C). The putative X chromosome of this species carries too few genes (1) to
determine homology to a Muller element. To investigate what might have happened to element
F in this clade , we first searched for element F genes in the 10 species of the family Muscidae
for which genomes were available, even if they did not have an assigned X chromosome in the
assembly. In two species of the genus *Hydrotaea* (Falk and Grzywacz 2024; Falk and Grzywacz
2024), small scaffolds (assigned as B1 and B2 in *H. diabolus*, and as chromosome 6 in *H.*
cyrtoneurina) were enriched for element F genes (Figure 3B and Figure S1). However, these
scaffolds were much smaller than chromosomes homologous to element F in other families,
with a total of only 17 and 20 genes mapped in the two species (of which 7 were common to
both).
Karyotypic data in other *Hydrotaea* suggests that in addition to having 5 large euchromatic
chromosomes, species are often polymorphic for small heterochromatic chromosomes, with
their number varying from 1 to 7 across individuals and species (Loeschcke et al. 1994). The
two small scaffolds of *H. diabolus* had reduced coverage in the sequenced male individual, and
were in light of this karyotypic data inferred to be supernumerary (B) chromosomes (Falk and
Grzywacz 2024). However, such a coverage pattern is equally likely to reflect a small X
chromosome (or two small X chromosomes), a simpler explanation to account for both reduced
coverage and homology to the ancestral X. It is further possible that such a small X could vary
in number across cells, as a similar pattern has been found for the dot chromosome in
Drosophila melanogaster (Mohr 1932). We therefore hypothesize that the ancestral X
underwent dramatic shrinking in the ancestor of Muscidae to form the minute X of *Hydrotaea*. In
other Muscidae species, no such small chromosomes with homology to element F were
reported.
290

291
292
293 **Figure 3: Extreme shrinking of Muller element F in Muscidae. A.** Genomic location of *D.*
294 *melanogaster* genes mapped to the genomes of *Sarcophaga caerulescens* (Sarcophagidae), colored
295 according to their Muller element in *D. melanogaster,* to show an example of a chromosome homologous
296 to element F (the X chromosome); **B.** the same for *Hydrotaea diabolus* and **C.** for *Polietes domitor*
297 (Muscidae). **D.** Model for the evolution of Muller element F in Muscidae.
298
299

To further investigate the evolution of element F in Muscidae, we selected genes that are on
F in *D. melanogaster* and on its homologous X in *Sarcophaga caerulescens* (Figure 3A), and
which therefore must have been on the ancestral X of the lineage leading up to Muscidae
(Calyptratae) (Table S5). Of these 55 genes, only 10 (18%) were found on the minute X
chromosomes of either *Hydrotaea* species. 18 genes (32%) cannot be found in any of the 10
Muscidae species, suggesting that the shrinking of Muller element F in the ancestor of this
group was accompanied by substantial gene loss. A similar pattern is observed when all *D.*
melanogaster genes from element F are mapped to each of the Muscidae species, with a
relatively large proportion missing (Figure S6: 49-62% of element F genes recovered, compared
to >80% in other comparably distant lineages). Finally, 27 of the 55 ancestrally X-linked genes
(49%) are detected in at least one Muscidae species, but are not found on the *Hydrotaea*
minute X, suggesting that extensive gene movement from the X to the autosomes also
contributed to shrinking element F.
The 10 minute X genes had different fates in the various Muscidae that did not have an
independent Muller element F (Figure 3D). In *Eudasyphora cyanicolor* and *Muscina levida*, the
majority of them were found on Muller element A (8/8 and 6/9, respectively), suggesting that a
minute X - A fusion occurred at least once. In the stable fly *Stomoxys calcitrans*, all 8 out of 8
minute X genes were instead found on Muller element D, supporting an independent fusion of
the minute X to this chromosome. These observations are in line with (Meisel et al. 2020), who
inferred that the X chromosomes of the horn fly and the stable fly correspond to A-F and D-F
fusions respectively. Interestingly, this work (which was based on partly assembled genomes)
already noted the small number of genes putatively assigned to element F, as well as the lack of
an observed homologue of element F in cytological studies. Finally, in *Polietes domitor,* only 2
out of 10 minute X genes were recovered (one on element D, and one on an unmapped
scaffold). While it is possible that a minute F is found in this species but failed to assemble, it is
also possible that this small chromosome was lost in this lineage (the BUSCO score for the
assembly is >99%, (Falk et al. 2024)). Muscidae lineages are very labile in their sex
chromosome complements (Meisel et al. 2020). The house fly *Musca domestica* is also noted
for its highly polymorphic sex chromosomes, with all of the 5 large chromosomes having been
found to carry the male-determining factor (Sharma et al. 2017). Muscidae may therefore
perfectly exemplify how an extreme reduction in gene content of the X can be accompanied by
highly variable sex determination systems, although more work to identify the X chromosome in
its various lineages is still needed to fully characterize turnover dynamics in this clade.
333

Figure 4: GC content proportion in 100,000 bp windows of the X and autosomes in Diptera species representative of lineages with derived large X chromosomes (p-value annotation legend according to a Mann-Whitney statistical test: **** for p <= 0.0001, *** for p <= 0.001, ** for p <= 0.01, * for p <= 0.05, ns for not significant).

4. High GC content on derived X chromosomes supports widespread male achiasmy

Most *Drosophila* species are male-achiasmatic, i.e., meiotic recombination typically does not

occur in males. Cytological observations in a few other dipteran species suggests that a lack of

crossovers in male meiosis may be widespread in this group (Satomura et al. 2019). Male
achiasmy dramatically influences sex chromosome evolution, as full sex linkage of a
chromosome is expected to initially favor turnover (by linking male-favorable mutations to
males), but should also promote rapid degeneration of whole Y chromosomes once turnover
has occurred (Satomura et al. 2019). Since measuring recombination rates in non-model
species remains challenging, how widespread male achiasmy is remains an open question,
which we address here indirectly. In particular, the GC content throughout the genome is
typically correlated with the local recombination rate (Charlesworth et al. 2020). In species
where recombination is similar in males and females, the X has lower recombination rates than
the autosomes because it does not recombine in males (where it is paired with the Y). However,
in male-achiasmatic species, the X is expected to have higher rates of recombination than the
autosomes, because it spends less time in nonrecombining males. We therefore predict that if
male achiasmy is widespread, novel X chromosomes should overall have acquired a higher GC
content than autosomes.

We compared the GC content of the X and autosomes of the 11 lineages that acquired one
of the large Muller elements as the X (either by itself or as a fusion to the ancestral X, as the
small size of element F means the signal will likely be driven by the neo-X chromosome).
Specifically, chromosomes were broken into 100kb windows, which were compared. It should
be noted subdividing chromosomes into windows could introduce pseudoreplication, and the
absolute p-values should therefore be interpreted with caution. For lineages for which several
assemblies were available, only one is represented in Figure 4 (see Supplementary Dataset 5
for GC content estimation in all the species). In 7 out of 11 lineages, the X had significantly
higher GC content than the autosomes, consistent with lack of recombination in males being
common in Diptera. In *D. melanogaster* and in *Sicus ferrugineus* (Conopidae) the GC content of
the X was similar to that of the autosomes, while in only two species it had significantly lower
GC content (Sciaridae: *Bradysia coprophila*, Bombyliidae: *Bombylius discolor*). At least one
Sciaridae species appears to be male-achiasmatic (Amabis et al. 1979). However, Sciaridae
males do not transmit the paternal copy of their genome (neither the X nor the autosomes)
(Baird et al. 2025), and sex-averaged rates of recombination should therefore be equal for all
chromosomes in this group. Bombyliidae, on the other hand, may represent a true case of loss
of male heterochiasmy, as low GC content on the X is consistent in the 4 available species. The
fact that *D. melanogaster*, the best known example of male achiasmy, does not have higher GC
content on the X also shows that other factors play a role (such as differences in female
recombination rates between the X and autosomes, or sex-biased mutation rates). This is in line

with previous data that recombination rate and GC content are negatively correlated on the X in

D. melanogaster, though why this occurs is not fully understood (Singh et al. 2005). Despite

these limitations, our data is therefore at least consistent with a prevalent role of male achiasmy

in shaping dipteran sex chromosome evolution.

5. Excessive movement out of most but not all X chromosomes

X chromosomes often have an unusual gene content. In many insects, they are enriched for

female-biased genes, but have a deficit of male-biased genes (Sturgill et al. 2007). How this

evolves is unclear, but one potential contributor is the excessive gene movement from the X to

the autosomes that has been observed in various clades (Betrán et al. 2002; Emerson et al.

2004; Baker and Wilkinson 2010; Toups and Hahn 2010; Miller et al. 2022). Such runaway

genes often have testis- or male-biased expression (Potrzebowski et al. 2008; Vibranovski et al.

2009), and their movement could therefore contribute to the demasculinization of the X (Sturgill

et al. 2007). Most of these analyses were done within clades that share the same X

chromosome, and relied on duplicates to infer movement, such that the original copy of the

genes still remained on the X. We therefore took advantage of the many X chromosomes of

dipterans to extend these analyses to see if an excess of genes fully move from the X to the

autosomes over longer evolutionary times.

We selected from the phylogeny trios of species where (1) a focal species has a newly

derived X that corresponds to one of Muller elements A to E; and (2) two independent outgroups

carry element F as their X. To avoid biases due to different qualities of annotations, we mapped

the *D. melanogaster* gene sequences and used their best mapping location in each of the other

genomes as a proxy for the location of their ortholog. We used the two outgroups to infer the

ancestral location of the genes (if it was found on homologous chromosomes in the two,

otherwise an ancestral location could not be inferred), and then inferred gene movement if in the

focal species the gene was on a different chromosome (see Table S6 for the number of gene

movements between chromosomes in each of the trios). Since the correspondence of Muller

elements and chromosomes becomes more difficult to assess with increasing distance to

Drosophila, we focused exclusively on new X chromosomes found in the section Schizophora

(in Diopsidae, Clusiidae, Hippoboscidae, Conopidae and Drosophilidae). We selected *D. virilis*

as representative species of the Drosophilids because each chromosome comprises a single

Muller element, making the analysis more straightforward. In 4 out of 6 species, a clear out-of-X

pattern could be seen (Figure 5A), with the X having lost a greater fraction of genes than the

other large elements even after controlling for chromosome size, which should affect the target

space of gene movement. In the other two, however, the X did not stand out from other

chromosomes. While this could reflect true biological variability, it is also possible that these

patterns are influenced by the age of these sex chromosomes, as gene loss is likely to be slow.

In particular, the strongest effect is seen in *Drosophila*, whose X chromosome has existed for

>60 millions of years (estimate of divergence time between *Drosophila* and *Phortica*, with whom

the X is shared). On the other hand, all new Conopidae X chromosomes arose since the split of

the genera *Conops* and *Myopa*, estimated at 44 million years ago (Kumar et al. 2017), and no

estimated age is available for Hippoboscidae. In the two lineages where element F reverted to

an autosome without fusing (Hippoboscidae and Drosophilidae), we detect a very strong out-of-

F effect, likely due to its history as an X during part of the history of the focal lineages.

We investigated whether genes that moved out of the X had unusual patterns of expression

in *D. melanogaster* (as a proxy for ancestral expression, with the exception of the Drosophilidae

analysis, where it largely represents current Drosophilid expression). For each of the tissues in

the FlyAtlas2 dataset, we compared the distribution of expression of genes that moved out of

the X to the expression distribution of all mapped genes. Genes that moved out of the X had

significantly higher expression in between 0 and 12 tissues (Wilcoxon tests). In 5 of the 6 trios,

genes that moved out of the X had significantly higher expression in testis than all genes (Figure

5B; the only species where this was not found, *Sicus ferrugineus*, also did not show an out-of-X

effect), in line with the idea that out-of-X gene movement may deplete X chromosomes of genes

with male-biased expression. This was not the case for genes that moved into the X, or between

autosomes (p>0.05 in all, Figure 5B). Ovary expression was also enriched in out-of-X genes in

3 lineages (Figure S7), but in two of those (Drosophilidae and Conopidae) less so than testis

expression. The one exception is Hippoboscidae, where out-of-X genes had much stronger

ovary expression than any other category of genes; this lineage also did not show an out-of-X

effect. Both testis and ovary enrichment patterns held when modENCODE expression values

were used instead (Figures S8-S10), except for the high ovary expression of out-of-X genes in

Drosophilidae, which was not detected in these data. Finally, we plotted expression distributions

for somatic controls (female and male head, female and male carcass, Figure S11-S14), where

out-of-X genes were never overexpressed compared to the full set of mapped genes. Taken

together, these results suggest that the high expression observed for gonads does not simply

represent broadly high expression, and that in lineages where an excess of movement from the

X is observed, these moved genes tend to have high testis expression.

449
Figure 5: Out-of-X gene movement in Schizophora species which have co-opted a large Muller
element as their X chromosome after turnover. A. Bar plots show the fraction of genes inferred to
have moved out of each chromosome in the focal species, using the trios shown in the phylogenetic trees
above each (focal species are in bold). Since smaller chromosomes have a higher chance of
experiencing gene loss (because their target space on other chromosomes is larger), we normalize the
fractions by dividing them by the fraction of genes found on other chromosomes. The bars representing
the X chromosome are in green. Asterisks denote a significant excess of movement out of the X
chromosome (*** for p<0.001, ** for p<0.01, * for p<0.05) without correcting for multiple comparisons. All

p-values remain significant after Bonferroni correction. **B. *D. melanogaster* testis expression of**
homologs inferred to have moved chromosomes, and that of all mapped genes, for each of the
trios (named after the focal lineage under study). X->Auto refers to genes that moved out of the X,
Auto->X refers to genes that moved into the X, Auto->Auto refers to genes that moved between
autosomes, and All genes refers to all mapped genes in each trio. Significance was assessed with
Wilcoxon tests (* for p<0.05, ** for p<0.01, *** for p<0.001, no asterisk for non significant).
464

We also performed a functional enrichment of genes that had moved out of the X using

stringDB with the *D. melanogaster* genome as background (Szklarczyk et al. 2023). In the two

lineages where no excess of gene movement could be detected, no enrichment was found

either. In the others, between 11 and 40 functional terms were enriched (FDR<0.05, see Table

S7-S11), again suggesting that the set of genes that move out of the X is not random.

Interestingly, several of these functional terms were shared by at least two species (Figure

S15). When the genes that were mapped in each trio were used as the respective background

instead (a more conservative approach to control for biases in what genes are mapped across

species), 4 terms related to metabolism were enriched in Drosophilidae (Table S10), and one in

Diopsidae (RNA polymerase, Table S7). This reduced number of functional enrichments may

reflect a loss of power due to the smaller background set, but also suggests some caution in

interpreting the larger sets of functional enrichments.   Taken together, these results show that

the out-of-X pattern is not unique to the lineages previously sampled, and that the specialized

set of genes that move is likely to shape X chromosome gene content over long evolutionary

time periods.

Conclusions
We analyzed more than 100 different chromosome-level genome assemblies of Dipterans, and

identified 4 new sex-chromosome transitions from the ancestral insect X in Diptera (in addition

to the 5 previously identified). Similar to previous analyses of *Drosophila* sex chromosomes,

transitions to new sex-chromosome systems did not preferentially involve a single Muller

element, suggesting there was no single element predisposed to sex-linkage. Interestingly, most

turnovers occurred within the Schizophora, which also has fewer genes on Muller element F

than lower Dipterans, suggesting that even on this scale reverting the ancestral X chromosome

to an autosome may be less deleterious with fewer genes. We indirectly tested for male

achiasmy in the species with novel X chromosomes. We found that most of these X have higher

GC content than the autosomes, supporting widespread male achiasmy in Diptera. Finally,

excess gene movement off the X is common and, over time, may contribute to the depletion of

testis-expressed genes that is typically observed on old X chromosomes.

Methods

Genome assemblies

We downloaded from NCBI (https://www.ncbi.nlm.nih.gov/assembly) the set of available Diptera

genome assemblies, and their respective list of chromosomes. We selected all species for

which "X" was found in a chromosome description, and manually checked that an X

chromosome had been annotated. While this search was first performed in February 2023,

further species were added to the dataset as they became available. The full list of analysed

genomes, as well as their respective publication when available, is provided in Table S1

(including scaffolded genomes and genomes with no annotated X in Muscidae and Conopidae).

Phylogenetic tree

A phylogenetic tree of Diptera comprising all but one of the families for which a genome was

analysed was adapted from (Sperling and Glover 2023), and visualized with iTOL, version 7.1

(Letunic and Bork 2024). iTOL was also used to edit the phylogenetic trees shown in Figure 2A

and Figure S3 and S4. The published phylogeny did not include branch lengths. In order to be

able to perform the phylogenetic inferences described in section "**2 The section Schizophora**

has reduced gene content on element F and increased turnover rates", we downloaded

protein annotations for one representative of each family from Ensembl (Darwin Tree of Life)

https://projects.ensembl.org/darwin-tree-of-life/ or other sources (Table S4) and extract the

longest sequence per protein using a custom Perl script (GetLongestCDS_V2.pl). When no

annotations were available (in the case of *Glossina fuscipes* (Glossinidae), *Liriomyza trifolii*

(Agromyzidae), *Cirrula hians* (Ephydridae species), *Sitodiplosis mosellanafa*, (Cecidomyiidae),

Coboldia fuscipes, (Scatopsidae), Trichoceridae species, *Coelopa pilipes* (Coelopidae),

Dryomyza anilis (Dryomyzidae), *Ornithomya fringillina* (Hippoboscidae) and *Rhagio lineola*

(Rhagionidae), we ran BUSCO version v5.8.2 (Manni et al. 2021) on the genome assembly

with the Diptera gene set, and used the resulting conserved genes instead (we used the default

gene mapper Miniprot (Li 2023) but for *C. hians*, we used Metaeuk (Levy Karin et al. 2020) as

the gene predictor. The predicted genes were then passed to HMMER (Eddy 2011)). In the

case of *C. hians*, we used an improved assembly that was obtained by running Megahit (Li et al.

2015) with default parameters on all available published Illumina genomic reads (NCBI

Bioproject accession: PRJNA268392). We also included the outgroup *Panorpa germanica*

(protein set from Ensembl, see Table S4). All protein sets were used as input for Orthofinder

(with standard parameters), which produced a rooted species tree.

Mapping of *Drosophila* genes to outgroup genomes

D. melanogaster genes were retrieved from Ensembl
(https://nov2020.archive.ensembl.org/Drosophila_melanogaster/Info/Index). We filtered for the
longest sequence of each CDS using a custom Perl script (GetLongestCDS_V2.pl). We then
used Standalone BLAT, version 36x2 (Kent 2002) to map the resulting gene sequences to the
dipteran genomes using default parameters but with translated DNA for the query and
database. We kept the best hit for each gene and kept only those genes that do not overlap or
overlap by <20 bp using two Perl scripts (1-besthitblat.pl and 2-redremov_blat_V2.pl). *D.
melanogaster* genes were mapped to all the dipteran genomes specified in Table S1. The set of
D. melanogaster genes and their assigned chromosomal location in each outgroup genome is
provided as Supplementary Dataset 7, and the graphical representation of the location of *D.
melanogaster* genes colored according to their original Muller element is provided in
Supplementary Dataset 2.

Detection of X-linked Muller elements in dipterans

We used the results from Blat (described above) to infer X-linked Muller elements in each
species. In the absence of homology, the expected number of genes that should be shared by a
pair of chromosomes of two species is:

$$N_{expected} = P_1 * P_2 * N_{total}$$

Where P1 is the proportion of genes found on the chromosome of species 1, P2 is the
proportion of genes found on the chromosome of species 2, and N_total is the total number of
genes found in both species. Chromosomes of outgroup species were inferred to be
homologous to *Drosophila* Muller elements when the number of genes they shared with the
Muller element was greater than 1.5-fold the expected number.

Detection of X-autosome and autosome-autosome fusions

Within the Schizophora, each Muller element typically corresponds to only 1 chromosome (one
exception is found in *Sicus ferrugineus*, where element C is found on two chromosomes; since it
is difficult to separate a fission from a misassembly, we consider the two to be a single
chromosome in our analyses). On the other hand, Diptera chromosomes can be enriched
(>1.5x) for more than 1 Muller element, showing that fusions have occurred. Most lineages only
had independent fusions that could be counted directly. Where fusions were present in more

than one species in a single family, parsimony was used to infer the number of times they

arose.

Conopidae tree

The Conopidae phylogenetic tree was inferred with Orthofinder, version 2.5.5 (Emms and Kelly

2019), using the multiple sequence alignment option (MSA). This option infers a maximum

likelihood tree, which uses MAFFT, version 7.505 (Katoh et al. 2002) to align the sequences

and FastTree (Price et al. 2010) to infer the tree. We used protein sequences from the six

Conopidae species and a Lauxaniidae (*Calliopum simillimum*) which was used as the outgroup.

The fasta files were downloaded from https://projects.ensembl.org/darwin-tree-of-life/ and we

filtered for the longest sequence of each protein with the Perl script GetLongestCDS_V2.pl.

Phylogenetic analyses

Comparison of turnover rates within and outside Schizophora The Phytools (Revell 2012)

functions fitMk and fitmultiMk were used to compare the log-likelihood of models that assume a

single turnover rate for the whole tree (fitMk) to models that assume two different turnover rates

inside and outside of Schizophora (fitmultiMk). Two types of models were used: ER assumes

turnover and turnover reversal happen at equal rates, while ARD allows for asymmetric turnover

and reversal rates. P-values were obtained with likelihood ratio tests.

Ancestral reconstruction of % of mapped genes on element F For each species in our

analysis, we detected the chromosome that was enriched for element F genes (>1.5 fold

relative to the expected number, see description of the approach for detecting homology to X). If

this chromosome was also enriched for genes from another Muller element, a fusion was

inferred, and the species was excluded from this analysis. For each species where no fusion

was found, we extracted the total number of genes mapped to the chromosome homologous to

F, and divided it by the total number of mapped genes to obtain a fraction of genes on F

(Supplementary Dataset 4). We then used the Phytools function FastAnc to perform ancestral

reconstruction of the percentage of mapped genes located on element F throughout the

dipteran phylogeny. Families for which only scaffold level assemblies were available, or where

element F was fused to one of the large chromosomes, were excluded from this analysis (as it

was not possible to infer the % of genes on F).

Estimation of GC content

GC content was estimated in the same way described in (Lasne et al. 2023). In short, for the

GC content we used the Python script GCcalc.py (https://github.com/WenchaoLin/GCcalc) in

100,000 bp window size and 100,000 bp step size along the genome of each species.

Gene movement inference and statistical testing

We focused on the section Schizophora to examine movement of genes between

chromosomes, as within this section homologous chromosomes are generally well conserved,

making inferences of gene movement reliable. For each species that coopted one of the large

Muller elements as an X chromosome, we manually selected two independent outgroups that

still had element F as the ancestral X. We first identified homologous chromosomes between

the outgroup and focal species. To do so, we compared the mapping location of *D.*

melanogaster genes on the chromosomes of each focal-outgroup pair of species, and classified

two chromosomes as homologous when they had a more than 1.5-fold excess of observed /

expected shared genes (where the expectation is simply: proportion of mapped genes on the

chromosome of species 1 * proportion of mapped genes on the chromosome of species 2 * total

number of mapped genes). When the location was not in agreement in the two outgroup

species, the gene was not considered further; when it was, this location was assigned as its

putative ancestral location. Gene movement was then inferred when the location in the focal

species was different from the inferred ancestral one.

In order to determine if there was a significant excess of movement out of the X for each

species, we computed the expected number of movements between chromosomes using the

following formula (taken from (Toups et al. 2011), originally proposed by (Betrán et al. 2002)):

$$\frac{\sum N_i L_j f_{ij}}{\sum_i \sum_{j \neq i} N_i L_j f_{ij}}$$

This formula accounts for the number of genes on the chromosome of the parent copy, the size

of the chromosome with the daughter copy, and whether movement occurred to an X

chromosome or an autosome. The indices i and j are for the chromosomes with the parent copy

and with the daughter copy, respectively, N_i is the number of genes on the chromosome with

the parent copy, L_j is the length of the chromosome of the daughter copy, and f_{ij} is 1 when j is

an autosome and 0.75 when j is an X chromosome. We computed the expected gene

movements for each chromosome, and then binned the movements as (1) autosome to

autosome (2) autosome to X and (3) X to autosome. We then determined significance using a

Chi-square goodness of fit test.

Gene expression analysis

A table containing all the expression values stored in FlyBase (Larkin et al. 2020) was

downloaded from their website

(https://s3ftp.flybase.org/releases/FB2025_04/precomputed_files/genes/gene_rpkm_matrix_fb_

2025_04.tsv.gz) and subsetted for columns containing FlyAtlas2 expression (Leader et al. 2018)

or for modENCODE expression (modENCODE Consortium et al. 2010). For each FlyAtlas2

tissue, the expression of *D. melanogaster* homologs of genes that moved out of the X in each

trio was compared to the expression of all mapped tissues with a Wilcoxon test (the resulting p-

values are provided in Supplementary Dataset 6). For tissues of interest (testis, ovary, male and

female head, male and female carcass), comparisons were also performed between genes that

moved into the X and the full mapped set, and between genes that moved between autosomes

and the full mapped set.

Functional enrichment on StringDB

The Flybase accessions of *D. melanogaster* homologs of genes that moved out of the X in each

of the focal species (previous section) were used as input for StringDB (Szklarczyk et al. 2023)

with multiple proteins. Either "*Drosophila melanogaster*" was selected as the reference species,

or the set of D. melanogaster genes that were mapped in each of the trios was given as the

reference background instead. The enriched functional categories obtained with the standard

StringDB analysis with default parameters (as of May 2025) were then downloaded (Table S7 to

S10). A similar analysis was performed with sets of genes inferred to have moved between

autosomes (Table S11 and 12 for Hippoboscidae and Drosophilidae; no enrichment was found

in other lineages).

Data availability

Supplementary datasets and Table S1, S2, S5 and S6 are available at:

https://seafile.ist.ac.at/d/6e69e6be57374552acfe/ [*a permanent URL will be provided upon*

acceptance]. Pipelines are available at https://git.ista.ac.at/llayanaf/transitions_diptera.

Author contributions

MAT and BV designed the original study. LL performed most analyses with support from MAT
and BV. LL, MAT and BV wrote the manuscript.
663

664 **Funding**
665 This work was supported by a grant from the Austrian Science Fund (FWF, grant number PAT
666 8748323) to BV.
667

668 **Conflict of interest statement**
669 The authors declare no conflict of interest.
670

671 **Acknowledgements**
672 We thank the Vicoso group for their feedback on an early version of the manuscript. We are
673 grateful to Kamil Jaron and Julia Gries for helpful discussions and for sharing their unpublished
674 work. Computational resources and support were provided by the Scientific Computing Unit at
675 ISTA.
676

References

Alencar RM et al. 2023. Unravelling the genome of the brackish water malaria vector Anopheles
aquasalis. Sci Rep. 13(1):20472 [accessed 2025 Jun 4].
https://www.nature.com/articles/s41598-023-47830-1. https://doi.org/10.1038/s41598-023-
47830-1

Amabis JM, Reinach FC, Andrews N. 1979. Spermatogenesis in Trichosia pubescens
(Diptera:Sciaridae). J Cell Sci. 36:199–213. https://doi.org/10.1242/jcs.36.1.199

Anderson NW, Hjelmen CE, Blackmon H. 2020. The probability of fusions joining sex
chromosomes and autosomes. Biol Lett. 16(11):20200648.
https://doi.org/10.1098/rsbl.2020.0648

Ansai S et al. 2022. Diversity of sex chromosomes in Sulawesian medaka fishes. J Evol Biol.
35(12):1751–1764. https://doi.org/10.1111/jeb.14076

Baird RB et al. 2025. Faster adaptation but slower divergence of X chromosomes under
paternal genome elimination. Nat Commun. 16(1):5288 [accessed 2025 Oct 23].
https://www.nature.com/articles/s41467-025-60114-8. https://doi.org/10.1038/s41467-025-
60114-8

Baker RH, Wilkinson GS. 2010. Comparative Genomic Hybridization (CGH) reveals a neo-X
chromosome and biased gene movement in stalk-eyed flies (genus Teleopsis). PLoS Genet.
6(9):e1001121. https://doi.org/10.1371/journal.pgen.1001121

Behrens KA et al. 2024. Turnover of sex chromosomes in the Lake Tanganyika cichlid tribe
Tropheini (Teleostei: Cichlidae). Sci Rep. 14(1):2471. https://doi.org/10.1038/s41598-024-
53021-3

Betrán E, Thornton K, Long M. 2002. Retroposed New Genes Out of the X in Drosophila.
Genome Res. 12(12):1854–1859 [accessed 2025 Apr 26].

701 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC187566/. https://doi.org/10.1101/gr.604902

702 Blackmon H, Adams R. 2015. EvobiR: tools for comparative analyses and teaching evolutionary
703 biology. https://github.com/coleoguy/evobir

704 Blaser O, Neuenschwander S, Perrin N. 2014. Sex-chromosome turnovers: the hot-potato
705 model. Am Nat. 183(1):140–146. https://doi.org/10.1086/674026

706 Bracewell R, Chatla K, Nalley MJ, Bachtrog D. 2019. Dynamic turnover of centromeres drives
707 karyotype evolution in Drosophila Henikoff S, Tautz D, Henikoff S, editors. eLife. 8:e49002
708 [accessed 2025 Jun 4]. https://doi.org/10.7554/eLife.49002. https://doi.org/10.7554/eLife.49002

709 Bracewell R, Tran A, Chatla K, Bachtrog D. 2024. Sex and neo-sex chromosome evolution in
710 beetles. PLoS Genet. 20(11):e1011477. https://doi.org/10.1371/journal.pgen.1011477

711 Chakraborty M et al. 2021. Evolution of genome structure in the Drosophila simulans species
712 complex. Genome Res. 31(3):380–396. https://doi.org/10.1101/gr.263442.120

713 Charlesworth D et al. 2020. Using GC Content to Compare Recombination Patterns on the Sex
714 Chromosomes and Autosomes of the Guppy, Poecilia reticulata, and Its Close Outgroup
715 Species. Mol Biol Evol. 37(12):3550–3562. https://doi.org/10.1093/molbev/msaa187

716 Charlesworth D, Charlesworth B. 1980. Sex differences in fitness and selection for centric
717 fusions between sex-chromosomes and autosomes. Genet Res. 35(2):205–214.
718 https://doi.org/10.1017/s0016672300014051

719 Chauhan P et al. 2021. Genome assembly, sex-biased gene expression and dosage
720 compensation in the damselfly Ischnura elegans. Genomics. 113(4):1828–1837.
721 https://doi.org/10.1016/j.ygeno.2021.04.003

722 Coffing GC et al. 2025. Cephalopod sex determination and its ancient evolutionary origin. Curr
723 Biol. 35(4):931-939.e4 [accessed 2025 Apr 26]. https://www.cell.com/current-
724 biology/abstract/S0960-9822(25)00005-3. https://doi.org/10.1016/j.cub.2025.01.005

725 Congrains C et al. 2024. Chromosome-scale genome of the polyphagous pest Anastrepha
726 ludens (Diptera: Tephritidae) provides insights on sex chromosome evolution in Anastrepha. G3
727 GenesGenomesGenetics. 14(12):jkae239 [accessed 2025 Jun 4].
728 https://doi.org/10.1093/g3journal/jkae239. https://doi.org/10.1093/g3journal/jkae239

729 van Doorn GS, Kirkpatrick M. 2007. Turnover of sex chromosomes induced by sexual conflict.
730 Nature. 449(7164):909–912. https://doi.org/10.1038/nature06178

731 Durkin SM et al. 2021. Behavioral and Genomic Sensory Adaptations Underlying the Pest
732 Activity of Drosophila suzukii. Mol Biol Evol. 38(6):2532–2546 [accessed 2025 Jun 4].
733 https://doi.org/10.1093/molbev/msab048. https://doi.org/10.1093/molbev/msab048

734 Eddy SR. 2011. Accelerated Profile HMM Searches. PLOS Comput Biol. 7(10):e1002195
735 [accessed 2025 Nov 4].
736 https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1002195.
737 https://doi.org/10.1371/journal.pcbi.1002195

El Taher A et al. Dynamics of sex chromosome evolution in a rapid radiation of cichlid fishes. Sci Adv. 7(36):eabe8215 [accessed 2025 Apr 26]. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8442896/. https://doi.org/10.1126/sciadv.abe8215

Ellison C, Bachtrog D. 2019. Recurrent gene co-amplification on Drosophila X and Y chromosomes. PLoS Genet. 15(7):e1008251. https://doi.org/10.1371/journal.pgen.1008251

Emerson JJ, Kaessmann H, Betrán E, Long M. 2004. Extensive gene traffic on the mammalian X chromosome. Science. 303(5657):537–540. https://doi.org/10.1126/science.1090042

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome Biol. 20(1):238 [accessed 2021 Apr 18]. https://doi.org/10.1186/s13059-019-1832-y. https://doi.org/10.1186/s13059-019-1832-y

Evans BJ et al. 2024. Rapid Sex Chromosome Turnover in African Clawed Frogs (Xenopus) and the Origins of New Sex Chromosomes. Mol Biol Evol. 41(12):msae234. https://doi.org/10.1093/molbev/msae234

Falk S, Grzywacz A. 2024. The genome sequence of a muscid fly, Hydrotaea diabolus (Harris, [1780]). Wellcome Open Res. [published online ahead of print] [accessed 2025 Oct 23]. https://doi.org/10.12688/wellcomeopenres.21211.1

Falk S, Sivell D, Webb J, Grzywacz A. 2024. The genome sequence of a muscid fly, Polietes domitor (Harris, 1780). Wellcome Open Res. 9:58 [accessed 2025 Oct 23]. https://pmc.ncbi.nlm.nih.gov/articles/PMC11411267/. https://doi.org/10.12688/wellcomeopenres.20648.1

Graves JAM. 2016. Evolution of vertebrate sex chromosomes and dosage compensation. Nat Rev Genet. 17(1):33–46. https://doi.org/10.1038/nrg.2015.2

Hemmer LW et al. 2020. Hybrid dysgenesis in Drosophila virilis results in clusters of mitotic recombination and loss-of-heterozygosity but leaves meiotic recombination unaltered. Mob DNA. 11(1):10 [accessed 2025 Jun 4]. https://doi.org/10.1186/s13100-020-0205-0. https://doi.org/10.1186/s13100-020-0205-0

Hill T, Koseva BS, Unckless RL. 2019. The Genome of Drosophila innubila Reveals Lineage-Specific Patterns of Selection in Immune Genes. Mol Biol Evol. 36(7):1405–1417 [accessed 2025 Jun 4]. https://doi.org/10.1093/molbev/msz059. https://doi.org/10.1093/molbev/msz059

Iwasaki M et al. 2021. Identification of the sex-determining factor in the liverwort Marchantia polymorpha reveals unique evolution of sex chromosomes in a haploid system. Curr Biol CB. 31(24):5522-5532.e7. https://doi.org/10.1016/j.cub.2021.10.023

Jayaprasad S et al. 2024. Orthopteran Neo-Sex Chromosomes Reveal Dynamics of Recombination Suppression and Evolution of Supergenes. Mol Ecol. 33(23):e17567. https://doi.org/10.1111/mec.17567

Jeffries DL et al. 2018. A rapid rate of sex-chromosome turnover and non-random transitions in true frogs. Nat Commun. 9(1):4088. https://doi.org/10.1038/s41467-018-06517-2

Katoh K, Misawa K, Kuma K, Miyata T. 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. 30(14):3059–3066
[accessed 2022 Apr 3]. https://doi.org/10.1093/nar/gkf436. https://doi.org/10.1093/nar/gkf436

Kent WJ. 2002. BLAT--the BLAST-like alignment tool. Genome Res. 12(4):656–664.
https://doi.org/10.1101/gr.229202

Konganti K et al. 2018. A Whole Genome Assembly of the Horn Fly, Haematobia irritans, and
Prediction of Genes with Roles in Metabolism and Sex Determination. G3
GenesGenomesGenetics. 8(5):1675–1686 [accessed 2025 Jun 4].
https://doi.org/10.1534/g3.118.200154. https://doi.org/10.1534/g3.118.200154

Kumar S, Stecher G, Suleski M, Hedges SB. 2017. TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. Mol Biol Evol. 34(7):1812–1819.
https://doi.org/10.1093/molbev/msx116

Larkin A et al. 2020. FlyBase: updates to the Drosophila melanogaster knowledge base. Nucleic
Acids Res. 49(D1):D899–D907 [accessed 2025 Nov 4].
https://pmc.ncbi.nlm.nih.gov/articles/PMC7779046/. https://doi.org/10.1093/nar/gkaa1026

Lasne C et al. 2023. The Scorpionfly (Panorpa cognata) Genome Highlights Conserved and
Derived Features of the Peculiar Dipteran X Chromosome. Mol Biol Evol. 40(12):msad245.
https://doi.org/10.1093/molbev/msad245

Leader DP et al. 2018. FlyAtlas 2: a new version of the Drosophila melanogaster expression
atlas with RNA-Seq, miRNA-Seq and sex-specific data. Nucleic Acids Res. 46(D1):D809–D815
[accessed 2022 Nov 3]. https://doi.org/10.1093/nar/gkx976. https://doi.org/10.1093/nar/gkx976

Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
tree display and annotation tool. Nucleic Acids Res. 52(W1):W78–W82.
https://doi.org/10.1093/nar/gkae268

Leung W et al. 2023. Long-read genome assemblies for the study of chromosome expansion:
Drosophila kikkawai, Drosophila takahashii, Drosophila bipectinata, and Drosophila ananassae.
G3 GenesGenomesGenetics. 13(10):jkad191 [accessed 2025 Jun 4].
https://doi.org/10.1093/g3journal/jkad191. https://doi.org/10.1093/g3journal/jkad191

Levy Karin E, Mirdita M, Söding J. 2020. MetaEuk-sensitive, high-throughput gene discovery,
and annotation for large-scale eukaryotic metagenomics. Microbiome. 8(1):48.
https://doi.org/10.1186/s40168-020-00808-x

Li D et al. 2015. MEGAHIT: an ultra-fast single-node solution for large and complex
metagenomics assembly via succinct de Bruijn graph. Bioinformatics. 31(10):1674–1676
[accessed 2022 Apr 3]. https://doi.org/10.1093/bioinformatics/btv033.
https://doi.org/10.1093/bioinformatics/btv033

Li H. 2023. Protein-to-genome alignment with miniprot. Bioinformatics. 39(1):btad014 [accessed
2025 Nov 4]. https://doi.org/10.1093/bioinformatics/btad014.
https://doi.org/10.1093/bioinformatics/btad014

Li X, Mank JE, Ban L. 2024. The grasshopper genome reveals long-term gene content

conservation of the X Chromosome and temporal variation in X Chromosome evolution. Genome Res. 34(7):997–1007. https://doi.org/10.1101/gr.278794.123

Loeschcke V, Nielsen BO, Andersen D. 1994. Relationship Among Hydrotaea Species Based on Allozymes, Karyotype and Morphology (Diptera: Muscidae). Hereditas. 121(1):103–111 [accessed 2025 Oct 23]. https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1601-5223.1994.00103.x. https://doi.org/10.1111/j.1601-5223.1994.00103.x

Lukyanchikova V et al. 2022. Anopheles mosquitoes reveal new principles of 3D genome organization in insects. Nat Commun. 13(1):1960 [accessed 2025 Jun 4]. https://www.nature.com/articles/s41467-022-29599-5. https://doi.org/10.1038/s41467-022-29599-5

Ma W-J, Veltsos P. 2021. The Diversity and Evolution of Sex Chromosomes in Frogs. Genes. 12(4):483. https://doi.org/10.3390/genes12040483

Mahajan S et al. 2018. De novo assembly of a young Drosophila Y chromosome using single-molecule sequencing and chromatin conformation capture. PLOS Biol. 16(7):e2006348 [accessed 2025 Jun 4]. https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.2006348. https://doi.org/10.1371/journal.pbio.2006348

Mai D, Nalley MJ, Bachtrog D. 2020. Patterns of Genomic Differentiation in the Drosophila nasuta Species Complex. Mol Biol Evol. 37(1):208–220 [accessed 2025 Jun 4]. https://doi.org/10.1093/molbev/msz215. https://doi.org/10.1093/molbev/msz215

Manni M et al. 2021. BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 38(10):4647–4654 [accessed 2022 Apr 3]. https://doi.org/10.1093/molbev/msab199. https://doi.org/10.1093/molbev/msab199

Marshall Graves JA, Peichel CL. 2010. Are homologies in vertebrate sex determination due to shared ancestry or to limited options? Genome Biol. 11(4):205. https://doi.org/10.1186/gb-2010-11-4-205

Massey JH et al. 2020. Co-evolving wing spots and mating displays are genetically separable traits in Drosophila. Evolution. 74(6):1098–1111 [accessed 2025 Jun 4]. https://onlinelibrary.wiley.com/doi/abs/10.1111/evo.13990. https://doi.org/10.1111/evo.13990

Matsumoto T, Kitano J. 2016. The intricate relationship between sexually antagonistic selection and the evolution of sex chromosome fusions. J Theor Biol. 404:97–108. https://doi.org/10.1016/j.jtbi.2016.05.036

Meisel RP et al. 2020. Sex Chromosome Evolution in Muscid Flies. G3 GenesGenomesGenetics. 10(4):1341–1352 [accessed 2025 Apr 26]. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7144080/. https://doi.org/10.1534/g3.119.400923

Meisel RP et al. 2023. New insights into immune genes and other expanded gene families of the house fly, Musca domestica, from an improved whole genome sequence. Arch Insect Biochem Physiol. 114(3):e22049 [accessed 2025 Jun 4]. https://onlinelibrary.wiley.com/doi/abs/10.1002/arch.22049. https://doi.org/10.1002/arch.22049

Meisel RP, Delclos PJ, Wexler JR. 2019. The X chromosome of the German cockroach,
Blattella germanica, is homologous to a fly X chromosome despite 400 million years divergence.
BMC Biol. 17(1):100. https://doi.org/10.1186/s12915-019-0721-x

Miller D et al. 2022. Retrogene Duplication and Expression Patterns Shaped by the Evolution of
Sex Chromosomes in Malaria Mosquitoes. Genes. 13(6):968.
https://doi.org/10.3390/genes13060968

modENCODE Consortium et al. 2010. Identification of functional elements and regulatory
circuits by Drosophila modENCODE. Science. 330(6012):1787–1797.
https://doi.org/10.1126/science.1198374

Mohr OL. 1932. GENETICAL AND CYTOLOGICAL PROOF OF SOMATIC ELIMINATION OF
THE FOURTH CHROMOSOME IN DROSOPHILA MELANOGASTER. Genetics. 17(1):60–80
[accessed 2025 Oct 23]. https://doi.org/10.1093/genetics/17.1.60.
https://doi.org/10.1093/genetics/17.1.60

Mora P et al. 2024. Sex-biased gene content is associated with sex chromosome turnover in
Danaini butterflies. Mol Ecol. 33(24):e17256. https://doi.org/10.1111/mec.17256

POKORNÁ M, KRATOCHVÍL L. 2009. Phylogeny of sex-determining mechanisms in squamate
reptiles: are sex chromosomes an evolutionary trap? Zool J Linn Soc. 156(1):168–183
[accessed 2025 Apr 26]. https://doi.org/10.1111/j.1096-3642.2008.00481.x.
https://doi.org/10.1111/j.1096-3642.2008.00481.x

Potrzebowski L et al. 2008. Chromosomal Gene Movements Reflect the Recent Origin and
Biology of Therian Sex Chromosomes. PLoS Biol. 6(4):e80 [accessed 2025 Apr 26].
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2276528/.
https://doi.org/10.1371/journal.pbio.0060080

Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees for
large alignments. PloS One. 5(3):e9490. https://doi.org/10.1371/journal.pone.0009490

Reinhardt JA et al. 2023. Impacts of Sex Ratio Meiotic Drive on Genome Structure and Function
in a Stalk-Eyed Fly. Genome Biol Evol. 15(7):evad118 [accessed 2025 Jun 4].
https://doi.org/10.1093/gbe/evad118. https://doi.org/10.1093/gbe/evad118

Renschler G et al. 2019. Hi-C guided assemblies reveal conserved regulatory topologies on X
and autosomes despite extensive genome shuffling. Genes Dev. 33(21–22):1591–1612.
https://doi.org/10.1101/gad.328971.119

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other
things). Methods Ecol Evol. 3(2):217–223 [accessed 2025 Oct 23].
https://onlinelibrary.wiley.com/doi/abs/10.1111/j.2041-210X.2011.00169.x.
https://doi.org/10.1111/j.2041-210X.2011.00169.x

Revell LJ, Schliep KP, Mahler DL, Ingram T. 2024. Testing for heterogeneous rates of discrete
character evolution on phylogenies. J Evol Biol. 37(12):1591–1602 [accessed 2025 Oct 23].
https://doi.org/10.1093/jeb/voae119. https://doi.org/10.1093/jeb/voae119

Richards S et al. 2005. Comparative genome sequencing of Drosophila pseudoobscura:

chromosomal, gene, and cis-element evolution. Genome Res. 15(1):1–18.
https://doi.org/10.1101/gr.3059305

Riddle NC, Elgin SCR. 2018. The Drosophila Dot Chromosome: Where Genes Flourish Amidst
Repeats. Genetics. 210(3):757–772 [accessed 2025 Jun 4].
https://doi.org/10.1534/genetics.118.301146. https://doi.org/10.1534/genetics.118.301146

Romine MG et al. 2022. The genome sequence of the avian vampire fly (Philornis downsi), an
invasive nest parasite of Darwin's finches in Galápagos. G3 GenesGenomesGenetics.
12(2):jkab414 [accessed 2025 Jun 4]. https://doi.org/10.1093/g3journal/jkab414.
https://doi.org/10.1093/g3journal/jkab414

Satomura K, Osada N, Endo T. 2019. Achiasmy and sex chromosome evolution. Ecol Genet
Genomics. 13:100046 [accessed 2025 Oct 23].
https://www.sciencedirect.com/science/article/pii/S2405985418300351.
https://doi.org/10.1016/j.egg.2019.100046

Schaeffer SW. 2018. Muller "Elements" in Drosophila: How the Search for the Genetic Basis for
Speciation Led to the Birth of Comparative Genomics. Genetics. 210(1):3–13 [accessed 2025
Apr 26]. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6116959/.
https://doi.org/10.1534/genetics.118.301084

Sharma A et al. 2017. Male sex in houseflies is determined by Mdmd, a paralog of the generic
splice factor gene CWC22. Science. 356(6338):642–645.
https://doi.org/10.1126/science.aam5498

Sigeman H et al. 2021. Avian Neo-Sex Chromosomes Reveal Dynamics of Recombination
Suppression and W Degeneration. Mol Biol Evol. 38(12):5275–5291.
https://doi.org/10.1093/molbev/msab277

Singh ND, Davis JC, Petrov DA. 2005. Codon bias and noncoding GC content correlate
negatively with recombination rate on the Drosophila X chromosome. J Mol Evol. 61(3):315–
324. https://doi.org/10.1007/s00239-004-0287-1

Small ST et al. 2020. Radiation with reticulation marks the origin of a major malaria vector. Proc
Natl Acad Sci. 117(50):31583–31590 [accessed 2025 Jun 4].
https://www.pnas.org/doi/10.1073/pnas.2018142117. https://doi.org/10.1073/pnas.2018142117

Sperling AL, Glover DM. 2023. Parthenogenesis in dipterans: a genetic perspective. Proc Biol
Sci. 290(1995):20230261. https://doi.org/10.1098/rspb.2023.0261

Sturgill D, Zhang Y, Parisi M, Oliver B. 2007. Demasculinization of X chromosomes in the
Drosophila genus. Nature. 450(7167):238–241. https://doi.org/10.1038/nature06330

Szklarczyk D et al. 2023. The STRING database in 2023: protein-protein association networks
and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Res.
51(D1):D638–D646. https://doi.org/10.1093/nar/gkac1000

Tandonnet S et al. 2023. A chromosomal-scale reference genome of the New World
Screwworm, Cochliomyia hominivorax. DNA Res. 30(1):dsac042 [accessed 2025 Jun 4].
https://doi.org/10.1093/dnares/dsac042. https://doi.org/10.1093/dnares/dsac042

The Darwin Tree of Life Project Consortium. 2022. Sequence locally, think globally: The Darwin
Tree of Life Project. Proc Natl Acad Sci. 119(4):e2115642118 [accessed 2025 Jun 4].
https://www.pnas.org/doi/full/10.1073/pnas.2115642118.
https://doi.org/10.1073/pnas.2115642118

Torosin NS et al. 2020. 3D genome evolution and reorganization in the Drosophila
melanogaster species group. PLOS Genet. 16(12):e1009229 [accessed 2025 Jun 4].
https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1009229.
https://doi.org/10.1371/journal.pgen.1009229

Toups MA, Hahn MW. 2010. Retrogenes reveal the direction of sex-chromosome evolution in
mosquitoes. Genetics. 186(2):763–766. https://doi.org/10.1534/genetics.110.118794

Toups MA, Pease JB, Hahn MW. 2011. No excess gene movement is detected off the avian or
lepidopteran Z chromosome. Genome Biol Evol. 3:1381–1390.
https://doi.org/10.1093/gbe/evr109

Toups MA, Rodrigues N, Perrin N, Kirkpatrick M. 2019. A reciprocal translocation radically
reshapes sex-linked inheritance in the common frog. Mol Ecol. 28(8):1877–1889 [accessed
2025 Apr 29]. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7558804/.
https://doi.org/10.1111/mec.14990

Toups MA, Vicoso B. 2023. The X chromosome of insects likely predates the origin of class
Insecta. Evol Int J Org Evol. 77(11):2504–2511. https://doi.org/10.1093/evolut/qpad169

Trinca V et al. 2023. Biocatalytic potential of Pseudolycoriella CAZymes (Sciaroidea, Diptera) in
degrading plant and fungal cell wall polysaccharides. iScience. 26(4):106449.
https://doi.org/10.1016/j.isci.2023.106449

Urban JM et al. 2021. High contiguity de novo genome assembly and DNA modification
analyses for the fungus fly, Sciara coprophila, using single-molecule sequencing. BMC
Genomics. 22(1):643 [accessed 2025 Jun 4]. https://doi.org/10.1186/s12864-021-07926-2.
https://doi.org/10.1186/s12864-021-07926-2

Vibranovski MD, Lopes HF, Karr TL, Long M. 2009. Stage-Specific Expression Profiling of
Drosophila Spermatogenesis Suggests that Meiotic Sex Chromosome Inactivation Drives
Genomic Relocation of Testis-Expressed Genes. PLOS Genet. 5(11):e1000731 [accessed 2022
Mar 28]. https://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1000731.
https://doi.org/10.1371/journal.pgen.1000731

Vicoso B. 2019. Molecular and evolutionary dynamics of animal sex-chromosome turnover. Nat
Ecol Evol. 3(12):1632–1641 [accessed 2022 Mar 28]. https://www.nature.com/articles/s41559-
019-1050-8. https://doi.org/10.1038/s41559-019-1050-8

Vicoso B, Bachtrog D. 2015. Numerous transitions of sex chromosomes in Diptera. PLoS Biol.
13(4):e1002078. https://doi.org/10.1371/journal.pbio.1002078

Wei KH-C, Mai D, Chatla K, Bachtrog D. 2022. Dynamics and Impacts of Transposable Element
Proliferation in the Drosophila nasuta Species Group Radiation. Mol Biol Evol. 39(5):msac080
[accessed 2025 Jun 4]. https://doi.org/10.1093/molbev/msac080.
https://doi.org/10.1093/molbev/msac080

Wiegmann BM et al. 2011. Episodic radiations in the fly tree of life. Proc Natl Acad Sci U S A.
108(14):5690–5695. https://doi.org/10.1073/pnas.1012675108

Wright CJ et al. 2024. Comparative genomics reveals the dynamics of chromosome evolution in
Lepidoptera. Nat Ecol Evol. 8(4):777–790. https://doi.org/10.1038/s41559-024-02329-4

Zamyatin A et al. 2021. Chromosome-level genome assemblies of the malaria vectors
Anopheles coluzzii and Anopheles arabiensis. GigaScience. 10(3):giab017 [accessed 2025 Jun
4]. https://doi.org/10.1093/gigascience/giab017. https://doi.org/10.1093/gigascience/giab017

Zdobnov EM et al. 2002. Comparative genome and proteome analysis of Anopheles gambiae
and Drosophila melanogaster. Science. 298(5591):149–159.
https://doi.org/10.1126/science.1077061