Manuscript title:

Patterns of aDNA Damage Through Time end Environments – lessons from herbarium

specimens

Authors:

Stefano Porrelli[1,*], ORCID: 0000-0003-3878-7745

Alice Fornasiero[2], ORCID: 0000-0001-6165-4233

Phuong Hong Le[1], ORCID: 0000-0002-2862-6502

Whenzhe Yin[3,4], ORCID: 0000-0002-4350-3578

Maria Navarrete Rodriguez[2], ORCID: 0000-0001-5052-3713

Nahed Mohammed[2], ORCID: 0000-0002-8857-3246

Axel Himmelbach[5], ORCID: 0000-0001-7338-0946

Andrew C. Clarke[3], ORCID: 0000-0003-2293-1423

Nils Stein[5,6], ORCID: 0000-0003-3011-8731

Paul J. Kersey[1], ORCID: 0000-0002-7054-800X

Rod A. Wing[2,7] ORCID: 0000-0001-6633-6226

Rafal M. Gutaker[1,*], ORCID: 0000-0001-9226-879X

[1] Trait Diversity and Function, Royal Botanic Gardens, Kew, Richmond, TW9 3AE, United

Kingdom.

[2] Plant Science Program, Biological and Environmental Science and Engineering Division,

King Abdullah University of Science and Technology (KAUST), Thuwal, Saudi Arabia.

[3] School of Biosciences, University of Nottingham, Sutton Bonington, LE12 5RD, United

Kingdom.

[4] State Key Laboratory of Emerging Infectious Diseases, School of Public Health, The

University of Hong Kong, Hong Kong, SAR, China.

[5] Genebank, Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Seeland,

06466, Germany.

[6] Institute of Agricultural and Nutritional Sciences, Martin Luther University of Halle-

Wittenberg, Halle (Saale), Germany.

[7] Arizona Genomics Institute, School of Plant Sciences, University of Arizona, Tucson, AZ,

USA.

* To whom correspondence should be addressed: R.Gutaker@kew.org; S.Porrelli2@kew.org

Abstract:

Herbarium collections are a vast but underutilized resource for ancient DNA research,

containing over 400 million specimens with detailed metadata and spanning centuries of global

biodiversity. Understanding patterns of DNA preservation in natural collections is crucial for

optimizing ancient DNA studies and informing future curation practices. We analysed genomic

data for 573 herbarium specimens from six plant species from the genera *Hordeum* and *Oryza*

collected from the Americas and Eurasia over 220 years using standardized laboratory

protocols and shotgun sequencing to quantify DNA degradation and elucidate factors that

accelerate it. We find significant age-dependent DNA fragmentation rates, indicating temporal

degradation processes not detected in prehistoric samples. In our analysis, DNA decay rates in

herbarium specimens were almost eight times faster than in moa bones, reflecting fundamental

differences in tissue composition and preservation environments. Environmental conditions at

the time of specimen collection emerged as the major determinants of post-mortem mutation

rates, with temperature being the dominant driver of cytosine deamination. We find no effect

of sample storage on DNA degradation. These findings provide insights into how climatic

origin, preservation environment, taxonomic identity and age influence DNA preservation

while highlighting opportunities for improving institutional preservation practices. Due to

standardized preservation conditions, museum collections can provide better insights into DNA

degradation over time than archaeological and paleontological samples.

Introduction:

Understanding preservation of DNA from old biological samples, commonly referred to as

ancient DNA (aDNA), has been at the core of the genomic revolution in archaeological research

[1,2]. There is a consensus among researchers that aDNA is defined by its degradation and not

by its age [3,4], though the two are correlated [5]. The two most prominent reactions associated

with DNA degradation are deamination (resulting with spontaneous substitutions of cytosine

residues to uracil) and depurination (breakage of the phosphodiester bond resulting in DNA

backbone fragmentation). Both reactions occur spontaneously in DNA in the absence of

enzymatic repair machinery [6,7]. Deamination leads to a characteristic pattern of 'C>T'

substitutions in DNA fragments, with increased frequency at fragment termini, while

depurination leads to DNA fragmentation towards very small molecules (~30-100 bp). Both

patterns are common in historical, archaeological and sedimentary samples [2,5,8] and make

bioinformatic processing and downstream analyses challenging [9]. Although C>T

misincorporations can be used to authenticate genuine aDNA sequences, they can bias variant

calling and phylogenetic analyses, potentially leading to incorrect inferences [3]. Similarly,

extensive DNA fragmentation reduces mapping efficiency and increases the likelihood of

spurious alignments [10,11]. These challenges are particularly acute for ancient samples from

hot and humid environments, where accelerated degradation processes further compromise

DNA integrity [12,13]. As a result, tools and approaches have been developed to improve

quality control, reads processing, mapping and downstream analyses in a quest to utilize even highly degraded samples [14–16]. In addition to chemical degradation, DNA can also be altered through biological processes such as microbial colonization. In effect, the *bona fide* DNA from the target species can be depleted and replaced with post-mortem microbial DNA [17].

While the vast majority of published aDNA research focuses on human skeletal remains [18,19], there are an increasing number of studies in other mammals [20], arthropods [8,21], and in plants [22,23]. Archaeobotanical materials, primarily seeds, are often used as sources of prehistoric DNA. While not older than 500 years, herbarium specimens number over 400 million [24], and have solid species identification and, frequently, collection metadata. They comprise a vast but underutilised resource for studying evolutionary and ecological changes in the last couple of centuries [25,26]. The leaf tissue is most often the target in DNA isolation protocols but does not possess the same structural isolation from the environment as bones or seeds. This is the most likely reason why DNA in herbarium samples decays at a rate roughly six times faster than in ancient Moa bones [27], but also more than twice as slow when compared to dry-pinned arthropod museum specimens [8]. On the other hand, herbarium specimens are commonly dried upon collection and are generally stored in standardized conditions that are favourable for DNA preservation, such as stable temperatures and low air humidity. Availability of materials, good geolocation information and relatively stable preservation condition ex situ make herbarium specimens a perfect system to study the effects of the environment at the point of collection on the subsequent preservation of DNA in biological samples.

One of the major challenges in studies trying to investigate the dynamics of DNA mutation, decay and fragmentation is lack of large datasets with a global distribution and consistent sampling and laboratory processing procedures. Previous studies were limited to one or two

species representing narrow geographical range [12,27]. Here, we present new sequencing data

for a total of 573 herbarium samples from 6 plant species, spanning the Americas and Eurasia,

processed with the same laboratory protocol and sequenced using a whole-genome shotgun

approach. Our main aim is to understand which environmental factors influence the rate of

DNA mutation and decay. The outcome of this investigation brings new insights into the

fundamental processes of DNA degradation that can inform future conservation of museum

specimens, as well as help researchers to prioritize materials for their studies.

Methods

Herbarium samples

Historical herbarium samples ($N = 573$) were obtained from seven herbaria in Europe and

North America (table 1): Royal Botanic Gardens, Kew (UK), Smithsonian Institute Herbarium

(US), Missouri Botanic Gardens (US), New York Botanic Gardens (US), National History

Museum (France), Harvard University Herbarium (US), and National History Museum (UK).

Passport information for each sample is included in the supplementary material (supplementary

table S1). The *Hordeum* samples were predominantly collected from temperate Europe to semi-

arid regions of the Middle East, with highest counts from countries such as Iraq, Israel and

Iran. The *Oryza* samples were predominantly collected from tropical and subtropical regions

of the Americas and Southeast Asia, particularly Brazil, Costa Rica, Colombia, Mexico and

Thailand (figure 1).

Table 1: Number of historical herbarium samples ($N = 573$) included in this study collected

from seven herbaria. RBGK: Royal Botanic Gardens Kew, UK; SIHUS: Smithsonian Institute

Herbarium, US; MBG: Missouri Botanic Gardens, US; NYBGUS: New York Botanic Gardens,

US; NHMFR: National History Museum, France; HUHUS: Harvard University Herbarium,

US; NHMUK: National History Museum, UK.

Species	RBGK	SIHUS	MBG	NYBGUS	NHMFR	HUHUS	NHMUK	Total	Age range
Hordeum spontaneum	146	0	0	0	0	0	0	146	1822-2017
Hordeum vulgare	82	0	0	0	0	0	0	82	1819-2015
Oryza alta	0	2	9	0	2	0	0	13	1913-2002
Oryza grandiglumis	1	30	1	0	6	0	0	38	1930-2002
Oryza latifolia	22	97	36	30	20	16	10	231	1797-2011
Oryza rufipogon	29	31	3	0	0	0	0	63	1909-1998
Total	280	160	49	30	28	16	10	573	1797-2017

Figure 1: Overview of the herbarium samples of barley (*Hordeum*) and rice (*Oryza*) species:

(a) geographical distribution based on country of collection; (b) source of herbarium of

Hordeum and *Oryza* species (RBGK: Royal Botanic Gardens Kew, UK; SIHUS: Smithsonian

Institute Herbarium, US; MBG: Missouri Botanic Gardens, US; NYBGUS: New York Botanic

Gardens, US; NHMFR: National History Museum, France; HUHUS: Harvard University

Herbarium, US; NHMUK: National History Museum, UK); (c) temporal distribution for each

species.

DNA extraction, library preparation and low throughput sequencing:

Laboratory steps for DNA extraction and library preparation for all 228 *Hordeum* (146 *H.*

spontaneum and 82 *H. vulgare)* and 58 Oryza (30 *O. latifolia,* 18 *O. rufipogon* and 10 *O.*

grandiglumis) (Table 1) samples were carried out in clean room facilities at Royal Botanic

Gardens, Kew (UK) following previously published best practices and aDNA protocol

described in [28] with adjusted volumes. Tissue samples (5-10 mg) were grinded in 2mL

PowerBead metal tubes (QIAGEN, 13117-50) with a bead mill homogeniser (Precellys

Evolution, P002511-PEVT0-A.0). 1mL of PTB-based mix was added to the homogenized

samples, which were then incubated on a rotor overnight at 37°C. gDNA was isolated and half

the volume (0.5 mL) was purified using the DNeasy Plant Mini Kit (QIAGEN, 69106) with

modifications described in [28], and quantified with Quantus™ fluorometer (Promega, E6150).

For a subset of the gDNA isolates (*N*=40, 10 *H. spontaneum*, 10 *H. vulgare*, 10 *O. rufipogon*,

and 10 *O. latifolia*; supplementary table S2), the genomic DNA fragment size distribution was

analysed with the 4200 TapeStation system (Agilent, G2991BA) using a D1000 ScreenTape

(Agilent, 5067-5582).

Double stranded and doubled indexed genomic libraries were constructed by blunt-end DNA

ligation [29,30] following [28]. Briefly, genomic DNA was blunt-end repaired, and universal

Illumina double-stranded adapters were ligated. This was followed by fill-in of adapters,

indexing with unique combinatorial Illumina indexes, and amplification. Between each step,

samples were purified with the MiniElute® PCR purification kit (QIAGEN, 28006). To

quantify the excess of C>T substitutions at reads termini caused by spontaneous deamination

of cytosines [31], and to validate aDNA authenticity [27,32], we did not perform enzymatic

removal of aDNA-associated DNA misincorporation [33]. Indexed and amplified libraries

were quantified with Quantus™ fluorometer (Promega, E6150). fragment size of each library

were estimated with a 4200 TapeStation System (Agilent, G2991BA) using a D1000

ScreenTape (Agilent, 5067-5582). Libraries were pooled equimolarly and sequenced in paired-

end mode. All *Hordeum* libraries were sequenced on a NovaSeq 6000 or NovaSeq X+ device

(2 x 62 bp) according to manufacturer's instructions (Illumina, San Diego, CA, USA) at the

Leibniz Institute of Plant Genetics and Crop Plant Research (IPK, Gatersleben, Germany). 40

Oryza libraries were sequenced at Macrogen (Macrogen, Europe), on an Illumina Novaseq X+

system (2 x 150 bp). A further 287 *Oryza* samples were obtained with fundings from the King

Abdullah University of Science and Technology (KAUST). These samples were processed in

a clean room facility at the Ancient and Environmental DNA Laboratory (ÆDNA) at the

University of Nottingham (UoN, UK) following the same protocol, and sequenced on an

Illumina Miseq platform (2 x 150 bp) according to manufacturer's instructions (Illumina, San

Diego, CA, USA) at the Deep Seq facility at UoN.

Bioinformatic screening and validation of aDNA authenticity

Bioinformatics protocols for read processing, quality assessment, screening and authentication

of aDNA-derived libraries follow [28]. Demultiplexed raw Illumina paired-end reads were

adapter trimmed and merged with AdapterRemoval v2.3.4 [34] with a minimum overlap of 11

bases,   and   quality-checked   with   FastQC   v0.11.8

(http://www.bioinformatics.bbsrc.ac.uk/projects/fastqc). Reference genome assemblies were

retrieved from the National Center for Biotechnology Information (NCBI) database (table 2)

and indexed with BWA v0.7.19 [35] and SAMtools v1.21 [36]. Trimmed and merged reads

were mapped to their respective reference genome with BWA-aln using aDNA-specific

parameters ("-l 1024"). PCR optical duplicate reads removed with DeDup [16]. Base

frequencies at DNA break points and nucleotide misincorporations were estimated for each

library using MapDamage2 v2.0.6 [37]. Mapping statistics of the additional 287 *Oryza* samples

were obtained by mapping sequencing reads to the following reference genomes: *O. alta*

(PRJNA1039467), *O. grandiglumis* (PRJNA737282), *O. latifolia* (PRJNA737486) [38]. The

O. rufipogon samples originally collected in the American continent were aligned to the *O.*

glumipatula genome reference (PRJNA48429).

Table 2: Reference genome assemblies used in this study with associated BioProject,

BioSample and Accession IDs.

Reference	BioProject	BioSample	Assembly Accession
Hordeum spontaneum	PRJEB57567	SAMEA112465237	GCA_949783245.1
Hordeum vulgare	PRJEB40589	SAMEA7384724	GCF_904849725.1
Oryza alta	PRJNA1039467	SAMN38217704	GCA_047899615.1
Oryza grandiglumis	PRJNA737282	SAMN19687255	GCA_048188845.1
Oryza latifolia	PRJNA737486	SAMN19696891	GCA_048174585.1
Oryza rufipogon	PRJNA1029807	SAMN40302812	GCA_037997075.1
Oryza glumipatula	PRJNA48429	SAMN02981440	GCA_000576495.2

Ancient DNA damage metrics and regression analyses

Four metrics were selected to quantify patterns of aDNA damage: (i) the proportion of

endogenous DNA content, (ii) the fragment length distribution, (iii) the damage fraction per

site (λ), and (iv) the frequencies of 5' C>T substitutions. The four metrics were analysed in

linear models as a function of collection year and sample age using the '*lm*' function in R [39].

The DNA decay rates (k) of *Hordeum* and *Oryza* samples were also determined independently.

Endogenous fraction

The percentage of post-quality trimmed and merged reads for each sample mapped to its

respective reference was used as a proxy for the proportion of endogenous DNA content.

Endogenous content was calculated with SAMtools "*flagstat*" tool [36] and plotted as a

function of collection year.

Fragment length

We quantified DNA fragmentation using two approaches. Firstly, to validate that the fragment

sizes of the merged reads reflects the original molecule length, we examined the relationship

between the fragment size distribution of isolated gDNA and that of the amplified libraries for

a subset of the samples using TapeStation profiles for both (*N*= 40; 10 *H. spontaneum*, 10 *H.*

vulgare, 10 *O. rufipogon*, and 10 *O. latifolia*; supplementary table S2). Library and gDNA peak

size measurements from the TapeStation profiles were compared with each other and with

bioinformatically-derived median fragment lengths of merged reads to assess the concordance

between direct physical measurements and computational estimates. In addition, we further

analysed the correlation between peak sizes of the gDNA isolates and age of the sample.

Secondly, for the whole dataset (*N*= 573), we fitted the fragment length distribution of the

mapped merged reads to a lognormal distribution using the 'fitdistr' function from the MASS

package [40] in R. We used the mean of this distribution (log-mean) to summarise fragment

length for each library and carried out regressions on the relationship between the log-mean of

fragment lengths and collection year. As suggested in [27], we used the median plotted on a

log-scaled *y*-axis for visualisation, as the latter is more intuitive to understand than the log-

mean.

Damage fraction per site (λ) and DNA decay rate (k)

The damage fraction per site (λ) was calculated for each sample using previously described

methods [12]. We fitted an exponential decay model to the empirical fragment length

distribution of the mapped reads and derived λ as the negative of the slope coefficient of the

linear regression. To identify samples deviating from the assumption of exponential decay of

fragment length, we assessed goodness of fit and statistical significance on a per-sample basis.

Model fit was considered satisfactory when $R^2 > 0.95$ and $p < 0.05$.

Having determined the damage fraction per site (λ), we calculated genus-specific DNA decay

rates (*k*) by plotting λ as a function of sample age and extrapolating *k* from the slope of the

linear regression, according to the linear relationship $k = \lambda/age$ [41].

Nucleotide misincorporations

The frequency of 5' C>T substitutions at first position was used as a proxy for 5' damage.

Although the depth of sequencing needed for reliable estimation of C>T frequencies is sample-

dependent, an order of magnitude of thousands of reads is considered sufficient [42]. Therefore,

only samples with >5,000 merged paired reads were retained for further analyses. Furthermore,

the exponential increase of C>T substitutions at reads termini is routinely used as a metric to

validate authenticity of aDNA [5]. We scored this pattern for each sample by fitting an

exponential model to the C>T frequencies for the first 20 bases at 5' terminus. We evaluated

the goodness of fit with a one-sided *t*-test on a per-sample basis, as described in [43], and only

retained samples that fitted the exponential model ($R^2 > 0.5$, $p < 0.05$).

Analysis of covariance

To investigate differences in aDNA damage patterns between *Hordeum* and *Oryza*, we

performed Analysis of Covariance (ANCOVA) using the 'anova' function in R. For each aDNA

damage metric, sample age was used as the covariate and the genus as the factor using the

model "*y ~ covariate* x *factor",* which also tested for possible interactions between sample age

and genus (i.e., differences in the slope of regression are dependent on genus). We tested this

model against a model of type "*y ~ covariate + factor"* to assess whether the removal of the

interaction influenced model fit. When the interaction was not significant ($p > 0.05$) we

accepted the simpler model without interaction and concluded that regression slopes did not

differ between genera, though intercepts might [27].

Climate analyses

We obtained climate data from the CHELSA V2.1 climate dataset [44] to investigate the

relationship between specimen preservation conditions, aDNA damage patterns, and climatic

variables. Four primary bioclimatic variables were extracted for each sample location based on

latitude and longitude metadata: (i) annual mean temperature (bio1), (ii) temperature

seasonality (bio4), (iii) annual precipitation (bio12), and (iv) precipitation seasonality (bio15).

Additionally, we extracted monthly temperature (tas_01 - tas_12) and monthly precipitation

(pr_01 - pr_12) data for each sample location to infer climatic conditions at the time of

specimen collection. To quantify the unique and shared contributions of multiple explanatory

variables to the total variance in aDNA damage metrics, we performed a variance partitioning

analysis using the 'varpart' function implemented in the VEGAN package [45]. For each aDNA

metric, we applied a "collection climate" model, where monthly climatic variables

(temperature and precipitation) were assigned to samples based on their location and month of

collection, and an "annual climate" model, where annual temperature and precipitation means

and their seasonality were assigned to samples based on their geographical location.

Additionally, we included in this analysis the genus as a possible confounding effect and

herbarium, which could be interpreted as different storage conditions. Since 'varpart' function

accommodates only up to four variables, for some analysis, temperature and precipitation were

merged into single climatic variable. For each response (endogenous DNA fraction, fragment

size, lambda and 5' C>T damage), we tested statistical significance of individual fractions and

combinations using redundancy analysis with the 'rda' function implemented in VEGAN, with

999 permutation followed by ANOVA.

Results:

A total of 573 herbarium specimens were sequenced, generating libraries sequenced to variable

read counts and depth (supplementary table S1). *Hordeum* samples ($N = 228$) processed at the

Royal Botanic Gardens, Kew (RBGK) generated an average of 12.08 ± 5.94 million reads per

library, with $86.2 \pm 24.3\%$ of reads mapping to the relevant reference genome. *Oryza* samples

($N = 58$) processed at the RBGK yielded an average of 1.41 ± 0.93 million reads per library,

with $86.5 \pm 14.6\%$ of reads mapping to the relevant refence genome. The remaining *Oryza*

samples processed at the University of Nottingham ($N = 287$) generated an average of $0.06 \pm$

0.09 million reads per library, with $78.0 \pm 26.2\%$ of reads mapping to the relevant reference

genome. Damage pattern analysis revealed characteristic aDNA signatures. *Hordeum* samples

exhibited an average 5' C>T frequency of $1.09 \pm 0.44\%$ at the first base of sequenced molecule,

whilst *Oryza* samples exhibited an average of $2.13 \pm 0.70\%$. These damage signatures,

combined with fragment length distributions, were used to authenticate samples and filter out

those inconsistent with genuine aDNA characteristics. Our filtering strategy identified samples

deviating from the assumption of exponential decay of fragment length, as well as samples that

did not display an exponential increase of C>T substitutions at read termini diagnostic of

aDNA. In total, 117 (20%) samples were removed, leaving a final dataset containing 456

samples.

Regression analyses

Endogenous fraction

We analysed the proportion of reads mapping to the reference genome as a proxy for

endogenous DNA content. The regression analyses revealed no statistically significant

relationship between the proportion of endogenous DNA and the sample collection year in

Hordeum ($R^2 = 0.003$, $p = 0.451$, $N = 211$), but a very weak yet significant relationship was

observed in *Oryza* ($R^2 = 0.04$, $p = 0.00167$, $N = 245$; figure 2a). As we aimed at investigating

the effect of genus on the rates of aDNA damage, we also carried the regression for all samples

irrespective of genera, which provided comparable results ($R^2 = 0.012$, $p = 0.0215$, $N = 456$,

supplementary figure S1a).

Fragment length

We measured DNA fragment size using two complementary approaches. First, we validated

our bioinformatic estimates of fragment size distributions using TapeStation profiles of both

gDNA and amplified libraries for a subset of samples ($N = 40$). Peak size of amplified libraries

strongly correlated with the bioinformatically-derived median fragment size of the merged

library reads ($R^2 = 0.614$, $p = 2.22 \times 10^{-9}$, supplementary figure S2a), compared to the weaker

correlation between the TapeStation peaks of libraries and gDNA origin, ($R^2 = 0.129$, $p = 0.029$,

supplementary figure S2b). A significant, albeit weaker correlation was also observed between

the gDNA peak size and the median fragment size of merged reads ($R^2 = 0.287$, $p = 6.35 \times 10^{-4}$, supplementary figure S2c), indicating that merging of overlapping reads of short insert

libraries reflects, at least in part, the original molecule length [29]. The weaker correlation was

expected, as library preparation and sequencing involve processing and purification steps that

can impact the fragment size distribution. Furthermore, we observed a strong and significant

relationship between gDNA peak size and collection year ($R^2 = 0.61$, $p = 3.04 \times 10^{-8}$, $N = 40$,

figure 3), but only after removing the two *Hordeum* outliers HV0061 and HV0081, which were

particularly old (collection years: 1842 and 1867, respectively) but showed gDNA size

distribution patterns inconsistent with their age, possibly due to modern contamination of

exogenous DNA. With the inclusion of these two outliers a significant but weaker relationship

($R^2 = 0.263$, $p = 1.19 \times 10^{-3}$, supplementary figure S3) was observed.

Second, we extended this analysis to the full dataset and examined the relationship between

bioinformatically-derived fragment size and age. We observed a statistically significant

relationship between the log-mean fragment length and the sample collection year for both

genera (figure 2b), with a stronger relationship for *Hordeum* ($R^2 = 0.27$, $p = 5.33 \times 10^{-16}$, $N = $

211) than *Oryza* ($R^2 = 0.112$, $p = 8.58 \times 10^{-8}$, $N = 245$). A statistically significant relationship

was also observed when analysing all samples irrespective of genera ($R^2 = 0.171$, $p = 3.16 \times$

10^{-20}, $N = 456$; supplementary figure S1b).

Damage fraction per site (λ) and DNA decay rate (k)

The slope of log-transformed exponential decline of fragment length frequencies in aDNA (λ)

describes the probability of bond breaking in DNA backbone [41]. We estimated the DNA

decay rate per year (*k*) for *Hordeum* and *Oryza* from the slope of the linear relationship between

λ and sample age (figure 2c). We observed a per nucleotide decay rate of $k = 2.64 \times 10^{-4}$ per

year for *Hordeum* ($R^2 = 0.208$, $p = 3.27 \times 10^{-12}$, $N = 211$), which was 1.5 times faster than the

decay rate of *Oryza* of k= 1.79 x 10^{-4} per year (R^2 = 0.101, p = 3.65 x 10^{-7}, N= 245). The overall

decay rate for all herbarium samples was k= 2.08 x 10^{-4} per year (R^2 = 0.129, p = 2.52 x 10^{-15},

N= 456, supplementary figure S1c), which is slightly faster than the decay rate of k= 1.66 x 10^{-4}

per year observed in *Arabidopsis* and *Solanum* herbarium specimens [27], approximately 2.2

times slower than the $k = 4.6 \times 10^{-4}$ per year observed in dry-pinned arthropod museum

specimens [8], and nearly eight times faster than the rate of k= 2.71 x 10^{-5} observed for ancient

moa bones [12].

Nucleotide misincorporations

Both genera displayed statistically significant increases in the frequencies of 5' C>T

substitutions at first position correlating with the age of the sample (figure 2d), with *Oryza*

starting from a higher baseline of damage when compared to *Hordeum* and displaying a

stronger relationship (R^2 = 0.303, p = 8.62 x 10^{-21}, N= 245 for *Oryza*, and R^2 = 0.207, p =3.63

x 10^{-12}, N= 211 for *Hordeum,* respectively). A slightly weaker yet significant relationship

between 5'C>T substitutions and sample age was also observed when analysing all samples

together (R^2 = 0.106, p = 1.11 x 10^{-12}, N= 456; supplementary figure S1d).

Figure 2: Regression analyses of aDNA damage metrics for *Hordeum* and *Oryza*: (a) Fraction

of endogenous DNA as a function of collection year. (b) Median fragment length of merged

reads as a function of collection year, with log-scaled y-axis to show exponential relationship.

(c) Damage fraction per site (λ) as a function of sample age, with the slope of regression

corresponding to the DNA decay rate per base per year for *Hordeum* ($k = 2.64 \times 10^{-4}$) and

Oryza ($k = 1.79 \times 10^{-4}$). (d) Frequencies of C>T substitutions at first base (5'-end) as a function

of sample age. Insets show regression statistics for each aDNA damage metric for each genus.

Figure 3: Regression between peaks of genomic DNA (gDNA) fragment size distribution

obtained from high resolution fractionation electrophoresis (TapeStation) and collection year

for a subset of samples ($N = 40$; 10 *H. spontaneum*, 10 *H. vulgare*, 10 *O. rufipogon*, and 10 *O.*

latifolia;) after outlier removal. Inset shows regression statistics.

Differences in rates of damage for *Hordeum* and *Oryza*

We compared differences in the regression slopes and intercepts for all aDNA damage metrics

between the genera *Hordeum* and *Oryza* with an analysis of covariance (ANCOVA) and

visualised this comparison with boxplots.

Endogenous fraction

The analysis of covariance revealed significant effects of both sample age (Pr(Sample age) =

0.0114, $N = 456$) and genus (Pr(Genus) = 0.0204, $N = 456$) on the fraction of endogenous

DNA (figure 4a). No significant interaction was observed between sample age and genus

(Pr(Sample age: Genus) = 0.176, $N = 456$), indicating that the rate of exogenous DNA

colonisation over time does not differ significantly between the two genera and is not time

dependent. This was further supported by ANOVA between the model including the interaction

and the model with no interaction, which confirmed that adding the interaction did not improve

model fit ($F = 1.841$, $Pr = 0.175$). The model with no interaction displayed a significant fit ($F =$

5.395, $Pr = 0.004835$) but explained only a small proportion of the variance ($R^2 = 0.02327$),

suggesting that factors beyond genus and sample age can substantially influence endogenous

DNA content.

Fragment length

The analysis of covariance revealed significant effects of sample age and genus on fragment

length (figure 4b) (Pr(Sample age) < 3.45×10^{-21}; Pr(Genus) = 0.0103; N = 456). The

interaction between sample age and genus was not significant (Pr(Sample age: Genus) =

0.0893; N = 456) and its inclusion did not significantly affect model fit (F = 2.8993, Pr =

0.0904). The model with no interaction explained 18% of the variance in fragment length (R^2

= 0.1827) and displayed a significant fit (F = 50.65, Pr < 2.2×10^{-16}), indicating that the genera

differ in their baseline fragment lengths, but not in their rate of fragmentation over time.

Damage fraction per site (λ) and DNA decay rate (k)

We observed a slower DNA decay rate of *Oryza* (k= 1.79×10^{-4} per nucleotide per year) than

that of *Hordeum* (k= 2.64×10^{-4} per nucleotide per year). The analysis of covariance revealed

significant effects of both sample age (Pr(Sample age) = 7.48×10^{-18}; N = 456) and genus

(Pr(Genus) = 5.18×10^{-8}; N = 456) on the rates of bond breaking (figure 4c). However, no

interaction between sample age and genus was observed (Pr(Sample age: Genus) = 0.0884; N

= 456) and including the interaction did not significantly improve model fit (F = 2.9152, Pr =

0.08843). The model with no interaction explained 18% of the variance in lambda values (R^2

= 0.1842) and displayed a highly significant fit (F = 51.14, Pr < 2.2×10^{-16}). Therefore, the

DNA decay rates (k) of *Hordeum* and *Oryza*, which correspond to the slopes of the regression

between damage fraction per site (λ) and age, are not significantly different.

Nucleotide misincorporations

We observed a significant effect of both sample age (Pr(Sample age) = 3.66×10^{-6}; N = 456)

and genus (Pr(Genus) = 3.14×10^{-4}; N = 456) on the rates of 5' C>T substitutions (figure 4d).

The interaction between sample age and genus was significant (Pr(Sample age: Genus) = 2.16

x 10^{-8}; N = 456), indicating that the rate of cytosine deamination over time differs substantially

between the two genera. The model with interaction explained almost 55% of the variance in 5'C>T damage ($R^2 = 0.5448$) and displayed a significant fit ($F = 180$, $Pr < 2.2 \times 10^{-16}$). This was further corroborated by the ANOVA analysis, which confirmed that including the interaction significantly improved model fit ($F = 32.49$, $Pr = 2.165 \times 10^{-8}$). Therefore, not only do *Hordeum* and *Oryza* samples differ in their baseline levels of cytosine deamination, but they also accumulate this type of damage at significantly different rates over time, with *Oryza* showing a steeper increase in deamination with age.

Figure 4: Analysis of Covariance (ANCOVA) of aDNA damage metrics for *Hordeum* and *Oryza*, with sample age as covariate and genus as factor. (a) Endogenous DNA fraction; (b) Fragment length (c) Damage fraction per site (λ); (d) 5' C>T damage frequencies. Significance values reported below each boxplot indicate the effects of age, genus and their interaction upon the analysed aDNA damage metric.

Effects of climatic variables upon rates of aDNA damage

We examined climatic effects using two complementary approaches: a "collection climate" model that assigned monthly temperature and precipitation values based on each sample's collection location and month, and an "annual climate" model that used annual means and seasonality measures based solely on geographical location. Variance partitioning analysis revealed distinct patterns of environmental and temporal control over different aspects of DNA preservation (figure 5). The relative importance of factors varied substantially among damage metrics, with some showing strong environmental sensitivity while others were primarily controlled by age-related processes. In addition to environmental factors, genus emerged as having significant explanatory power, possibly as a variable confounded with climate due to distinct distributions of the *Hordeum* and *Oryza* genera from temperate and tropical climates respectively (figure 1 and 5).

Endogenous fraction

Endogenous DNA content showed little association with the tested variables in the variance partitioning analysis (figure 5a). Only the collection climate model displayed statistical significance, but the fraction of variance explained by all analysed predictors was negligible (Adj. R^2 = 0.024, p = 0.007). The unique contributions of temperature and genus and the combined contributions of temperature, precipitation, age and genus showed marginal statistical significance ($p \leq 0.05$, supplementary table S3a). However, their poor performance in explaining the fraction of variance (Adj. R^2 ranging from 0.001 to 0.01) indicates that endogenous DNA fraction is largely determined by factors not captured by climatic variable nor age. Removing genus from the variance partitioning analysis did not significantly alter the results (supplementary figure S4a, supplementary table S4a). Similarly, inclusion of herbarium as a proxy of institutional preservation practices also did not significantly alter the results (supplementary figure S5a, supplementary table S5a).

Fragment length

Our models had more explanatory power when it came to fragment length, with both collection

and annual climate factors being statistically significant ($p \leq 0.001$) and performing similarly

(figure 5b, Adj. $R^2 = 0.227$ and 0.228, respectively). Sample age was revealed as the largest

contributing factor in both collection and annual models (Adj. $R^2 = 0.199$, 0.193 respectively),

consistent with temporal degradation processes. The shared variance between climatic

variables and age were small, indicating that age effects on fragment size are largely

independent of environmental variables. Indeed, whilst climatic variables were significant in

both models (supplementary table S3b), their unique and combined variance explained were

negligible (Adj. $R^2 \leq 0.01$). Removal of genus from the analysis marginally improved the

unique and combined fraction of variance explained by temperature and precipitation in the

collection and annual model respectively (supplementary figure S4b, supplementary table

S4b), implying collinearity between genus and climatic variables, whereby further inclusion of

herbarium had minimal effects (supplementary figure S5b, supplementary table S5b).

Damage fraction per site (λ)

Models predicting rates of DNA bond breaking (damage fraction per site, λ) from climatic and

temporal variables explained up to 23% of the total variance in λ (figure 5c; Adj. $R^2 = 0.231$,

0.225 in the collection and annual models, respectively). Sample age contributed the largest

unique fraction in both models (Adj. $R^2 =: 0.170$, 0.163), reflecting the fundamental

relationship between specimen age and DNA backbone degradation. Temperature and

precipitation showed minimal unique contributions and minimal shared variance with genus in

the collection climate model (Adj. R^2 ranging from 0.01 to 0.03), and moderate shared variance

with genus in the annual climate model (Adj. R^2 ranging from 0.03 to 0.04), suggesting indirect

effects through correlations between genus and climatic variables. Indeed, removal of genus

from the analysis did not strongly affect the overall variance explained by each model but

inflated the unique and shared variance explained by temperature and precipitation

(supplementary figure S4c).

Nucleotide misincorporations

The 5' C>T damage metric showed the highest explained variance among all damage metrics,

with both collection and annual climate models being significant ($p \leq 0.001$) and explaining

approximately 49% of the total variance (figure 5d). Whilst the unique contribution of age

emerged as the strongest predictor of 5' damage (Adj. R^2 = 0.120, 0.119 in the collection and

annual models, respectively), most explanatory power derived from shared effects among

predictors rather than unique contributions. The combined effects of temperature and genus

explained a substantial proportion of the total variance (Adj. R^2 = 0.135, 0.097). A similar

pattern was observed for the combined effect of precipitation and genus (Adj. R^2 = 0.047,

0.008) and for the combined effects of temperature, precipitation and genus (Adj. R^2 = 0.096,

0.213), suggesting a strong correlation between climatic variables and genus. Removal of genus

from the analysis (supplementary figure S4d) revealed temperature as the largest unique

contributing factor in the collection model (Adj. R^2 = 0.134), followed by sample age (Adj. R^2

= 0.121) and precipitation (Adj. R^2 = 0.047), whilst the combined effect of temperature and

precipitation explained almost 10% of the variance (R^2 = 0.096). Notably, the variance

explained by the combined effect of temperature and precipitation was inflated in the annual

climate model (Adj. R^2 = 0.214, supplementary figure S4d) and emerged as the largest

contributor, followed by unique contribution of sample age (Adj. R^2 = 0.109) and temperature

(Adj. R^2 = 0.096). The inclusion of herbarium did not improve the total fraction of variance

explained in neither model (supplementary figure S5d), with the unique and shared fraction of

explained variance by herbarium and other variables (age, climate and genus) being negligible

(Adj. R^2 ranging from 0.0003 to 0.01).

Temperature effects on 5' C>T substitution frequencies

Given the predictive power of temperature in explaining 5' C>T damage patterns, we conducted

post-hoc regression analyses to examine the relationships between temperature and cytosine

deamination. We observed a strong positive relationship between temperature and 5' C>T

substitutions (figure 6). The annual temperature model explained 18.4% of variance ($R^2 =$

0.184, p = 1.93 x 10^{-18}, N= 456, figure 6a), while the collection temperature model showed a

slightly higher explanatory power (R^2 = 0.201, p = 3.74 x 10^{-20}, N= 456, figure 6b). However,

this relationship was only observed when analysing the genera together and disappeared when

analysing the genera on an individual basis (figure 6c, d).

Figure 5: Climate influences on aDNA damage metrics in herbarium specimens. Maps (top

panels) show the distribution of sampling locations for *Hordeum* and *Oryza* specimens overlaid

on mean annual temperature (left) and annual precipitation (right) from the WorldClim climate

dataset [46]. Venn diagrams (bottom panels) display the unique and shared contributions of the

explanatory variables to the total variance in aDNA damage metrics.: (a) endogenous DNA

fraction, (b) fragment size, (c) damage fraction per site (lambda), and (d) 5' C>T substitution

frequencies at first base. Each metric is analysed using two models: collection climate (left)

and annual climate (right). Adjusted R^2 values for each model are shown above the plots.

Asterisks indicate statistical significance for the overall models and for each unique predictor

and combination of predictors (*$p \leq 0.05$, **$p \leq 0.01$, ***$p \leq 0.001$).

Figure 6: Relationship between temperature and deamination rates (5' C>T substitution

frequencies) in herbarium specimens for (a) annual mean temperature model, (b) collection

temperature model, (c) annual mean temperature model for *Hordeum* and *Oryza*, and (d)

collection temperature model for *Hordeum* and *Oryza*. Insets show regression statistics.

Discussion:

Genomic inferences from preserved samples allows scientists to directly add temporal scales

to the studies of evolutionary and ecological histories of species, and of fundamental questions

about the mechanisms and tempo of evolution. The advent of high-throughput sequencing

technologies fuelled the revolutionary growth of the field of archaeo- and paleo-genomics. In

2022, the number of ancient human genomes passed 10,000 with over 200 papers published

[19]. The discovery of extremally well-preserved samples [47] and development of new

approaches [48–51] have pushed the limits of DNA recovery and sequencing. For plants, owing

to the great diversity of wild and cultivated taxa, it is not possible to know the exact number of

sequenced preserved specimens. In recent publication of 'Plant Tree of Life' alone, genome-

wide enrichment sequences have been generated for over 2,500 plant genera from herbarium

specimens [52]. Current projects focusing on crops and their wild relatives routinely sequence

whole genomes for hundreds of historical specimens [53–55]. RGB Kew is currently aiming

to sequence genomes of 7,000 preserved specimens of fungi from its fungarium collection [56].

With increasing scale of aDNA research across all kingdoms, it is fundamentally important to

improve our understanding of DNA preservation in historical and archaeological samples.

The focus of aDNA preservation research has been the impact of time on DNA degradation.

Thanks to two well-researched examples, we indeed know that DNA deamination and

depurination is correlated with sample age [12,27]. However, it has been noted that

environmental conditions pre- and post-mortem should modulate the pace at which DNA is

degraded [2]. Emphasis has been put on temperature and humidity, two parameters that have

been suggested as reasons for difficulties in sequencing ancient genomes from the tropics [2].

Our work on herbarium specimens has allowed us to quantify the impact of age, environment

and storage conditions on DNA preservation, revealing complex genus-specific differences

extending beyond age-related degradation. Our regression analyses showed relationships

between sample age and different damage metrics, suggesting that while some aspects of aDNA

degradation follow predictable temporal trajectories, others are more heavily influenced by

environmental and biological factors.

Age-dependent aDNA degradation patterns

We further confirmed the highly fragmented nature of aDNA retrieved from herbarium

specimens, with the median fragment size of all analysed samples averaging 46.24 bp (SD =

6.97). This is consistent with previous studies on herbarium samples of a similar age range to

that of the current study [27] and on dry-pinned arthropods museum collections [8], but also

comparable to fragment sizes observed for animal remains that are several orders of magnitude

older, from a few hundred up to thousands of years old [5,12]. Possibly owing to lower levels

of environmental variation experienced by herbarium samples [27], we were able to detect a

weak yet significant relationship between median fragment length and collection year (figure

2b). For a subset of samples ($N = 40$), we directly measured gDNA fragment size distributions

prior to library preparation with high-resolution capillary electrophoresis (TapeStation),

allowing us to assess DNA fragmentation independent of any library construction artifacts or

bias. When examining the relationship between gDNA fragment size (TapeStation) and

collection year, we found an even stronger correlation with age (figure 3). This contrasts with

findings from animal bone studies, where no correlation between DNA fragmentation and

sample age has been observed [5,12,57], even when controlling for environmental variables

[2]. This fundamental difference can be explained by the two-step process of DNA decay and

degradation: a first rapid phase of enzymatic decomposition occurring immediately after host

death, driven by endogenous nucleases and proteases [58], and microbial digestion [59]. A

second phase of chemical decomposition driven by hydrolytic and oxidative reactions,

occurring at much lower rates [6,7]. While the environment during sampling of herbarium

specimens is variable, the standardized preparation and storage procedures used in herbaria

reduce environmental variation during storage compared to the highly variable burial

conditions experienced by animal bones. Indeed, the source herbaria (a proxy for storage

conditions) have very limited explanatory power for DNA degradation (figure 5). The lower

levels of environmental variation experienced by herbarium specimens allows detection of the

underlying temporal fragmentation process that occurs during the second phase of chemical

degradation [27,58]. In archaeological contexts, the influence of variable environmental,

physical and chemical conditions, as well as tissue types and sample excavation and storage

[2,12], may mask more subtle age-related fragmentation patterns that become detectable under

the controlled conditions of herbarium storage.

Differences in physical and chemical tissue characteristics can explain the variation in DNA

decay rates observed across different biological materials. Our herbarium samples exhibited

decay rates ($k = 2.08 \times 10^{-4}$) nearly eight times faster than that observed in bones ($k = 2.71 \times$

10^{-5};[12]), suggesting a higher susceptibility of *post-mortem* enzymatic and chemical DNA

damage in herbarium samples compared to bones [27]. Interestingly, arthropod museum

specimens show even faster decay ($k = 4.6 \times 10^{-4}$,[8]), approximately twice the rate observed

in our herbarium samples. This variation likely reflects differences in tissue composition

(cellulose/lignin in plants, chitin/protein in arthropods and hydroxyapatite/collagen in bones),

storage methods (pressed herbarium sheets, pinned insects, and buried remains), and the

protective properties of these different methods. The intermediate position of herbarium

specimens between museum arthropods collections and ancient bones in terms of decay rate

suggests that DNA fragmentation is influenced by both intrinsic tissue properties and storage

conditions.

Furthermore, the 5' C>T damage patterns also showed significant correlations with age in both

genera (figure 2d). However, while both genera show similar rates of DNA fragmentation, their

divergent responses to chemical modifications suggest different susceptibilities to oxidative

damage and deamination. The higher baseline damage and steeper accumulation rates in *Oryza*

species indicate that the genus is more susceptible to *post-mortem* deamination, possibly due

them growing in different environments.

While aDNA damage metrics such as fragmentation and deamination were observed to be time-

dependent processes, the fraction of endogenous DNA content showed no correlation with age

(figure 2a), suggesting that microbial colonization and subsequent displacement of endogenous

DNA is largely independent of specimen age. Biological degradation processes are thus

primarily driven by factors other than time, such as initial microbial load, tissue characteristics,

and individual storage conditions. However, whilst we used species-specific reference

genomes, increasing evolutionary distance between query and reference might have impacted

this metric [8]. We therefore suggest caution when interpreting such results.

Environmental controls on aDNA preservation

Our variance partitioning analysis shows that climatic effects on aDNA damage are largely

mediated through complex interactions among predictors rather than independent

contributions. The dominance of shared variance fractions indicates that temperature,

precipitation and genus effects are highly correlated in our dataset, reflecting the geographic

and temporal sampling patterns of herbarium collections.

For metrics reflecting DNA backbone integrity (fragment size and lambda), age consistently

contributed the largest unique variance fractions, confirming that temporal degradation

processes are the primary drivers of DNA fragmentation. The relatively large unique age

contributions suggest that these physical degradation processes proceed independently of

environmental and taxonomic factors once specimens enter standardized herbarium

preparation methods storage practices. This age-dependent fragmentation is a distinctive

feature of herbarium specimens that contrasts with the patterns observed in both archaeological

animal bones [5,12] and arthropod museum collections [8], where environmental variation

during deposition or storage can mask temporal patterns.

The strong correlation between temperature and 5' C>T damage provides mechanistic insight

into why *Oryza* specimens, predominantly from tropical and sub-tropical regions, consistently

show higher damage levels compared to temperate *Hordeum* specimens (figure 6). Elevated

temperatures accelerate the spontaneous deamination of cytosine residues, explaining both the

higher baseline damage observed in *Oryza* and the overall temperature-damage relationship

observed across all herbarium samples. Precipitation variables as a proxy of humidity also

showed significant effects, indicating that deamination is a time-dependent process modulated

by temperature and humidity [2]. Overall, the complex interplay between temperature,

humidity and their seasonality and damage metrics suggests that specimens from regions with

less seasonal climates may experience different preservation trajectories than those from highly

seasonal climates. The substantial unique genus contributions to 5' C>T damage confirm

differential susceptibility to cytosine deamination between *Hordeum* and *Oryza*. However, the

large shared variance fractions involving genus indicate that taxonomic differences are partially

confounded with climatic variables, reflecting the contrasting geographic origins of these

genera (temperate vs. tropical; figure 5). It is possible that genus captures other environmental

factors that were not included in temperature and precipitation. Precipitation is a decent proxy

for humidity, but there are multiple other factors influencing it, and high humidity might be

accelerating the deamination process. Alternatively, the differences observed due to tropical

and temperate samples (captured in genus) could be explained by different sample processing

approaches in the two areas, with tropical specimens being more often oven-dried/baked.

Indeed, baking has been shown to substantially affect DNA degradation [60]. Additionally, in

the tropics, alcohol treatment used to be common to prevent moulding, and this has been shown

to limit the success in DNA amplification, presumably due to rapid degradation [61].

The negligible variance explained for endogenous fraction across all models supports our

hypothesis that microbial colonization is largely independent of the measured environmental

and biological predictors. This finding suggests that endogenous DNA loss is driven by

stochastic factors such as initial contamination loads, handling procedures, or unmeasured

specimen-specific characteristics.

Conclusions and implications for herbarium curation and ancient DNA research

We show that ancient DNA damage patterns result from complex interactions between temporal

degradation processes, environmental conditions during sampling, biological properties and

storage condition. Herbarium specimens exhibit age-dependent DNA fragmentation patterns

that are not observed in animal bone studies, indicating that standardized preservation

conditions can reveal underlying temporal degradation processes masked by environmental

variation in archaeological contexts. We identified significant genus-specific differences in

aDNA damage susceptibility and established temperature as the dominant environmental driver

of cytosine deamination.

Environmental effects on DNA damage operate primarily through complex interactions with

taxonomic and temporal factors rather than direct independent contributions. Age-related

degradation dominates physical DNA breakdown, while genus-specific differences in cytosine

deamination susceptibility reflect both environmental adaptation and intrinsic biochemical

properties. The comparison across biological materials reveals that tissue composition

fundamentally constrains preservation potential, while storage conditions modulate decay

rates.

These findings highlight the importance of considering specimen age, climatic origin and

storage conditions when selecting herbarium specimens for ancient DNA analyses. The

mechanisms underlying genus-specific differences in aDNA damage accumulation warrant

detailed biochemical investigation. The significant, although small effect of herbarium-specific

factors on damage metrics indicates that institutional preservation practices vary and represent

an underexplored factor in ancient DNA research. Future studies should provide insights into

the biological processes driving DNA degradation, including detailed microbiome analyses.

Understanding which microbial taxa are most problematic for DNA preservation and how their

colonization is influenced by specimen characteristics could inform targeted preservation

strategies. As the scale of ancient DNA research continues to expand, such findings will be

essential for maximizing the scientific value of the world's natural collections and informing

evidence-based approaches to specimen sampling and preservation.

Availability of code and requirements:

Project name: Herbaria aDNA Damage

Project home page: https://github.com/Stefano-Porrelli/Herbaria_aDNA_Damage

Operating system(s): Linux/Unix (tested on SLURM-based HPC systems)

Programming language: Bash, R (version $\geq$4.0)

Other requirements:

•   Conda/Miniconda,

•   AdapterRemoval (≥2.0), BWA (≥0.7.17), FastQC (≥0.11), SAMtools (≥1.10), DeDup

(≥0.12), mapDamage2 (≥2.0), MultiQC (≥1.8), Preseq (≥2.0), AMBER.

•   R packages: dplyr, tidyr, purrr, stringr, readr, MASS, vegan, car, ggplot2, colorspace,

viridis, ggrepel, ggtext, ggpubr, gridExtra, cowplot, sf, geodata, terra, maps

•   External data: CHELSA V2.1 climate data (https://chelsa-climate.org/)

License: MIT licence

Any restrictions to use by non-academics: None

Data availability:

Raw FASTQ DNA sequences of all samples processed at the Royal Botanic Gardens, Kew

(UK) are deposited on the sequence reads archive (SRA) on NCBI: *Hordeum vulgare*

sequences are deposited under BioProject PRJNA1288534. *Hordeum spontaneum* sequences

are deposited under BioProject PRJNA1289164. *Oryza rufipogon* sequences are deposited

under BioProject PRJNA1288425. *Oryza grandiglumis* sequences are deposited under

BioProject PRJNA1288424. *Oryza latifolia* sequences are deposited under BioProject

PRJNA1288423. All remaining raw FASTQ DNA sequences of *Oryza* are deposited on the

SRA under BioProject PRJNA1302186. Scripts and dataset to reproduce the analyses are

available at https://github.com/Stefano-Porrelli/Herbaria_aDNA_Damage (EMBARGOED).

Supplementary material:

Supplementary material is available at XXXXXXX

Authors' contributions:

S.P. and R.M.G. conceived the project and designed the study; R.M.G. and S.P. designed the

sampling strategy, with contributions from A.F. and P.H.L.; P.J.K and N.S. overseen and

supervised the generation of sequencing data for *Hordeum* samples; S.P. sampled herbaria and

performed the aDNA laboratory work and bioinformatic screening for the *Hordeum* samples

processed at RBGK; A.H. sequenced the *Hordeum* libraries at IPK; P.H.L. sampled herbarium

samples and performed the aDNA laboratory work and bioinformatic screening for the *Oryza*

samples processed at RBGK; A.F. selected herbarium specimens, A.F., M.N.R., and R.A.W.

performed the sampling, A.F., W.Y., N.M., M.N.R., and A.C.C. isolated aDNA and prepared

aDNA libraries, and A.F. performed screening and validation for the *Oryza* samples processed

at KAUST/UoN; S.P. analysed the historical data and interpreted the results with the

contribution of R.M.G. and P.H.L; S.P. and R.M.G. wrote the manuscript with contributions

from all authors.

Acknowledgements:

We thank curators for providing herbarium specimens: Anna Haigh and Sue Zmartzy (Royal

Botanic Gardens, Kew); Paul Peterson and Robert Soreng (Smithsonian Institute Herbarium);

Jordan K. Teisher (Missouri Botanic Gardens); Miriam Gaudeul (National History Museum,

France); Michaela Schmull and Anthony R. Brach (Harvard Herbarium); Matthew C. Pace

(New York Botanic Gardens). We acknowledge the expert technical assistance of Ines Walde

and Jacqueline Pohl during DNA sequencing of *Hordeum* samples at the Leibniz Institute of

Plant Genetics and Crop Plant Research (IPK). We also thank Anne Fiebig for her expert

technical assistance with data submission of *Hordeum* low-throughput SRA sequences at IPK.

Funding:

This work was supported by the European Union Horizon 2020 research and innovation

programme under grant agreement No. 862613 (AGENT - Activated GEnebank NeTwork),

which funded the processing and low-coverage sequencing of *Hordeum* samples at the Royal

Botanic Gardens, Kew and the Leibniz Institute of Plant Genetics and Crop Plant Research (IPK). UK Research and Innovation grant EP/X022404/1 funded the processing and low-coverage sequencing of *Oryza* samples at the Royal Botanic Gardens, Kew, while King Abdullah University of Science and Technology (KAUST) grant ORA-CRG10-2021-4734 to R.A.W. supported the processing and low-coverage sequencing of *Oryza* samples at the University of Nottingham.

Ethics and permissions:

Historical herbarium specimens were obtained from established herbaria following institutional specimen access protocols. All sampling was conducted under standard museum loan agreements and followed institutional guidelines for destructive sampling of herbarium material. No additional ethical permissions were required as the study involved only preserved specimens collected under historical botanical collecting practices prior to CBD, but authors adhere to the ethical principles for Access and Benefit Sharing.

Conflict of interest:

The authors declare no conflicts of interest.

References:

1. Kristiansen K. Towards a new paradigm? The third science revolution and its possible consequences in archaeology. *Curr Swed Archaeol*. Svenska Arkeologiska Samfundet; 2021; doi: 10.37718/csa.2014.01.

2. Kistler L, Ware R, Smith O, Collins M, Allaby RG. A new model for ancient DNA decay based on paleogenomic meta-analysis. *Nucleic Acids Res*. 2017; doi: 10.1093/nar/gkx361.

3. Orlando L, Allaby R, Skoglund P, Der Sarkissian C, Stockhammer PW, Ávila-Arcos MC, et

al.. Ancient DNA analysis. *Nat Rev Methods Primers*. Springer Science and Business Media

LLC; 2021; doi: 10.1038/s43586-020-00011-0.

4. Dabney J, Meyer M, Pääbo S. Ancient DNA damage. *Cold Spring Harb Perspect Biol*. Cold

Spring Harbor Laboratory; 2013; doi: 10.1101/cshperspect.a012567.

5. Sawyer S, Krause J, Guschanski K, Savolainen V, Pääbo S. Temporal patterns of nucleotide

misincorporations and DNA fragmentation in ancient DNA. *PLoS One*. Public Library of

Science (PLoS); 2012; doi: 10.1371/journal.pone.0034131.

6. Lindahl T, Nyberg B. Rate of depurination of native deoxyribonucleic acid. *Biochemistry*.

American Chemical Society (ACS); 1972; doi: 10.1021/bi00769a018.

7. Lindahl T. Instability and decay of the primary structure of DNA. *Nature*. Springer Science

and Business Media LLC; 1993; doi: 10.1038/362709a0.

8. Mullin VE, Stephen W, Arce AN, Nash W, Raine C, Notton DG, et al.. First large-scale

quantification study of DNA preservation in insects from natural history collections using

genome-wide sequencing. *Methods Ecol Evol*. Wiley; 2023; doi: 10.1111/2041-210x.13945.

9. Lan T, Lindqvist C. Technical advances and challenges in genome-scale analysis of ancient

DNA. *Population Genomics*. Cham: Springer International Publishing;

10. Schubert M, Ginolhac A, Lindgreen S, Thompson JF, Al-Rasheid KAS, Willerslev E, et al..

Improving ancient DNA read mapping against modern reference genomes. *BMC Genomics*.

Springer Science and Business Media LLC; 2012; doi: 10.1186/1471-2164-13-178.

11. Dolenz S, van der Valk T, Jin C, Oppenheimer J, Sharif MB, Orlando L, et al.. Unravelling

reference bias in ancient DNA datasets. *Bioinformatics*. Oxford University Press (OUP);

2024; doi: 10.1093/bioinformatics/btae436.

12. Allentoft ME, Collins M, Harker D, Haile J, Oskam CL, Hale ML, et al.. The half-life of

DNA in bone: measuring decay kinetics in 158 dated fossils. *Proc Biol Sci*. The Royal

Society; 2012; doi: 10.1098/rspb.2012.1745.

13. Hofreiter M, Paijmans JLA, Goodchild H, Speller CF, Barlow A, Fortes GG, et al.. The

future of ancient DNA: Technical advances and conceptual shifts. *Bioessays*. Wiley; 2015;

doi: 10.1002/bies.201400160.

14. Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of next generation

sequencing data. *BMC Bioinformatics*. Springer Nature; 2014; doi: 10.1186/s12859-014-

0356-4.

15. Schubert M, Ermini L, Der Sarkissian C, Jónsson H, Ginolhac A, Schaefer R, et al..

Characterization of ancient and modern genomes by SNP detection and phylogenomic and

metagenomic analysis using PALEOMIX. *Nat Protoc*. Springer Science and Business

Media LLC; 2014; doi: 10.1038/nprot.2014.063.

16. Peltzer A, Jäger G, Herbig A, Seitz A, Kniep C, Krause J, et al.. EAGER: efficient ancient

genome reconstruction. *Genome Biol*. Springer Science and Business Media LLC; 2016;

doi: 10.1186/s13059-016-0918-z.

17. Willerslev E, Cooper A. Ancient DNA. *Proc Biol Sci*. The Royal Society; 2005; doi:

10.1098/rspb.2004.2813.

18. Liu Y, Bennett EA, Fu Q. Evolving ancient DNA techniques and the future of human

history. *Cell*. Elsevier BV; 2022; doi: 10.1016/j.cell.2022.06.009.

19. Mallick S, Micco A, Mah M, Ringbauer H, Lazaridis I, Olalde I, et al.. The Allen Ancient

DNA Resource (AADR) a curated compendium of ancient human genomes. *Sci Data*.

Springer Science and Business Media LLC; 2024; doi: 10.1038/s41597-024-03031-7.

20. Frantz LAF, Bradley DG, Larson G, Orlando L. Animal domestication in the era of ancient

genomics. *Nat Rev Genet*. Springer Science and Business Media LLC; 2020; doi:

10.1038/s41576-020-0225-0.

21. Peris D, Janssen K, Barthel HJ, Bierbaum G, Delclòs X, Peñalver E, et al.. DNA from resin-

embedded organisms: Past, present and future. *PLoS One*. Public Library of Science (PLoS);

2020; doi: 10.1371/journal.pone.0239521.

22. Kistler L, Bieker VC, Martin MD, Pedersen MW, Ramos Madrigal J, Wales N. Ancient

plant genomics in archaeology, herbaria, and the environment. *Annu Rev Plant Biol*. Annual

Reviews; 2020; doi: 10.1146/annurev-arplant-081519-035837.

23. Estrada O, Breen J, Richards SM, Cooper A. Ancient plant DNA in the genomic era. *Nat

Plants*. Springer Science and Business Media LLC; 2018; doi: 10.1038/s41477-018-0187-

9.

24. : Index herbariorum - the William & Lynda Steere herbarium. The William & Lynda Steere

Herbarium. http://sweetgum.nybg.org/ih/ (2018). Accessed 2025 Oct 23.

25. Burbano HA, Gutaker RM. Ancient DNA genomics and the renaissance of herbaria.

Science. 2023; doi: 10.1126/science.adi1180.

26. Eckert L, Eckert I, Rahn O, So CP, Barrett RDH. Using herbarium collections to study

genetic responses to global change. *New Phytol*. 2025; doi: 10.1111/nph.70454.

27. Weiß CL, Schuenemann VJ, Devos J, Shirsekar G, Reiter E, Gould BA, et al.. Temporal

patterns of damage and decay kinetics of DNA retrieved from plant herbarium specimens.

R Soc Open Sci. The Royal Society; 2016; doi: 10.1098/rsos.160239.

28. Latorre SM, Lang PLM, Burbano HA, Gutaker RM. Isolation, library preparation, and

bioinformatic analysis of historical and ancient plant DNA. *Curr Protoc Plant Biol*. Wiley;

2020; doi: 10.1002/cppb.20121.

29. Kircher M, Sawyer S, Meyer M. Double indexing overcomes inaccuracies in multiplex

sequencing on the Illumina platform. *Nucleic Acids Res*. Oxford University Press (OUP);

2012; doi: 10.1093/nar/gkr771.

30. Meyer M, Kircher M. Illumina sequencing library preparation for highly multiplexed target

capture and sequencing. *Cold Spring Harb Protoc*. Cold Spring Harbor Laboratory; 2010;

doi: 10.1101/pdb.prot5448.

31. Hofreiter M, Jaenicke V, Serre D, Haeseler A, Pääbo S. DNA sequences from multiple

amplifications reveal artifacts induced by cytosine deamination in ancient DNA. *Nucleic*

Acids Res. 2001; doi: 10.1093/NAR/29.23.4793.

32. Gutaker RM, Reiter E, Furtwängler A, Schuenemann VJ, Burbano HA. Extraction of

ultrashort DNA molecules from herbarium specimens. *Biotechniques*. 2017; doi:

10.2144/000114517.

33. Briggs AW, Stenzel U, Meyer M, Krause J, Kircher M, Pääbo S. Removal of deaminated

cytosines and detection of in vivo methylation in ancient DNA. *Nucleic Acids Res*. Oxford

University Press (OUP); 2010; doi: 10.1093/nar/gkp1163.

34. Schubert M, Lindgreen S, Orlando L. AdapterRemoval v2: rapid adapter trimming,

identification, and read merging. *BMC Res Notes*. Springer Nature; 2016; doi:

10.1186/s13104-016-1900-2.

35. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv [q-bio.GN].

36. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al.. Twelve years of

SAMtools and BCFtools. *Gigascience*. Oxford University Press (OUP); 2021; doi:

10.1093/gigascience/giab008.

37. Jónsson H, Ginolhac A, Schubert M, Johnson PLF, Orlando L. mapDamage2.0: fast

approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics*.

Oxford University Press (OUP); 2013; doi: 10.1093/bioinformatics/btt193.

38. Fornasiero A, Feng T, Al-Bader N, Alsantely A, Mussurova S, Hoang NV, et al.. Oryza

genome evolution through a tetraploid lens. *Nat Genet*. Springer Science and Business

Media LLC; 2025; doi: 10.1038/s41588-025-02183-5.

39. . R: language environment statistical computing. *R Foundation Statistical Computing*.

Vienna, Austria;

40. Venables WN, Ripley BD, editors. Modern applied statistics with S-plus. New York, NY:

Springer;

41. Deagle BE, Eveson JP, Jarman SN. Quantification of damage in DNA recovered from

highly degraded samples--a case study on DNA in faeces. *Front Zool*. Springer Science and

Business Media LLC; 2006; doi: 10.1186/1742-9994-3-11.

42. Gutaker RM, Burbano HA. Reinforcing plant evolutionary genomics using ancient DNA.

Curr Opin Plant Biol. Elsevier BV; 2017; doi: 10.1016/j.pbi.2017.01.002.

43. Weiß CL, Dannemann M, Prüfer K, Burbano HA. Contesting the presence of wheat in the

British Isles 8,000 years ago by assessing ancient DNA authenticity from low-coverage data.

Elife. eLife Sciences Publications, Ltd; 2015; doi: 10.7554/eLife.10005.

44. Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, et al.. Climatologies

at high resolution for the earth's land surface areas. *Sci Data*. Springer Science and Business

Media LLC; 2017; doi: 10.1038/sdata.2017.122.

45. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al.. Vegan:

community ecology package. *R package version 25-4*. 2019;

46. Fick SE, Hijmans RJ. WorldClim 2: new 1-km spatial resolution climate surfaces for global

land areas. *Int J Climatol*. Wiley; 2017; doi: 10.1002/joc.5086.

47. van der Valk T, Pečnerová P, Díez-Del-Molino D, Bergström A, Oppenheimer J, Hartmann
S, et al.. Million-year-old DNA sheds light on the genomic history of mammoths. *Nature*.
Springer Science and Business Media LLC; 2021; doi: 10.1038/s41586-021-03224-9.

48. Dabney J, Knapp M, Glocke I, Gansauge M-T, Weihmann A, Nickel B, et al.. Complete
mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from
ultrashort DNA fragments. *Proc Natl Acad Sci U S A*. Proceedings of the National Academy
of Sciences; 2013; doi: 10.1073/pnas.1314445110.

49. Dabney J, Meyer M. Extraction of highly degraded DNA from ancient bones and teeth.
Methods in Molecular Biology. New York, NY: Springer New York;

50. Gansauge M-T, Meyer M. Single-stranded DNA library preparation for the sequencing of
ancient or damaged DNA. *Nat Protoc*. Springer Science and Business Media LLC; 2013;
doi: 10.1038/nprot.2013.038.

51. Slon V, Hopfe C, Weiß CL, Mafessoni F, de la Rasilla M, Lalueza-Fox C, et al.. Neandertal
and Denisovan DNA from Pleistocene sediments. *Science*. American Association for the
Advancement of Science (AAAS); 2017; doi: 10.1126/science.aam9695.

52. Zuntini AR, Carruthers T, Maurin O, Bailey PC, Leempoel K, Brewer GE, et al..
Phylogenomics and the rise of the angiosperms. *Nature*. Springer Science and Business
Media LLC; 2024; doi: 10.1038/s41586-024-07324-0.

53. Kistler L, de Oliveira Freitas F, Gutaker RM, Maezumi SY, Ramos-Madrigal J, Simon MF,
et al.. Historic manioc genomes illuminate maintenance of diversity under long-lived clonal
cultivation. *Science*. American Association for the Advancement of Science (AAAS); 2025;
doi: 10.1126/science.adq0018.

54. Alsantely A, Gutaker R, Navarrete Rodríguez ME, Arrieta-Espinoza G, Fuchs EJ, Costa de
Oliveira A, et al.. The International Oryza Map Alignment Project (IOMAP): the Americas-

past achievements and future directions. *J Exp Bot*. Oxford University Press (OUP); 2023; doi: 10.1093/jxb/erac490.

55. Kreiner JM, Latorre SM, Burbano HA, Stinchcombe JR, Otto SP, Weigel D, et al.. Rapid weed adaptation and range expansion in response to agriculture over the past two centuries. *Science*. American Association for the Advancement of Science (AAAS); 2022; doi: 10.1126/science.abo7293.

56. Varga T, Woods R, Pitsillides F, Hill R, Biketova AY, Llewellyn T, et al.. Whole genome sequencing of historical specimens from the world's largest fungal collection yields high-quality assemblies. *New Phytol*. Wiley; 2025; doi: 10.1111/nph.70472.

57. Pääbo S. Ancient DNA: extraction, characterization, molecular cloning, and enzymatic amplification. *Proc Natl Acad Sci U S A*. Proceedings of the National Academy of Sciences; 1989; doi: 10.1073/pnas.86.6.1939.

58. Molak M, Ho SYW. Evaluating the impact of post-mortem damage in ancient DNA: a theoretical approach. *J Mol Evol*. Springer Science and Business Media LLC; 2011; doi: 10.1007/s00239-011-9474-z.

59. Eglinton G, Logan GA. Molecular preservation. *Philos Trans R Soc Lond B Biol Sci*. The Royal Society; 1991; doi: 10.1098/rstb.1991.0081.

60. Staats M, Cuenca A, Richardson JE, Vrielink-van Ginkel R, Petersen G, Seberg O, et al.. DNA damage in plant herbarium tissue. *PLoS One*. Public Library of Science (PLoS); 2011; doi: 10.1371/journal.pone.0028448.

61. Särkinen T, Staats M, Richardson JE, Cowan RS, Bakker FT. How to open the treasure chest? Optimising DNA extraction from herbarium specimens. *PLoS One*. Public Library of Science (PLoS); 2012; doi: 10.1371/journal.pone.0043808.