Deep-learning deconvolution and segmentation of fluorescent membranes for high-precision bacterial cell-size profiling

Octavio Reyes-Matte[1,*], Carsten Fortmann-Grote[1], Beate Gericke[1], Nadja Hüttmann[1], Nikola Ojkic[2], Javier Lopez-Garrido[1,*]

[1] Max Planck Institute for Evolutionary Biology, Plön, Germany

[2] School of Biological and Behavioral Sciences, Queen Mary University of London, UK

Correspondence: lopezgarrido@evolbio.mpg.de; reyesmatte@evolbio.mpg.de

Keywords: image segmentation, image deconvolution, bacterial cell size, PBP1, *Priestia megaterium*, *Bacillus*

ABSTRACT

Evolutionary studies in bacteria have emphasized genetic and metabolic diversity, while cell-size
variation has received less attention. Here we introduce MEDUSSA, a high-throughput method
for precise bacterial cell-size profiling based on automatic segmentation of fluorescent membrane
images, well suited to studying cell-size diversity. The approach uses deep-learning-based mem-
brane deconvolution, segmentation models fine-tuned for fluorescent-membrane images, and er-
ror-corrected cell measurement to extract accurate sizes from individual bacterial cells regardless
of shape, size, chaining, or clustering. Our method overcomes limitations of phase-contrast seg-
mentation, yielding reliable single-cell dimensions. We applied MEDUSSA to six strains of *Priestia*
megaterium and found over twofold differences in cell volume across strains, largely driven by dif-
ferences in cell width. We identified a partially-functional PBP1 allele that underlies the reduced
width of one strain. Together, these results demonstrate the power of comparative analyses in
bacterial cell biology and expand the toolkit to investigate the evolution of bacterial cell size.

INTRODUCTION

Bacterial cells are often perceived as uniformly small, which has led to a relative lack of attention to the evolution of bacterial cell size. Yet, careful microscopic observation has revealed an incredible diversity of bacterial shapes and sizes, with cell volumes spanning multiple orders of magnitude (Schavemaker & Munoz-Gomez, 2022; Schober et al., 2025). Even among closely related species, cell size can vary significantly. For instance, strains from two *Bacillus* species, *B. thuringiensis* and *B. pumilus*, can differ in volume by nearly an order of magnitude (Lopez-Garrido et al., 2018), suggesting that cell size is an evolvable trait capable of changing over relatively short evolutionary timescales.

Despite this diversity, our appreciation of bacterial cell size variation at finer evolutionary scales remains limited. Systematic studies have focused on mutant libraries in a few model strains (Juillot et al., 2021; Ursell et al., 2017). However, comparative studies analyzing the size of different bacterial strains or species are rare (Delaby et al., 2025; Gray et al., 2019), and most available data come from independent studies, often using different imaging methods and manual measurements of a few representative cells cultured in different conditions. While this highlights bacterial size diversity, methodological inconsistencies make cross-study comparisons difficult. As a result, we lack a clear picture of how bacterial cell size vary among closely related species and strains.

Recent advances in image analysis pose a unique opportunity for the standardization of methods to measure bacterial cell size and may help to address these challenges. In particular, the incorporation of machine-learning into image analysis pipelines have enabled the automatic identification and precise delineation of boundaries of individual bacterial cells in microscopy images. This process, known as image segmentation, generates masks of individual cells that can be used for accurate cell size measurements. The U-Net, a convolutional neural network architecture originally designed for the analysis of biomedical images (Falk et al., 2019; Ronneberger et al., 2015), is behind some of the most powerful image segmentation algorithms currently available, such as StarDist (Schmidt et al., 2018), DeLTA (Lugagne et al., 2020), Cellpose (Stringer et al., 2021), MiSiC (Panigrahi et al., 2021), Omnipose (Cutler et al., 2022), DeepBacs (Spahn et al., 2022) and DistNet (Ollion et al., 2024), to name a few. More recently, vision transformers (Dosovitskiy et al., 2020) have enabled the development of foundation models (Kirillov et al., 2023; Zou et al., 2023), which support strong off-the-shelf performance and facilitate fine-tuning on custom data.

60 Several models have leveraged large public microscopy datasets to achieve high flexibility, includ-
61 ing microSAM (Archit et al., 2025), Cellpose-SAM (Pachitariu et al., 2025) and CellSAM (Marks
62 et al., 2025). These tools are significantly improving the precision and automation of bacterial cell
63 segmentation, paving the way for more consistent and reproducible cell size measurements.
64
Here, we have fine-tuned and benchmarked some these models for segmenting fluorescent mem-
branes. Building on this, we developed a pipeline for extracting precise single-cell size information
from fluorescent membrane images, which allows the characterization of individual cell sizes re-
gardless of clustering or chaining patterns. We applied this pipeline, that we call MEDUSSA
(MEmbrane DeconvolUtion and Segmentation for Size Analyses), to investigate cell-size diversity
across six strains of *Priestia megaterium*, a Gram-positive bacterium commonly used for physiological
studies and biotechnological applications (Logan & Vos, 2015). We found significant differences
in volume between the strains—spanning more than twofold—underpinned by consistent differ-
ences in cell length and width. We further link width differences between two of the strains to a
specific mutation in *ponA*, which encodes penicillin binding protein 1 (PBP1), resulting in a hypo-
morphic PBP1 allele.

RESULTS

Limitations of phase contrast segmentation for single cell studies.

Deep-learning-based segmentation algorithms are versatile and can segment different types of im-

ages, as long as they are trained with appropriately labeled datasets for the specific task. However,

phase-contrast has become the default imaging modality for bacterial segmentation, as it provides

excellent contrast between cells and background without the need for staining or genetic manipu-

lation (Cutler et al., 2022; Ducret et al., 2016; O'Connor et al., 2022; Ollion et al., 2024; Paintdakhi

et al., 2016; Stylianidou et al., 2016; Ursell et al., 2017; Van Valen et al., 2016). Despite its ad-

vantages, phase contrast images have limitations for cell size estimation. Firstly, they lack clear

visual references to precisely define cell boundaries. Secondly, phase contrast does not allow the

distinction of cells that remain attached after division, complicating the identification of individual

cells in bacterial species that form cellular chains (Holt et al., 1994; Syvertsson et al., 2016). For

instance, *Bacillus subtilis* and related species form heterogeneous populations in which single cells

coexist with long chains consisting of dozens of cells (Kearns & Losick, 2005; Margulis et al.,

1998). Because the separation of individual cells within a chain is not visible in phase-contrast

images, state-of-the-art segmentation models optimized for phase-contrast segmentation, such as

the Omnipose "bact_phase_omni" model (Cutler et al., 2022), segment chains as very long indi-

vidual cells, making it difficult to extract information at the single cell level (Fig. 1a,c).

This issue can be circumvented by staining the cell membranes with lipophilic fluorescent dyes

such as FM 4-64 (Pogliano et al., 1999) or Nile Red (Strahl et al., 2014), which allows the visuali-

zation of the septal membranes that separate individual cells within chains (Fig. 1b,d). The fluo-

rescence intensity profile of FM 4-64-stained membranes shows a sharp peak that can be traced

around the cell (Fig 1e,f), providing an unambiguous guide to delimit the cell boundaries. We

therefore explored the performance of different segmentation models on fluorescent membrane

images.

104
Figure 1. Phase-contrast versus fluorescent-membrane segmentation. a, b, Phase-contrast (**a**) and fluorescent
membrane images (**b**) of *B. subtilis* growing vegetatively. Membranes were stained with FM 4-64. The insets show
zoomed-in views of the boxed regions; the red lines illustrate the paths along which intensity profiles are calculated
in panels **e** and **f**. **c, d,** Segmentation masks for the phase-contrast (**c**) and fluorescent membrane (**d**) images shown
in **a** and **b**. The phase-contrast image was automatically segmented with Omnipose (Cutler et al., 2022) and the fluo-
rescent-membrane image was manually segmented with JFilament (Smith et al., 2010) (see also panel **g**). Every mask
has a different color. **e, f,** Phase-contrast (**e**) and fluorescent membrane (**f**) intensity profiles of 39 cells, measured
along a 24-pixel line perpendicular to their long axis, centered at midcell (illustrated in insets in panels **a** and **b**). Pixel
intensities were normalized to the maximum intensity observed across the lines for each imaging modality. **g,** Example
of manual fluorescent-membrane segmentation using the active contour software JFilament (Smith et al., 2010).
115

High-throughput segmentation of fluorescent membranes.

We fine-tuned existing deep-learning segmentation models for segmenting fluorescent membrane images of bacterial cells, and benchmarked their performance. We focused on Cellpose3 (Stringer & Pachitariu, 2025), Omnipose (Cutler et al., 2022), and the SAM-based foundation models microSAM (Archit et al., 2025) and Cellpose-SAM (Pachitariu et al., 2025). Omnipose has been reported to perform better on elongated cells than other segmentation algorithms (Cutler et al., 2022). The models were fine-tuned using over seven-thousand FM 4-64-stained *B. subtilis* cells annotated with JFilament active-contour software (Fig. 1g) (Smith et al., 2010), which allows the semiautomated generation of snakes that track the fluorescent membrane peak around the cell, thereby diminishing user annotation bias. JFilament snakes precisely delimit cell contours and provide accurate cell dimensions (Supplementary Fig. 1). The training dataset included single cells, cells within chains and cells within clusters (see Methods for details). We tested the fine-tuned models with images of three bacilli of different sizes: *B. subtilis*, *B. thuringiensis*, and *Priestia megaterium* (Fig. 2a; Supplementary Fig. 2a). We also tested Morphometrics, a non-deep-learning segmentation method that has shown good performance on fluorescent membrane images (Ursell et al., 2017). Cellpose-SAM showed the best performance in benchmarking tests (mean F1 at IoU 0.8, 0.86±0.08), followed by Omnipose (0.76±0.16), Morphometrics (0.64±0.22), Cellpose3 (0.59±0.17) and microSAM (0.30±0.16) (Fig. 2a). Because segmentation algorithms are known to often fail on elongated cells, we further tested the performance of the fine-tuned models on a *Lysinibacillus* strains that naturally produces long cells under laboratory conditions (Supplementary Fig. 2b). Only Omnipose segmented long cells appropriately. The other models oversegmented them. Since our goal was to develop a pipeline for cell size analysis across different strains and species, many of which may naturally form long cells, we further analyzed Omnipose segmentation to improve its performance.

The fine-tuned Omnipose model produced segmentation masks lacking aberrations, truncations, and protrusions for isolated cells and individual cells within a chain, maintaining a proper rod-shaped morphology. However, it struggled with clustered cells, often producing indented or truncated masks (Supplementary Fig. 2a). We reasoned that this limitation might reflect the broader and less well-defined membrane signal in raw fluorescent images, particularly in densely packed regions (Hardo et al., 2024). To test whether improving membrane definition could enhance segmentation accuracy, we retrained Omnipose using deconvolved images. Deconvolution reduces out-of-focus blur and produces sharper membranes (Agard, 1984; Scalettar et al., 1996). When evaluated on deconvolved membrane images, the retrained model, FMSeg_Omni (hereafter

FMSeg), performed better than Omnipose retrained and evaluated on the corresponding raw
membrane images, and reached a performance better than that of Cellpose-SAM retrained and
evaluated on raw membrane images (mean F1 score at IoU 0.8, 0.88 ± 0.09) (Fig. 2a). More broadly,
deconvolution affected model performance in a model-dependent manner (Fig. 2a; Supplementary
Fig. 2; Supplementary Fig. 3): Omnipose, microSAM and Morphometrics performed better on
deconvolved membrane images than on raw membrane images, whereas Cellpose-SAM and Cell-
pose3 showed the opposite trend, despite being fine-tuned separately for each image type.
158
159 FMSeg successfully identified and segmented single cells within groups, isolated cells, chains, and
160 even extremely elongated cells, regardless of the species or size (Fig. 2b,c; Supplementary Fig. 3).
161 Additionally, FMSeg successfully segmented yeast and cocci of various sizes, both isolated and
162 clustered, and worked with alternative membrane labels, including Mitotracker Green and GFP
163 fused to a homogeneously distributed membrane protein (QoxB) (Fig. 2d; Supplementary Fig. 4).

a Performance quantification of segmentation models

b Test data example

c Segmentation in a complex image (*B. subtilis*)

d Segmentation of different cell types and membrane dyes

Figure 2. Evaluation of models for fluorescent-membrane segmentation. a, Quantitative benchmarking of Om-
nipose (pink), CellposeSAM (blue), microSAM (yellow), Cellpose3 (green) and Morphometrics (gray) on fifty-six im-
ages of *B. subtilis*, *B. thuringiensis* and *P. megaterium*. Omnipose, CellposeSAM, microSAM, and Cellpose3 were fine-
tuned separately on raw and deconvolved fluorescent membrane images; models fine-tuned on raw images were eval-
uated on raw images, whereas models fine-tuned on deconvolved images were evaluated on deconvolved images. Top,
F1 score across Intersection over Union (IoU) thresholds. Lines are the mean F1 scores of 56 images. Dashed lines
correspond to model performance on raw fluorescent membrane images, and solid lines on deconvolved images.
Bottom, F1 at IoU of 0.8. The first violin plot of each pair (lighter color and delineated by a dashed line) corresponds
to model performance on raw images, and the second violin plot (darker color) to model performance on deconvolved
images. The horizontal dashed line in the violin plots indicate the median, and the dotted lines the first and third
quartiles. We called Omnipose model fine-tuned on deconvolved membranes FMSeg_omni (FMSeg) **b,** Examples of
images showing chains, clusters, and heavily elongated cells. Top, deconvolved micrographs of *B. subtilis*, *B. thuringiensis*
and *P. megaterium* cells stained with FM 4-64; bottom, FMSeg masks. F1 scores at IoU 0.8 are indicated at the top-right
corner of each image. Scale bar, 15 μm. **c,** FMSeg segmentation of a *B. subtilis* microcolony. The image is split into
three sectors. Segmentation results are shown in the middle sector, and FM 4-64 stained membranes in the flanking
sectors. Scale bar, 30 μm. **d,** Segmentation of cells of different shapes and sizes, with membranes labeled with FM 4-
64 or other fluorophores. Top, deconvolved fluorescent images; bottom, FMSeg segmentation. *Saccharomyces cerevisiae*,
Micrococcus sp., *Lactococcus lactis* and *B. pumilus* cells were stained with FM 4-64 (red). *B. subtilis* was stained with Mito-
Tracker Green (MTG). We also imaged a *B. subtilis* strain producing a GFP fused to the membrane protein QoxB
(QoxB-sfGFP, green), which is homogenously distributed in the membrane. See Supplementary Fig. 4 for bench-
marking of MTG and QoxB-GFP segmentation. Scale bars, 10 μm.

To facilitate the segmentation of fluorescent membrane images acquired with different microscopes, including systems without deconvolution capabilities, we trained an image restoration model to predict deconvolution from non-deconvolved images (FM2FM model; Fig. 3a). When direct deconvolution of membrane images is possible, it can be used directly. FM2FM provides an alternative for datasets in which this is not feasible. FM2FM was trained using the CARE framework (Weigert et al., 2018), in which a supervised U-Net predicts pixelwise fluorescence values while accounting for spatial context (i.e. neighboring pixel intensities). The fluorescence profiles of the predicted images closely matched those of deconvolved images (Fig. 3a-c), and the model could generate images amenable for segmentation from different microscopes (Supplementary Fig. 5).

Figure 3. Deconvolution prediction. **a**, Example of deconvolution prediction with FM2FM. Top, input raw fluorescent membrane images (Raw FM 4-64); middle row, true deconvolved fluorescent membrane images (Deconvolved FM 4-64); bottom, FM2FM-predicted image. White squares mark regions zoomed in at right (Block 1 and Block 2). Diagonal dotted lines indicate profile traces in **b**. Scale bars: full images, 10 μm; zoomed-in blocks, 5 μm. **b**, Fluorescence profiles across the dotted lines in Block 1 (top) and Block 2 (bottom). Light blue, raw FM 4-64; orange, deconvolved FM 4-64; burgundy, FM2FM prediction. **c**, Structural similarity index measure (SSIM) between FM2FM predicted images and deconvolved images. The distribution of SSIM values across 74 image crops is shown. **d**, Violin plots of cell width, length, surface area, volume, cross-sectional area, convex hull area, eccentricity and solidity calculated from deconvolved FM 4-64 images (orange) and FM2FM-predicted images (burgundy). The white dots and vertical lines within the violin plots represent the medians and standard deviations. P values from statistical comparisons between distributions (ANOVA or Kruskal test; see methods for details) are indicated in each panel. Twenty images (over 1000 cells) were processed and segmented with FMSeg. See the Methods for size calculation details.

To assess whether deconvolution prediction affects segmentation, we compared cell size parame-
ters—width, length, cross-sectional area, surface area and volume (see next section and methods
for details)—extracted from masks generated using deconvolved and FM2FM-predicted images.
We used multiple size metrics to verify that information relevant to downstream analyses is pre-
served across processing scenarios, as single metrics often fail to predict how image-derived meas-
urements propagate into downstream results (Gunawan et al., 2025). We also compared shape
descriptors not directly related to cell size, such as convex hull area, eccentricity and solidity. The
distributions were very similar, with no statistically significant differences (Fig. 3d), indicating that
membrane deconvolution prediction from raw membrane images has a minimal effect on segmen-
tation and does not impact cell size calculations.
219
Additionally, we trained a fluorescence translation model to infer deconvolved membranes from
cytoplasmic fluorescence (FP2FM model; Supplementary Fig. 6), which might be useful when
membrane staining is not feasible. Previously developed CARE restoration models for the predic-
tion of superresolution membranes can also be used (Spahn et al., 2022). Although these ap-
proaches expand the applicability of the method, they may introduce biases on cell size measure-
ments (Supplementary Fig. 6).
226
227 Altogether, our fine-tuned segmentation and deconvolution prediction models offer a robust ap-
228 proach for the high-throughput segmentation of individual bacterial cells with fluorescent mem-
229 branes, regardless of their size, shape and chaining pattern.
230
MEDUSSA, an analytical pipeline for extracting cell size from fluorescent membrane im-
ages.
We designed an algorithm to extract precise 2D cell dimensions—width and length—and extrap-
olate 3D measurements such as membrane surface area and volume (Fig. 4a). Starting from
FMSeg-generated masks, we applied a distance transform to compute, for each pixel, the shortest
distance to the mask periphery. We then overlaid a medial skeleton on this transform and sampled
the radius along the skeleton to obtain cell width across the long axis. Radii at the skeleton end-
points were used to estimate the size of the hemispherical caps, yielding length and width profiles
for each cell. Assuming rotational symmetry and imaging in the medial focal plane, we estimated
volume and surface area by summing cylindrical segments along the skeleton and adding hemi-
spherical ends. Additional details are provided in the Methods section.
242

Integrating segmentation with size calculations revealed two key factors critical for reliable measurements (Fig. 4b-d; Supplementary Fig. 7; Supplementary Fig. 8). The first is that the size of the segmentation masks is sensitive to the focal plane (Fig. 4b,c; Supplementary Fig. 7). For rod-shaped cells, width is maximal at the mid-cell focal plane but decreases above and below it (Fig. 4c) (Ursell et al., 2017). It is therefore critical to ensure that size measurements are taken from mid-cell focal planes. However, due to minor aberrations in microscopy slides, cells within a field of view often reside in slightly different focal planes (Fig. 4b; Supplementary Fig. 7). To address this, we acquired z-stacks and projected them onto a single plane before predicting deconvolution with FM2FM and segmentation (Fig. 4b). This approach yielded size estimations equivalent to those obtained from mid-cell focal planes (Fig. 4c), enabling accurate measurements across entire microscopy field, regardless of focal plane differences.

Figure 4. Cell size estimation from fluorescent-membrane segmentation masks. a, Cell size calculation work-flow. **b**, Representative z-slices from an optical-sectioning stack of a 14-cell *B. subtilis* chain stained with FM 4-64. Different cells are in focus in different slices. Circled cells are analyzed in **c**. The last image is a maximum-intensity projection of whole stack, processed FM2FM, where all cells appear in focus. Scale bar, 15 μm. Full stack in Supplementary Fig. 3. **c**, Width profiles (solid blue lines) of the cells circled in **b** on different z-slices across the optical sectioning stack: Cell 1 (right graph), Cell 7 (middle graph) and Cell 14 (right graph). Width is maximized on the slice in which the cell is in focus (vertical dotted lines). Cell width calculated from projected images is indicated by hori-zontal dashed lines. Results were equivalent regardless of the projection type (maximun intensity, red dashed line; average intensity, orange dashed line; sum, purple dashed line). Cells were segmented using JFilament in every z-slice and projections with FMSeg. **d**, FMSeg masks are slightly wider than manual JFilament masks. Density plots show the width distribution of 1100 bacilli cells. The top plot compares JFilament masks (blue) to FMSeg masks (green). After transforming the data with our sampling strategy (transformed masks, bottom plot), the distributions align bet-ter. The similarity of the distributions was quantitatively compared using Kullback–Leibler Divergence (KL div, values in insets within the graphs). See Supplementary Fig. 8 and the Methods section for details. **e**, Widths of wild-type (WT, blue) and Δ*ponA* (orange) cells from sporulating *B. subtilis* cultures, measured from deconvolved fluorescent-membrane images segmented with FMSeg (FM 4-64 deconvolved; 326 and 617 cells for WT and Δ*ponA*, respectively), or from CryoEM images (24 and 51 cells for WT and Δ*ponA*, respectively). Dashed lines mark the median: WT 0.70 μm (FM) vs 0.81 μm (cryo-EM); Δ*ponA* 0.54 μm (FM) vs 0.63 μm (cryo-EM). FM widths were ~14% lower than cryo-EM, but the WT/ΔponA ratio was nearly identical (1.300 vs 1.304; <1% difference). Cryo-EM images courtesy of E. Tocheva (Δ*ponA*) and E. Villa & K. Khanna (WT).

Second, we observed that the FMSeg model consistently generated masks slightly wider than ground truth masks (Fig. 4d; Supplementary Fig. 8a). This bias propagated into surface area and volume calculations (Supplementary Fig. 8b). Fortunately, the overestimation was independent of cell size, allowing us to apply a systematic correction (Fig. 4d; Supplementary Fig. 8b). We devel-oped a size correction pipeline that allowed us to predict the "true" cell size based on the dimen-sions of FMSeg-generated masks (Supplementary Fig. 8; see methods for details). We validated this approach by correcting the segmentation outputs for *B. subtilis*, *B. thuringiensis* and *P. megaterium* cells used in our benchmarking tests (Fig. 2a). As shown in Fig. 4d, FMSeg-mask width distribution was slightly shifted compared to that obtained from ground-truth masks generated using JFila-ment. After applying the correction, both distributions became nearly identical (Fig. 4d).

To assess the accuracy of our cell size measurements, we compared estimates from our pipeline with those obtained from plunged-frozen *B. subtilis* imaged by cryo-electron microscopy (cryoEM), a technique that provides high-resolution images of cells preserved in their native physiological state. We focused on sporulating *B. subtilis* cells, as we could access sets of cryoEM images of these cells, kindly provided by Elitza Tocheva (Tocheva et al., 2013), and Elizabeth Villa and Kanika Khanna (Khanna et al., 2021; Khanna et al., 2019; Lopez-Garrido et al., 2018). To test robustness across different cell sizes, we measured both wild type cells and cells lacking PBP1 (Δ*ponA*), which are ~25% thinner than wild type cells (Popham & Setlow, 1996). Our pipeline reproduced the difference in width between wild-type and Δ*ponA* cells observed by cryo-EM, although the abso-lute width estimates were on average ~14% lower than those obtained by cryo-EM (Fig. 4e). This discrepancy may reflect a systematic bias in our fluorescence-based measurements, but it could

298 also arise from differences in sample preparation, imaging modality, or growth conditions between
299 the two datasets. These results indicate that our pipeline provides realistic cell size estimates and
300 is robust to biologically relevant differences in cell width.
301
302 Overall, our fine-tuned segmentation model, coupled with deconvolution prediction and size cor-
303 rection, enables high-throughput, reliable cell size measurements from fluorescent-membrane mi-
304 croscopy images. We refer to the full pipeline for bacterial cell-size measurement as MEDUSSA
305 (MEmbrane DeconvolUtion and Segmentation for Size Analyses) (Supplementary Fig. 9).
306
Insight into intraspecific cell-size diversity in *P. megaterium*.
We applied MEDUSSA to study variation in cell size across strains of *P. megaterium* (de Bary, 1884),
a member of the *Bacillaceae* generally considered to produce large cells but for which intraspecific
morphological diversity has been reported (Logan & Vos, 2015; López & Alippi, 2009; Murumkar
et al., 2012). We focused on six strains widely used in physiology and biotechnology: the type strain
DSM 32 (ATCC 14581) (Arya et al., 2014) from the German Collection of Microorganisms and
Cell Cultures (DSMZ); the classic multi-plasmid strain QM B1551 (Eppinger et al., 2011); the
naturally plasmidless strain DSM 319 (Eppinger et al., 2011; Stahl & Esser, 1983); KM, used in
early bacteriophage studies; 899, also used in early bacteriophage and bacteriocin studies; and
WH320 (Rygus et al., 1991), a Lac⁻ derivative of DSM 319 commercially available for protein-
expression. We sampled cultures of each strain during exponential and stationary phases in LB
medium (Supplementary Fig. 10), imaged them in agarose pads supplemented with FM 4-64 for
membrane staining, and processed the images through our complete pipeline to obtain estimations
of the sizes of individual cells (Fig. 5; Supplementary Fig. 11a). Although all strains tended to form
chains (Fig. 5a), MEDUSSA could resolve single cells with high precision. Single-cell dimensions
were generally consistent across experimental runs, although we observed variation in population
averages of some strains (up to 27% for cell length and 12% for cell width; Supplementary Fig.
11a) likely due to day-to-day differences in conditions outside experimental control. Technical
variation among samples from the same culture was low (2-3% at most for cell width and length)
(Supplementary Fig. 11b).
327
We observed striking between-strain diversity in cell width: during exponential growth, median
width ranged from 0.92 μm in WH320 to 1.39 μm in 899 (Fig. 5b; Supplementary Figure 11a).
Most strains were slightly thinner in stationary phase, except KM, whose width distribution broad-
ened without a clear shift in the median (Fig. 5b). Although width showed a smaller coefficient of

variation (CV) than length—consistent with tighter control—the width CVs of *P. megaterium* strains

were generally higher than those reported for *B. subtilis* (Juillot et al., 2021; Nordholt et al., 2020;

Sauls et al., 2019; Taheri-Araghi et al., 2015; also reproduced by us in Supplementary Fig. 11a),

with some strains reaching ~0.15 in exponential or stationary phase (Supplementary Fig. 11a).

Notably, cells of different widths were sometimes present within a single chain (arrowheads in Fig.

5a).

Figure 5. Morphological characterization of six *P. megaterium* strains. a, Fluorescence micrographs of FM 4-
64-stained cells of different *P. megaterium* strains sampled form exponential and stationary phase LB cultures: WH320,
DSM 32, KM, QM B1551, DSM 319 and 899. Arrowheads in the KM stationary image point to thicker (white) and
thinner (yellow) cells within a chain. Scale bars, 10 µm. **b-e,** Distributions of cell with (**b**), length (**c**), volume (**d**) and
surface to volume (**e**) for the indicated strains (strain names in panel **b** and **d**; same order maintained in the other
panels) during exponential (blue) and stationary (orange) phase. The dashed line in the violin plots indicate the median,
and the dotted lines the first and third quartiles. The inset in panel **c** shows a micrograph of a long cell produced by
DSM 32 during exponential growth (membranes, red; scale bar, 20 µm). The following number of cells were analyzed
for each strain in exponential and stationary phase: WH320, 153 and 961 cells; DSM 32, 192 and 904 cells; KM, 149
and 856 cells; QM B1551, 292 and 447 cells; DSM 319, 195 and 471 cells; 899, 758 and 548 cells. The central 95% of
observed values are represented. See Supplementary Fig. 11 for comparison between experimental runs. Prior to
measurement, segmentation output was manually curated to remove aberrant masks and truncated masks at image
edges.

We also observed significant differences in cell length across strains: in exponential phase, the

median ranged from 4.06 µm in DSM 32 to 5.76 µm in strain 899, and cells shortened in stationary

phase, as expected (Fig. 5a,c). Within-strain variation in cell length was high in some cases (Figure

5c; Supplementary Fig. 11a), with rare, extremely elongated cells occurring in several strains; the

more dramatic cases were in DSM 32, where approximately 10% of the cells were over 10 µm long

(Fig. 5c, inset). These extreme elongation events were not observed in stationary phase (Figure

5c), suggesting that it is a phenomenon associated with active growth and that these filaments

septate into multiple cells or lyse upon entry into stationary phase.

These width and length differences underpinned differences in surface-to-volume ratios—thinner

cells exhibited higher surface-to-volume ratios than thicker cells (Ojkic et al., 2019, 2022)—and

contributed to the diversity of cell volumes observed between strains, spanning nearly threefold

(Fig. 5d-e).

***P. megaterium* WH320 carries a partially functional version of PBP1.**

Most accounts of *P. megaterium* cell width report values of ~1.2-1.5 µm (Logan & Vos, 2015).

However, two of our strains, DSM 32 and WH320, had widths below these values. We were par-

ticularly surprised by WH320, whose median width during exponential growth was ~0.9 µm, as

this strain is a derivative of DSM 319, one of the thickest strains in our panel (median width in

exponential phase ~1.4 µm).

WH320 was generated by ethyl-metanosulfonate (EMS) mutagenesis of DSM 319, selecting for

Lac⁻ clones (Rygus et al., 1991). One potential explanation for the reduced cell width of WH320

is that an additional mutation introduced during EMS mutagenesis or at some point after the two

strains diverged affected cell width. To explore this possibility, we sequenced the genome of

WH320 and compared it to the parental DSM 319 reference genome (Eppinger et al., 2011). The

two strains differed by 37 single nucleotide polymorphisms, with the majority of them being

G·C→A·T transitions, consistent with EMS mutagenesis (Sega, 1984) (Supplementary Data in

Zenodo repository). We also observed other mutations including transversions and indels, which

might have occurred during transfer of the strains in the lab before we acquired them. One of the

transitions caused a glycine-to-arginine amino acid substitution in the catalytic site of *P. megaterium*

β-galactosidase, likely responsible for the Lac⁻ phenotype. We also found a C to A transversion in

ponA that produced an alanine-to-aspartic acid change at residue 214 in the transglycosylase do-

main of the classA penicillin binding protein PBP1 (Supplementary Data in Zenodo repository).

Figure 6. WH320 carries a hypomorphic PBP1 allele. **a,** Representative FM 4-64-stained cells corresponding to
panel b. Scale bar, 10 μm. **b,** Cell width of a *B. subtilis* PY79 *ponA* mutant (Δ*ponA*, blue) and the same mutant expressing
P. megaterium DSM 319 *ponA* (Δ*ponA* P_spank-*ponA*_DSM319, orange) or *P. megaterium* WH320 *ponA* (Δ*ponA* P_spank-*ponA*_WH320,
green) from the IPTG-inducible P_spank promoter inserted at the *amyE* locus. Violin plots show width distributions at
different IPTG concentrations. Central 99% of data points was sampled for each strain in each condition. The dashed
line indicates the median; dotted lines mark the first and third quartiles. The number of cells for each violin is shown
in the plot under each violin. Prior to measurement, segmentation output was manually curated to remove aberrant
masks and truncated masks at image edges. Width distributions of the three strains were statistically different from
each other at all IPTG concentrations (Kruskal test).

As noted above, PBP1 modulates cell width in *B. subtilis*, with *ponA* mutants being ~25% thinner

that the wild type and *ponA* overexpression producing cells thicker than the wild type (Fig. 4e)

(Dion et al., 2019; Popham & Setlow, 1996). We therefore hypothesized that the *ponA* mutation

reduces PBP1 activity and underlies the thin phenotype of WH320. To get insight into this possi-

bility, we asked whether the *ponA* alleles from DSM 319 and WH320 could complement a *B. subtilis*

ponA mutant when produced ectopically. We placed both alleles under the IPTG-inducible pro-

moter P_{spank} at the ectopic *amyE* locus, and measured cell width after incubation with varying IPTG

concentrations. As shown in Fig. 6, expression of DSM319 *ponA* produced a marked increase in

cell width, even at low IPTG concentrations. In contrast, expression of WH320 *ponA* caused a

milder increase in cell width and never reached the values observed with $ponA_{DSM319}$, even at high

induction levels. These results indicate that the alanine-to-aspartate substitution in *ponA* reduces

PBP1 function.

DISCUSSION

Here, we introduce MEDUSSA, an image-analysis pipeline that extracts accurate single-cell size measurements from fluorescent-membrane microscopy images, and demonstrate its utility for dissecting mechanisms of cell-size diversity among bacterial strains. The workflow combines deep-learning–based image restoration for deconvolution prediction with automated segmentation to delineate cell boundaries from membrane signal, followed by mask analysis to estimate cell size. Because each step is modular, the restoration (FM2FM), segmentation (FMseg), and measurement components can be used independently to suit specific study needs (Supplementary Fig. 9). Applied across strains and species, MEDUSSA yields cell-size measurements that can be further interrogated to identify underlying mechanisms.

MEDUSSA is well suited for comparative studies because it addresses key shortcomings of previous methods. First, fluorescent membrane imaging enables precise delineation of cell boundaries by tracking the peak membrane signal around each cell (Fig. 1). This supports consistent training-set annotation and yields more reliable cell-size estimates than phase-contrast-based approaches (Hardo et al., 2024) (Fig. 1). Second, membrane dyes reveal septal membranes, allowing clear separation of daughter cells that remain attached after division and thus enabling reliable length measurements for both isolated cells and chains—something that is difficult with phase contrast (Fig. 1) (Syvertsson et al., 2016). Third, the pipeline corrects focal-plane differences among cells within the same image, which would otherwise bias measurements because focal-plane shifts alter mask size (Fig. 4b-c) (Ursell et al., 2017). Specifically, projecting fluorescence images from a z-stack and applying a deconvolution-prediction restoration algorithm (FM2FM) produces segmentation-ready images that yield mid-focal-plane masks. Thus, this standardized protocol of restoration, segmentation, and measurement converts membrane-fluorescence images into population-level size distributions.

A trade-off is the need for fluorescent labeling. Several membrane and cell-wall dyes are routinely used in bacterial cell biology (Greenspan & Fowler, 1985; Herman et al., 1994; Kuru et al., 2012; Lucidi et al., 2024; Monteiro et al., 2015; Pogliano et al., 1999; Sharp & Pogliano, 1999). Most experiments presented here use FM 4-64, a lipophilic styryl fluorophore that associates with the outer leaflet of the cytoplasmic membrane without disrupting membrane function (Pogliano et al., 1999). FM 4-64 is broadly applicable across bacterial species and compatible with live-cell imaging (Calvez et al., 2019; Dalebroux et al., 2015; Ma et al., 2009; Maloney et al., 2009; O'Hara et al., 2017; Ojkic et al., 2016; Shin et al., 2015; Specht et al., 2013; Studer et al., 2016; Sun & Margolin,

1998; Tsuge et al., 2008; Yeh et al., 2010). However, our segmentation model also works with other labels—including the membrane-permeable MitoTracker Green and with GFP fused to a membrane protein (Fig. 2d; Supplementary Fig. 4)—and, in principle, with any dye that provides uniform signal along the cell periphery. In addition, we provide a transformation model, FP2FM, that predicts membrane-like boundaries suitable for segmentation from cytoplasmic fluorescence, adding flexibility (Supplementary Fig. 6). Altogether, the available dyes, combined with the transformation tools provided here and elsewhere (Spahn et al., 2022), extend the applicability of the MEDUSSA method to a broad range of species. For direct comparisons, we recommend using the same staining modality and image transformation procedures, because different labels or transformation models may slightly affect size estimations. For example, FM 4-64 labels the outer leaflet of the cytoplasmic membrane, whereas GFP fused to a membrane protein is displaced toward the cytoplasmic side. Consequently, boundaries inferred from GFP-based signals may appear slightly inward relative to those inferred from FM 4-64, leading to systematic differences in estimated cell size (Supplementary Fig. 12).

Although MEDUSSA addresses several limitations associated with cell-size estimation from phase-contrast images, it does not overcome challenges inherent to the imaging of small bacterial cells by optical microscopy. A single pixel in a fluorescence microscopy image may amount to 10% of the width of a typical bacterial cell, so even minor segmentation imprecisions can translate into appreciable differences in measured size. For reliable comparisons between strains and replicates, it is therefore important to quantify sufficiently large numbers of cells, and compare the size distributions of the populations. Bacterial cell size also depends on growth conditions (Schaechter et al., 1958), and biological variation between cultures can be substantial depending on media and growth stage (Bakshi et al., 2021; Cylke & Banerjee, 2023; Kellogg & Levin, 2022). We therefore recommend to keep growth conditions as constant as possible to obtain biologically meaningful and reproducible results.

Segmentation accuracy also depends on image quality. In particular, images with low signal-to-noise ratios can result in lower segmentation accuracy (Supplementary Fig. 13). To mitigate this limitation, we trained an additional CARE restoration model (FM2FM-HiSNR; see methods for details) that predicts high signal-to-noise deconvolved membranes across a range of input signal-to-noise levels. Under our imaging conditions, restoration with this model improves the segmentation quality of low signal-to-noise images (Supplementary Fig. 13), although extremely low signal levels will ultimately limit segmentation accuracy.

In addition, projection artifacts can lead to cell-size misestimation from fluorescent images, an

effect that may be particularly pronounced for smaller cells (Hardo et al., 2024; Ursell et al., 2017).

Hardo et al. reported marker-dependent biases, with membrane fluorescence tending to underes-

timate cell width and cytoplasmic fluorescence tending to overestimate it, and with proportionally

larger effects in thinner cells (below ~1 μm) than in thicker cells. To probe for this effect under

our imaging and analysis conditions, we compared cell width estimates derived from independently

segmented membrane and cytoplasmic fluorescence across a controlled width range (*ponA* under

IPTG; Supplementary Fig. 14). If projection artifacts were prominent under our conditions, we

would expect cytoplasmic-fluorescence-based estimates to be relatively larger than membrane-

based estimates for thin cells, with the discrepancy decreasing as cells become thicker. However,

width scaled almost identically when estimated from membrane and cytoplasmic fluorescence

across the tested width range (~0.6-1.2 μm) (Supplementary Fig. 14b). These results suggest that

projection artifacts, as described by Hardo et al., do not measurably bias our width estimates under

these conditions, although they may become more relevant in cells thinner than those examined

here or under different imaging settings.

In this study, we applied MEDUSSA to quantify the size of six strains of *P. megaterium*. Our

measurements reveal remarkable diversity in cell size both among strains and among individual

cells within a strain (Fig. 5). Although all strains produce comparatively large cells by bacterial

standards, some produce cells that are, on average, over twofold larger in volume than others. We

also observed that some strains formed extremely elongated cells during exponential growth,

which were not observed in stationary phase. To our knowledge, filamentation has not been

reported previously in *P. megaterium*—likely because membrane dyes are not routinely used in

microscopy. The mechanism underlying this filamentation, and whether such filaments occur in

nature or are artifacts of laboratory culture, remains to be determined.

Cell width in bacteria is generally assumed to be tightly controlled within a species, with little cell-

to-cell variation in steady state cultures (Campos et al., 2014; Chien et al., 2012; Ojkic & Banerjee,

2021; Ojkic et al., 2022; Sauls et al., 2019; Taheri-Araghi et al., 2015). Our dataset, however, reveals

substantial width variation both within and between strains. Within strains, width values were

approximately normally distributed, and some strains showed wide cell-to-cell variation; for

example, in strain KM during stationary phase, the thinnest cells were up to twofold thinner than

the thickest cells (Fig. 5). The coefficient of variation for width reached up to 0.15 (Supplementary

Fig. 11)—more than twice that of *B. subtilis*—suggesting that mechanisms of width homeostasis

in this bacterium are less precise than in *B. subtilis* (Nordholt et al., 2020; Sauls et al., 2019).

Between strains, width distributions were shifted, with median widths ranging from ~0.9 to 1.4

μm (Fig. 5; Supplementary Fig. 11a), indicating variation in width regulation across strains. We

further examined differences between the thick strain DSM 319 and its Lac⁻ derivative WH320,

which is ~0.5 μm thinner. Genome comparison revealed an alanine-to-aspartic acid substitution

in the transglycosylase domain of PBP1. An alignment of PBP1 sequences from 854 strains

indicates that alanine at position 214 is largely conserved (~75%); ~24% carry serine at this

position, and fewer than 1% contain other residues (e.g., cysteine, threonine, valine)

(Supplementary Data in Zenodo repository). Notably, no sequence harbors an aspartic acid or any

other charged residue at this position, suggesting that introducing charge there impairs PBP1

activity. Because PBP1 mutants yield thinner cells in *B. subtilis* (Popham & Setlow, 1996), it is

plausible that the hypomorphic PBP1 allele in *P. megaterium* WH320 contributes—at least in part—

to its reduced width relative to the DSM 319 parental strain.

Unlike the mutation in *lacZ*, the *ponA* mutation in *P. megaterium* WH320 is a G→T transversion.

Because EMS typically induces G·C→A·T transitions (Sega, 1984), it is unlikely that the *ponA*

mutation arose directly from EMS mutagenesis. Instead, we suspect that the *ponA* mutation in

strain WH320 was either already present in the DSM 319 clone used for EMS treatment or was

acquired during subsequent propagation of WH320 before we received the strain.

Overall, these results illustrate how MEDUSSA can be used to explore within- and between-strain

diversity in bacterial cell size and to reveal differences that can be pursued to identify underlying

mechanisms. Future work will expand our strain set and survey additional species and

environmental isolates to advance our understanding of the evolution of bacterial cell size and

contribute to the emerging field of evolutionary cell biology (Jiang et al., 2015; Lynch et al., 2014).

METHODS

Strains and plasmids

A list of strains is provided in Supplementary Table 1. Plasmid and primer lists are in Supplementary Tables 2 and 3, along with detailed descriptions of plasmid construction.

Culture media and growth conditions

Bacteria were routinely grown on LB agar plates (Lennox formulation, 5 g NaCl/L) overnight at 30 °C, with the exception of *Lactococcus lactis*, which was grown on MRS medium at 28 °C. *Saccharomyces cerevisiae* was grown on YPD at 37°C.

For size quantification of *B. subtilis* and *P. megaterium* strains, a single colony was inoculated into 5 mL fresh LB Lennox and grown overnight in a roller drum at 30 °C with constant rotation. The next morning, cultures were diluted 1:1000 into 30 mL of the appropriate medium in 250-mL baffled flasks and incubated in a water bath at 30 °C with shaking at 220 rpm. Samples were taken after five hours (exponential; Supplementary Fig. 10) and after ten hours (stationary; Supplementary Fig. 10) for microscopy (see next section).

For the induction experiments shown in Fig. 6 and Supplementary Fig. 14, a single colony was inoculated into LB Lennox diluted 1/4, then grown at 25 °C in a shaking incubator at 220 rpm. The next morning, cultures were diluted to an OD600 of 0.02 in 1/4-diluted LB Lennox with the indicated IPTG concentration (0, 25, 50, 100, 500, or 1000 μM). Cultures were grown at 30 °C in a shaking incubator to an OD600 of 0.2–0.5 and then samples were taken for microscopy.

Sporulation (Fig. 4e) was induced by resuspension, as described elsewhere (Riley et al., 2018).

Microscopy

For imaging, 10 μL culture samples were imaged on 1.2% agarose pads in 1/4-diluted LB-Lennox supplemented with 0.5 μg/mL FM4-64. The pad was covered with a #1.5 glass coverslip. Ten-hour samples were diluted tree times to avoid overcrowded fields.

For cell-size estimation experiments for the different *B. subtilis* and *P. megaterium* strains, cells were imaged using a Leica Thunder Imager live cell with a Leica DFC9000 GTC sCMOS camera and a 100X HC PL APO objective (Numerical aperture 1.4, Refractive index 1.518 objective). FM 4-64 images were taken using a 510 nm excitation LED light source, with 70% intensity and 0.1 s exposure, and a custom filter cube produced by Chroma Technology GmbH (Olching, Germany)

with the following components: excitation filter, ET505/20x; dichroic mirror, T660lpxr; emission

filter, ET700lp. Training images for the restoration (FM2FM and FM2FM-HiSNR) and translation

(FP2FM) models were taken with a DeltaVision Ultra microscope with a PCO Edge sCMOS

camera and an Olympus 100X UPLX APO objective (Numerical aperture 1.45, Refractive index

1.518 Ph3). FM 4-64 and mVenusQ69M were visualized with the following filters: excitation 542

nm and emission 679 nm for FM 4-64, and excitation 475 nm and emission 525 nm for

mVenusQ69M, with light transmission at 100% and an exposure time of 0.2 s for both channels.

Training images for the segmentation models (FMSeg, RawFMSeg) were taken using the same

illumination conditions as the restoration model. For the membrane images of *E. coli*, the FM4-64

was captured with light transmission of 40% and an exposure time of 0.1 s.

For phase-contrast microscopy, the aforementioned DeltaVision microscope was used, light

transmission was set to 100% and exposure time to 0.1 s.

For microscopy comparisons, we also used a Zeiss Axio Imager Z2 microscope with a Hamamatsu

sCMOS camera and a Zeiss 100x Plan-Apochromat objective (Numerical aperture 1.40, Refractive

index 1.518 Ph3). FM 4-64 was visualized with an excitation filter of 506 nm and emission filter

of 751 nm, with light transmission at 100% and an exposure time of 0.2 s.

Ground truth image annotation using active contours

We annotated cells in our ground-truth images using the ImageJ plugin JFilament (Smith et al.,

2010), which allows the semi-automatic generation of sub-pixel resolution active contours (snakes)

that follow the ridges of maximum fluorescence membrane intensity around the cell (Fig. 1f; Sup-

plementary Fig. 1) (Lopez-Garrido et al., 2018; Ojkic et al., 2014, 2016). Default parameters were

used except for Gamma, which we increased when the preset value caused excessive deformation

of the snakes. For cell clusters, snakes were typically manually corrected because contour edges

tended to deform toward neighboring cells. When membrane blobs distorted cell contours, we

also corrected the snakes to follow an appropriate straight line. These steps ensured consistent

single-cell segmentation for model training and improved robustness of the segmentation model

to such confounding artifacts.

Training and evaluation of models for segmenting fluorescent membranes.

We annotated 143 raw and SoftWorx (Cytiva Life Sciences)-deconvolved FM 4-64 membrane

images using JFilament (Smith et al., 2010), as described above, yielding over 7000 cells for train-

ing. Training images were maximum-intensity projections of FM4-64 z-stacks. Following the Om-
nipose documentation (Cutler et al., 2022), we trained the models from scratch, i.e., not using any
existing model parameters as starting points. In contrast, for Cellpose3 (Stringer & Pachitariu,
2025), Cellpose-SAM (Pachitariu et al., 2025) and microSAM (Archit et al., 2025), the existing base
models were fine-tuned with the existing data. All trainings were done following the suggestions
found in their respective documentations. Omnipose models were trained with a batch size of 8
images for 4000 epochs using a learning rate of 0.1 and a Rectified Adam (RAdam) optimizer (Liu
et al., 2019). Both models (raw and deconvolved) were trained using the same parameters. Cell-
pose3 models were trained with a batch size of 1 image for 100 epochs using a learning rate of 0.1,
a weight decay of $1x10^{-4}$, and a Stochastic Gradient Descent (SGD) optimizer (Sutskever et al.,
2013). Both models (raw and deconvolved) were trained using the same parameters. Cellpose-
SAM models were trained with a batch size of 1 image for 100 epochs using a learning rate of
$1x10^{-5}$, a weight decay of 0.1, a tile size of 256x256 px, and an AdamW optimizer (Loshchilov &
Hutter, 2017). Both models (raw and deconvolved) were trained using the same parameters. Mi-
croSAM models were trained with a batch size of 1 image for 10 epochs of 1788 iterations per
epoch, using a learning rate of $1x10^{-5}$, a weight decay of 0.1, a tile size of 512x512 px, and an
AdamW optimizer. Segmentation decoder was also trained to allow for automatic instance seg-
mentation. Both models (raw and deconvolved) were trained using the same parameters.

Training was done in the "deepbio" computing cluster at the Max Planck Institute for Evolution-
ary Biology, using a single NVIDIA(R) GeForce GTX 2080 Ti GPU with 12GB memory.

For accuracy assessment, we annotated 56 images with JFilament. F1 score (Caicedo et al., 2019;
Maier-Hein et al., 2024; Taha & Hanbury, 2015) was calculated at eleven Intersection over Union
(IoU) thresholds spanning 0.5 to 1, in 0.05 intervals. True positives, false positives, and false neg-
atives were calculated using Omnipose library function "average_precision". F1 was then com-
puted using these values following Caicedo et al. 2019 (Caicedo et al., 2019).

Simulation of fluorescent membranes
Synthetic rod-shaped cells were generated first by generating spheres of defined radii and then
stretching them in the respective center point in the X-direction only. Contours were then obtained
in each plane of each synthetic cell to mimic membranes. Contour spatial coordinates were ex-
tracted for simulating microscopy images using BlurLab (Ursell et al., 2017). For each contour

point, an intensity value was sampled from a normal distribution of mean 1000 and standard de-
viation 50. For simulating light spread, a Point-Spread Function (PSF) was simulated in BlurLab
using our image acquisition parameters: Numerical aperture of 1.4, refractive index of 1.518, emis-
sion wavelength of 679nm and magnification of 100X. Gaussian noise of mean 100 and standard
deviation 50 was added to the simulations. We then simulated Z-stacks of 4000 nm with 50 equi-
distant Z-slices for quantification.

Training and evaluation of models for deconvolution prediction (FM2FM) and fluores-
cence translation (FP2FM)
Deconvolution-prediction and fluorescence-translation models were trained using the Content
Aware Restoration (CARE) framework (Weigert et al., 2018). For FM2FM and FP2FM model,
200 images (2048x2048 pixels) of vegetatively growing *B. subtilis* constitutively expressing
mVenusQ69M (Cox et al., 2010) were acquired using a DeltaVision Ultra microscope with a PCO
Edge sCMOS camera, capturing FM4-64 membrane fluorescence, and mVenusQ69M cytoplasmic
fluorescence. Images were deconvolved in the DeltaVision software (SoftWorx) and then cropped
to remove deconvolution edge artifacts, yielding 1914x1914 px image outputs. Non-deconvolved
training inputs were cropped to the same size. Training input images were single-plane images of
raw FM4-64 (for the FM2FM model) or mVenusQ69M (for the FP2FM model) z-stacks, and
target images were the deconvolved FM4-64 images for both models. For training, 16 patches of
512x512 px were extracted from each image pair. Models were then trained using a batch size of
32 images for 100 epochs, with 100 training steps per epoch. Training was performed in the "deep-
bio" computing cluster at the Max Planck Institute for Evolutionary Biology, as above. Model
parameters and training hardware was the same for both models.

To compare model performance against the membrane deconvolution target (SoftWorx), we con-
ducted qualitative assessments of output images intensity profiles. We quantified the Structural
Similarity Index Measure (SSIM) between extracted patches of the images. Patches were selected
to maximize number of cells and minimize background. SSIM was calculated using the Python
scikit-image library (van der Walt et al., 2014). We then statistically compared cell-size estimates
using membrane images deconvolved with SoftWorx or predicted with FM2FM or FP2FM (Fig.
3; Supplementary Fig. 6). Width, Length, Surface Area, and Volume were measured using the
skeletons as explained further down. Cross-sectional area, convex hull area, eccentricity, and so-
lidity, were directly obtained from segmentation masks using the Python scikit-image library (van
der Walt et al., 2014).

Training and evaluation of a deconvolution-prediction model that increases the signal-to-noise ratio (SNR) of predicted images (FM2FM-HiSNR).

We use CARE (Weigert et al., 2018) to train an additional deconvolution prediction model that produces high signal-to-noise ratio (SNR) images from low SNR non-deconvolved fluorescent membrane images. We named the model FM2FM-HiSNR. FM 4-64 images of vegetatively growing *B. subtilis* were acquired at four different illumination conditions using a DeltaVision Ultra microscope with a PCO Edge sCMOS camera. The illumination conditions were as follows (exposure time and percentage of transmitted light listed after their names in quotation marks): "Noisy", 0.01s, 10%; "i1", 0.01s, 50%; "i2", 0.025s, 10%; "Target", 0.2s, 100%. Images were deconvolved using the DeltaVision software (SoftWorx) and then cropped to remove deconvolution edge artifacts, yielding 1914x1914 px image outputs. Non-deconvolved training inputs were cropped to the same size. Training input images were single-plane images of raw FM4-64 z-stacks from all different illumination conditions, and target images were the deconvolved FM4-64 images at the "Target" illumination condition. In total, 98 input images were used. For training, 16 patches of 512x512 px were extracted from each image pair. Model was then trained using a batch size of 32 images for 100 epochs, with 100 training steps per epoch. Training was performed in the "deep-bio" computing cluster at the Max Planck Institute for Evolutionary Biology, as above.

To explore the relationship between SNR and segmentation accuracy, and to test whether restoration with FM2FM-HiSNR improve segmentation accuracy of low SNR images, FM 4-64 images of vegetatively growing *B. subtilis* were acquired at 12 different illumination conditions using a DeltaVision Ultra microscope, under the following illumination conditions (exposure time and percentage of transmitted light listed after their names in quotation marks): "Noisy", 0.01s, 10%; "i1", 0.01s, 50%; "i2", 0.025s, 10%; "i3", 0.025s, 50%; "i4", 0.05s, 10%; "i5", 0.05s, 50%; "i6", 0.1s, 10%; "i7", 0.1s, 50%; "i8", 0.1s, 100%; "i9", 0.2s, 10%; "i10", 0.2s, 50%; "Target", 0.2s, 100%. Images were deconvolved using the DeltaVision software (SoftWorx) or restored with FM2FM-HiSNR, and segmentation accuracy was calculated (Supplementary Fig. 13).

Extracting cell size information from masks of rod-shaped cells.

For rod-shaped bacteria, we assumed a sphero-cylindrical geometry with hemispherical caps (Ojkic et al., 2019). First, we computed the minimum Euclidean distance from each pixel inside the mask to the closest pixel on the mask boundary (Borgefors, 1986), which is commonly referred to as the distance transform. Next, we generated a one-dimensional centerline (commonly known as "skel-

eton") (Zhang & Suen, 1984), overlaid it on the distance transform map, and determined the dis-

tance value at each skeleton pixel, thereby obtaining the local radius of the mask along the skeleton

line (Fig. 4a). An alternative one-dimensional representation is the medial axis (Blum, 1967), also

known as the "topological skeleton" (Supplementary Fig. 15). We compared size estimates ob-

tained with the two representations and found significant differences in some size metrics in a set

of *E. coli* cells (Supplementary Fig. 15d). Inspection of individual masks revealed that some medial-

axis representations tended to over-branch (Supplementary Fig. 15e). If common across many

cells, such over-branching would lead to inaccurate size estimates. We therefore proceeded with

the skeleton representation.

To estimate the extend of the hemispherical caps, we took the distance between each skeleton

endpoint and the corresponding mask tip as the cell radius at the skeleton endpoints. This yielded

precise estimates of mask length and with profiles at every pixel along the skeleton. Assuming

rotational symmetry and that the masks derive from medial focal-plane images, we estimated cell

volume and surface area by adding the volumes and lateral surface areas of cylindrical segments

generated by circularly projecting the radius of the mask at each pixel of the skeleton, plus the

hemispherical caps (Fig. 4a). This approach accurately handles elongated masks, provides local

morphological information of cell morphology, and is robust to cellular curvature. While tailored

to rods, the framework can be adapted to other morphologies.

Dealing with branching skeletons

Due to the mask geometry, it is possible that the skeleton branches into multiple endpoints (Sup-

plementary Fig. 15c). To handle both branched and non-branched cases, we developed the fol-

lowing workflow. First, we treat the skeleton as a graph, with each pixel considered a node. As-

suming connectivity only between neighboring pixels (horizontal, vertical, or diagonal), we define

edges for pairs of pixels whose Euclidean distance is less than 1.5 pixels, thereby retaining con-

nected neighbors. Next, we implement a Depth-first search (DFS) algorithm to find the longest

possible path connecting the opposite endpoints in the graph. If the skeleton is not branched, this

path corresponds to the skeleton. For branched skeletons, multiple candidate paths exist. For each

candidate path we compute the cell size metrics described in the next section and take the median

of each metric as the cell size (Supplementary Fig. 15).

Cell size measurements:

We combined the mask skeleton and distance transform to derive cell measurements under a spherocylindrical geometry with hemispherical caps and rotational symmetry. The measurements are implemented in a custom code written in Python3, which can be found in the GitHub repository linked in the "Software and data availability section."

Cell length (L) is the sum of Euclidean distances d between two successive points in the mask skeleton, plus the distance from each skeleton endpoint and the corresponding mask tip, which corresponds to the local radius at the skeleton endpoint r_s. With that, L follows the equation:

$$L = r_{s1} + r_{s2} + \sum_{i=0}^{n-1} d(i,i+1). \qquad \text{Eq. (1)}$$

Cell width (w_i) is calculated as twice the value of the radius at each point in the skeleton (r_i). Mean w can then be calculated as:

$$\bar{w} = \frac{2}{n} \sum_{i=0}^{n} r_{c\,i}. \qquad \text{Eq. (2)}$$

Cell surface area (S) is calculated as the sum of lateral surface areas of the cylinders projected at each point in the skeleton, with radius r_c and height h of 1 pixel, plus the lateral surface area of the hemispherical caps with radius r_s. Thus, S is calculated as:

$$S = 2\pi \left(r_{s1}^2 + r_{s2}^2 + h \sum_{i=0}^{n} r_{c\,i} \right). \qquad \text{Eq. (3)}$$

Cell volume (V) is obtained similarly to S, as it adds the volume of each cylinder and the hemispherical caps (with same parameters of radius and height as S). V is then obtained from:

$$V = \pi \left(\frac{2}{3} r_{s1}^3 + \frac{2}{3} r_{s2}^3 + h \sum_{i=0}^{n} r_{c\,i}^2 \right). \qquad \text{Eq. (4)}$$

Measurement transformation

We segmented 864 single cells of *Escherichia coli* stained with FM 4-64 using either JFilament (ground-truth) or FMseg. Cells were from early exponential phase (OD_{600} = 0.1-0.3) cultures in either LB Lennox, or MOPS minimal medium with 0.1% glucose, providing a range of cell widths from 0.53 to 1.04 μm, and a range of cell lengths from 0.87 to 9.61 μm (based on measurements from ground-truth masks) (Supplementary Figure 8a). Measurements from ground-truth masks were plotted against the values obtained from on the FMseg-generated masks, revealing a consistent size overestimation for FMseg masks (Supplementary Figure 8b). The data shows a linear relationship for the overestimation, which allows us to leverage statistical tools to compute the

necessary parameters to transform the data between the automatic segmentation results to the ground truth results. A linear regression would be a direct way to do this, extracting the values for the slope (*m*) and intercept (*n*) necessary to transform the data. While linear regression returns a coefficient of error that would allow us to simulate distributions to sample *m* and *n*, there is the chance that we may over or under-estimate these values, as well as sample pairs that do not make sense in the context of our data. Thus, we leveraged the PYMC library (Abril-Pla et al., 2023) and its use of Bayesian statistics for sampling the necessary parameters, which are *m*, the fit slope, *n*, the fit intercept, and *σ*, the standard deviation of our sampled observations. Since our data behaves linearly, we used a Bayesian Generalized Linear Model (GLM), assuming that our parameters (slope, intercept, size metric, and its standard deviation) follow certain types of distributions. For *σ*, we assume a Half-Cauchy distribution, which only returns positive, non-zero values, which is the only type of values a standard deviation can have. For *m* and *n*, we sample from normal distributions with mean 0 and standard deviation 20, giving us a wide range of possible parameters. Then, for computing the observed values, we also assume a normal distribution. PYMC performs a series of draws to fit the data, and thus after successive draws we end with the distributions of *m*, *n* and *σ*. In our case, we performed four chains of 3000 draws each, giving us a total of 12000 sampled values. We can then randomly sample pairs from these distributions in order to transform the data, as well as to include the uncertainty of the transformation in the analysis (Supplementary Figure 8b). For the examples shown in Figure 5, Figure 6, and Supplementary Figure 11, we sampled 250 pairs of m and n for each individual cell measurement, thus allowing us to compute new metrics.

Statistical analyses

For the statistical tests in Fig. 3, Fig. 6, Supplementary Fig. 6, and Supplementary Fig. 15, Levene's test was computed for the distributions of interest. If the p value was greater than 0.05, one-way ANOVA was performed for obtaining the p values and defining if the difference between distributions was significant or not. In cases where Levene's test was significant ($p < 0.05$), Kruskal test was used instead. All three tests were used as they are implemented in SciPy (Virtanen et al., 2020).

CryoEM image analysis

Because the high magnification of the cryo-EM images typically did not capture entire cells, we estimated cell width as the straight-line distance between the mother-cell lateral membranes, measured perpendicular to the cell longitudinal axis. In cryo-FIB tomograms of wild-type cells, only a

portion of the cellular volume was retained: focused ion-beam milling produced a thin lamella that

sampled a section of the cell. We analyzed only tomograms in which this section included the mid-

cell region. To verify this, we measured width across successive z-slices and selected tomograms

showing a clear width peak at mid-cell that decreased in slices above and below.

Whole genome sequencing

For whole-genome sequencing, *Priestia megaterium* WH320 was streaked in an LB-Lennox agar plate

and grown overnight at 30°C. Next morning, a single colony was picked and re-streaked in an LB-

Lennox agar plate and grown overnight at 30°C. Cells were collected from the plate and genomic

DNA was extracted with the MasterPure Gram Positive DNA Purification Kit (Biosearch tech-

nologies). Extracted genome was then sequenced using an Illumina NextSeq 500 sequencer. Raw

pair-end sequences were loaded into Geneious prime version 2025.1.2 for sequence trimming and

pairing with an insert size of 500 base pairs. Paired sequence list was then exported in fastq format

for analysis using breseq (Deatherage & Barrick, 2014).

Software and data availability

The fine-tuned segmentation models for raw and deconvolved images (for Omnipose, Cellpose-

SAM, microSAM, and Cellpose3), deconvolution prediction (FM2FM), deconvolution prediction

and SNR increase (FM2FM-HiSNR2), and fluorescence translation (FP2FM) models can be found

in the Zenodo (https://zenodo.org/records/18978187). Segmentation model training and bench-

marking data can be found at the BioImage Archive repository https://www.ebi.ac.uk/biostud-

ies/studies/S-BIAD2350. Deconvolution model training and benchmarking data can be found at

the BioImage Archive repository https://www.ebi.ac.uk/biostudies/studies/S-BIAD2353. Im-

ages of individual *E. coli* cells used for measurement transformation with their corresponding an-

notations and automatic segmentations be found at the BioImage Archive repository

https://www.ebi.ac.uk/biostudies/studies/S-BIAD2352. Segmentation of *B. subtilis* and *P. mega-*

terium strains (wild-type and induction constructs) be found at the BioImage Archive repository

https://www.ebi.ac.uk/biostudies/studies/S-BIAD2354.

The MEDUSSA GitHub repository contains all software-related material

(https://github.com/OReyesMatte/MEDUSSA). The "MEDUSSA" folder contains the func-

tions for size measurement (measure.py), Bayesian sampling, and measurement transformation

(both in transform.py). The "CARE" folder contains the environment file which was used for

training the deconvolution models, as well as the corresponding notebooks for preparing the data,

train the model, and test it. The "Segmentation" folder contains the environment files for the

different segmentation frameworks, as well as examples for both training commands and upload-

ing and running the models. The "Figures" folder contains notebooks to reproduce the plots

shown in the manuscript. The "Raw data" folder contains all data used to make the manuscript

plots. *P. megaterium* WH320 sequencing data, *ponA* protein sequence alignment, and results from

breseq (Deatherage & Barrick, 2014) are available in Zenodo (https://zenodo.org/up-

loads/17392073).

ACKNOWLEDGMENTS

We are grateful to Elitza Tocheva, Elizabeth Villa and Kanika Khanna for kindly sharing CryoEM

images of *B. subtilis* sporulating cells, Kaumudi H. Prabhakara for providing the *Saccharomyces cere-*

visiae and *Lactococcus lactis* strains used in Fig. 2d, and Felipe Sinhorini for helping with the acquisi-

tion of the BGSC strains. We thank members of the Evolutionary Cell Biology group, the Ojkic

lab, and the Microbial Population Biology department for continuous discussion and feedback.

We thank Elisa Brambilla and Michael Schwarz for helping with the acquisition of images with the

Zeiss Microscope. We also thank the members of the Scientific IT group at the Max Planck Insti-

tute for Evolutionary Biology for computational support. Work in JLG lab was supported by ERC

starting grant 853323. NO was supported by BBSRC grant (BB/Y009002/1).

AUTHOR CONTRIBUTIONS

Conceptualization, ORM, NO, JLG; Methodology, ORM, JLG; Software, ORM, CFG, BG; Vali-

dation, ORM; Formal analysis, ORM, CFG, BG; Investigation, ORM, NH; Resources, NH; Data

Curation, ORM; Writing - Original Draft, JLG; Writing - Review & Editing, All authors; Visuali-

zation, ORM, NO, JLG; Supervision, NO, JLG; Project administration, ORM, JLG; Funding ac-

quisition, JLG.

854
855 **COMPETING INTERESTS**

856
857 The authors declare no competing interests.

858
859

REFERENCES

Abril-Pla, O., Andreani, V., Carroll, C., Dong, L., Fonnesbeck, C. J., Kochurov, M., Kumar, R., Lao, J., Luhmann, C. C., Martin, O. A., Osthege, M., Vieira, R., Wiecki, T., & Zinkov, R. (2023). PyMC: a modern, and comprehensive probabilistic programming framework in Python. *PeerJ Computer Science*, *9*, e1516. https://doi.org/10.7717/peerj-cs.1516

Agard, D. A. (1984). Optical sectioning microscopy: cellular architecture in three dimensions. *Annual Review of Biophysics and Bioengineering*, *13*, 191-219. https://doi.org/10.1146/annurev.bb.13.060184.001203

Archit, A., Freckmann, L., Nair, S., Khalid, N., Hilt, P., Rajashekar, V., Freitag, M., Teuber, C., Spitzner, M., Tapia Contreras, C., Buckley, G., von Haaren, S., Gupta, S., Grade, M., Wirth, M., Schneider, G., Dengel, A., Ahmed, S., & Pape, C. (2025). Segment Anything for Microscopy. *Nat Methods*, *22*(3), 579-591. https://doi.org/10.1038/s41592-024-02580-4

Arya, G., Petronella, N., Crosthwait, J., Carrillo, C. D., & Shwed, P. S. (2014). Draft genome sequence of *Bacillus megaterium* type strain ATCC 14581. *Genome Announcements*, *2*(6), e01124-01114. https://doi.org/10.1128/genomeA.01124-14

Bakshi, S., Leoncini, E., Baker, C., Canas-Duarte, S. J., Okumus, B., & Paulsson, J. (2021). Tracking bacterial lineages in complex and dynamic environments with applications for growth control and persistence. *Nat Microbiol*, *6*(6), 783-791. https://doi.org/10.1038/s41564-021-00900-4

Blum, H. (1967). A transformation for extracting new descriptors of shape. In W. Walthen-Dunn (Ed.), *Models for the Perception of Speech and Visual Form* (pp. 362-380). MIT Press.

Borgefors, G. (1986). Distance transformations in digital images. *Computer Vision, Graphics, and Image Processing*, *34*(3), 344-371. https://doi.org/https://doi.org/10.1016/S0734-189X(86)80047-0

Caicedo, J. C., Roth, J., Goodman, A., Becker, T., Karhohs, K. W., Broisin, M., Molnar, C., McQuin, C., Singh, S., Theis, F. J., & Carpenter, A. E. (2019). Evaluation of deep learning strategies for nucleus segmentation in fluorescence images. *Cytometry*, *95*(9), 952-965. https://doi.org/10.1002/cyto.a.23863

Calvez, P., Jouhet, J., Vie, V., Durmort, C., & Zapun, A. (2019). Lipid phases and cell geometry during the cell cycle of *Streptococcus pneumoniae*. *Frontiers in Microbiology*, *10*, 351. https://doi.org/10.3389/fmicb.2019.00351

Campos, M., Surovtsev, Ivan V., Kato, S., Paintdakhi, A., Beltran, B., Ebmeier, Sarah E., & Jacobs-Wagner, C. (2014). A constant size extension drives bacterial cell size homeostasis. *Cell*, *159*(6), 1433-1446. https://doi.org/10.1016/j.cell.2014.11.022

Chien, A. C., Hill, N. S., & Levin, P. A. (2012). Cell size control in bacteria. *Current Biology*, *22*(9), R340-R349. https://doi.org/10.1016/j.cub.2012.02.032

Cox, R. S., 3rd, Dunlop, M. J., & Elowitz, M. B. (2010). A synthetic three-color scaffold for monitoring genetic regulation and noise. *Journal of Biological Engineering*, *4*, 10. https://doi.org/10.1186/1754-1611-4-10

Cutler, K. J., Stringer, C., Lo, T. W., Rappez, L., Stroustrup, N., Brook Peterson, S., Wiggins, P. A., & Mougous, J. D. (2022). Omnipose: a high-precision morphology-independent solution for bacterial cell segmentation. *Nature Methods*, *19*(11), 1438-1448. https://doi.org/10.1038/s41592-022-01639-4

Cylke, A., & Banerjee, S. (2023). Super-exponential growth and stochastic size dynamics in rod-like bacteria. *Biophys J*, *122*(7), 1254-1267. https://doi.org/10.1016/j.bpj.2023.02.015

Dalebroux, Z. D., Edrozo, M. B., Pfuetzner, R. A., Ressl, S., Kulasekara, B. R., Blanc, M. P., & Miller, S. I. (2015). Delivery of cardiolipins to the *Salmonella* outer membrane is necessary for survival within host tissues and virulence. *Cell Host Microbe*, *17*(4), 441-451. https://doi.org/10.1016/j.chom.2015.03.003

de Bary, A. (1884). *Vergleichende Morphologie und Biologie der Pilze, Mycetozoen und Bacterien.* Engelmann.

Deatherage, D. E., & Barrick, J. E. (2014). Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using breseq. *Methods in Molecular Biology, 1151*, 165-188. https://doi.org/10.1007/978-1-4939-0554-6_12

Delaby, M., Yang, L., Jacq, M., Gallagher, K. A., Kysela, D. T., Hughes, V., Pulido, F., Veyrier, F. J., VanNieuwenhze, M. S., & Brun, Y. V. (2025). Phenotypic plasticity in cell elongation among closely related bacterial species. *Nature Communications, 16*(1), 5099. https://doi.org/10.1038/s41467-025-60005-y

Dion, M. F., Kapoor, M., Sun, Y., Wilson, S., Ryan, J., Vigouroux, A., van Teeffelen, S., Oldenbourg, R., & Garner, E. C. (2019). *Bacillus subtilis* cell diameter is determined by the opposing actions of two distinct cell wall synthetic systems. *Nature Microbiology, 4*(8), 1294-1305. https://doi.org/10.1038/s41564-019-0439-0

Dosovitskiy, A., Beyer, L., Kolesnikov, A., Weissenborn, D., Zhai, X., Unterthiner, T., Dehghani, M., Minderer, M., Heigold, G., & Gelly, S. (2020). An image is worth 16x16 words: Transformers for image recognition at scale. *arXiv preprint arXiv:2010.11929*.

Ducret, A., Quardokus, E. M., & Brun, Y. V. (2016). MicrobeJ, a tool for high throughput bacterial cell detection and quantitative analysis. *Nature Microbiology, 1*(7), 16077. https://doi.org/10.1038/nmicrobiol.2016.77

Eppinger, M., Bunk, B., Johns, M. A., Edirisinghe, J. N., Kutumbaka, K. K., Koenig, S. S., Creasy, H. H., Rosovitz, M. J., Riley, D. R., Daugherty, S., Martin, M., Elbourne, L. D., Paulsen, I., Biedendieck, R., Braun, C., Grayburn, S., Dhingra, S., Lukyanchuk, V., Ball, B.,…Vary, P. S. (2011). Genome sequences of the biotechnologically important *Bacillus megaterium* strains QM B1551 and DSM319. *Journal of Bacteriology, 193*(16), 4199-4213. https://doi.org/10.1128/JB.00449-11

Falk, T., Mai, D., Bensch, R., Cicek, O., Abdulkadir, A., Marrakchi, Y., Bohm, A., Deubner, J., Jackel, Z., Seiwald, K., Dovzhenko, A., Tietz, O., Dal Bosco, C., Walsh, S., Saltukoglu, D., Tay, T. L., Prinz, M., Palme, K., Simons, M.,…Ronneberger, O. (2019). U-Net: deep learning for cell counting, detection, and morphometry. *Nature Methods, 16*(1), 67-70. https://doi.org/10.1038/s41592-018-0261-2

Gray, W. T., Govers, S. K., Xiang, Y., Parry, B. R., Campos, M., Kim, S., & Jacobs-Wagner, C. (2019). Nucleoid size scaling and intracellular organization of translation across bacteria. *Cell, 177*(6), 1632-1648.e1620. https://doi.org/10.1016/j.cell.2019.05.017

Greenspan, P., & Fowler, S. D. (1985). Spectrofluorometric studies of the lipid probe, nile red. *Journal of lipid research, 26*(7), 781-789. https://www.ncbi.nlm.nih.gov/pubmed/4031658

Gunawan, I., Marsh, R. J., Aggarwal, N., Meijering, E., Cox, S., Lock, J. G., & Culley, S. (2025). Image quality metrics fail to accurately represent biological information in fluorescence microscopy. *bioRxiv*, 2025.2008.2005.668508. https://doi.org/10.1101/2025.08.05.668508

Hardo, G., Li, R., & Bakshi, S. (2024). Quantitative microbiology with widefield microscopy: navigating optical artefacts for accurate interpretations. *Npj Imaging, 2*(1), 26. https://doi.org/10.1038/s44303-024-00024-4

Herman, P., Konopasek, I., Plasek, J., & Svobodova, J. (1994). Time-resolved polarized fluorescence studies of the temperature adaptation in *Bacillus subtilis* using DPH and TMA-DPH fluorescent probes. *Biochimica et biophysica acta, 1190*(1), 1-8. https://doi.org/10.1016/0005-2736(94)90028-0

Holt, J. H., Krieg, N. R., Sneath, P. H. a., Staley, J. T., & Williams, S. T. (1994). *Bergey's manual of determinative bacteriology ninth edition* (9th ed.). Lippincott Williams & Wilkins. https://doi.org/10.1016/j.ejpn.2008.10.006

Jiang, C., Caccamo, P. D., & Brun, Y. V. (2015). Mechanisms of bacterial morphogenesis: evolutionary cell biology approaches provide new insights. *BioEssays*, *37*(4), 413-425. https://doi.org/10.1002/bies.201400098

Juillot, D., Cornilleau, C., Deboosere, N., Billaudeau, C., Evouna-Mengue, P., Lejard, V., Brodin, P., Carballido-Lopez, R., & Chastanet, A. (2021). A high-content microscopy screening identifies new genes involved in cell width control in *Bacillus subtilis*. *mSystems*, *6*(6), e0101721. https://doi.org/10.1128/mSystems.01017-21

Kearns, D. B., & Losick, R. (2005). Cell population heterogeneity during growth of *Bacillus subtilis*. *Genes and Development*, *19*(24), 3083-3094. https://doi.org/10.1101/gad.1373905

Kellogg, D. R., & Levin, P. A. (2022). Nutrient availability as an arbiter of cell size. *Trends Cell Biol*, *32*(11), 908-919. https://doi.org/10.1016/j.tcb.2022.06.008

Khanna, K., Lopez-Garrido, J., Sugie, J., Pogliano, K., & Villa, E. (2021). Asymmetric localization of the cell division machinery during *Bacillus subtilis* sporulation. *eLife*, *10*, e62204. https://doi.org/10.7554/eLife.62204

Khanna, K., Lopez-Garrido, J., Zhao, Z., Watanabe, R., Yuan, Y., Sugie, J., Pogliano, K., & Villa, E. (2019). The molecular architecture of engulfment during *Bacillus subtilis* sporulation. *eLife*, *8*, e45257. https://doi.org/10.7554/eLife.45257

Kirillov, A., Mintun, E., Ravi, N., Mao, H., Rolland, C., Gustafson, L., Xiao, T., Whitehead, S., Berg, A. C., & Lo, W.-Y. (2023). Segment anything. Proceedings of the IEEE/CVF international conference on computer vision,

Kuru, E., Hughes, H. V., Brown, P. J., Hall, E., Tekkam, S., Cava, F., de Pedro, M. a., Brun, Y. V., & VanNieuwenhze, M. S. (2012). In Situ probing of newly synthesized peptidoglycan in live bacteria with fluorescent D-amino acids. *Angewandte Chemie (International ed. in English)*, *51*(50), 12519-12523. https://doi.org/10.1002/anie.201206749

Liu, L., Jiang, H., He, P., Chen, W., Liu, X., Gao, J., & Han, J. (2019). On the variance of the adaptive learning rate and beyond. *arXiv*, 1908.03265.

Logan, N. A., & Vos, P. D. (2015). *Bacillus*. In *Bergey's Manual of Systematics of Archaea and Bacteria* (pp. 1-163). https://doi.org/10.1002/9781118960608.gbm00530

López, A. C., & Alippi, A. M. (2009). Diversity of *Bacillus megaterium* isolates cultured from honeys. *LWT - Food Science and Technology*, *42*(1), 212-219. https://doi.org/10.1016/j.lwt.2008.05.001

Lopez-Garrido, J., Ojkic, N., Khanna, K., Wagner, F. R., Villa, E., Endres, R. G., & Pogliano, K. (2018). Chromosome translocation inflates *Bacillus* forespores and impacts cellular morphology. *Cell*, *172*(4), 758-770.e714. https://doi.org/10.1016/j.cell.2018.01.027

Loshchilov, I., & Hutter, F. (2017). Decoupled weight decay regularization. *arXiv preprint arXiv:1711.05101*.

Lucidi, M., Capecchi, G., Visaggio, D., Gasperi, T., Parisi, M., Cincotti, G., Rampioni, G., Visca, P., & Kolmakov, K. (2024). Expanding the microbiologist toolbox via new far-red-emitting dyes suitable for bacterial imaging. *Microbiology Spectrum*, *12*(1), e0369023. https://doi.org/10.1128/spectrum.03690-23

Lugagne, J. B., Lin, H., & Dunlop, M. J. (2020). DeLTA: Automated cell segmentation, tracking, and lineage reconstruction using deep learning. *PLoS Comput Biol*, *16*(4), e1007673. https://doi.org/10.1371/journal.pcbi.1007673

Lynch, M., Field, M. C., Goodson, H. V., Malik, H. S., Pereira-Leal, J. B., Roos, D. S., Turkewitz, A. P., & Sazer, S. (2014). Evolutionary cell biology: Two origins, one objective. *Proceedings of the National Academy of Sciences*, *111*(48), 16990-16994. https://doi.org/10.1073/pnas.1415861111

Ma, L., Conover, M., Lu, H., Parsek, M. R., Bayles, K., & Wozniak, D. J. (2009). Assembly and development of the *Pseudomonas aeruginosa* biofilm matrix. *PLoS Pathogens*, *5*(3), e1000354. https://doi.org/10.1371/journal.ppat.1000354

Maier-Hein, L., Reinke, A., Godau, P., Tizabi, M. D., Buettner, F., Christodoulou, E., Glocker, B., Isensee, F., Kleesiek, J., Kozubek, M., Reyes, M., Riegler, M. A., Wiesenfarth, M., Kavur, A. E., Sudre, C. H., Baumgartner, M., Eisenmann, M., Heckmann-Notzel, D., Radsch, T.,…Jager, P. F. (2024). Metrics reloaded: recommendations for image analysis validation. *Nature Methods*, *21*(2), 195-212. https://doi.org/10.1038/s41592-023-02151-z

Maloney, E., Madiraju, M., & Rajagopalan, M. (2009). Overproduction and localization of *Mycobacterium tuberculosis* ParA and ParB proteins. *Tuberculosis (Edinb)*, *89 Suppl 1*(Suppl 1), S65-69. https://doi.org/10.1016/S1472-9792(09)70015-0

Margulis, L., Jorgensen, J. Z., Dolan, S., Kolchinsky, R., Rainey, F. A., & Lo, S.-C. (1998). The *Arthromitus* stage of *Bacillus cereus:* Intestinal symbionts of animals. *Proceedings of the National Academy of Sciences*, *95*, 1236-1241.

Marks, M., Israel, U., Dilip, R., Li, Q., Yu, C., Laubscher, E., Iqbal, A., Pradhan, E., Ates, A., Abt, M., Brown, C., Pao, E., Li, S., Pearson-Goulart, A., Perona, P., Gkioxari, G., Barnowski, R., Yue, Y., & Van Valen, D. (2025). CellSAM: a foundation model for cell segmentation. *Nat Methods*, *22*(12), 2585-2593. https://doi.org/10.1038/s41592-025-02879-w

Monteiro, J. M., Fernandes, P. B., Vaz, F., Pereira, A. R., Tavares, A. C., Ferreira, M. T., Pereira, P. M., Veiga, H., Kuru, E., Vannieuwenhze, M. S., Brun, Y. V., Filipe, S. R., & Pinho, M. G. (2015). Cell shape dynamics during the staphylococcal cell cycle. *Nature Communications*, *6*, 1-12. https://doi.org/10.1038/ncomms9055

Murumkar, D. R., Borkar, S. G., & Chimote, V. P. (2012). Diversity for cell morphology, in-vitro phosphate solubilization and DNA profile of *Bacillus megaterium* isolates present in soils of Maharashtra. *World of Applied Sciences Journal*, *17*(6), 776-785.

Nordholt, N., van Heerden, J. H., & Bruggeman, F. J. (2020). Biphasic cell-size and growth-rate homeostasis by single *Bacillus subtilis* cells. *Current Biology*, *30*(12), 2238-2247 e2235. https://doi.org/10.1016/j.cub.2020.04.030

O'Connor, O. M., Alnahhas, R. N., Lugagne, J. B., & Dunlop, M. J. (2022). DeLTA 2.0: A deep learning pipeline for quantifying single-cell spatial and temporal dynamics. *PLoS Computational Biology*, *18*(1), e1009797. https://doi.org/10.1371/journal.pcbi.1009797

O'Hara, B. J., Barth, Z. K., McKitterick, A. C., & Seed, K. D. (2017). A highly specific phage defense system is a conserved feature of the *Vibrio cholerae* mobilome. *PLoS Genetics*, *13*(6), e1006838. https://doi.org/10.1371/journal.pgen.1006838

Ojkic, N., & Banerjee, S. (2021). Bacterial cell shape control by nutrient-dependent synthesis of cell division inhibitors. *Biophysical Journal*, *120*(11), 2079-2084. https://doi.org/10.1016/j.bpj.2021.04.001

Ojkic, N., López-Garrido, J., Pogliano, K., & Endres, R. G. (2014). Bistable forespore engulfment in *Bacillus subtilis* by a zipper mechanism in absence of the cell wall. *PLoS Computational Biology*, *10*(10), e1003912. https://doi.org/10.1371/journal.pcbi.1003912

Ojkic, N., López-Garrido, J., Pogliano, K., & Endres, R. G. (2016). Cell-wall remodeling drives engulfment during *Bacillus subtilis* sporulation. *eLife*, *5*, e18657. https://doi.org/10.7554/eLife.18657

Ojkic, N., Serbanescu, D., & Banerjee, S. (2019). Surface-to-volume scaling and aspect ratio preservation in rod-shaped bacteria. *eLife*, *8*, e47033. https://doi.org/10.7554/eLife.47033

Ojkic, N., Serbanescu, D., & Banerjee, S. (2022). Antibiotic resistance via bacterial cell shape-shifting. *mBio*, *13*(3), e0065922. https://doi.org/10.1128/mbio.00659-22

Ollion, J., Maliet, M., Giuglaris, C., Vacher, É., & Deforet, M. (2024). DiSTNet2D: Leveraging long-range temporal information for efficient segmentation and tracking. *PRX Life*, *2*(2), 023004. https://doi.org/10.1103/PRXLife.2.023004

Pachitariu, M., Rariden, M., & Stringer, C. (2025). Cellpose-SAM: superhuman generalization for cellular segmentation. *bioRxiv*, 2025.2004.2028.651001. https://doi.org/10.1101/2025.04.28.651001

Paintdakhi, A., Parry, B., Campos, M., Irnov, I., Elf, J., Surovtsev, I., & Jacobs-Wagner, C. (2016). Oufti: an integrated software package for high-accuracy, high-throughput quantitative microscopy analysis. *Molecular Microbiology*, *99*(4), 767-777. https://doi.org/10.1111/mmi.13264

Panigrahi, S., Murat, D., Le Gall, A., Martineau, E., Goldlust, K., Fiche, J. B., Rombouts, S., Nollmann, M., Espinosa, L., & Mignot, T. (2021). Misic, a general deep learning-based method for the high-throughput cell segmentation of complex bacterial communities. *eLife*, *10*, e65151. https://doi.org/10.7554/eLife.65151

Pogliano, J., Osborne, N., Sharp, M. D., Abanes-De Mello, a., Perez, a., Sun, Y. L., & Pogliano, K. (1999). A vital stain for studying membrane dynamics in bacteria: a novel mechanism controlling septation during *Bacillus subtilis* sporulation. *Molecular Microbiology*, *31*(4), 1149-1159.

Popham, D. L., & Setlow, P. (1996). Phenotypes of *Bacillus subtilis* mutants lacking multiple class A high-molecular-weight penicillin-binding proteins. *Journal of Bacteriology*, *178*(7), 2079-2085.

Riley, E. P., Trinquier, A., Reilly, M. L., Durchon, M., Perera, V. R., Pogliano, K., & Lopez-Garrido, J. (2018). Spatiotemporally regulated proteolysis to dissect the role of vegetative proteins during *Bacillus subtilis* sporulation: cell-specific requirement of σ^H and σ^A. *Molecular Microbiology*, *108*(1), 45-62. https://doi.org/10.1111/mmi.13916

Ronneberger, O., Fischer, P., & Brox, T. (2015). U-Net: Convolutional networks for biomedical image segmentation. In N. Navab, J. Hornegger, W. M. Wells, & A. F. Frangi, *Medical Image Computing and Computer-Assisted Intervention – MICCAI 2015* Cham.

Rygus, T., Scheler, A., Allmansberger, R., & Hillen, W. (1991). Molecular cloning, structure, promoters and regulatory elements for transcription of the *Bacillus megaterium* encoded regulon for xylose utilization. *Archives of Microbiology*, *155*(6), 526-534. https://doi.org/10.1007/BF00245346

Sauls, J. T., Cox, S. E., Do, Q., Castillo, V., Ghulam-Jelani, Z., & Jun, S. (2019). Control of *Bacillus subtilis* replication initiation during physiological transitions and perturbations. *mBio*, *10*(6), 10.1128/mbio.02205-02219. https://doi.org/10.1128/mBio.02205-19

Scalettar, B. A., Swedlow, J. R., Sedat, J. W., & Agard, D. A. (1996). Dispersion, aberration and deconvolution in multi-wavelength fluorescence images. *Journal of Microscopy*, *182*(1), 50-60. https://doi.org/10.1046/j.1365-2818.1996.122402.x

Schaechter, M., MaalOe, O., & Kjeldgaard, N. O. (1958). Dependency on Medium and Temperature of Cell Size and Chemical Composition during Balanced Growth of Salmonella typhimurium. *Journal of General Microbiology*, *19*(3), 592-606. https://doi.org/10.1099/00221287-19-3-592

Schavemaker, P. E., & Munoz-Gomez, S. A. (2022). The role of mitochondrial energetics in the origin and diversification of eukaryotes. *Nature Ecology and Evolution*, *6*(9), 1307-1317. https://doi.org/10.1038/s41559-022-01833-9

Schmidt, U., Weigert, M., Broaddus, C., & Myers, G. (2018). Cell detection with star-convex polygons. In A. F. Frangi, J. A. Schnabel, C. Davatzikos, C. Alberola-López, & G. Fichtinger, *Medical Image Computing and Computer Assisted Intervention – MICCAI 2018* Cham.

Schober, I., Koblitz, J., Sarda Carbasse, J., Ebeling, C., Schmidt, M. L., Podstawka, A., Gupta, R., Ilangovan, V., Chamanara, J., Overmann, J., & Reimer, L. C. (2025). BacDive in 2025: the core database for prokaryotic strain data. *Nucleic Acids Research*, *53*(D1), D748-D756. https://doi.org/10.1093/nar/gkae959

Sega, G. A. (1984). A review of the genetic effects of ethyl methanesulfonate. *Mutation Research*, *134*(2-3), 113-142. https://doi.org/10.1016/0165-1110(84)90007-1

Sharp, M. D., & Pogliano, K. (1999). An in vivo membrane fusion assay implicates SpoIIIE in the final stages of engulfment during *Bacillus subtilis* sporulation. *Proceedings of the National Academy of Sciences of the United States of America*, *96*(25), 14553-14558.

Shin, J. Y., Lopez-Garrido, J., Lee, S.-H. H., Diaz-Celis, C., Fleming, T., Bustamante, C., & Pogliano, K. (2015). Visualization and functional dissection of coaxial paired SpoIIIE channels across the sporulation septum. *eLife*, *4*(MAY), e06474. https://doi.org/10.7554/eLife.06474

Smith, M. B., Li, H., Shen, T., Huang, X., Yusuf, E., & Vavylonis, D. (2010). Segmentation and tracking of cytoskeletal filaments using open active contours. *Cytoskeleton*, *67*(11), 693-705. https://doi.org/10.1002/cm.20481

Spahn, C., Gomez-de-Mariscal, E., Laine, R. F., Pereira, P. M., von Chamier, L., Conduit, M., Pinho, M. G., Jacquemet, G., Holden, S., Heilemann, M., & Henriques, R. (2022). DeepBacs for multi-task bacterial image analysis using open-source deep learning approaches. *Communications Biology*, *5*(1), 688. https://doi.org/10.1038/s42003-022-03634-z

Specht, M., Dempwolff, F., Schatzle, S., Thomann, R., & Waidner, B. (2013). Localization of FtsZ in *Helicobacter pylori* and consequences for cell division. *Journal of Bacteriology*, *195*(7), 1411-1420. https://doi.org/10.1128/JB.01490-12

Stahl, U., & Esser, K. (1983). Plasmid heterogeneity in various strains of *Bacillus megaterium*. *European journal of applied microbiology and biotechnology*, *17*(4), 248-251. https://doi.org/10.1007/BF00510424

Strahl, H., Burmann, F., & Hamoen, L. W. (2014). The actin homologue MreB organizes the bacterial cell membrane. *Nat Commun*, *5*, 3442. https://doi.org/10.1038/ncomms4442

Stringer, C., & Pachitariu, M. (2025). Cellpose3: one-click image restoration for improved cellular segmentation. *Nature Methods*, *22*(3), 592-599. https://doi.org/10.1038/s41592-025-02595-5

Stringer, C., Wang, T., Michaelos, M., & Pachitariu, M. (2021). Cellpose: a generalist algorithm for cellular segmentation. *Nature Methods*, *18*(1), 100-106. https://doi.org/10.1038/s41592-020-01018-x

Studer, P., Staubli, T., Wieser, N., Wolf, P., Schuppler, M., & Loessner, M. J. (2016). Proliferation of *Listeria monocytogenes* L-form cells by formation of internal and external vesicles. *Nature Communications*, *7*, 13631. https://doi.org/10.1038/ncomms13631

Stylianidou, S., Brennan, C., Nissen, S. B., Kuwada, N. J., & Wiggins, P. A. (2016). SuperSegger: robust image segmentation, analysis and lineage tracking of bacterial cells. *Molecular Microbiology*, *102*(4), 690-700. https://doi.org/10.1111/mmi.13486

Sun, Q., & Margolin, W. (1998). FtsZ dynamics during the division cycle of live *Escherichia coli* cells. *Journal of Bacteriology*, *180*(8), 2050-2056. https://doi.org/10.1128/JB.180.8.2050-2056.1998

Sutskever, I., Martens, J., Dahl, G., & Hinton, G. (2013). *On the importance of initialization and momentum in deep learning* Proceedings of the 30th International Conference on Machine Learning, Proceedings of Machine Learning Research.

Syvertsson, S., Vischer, N. O., Gao, Y., & Hamoen, L. W. (2016). When phase contrast fails: ChainTracer and NucTracer, two ImageJ methods for semi-automated single cell analysis using membrane or DNA staining. *PLoS ONE*, *11*(3), e0151267. https://doi.org/10.1371/journal.pone.0151267

Taha, A. A., & Hanbury, A. (2015). Metrics for evaluating 3D medical image segmentation: analysis, selection, and tool. *BMC Medical Imaging*, *15*, 29. https://doi.org/10.1186/s12880-015-0068-x

Taheri-Araghi, S., Bradde, S., Sauls, J. T., Hill, N. S., Levin, P. A., Paulsson, J., Vergassola, M., &
Jun, S. (2015). Cell-size control and homeostasis in bacteria. *Current Biology*, *25*(3), 385-
391. https://doi.org/10.1016/j.cub.2014.12.009
Tocheva, E. I., López-Garrido, J., Hughes, H. V., Fredlund, J., Kuru, E., Vannieuwenhze, M. S.,
Brun, Y. V., Pogliano, K., & Jensen, G. J. (2013). Peptidoglycan transformations during
Bacillus subtilis sporulation. *Molecular Microbiology*, *88*(4), 673-686.
https://doi.org/10.1111/mmi.12201
Tsuge, Y., Ogino, H., Teramoto, H., Inui, M., & Yukawa, H. (2008). Deletion of cgR_1596 and
cgR_2070, encoding NlpC/P60 proteins, causes a defect in cell separation in
Corynebacterium glutamicum R. *Journal of Bacteriology*, *190*(24), 8204-8214.
https://doi.org/10.1128/JB.00752-08
Ursell, T., Lee, T. K., Shiomi, D., Shi, H., Tropini, C., Monds, R. D., Colavin, A., Billings, G.,
Bhaya-Grossman, I., Broxton, M., Huang, B. E., Niki, H., & Huang, K. C. (2017). Rapid,
precise quantification of bacterial cellular dimensions across a genomic-scale knockout
library. *BMC Biology*, *15*(1), 17. https://doi.org/10.1186/s12915-017-0348-8
van der Walt, S., Schonberger, J. L., Nunez-Iglesias, J., Boulogne, F., Warner, J. D., Yager, N.,
Gouillart, E., Yu, T., & scikit-image, c. (2014). scikit-image: image processing in Python.
PeerJ, *2*, e453. https://doi.org/10.7717/peerj.453
Van Valen, D. A., Kudo, T., Lane, K. M., Macklin, D. N., Quach, N. T., DeFelice, M. M.,
Maayan, I., Tanouchi, Y., Ashley, E. A., & Covert, M. W. (2016). Deep learning
automates the quantitative analysis of individual cells in live-cell imaging experiments.
PLoS Computational Biology, *12*(11), e1005177.
https://doi.org/10.1371/journal.pcbi.1005177
Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,
Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M.,
Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson,
E.,…SciPy, C. (2020). SciPy 1.0: fundamental algorithms for scientific computing in
Python. *Nat Methods*, *17*(3), 261-272. https://doi.org/10.1038/s41592-019-0686-2
Weigert, M., Schmidt, U., Boothe, T., Muller, A., Dibrov, A., Jain, A., Wilhelm, B., Schmidt, D.,
Broaddus, C., Culley, S., Rocha-Martins, M., Segovia-Miranda, F., Norden, C.,
Henriques, R., Zerial, M., Solimena, M., Rink, J., Tomancak, P., Royer, L.,…Myers, E.
W. (2018). Content-aware image restoration: pushing the limits of fluorescence
microscopy. *Nature Methods*, *15*(12), 1090-1097. https://doi.org/10.1038/s41592-018-
0216-7
Yeh, Y. C., Comolli, L. R., Downing, K. H., Shapiro, L., & McAdams, H. H. (2010). The
Caulobacter Tol-Pal complex is essential for outer membrane integrity and the positioning
of a polar localization factor. *Journal of Bacteriology*, *192*(19), 4847-4858.
https://doi.org/10.1128/JB.00607-10
Zhang, T. Y., & Suen, C. Y. (1984). A fast parallel algorithm for thinning digital patterns.
Communications of the ACM, *27*(3), 236–239. https://doi.org/10.1145/357994.358023
Zou, X., Yang, J., Zhang, H., Li, F., Li, L., Wang, J., Wang, L., Gao, J., & Lee, Y. J. (2023).
Segment everything everywhere all at once. *Advances in neural information processing systems*,
36, 19769-19782.

1202