1
12    <u>A20 restriction of nitric oxide production restores macrophage bioenergetic balance</u>
23    Mei-po Chan[1], Rommel Advincula[1], Philip Achacoso[1], Elizabeth A. Grossman[2], Daniel K.
24    Nomura[2], Barbara A. Malynn[1], and Averil Ma[1*]
25
26
27
28
29    [1]Department of Medicine, University of California, San Francisco, San Francisco, CA. 94143

31    [2]Departments of Chemistry, Molecular and Cell Biology, and Nutritional Sciences and
32    Toxicology, University of California, Berkeley, Berkeley, CA 94720

34    *Correspondence to: Averil Ma, MD, 513 Parnassus Ave, S-1032C, University of California, San
35    Francisco, San Francisco, CA. 94143. Email: Averil.ma@ucsf.edu
36
37
38
39
40    Short running title: A20 preserves mitochondrial function

**Abstract**

Macrophages play critical roles in regulating host responses to microbial pathogens and other forms of tissue stress and can acquire pro-inflammatory or tissue reparative phenotypes. A20, or TNFAIP3, is a potent regulator of innate immune cell functions and is extensively linked to human inflammatory and autoimmune diseases. We now find that A20 is a powerful regulator of both glycolytic and mitochondrial respiration in macrophages. Differentiated macrophages that are acutely rendered A20 deficient exhibit increased glycolytic activity and markedly decreased mitochondrial respiration after LPS stimulation. These cells are unable to repolarize from an M1-like to M2-like phenotype. Compromised mitochondrial oxygen consumption in A20 deficient macrophages is caused by increased nitric oxide production. Inhibition or genetic ablation of inducible nitric oxide synthase (iNOS) restores mitochondrial oxidative phosphorylation and lactate production in these cells. These metabolic perturbations occur independently of exaggerated cytokine production and despite robust production of IL-10. Therefore, A20 prevents Warburg-like aerobic glycolysis and restores macrophage homeostasis.

## Introduction

Activation of immune cells is associated with a rapid burst in glycolytic activity along with mitochondrial remodeling (Buck et al., 2016). Several days post-activation, T cells return to more basal states associated with lower glycolysis and persistently increased mitochondrial respiration (Makowski et al., 2020). This latter transition may be important for maintaining an epigenetic state that retains physiological memory of past cellular exposures. It may also facilitate feedback mechanisms that control the duration and longitudinal character of immune responses.

Though less studied than T cells, macrophages appear to undergo similar metabolic changes during activation with ligands such as LPS (O'Neill and Pearce, 2016; Galván-Peña and O'Neill, 2014). Initial LPS stimulation of these cells triggers a glycolytic surge that provides energy for cytokine transcription, translation, and secretion. Subsequent subsiding of glycolysis with a relative shift back to mitochondrial oxidative phosphorylation coincides with transitions to relative LPS tolerance or to cells that express anti-inflammatory molecules associated with wound repair and tissue restoration, e.g., M2 macrophages. Importantly, while selected metabolic enzymes have been implicated in these processes (Mills et al., 2016), the cell biological mechanisms that couple immune cell activation with metabolic shifts are incompletely understood.

The ubiquitin editing protein A20, also known as TNF-α-induced protein 3 (TNFAIP3), is a

susceptibility gene for multiple diseases including rheumatoid arthritis, Crohn's disease,

systemic lupus erythematosus, psoriasis, multiple sclerosis, scleroderma, and asthma (Ma and

Malynn, 2012). In addition, patients with mono-allelic loss-of-function mutations of A20

develop mucosal and skin ulcers resembling Behçet's disease as well as a growing number of

other autoinflammatory and autoimmune disorders (Zhou et al., 2016; Ohnishi et al., 2017;

Takagi et al., 2017; Aeschlimann et al., 2018). In parallel, A20 deficiency or mutations in mice

leads to the development of spontaneous inflammation and autoimmune phenotypes (Boone

et al., 2004; Lee et al., 2000; Tavares et al., 2010; Razani et al., 2020). Mice lacking A20 in

selected innate immune cell types exhibit phenotypes that highlight the importance of A20

functions in macrophages, dendritic cells, and innate lymphoid cells (Hammer et al., 2011; Kool

et al., 2011; Matmati et al., 2011; Schneider et al., 2018). A20 also regulates myelopoiesis, and

perturbed myeloid differentiation may contribute to altered functions of mature cells (Muto et

al., 2020). We have now explored the role of A20 in regulating metabolic transitions of

macrophages, using a tamoxifen inducible system that allows myeloid cells to undergo normal

hematopoiesis prior to A20 deletion. We find that A20 is a potent regulator of both glycolytic

and mitochondrial activity in activated macrophages.

**Results and Discussion**

<u>Induced deletion of A20 in macrophages promotes aerobic glycolysis</u>

To understand how A20 regulates macrophage functions independently from developmental

events, we interbred A20$^{\text{Flox/Flox}}$ mice with mice bearing the ROSA/ER-Cre gene (Tavares et al.,

2010). We generated bone marrow derived macrophages (BMDMs) from the resulting

A20$^{Flox/Flox}$ ROSA/ER-Cre+, A20$^{Flox/+}$ ROSA/ER-Cre-, and A20$^{Flox/Flox}$ ROSA/ER-Cre- mice, and

established conditions for deleting A20 from A20$^{Flox/Flox}$ ROSA/ER-Cre+ bone marrow cells *in*

*vitro*. We call these cells A20$^{tiKO}$ BMDMs. While Cre negative BMDMs induced expression of A20

protein within one hour after LPS stimulation, which gradually returned to basal levels over 16

hours, stimulation of A20$^{tiKO}$ BMDMs expressed no detectable protein (**Fig. 1A**). LPS stimulation

of A20$^{tiKO}$ cells caused increased production of pro-inflammatory cytokines such as TNF and IL-6

when compared with control BMDMs, consistent with prior findings with A20$^{-/-}$ BMDMs (**Fig.**

**1B**) (Boone et al., 2004; Turer et al., 2008). As macrophages develop tolerance to LPS and are

thought to restore homeostatic functions after 16-48 hours (Seeley and Ghosh, 2017; Kotwal

and Chien, 2017), we studied macrophages 24 hours after LPS stimulation. Remarkably, LPS

stimulation of A20$^{tiKO}$ BMDMs led to visibly changed media when compared to control cells (**Fig.**

**1C**). As culture medium acidification is most commonly caused by a change in lactate

production, lactate levels in the culture supernatants were measured. A20$^{tiKO}$ BMDMs produced

markedly more lactate than control cells at 18 and 24 hours after LPS stimulation (**Fig. 1D**). This

exaggerated lactate production was not observed after stimulation with other TLR ligands (**Fig.**

**1E**). FACS analyses revealed that cell survival was similar in A20$^{tiKO}$ and WT BMDMs at 24 hours

post LPS stimulation (**Fig. 1F**). As lactate is an end product of glycolysis, we measured glycolytic

activity of these cells via Seahorse extracellular proton flux assay. Prior to LPS stimulation, both

A20$^{WT}$ and A20$^{tiKO}$ BMDMs exhibited a similarly modest extracellular acidification rate (ECAR)

(**Fig. 1G**). Consistent with prior studies (Galván-Peña and O'Neill, 2014), LPS caused increased

glycolysis in normal macrophages (**Fig. 1G**). Importantly, after LPS stimulation, the ECAR of

A20$^{tiKO}$ cells was markedly higher than in LPS stimulated WT cells (**Fig. 1G**). After

rotenone/antimycin A mediated inhibition of mitochondrial respiration, both A20$^{tiKO}$ and WT

cells enhanced proton efflux, with A20$^{tiKO}$ cells showing greater glycolytic capacity (**Fig. 1G**). The

differential ECAR between maximal glycolytic capacity and basal rate was slightly greater in WT

cells, suggesting diminished glycolytic reserve in A20$^{tiKO}$ cells (**Fig. 1G**).

<u>Figure 1: Inducible deletion of A20 from BMDMs leads to enhanced expression of glycolytic enzymes and glycolytic</u>

<u>activity after LPS stimulation.</u> **(A)** Immunoblot of A20 expression in A20$^{Flox/Flox}$ Rosa/ER-Cre ("ER-Cre +" or A20$^{tiKO}$)

and A20$^{Flox/Flox}$ ("ER-Cre –") BMDMs after 4-OH-T mediated deletion of A20 followed by LPS stimulation for the

indicated number of hours. Data are representative of three independent experiments. **(B)** ELISA of TNF and IL-6

secretion from A20$^{tiKO}$ (red columns) and WT (black columns) BMDMs after LPS. Mean values ± SD are shown. ***

indicates $p<0.001$ by Student's T test. Data are representative of three independent experiments. **(C)** Gross images

of culture media from A20$^{tiKO}$ and WT BMDMs with or without LPS stimulation. Data are representative of at least

three experiments. **(D)** Lactate secretion from BMDMs of indicated genotypes displayed as fold induction of LPS

stimulated cells at indicated time points after LPS relative to unstimulated cells. Pooled results from six

independent experiments are shown. Mean values ± SD are shown. ** indicates $p<0.01$ and *** indicates $p<0.001$

by one way ANOVA. **(E)** Lactate secretion from A20$^{tiKO}$ (red columns) and WT (black columns) BMDMs after

stimulation with indicated TLR ligands. Mean values ± SD are shown. ** indicates $p<0.01$ by Student's T test. Data

are representative of two independent experiments. **(F)** Flow cytometric quantification of BMDM cell viability via

live-dead staining. Data are representative of two independent experiments. Mean values ± SD are shown. **(G)**

Extracellular acidification rate (ECAR) of A20$^{tiKO}$ and WT BMDMs with and without LPS stimulation analyzed by

Seahorse proton flux assay. Rotenone/antimycin A and 2-deoxyglucose (2-DG) added at indicated times. Mean

values ± SD are shown. Data are representative of three independent experiments. **(H)** Real-time quantitative PCR

analyses of *glut1, hk2, ldha, gpi1, pkm2* and *hif1a* mRNA expression relative to *actin* mRNA in indicated genotypes

of cells at indicated times after LPS stimulation. Mean values ± SD are shown. Data are representative of two

independent experiments. All TLR stimulations were conducted for 24 hours unless otherwise stated.

The glycolytic surge in activated lymphocytes is supported by multiple molecular adaptations.

Glut1 is a glucose transporter which is transcriptionally induced in CD4[+] T cells, and is essential

for their activation and differentiation into effector cells (Macintyre et al., 2014). Hexokinase2

phosphorylates intracellular glucose, preventing its transport out of the cell, and represents the

first step of glycolysis. Lactate dehydrogenase (LDH) converts pyruvate to lactate and

regenerates nicotinamide adenine dinucleotide (NAD) to support glycolysis. Glucose 6

phosphate isomerase (GPI) converts glucose-6-phosphate to fructose-6-phosphate and

represents the second step of glycolysis. Pyruvate kinase M2 (PKM2) converts

phosphoenolpyruvate to pyruvate. HIF1$\alpha$ is a transcription factor that drives expression of

several glycolytic enzymes. After LPS stimulation, expression of mRNAs for all these molecules

was markedly increased in A20 deficient macrophages when compared to control cells (**Fig. 1H**).

Hence, A20 restrains multiple facets of glycolysis that collaborate to limit the glycolytic surge

accompanying LPS stimulation. This increased glycolytic activity may support enhanced

production and secretion of cytokines by A20 deficient macrophages.

<u>Induced deletion of A20 in macrophages suppresses mitochondrial respiration and prevents M2</u>

<u>repolarization after LPS</u>

After LPS stimulation, macrophages can assume tissue reparative functions, reduction of

glycolysis, and a return to predominantly mitochondrial oxidative phosphorylation for ATP

production. We thus assayed mitochondrial respiration by measuring the oxygen consumption

rate (OCR) of A20$^{tiKO}$ and control BMDMs via Seahorse assays. Consistent with prior work, the

OCR of WT BMDMs increased after LPS stimulation (**Fig. 2A**). By contrast, A20$^{tiKO}$ BMDMs, which

contained mildly elevated levels of OCR when compared to unstimulated WT cells, exhibited

markedly decreased OCR after LPS stimulation (**Fig. 2A**, see basal OCR). Oligomycin inhibits ATP

synthase and the OCR in oligomycin treated cells reflects the amount of ATP generated by the

mitochondrial electron transport chain. A20$^{tiKO}$ BMDMs exhibited negligible decrements of OCR

after oligomycin (**Fig. 2A**, see ATP production). FCCP uncouples the proton gradient generated

by the electron transport chain, unveiling the maximal oxygen consumption potential of

mitochondria in normal cells. While WT cells and unstimulated A20$^{tiKO}$ BMDMs increased their

OCR to maximal levels, LPS stimulated A20$^{tiKO}$ BMDMs completely failed to increase their OCR in

response to FCCP (**Fig. 2A**, see maximal). Therefore, LPS stimulation cripples the ability of

mitochondrial electron transport complexes in A20$^{tiKO}$ BMDMs to consume oxygen even when

these complexes are uncoupled from their electron transport functions. The difference

between FCCP induced maximal OCR and basal OCR, termed the spare respiratory capacity

(SRC), reflects the cell's ability to marshal reserves for energy production and is an indicator of

cellular fitness (**Fig. 2A**, see SRC). In this regard, A20$^{tiKO}$ BMDMs displayed negligible SRC. Finally,

inhibition of complexes I and III with rotenone and antimycin A blocks mitochondrial respiration

and unveils oxygen consumption from non-mitochondrial sources. These treatments revealed

modestly increased OCR in A20$^{tiKO}$ BMDMs, consistent with the increased glycolysis observed in

these cells. These respiratory defects in A20$^{tiKO}$ BMDMs were triggered by LPS stimulation but

not by other TLR ligands such as Pam$_3$CSK$_4$ and polyI:C (**Figs. 2B, 2C**). Hence, while A20$^{tiKO}$

BMDMs conduct increased glycolysis after LPS stimulation, these cells perform markedly less

mitochondrial respiration, thereby exhibiting a profound bioenergetic bias toward aerobic

glycolysis. This dramatic metabolic shift from oxidative phosphorylation to aerobic glycolysis in

A20 deficient macrophages constitutes a potent version of the Warburg effect initially

described in cancer cells. Importantly, as A20 is a tumor suppressor (Schmitz et al., 2009; Kato

et al., 2009; Compagno et al., 2009), the loss of this protein could drive aerobic glycolysis in

some cancer cells as well.

<u>Figure 2:  LPS activation cripples mitochondrial respiration and prevents M2 repolarization in A20$^{tiKO}$ BMDMs.</u> **(A)**

Oxygen consumption rate (OCR) measured by Seahorse flux assay of LPS stimulated A20 deficient (red circles) and

WT (black circles) BMDMs. Open circles represent unstimulated cells; Closed circles represent LPS stimulated cells.

Mitochondrial stress tests with oligomycin (Oligo), FCCP, and rotenone/antimycin A injected at indicated times

(arrows). Calculated levels of basal and maximal mitochondrial respiration, spare respiratory capacity (SRC) and

ATP production shown at right. Mean values ± SD are shown. *** indicates p<0.001 by unpaired two-tailed T test.

Data are representative of at least five independent experiments. **(B and C)** Mitochondrial stress tests of A20$^{WT}$

and A20$^{tiKO}$ BMDMs after stimulation with Pam$_3$CSK$_4$ **(B)** or polyI:C **(C)**. Mean values ± SD are shown. Data are

representative of three independent experiments. **(D)** Schematic representation of experimental design of M2

repolarization assay. On day 7 of BMDM differentiation, cells were treated with LPS for 24 hours, washed and

stimulated with IL-4 or medium for another 24 hours. Flow cytometric measurement of M2 surface markers were

assayed on day 9. **(E)** Representative flow cytometric analyses of CD301 and CD206 cell surface expression (M2

markers) on A20$^{tiKO}$ and WT BMDMs after IL-4 mediated repolarization as described in (D). Mean values ± SD are

shown. Data are representative of two independent experiments. **(F)** Oil red O staining of indicated genotypes of

BMDMs after stimulation with LPS. Note lipid droplets inside A20$^{tiKO}$ BMDMs. Scale bars represent 2 μm. Data are

representative of three independent experiments.

The major source of ATP in M2 macrophages derives from oxidative phosphorylation by

mitochondria (Jha et al., 2015; Vats et al., 2006). Hence, A20's preservation of macrophage

mitochondrial function might support M2 repolarization after LPS stimulation. We tested the

ability of LPS stimulated A20$^{tiKO}$ BMDMs to repolarize to an M2 phenotype after IL-4

stimulation. We stimulated A20$^{tiKO}$ and A20$^{WT}$ BMDMs with LPS for 24 hours, treated these cells

with IL-4 for another 24 hours, and finally analyzed the expression of M2 markers by FACS (**Fig.**

**2D**). A20$^{tiKO}$ BMDMs were largely unable to express the M2 markers CD301 and CD206 after

repolarization with IL-4 (**Fig. 2E**). Instead, A20$^{tiKO}$ BMDMs were strongly stained with Oil Red O,

indicating enhanced lipid droplet formation and accumulation after LPS stimulation (**Fig. 2F**), a

hallmark of inflammatory M1 macrophages (Rosas-Ballina et al., 2020; Wen et al., 2011). Such

lipid droplets have been described as sites of eicosanoid synthesis, enzyme localization and

cytokine storage in inflammatory leukocytes (Bozza and Viola, 2010; Jarc and Petan, 2020).

Hence, A20 is required for the ability of activated macrophages to express type 2 markers and

potentially evolve into cells supporting tissue repair. The persistence of M1 associated proteins

likely contributes to the ability of these cells to drive inflammatory diseases.

<u>Mitochondria in A20 deficient macrophages exhibit defective membrane potential and</u>

<u>compromised turnover</u>

Healthy mitochondria generate an electrochemical proton gradient that fuels the electron

transport chain and ATP generation. To better understand how A20 maintains mitochondrial

respiration in LPS stimulated macrophages, we first measured the mitochondrial membrane

potential in A20$^{WT}$ and A20$^{tiKO}$ cells. Consistent with our Seahorse assays (**Fig. 2A**), A20$^{tiKO}$

BMDMs exhibited decreased transmembrane potential after LPS stimulation (**Fig. 3A**). This

decreased membrane potential was not associated with reduced mitochondrial mass, as the

ratio of mitochondrial DNA to nuclear DNA was preserved in A20$^{tiKO}$ BMDMs (**Fig. 3B**).

Immunoblotting analyses of the mitochondrial protein Tomm20 revealed normal amounts of

this protein in both unstimulated and LPS stimulated A20$^{tiKO}$ BMDMs, reinforcing the notion

that mitochondrial mass is preserved in these cells (**Fig. 3C**). Taken together, these results

indicate that mitochondria are structurally intact, but functionally compromised in LPS

stimulated A20$^{tiKO}$ BMDMs.

<u>Figure 3:  Mitochondria in A20$^{tiKO}$ BMDMs exhibit lower membrane potential and decreased turnover.</u> **(A)** FACS

mediated measurement of mitochondrial transmembrane potential using tetramethylrhodamine methyl ester

(TMRM) labeling. Representative histograms of fluorescent TMRM expression in BMDMs of the indicated

genotypes shown at left. Percentage change in TMRM staining calculated at right from three independent

experiments. Mean values ± SD are shown. ** indicates p< 0.01 by unpaired T test. **(B)** Real time qPCR analyses

of mitochondrial and nuclear DNA from the indicated BMDMs. Mean values ± SD are shown. *** indicates

p<0.001. Data are representative of two independent experiments. **(C)** Immunoblot analysis of mitochondrial

Tomm20 protein expression in indicated BMDMs. GAPDH expression shown below as loading control. Data are

representative of three independent experiments. **(D)** FACS analyses of mitochondrial turnover using differential

labeling. BMDMs were first labeled with MitoTracker Green FM (MTG), then stimulated with LPS, and

subsequently labeled with MitoTracker Red (MTR) prior to FACS analysis. In this format, MTG represent pre-

existing (old) mitochondria; MTR represent all mitochondria present after LPS stimulation. Note that A20tiKO cells

retain a higher proportion of MTG positive (old) mitochondria than WT cells, particularly after LPS stimulation

(52.7% vs. 27.7%). Data are representative of three independent experiments. **(E)** Mass spectrometric

quantification of indicated tricarboxylic acid (TCA) cycle intermediates in A20$^{tiKO}$ (red) and WT (black) BMDMs after

LPS stimulation. Counts reflect average of five replicates. **(F)** Fluorometric assay of citrate synthase (CS) activity of

indicated cells after LPS stimulation was measured. **(G)** FACS analyses of mitochondrial superoxide production in

A20$^{tiKO}$ and WT BMDMs using using mitoSOX dye (mSOX). Data are representative of three independent

experiments.

During metabolically demanding cellular activation events, damaged mitochondria undergo

conformational changes and turnover to preserve the quantity and quality of these organelles

(Gkikas et al., 2018). We thus assessed mitochondrial turnover in A20$^{WT}$ and A20$^{tiKO}$ BMDMs.

We utilized differential labeling of mitochondria using MitoTracker Green (MTG) and

MitoTracker Red (MTR) before and after LPS stimulation, to quantitate pre-existing and newly

generated mitochondria. A20$^{tiKO}$ BMDMs contained a larger fraction of older, pre-existing

(MTG+) mitochondria than WT BMDMs without LPS, suggesting that basal mitochondrial

turnover is compromised in A20$^{tiKO}$ cells (**Fig. 3D**). This difference became much more

pronounced after LPS stimulation, indicating that A20 may be particularly important for

replacing senescent mitochondrial in activated macrophages (**Fig. 3D**). A20 dependent

mitochondrial replacement thus appears to collaborate with A20's ability to preserve

mitochondrial OCR in a coordinated effort to preserve cellular energetic homeostasis.

To further investigate the defective mitochondrial respiration in A20 deficient macrophages, we

performed targeted metabolomic studies on LPS stimulated cells. These assays revealed that

A20$^{tiKO}$ BMDMs contain similar levels of the tricarboxylic acid (TCA) cycle intermediates,

succinate, citrate and $\alpha$-ketoglutarate ($\alpha$-KG) as control cells, suggesting that TCA cycling

proceeds at sufficient rates in these cells to generate electron donors and substrates to the

mitochondrial electron transport chain (**Fig. 3E**). We also directly assayed the activity of citrate

synthase, the enzyme catalyzing the condensation between oxaloacetate and acetyl Co-A to

generate citrate. This enzyme is considered the first step of the TCA cycle. These tests showed

that citrate synthase activity was preserved in A20$^{tiKO}$ BMDMs throughout 24 hours of LPS

stimulation (**Fig 3F**). As the levels of citrate and succinate were comparable in WT and A20

deficient macrophages, previously described inhibition of these TCA cycle steps do not appear

to depend upon A20 (Tannahill et al., 2013; Liu et al., 2017; Jha et al., 2015). Finally, as

increased release of mitochondrial superoxide can cause dysfunctional mitochondria with

augmented electron leaks, we measured oxidation of the superoxide indicator, MitoSOX (**Fig.**

**3G**). During the 24 hours of LPS stimulation, MitoSOX levels remained similar between A20$^{WT}$

and A20$^{tiKO}$ cells, implying that impaired mitochondrial respiration in A20$^{tiKO}$ cells was not

caused by excessive induction of mitochondrial superoxide. Taken together, these data suggest

that acute A20 deficiency in macrophages causes defective mitochondrial respiration despite

the availability of comparable TCA substrates and structurally intact mitochondria.

<u>A20 deficient macrophages accumulate increased reactive oxygen and nitrogen species, and</u>

<u>exaggerated inducible nitric oxide synthase (iNOS) expression compromises mitochondrial</u>

<u>respiration</u>

Elevated cellular reactive oxygen species (ROS) and reactive nitrogen species (RNS) can

compromise mitochondrial function by inhibiting electron transport chain enzymes (Kaludercic

and Giorgio, 2016; Kowalska et al., 2020). Our metabolomic analyses revealed markedly

decreased glutathione (GSH) levels in LPS stimulated A20$^{tiKO}$ BMDMs (**Fig. 4A**). In addition,

glycine, an essential and rate-limiting precursor of GSH, was also significantly lower in these

BMDMs (**Fig. 4A**). As GSH is a major cellular antioxidant that reacts with and scavenges

potentially deleterious ROS and RNS, we hypothesized that A20 deficient macrophages might

accumulate excessive amounts of these reactive species. We measured cellular ROS/RNS levels

using the fluorescent indicator dichlorodihydrofluorescein diacetate (DCFDA), and found that

A20$^{tiKO}$ BMDMs exhibited significantly higher levels of DCFDA staining after LPS stimulation (**Fig.**

**4B**). Among RNS, nitric oxide (NO) and its metabolite peroxynitrite inhibit electron transport

chain enzymes by binding to $O_2$ binding sites on these proteins (Stewart and Heales, 2003). We

assayed NO levels and found that A20$^{tiKO}$ BMDMs produced markedly more NO than control

cells after LPS (**Fig. 4C**). This exaggerated NO production was not observed after stimulation

with other TLR ligands (**Fig. 4C**). We next tested whether abrogating production of these species

might restore mitochondrial defects. While allopurinol, a xanthine oxidase inhibitor, did not

rescue OCR defects in LPS stimulated A20$^{tiKO}$ BMDMs (**Fig. 4D**), the iNOS inhibitor 1400w

effectively restored mitochondrial respiration (**Fig. 4E**) and M2 repolarization (**Suppl. Fig. 1**).

Figure 4: A20$^{tiKO}$ BMDMs produce exaggerated amounts of nitric oxide (NO), and impaired mitochondrial

respiration is rescued by inhibition of nitric oxide synthase (iNOS). **(A)** Mass spectrometric quantification of

reduced and oxidized forms of glutathione (GSH) (left graph) and glycine (right graph) in A20$^{tiKO}$ (red) and WT

(black) BMDMs after LPS stimulation. * indicates p<0.05, ** indicates p<0.01, *** indicates p<0.001. Data are

representative of 5 independent samples. **(B)** Flow cytometric analyses of cytosolic ROS/RNS species in LPS treated

BMDMs of the indicated genotypes using dichlorodihydrofluorescein diacetate (DCFDA) dye. Y-axis represents

geometric mean fluorescence intensity (gMFI) of each sample relative to untreated cells at the indicated

timepoints after LPS stimulation. * indicates p<0.05, ** indicates p<0.01.  Mean values ± SD are shown. Pooled

results from five independent experiments are shown. **(C)** Calorimetric determination of nitric oxide (NO)

production from A20$^{tiKO}$ (red) and WT (black) BMDMs after stimulation with the indicated TLR ligands. Mean values

± SD are shown. *** indicates p<0.001. Data are representative of three independent experiments. **(D)** Seahorse

mitochondrial stress test of A20$^{tiKO}$ (red) and WT (black) BMDMs with or without xanthine oxidase inhibitor

allopurinol (allo). Mean values ± SD are shown. Data are representative of three independent experiments. **(E)**

Seahorse mitochondrial stress test of A20$^{tiKO}$ (red) and WT (black) BMDMs with or without iNOS inhibitor 1400w.

Basal and maximal respiratory rates, spare respiratory capacity (SRC), and ATP production are shown at right.

Mean values ± SD are shown. *** indicates p<0.001. Data are representative of three independent experiments. In

all panels, * indicates p<0.05, ** indicates p<0.01, *** indicates p<0.001 using unpaired two-tailed t test.

NO is produced by several nitric oxide synthase enzymes that are expressed in multiple cell

types. To understand why A20$^{tiKO}$ BMDMs secreted excessive amount of NO after LPS

stimulation, we measured the expression of iNOS (*nos2*), the inducible isoform of NOS that is

robustly expressed in macrophages and generates NO in a Ca$^{++}$ independent fashion

(Förstermann and Sessa, 2012). While resting BMDMs express negligible levels of iNOS mRNA

and protein, these species were both expressed at markedly higher levels in A20$^{tiKO}$ BMDMs

than control cells after LPS stimulation (**Figs. 5A, B**). To define the contribution of iNOS to

mitochondrial dysfunction in A20$^{tiKO}$ BMDMs, we interbred A20$^{Flox/Flox}$ ROSA/ER-Cre mice with

iNOS$^{-/-}$ mice (Laubach et al., 1995) and tested LPS responses of the compound mutant

macrophages derived from these mice. Immunoblot analyses of A20 and iNOS expression

confirmed elimination of these proteins in the appropriate genotypes (**Suppl. Fig. 2**).

Stimulation of A20$^{tiKO}$ iNOS$^{-/-}$ BMDMs with LPS resulted in normalization of cellular ROS/RNS

levels when compared with A20$^{tiKO}$ BMDMs (**Fig. 5C**). Hence, iNOS is predominantly responsible

for the exaggerated ROS/RNS species generated in LPS stimulated A20$^{tiKO}$ BMDMs. Similar

results were obtained with the iNOS inhibitor 1400w (**Suppl. Fig. 3**). Remarkably, extracellular

flux assays revealed that ablation of iNOS completely restored mitochondrial respiration in

A20$^{tiKO}$ BMDMs (**Fig. 5D**). These A20$^{tiKO}$ iNOS$^{-/-}$ BMDMs also exhibited normal levels of ATP

production and spare respiratory capacity (**Fig. 5D**). In addition to restoring deficient

mitochondrial respiration, abrogation of iNOS expression also normalized the exaggerated

lactate production observed in A20$^{tiKO}$ BMDMs (**Fig. 5E**). Therefore, iNOS expression is a

dominant mechanism regulating the balance between glycolytic and mitochondrial respiration.

<u>Figure 5:  Expression of iNOS is greatly induced in A20$^{tiKO}$ BMDMs, and genetic ablation of *nos2* restores

mitochondrial bioenergetic balance</u>. (**A**) Real-time qPCR analyses of *nos2* mRNA expression relative to *actin* mRNA

in A20$^{tiKO}$ (red) and A20$^{WT}$ (black) BMDMs at indicated timepoints after LPS stimulation. Mean values ± SD are

shown. *** indicates p<0.001 by unpaired T test. Data are representative of three independent experiments. (**B**)

Immunoblot analyses of iNOS protein expressions in indicated BMDMs at indicated timepoints after LPS

stimulation. GAPDH protein expression shown below as loading control. The iNOS inhibitor 1400w inhibits iNOS

activity rather than iNOS expression and is shown as a control. Data are representative of three independent

experiments. (**C**) Flow cytometry analysis of ROS/RNS species by DCFDA staining in the indicated genotypes of

BMDMs after LPS stimulation. DCFDA staining intensity is plotted relative to unstimulated cells. Note that

increased ROS/RNS in LPS stimulated A20$^{tiKO}$ cells (red) is normalized in LPS stimulated A20$^{tiKO}$ iNOS$^{-/-}$ cells (blue).

Mean values ± SD are shown.  * indicates p<0.05, ** indicates p<0.01. Data are representative of three

independent experiments. (**D**) Seahorse mitochondrial stress test of BMDMs from the indicated genotypes of mice.

Note that depressed basal and maximal oxygen consumption rates (OCR) in A20$^{tiKO}$ cells (red) is normalized in LPS

stimulated A20$^{tiKO}$ iNOS$^{-/-}$ cells (blue).  Graphic representation of basal and maximal OCRs, spare respiratory

capacity (SRC), and ATP production are shown below. Mean values ± SD are shown. *** indicates p<0.001. Data

are representative of four independent experiments. (**E**) Colorimetric measurements of lactate secretion by

indicated genotypes of cells. Mean values ± SD are shown. * indicates p< 0.05, ** indicates p<0.01 by unpaired T

test. Data are representative of three independent experiments. (**F**) Real-time quantitative PCR analyses of *tnf, il-6*

*and ifnb1* mRNA expression relative to *actin* in the respective BMDMs after LPS stimulation. Mean values ± SD are

shown. Data are representative of three independent experiments. **(G)** ELISA of IL-1β secretion by A20$^{WT}$ and

A20$^{tiKO}$ and WT BMDMs after indicated treatments. Mean values ± SD are shown. * indicates p<0.05, *** indicates

p<0.001. Data are representative of two independent experiments. **(H)** ELISA of IL-10 secretion by A20$^{WT}$ and

A20$^{tiKO}$ and WT BMDMs after indicated treatments. Mean values ± SD are shown. *** indicates p<0.001 by

unpaired T test. Data are representative of two independent experiments.

<u>Resuscitation of mitochondrial respiration in A20$^{tiKO}$ iNOS$^{-/-}$ BMDMs occurs independently from</u>

<u>A20 regulated inflammatory cytokines and inflammasome activation</u>

A20 deficient macrophage express greater amounts of inflammatory cytokines than control

cells in response to LPS stimulation (Boone et al., 2004; Turer et al., 2008). These cytokines

could theoretically contribute to mitochondrial dysfunction in A20$^{tiKO}$ BMDMs. We tested

cytokine responses in A20$^{tiKO}$ and A20$^{tiKO}$ iNOS$^{-/-}$ BMDMs after LPS stimulation and found that

exaggerated production of TNF, IL-6, and IFN-β mRNAs by A20$^{tiKO}$ BMDMs was not abrogated by

elimination of iNOS expression in A20$^{tiKO}$ iNOS$^{-/-}$ BMDMs (**Fig. 5F**). Therefore, iNOS deficiency

decouples mitochondrial dysfunction in A20$^{tiKO}$ BMDMs from exaggerated secretion of pro-

inflammatory cytokines.

ROS have been described to mediate a second step to activate NLRP3 inflammasomes (Zhou et

al., 2011), and A20$^{-/-}$ BMDMs exhibit aberrant spontaneous NLRP3 inflammasome activation,

i.e., secretion of mature IL-1β after LPS without ATP stimulation (Walle et al., 2014; Duong et

al., 2015). Hence, exaggerated NO produced in A20$^{tiKO}$ BMDMs might drive aberrant NLRP3

inflammasome activation by generating ROS that activate these protein complexes. We

measured IL-1β secretion by ELISA, and found that A20$^{tiKO}$ BMDMs secreted IL-1β even in the

presence of the iNOS inhibitor 1400w that restored mitochondrial respiration (**Fig. 5G**). Hence,

A20 regulates mitochondrial respiration via a distinct cell biological pathway from its regulation

of inflammasome activation.

IL-10 deficient macrophages exhibit defects in mitochondrial function after LPS stimulation (Ip

et al., 2017). Hence, A20 deficient macrophages might develop mitochondrial dysfunction due

to defective IL-10 production. We assayed IL-10 secretion from LPS stimulated A20$^{tiKO}$ and

control BMDMs, and found that A20$^{tiKO}$ cells secreted significantly more IL-10 than control cells

(**Fig. 5H**). In addition, while mitochondrial defects in IL-10 deficient macrophages do not depend

upon NO (Ip et al., 2017), this reactive nitrogen species is critical for compromising

mitochondrial respiration in A20 deficient cells. Thus, A20 and IL-10 appear to control

mitochondrial function via distinct mechanisms. Nevertheless, the shared metabolic defects of

IL-10 deficient and A20 deficient macrophages may underlie overlapping links of these two

molecules to early onset inflammatory bowel disease and other inflammatory diseases.

In summary, our findings reveal a profound role for A20 in restraining aerobic glycolysis and

supporting metabolic reprogramming toward oxidative phosphorylation. The metabolic impact

of A20 on macrophage homeostasis is mediated by iNOS and NO. A20 deficient macrophages

share some mitochondrial defects with IL-10 deficient macrophages, but are distinguished by

the secretion of high levels of IL-10 and reliance upon iNOS. As A20 deficient macrophages may

contribute to inflammatory processes in an expanding array of human diseases, including

haploinsufficient A20 patients, understanding the metabolic defects found in these cells should

open novel avenues toward controlling these clinical syndromes. Finally, as A20 is a tumor

suppressor in both mice (Shao et al., 2013) and humans (Kato et al., 2009; Compagno et al.,

2009), the severe skewing toward aerobic glycolysis in A20 deficient macrophages suggests that

loss of A20 may also support tumorigenesis via metabolic mechanisms as well as via regulation

of cell survival pathways.

430
431     **Materials and Methods**
432
<u>Mice</u>

$A20^{Flox}$ mice were generated in our laboratory and described previously (Tavares et al., 2010).

Rosa/ER-Cre mice and $iNOS^{-/-}$ mice were purchased from Jackson Laboratories. All animal

research was approved by University of California San Francisco (UCSF) Office of Ethics and

Compliance.

<u>Generation of bone marrow derived macrophages and treatments</u>

Bone marrow derived macrophages were prepared as previously described (Duong et al., 2015).

Deletion of A20 *in vitro* was performed by adding 60nM 4-hydroxytamoxifen (4-OHT) 2 days

before harvested. The doses of ligands used were as follows: lipopolysaccharide 100 ng/ml;

poly(I:C) HMW 20 μg/ml; $Pam_3CSK_4$ 1 μg/ml; CpG-A 2 μM; CpG-B 250 nM, Gardiquimod 1

μg/ml; PolyU (complexed with lipofectamine) 0.5 μg/ml and flagellin (from *S. typhimurium*) 2

μg/ml (Invivogen). For pharmacological inhibitors: allopurinol 25 μM and 1400w 100 μM

(Sigma). For macrophage repolarization, cells were stimulated with LPS for 24 hours and

washed twice before the addition of 200 ng/ml IL-4 for another 24 hours (PeproTech). The

resulting cells were analyzed by flow cytometry (BD Fortessa) at the UCSF Parnassus Flow Core,

RRID:SCR_01826, and the software FlowJo (BD).

<u>NO, ROS and lactate assays</u>

Culture supernatants from stimulated macrophages were briefly centrifugated to remove any

cell debris. Nitric oxide (NO) was measured using a Total Nitric Oxide Assay kit (Invitrogen). To

measure intracellular ROS/RNS, macrophages were loaded with 25 µM $H_2DCFDA$ (Sigma) in

serum-free DMEM medium and incubated for 30 min at 37 °C. Cells were then collected,

washed once with PBS and resuspended in FACS buffer (2.5% FBS, 0.1% sodium azide, 0.02%

sodium bicarbonate in PBS) supplemented with 7-AAD at a working concentration of 2.5 µg/ml

for the exclusion of dead cells, and analyzed by flow cytometry (BD Fortessa) at the UCSF

Parnassus Flow Core, RRID:SCR_01826, and the software FlowJo (BD). Lactate secretion was

measured using a colorimetric Lactate Assay kit (BioVision). All assays were performed

according to manufacturer's instructions.

<u>Mitochondrial assays</u>

Mitochondrial membrane potential was measured using tetramethylrhodamine methyl ester

(TMRM) (Invitrogen) according to manufacturer's instruction. TMRM (20 nM) was added and

incubated for 30 minutes in 37°C $CO_2$ incubator before cell harvest. Mitochondrial turnover was

examined by first labeling mitochondria in unstimulated cells with MitoTracker Green FM

(MTG) (Invitrogen) for 30 minutes according to manufacturer's instructions. Cells were then

washed twice with PBS and stimulated with LPS for 24 hours. MitoTracker Red CM-$H_2$Xros

(MTR) (Invitrogen) was then added to the culture 30 minutes before harvest in 37°C $CO_2$

incubator. Dead cells were excluded with LIVE/DEAD Fixable Near-IR Viability Dye (Invitrogen)

according to manufacturer's instructions. Fluorescent signals were analyzed with flow

cytometry (BD Fortessa) at the UCSF Parnassus Flow Core, RRID:SCR_01826, and the software

FlowJo (BD). Mitochondrial DNA (mDNA) to nuclear DNA (nDNA) ration was determined as

described in details by Malik and colleagues (Malik et al., 2016) using Quantifast SYBR Master

Mix (Qiagen) and analyzed with quantitative real-time PCR (Quantstudio 6 Flex Real-Time PCR

Systems from Thermo). To measure the mitochondrial protein Tomm20, cell palettes were

subjected to lysis in 1x RIPA lysis buffer (Alfa Aesar) supplemented with complete EDTA-Free

Protease Inhibitor Cocktail (Roche), phosphatase inhibitors (1 mM $Na_3VO_4$ and 10 mM NaF) and

10 mM N-ethylmaleimide (Sigma), and centrifuged for 20 minutes (21,000 x g at 4°C).

Mitochondrial citrate synthase activity was quantitated using the Citrate Synthase Activity

Assay kit (BioVision) according to the manufacturer's instruction. To measure mitochondrial

derived superoxide, the superoxide indicator MitoSOX Red (Invitrogen) was used according to

manufacturer's instructions. Mitochondrial TCA cycle metabolites, GSH and glycine were

measured via mass spectrometry as previously described (Louie et al, 2016; Kohnz et al, 2016).

Quantitative real-time PCR

Total RNA was extracted from BMDMs after treatment using the RNeasy Mini kit (Qiagen)

according to the manufacturer's instructions with on-column DNase I digestion (Qiagen). cDNA

was synthesized using the PrimeScript RT-PCR kit (Takara Bio) according to manufacturer's

instructions. Quantitative real-time PCR was performed using TaqMan probes on Quantstudio 6

Flex Real-Time PCR Systems from Thermo.

### Seahorse assays

494 BMDMs were seeded 2 hours before stimulation on the 24-well Seahorse cell culture plates

495 (Agilent). Cells were stimulated for 24 hours in 37°C $CO_2$ incubator, after which glycolytic

496 activity was analyzed using the XF Glycolytic Rate Assay Kit (Agilent). For mitochondrial oxygen

497 consumption, the XF Cell Mito Stress Test Kit was used (Agilent). Both tests were performed

498 according to manufacturer's instructions.

### Oil Red O staining

BMDMs were seeded on Lab-Tek II 8-well Chamber Slides (Thermo) and stimulated 2 hours

after attachment. Cells were harvested by washing in PBS twice and then fixed in 10% NBF

(Sigma) at room temperature for 10 minutes. Fixative was removed and fresh 10% NBF was

added and allow fixation for another 3 hours. After fixation, cells were washed with $ddH_2O$

twice and subsequently with 60% isopropanol for 5 minutes at room temperature. The samples

were left to dry completely overnight. The next day, samples were stained and incubated with

Oil Red O solution (0.21% Oil Red O powder, Sigma, dissolved in isopropanol and filtered, let set

at room temperature for 20 minutes before use) for 10 minutes at room temperature.

Afterwards the dye was removed and $ddH_2O$ immediately added. Cells were washed 4 times

with $ddH_2O$. The slides were allowed to dry and mounted in ProLong Gold Antifade Mountant

(Invitrogen) before acquiring images under the microscope (Zeiss Axioscope).

512
513
**Acknowledgments**

This work was supported by NIH RO1 grants, the UCSF Nutrition and Obesity Research Center

(P30DK098722) and the UCSF Liver Center (P30DK026743) from the National Institute of

Diabetes and Digestive and Kidney Diseases. Author contributions: M. Chan, B. Malynn, A. Ma

conceptualized the study, designed and interpreted the experiments, and wrote the

manuscript. M. Chan performed and analyzed the experiments. E. Grossman and D. Nomura

performed and analyzed metabolomic experiments. R. Advincula and P. Achacoso assisted with

mouse breeding and experiments.  The authors declare no competing financial interests.

**References**

Aeschlimann FA, Batu ED, Canna SW, Go E, Gül A, Hoffmann P, et.al. A20 haploinsufficiency
(HA20): clinical phenotypes and disease course of patients with a newly recognised NF-
kB-mediated autoinflammatory disease. Ann Rheum Dis. 2018;77: 728–735.

Boone DL, Turer EE, Lee EG, Ahmad RC, Wheeler MT, Tsui C, et. al. The ubiquitin-modifying
enzyme A20 is required for termination of Toll-like receptor responses. Nat Immunol.
2004;5:1052–1060.

Bozza PT, and Viola JPB. Lipid droplets in inflammation and cancer. Prostaglandins Leukot
Essent Fatty Acids. 2010;82:243–250.

Buck MD, O'Sullivan D, Klein Geltink RI, Curtis JD, Chang CH, Sanin DE, et al. Mitochondrial
dynamics controls T cell fate through metabolic programming. Cell. 2016;166:63–76.

Compagno M, Lim WK, Grunn A, Nandula SV, Brahmachary M, Shen Q, et. al. Mutations of
multiple genes cause deregulation of NF-kappaB in diffuse large B-cell lymphoma.
Nature. 2019;459:717–721.

Duong BH, Onizawa M, Oses-Prieto JA, Advincula R, Burlingame A, Malynn BA, et al. A20
restricts ubiquitination of pro-interleukin-1β protein complexes and suppresses NLRP3
inflammasome activity. Immunity. 2015;42:55–67.

Förstermann U, and Sessa WC. Nitric oxide synthases: regulation and function. Eur Heart J.
2012;33:829–837.

Galván-Peña S, and O'Neill LAJ. Metabolic reprograming in macrophage polarization. Front
Immunol. 2014;5:420.

Gkikas I, Palikaras K, and Tavernarakis N. The role of mitophagy in innate immunity. Front.
Immunol. 2018;9-21.

Hammer GE, Turer EE, Taylor KE, Fang CJ, Advincula R, Oshima S, et.al. Expression of A20 by
dendritic cells preserves immune homeostasis and prevents colitis and
spondyloarthritis. Nat Immunol. 2011;12:1184–1193.

Ip WKE, Hoshi N, Shouval DS, Snapper S, and Medzhitov R. Anti-inflammatory effect of IL-10
mediated by metabolic reprogramming of macrophages. Science. 2017;356:513–519.

Jarc E, and Petan T. A twist of FATe: Lipid droplets and inflammatory lipid mediators. Biochimie.
2020;169:69–87.

Jha AK, Huang SCC, Sergushichev A, Lampropoulou V, Ivanova Y, Loginicheva E, et. al. Network
integration of parallel metabolic and transcriptional data reveals metabolic modules
that regulate macrophage polarization. Immunity. 2015;42:419–430.

Kaludercic N, and Giorgio V. 2016. The dual function of reactive oxygen/nitrogen species in
bioenergetics and cell death: the role of ATP synthase. Oxidative Medicine and Cellular
Longevity. 2016:e3869610.

Kato M, Sanada M, Kato I, Sato Y, Takita J, Takeuchi K, et. al. Frequent inactivation of A20 in B-
cell lymphomas. Nature. 2009;459:712–716.

Kohnz RA, Roberts LS, DeTomaso D, Bideyan L, Yan P, Bandyopadhyay, et. al. 2016. Protein
sialylation regulates a gene expression signature that promotes breast cancer cell
pathogenicity. ACS Chem Biol. 2016;11:2131-2139.

Kool M, van Loo G, Waelput W, De Prijck S, Muskens F, Sze M, et. al. The ubiquitin-editing
protein A20 prevents dendritic cell activation, recognition of apoptotic cells, and
systemic autoimmunity. Immunity. 2010;35:82–96.

Kotwal GJ, and Chien S. Macrophage differentiation in normal and accelerated wound healing.
Results Probl Cell Differ. 2017;62:353–364.

Kowalska M, Piekut T, Prendecki M, Sodel A, Kozubski W, and Dorszewska J. Mitochondrial and
nuclear DNA oxidative damage in physiological and pathological aging. DNA and Cell
Biology. 2020;39:1410–1420.

Laubach VE, Shesely EG, Smithies O, and Sherman PA. Mice lacking inducible nitric oxide
synthase are not resistant to lipopolysaccharide-induced death. Proc Natl Acad Sci U S A.
1995;92:10688–10692.

Lee EG, Boone DL, Chai S, Libby SL, Chien M, Lodolce JP, and Ma A. Failure to Regulate TNF-
Induced NF-κB and Cell Death Responses in A20-Deficient Mice. Science.
2000;289:2350–2354.

Liu PS, Wang H, Li X, Chao T, Teav T, Christen S, et. al. α-ketoglutarate orchestrates macrophage
activation through metabolic and epigenetic reprogramming. Nat Immunol.
2017;18:985–994.

Louie SM, Grossman EA, Crawford LA, Ding L, Camarda R, Huffman TR, et al.. GSTP1 Is a driver
of triple-negative breast cancer cell metabolism and pathogenicity. Cell Chem Biol.
2016;23:567-578.
585
Ma A, and Malynn BA. A20: linking a complex regulator of ubiquitylation to immunity and
human disease. Nat Rev Immunol. 2012;12:774–785.

Macintyre AN, Gerriets VA, Nichols AG, Michalek RD, Rudolph MC, Deoliveira D, et. al. The
glucose transporter Glut1 is selectively essential for CD4 T cell activation and effector
function. Cell Metab. 2014;20:61–72.

Makowski L, Chaib M, and Rathmell JC. Immunometabolism: From basic mechanisms to
translation. Immunol Rev. 2020;295:5–14.

Malik AN, Czajka A, and Cunningham P. Accurate quantification of mouse mitochondrial DNA
without co-amplification of nuclear mitochondrial insertion sequences. *Mitochondrion*.
2016;29:59–64.

Matmati M, Jacques P, Maelfait J, Verheugen E, Kool M, Sze M, et. al. A20 (TNFAIP3) deficiency
in myeloid cells triggers erosive polyarthritis resembling rheumatoid arthritis. Nat
Genet. 2011;43:908–912.

Mills EL, Kelly B, Logan A, Costa ASH, Varma M, Bryant CE, et. al. Succinate Dehydrogenase
Supports Metabolic Repurposing of Mitochondria to Drive Inflammatory Macrophages.
Cell. 2016;167:457-470.e13.

Mulvihill MM, and Nomura DK. Metabolomic strategies to map functions of metabolic
pathways. Am J Physiol Endocrinol Metab. 2014;307:E237-244.

Muto T, Walker CS, Choi K, Hueneman K, Smith MA, Gul Z, et al. Adaptive response to
inflammation contributes to sustained myelopoiesis and confers a competitive
advantage in myelodysplastic syndrome HSCs. Nat Immunol. 2020;21:535–545.

Ohnishi H, Kawamoto N, Seishima M, Ohara O, and Fukao T. A Japanese family case with
juvenile onset Behçet's disease caused by TNFAIP3 mutation. Allergology International.
2017;66:146–148.

O'Neill LAJ and Pearce EJ. Immunometabolism governs dendritic cell and macrophage function.
J Exp Med. 2016;213:15–23.

Onizawa M, Oshima S, Schulze-Topphoff U, Oses-Prieto JA, Lu T, Tavares R, et al. The ubiquitin-
modifying enzyme A20 restricts ubiquitination of the kinase RIPK3 and protects cells
from necroptosis. Nat Immunol. 2015;16:618–627.

Razani B, Whang MI, Kim FS, Nakamura MC, Sun X, Advincula R, et al. Non-catalytic ubiquitin
binding by A20 prevents psoriatic arthritis–like disease and inflammation. Nat Immunol.
2020;21:422–433.

Rosas-Ballina M, Guan XL, Schmidt A, and Bumann D. Classical activation of macrophages leads
to lipid droplet formation without de novo fatty acid synthesis. Front Immunol. 2020;11.

Schmitz R, Hansmann M-L, Bohle V, Martin-Subero JI, Hartmann S, Mechtersheimer G, et al.
TNFAIP3 (A20) is a tumor suppressor gene in Hodgkin lymphoma and primary
mediastinal B cell lymphoma. J Exp Med. 2009;206:981–989.

Schneider C, O'Leary CE, von Moltke J, Liang HE, Ang QY, Turnbaugh PJ, et al. A metabolite-
triggered tuft cell-ILC2 circuit drives small intestinal remodeling. Cell. 2018;174:271-
284.e14.

Seeley JJ and Ghosh S. Molecular mechanisms of innate memory and tolerance to LPS. Journal
of Leukocyte Biology. 2017;101:107–119.

Shao L, Oshima S, Duong B, Advincula R, Barrera J, Malynn BA, and Ma A. A20 restricts wnt
signaling in intestinal epithelial cells and suppresses colon carcinogenesis. PLoS One.
2013;8:e62223. doi:10.1371/journal.pone.0062223.

Stewart VC and Heales SJR. Nitric oxide-induced mitochondrial dysfunction: implications for
neurodegeneration. Free Radical Biology and Medicine. 2003;34:287–303.

Takagi M, Ogata S, Ueno H, Yoshida K, Yeh T, Hoshino A, et al. Haploinsufficiency of TNFAIP3
(A20) by germline mutation is involved in autoimmune lymphoproliferative syndrome.
Journal of Allergy and Clinical Immunology. 2017;139:1914–1922.

Tannahill G, Curtis A, Adamik J, Palsson-McDermott E, McGettrick A, Goel G, et al. Succinate is a
danger signal that induces IL-1β via HIF-1α. Nature. 2013;496:238–242.

Tavares RM, Turer EE, Liu CL, Advincula R, Scapini P, Rhee L, et al. The ubiquitin modifying
enzyme A20 restricts B cell survival and prevents autoimmunity. Immunity.
2010;33:181–191.

Turer EE, Tavares RM, Mortier E, Hitotsumatsu O, Advincula R, Lee B, et al. The Journal of
Experimental Medicine. 2008;205:451–464.

Vats D, Mukundan L, Odegaard JI, Zhang L, Smith KL, Morel CR, et al. Oxidative metabolism and
PGC-1β attenuate macrophage-mediated inflammation. Cell Metabolism. 2006;4:13–24.

Walle LV, Opdenbosch NV, Jacques P, Fossoul A, Verheugen E, Vogel P, et al Negative regulation
646            of the NLRP3 inflammasome by A20 protects against arthritis. Nature. 2014;512:69–73.

647     Wen H, Gris D, Lei Y, Jha S, Zhang L, Huang MT, et al. Fatty acid–induced NLRP3-ASC
648            inflammasome activation interferes with insulin signaling. Nature Immunology.
649            2013;12:408–415. doi:10.1038/ni.2022.

650     Zhou Q, Wang H, Schwartz DM, Stoffels M, Park YH, Zhang Y, et al. Loss-of-function mutations
651            in TNFAIP3 leading to A20 haploinsufficiency cause an early onset autoinflammatory
652            syndrome. Nat Genet. 2016;48:67–73.

653     Zhou R, Yazdi AS, Menu P, and Tschopp J. A role for mitochondria in NLRP3 inflammasome
654            activation. Nature. 2011;469:221–225.

# Fig 3. : Mitochondria in A20-deficient BMDMs have lower membrane potential and turnover rate, but are mostly intact after LPS stimulation

Fig 4.