A trait syndrome ties cell morphology to glycolysis across the yeast subphylum

Linda C. Horianopoulos[1,2*], Christina M. Chavez[3,4*], Antonis Rokas[3,4–] and Chris Todd Hittinger[1–]

[1] *Laboratory of Genetics, DOE Great Lakes Bioenergy Research Center, Wisconsin Energy*

Institute, Center for Genomic Science Innovation, J.F. Crow Institute for the Study of Evolution,

University of Wisconsin-Madison, WI 53726, USA

[2] *Department of Food Science, University of Guelph, Guelph, ON, N1G 2W1, Canada*

[3] *Department of Biological Sciences, Vanderbilt University, Nashville, TN 37235, USA*

[4] *Evolutionary Studies Initiative, Vanderbilt University, Nashville, TN 37235, USA*

* These authors contributed equally to this work

⬚Corresponding authors: cthittinger@wisc.edu & antonis.rokas@vanderbilt.edu

Keywords: yeast, cell morphology, metabolism, genome size, cell size, surface area-to-volume

Running title: Metabolism and cell morphology in yeasts

Summary

Traits that co-vary across species can provide fundamental insights into the trade-offs and constraints that govern their evolution. In a recent article in *Current Biology*, Li et al.[1] reported that glucose uptake rates (GUR) are inversely correlated with the cell surface area-to-volume (SA:V) ratio across 11 yeast species. Here we substantially expand this analysis to 282 species to test whether the GUR-SA:V correlation generalizes across the ancient Saccharomycotina yeast subphylum and to determine the contribution of shared evolutionary history to the co-variation of these two traits. Using regression models that account for co-variation due to phylogeny, we found that extracellular acidification rates (ECAR, which we used as a proxy for GUR) had a significant correlation with SA:V across Saccharomycotina. In contrast to Li et al.[1], our increased sample sizes provided statistical power to reveal additional significant correlations of ECAR with genome sizes and growth rates. Our findings dramatically extend and expand those of Li et al.[1] and suggest that a trait syndrome governs several metabolic, genomic, and morphological traits across yeasts.

Main Text

The Saccharomycotina yeast subphylum includes over a thousand species and displays impressive genetic and phenotypic diversity.[2,3] The ability of most yeast species to grow well in axenic culture enables both the characterization of many traits and the exploration of biological principles that govern their variation.[3] Such studies have revealed that certain traits are consistently associated with each other and may collectively contribute to trait syndromes.[4] These trait syndromes can be driven by intrinsic factors (e.g. genomic characteristics) or extrinsic factors (e.g. varying resource availability).[3,5]

Recently, Li et al.[1] reported the interesting finding that glucose uptake rate (GUR) is inversely correlated with surface area-to-volume ratio (SA:V) across 16 yeast strains in 10 Ascomycota species (subphyla Saccharomycotina and Taphrinomycotina) and 1 Basidiomycota

species (subphylum Pucciniomycotina). Parallel to and independently from Li et al., we have

measured glucose-induced extracellular acidification rates (ECAR) as a proxy for glycolytic

rates[6] and characterized yeast phenotypic variation by compiling data on asexual budding cell

size and shape[7] of 282 diverse yeast species within Saccharomycotina. Our dataset is uniquely

suited to test whether the correlation reported by Li et al.[1] holds true across an entire yeast

subphylum. We also considered two additional traits, genome size and rates of growth on

glucose, which we previously collected for these yeast species as part of the Y1000+ Project.[3]

First, we calculated the surface area, volume, and SA:V of each yeast species (Table

S1). We then calculated the phylogenetic signal of each trait using Pagel's lambda

(Supplemental methods) and found that all traits had high and significant phylogenetic signal

values (ECAR: λ = 0.815, p-value = $1.14E^{-49}$; SA:V ratio: λ = 0.846, p-value = $1.23E^{-19}$; genome

size: λ = 0.975, p-value = $1.07E^{-42}$; and growth rate on glucose: λ = 0.725, p-value = $5.78E^{-7}$).

The high phylogenetic signal values suggest that some of the variation in each trait can be

attributed to shared evolutionary history. Thus, we applied phylogenetically-corrected regression

models (Supplemental methods) to determine the relationships that exist between ECAR and

SA:V and genome size (n = 282 species), as well as between ECAR and growth rate on

glucose (n = 231 species). We found that ECAR was positively correlated with SA:V

(phylogenetic signal λ of the model = 0.808, R^2 = 0.565, p-value = 0.0001, Figure 1A). Since

ECAR is negative (faster glucose-induced acidification) when GUR is high, these results support

the main finding of Li et al.[1] and reveal that the inverse correlation between GUR and SA:V

holds across a much larger group of yeast species and that it cannot be explained solely

through phylogenetic co-variance.

Figure 1. Glycolytic rate is significantly associated with cell morphology, genome size, and growth across the Saccharomycotina yeasts. (A-C) Scatter plots show significant correlations between ECAR (extracellular acidification rates, measured as a proxy for glycolytic rates) and: (A) cell surface area-to-volume ratio; (B) genome size; and (C) growth rate on glucose. Each dot represents a single yeast species colored according to its taxonomic order as labeled (B). *S. cerevisiae* is represented by a triangle in panels (A) and (B). The R^2 and p-values are based on a phylogenetic generalized least squares model, which accounts for correlation stemming from shared phylogenetic history. (D) A schematic model proposes a trait syndrome associated with a rapid glucose uptake rate among diverse yeast species.

Typically, smaller cells with larger SA:V ratios have been found to exhibit higher

metabolic rates due to increased transport capacity at the cell surface.[8] However, an alternative

model suggests that efficient scaling of biosynthesis and sufficient protein production are key to

cell enlargement and rapid metabolism.[9] Our findings are largely consistent with this second

model as cells with lower SA:V ratios generally had higher metabolic rates. We speculate that,

in yeasts, larger cells have more space in the cytosol for the catabolism of glucose, thereby

enabling a stronger pull through glycolysis and consequently rapid GUR and growth rates. This

glycolytic effect may be particularly applicable to Crabtree/Warburg-positive yeasts, which

ferment in the presence of oxygen and have been proposed as an exception to the classic view

that high SA:V is necessary for high GUR.[8] Overall, this model proposes that the metabolic rate

of yeasts growing on glucose is more influenced by enhanced glycolytic flux enabled by

increased cytosolic volume than by nutrient import.

In contrast to the findings of Li et al.[1], our much larger dataset also showed significant

negative correlations between ECAR and genome size ($\lambda = 0.826$, $R^2 = 0.550$, p-value =

0.0182, Fig 1B), and between ECAR and growth rate on glucose ($\lambda = 0.825$, $R^2 = 0.556$, p-value

= 0.0031, Fig 1C). The contrast between our findings and those of Li et al.[1] is likely explained by

the enhanced statistical power of our dataset. Previous work showed that a larger genome size

was associated with a generalist phenotype in Saccharomycotina yeasts, wherein generalists

grew on a larger range of carbon sources and had faster growth rates.[3] Our data further show

that yeast species with larger genomes have increased glucose uptake rates and faster rates of

growth on glucose. Therefore, yeasts with larger genomes, such as those with whole genome

duplications[10], could be promising for industrial applications requiring rapid metabolism.

Altogether our results and previous work lead us to propose a trait syndrome in which

glycolytic rate is associated with multiple morphological and genomic features across the yeast

subphylum. Furthermore, the association between metabolism and genome size suggests that

gene acquisitions, gene family expansions, or genome duplications may be, at least in part,

influencing glucose uptake rates and cell size. Finally, our results raise important questions with

respect to the mechanisms that link primary carbon metabolism to the regulation of cell

morphology, whether genomic features other than size contribute to the relationship between

these traits, and the environments that select for this trait syndrome. The findings reported here

and by Li et al.[1], alongside the rich genomic dataset available for the Saccharomycotina

subphylum[3], are poised to guide experiments that will uncover these mechanisms.

Acknowledgements

We would like to thank members of the Y1000+ Project (http://y1000plus.org) for helpful

discussions. This project was supported by the National Science Foundation under Grants Nos.

DEB-2110403 (to C.T.H.) and DEB-2110404 (to A.R.); in part by the Great Lakes Bioenergy

Research Center, U.S. Department of Energy, Office of Science, Biological and Environmental

Research Program under Award Number DE-SC0018409 (of which C.T.H. is a co-investigator);

and the National Institute of Food and Agriculture, United States Department of Agriculture,

Hatch project 7005101 (to C.T.H.). Research in A.R.'s lab is also supported by the National

Institutes of Health/National Institute of Allergy and Infectious Diseases (R01 AI153356). L.C.H.

was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC)

postdoctoral fellowship.

Author contributions

132 L.C.H, C.M.C, A.R., and C.T.H. conceptualized this project. L.C.H and C.M.C. developed the

133 methodology, curated and analyzed the data, and prepared the figures under the supervision

134 and guidance of A.R. and C.T.H. L.C.H and C.M.C wrote the manuscript, and A.R. and C.T.H.

135 reviewed edited and approved the manuscript. L.C.H., A.R., and C.T.H. acquired funding for this

136 project.

Conflicts of Interest

139 A.R. is a scientific consultant of LifeMine Therapeutics, Inc. and a member of the Advisory

140 Board of *Current Biology.* All other authors declare no competing interests.

References

1.   Li, X., de Assis Souza, R., and Heinemann, M. (2025). The rate of glucose metabolism
sets the cell morphology across yeast strains and species. Current Biology *35*, 788-
798.e4. https://doi.org/10.1016/j.cub.2024.12.039.

2.   Shen, X.X., Opulente, D.A., Kominek, J., Zhou, X., Steenwyk, J.L., Buh, K. V., Haase,
M.A.B., Wisecaver, J.H., Wang, M., Doering, D.T., et al. (2018). Tempo and Mode of
Genome Evolution in the Budding Yeast Subphylum. Cell *175*, 1533-1545.e20.
https://doi.org/10.1016/j.cell.2018.10.023.

3.   Opulente, D.A., LaBella, A.L., Harrison, M.C., Wolters, J.F., Liu, C., Li, Y., Kominek, J.,
Steenwyk, J.L., Stoneman, H.R., VanDenAvond, J., et al. (2024). Genomic factors shape
carbon and nitrogen metabolic niche breadth across Saccharomycotina yeasts. Science
384, eadj4503. https://doi.org/10.1126/science.adj4503.

4.   Opulente, D.A., Rollinson, E.J., Bernick-Roehr, C., Hulfachor, A.B., Rokas, A., Kurtzman,
C.P., and Hittinger, C.T. (2018). Factors driving metabolic diversity in the budding yeast
subphylum. BMC Biol *16*, 26. https://doi.org/10.1186/s12915-018-0498-3.

5.   Aspinwall, M.J., Lowry, D.B., Taylor, S.H., Juenger, T.E., Hawkes, C. V., Johnson, M.V. V.,
Kiniry, J.R., and Fay, P.A. (2013). Genotypic variation in traits linked to climate and
aboveground productivity in a widespread C4 grass: Evidence for a functional trait
syndrome. New Phytologist *199*. https://doi.org/10.1111/nph.12341.

6.   Horianopoulos, L.C., Rokas, A., and Hittinger, C.T. (2025). Convergent evolution of
aerobic fermentation through divergent mechanisms acting on key shared glycolytic
genes. bioRxiv, 2025.10.02.679963. https://doi.org/10.1101/2025.10.02.679963.

7.   Chavez, C.M., Groenewald, M., Hulfachor, A.B., Kpurubu, G., Huerta, R., Hittinger, C.T.,
and Rokas, A. (2024). The cell morphological diversity of Saccharomycotina yeasts.
FEMS Yeast Res *24*. https://doi.org/10.1093/femsyr/foad055.

8.   Czarnolesk, M., and Verberk, W.C.E.P. (2025). Cell size matters: a unifying theory across
the tree of life. Trends Ecol Evol *0*. https://doi.org/10.1016/J.TREE.2025.09.003.

9.   Cadart, C., and Heald, R. (2022). Scaling of biosynthesis and metabolism with cell size.
https://doi.org/10.1091/mbc.E21-12-0627 *33*. https://doi.org/10.1091/MBC.E21-12-0627.

10.   David, K.T., Horianopoulos, L., Gonçalves, C., Steenwyk, J.L., Pontes, A., Gonçalves, P.,
Hittinger, C.T., Pennell, M., and Rokas, A. (2025). Discovery of additional ancient genome
duplications in yeasts. bioRxiv, 2025.08.31.673279.
https://doi.org/10.1101/2025.08.31.673279.

Supplemental methods

To investigate the role of shared evolutionary history in the variation of each trait, we calculated Pagel's lambda for each trait using the *R* package phytools[1] version 2.5-2 phylosig() with a set seed. Since these traits each had high phylogenetic signal values, we applied phylogenetically corrected regression models to determine their relationships using the *R* package nlme version 3.1-168 gls() function with model set to 'corPagel' from the package ape version 5.8-1.[2-4]

Supplemental references

1. Revell, L.J. (2024). phytools 2.0: an updated *R* ecosystem for phylogenetic comparative methods (and other things). PeerJ *12*. https://doi.org/10.7717/peerj.16505.

2. Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., and Van Willigen, B. (2020). Package nlme: Linear and Nonlinear Mixed Effects Models. *R* package.

3. Paradis, E., and Schliep, K. (2019). Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in *R*. Bioinformatics *35*. https://doi.org/10.1093/bioinformatics/bty633.

4. Pinheiro, J.C., and Bates, D.M. (2000). Mixed-Effects Models in S and S-PLUS (Statistics and Computing) (Springer Science & Business Media).