1   **Integrating migratory marine connectivity into shark conservation**
2
3   **Abstract**
4

Understanding migratory connectivity is important for the conservation of highly
mobile marine species that face escalating threats across the globe. Establishing baseline
information on migratory connectivity is therefore needed to identify regions of
conservation focus. Despite efforts to track migratory sharks and rays, information on
transboundary movements is limited and often inaccessible to managers and
policymakers. Here, we synthesised multimethod movement data for migratory
Australian shark and ray species, investigating which species require international
engagement to support their population recovery. Based on data from a systematic
literature review, we built connectivity networks from telemetry and mark-recapture
studies that provide a first baseline for transboundary migratory connectivity for
Australian sharks and rays. Of the 31 shark and ray species reviewed, we identified 6
species that link the Australian Exclusive Economic Zone to other national jurisdictions
via multispecies migratory connections through the Tasman Sea to New Zealand,
through the Tasman and Coral Sea to New Caledonia, and north across the Timor Sea
and Torres Strait to Indonesia and Papua New Guinea. White sharks (*Carcharhinus
carcharias*) and whale sharks (*Rhincodon typus*) were the most data rich, whereas 14
shark and ray species had no movement information. There is a grave deficiency in
available information for endangered or critically endangered migratory shark and ray
populations, with 76% having only one or no published studies. This work supports
future conservation strategies for migratory sharks that require robust international
collaboration and the adoption of integrated and dynamic management approaches.
26
27   **Keywords:** Chondrichthyans; elasmobranchs; satellite telemetry; acoustic telemetry;
28   biogeography; movement ecology; photo-identification

Introduction

Migration, the broad-scale movement of populations (Wiens 1997), can transcend ecosystem, national, economic, and cultural boundaries (Harrison et al. 2018; McLean et al. 2023). By connecting distant habitats, migrations contribute to ecosystem function and resilience (Alerstam & Bäckman 2018; Davidson et al. 2020). While migrations can facilitate access to seasonally abundant resources required for key biological processes (Alerstam et al. 2003), they also expose animals to a variety of threats and uncoordinated governance regimes (Dunn et al. 2019; Roberson et al. 2021). This often results in population declines (Teitelbaum & Mueller 2019; Kubelka et al. 2022). Nearly half of all migratory species are in decline, and 20% are at risk of extinction (UNEP-WCMC 2024). Improved management of migratory species is thus urgently needed.

Migratory sharks and rays often exhibit breeding-site philopatry, in which they return to specific regions for parturition or mating after long-distance migrations (Dudgeon et al. 2013; Chapman et al. 2015). These movements link biologically important areas that often span national boundaries, either between adjacent countries or between distant regions connected by extensive transnational migrations (Harrison et al. 2018; Thorburn et al. 2024). This phenomenon, of the geographic linking of individuals and populations across their migratory cycles, is known as migratory connectivity (Webster et al. 2002). Understanding migratory connectivity provides insights into the various stressors an animal encounters during its migration, and how these stressors may accumulate and shape population structure, genetic diversity, and survival rates (Heupel et al. 2015; Somveille et al. 2021; Flack et al. 2022). Therefore, enhancing our knowledge of migratory connectivity underpins the development of effective conservation and management practices aimed at mitigating the decline of migratory species. This is particularly true for elasmobranchs (sharks and rays) – many of which are both highly migratory and under unprecedented threat (Pacoureau et al. 2021; Juan-Jordá et al. 2022), with many populations in decline (Roff et al. 2018; Dulvy et al. 2021; Yan et al. 2021; Simpfendorfer et al. 2023). Globally, approximately one-third of sharks and rays are threatened with extinction (Dulvy et al. 2021). Oceanic populations are more threatened compared to species in other habitats and have declined by 71% over recent decades (Pacoureau et al. 2021).

The Indo-West Pacific, and Australia in particular, is a global biodiversity hotspot for sharks and rays (Pimiento et al. 2023). Australia is frequently cited as a high-priority country for conservation due to its diversity, species richness and the threatened status of its sharks (Dulvy et al. 2017; Stein et al. 2018; Neubauer et al. 2024). Many of these species are dependent on specific habitats in disparate locations and often migrate both within the Australian Exclusive Economic Zone (EEZ) (Stevens et al. 2010; Heupel et al. 2015) and beyond (Chin et al. 2017; McMillan et al. 2019).

Within Australian waters, the Environment Protection and Biodiversity Conservation (EPBC) Act of 1999 is the primary environmental legislation designed to protect and manage flora, fauna, and ecological communities. International cooperation on the conservation of migratory species is coordinated through bilateral and multilateral agreements, and more broadly via the Convention on the Conservation of Migratory Species of Wild Animals (CMS). CMS Appendix I requires binding protection measures with limited scope for exceptions, while Appendix II, where most sharks and rays are listed, only requires cooperation from signatories on management

(Lawson & Fordham 2018). The EPBC Act lists eight shark and ray species as
threatened, including the whale shark, white shark, grey nurse shark, two river shark
species, three sawfish species and Maugean skate (*Zearaja maugean*). The Act further
lists 17 species of sharks and rays from the CMS Appendices as migratory, but does not
include 12 other CMS-listed species because Australia took a formal exception to those
species being listed under CMS (and thus is not compelled to address them as it does
other CMS-listed species). While the EPBC Act provides the overarching environmental
protection for threatened sharks and rays in Australia, managing their major threat
(fishing), including sustainable harvesting and bycatch regulation, is undertaken by state
agencies and the Australian Fisheries Management Authority at a national level, and
through regional fisheries management organisations (RFMOs) for species caught in
shared-stock fisheries in areas beyond national jurisdictions (ABNJ).
92
92
Listings of species under CMS and the EPBC Act, and management of sharks and
rays by relevant fisheries organisations, are dependent on information on the movement
of animals and status of populations being monitored and provided to management and
policy fora in an accessible and actionable manner. However, understanding
elasmobranch movement at large scales is hampered by major sampling obstacles.
Conventional mark-recapture and photo-identification studies are cost-effective and
often have large samples of tagged/sighted animals, but the likelihood of recapturing
tagged individuals or resighting identifiable individuals is low, and often biased towards
highly populated areas and areas of high tourism (Kohler & Turner 2001; Rowat et al.
2009; Mas et al. 2022). Acoustic telemetry can capture local movements, particularly in
estuarine or reef ecosystems, but detection probabilities are low across vast areas
(Bradford et al. 2019; Spaet et al. 2020). When migratory locations or corridors are
unknown, or when species are highly pelagic and travel far from the coastline where
receivers are placed, it is usually necessary to undertake expensive satellite tracking or
low-resolution archival pop-up light-level tags to capture these movements (Jaine et al.
2014). Despite transnational collaborations expanding acoustic receiver networks, mark-
recapture, satellite, and archival pop-up tags remain the main sources of information on
large-scale connectivity for elasmobranchs. Thus, difficulties in collecting data on
elasmobranch movements persist and have limited our understanding of migration
pathways and areas of importance for various life-history stages, which impedes their
management and conservation both nationally and internationally. These obstacles to
data collection make the available data all the more valuable and increases the
importance of aggregating and making use of all available information on shark and ray
connectivity.
117
Here we review and quantify published information on migratory connectivity for
shark and ray species listed under the CMS to aggregate evidence of migratory
behaviour and transboundary movement requiring international collaboration, and to
support more effective management. We build on the Action Plan for Australian Sharks
and Rays 2021 that reviewed the extinction risk of all 328 Australian species (Kyne et
al. 2021). We conducted a data synthesis of over 30 years of literature on biotelemetry,
mark-recapture, and photo-identification studies to establish the first baseline for
migratory connectivity of globally at-risk sharks and rays in Australian waters. We also
produced network models for migratory shark and ray species, compiling movement
information to reveal species-level migratory pathways (Bentley et al. 2024). Our aim is
to establish an initial baseline of migratory connectivity for Australian sharks and rays
to guide their effective conservation and management. We hope that this will support

Australia's commitments and reporting obligations under international conservation
agreements, including the CMS, and the Convention on Biological Diversity. This
baseline will determine areas supporting migrations of multiple species, highlight
regional and taxonomic data gaps, and help identify nations whose cooperation is
needed to improve the conservation and management of individual species.
135
136 **Methods**
137
138 *Species selection criteria*
139
Species selection was based on sharks and rays listed under CMS Appendices I and
II, and the Memorandum of Understanding on the Conservation of Migratory Sharks
(Shark MoU) (Table 1). The spiny dogfish (*Squalus acanthias*) was not included in our
study because only the northern hemisphere populations are listed under CMS. Listing
of species under CMS and the Shark MoU is not a systematic process and there are
likely to be many species that have not been included yet due to lack of capacity by a
CMS member state to nominate them, or lack of available data. We included two further
species due to their known large-scale movements and the many individuals tagged in
Australasia; the tiger shark (*Galeocerdo cuvier*) and the bull shark (*Carcharhinus
leucas*), leading to a potential total of 31 species included in this review.
150
151 *Systematic Literature Review Protocol*
152
We followed the PRISMA (Preferred Reporting Items for Systematic Reviews and
Meta-Analyses) framework to select and screen peer-reviewed literature for this study
(Moher et al. 2009). The PRISMA framework outlines a series of steps, beginning with
the identification of relevant studies through comprehensive database searches.
Subsequently, references meeting inclusion criteria undergo screening and eligibility
assessment based on predefined criteria (here, studies with a migratory connection with
Australia). Data are extracted only from selected references (Moher et al. 2009). For this
study, we compiled peer-reviewed studies that used biotelemetry, mark-recapture, and
photo-identification methods or derived products (e.g., kernel density estimates or state
space models) to describe the movements of the 31 candidate species.
163
All searches for the collection of references were conducted between June and
November 2023. We tested the relevance of search strings for a subset of shark and ray
species using the Web of Science and Scopus abstract and citation databases (Appendix
1). We noted whether relevant reference ("control") literature was captured, the total
number of references, and how many of the initial papers identified would meet the
screening criteria. Within the Web of Science database, all search strings were used to
conduct topic searches in English for indexed literature within subscribed databases
(WOS, BCI, CCC, DRCI, DIIDW, KJD, MEDLINE, RSCI, SCIELO, ZOOREC),
published until June 2023. Within Scopus, all search strings were used to search by
keyword, title, and subject area.
174
175 *Site aggregations and network models*
176
To determine the connectivity of migratory sharks and rays, the information from
the systematic literature review was aggregated into network models following the
methods used to develop the MiCO system (www.mico.eco/system; Bentley et al. 2024,

Dunn et al. 2019). The general approach is to use network models to aggregate multiple
data types describing connectivity. In this approach, individuals reviewing the literature
identified areas used by a tagged/visually identified animal (hereafter "sites"), along
with "connections" to other sites. The size, period, behaviour, number of animals, sex,
and age-class were recorded for each site and connection, if available. The methodology
is meant to support description of regional connectivity and sites are categorised into
three coarse size classes having 1°, 5° or 10° radii. If no specific animal behaviour was
described in the reference reviewed, it was listed as an "observation".

To generate network models from the many sites and connections identified across
various studies, we aggregated overlapping sites into "metasites" (sites made up of
multiple sites) and removed studies that appeared to use data employed in other studies.
The aggregation and data deduplication process followed Bentley et al. (2024), and has
the same four objectives: 1) Integrate data from multiple sites that capture species
activity in overlapping locations; 2) Preserve information on large-scale connections
(>500 km) resulting from animal movements; 3) Retain high-resolution data on critical
reproductive life history stages; and 4) Ensure a minimum sample size of known
individuals both within and between sites.

The process of generating network models for shark species entailed five key
stages: data preparation, site selection, metasite centroid definition, data duplication
checks, and metasite aggregation. In the data preparation stage, sites for each species are
compiled and organised based on behaviour, such as reproduction or feeding, as well as
factors such as size and number of nearby sites. The ordering of sites ensures that the
resolution of reproductive sites is maintained, but that overlapping foraging areas are
merged to simplify networks. Subsequently, during site selection, sites are iteratively
chosen (i.e., the script examines one at a time based on the ordering) to identify
overlapping sites. The metasite centroid definition stage iteratively calculates a new
centroid for sites to be aggregated, and then removes any sites whose centroid is no
longer within the radii of the new metasite. The centroid is repeatedly recalculated
whenever a site is dropped until no further sites are dropped from the metasite. To
maintain long distance connections, any sites that have a connection of >500 km were
not aggregated. The data duplication stage ensures the integrity of the dataset by
checking for similarities such as identical coordinates, route connectivity, authorship,
and temporal scope. Duplicate sites informed the location of a centroid of the new
metasite but were removed from calculations of the minimum numbers of individuals in
a given metasite in the final metasite aggregation stage. Once all sites had been
aggregated to a metasite, connections between sites were assigned to their respective
metasites, resulting in metaconnections (a connection made up of connections). All code
used to develop the network models is available on (see Cover Page below).
.
We exported the metasites and metaconnections from the MiCO system
(www.mico.eco/system), where they are freely available to the public. All subsequent
analyses were undertaken in R version 4.3.0, using the packages tidyverse, circlize,
ggOceanMaps, jsonlite, and ggExtra (Wickham et al. 2019; Attali & Baker 2022;
RTeam et al. 2023). The geodatabase files from the system were used to develop maps
of migratory connectivity for all shark and ray species where sufficient information was
available. Metasites were also allocated to an EEZ based on the latitude and longitude
of the metasite centroid. Adjacency matrices were then constructed from the data on

metaconnections that crossed EEZs and were used to generate chord diagrams
describing regional connectivity.
229
230 **Results**
231
Our data synthesis of 31 shark and ray species that use Australian waters identified
91 relevant peer-reviewed references for 18 species (Table 1, S1). Across these species,
we identified 162 metasites and 193 metaconnections based on individual movement
data (Table 2). Satellite and acoustic telemetry methods were the primary sources of
detailed movement data, with photo-identification and mark-recapture methods
providing a minority of evidence. The number of sites and routes identified rose
proportionally to tagging efforts and were heavily biased towards iconic species,
including white sharks, whale sharks, shortfin mako, tiger sharks and reef manta rays,
which accounted for approximately 70% of the sampled individuals.
241
Thirteen shark and ray species in our review had no published movement
information, and another eight had only one relevant publication, which provided
movement data specifically within Australian waters (Table 1). Among the 22
endangered species in this study (i.e., those classified as Endangered or Critically
Endangered globally by the IUCN Red List; Table 1), 55% (n = 12) had no published
movement information. Another five endangered species (23%) had only one published
movement study, primarily focused on fine-scale movements in estuaries. Despite the
number of data deficient species, our study identified 172 individual sharks of 6 species
that engaged in migratory movements either into or out of Australian waters to seven
other national jurisdictions and ABNJ (Figure 2).
252
253 *International transboundary movement of Australian sharks*
254
While we did not find any instances in the published literature of any ray species
crossing an international border with Australia, the six migratory shark species in
Australian waters that exhibited international transboundary movements were shortfin
mako (*Isurus oxyrinchus*), white sharks (*Carcharodon carcharias*), tiger sharks
(*Galeocerdo cuvier*), whale sharks (*Rhincodon typus*), blue sharks (*Prionace glauca*)
and school sharks (*Galeorhinus galeus*). These species demonstrated complex patterns
of habitat use, moving across various habitats, from coastal areas to oceanic habitats,
and moving through numerous governance regimes. New Caledonia had the highest
number of species with connections shared with Australia, which included the shortfin
mako, blue shark, white shark, and tiger shark. White sharks and tiger sharks were the
most connected species, with movements linking multiple jurisdictions: New Zealand,
New Caledonia, Papua New Guinea, Indonesia, South Africa and Solomon Islands. A
total of 34 individual sharks of various species moved between Australia and New
Zealand, indicating strong trans-Tasman migratory connectivity.
269
<u>White sharks</u>
For the white sharks, strong white shark connectivity was found between eastern
Australia and New Zealand, with 18 individuals migrating to New Zealand and 14
individuals migrating to Australia. White sharks also showed transboundary
connectivity between Australia and Papua New Guinea, New Caledonia, and South
Africa (Figure 2). Our network models for white shark migratory connectivity show

broad-scale movements along the Australian east and west coasts (Figure 3a), and, in general, wide-ranging longitudinal and latitudinal movements.

Shortfin mako

This species demonstrated extensive and diverse movements: some migrated to New Zealand and New Caledonia, while one individual travelled from the Bonney Upwelling Region in southern Australia to the Southwest Indian Ridge in ABNJ. Extensive movements within Australian waters were observed.

Tiger sharks

This species frequently traversed coral reef and seagrass ecosystems in coastal environments and demonstrated wide-ranging migratory movements, extending into the Coral Sea region and ABNJ, and crossing multiple national jurisdictions including Papua New Guinea and New Caledonia (Figure 3c).

Whale sharks

Published studies on whale shark movements centred around the coastal hotspot of Ningaloo National Marine Park in Western Australia, where they aggregate seasonally to feed (Figure 3d). There were migratory movements into ABNJ of the Indian Ocean, with 27 individuals that exhibited transboundary movements between Australia and Indonesia, Papua New Guinea and overseas Australian territories. Additionally, individuals were recorded migrating from Ningaloo Reef and the Great Barrier Reef to Micronesia.

Blue sharks and school sharks

Blue sharks were observed migrating to Indonesia, New Caledonia and ABNJ (Figure 3e). School sharks made extensive long-distance partial migrations using protected pupping areas along the south coast of Australia, with documented movements reaching New Zealand recorded by mark-recapture, acoustic and satellite telemetry methods (Figure 3f).

Large-scale movements within Australian waters

Movements of acoustically tagged bull sharks (n=73) were restricted to euryhaline and coastal ecosystems, extending from far North Cape York Peninsula to Tollgate Islands in Southeast Australia, and were not recorded moving into other national jurisdictions or ABNJ. Reef manta rays along the Great Barrier Reef in Eastern Australia to northern New South Wales and Ningaloo Marine Park in Western Australia are also well-studied and exhibited well-defined migration patterns that remain within Australian national waters. These populations engaged in seasonal movements between northern and southern reef sections (Figure S1b). Despite these insights, tracking data for other Mobulidae remains limited, with no data on their broader oceanic connectivity and no examples of transboundary movements beyond Australian waters in peer-reviewed literature. A total 473 (out of n = 2478) dusky sharks migrated between western and southern Australian waters, and eight individuals displayed shorter-range movements between Queensland and New South Wales**.** Whale sharks cross ABNJ to other Australian remote oceanic islands, such as Christmas Island and Cocos Keeling Island, distant from the mainland EEZ.

Multi-species connections

We identified several key transboundary connections that were important for multiple species. Firstly, the Trans-Tasman pathway represents a key migratory connection for several sharks (white shark, shortfin mako, and school shark). These species migrate across the Great Australian Bight and Bass Strait, extending their migratory pathways through the Tasman Sea to New Zealand. Secondly, shortfin mako, white sharks, and blue sharks shared similar connections extending from the trans-Tasman region to New Caledonia (Figure 3). In addition to these species, tiger sharks exhibited longitudinal migratory pathways further north, transiting the Coral Sea to New Caledonia. Lastly, an area of multispecies migratory connections was found in northern Australia, particularly between Western Australia and Indonesia across the Timor Sea. Both tiger sharks and whale sharks migrated throughout the Timor Sea, travelling from Ningaloo Reef to Indonesia. Moreover, there were connections from waters south of Australia to Indonesia for both shortfin mako and blue sharks.

Discussion

We provide a baseline for the movement of globally at-risk migratory shark species to support conservation policy in the Indo-West Pacific. Among the nine species requiring international engagement to manage their populations mentioned in The Action Plan for Australian Sharks and Rays (Kyne et al. 2021), only whale sharks and shortfin mako have sufficient data to develop migratory connectivity networks. By synthesising information from 91 published references with movement data from ~2,900 animals, we identified transboundary movements for six species: white sharks, whale sharks, shortfin mako, blue sharks, school sharks, and tiger sharks. These species visited the EEZs of seven countries or territories, as well as ABNJ. The regional and global scales of these shark migrations necessitate an international collaborative approach to their conservation and management.

Identification of multispecies migratory connections

The trans-Tasman migratory pathway was identified as important for multiple species in this review. Our network models align with our knowledge of genetic structuring in both white and tiger shark populations across the Tasman Sea and broader West Pacific (Holmes et al. 2017; Bradford et al. 2019). However, genetic information cannot confirm contemporary connectivity or inform spatial management to reduce mortality during migrations, underscoring the importance of collecting individual-based animal movement data and synthesising connectivity data to highlight shared migratory pathways (Figure 3) (Devloo-Delva et al. 2019; McMillan et al. 2019). Multiple species also continued from the Tasman Sea through to New Caledonia. The region has been recognised under the Important Shark and Ray Areas (ISRAs) Subcriterion C4 as a critical movement area linking Norfolk Island and New Caledonia (Jabado et al. 2024). The number of shark species and individuals migrating across the Tasman Sea suggests the need for strengthened bilateral management between Australia and New Zealand.

Multispecies use of the waters from northern Australian through the Timor Sea was also evident. Despite the presence of the Timor Trench, which is up to 3300 m deep and limits the dispersal and connectivity of shark populations (Hirschfeld et al. 2021), both tiger sharks and whale sharks migrated from Ningaloo Reef to Indonesia. Further

research could elucidate the degree that the Timor Trench may act as a barrier for
movements of coastal sharks, particularly in comparison to the Torres Strait, which may
represent a migratory corridor for species that prefer shallow waters but act as a barrier
for deep-dwelling species. However, data on movement behaviour in the Timor Sea and
Torres Strait are scarce. Although genetic data have provided valuable insights into
trans-national elasmobranch populations across the Torres Strait (Chin et al. 2017),
tracking data remain limited. There is potential for the movement of multiple species
across the strait, as indicated by the distribution, geographical proximity between
populations, and genetic studies, such as the critically endangered largetooth sawfish
(*Pristis pristis*) (Feutry et al. 2015). Prioritising data collection for poorly studied
species in this region is imperative to better understand connectivity, given that
Indonesia is one of the world's largest shark-fishing nations (Sherman et al. 2023).
Given the narrow and shallow nature of the Torres Strait, we recommend strategic
placement of acoustic receivers throughout this region to capture the potential
movement of shallow-obligate sharks and rays moving between Australia and Papua
New Guinea. While this approach undertaken in consultation with Traditional Owners
and other local stakeholders should work in that region, it is less suitable for the deeper
and larger Timor Sea nor for the open ocean areas in the West Pacific. For those regions
we recommend deploying satellite and archival tag technology on species caught and
retained by fisheries managed through the Western & Central Pacific Fisheries
Commission (WCPFC).
395
We found reduced evidence for multispecies connections with Melanesia and
islands east of New Caledonia, though this may be due to lower sampling effort in these
regions rather than an actual lack of connectivity. Whale sharks were identified to move
from southwest of Papua New Guinea to Kirabati, indicating that while no connection
directly from Australian waters was identified, there are movements from the region to
the Central Pacific (Womersley et al. 2022). The lack of information on blue shark
movement may also be concealing larger-scale movement in this region. Although there
were few references in the literature for blue shark in Australian waters, there is
extensive information on their ontogenetic habitat shifts across large spatial scales in
multiple ocean basins (Maxwell et al. 2019; Fujinami et al. 2021; Madigan et al. 2021;
Elliott et al. 2022; Poisson et al. 2024). Similarly, there is a lack of understanding of
shortfin mako movements in the region, leading the WCPFC to describe the degree of
connectivity as "unclear" and to suggest that the population may comprise
"geographically relatively distinct stocks or migratory contingents" (Large et al. 2022).
Further research, particularly tracking studies, could address limitations of stock
assessments and may provide a more comprehensive understanding of these species'
distributions.
413
414 *Transboundary management of migratory sharks*
415
Regional baselines are useful tools for understanding the international
transboundary movements of migratory species and, consequently, shared responsibility
for the management of migratory populations (Harrison et al. 2018). One existing
pathway for multinational collaboration on the conservation of migratory sharks is the
CMS and the Sharks Memorandum of Understanding (MoU) that sits beneath it. Of the
countries connected by shark movements identified in our study, Australia, New
Zealand, New Caledonia (France), and South Africa are signatories to the CMS and the
Shark MoU, while Indonesia, Papua New Guinea, the Federated States of Micronesia,

and the Solomon Islands are not, nor are they signatories to the Shark MoU. An
important step toward better governance of migratory sharks in the region would be for
all countries to develop mechanisms to cooperate (e.g., becoming parties to CMS or
signatories to the Shark MOU) or establish bilateral agreements where they do not exist.
This could then provide political momentum for greater support of research into
regional connectivity for sharks and rays. Although the CMS signatory status alone
cannot fully address governance gaps for migratory sharks and rays, it plays a
constructive role in driving efforts to implement agreed-upon conservation measures.
432
Effective transboundary management of migratory sharks requires robust data
sharing, collaborative research, and policy alignment among nations. Comprehensive
data on shark movements, population dynamics, and habitat use are vital to inform
management decisions. However, existing research shows a marked bias, with most data
focusing on iconic species in wealthier countries, such as Australia, New Zealand, and
New Caledonia (France). In contrast, less-studied species in poorer countries often lack
sufficient data and have weaker management frameworks. To support the
implementation of CMS agreements, it is crucial to prioritise research on the 22 shark
and ray species listed under the CMS, many of which currently have few or no
connectivity studies in the Indo-West Pacific. With its substantial research capacity and
economic resources in the region, Australia is well-placed to lead efforts in addressing
these disparities by generating critical data domestically and supporting research
initiatives in neighbouring countries with limited resources and capacity. The Australian
government could focus on the seventeen shark and ray species listed under the EPBC
Act as migratory. Particular attention could be paid to the three sawfish species listed as
Threatened under the Act that had only four connectivity studies among them and to
hammerhead species. This compares with the other two sharks listed as threatened
(white sharks and whale sharks), for which there were 38 studies. An emphasis towards
studies that could clarify movement across the Timor Sea and the Torres Strait would
also reduce geographic bias towards wealthier regions.
453
RFMOs play a central role in monitoring and managing the sustainability of
fisheries across vast oceanic areas. They implement a variety of measures to mitigate
threats to these species, such as setting catch limits, enacting fishery closures, enforcing
gear restrictions to reduce bycatch, and designating "no retention" species to prohibit
their landing. All West Pacific countries for which transboundary connections with
Australia were identified in this study are members of the WCPFC. The WCPFC has a
Conservation and Management Measure for Sharks that specifically prohibits: shark
finning; retention of whale sharks, oceanic whitetip sharks or silky sharks; the use of
shark lines in longlining; and the setting of purse seines on tuna associated with whale
sharks (Neubauer et al. 2024). In addition to the Conservation and Management
Measure for Sharks, the WCPFC also has a 2021-2030 Shark Research Plan that
provides a timeline for assessment of various shark species and a list of prioritised
projects (Brouwer & Hamer 2024). Of interest, the Shark Research Plan includes a
scoping study to understand stock structure and natal homing across all shark species.
Similarly, to address uncertainty in the stock assessment for silky sharks, there is a call
for more satellite tagging studies to be undertaken to "resolve fundamental questions…
about the degree of natal homing and limited mixing of the stock" (Neubauer et al.
2024). For the same reasons, the stock assessment for shortfin mako recommends
"additional tagging should be carried out… especially [in] known nursery grounds off
southeast Australia and New Zealand, as well as high seas areas to the north and east of

New Zealand, where catch-rates are high." The baseline connectivity models presented
here can enhance the understanding of metapopulation dynamics critical to RFMO
management, as well as underpin future risk-assessments by allowing cumulative
impacts to be accounted for across the known migratory movements of these species.
They also provide a framework for how information from various sampling methods
can be synthesised to describe regional connectivity for sharks and rays.
480
It is important to note that while, in this study, we use the term "transboundary" to
refer to movements that cross international borders, animals moving across multiple
management jurisdictions at any scale face similar challenges. For example, bull sharks
demonstrate large movements across Australian state boundaries. As bull shark
management is inconsistent across Australian states and territories, these movements are
a microcosm of the broader challenge of effectively managing highly migratory species.
Further, the scale of these movements could easily have resulted in transboundary
movement in regions with more or smaller EEZs. Indeed, while the published
movement data for bull sharks and reef manta rays did not show evidence of
international transboundary movements, this may be influenced by the limited use of
satellite telemetry for bull sharks and/or the predominantly coastal behaviour in reef
manta rays (confirmed through aggregated photo-identification data (Armstrong et al.
2019)).
494
495 *Using network models to understand migratory connectivity*
496
While information on migratory connectivity can be gleaned from many sampling
methods, findings are rarely integrated (Bentley et al. 2024). Our approach identifies
sites and connections across methods and integrates them into a single network of the
minimum known connectivity for each species. For example, acoustic telemetry data
were previously used to describe school shark movement in national waters throughout
southern New South Wales, Victoria and eastern Tasmania (Lédée et al. 2021). Our
network connectivity models, incorporating satellite telemetry studies, extend these
findings by describing transboundary migrations from South Australia to New Zealand
waters, and are corroborated by a vast amount of mark-recapture tag data (Hurst et al.
1999). This supports the recent listing of school sharks (also known as tope) under
Appendix II of the CMS. Establishing a benchmark for migratory connectivity
facilitates international collaboration and supports the identification of interested
parties, from national stakeholders to intergovernmental organisations, with
management mandates related to sharks and rays. These benchmarks also identify the
regions and species for which data gaps persist, providing evidence to guide the
prioritisation of future research efforts. Benchmarks for well-studied species (e.g., the
white shark in southeastern Australia) also demonstrate the power and feasibility of this
method. The approach adopted here offers a transferable model to help develop
baselines for migratory connectivity in sharks and rays in other regions, which is critical
to conserving these threatened taxa.
517
Some caveats should also be considered when interpreting our results. Network
models demonstrate movement patterns from known data, but also reflect biases in the
location of tagging or acoustic receivers, which can potentially skew results. Our results
mostly relied on satellite telemetry data and passive acoustic telemetry studies, methods
that are commonly deployed in coastal regions (Hussey et al. 2015). Another key
limitation is that network models do not capture the precise routes individuals take

between nodes. Connections between nodes represent observed migratory links, not a precise movement path. As such, the models provide a simplified representation of complex migratory behaviours and cannot readily inform spatial management of migratory corridors without further interrogation of the underlying data.

It is important to acknowledge that unpublished data, including telemetry studies not yet available in the public domain, were not included here but could provide additional insights into migratory patterns and further inform our understanding of species connectivity. Many research efforts are never published, or are published in un-indexed grey literature that was not reviewed in this study, potentially resulting in apparent "gaps" in which studies have indeed been undertaken. As such, this study should be viewed as an assessment of the minimum known transboundary connectivity for Australian sharks and rays, and further work should be undertaken to incorporate any available information on shark and ray movements. Together with improvements in spatial resolution and gap-filling to address geographic biases, these measures could offer a more comprehensive view of shark and ray migratory connectivity.

Further tagging efforts are needed to address the significant data gaps that persist within Australia and neighbouring countries. Critically, given the transboundary nature of many shark and ray migrations and distributions, regional funding to raise capacity across all neighbouring countries is essential to support Australian shark and ray populations and their important habitats. Our synthesis of available evidence identifies several key opportunities for international partnerships to strengthen migratory sharks and ray conservation in the Indo-West Pacific. Of particular interest are multispecies migratory connections between countries shared by multiple shark and ray species, such as Australia and Indonesia, Papua New Guinea, and New Zealand. Aggregation of such shared connections in unified models of migratory connectivity could help foster the cooperative transboundary management required to sustain shark and ray populations.

References

Alerstam T, Bäckman J. 2018. Ecology of animal migration. Current Biology.

Alerstam T, Hedenström A, Åkesson S. 2003. Long-distance migration: Evolution and determinants. Oikos.

Armstrong AO, Armstrong AJ, Bennett MB, Richardson AJ, Townsend KA, Dudgeon CL. 2019. Photographic identification and citizen science combine to reveal long distance movements of individual reef manta rays *Mobula alfredi* along Australia's east coast. Marine Biodiversity Records **12**.

Attali D, Baker C. 2022. ggExtra: Add Marginal Histograms to "ggplot2", and More "ggplot2" Enhancements. The R Journal.

Bentley L et al. 2024, June 12. Marine megavertebrate migrations connect the global oceans. Available from https://www.researchsquare.com/article/rs-4457815/v1.

Bradford BD, Harasti D, Lee K, Gallen C, Bradford R. 2019. Broad-scale movements of juvenile white sharks *Carcharodon carcharias* in eastern Australia from acoustic and satellite telemetry. Marine Ecology Progress Series **619**.

Brouwer S, Hamer P. 2024. The Commission for the Conservation and Management of Highly Migratory Fish Stocks in the Western and Central Pacific Ocean. Scientific Committee Twentieth Regular Session. WCPFC-SC20-2024:1–251.

Chapman DD, Feldheim KA, Papastamatiou YP, Hueter RE. 2015. There and back
again: A review of residency and return migrations in sharks, with implications for
population structure and management. Annual Review of Marine Science **7**.
Chin A, Simpfendorfer CA, White WT, Johnson GJ, McAuley RB, Heupel MR. 2017.
Crossing lines: A multidisciplinary framework for assessing connectivity of
hammerhead sharks across jurisdictional boundaries. Scientific Reports **7**.
Davidson SC et al. 2020. Ecological insights from three decades of animal movement
tracking across a changing Arctic. Science **370**.
Dudgeon CL, Lanyon JM, Semmens JM. 2013. Seasonality and site fidelity of the zebra
shark, *Stegostoma fasciatum*, in southeast Queensland, Australia. Animal
Behaviour **85**:471–481.
Dulvy NK et al. 2021. Overfishing drives over one-third of all sharks and rays toward a
global extinction crisis. Current Biology **31**.
Dulvy NK, Simpfendorfer CA, Davidson LNK, Fordham S V., Bräutigam A, Sant G,
Welch DJ. 2017. Challenges and Priorities in Shark and Ray Conservation.
Dunn DC et al. 2019. The importance of migratory connectivity for global ocean policy.
Proceedings of the Royal Society B: Biological Sciences **286**.
Elliott RG, Montgomery JC, Penna A Della, Radford CA. 2022. Satellite tags describe
movement and diving behaviour of blue sharks *Prionace glauca* in the southwest
Pacific. Marine Ecology Progress Series **689**:77–94.
Feutry P, Kyne PM, Pillans RD, Chen X, Marthick JR, Morgan DL, Grewe PM. 2015.
Whole mitogenome sequencing refines population structure of the Critically
Endangered sawfish *Pristis pristis*. Marine Ecology Progress Series **533**.
Flack A et al. 2022. New frontiers in bird migration research.
Fowler S. 2014. The Conservation Status of Migratory Sharks. UNEP/CMS Secretariat,
Memorandum of Understanding on the Conservation of Migratory Sharks.
Fujinami Y, Shiozaki K, Hiraoka Y, Semba Y, Ohshimo S, Kai M. 2021. Seasonal
migrations of pregnant blue sharks *Prionace glauca* in the northwestern Pacific.
Marine Ecology Progress Series **658**.
Harrison AL et al. 2018. The political biogeography of migratory marine predators.
Nature Ecology and Evolution **2**.
Heupel MR, Simpfendorfer CA, Espinoza M, Smoothey AF, Tobin A, Peddemors V.
2015. Conservation challenges of sharks with continental scale migrations.
Frontiers in Marine Science **2**.
Hirschfeld M, Dudgeon C, Sheaves M, Barnett A. 2021. Barriers in a sea of
elasmobranchs: From fishing for populations to testing hypotheses in population
genetics.
Holmes BJ, Williams SM, Otway NM, Nielsen EE, Maher SL, Bennett MB, Ovenden
JR. 2017. Population structure and connectivity of tiger sharks (*Galeocerdo cuvier*)
across the Indo-Pacific Ocean basin. Royal Society Open Science **4**.
Hurst RJ, Baglet NW, McGregor GA, Francis MP. 1999. Movements of the New
Zealand school shark, *Galeorhinus galeus*, from tag returns. New Zealand Journal
of Marine and Freshwater Research **33**:29–48.
Hussey NE et al. 2015. Aquatic animal telemetry: A panoramic window into the
underwater world. Science **348**.
Jaine FRA, Rohner CA, Weeks SJ, Couturier LIE, Bennett MB, Townsend KA,
Richardson AJ. 2014. Movements and habitat use of reef manta rays off eastern
Australia: Offshore excursions, deep diving and eddy affinity revealed by satellite
telemetry. Marine Ecology Progress Series **510**.

Juan-Jordá MJ, Murua H, Arrizabalaga H, Merino G, Pacoureau N, Dulvy NK. 2022.
Seventy years of tunas, billfishes, and sharks as sentinels of global ocean health.
Science **378**.
Kohler NE, Turner PA. 2001. Shark tagging: A review of conventional methods and
studies.
Kubelka V, Sandercock BK, Székely T, Freckleton RP. 2022. Animal migration to
northern latitudes: environmental changes and increasing threats.
Kyne PM, Heupel MR, White WT, Simpfendorfer C. 2021. The Action Plan for
Australian Sharks and Rays 2021. National Environmental Research Program
Marine Biodiversity Hub. **1 ed.**
Large K, Neubauer P, Brouwer S. 2022. SCIENTIFIC COMMITTEE EIGHTTEENTH
REGULAR SESSION ELECTRONIC MEETING Stock assessment of Southwest
Pacific Shortfin Mako shark Dragonfly Data Science 2 Saggitus Consulting Stock
assessment of Southwest Pacific Shortfin Mako shark.
Lédée EJI et al. 2021. Continental-scale acoustic telemetry and network analysis reveal
new insights into stock structure. Fish and Fisheries **22**.
Madigan DJ, Shipley ON, Carlisle AB, Dewar H, Snodgrass OE, Hussey NE. 2021.
Isotopic Tracers Suggest Limited Trans-Oceanic Movements and Regional
Residency in North Pacific Blue Sharks (*Prionace glauca*). Frontiers in Marine
Science **8**.
Mas F, Cortés E, Coelho R, Defeo O, Forselledo R, Jiménez S, Miller P, Domingo A.
2022. Shedding rates and retention performance of conventional dart tags in large
pelagic sharks: Insights from a double-tagging experiment on blue shark (*Prionace*
glauca). Fisheries Research **255**.
Maxwell SM, Scales KL, Bograd SJ, Briscoe DK, Dewar H, Hazen EL, Lewison RL,
Welch H, Crowder LB. 2019. Seasonal spatial segregation in blue sharks (*Prionace*
glauca) by sex and size class in the Northeast Pacific Ocean. Diversity and
Distributions **25**.
McLean M, Warner B, Markham R, Fischer M, Walker J, Klein C, Hoeberechts M,
Dunn DC. 2023. Connecting conservation & culture: The importance of
Indigenous Knowledge in conservation decision-making and resource management
of migratory marine species. Marine Policy **155**.
McMillan MN, Huveneers C, Semmens JM, Gillanders BM. 2019. Partial female
migration and cool-water migration pathways in an overfished shark. ICES Journal
of Marine Science **76**.
Moher D et al. 2009. Preferred reporting items for systematic reviews and meta-
analyses: The PRISMA statement.
Neubauer P, Kim K, Large K, Brouwer S. 2024. SCIENTIFIC COMMITTEE
TWENTIETH REGULAR SESSION Manila, Philippines Stock Assessment of
Silky Shark in the Western and Central Pacific Ocean 2024.
Pacoureau N et al. 2021. Half a century of global decline in oceanic sharks and rays.
Nature **589**.
Pacoureau N et al. 2023. Conservation successes and challenges for wide-ranging
sharks and rays. Proceedings of the National Academy of Sciences of the United
States of America **120**.
Pimiento C, Albouy C, Silvestro D, Mouton TL, Velez L, Mouillot D, Judah AB,
Griffin JN, Leprieur F. 2023. Functional diversity of sharks and rays is highly
vulnerable and supported by unique species and locations worldwide. Nature
Communications **14**.

Poisson F, Demarcq H, Coudray S, Bohn J, Camiñas JA, Groul JM, March D. 2024.
Movement pathways and habitat use of blue sharks (*Prionace glauca*) in the
Western Mediterranean Sea: Distribution in relation to environmental factors,
reproductive biology, and conservation issues. Fisheries Research **270**.
Roberson LA et al. 2021. Multinational coordination required for conservation of over
90% of marine species. Global Change Biology **27**.
Roff G, Brown CJ, Priest MA, Mumby PJ. 2018. Decline of coastal apex shark
populations over the past half century. Communications Biology **1**.
Rowat D, Speed CW, Meekan MG, Gore MA, Bradshaw CJA. 2009. Population
abundance and apparent survival of the Vulnerable whale shark *Rhincodon typus* in
the Seychelles aggregation. Page ORYX.
Runge CA, Gallo-Cajiao E, Carey MJ, Garnett ST, Fuller RA, McCormack PC. 2017.
Coordinating Domestic Legislation and International Agreements to Conserve
Migratory Species: A Case Study from Australia. Conservation Letters **10**.
Sherman CS, Digel ED, Zubick P, Eged J, Haque AB, Matsushiba JH, Simpfendorfer
CA, Sant G, Dulvy NK. 2023. High overexploitation risk due to management
shortfall in highly traded requiem sharks. Conservation Letters **16**.
Simpfendorfer CA et al. 2023. Widespread diversity deficits of coral reef sharks and
rays. Science **380**.
Somveille M, Bay RA, Smith TB, Marra PP, Ruegg KC. 2021. A general theory of
avian migratory connectivity. Ecology Letters **24**.
Spaet JLY, Patterson TA, Bradford RW, Butcher PA. 2020. Spatiotemporal distribution
patterns of immature Australasian white sharks (*Carcharodon carcharias*).
Scientific Reports **10**.
Stein RW, Mull CG, Kuhn TS, Aschliman NC, Davidson LNK, Joy JB, Smith GJ,
Dulvy NK, Mooers AO. 2018. Global priorities for conserving the evolutionary
history of sharks, rays and chimaeras. Nature Ecology and Evolution **2**.
Stevens JD, Bradford RW, West GJ. 2010. Satellite tagging of blue sharks (Prionace
glauca) and other pelagic sharks off eastern Australia: Depth behaviour,
temperature experience and movements. Marine Biology **157**.
Team RC. 2023. R Core Team 2023 R: A language and environment for statistical
computing. R foundation for statistical computing. https://www.R-project.org/. R
Foundation for Statistical Computing.
Teitelbaum CS, Mueller T. 2019. Beyond Migration: Causes and Consequences of
Nomadic Animal Movements.
Thorburn J et al. 2024. Assessing the potential of acoustic telemetry to underpin the
regional management of basking sharks (*Cetorhinus maximus*). Animal
Biotelemetry **12**. BioMed Central Ltd.
UNEP-WCMC. 2024. State of the World's Migratory Species. Convention on the
Conservation of Migratory Species of Wild Animals:1–88. Available from
www.unep-wcmc.org.
Webster MS, Marra PP, Haig SM, Bensch S, Holmes RT. 2002. Trends in Ecology &
Evolution **17**.
Wickham H et al. 2019. Welcome to the Tidyverse. Journal of Open Source Software **4**.
Wiens JA. 1997. Metapopulation Dynamics and Landscape Ecology. Page
Metapopulation Biology.
Womersley FC et al. 2022. Global collision-risk hotspots of marine traffic and the
world's largest fish, the whale shark. Proceedings of the National Academy of
Sciences of the United States of America **119**.

Yan HF, Kyne PM, Jabado RW, Leeney RH, Davidson LNK, Derrick DH, Finucci B,
Freckleton RP, Fordham S V., Dulvy NK. 2021. Overfishing and habitat loss
drives range contraction of iconic marine fishes to near extinction. Science
Advances **7**.
723
724

Table 1. Summary of the 31 species reviewed: their IUCN Red List Categories (Near Threatened (NT), Vulnerable (VU), Endangered (EN), and Critically Endangered (CR)); whether the species is included (marked by an 'x') in the Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act), inclusion in the Memorandum of Understanding on the Conservation of Migratory Sharks (Shark MoU) or mentioned as requiring international cooperation in the Action Plan for Australian Sharks and Rays 2021; if and which CMS instrument the species was included in; the total number of references describing connections to Australian water; and the Habitat Use of a species (Euryhaline (E), Coastal (C), or Oceanic (O)).

Order	Family	Species name	Common name	IUCN Status	EPBC Act Threatened	EPBC Act Migratory	CMS	Shark MoU	Action Plan Int'l Coop	No. Aus References	Habitat Use
Lamniformes	Lamnidae	*Carcharodon carcharias*	White Shark	VU	x	x	I/II	x		20	C/O
		Isurus oxyrinchus	Shortfin Mako	EN		x	II	x	x	5	O
		Isurus paucus	Longfin Mako	EN		x	II	x	x	1	O
		Lamna nasus	Porbeagle	VU		x	II	x		1	C/O
	Alopiidae	*Alopias pelagicus*	Pelagic Thresher	EN			II		x	0	O
		Alopias superciliosus	Bigeye Thresher	VU			II		x	1	O
	Cetorhinidae	*Cetorhinus maximus*	Basking Shark	EN		x	I/II	x	x	0	O
Orectolobiformes	Rhincodontidae	*Rhincodon typus*	Whale Shark	EN	x	x	I/II	x	x	16	C/O
Carcharhiniformes	Carcharhinidae	*Carcharhinus leucas*	Bull Shark	VU						9	E/C
		Galeocerdo cuvier	Tiger Shark	NT						8	C/O
		Carcharhinus falciformis	Silky Shark	VU		x	II		x	7	C/O
		Carcharhinus obscurus	Dusky Shark	EN			II			6	C/O
		Prionace glauca	Blue Shark	NT			II			1	O
		Carcharhinus longimanus	Oceanic Whitetip Shark	CR		x	I		x	0	O
	Sphyrnidae	*Sphyrna mokarran*	Great Hammerhead Shark	CR			II			1	
		Sphyrna lewini	Scalloped Hammerhead	CR			II			0	C/O
		Sphyrna zygaena	Smooth Hammerhead Shark	VU			II			0	C/O
	Triakidae	*Galeorhinus galeus*	School Shark	CR			II			3	C/O
Rhinopristiformes	Pristidae	*Pristis pristis*	Largetooth Sawfish	CR	x	x	I/II			3	E/C/O
		Pristis zijsron	Green Sawfish	CR	x	x	I/II			1	E/C/O
		Pristis clavata	Dwarf Sawfish	CR	x	x	I/II			0	E/C

		Pristis pectinata	Smalltooth Sawfish	CR		I/II		0	E/C
	Anoxypristidae	*Anoxypristis cuspidata*	Narrow Sawfish	CR	x	I/II		0	E/C/O
	Rhinidae	*Rhynchobatus australiae*	Bottlenose Wedgefish	CR		II		0	E/C
Myliobatiformes	Mobulidae	*Mobula alfredi*	Reef Manta Ray	VU	x	I/II		6	C/O
		Mobula birostris	Oceanic Manta Ray	EN	x	I/II	x	1	O
		Mobula mobular	Spinetail Devil Ray	EN	x	I/II		1	C/O
		Mobula kuhlii	Shortfin Devil Ray	EN		I/II		0	C/O
		Mobula tarapacana	Sicklefin Devil Ray	EN		I/II		0	C/O
		Mobula thurstoni	Bentfin Devil Ray	EN	x	I/II		0	C/O
		Mobula eregoodoo	Longhorned Pygmy Devil Ray	EN	x	I/II		0	C/O

733 **Table 2.** The proportion of individuals sampled by sex and life stage, and the number of metasites generated, for the six species demonstrating transboundary
734 movement. The total number of individuals includes those sampled using acoustic and satellite telemetry, and mark-recapture studies.

Common name	Total number of individuals sampled	Male	Female	Unknown sex	Adult	Subadults	Juvenile	Unknown life stage	Total number of metasites
White Shark	917	30.8%	29.1%	13.5%	5.4%	7.7%	27.2%	32.9%	29
Shortfin Mako	142	17.6%	58.4%	44.6%	2.8%	0	96.4%	1.4%	16
Whale Shark	1711	36.9%	11.4%	51.4%	0	0	4.6%	95.2%	27
Tiger Shark	68	26.0%	73.9%	0	30.4%	2.2%	64.2%	3.0%	28
Blue Shark	24	33.3%	41.6%	25%	29.1%	0	8.3%	62.5%	18
School Shark	57	8.7%	87.7%	0	63.1%	0	0	36.8%	8

736 *Figure 1. Schematic of the systematic literature review screening protocol illustrated using*
737 *the white shark (Carcharodon carcharias).*
738
739
740

741 *Figure 2. Species with known transboundary movements between the Australian EEZ and*
742 *other jurisdictions (including areas beyond national jurisdiction; ABNJ). Diagram*
743 *generated using R package circlize (Gu et al. 2014).*

Figure 3. Network connectivity models showing the transboundary movements of six shark species: (a) White shark (b) Shortfin mako, (c) Tiger shark, (d) Whale shark), (e) Blue shark, and (f) School shark. These models depict connectivity across Australia's Exclusive Economic Zone (EEZ) and into adjacent international waters. Maps were

769 *generated using additional data from rnaturalearth (Massicotte and Young, 2023) and*
770 *EEZ boundaries from Flanders Marine Institute (2023).*
771
772