Expanding on the portuarization syndrome from an ecological perspective: eDNA reveals

rich diversity, non-indigenous hotspots, and biotic homogenization in ports

Authors

Ginevra Lilli*[1,2,3], Annaëlle Caillarec-Joly[2,3], Clément Violet[4], Marc Bouchoucha[5], Xavier Turon[1],

Sophie Arnaud-Haond[2], Frédérique Viard*[3]

Affiliations

1 Centre for Advanced Studies of Blanes (CEAB-CSIC), Dept. of Marine Ecology, 17300 Blanes, Spain

2 MARBEC, IFREMER, IRD, CNRS, Univ Montpellier, Sète, France

3 ISEM, University of Montpellier, CNRS, IRD, Montpellier, France

4 Ifremer, Centre de Bretagne, DYNECO LEBCO, Plouzané, France, F-29280, Plouzane, France

5 Ifremer, LITTORAL, F-83500 La Seyne-sur-Mer, France

Corresponding authors: ginevralilli93@gmail.com; frederique.viard@umontpellier.fr

Abstract

Ports are well-known entry points for marine non-indigenous species (NIS), which arrive as hitchhikers on ships. Ports are also expected to be gateways for the spread of NIS in the wild and resemble each other more than communities outside due to their singular characteristics. However, the uniqueness of species assemblages in ports and how they differ from natural habitats have only been marginally investigated, notably at regional scale. Using eDNA metabarcoding, we obtained a comprehensive and standardized overview of metazoan community diversity in 12 paired ports and adjacent natural areas along the northwestern Mediterranean Sea. As expected, we found that NIS are more abundant in ports than in natural habitats, and that the species assemblages in ports differ from those in natural habitats. In addition, we observed that communities in ports are far more homogeneous than their natural counterparts. This finding supports the hypothesis of biotic homogenization in highly anthropized habitats. We also observed a pattern that had previously been documented mainly in fish, but that we identified here in every phylum studied except Arthropoda: species richness detected in ports is comparable to, and in some case even greater than, that observed in natural habitats. Overall, our findings broaden, through an ecological perspective, the "portuarization syndrome" concept, which originally defined ports as unique replicated environments that promote specific evolutionary processes.

Keywords

biological invasions, biotic homogenization, ports, artificial habitat, coastal habitats, metabarcoding

Introduction

Since the beginning of the nineteenth century, urban areas and megacities have increased in size and number

as part of the urban transition (Day et al., 2021), notably affecting coastal zones (Creel, 2003; Day et al., 2021).

Despite their defined borders, urban areas significantly impact the surrounding environment (Elmqvist et al.,

2013) and create modified or entirely new habitats. The ecological and evolutionary processes of these habitats

are only just beginning to be understood through the field of "urban science" (McKinney, 2002; Rivkin et al.,

2019; Szulkin et al., 2020), which has been so far largely focused on terrestrial ecosystems. However, coastal

urbanization substantially impacts marine coastal habitats and ecosystems (Aguilera et al., 2020; Bishop et al.,

2017; Mayer-Pinto et al., 2018), which calls for strengthening marine urban sciences (Rega-Brodsky et al., 2022;

Todd et al., 2019). In addition to their key ecological roles, coastal ecosystems support numerous economic

sectors, including fisheries, tourism, wind energy production, and mineral resource extraction (Martínez et al.,

2007; Seitz et al., 2014; Sempere-Valverde et al., 2022), leading to the worldwide proliferation of ports (Jouffray

et al., 2020).

Ports provide new marine habitats characterized by particular abiotic properties (Alter et al., 2021; Touchard

et al., 2023). For example, due to intense human activity and their (semi-)enclosed structure, they are

characterized by a high concentration of waste and pollutants, stagnant water, turbidity and an irregular light

regime (Cassi et al., 2008; Cutroneo et al., 2017; Heery et al., 2017; Luna et al., 2019; Schaefer et al., 2025). In

contrast, their biotic properties remain largely unexplored (Madon et al., 2023), though ports are recognized

as hotspots of biological introductions, with a high proportion of non-indigenous species (NIS) (e.g. Ferrario et

al., 2017; O'Shaughnessy et al., 2020). This issue is noteworthy since the escape of NIS from ports is recognised

as a primary vector of their spread into the natural environment (Afonso et al., 2020; Zenetos & Galanidi, 2020).

Nevertheless, NIS occurrence in natural habitats surrounding ports has only been marginally investigated (Diem

et al., 2023; Sghaier et al., 2019; Soares et al., 2018; Zarcero et al., 2024) as compared to their occurrence in

ports (e.g. Tempesti et al., 2020). Yet, it could provide useful information for better understanding the

conditions under which NIS expand their range from their primary sites of introduction.

Other biotic features of port ecosystems have been linked to reduced richness of "resident" species (i.e., term

used hereafter to refer to species not classified as NIS) (Tanasovici et al., 2025): a pattern often explained by

the low tolerance of many species to anthropogenic disturbances (Chapman, 2003; Chatzinikolaou et al., 2018;

Schaefer et al., 2025). So far, fish seem to be the only exception. Fish biodiversity tends to be higher in ports

than in nearby natural habitats because fish use ports to forage, escape predators, and establish nursery

grounds (Bouchoucha et al., 2016; Manel et al., 2024). Despite this exception, port communities are generally

considered to consist primarily of stress-tolerant, synanthropic, and non-indigenous species, which are

"filtered" by port abiotic properties (Tempesti et al., 2020). This may result in communities within ports that

resemble each other more than communities outside (i.e., biotic homogenisation, Olden et al. 2004, Todd et

al., 2019). This global consequence has only been marginally explored (Lokatis & Jeschke, 2022; McKinney &

Lockwood, 1999). The singularity of ports, acting as replicated environments, led to the definition of the

"biological portuarization" syndrome (Touchard et al., 2023), defined as "the evolution of marine species in

port ecosystems under human-altered selective pressures".

The present work aims to further investigate the portuarization syndrome through an ecological perspective

on animal marine communities and explore the aforementioned gaps, by comparing ports and their nearby

natural habitats at a regional scale in the northwestern Mediterranean Sea. Due to the growing recreational

boating industry, the number of small ports in the Mediterranean Sea has increased in recent years. Nowadays,

there are more than 900 marinas in this basin, over half of which located in the northwest Mediterranean Sea

(Carreño & Lloret, 2021). In our study area along the coast of France (continental and insular, including Corsica),

65 recreational ports with 39,900 berths were reported in 2018 (MTES, 2018), providing numerous sailing

opportunities and port connections. Mediterranean ports are indeed characterized by intense maritime traffic

connecting them at various scales (Ferrario et al., 2024; Ulman et al., 2019a). Additionally, previous studies

have shown the high prevalence of NIS in these ports (Ulman et al., 2017). However, the diversity of resident

species has been less investigated than that of NIS, leaving a gap in our understanding of the overall biodiversity

inhabiting these ports. This partly stems from the traditional visual methods used for investigating marine

biodiversity, which often involve a trade-off between depth and breadth. This restricts investigations to either

a few selected taxa across multiple sites or to multiple taxa in a limited spatial context. Environmental DNA

(eDNA) metabarcoding is a standardized approach that can overcome some of these limitations and

complement or substitute traditional methods for biodiversity assessment and NIS detection (Couton et al.,

2019; Fonseca et al., 2023; Keck et al., 2022; Stat et al., 2017).

We used eDNA metabarcoding of seawater samples to assess the diversity and composition of animalia

communities in 12 pairs of northwestern Mediterranean ports and neighbouring natural areas. Based on the

previous studies cited above, we tested the following hypotheses: species assemblages in ports differ from

those in natural habitats (H1) and show biotic homogenization across a regional scale (H2); communities in

ports are characterized by greater abundance of NIS (H3), lower species richness than natural habitats (H4),

and taxonomic and functional traits distinct from communities from the nearby natural habitats (H5).

92   Materials and Methods

93   Data collection

Seawater samples used for this study were collected, during summer 2021 (June-August) within the framework

of the SUCHIMed campaign (https://doi.org/10.17600/18001619), in seven small ports (i.e., marinas) along the

French Mediterranean continental coast and five along the Corsican coast (Figure 1A; Supplementary material).

Twelve sites in natural habitats, outside each study port, were also sampled at the same time. Seawater was

collected in triplicates from three pontoons spanning the port and from a boat in three spots within a few

hundred meters from its mouth. Two litres of seawater were collected at each sampling site over a maximum

depth of 50 cm and immediately filtered on site or on boat by using 0.22 μm Sterivex filters and a VAMPIRE

peristaltic pump (Bürkle GmbH, Deutschland). Filters were filled with 2 mL of preservation buffer (sucrose 0.75

M, Tris 0.05 M pH 8, EDTA 0.04 M) and stored at -20°C until DNA extraction. From the boat and in the ports,

field controls were obtained by filtering ultrapure water with the equipment used on site. In total, 72 samples

were collected across the 12 localities.

DNA was extracted from Sterivex filters using the NucleoSnap® Finisher Midi extraction kit (Macherey-Nagel)

following the protocol detailed in Supplementary Protocol 1 in Couton et al. (2022). Fragments of COI, 18S rRNA

and 12S rRNA genes were each amplified 9 times from each DNA sample. Information about primers sequences,

PCR conditions and library preparation are reported in Supplementary material. PCR libraries were sequenced

(2 x 250bp paired-end) by Novogene company, using NovaSeq 6000 SP Reagent Kit (500 cycles) and a Novaseq

6000 SP flow cell sequencer. Negative controls were included both during extraction and library preparation.

Figure 1: A) French localities where seawater was collected at three sites inside ports and three sites in nearby

natural habitats. An example is shown for Ajaccio in the small panel. Basemap: GEBCO_Grid, colour-shaded for

elevation (GEBCO Compilation Group, 2023); B) Proportion of resident species (i.e., autochthonous; RESS) (left

panel) and NIS (right panel) per phylum per locality and habitat type (i.e. Port vs Natural habitat). The number

of RESS and NIS recorded in each locality and habitat is reported at the top of each bar.

Bioinformatic analyses and taxonomic assignment

Paired-end raw reads were pre-processed using cutadapt v1.16 and bbMap v38.22, followed by a modified Unix

shell script based on Brandt et al. (2021) (https://gitlab.ifremer.fr/abyss-project/abyss-pipeline) for denoising

(DADA2 v1.10;Callahan et al., 2016) and clustering (Swarm v3.1.0 with d=1 for 12S and 18S and d=3 for COI;

Mahé et al., 2014). OTUs tables were decontaminated using the package decontam (Davis et al., 2018) and a

correction for tag switching was applied (Wangensteen et al., 2018). For COI data, mumu (Mahé, 2023, adapted

and modified from Lulu, Frøslev et al., 2017) was applied to minimize the bias due to the existence of paralogs

and pseudogenes (particularly nuclear degenerated copies, *numts*). OTUs sequences were aligned with those

in reference databases using BLAST v2.12.0 (Altschul et al., 1990). Sequence coverage was set at a minimum of

70% for 12S and 80% for COI and 18S. Minimum identity percentages with sequences in reference databases

were set according to Couton et al., (2022): 92% for COI and 90% for 12S (MIDORI2 database) and 99% for 18S

(Silva v138 database).

OTU tables were rarefied to the minimum number of reads per sample (i.e., 44,798 for COI, 107,876 for 18S,

44,798 for 12S). WoRMS package version 0.4.3 in R (Chamberlain & Vanhoorne, 2023) was used to correct the

nomenclature and remove non-marine taxa; chordates were split into Vertebrata and Tunicata due to the

important ecological differences between these two subphyla.

The OTU tables from the three markers were combined to generate a non-redundant taxonomy table. Taxa not

resolved to the species level with a particular marker were removed if they co-occur in the same sample with

an OTU assigned with higher resolution with another marker; otherwise, they were retained. The final dataset

(i.e., Species occurrence table) was converted to a species-level presence/absence matrix, excluding all OTUs

not identified at species level, to enable determining the NIS vs. resident status.

NIS identification

The final dataset including all OTUs identified at species level (hereafter "COMM" dataset, i.e. whole

community) was screened to identify non-indigenous species (NIS), by using the NIS database for France

established by Massè et al (2023). The ANIS-E database (Violet et al., *submitted*), which is a custom database

including all NIS reported in European Seas from 1492 to 2023, was also used for identifying NIS reported in

Europe. Species found in these databases were classified as "Non-Indigenous-Med Sea" when reported as NIS

in the Mediterranean Sea and "Non-Indigenous-Europe" when reported as NIS in other European Seas and not

native to the Mediterranean Sea. Species not found in these databases were classified by default as resident

species (RESS). RESS included species native to Mediterranean Sea, albeit some species presumably from

elsewhere but not yet reported as NIS in Europe may be included as well. We did not further screen the RESS

for potential novel NIS because of the risk of false positives (false NIS identification) when using metabarcoding

alone (Couton et al. 2022).

Functional traits classification

Functional traits can help elucidate the ecological processes that determine the presence of certain species in

specific environments (Mayer-Pinto et al., 2018) or those associated with biological invasions (Leclerc et al.,

2023). Hence, we characterized species using two functional traits, "motility" and "ecology", selected because

they provide information about the dispersal potential of NIS (mobile NIS have greater dispersal potential than

sessile ones) and the filtering effect of ports (ports can act as positive filters on species colonizing surfaces and

inhabiting the benthos). Motility of species was coded as "sessile" and "mobile". Ecology was coded for fish

using Fishbase (Froese & Pauly, 2000), following Rey et al., (2023) with slight modification to get a more even

distribution, as follows: "demersal" and "bathydemersal" species were grouped into "demersal", and "pelagic-

neritic", "pelagic-oceanic" and "bathypelagic" species into "pelagic"; "benthopelagic" and "reef-associated"

classes were left unmodified. For other species not classified as fish, the modalities for the "ecology" trait were

retrieved from the Functional group section available in WoRMS (WoRMS Editorial Board, 2025): species

classified as "benthos" and "benthos> macrobenthos" at adult stage were reported as "benthos"; species

classified as "plankton", "plankton>zooplankton" and "nekton" (only *Tursiops truncatus*) at adult stage were

classified as "pelagic-non-fish". An additional trait was considered for fishes, namely "Fishing vulnerability",

which could be related to the greater occurrence of fishes in ports, because of fishing restriction in ports and

presence of fishing discards (from recreational fishing boats moored in the port). Vulnerability scores were

retrieved from FishBase, where this trait is expressed as a numerical value from 0 (low vulnerability) to 100

(high vulnerability).

Diversity analyses

Diversity analyses were implemented in R studio (R Core Team, 2021). With presence-absence data, alpha-

diversity was estimated by species richness. As we expected to have more NIS (H3) but lower species richness

in ports (H4), t-tests or Mann Withey tests were performed. NIS proportion and richness were also examined

with one-way ANOVA (or Kruskal-Wallis test) between localities.

For beta-diversity analyses (H1 and H2), the Jaccard dissimilarity was used. Principal Coordinates Analyses

(PCoA) was used to visualize community's dissimilarities in an unconstrained ordination space. The possibility

that communities from ports and natural habitats displayed distinct species composition (H1) was tested

performing Partial Distance based Redundancy Analysis (dbRDA) on Jaccard dissimilarities with habitat type as

explanatory variable; to control for potential effect of locality, this factor was included in the partial dbRDA as

covariate. The statistical significance of this constrained ordination was assessed by ANOVA-like test

(*anova.cca*() function of vegan package). In addition, nestedness and turnover components of beta diversity

were estimated using the R package betapart v. 1.6 (Baselga et al., 2023) to define whether differences in

species composition between habitats were driven by species gain/loss or species replacement. Permutational

Analysis of Multivariate Dispersion (*PERMDISP*) was performed through the *betadispers*() function from the

vegan package in R (Oksanen et al., 2013) to test for different dispersion of Jaccard dissimilarities between ports

and between natural habitats. The hypothesis of greater biotic homogenization in ports (H3) was considered

verified if, in case of significant PERMDISP, this habitat type showed smaller dispersion of Jaccard dissimilarities

between samples than natural habitats.

To examine the influence of taxonomical and functional traits in determining species composition in different

habitats (H5), species significantly associated with either type of habitat were identified by Indicator Species

Analysis using R package indicspecies (De Cáceres & Legendre, 2009). In addition, species scores from the partial

dbRDA first axis were compared between taxonomical (i.e. Phylum and Class) and functional (i.e. motility and

ecology) traits by either t-tests or ANOVAs (or their non-parametric equivalents) depending on number of levels

of the trait. In case of significant ANOVA, post-hoc Tuckey Test (or their non-parametric equivalents) with

Benjamini-Hochberg ("*bh*") correction were implemented for pairwise comparisons between trait levels.

Spearman rank correlation was performed between the distribution of fish species scores along the first partial

dbRDA axis and values of fishing vulnerability.

Results

The outcome of the bioinformatics pipeline analysis is provided in the Supporting Information. The combination

of the three markers (Supplementary Figure 1) allowed to identify 538 taxa at the species level (i.e., Species

occurrence table) belonging to 18 phyla (Figure 1B). An average of 125 ± 23 (mean ± SE) species was recorded

in each locality (Figure 1B).

Out of the 538 species (i.e. whole community, COMM), 497 were classified as resident species (RESS), of which

327 were known as native to the Mediterranean Sea, leaving 120 species with unknown origin, possibly

including unreported Non-Indigenous species (NIS) in European seas. Finally, 41 species were identified as

unequivocal NIS, meaning they had previously been reported as such in Europe. Of these, eight were

cryptogenic and five had been reported as NIS in European Seas though not yet in the Mediterranean Sea

(Figure 2). The average percentage of NIS per locality (regardless of habitat type) was 6.8± 4.6% (mean± SE),

reaching up to 12.6 ± 2.03% and 9.7 ± 3.7% in Sete_1 and Sete_2, respectively (Supplementary Figure 2). This

contribution was not significantly different across the 12 localities (Kruskal-Wallis test, P value = 0.07).

Figure 2: NIS (n=41) presence per locality and attributes. The heatmap displays the number of replicates (out of

n=6, with 3 inside and 3 outside ports) where each NIS was recorded. Coloured bars on the right display habitat

and motility trait for each NIS. Symbols are added to highlight their status as cryptogenic (star) and/or as NIS

observed in Europe but not in yet in the Mediterranean (triangle).

Ports display species assemblages different from those in natural habitats and homogeneous at a regional

scale

Across the sampling range, communities from ports were distinct from those in natural habitats (as reported

by unconstrained and constrained ordinations respectively in Supplementary Figure 3A and Figure 3A). This

observation was supported statistically by ANOVA.cca on partial dbRDA on habitat type (F-statistic=2.70, p-

value = 0.001; R^2-adjusted=2.73%), thus confirming H1. The observed dissimilarities were mainly driven by

species replacement in the two habitats (turnover) rather than by species loss in either one of the habitats

(nestedness) (Figure 3B). Furthermore, lower dispersion of Jaccard dissimilarity values were observed for ports

than for natural habitats (Figure 3C), supporting the hypothesis of greater biotic homogenization in ports than

in natural habitats (H2) (*PERMDISP*, p-value = 0.001).

Testing H1 and H2 on RESS and NIS separately revealed a different behaviour of the two communities. Both

communities displayed different species in ports than natural habitats (Supplementary Figure 3B/C), and H1

was statistically confirmed by ANOVA.cca on partial dbRDA (RESS: ANOVA.cca on partial dbRDA: F-

statistic=2.63, p-value = 0.001; R^2-adjusted=2.6%; NIS: ANOVA.cca on partial dbRDA: F-statistic=4.17, p-value

0.001; R^2-adjusted=5.35%; Supplementary Figure 4A). However, RESS dissimilarities were driven by turnover

while those of NIS have an even contribution of both beta-diversity components (Supplementary Figure 4B).

Biotic homogenization in ports (H2) was confirmed for RESS (*PERMDISP*, p-value = 0.004) but not for NIS

(*PERMDISP*, p-value = 0.18) (Supplementary Figure 4C).

Figure 3: Jaccard dissimilarities based on species composition of communities (COMM) in ports and in natural

habitats. A) Partial dbRDA performed to test the effect of habitat on COMM species composition accounting for

the effect of locality. The analysis produced one constrained axis (CAP 1) and an unconstrained one (PCoA 1) due

to the inclusion of only one explanatory variable; B) boxplots displaying partitioning of Jaccard dissimilarities in

its nestedness and turnover components with comparisons between samples from inside ports and from natural

sites; C) boxplot showing Jaccard dissimilarities between samples collected inside ports (blue) and in natural

habitats (yellow).

NIS are overrepresented in ports compared to natural habitats

247 In line with our working hypothesis (H3), the proportion of NIS in ports (9.2 ± 3.7%) was more than twice that

248 in natural habitats (4.6 ± 3.5%). With two exceptions (Bonifacio and Sete_1, Figure 4), this trend was observed

249 in every locality and was significant in three of them, namely Seyne-sur-Mer, Porto-Vecchio and Nice (in the

250 later there was no NIS outside the port).

Figure 4: Barplot showing mean and standard error of the percentage of NIS in communities sampled in ports

and in natural habitats in each locality. Localities that showed a significant difference in the proportion of NIS

*inside and outside marinas are labelled with * (Mann-Whitney test, one-tailed, P value = 0.01 for Porto-Vecchio*

and Seyne-sur-mer; no NIS recorded outside Nice marina).

253
254
255
256

The number of NIS was also higher in ports across all localities (Mann Whitney tests, one-tailed, p-value < 0.01)

(Figure 5) and per locality (Supplementary Figure 5). In addition, six times more NIS were observed exclusively

in ports (23 NIS) than in natural habitats (4 NIS) (Figure 5). It is however worth noting that ca. 40% of NIS

detected in ports were also present in natural habitats (14 NIS; Figure 5).

The greatest NIS richness inside and outside ports was found in both Sete ports (Sete_1: average 10 ± 2 NIS

inside and 4 ± 1 NIS outside; Sete_2: average 11 ± 2 NIS inside and 3 ± 1 NIS outside).

Figure 5: Venn diagrams showing the total number of species exclusive or shared in ports and natural habitats,

considering the whole community (COMM), non –indigenous species (NIS) only and resident species (RESS) only.

Boxplots of species richness per sample calculated for each habitat type are also provided; significant difference

between the two habitats is indicated with asterisks (Mann-Whitney test, P value < 0.001).

Ports host rich communities regardless of high NIS occurrence

Contrary to our working hypothesis (H4), ports considered in this study did not display lower species richness

than nearby natural habitats (COMM, Figure 5, Supplementary Table 1). In fact, the reverse was true overall

(Mann-Whitney test, p-value < 0.001). Higher richness inside port was also observed in each locality separately

(Supplementary Figure 6), with significant differences in Calvi, Seyne-sur-Mer and Porto-Vecchio

(Supplementary Table 1). Considering phyla separately, H4 was confirmed only for Arthropoda (Mann-Whitney,

p-value = 0.04; Supplementary Figure 6); instead, nine phyla displayed greater species richness in ports than in

natural habitats: Vertebrata, Tunicata, Bryozoa, Cnidaria, Annelida, Porifera, Nemertea and Rotifera

(Supplementary Table 1 and Supplementary Figure 6).

The above-mentioned observations were not driven by the higher NIS occurrence in ports: testing H4 on RESS

community alone produced similar results of higher richness in ports that nearby natural habitats (Figure 5)

also across all locations (p-value > 0.05 for all comparisons).

Taxonomical and functional traits drive differential species occurrence in ports and natural habitats

Community dissimilarities between ports and natural habitats were explained by the differential occurrence of

60 species (including RESS and NIS) identified by Indicator Species Analysis (IndVal.g, p-value < 0.05,

Supplementary Table 2, Figure 6A). A higher number of species (53) are indicators of ports than of natural

habitats (7 species), suggesting again a more homogeneous and distinct community in the former. Most

indicator species were Teleostei (33.9% of indicator species of ports and 42.8% of natural habitats), followed

by Polychaeta in ports (18.8%) and Copepoda (42.8%) in natural habitats. NIS represented 16% of indicators of

ports while none was significantly associated with natural habitats (Figure 6A).

The distribution of species scores along the first axis of the partial dbRDA was analysed to determine if

differences in species occurrence between habitats were related to taxonomic or functional traits (Figure 6B).

Species scores were differentially distributed along the axis according to Phylum (Kruskal-Wallis test, p-value =

0.002), Class (Kruskal-Wallis, p-value =0.04), motility (Mann-Whitney test, P value = 0.003), ecology (for non-

fish species, Mann-Whitney test, p-value < 0.001; for fish species, Kruskal-Wallis test, P value = 0.03) and their

NIS status (Mann-Whitney test, P value = 0.002) (Supplementary Table 3). In contrast, no significant correlation

was found between fishing vulnerability and fish species scores along the first dbRDA axis (Spearman

correlation, p-value = 0.06), suggesting no relationship between the two factors.

As reported in Figure 6B and Supplementary Table 3, Vertebrata and Tunicata tended to be associated with

ports (i.e. species scores displaying positive values of dbRDA axis), and Arthropoda and Echinodermata with

natural habitats (i.e. negative values of dbRDA axis) (Post-hoc Dunn tests, p-value < 0.05). At Class level, species

in ports were mostly classified as Teleostei and Ascidiacea while those outside as Copepoda. Ports were also

associated with sessile and benthic non-fish species, as well as with demersal and benthopelagic fishes. In

contrast, mobile species, pelagic-non-fish and fishes classified as "reef-associated" and "pelagic" characterized

natural habitats.

Figure 6: Species scores of the partial dbRDA performed to test the effect of habitat on COMM species

composition accounting for locality; A) species scores of the 60 indicator species identified for each habitat by

Indicator Species Analysis: species labels are displayed for all 7 indicator species of natural habitats (yellow) and

only for the 20 indicator species of ports (blue) displaying the highest statistics in the permutation test (Inval.g);

NIS are reported in red; B) average values of species scores of partial dbRDA first axis obtained for each

functional and taxonomic group and for their classification as NIS and RESS: traits that significantly influenced

the distribution of species scores in the first dbRDA axis are flagged with an asterisk (for tests performed,

statistics and p-values see Supplementary Table 3). Values positioned toward the right side of the first dbRDA

axis are more strongly associated with ports while those toward the left side with natural habitats.

As reported in Supplementary Table 3, testing the influence of traits on RESS dissimilarities yielded results

similar to COMM, except for the phyla linked to each habitat type: only Vertebrata (Teleostei) and Arthropoda

(Copepoda) were mainly found, respectively, inside and outside ports (Post-hoc Dunn test, > 0.05). In contrast,

for NIS, no taxonomical or functional trait characterized NIS associated with either ports or natural habitats.

Discussion

Are ports acting as anthropogenic biodiversity reservoirs?

Using eDNA metabarcoding on 72 seawater samples collected in 12 pairs of ports and adjacent natural habitats

along the French Mediterranean coast, we identified 538 animal species, including 41 NIS. Our study revealed

marked differences between ports and natural habitats in community richness, structure and functional traits.

Beyond the expected higher prevalence of NIS in ports, a striking and not anticipated result was the overall

greater species richness in ports than in natural habitats across nearly all phyla.

Several factors may be considered to explain this observation. First, ports display a heterogeneous mosaic of

substrates, with infrastructure most often built on formerly soft-sediment habitats. The variety of materials and

structure used in ports creates a mosaic of microhabitats, likely suitable to support a diversity of species

assemblages (Connell & Glasby, 1999; Romoth et al., 2023; Todd et al., 2019). Second, the nutrient enrichment

in ports (García-Pintado et al., 2007; Guerra-García et al., 2021; Karakassis et al., 2005; Sebastiá & Rodilla, 2013)

may enhance primary production (Scott, 1978), and, under certain conditions, secondary production (Leslie et

al., 2005), thereby increasing taxonomic diversity (Whittaker et al., 2003), as observed for Polychaeta by

Dafforn et al., (2013). Third, the (semi-)enclosed structure of ports may benefit some taxa by providing still

waters suitable for foraging and nursery grounds, particularly for fishes, for which ports are also no-take areas.

In some cases, fish communities are as diverse –or even richer- than those in natural habitats (Bouchoucha et

al., 2016; Lapinski et al., 2014; Manel et al., 2024; Pérez-Ruzafa et al., 2006). In addition, greater fish presence

may also exert top-down control on space-dominant competitors among epibiota, enhancing community

diversity (Rivero et al., 2013). Intermediate levels of disturbance, common in small ports such as marinas, may

further promote diversity by enabling the coexistence of both subordinate and dominant species (Boulanger et

al., 2021; Dial & Roughgarden, 1998; Santos et al., 2021; Turon et al., 2022).

Lastly, methodological aspects of our study could also partly explain our results: although seawater samples

were collected in natural sites with a maximum depth of 50 meters, marinas are shallower and retain water

longer, potentially amplifying the eDNA signal from benthic species compared to more open natural habitats.

This explanation is in line with the exception observed in arthropods (i.e., lower species richness in ports), which

are dominated by pelagic/mobile species. However, fish diversity was also greater in ports, and our findings

thus aligned with others comparing Operational Taxonomic Units between ports and sites located in no- take

marine reserves (Macé et al., 2024; Manel et al., 2024). This methodological aspect is thus unlikely to explain

the results obtained overall. Nevertheless, further targeted studies on substrate-specific assemblages should

be carried out to make a more detailed comparison between the two types of habitats.

Port similarities and singularities: the ecological side of the "portuarization syndrome"

We found a greater community similarity between ports than between natural habitats at a regional scale, a

finding supporting the hypothesis of biotic homogenization. This also allows broadening, from an ecological

perspective, the concept of "portuarization syndrome" proposed by Touchard et al. (2023), which defines

specific port-induced evolutionary processes resulting from their distinctive biotic and abiotic properties and

their replication across a large spatial scale. This homogenization likely results from two processes:

environmental filtering of autochthonous species (Todd et al., 2019) and high connectivity among them due to

boating, which uniformizes species assemblages across scales, notably NIS (Touchard et al., 2023; Ulman, et al.,

2019a; Ulman et al., 2019b). To our knowledge, this is the first study to provide evidences for biotic

homogenization in ports at a regional scale. The portuarization syndrome is also visible through species

assemblages in ports differing significantly from those in nearby natural areas. Interestingly, and in line with

the reservoir effects discussed above, this dissimilarity between port and dissimilar habitats was not primarily

due to simple species gains or losses (nestedness) between natural sites and marinas, but rather to extensive

species turnover.

This pattern scales up to the Class level, with Teleostei and Copepoda exhibiting contrasted associations with

the two habitats, as revealed by the species scores of partial dbRDA. By being predated both by fish larvae and

adults, zooplankton populations are strongly regulated by fish (Gliwicz & Pijanowska, 1989). High fish density

in ports may reduce copepods species richness, through species-specific and size-dependent selective

predation (Fiksen et al., 2005; Lehtiniemi et al., 2007). Moreover, zooplankton is highly sensitive to water

temperature, transparency and oxygen concentration (Stoch et al., 2023; Witalis et al., 2024), which are often

altered in ports. This may further reduce the occurrence of zooplankton populations.

A second important factor distinguishing port communities from naturals sites was the studied functional traits,

namely species motility and ecology. Functional trait analyses confirmed the expectation of a rich and abundant

fouling community in ports (Tempesti et al., 2022) characterized by sessile and benthic species. The higher

association of demersal and benthopelagic fishes than pelagic ones with ports was also expected given the

reliance of the former species on benthic preys (Bergstad, 2019). In contrast, reef-associated fishes were less

represented, likely reflecting the lack of complex surfaces (e.g. crevices) within ports.

Interestingly, this role of ecological filter of ports that appears clearly for RESS is not obvious for NIS. No specific

beta-diversity component seems to drive NIS-community dissimilarities, with a balanced contribution of

turnover and nestedness. While only a portion of NIS may have spread and establish in nearby natural habitats

(15 shared detection across 42 NIS species), a little less than 10% (4) were detected only outside marinas. This

may reflect the difficulties in detecting all species with a limited sampling volume in a rich substrate, or to the

existence of alternative introduction vectors beyond shipping and boating – such as aquaculture, a particularly

relevant vector in France (Massé et al., 2023). Besides, none of the study traits provided insights into the

habitats where NIS were more likely to occur. Although it is challenging because many NIS are poorly studied

(but see e.g., Leclerc et al. 2023), future trait-based analyses of NIS should expand beyond mobility and ecology

traits, to examine in particular tolerance to environmental conditions including pollutants, foraging strategies,

dispersal traits and growth rates, with the aim of better predicting their establishment and spread likelihood.

Are NIS over-looked in the wild?

Across all localities, 41 non-indigenous species (NIS) were identified including five not yet recorded in the

Mediterranean but known as NIS in other European Seas. Although it is not possible to confirm their presence

with eDNA only (i.e., false positives may occur, see Couton et al., 2022), our result identifies localities in which

further investigation should be prioritized for targeted molecular investigation and/or surveys based on direct

observation.

As expected, most NIS were found in ports (and 23 exclusively) while 18 occurred in natural habitats (4 being

private to them) and the same was observed for NIS proportions. Nevertheless, in contrast with the general

pattern observed, a high proportion of NIS was observed in the Thau lagoon (higher than found in some ports),

which is the natural site paired with the port at Sete_1 and also close to the port of Sete_2. Despite being

recognized for its rich environment – supporting its classification as "Site of Community Importance" under the

Habitat directive of the Nature 2000 framework –, the Thau lagoon is also a well-known introduction hotspot

in France. This is due to a massive import of Japanese oysters in the seventies (Boudouresque et al., 2010),

usually travelling with their associated species settled on shells and in the material used to collect and transport

spats. In addition, the Thau lagoon is surrounded by many small cities and their associated marinas, as well as

diverse recreational activities adding to the port's effect. This exemplifies the tights link between the diversity

of introduction vectors/pathways and NIS establishment (Ojaveer et al., 2018).

The proportion of NIS in ports, though generally higher, was however not dramatically greater than in natural

habitats, with only three localities (Seyne-sur-Mer, Porto-Vecchio, and Nice) showing a significant increase in

NIS proportions in port. So far, little attention has been given to natural habitats near ports, with NIS being

primarily investigated only inside ports (Madon et al., 2023; Ulman et al., 2017) or in distant and pristine sites

such as marine protected areas (Borriglione et al., 2025; Mannino et al., 2017). Our results suggest that natural

habitats adjacent to ports may play a critical but underappreciated role as stepping-stone locations for the

colonization of more distant and pristine habitats, especially if this transient establishment allows NIS to

acclimate or adapt to local natural conditions. Natural habitats nearby ports are likely to be under the influence

of human activities. These disturbed conditions may lower competitive resistance from native species (Ardura

et al., 2016; Giakoumi & Pey, 2017), making these habitats particularly vulnerable to secondary establishment.

Therefore, it is crucial to survey NIS presence in these locations not only because it can inform about NIS capable

to spread further but also because they represent the majority of coastal zones in the highly urbanized

Mediterranean Sea.

Overall, our study revealed that ports are introduction hotspots harbouring unexpectedly high diversity across

animal phyla forming rather homogeneous communities at the regional scale. This provides empirical support

for the "portuarization syndrome". The substantial NIS presence in adjacent natural habitats simultaneously

underscores the need for expanding monitoring and NIS studies beyond ports to better understand the spatial

dynamics of marine bioinvasions and the interplay between the expanding urban sprawl and their natural

counterparts in coastal zones.

References

Afonso, I., Berecibar, E., Castro, N., Costa, J. L., Frias, P., Henriques, F., Moreira, P., et al., (2020).

Assessment of the colonization and dispersal success of non-indigenous species introduced in recreational

marinas along the estuarine gradient. Ecological Indicators, 113.

https://doi.org/10.1016/j.ecolind.2020.106147

Aguilera, M. A., Tapia, J., Gallardo, C., Núñez, P., & Varas-Belemmi, K. (2020). Loss of coastal

ecosystem spatial connectivity and services by urbanization: Natural-to-urban integration for bay

management. Journal of Environmental Management, 276. https://doi.org/10.1016/j.jenvman.2020.111297

Alter, E. S., Tariq, L., Creed, J. K., & Megafu, E. (2021). Evolutionary responses of marine organisms to

urbanized seascapes. Evolutionary Applications, 14(1), 210–232. https://doi.org/10.1111/eva.13048

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search

tool. Journal of Molecular Biology, 215, 403–410. https://doi.org/10.1016/s0022-2836(05)80360-2

Ardura, A., Juanes, F., Planes, S., & Garcia-Vazquez, E. (2016). Rate of biological invasions is lower in

coastal marine protected areas. Scientific Reports, 6. https://doi.org/10.1038/srep33013

Baselga, A., Orme, D., Villeger, S., De Bortoli, J., Leprieur, F., Logez, M., Martinez-Santalla, S. et al.,

(2012). betapart: an R package for the study of beta diversity. Methods in Ecology and Evolution, 3: 808-812.

https://doi.org/10.1111/j.2041-210X.2012.00224.x

Bergstad, OA. (2019). Fish: demersal fish (life histories, behavior, adaptations). In Encyclopedia of

Ocean Sciences, 3rd ed., Vols. 1–5. Elsevier Ltd. https://doi.org/10.1016/b978-0-12-813081-0.00673-x

Bishop, M. J., Mayer-Pinto, M., Airoldi, L., Firth, L. B., Morris, R. L., Loke, L. H. L., Hawkins, S. J et al.,

(2017). Effects of ocean sprawl on ecological connectivity: impacts and solutions. Journal of Experimental

Marine Biology and Ecology, 492, 7–30. https://doi.org/10.1016/j.jembe.2017.01.021

Borriglione, M., Ruitton, S., Blanfuné, A., Perret-Boudouresque, M., Thouroude, B., Verlaque, M.,

Boudouresque, C. F., & Thibaut, T. (2025). Non-indigenous species of macroalgae in French Mediterranean

marine protected areas: distribution and trends. in journal of marine science and engineering (Vol. 13).

Multidisciplinary Digital Publishing Institute (MDPI). https://doi.org/10.3390/jmse13020374

Bouchoucha, M., Darnaude, A. M., Gudefin, A., Neveu, R., Verdoit-Jarraya, M., Boissery, P., &

Lenfant, P. (2016). Potential use of marinas as nursery grounds by rocky fishes: Insights from four *Diplodus*

species in the Mediterranean. Marine Ecology Progress Series, 547, 193–209.

https://doi.org/10.3354/meps11641

Boudouresque, C. F., Klein, J., Ruitton, S., & Verlaque, M. (2010). Biological invasion: the Thau

lagoon, a Japanese biological island in the Mediterranean sea. In: Ceccaldi, HJ., Dekeyser, I., Girault, M., Stora,

G. (eds) Global Change: Mankind-Marine Environment Interactions. Springer, Dordrecht.

https://doi.org/10.1007/978-90-481-8630-3_27

Boulanger, E., Loiseau, N., Valentini, A., Arnal, V., Boissery, P., Dejean, T., Deter, J. et al., (2021).

Environmental DNA metabarcoding reveals and unpacks a biodiversity conservation paradox in

Mediterranean marine reserves. Proceedings of the Royal Society B: Biological Sciences, 288.

https://doi.org/10.1098/rspb.2021.0112

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016).

DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods, 13(7), 581–583.

https://doi.org/10.1038/nmeth.3869

Carreño, A., & Lloret, J. (2021). Environmental impacts of increasing leisure boating activity in

Mediterranean coastal waters. Ocean and Coastal Management, 209.

https://doi.org/10.1016/j.ocecoaman.2021.105693

Cassi, R., Tolosa, I., & Mora, S. de. (2008). A survey of antifoulants in sediments from Ports and

Marinas along the French Mediterranean coast. Marine Pollution Bulletin, 56(11), 1943–1948.

https://doi.org/10.1016/j.marpolbul.2008.07.011

Chamberlain, S., & Vanhoorne, B. (2023). _worrms: World Register of Marine Species (WoRMS)

Client_. R package version 0.4.3. In https://CRAN.R-project.org/package=worrms

Chapman, M. G. (2003). Paucity of mobile species on constructed seawalls: effects of urbanization

on biodiversity. Marine Ecology Progress Series, 264, 21–29. https://doi.org/10.3354/meps264021

Chatzinikolaou, E., Mandalakis, M., Damianidis, P., Dailianis, T., Gambineri, S., Rossano, C., Scapini,

F.,et al., (2018). Spatio-temporal benthic biodiversity patterns and pollution pressure in three Mediterranean

touristic ports. Science of the Total Environment, 624, 648–660.

https://doi.org/10.1016/j.scitotenv.2017.12.111

Connell, S. D., & Glasby, T. M. (1999). Do urban structures influence local abundance and diversity of

subtidal epibiota? A case study from Sydney Harbour, Australia. Marine Environmental Research, 47, 373–

387. https://doi.org/10.1016/s0141-1136(98)00126-3

Couton, M., Comtet, T., Le Cam, S., Corre, E., & Viard, F. (2019). Metabarcoding on planktonic larval

stages: an efficient approach for detecting and investigating life cycle dynamics of benthic aliens.

Management of Biological Invasions, 10(4), 657–689. https://doi.org/10.3391/mbi.2019.10.4.06

Creel, L. (2003). Ripple effects: population and coastal regions. Washington, DC: Population

Reference Bureau.

Cutroneo, L., Carbone, C., Consani, S., Vagge, G., Canepa, G., & Capello, M. (2017). Environmental

complexity of a port: evidence from circulation of the water masses, and composition and contamination of

bottom sediments. Marine Pollution Bulletin, 119(1), 184–194.

https://doi.org/10.1016/j.marpolbul.2017.03.058

Dafforn, K. A., Kelaher, B. P., Simpson, S. L., Coleman, M. A., Hutchings, P. A., Clark, G. F., Knott, N. A

et al, (2013). Polychaete richness and abundance enhanced in anthropogenically modified estuaries despite

high concentrations of toxic contaminants. PLoS ONE, 8. https://doi.org/10.1371/journal.pone.0077018

Davis, N. M., Proctor, Di. M., Holmes, S. P., Relman, D. A., & Callahan, B. J. (2018). Simple statistical

identification and removal of contaminant sequences in marker-gene and metagenomics data. Microbiome,

6(1). https://doi.org/10.1186/s40168-018-0605-2

Day, J. W., Gunn, J. D., & Burger, J. R. (2021). Diminishing opportunities for sustainability of coastal

cities in the Anthropocene: a review. Frontiers in Environmental Science, 9, 663275.

https://doi.org/10.3389/fenvs.2021.663275/bibtex

De Cáceres, M., & Legendre, P. (2009). Associations between species and groups of sites: Indices and

statistical inference. Ecology, 90(12), 3566–3574. https://doi.org/10.1890/08-1823.1

Dial, R., & Roughgarden, J. (1998). Theory of marine communities: The intermediate disturbance

hypothesis. Ecology, 79, 1412–1424. https://doi.org/10.1890/0012-9658(1998)079[1412:tomcti]2.0.co;2

Diem, A., Ramalhosa, P., Cacabelos, E., Ferrario, J., Castro, N., Henriques, F., Monteiro, J. G. et al.,

(2023). Monitoring non-indigenous species with passive sampling methods in an oceanic island. Journal of

Marine Science and Engineering, 11(2), 264. https://doi.org/10.3390/jmse11020264/s1

Elmqvist, T., Sukhdev, P., Goodness, J., Güneralp, B., Marcotullio, P. J., Mcdonald, R. I., Parnell, S. et

al., (2013). Urbanization, biodiversity and ecosystem services: challenges and opportunities A Global

Assessment (Springer Nature, Ed.).

Ferrario, J., Caronni, S., Occhipinti-Ambrogi, A., & Marchini, A. (2017). Role of commercial harbours

and recreational marinas in the spread of non-indigenous fouling species. Biofouling, 33(8), 651–660.

https://doi.org/10.1080/08927014.2017.1351958

Ferrario, J., Marchini, A., Gazzola, F., & Occhipinti-Ambrogi, A. (2024). The influence of recreational

boat traffic in the introduction of non-indigenous fouling species in three Ligurian marinas (Mediterranean

Sea, Italy). Estuarine, Coastal and Shelf Science, 303, 108810. https://doi.org/10.1016/j.ecss.2024.108810

Fiksen, Ø., Eliassen, S., & Titelman, J. (2005). Multiple predators in the pelagic: modelling behavioural

cascades. Journal of Animal Ecology, 74, 423–429. https://doi.org/10.1111/j.1365-2656.2005.00937.x

Fonseca, V. G., Davison, P. I., Creach, V., Stone, D., Bass, D., & Tidbury, H. J. (2023). The application of

eDNA for monitoring aquatic non-indigenous species: practical and policy considerations. In Diversity (Vol. 15,

Issue 5). MDPI. https://doi.org/10.3390/d15050631

Froese, R. and D. Pauly, Editors. 2000. FishBase 2000: concepts, design and data sources. ICLARM,

Los Baños, Laguna, Philippines. 344 pp. https://digitalarchive.worldfishcenter.org/handle/20.500.12348/2428

Frøslev, T. G., Kjøller, R., Bruun, H. H., Ejrnæs, R., Brunbjerg, A. K., Pietroni, C., & Hansen, A. J. (2017).

Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. Nature

Communications. 8(1), 1–11. https://doi.org/10.1038/s41467-017-01312-x

García-Pintado, J., Martínez-Mena, M., Barberá, G. G., Albaladejo, J., & Castillo, V. M. (2007).

Anthropogenic nutrient sources and loads from a Mediterranean catchment into a coastal lagoon: Mar

Menor, Spain. Science of the Total Environment, 373, 220–239.

https://doi.org/10.1016/j.scitotenv.2006.10.046

Giakoumi, S., & Pey, A. (2017). Assessing the effects of marine protected areas on biological

invasions: A global review. In Frontiers in Marine Science, 4:49. https://doi.org/10.3389/fmars.2017.00049

Gliwicz, Z. M., & Pijanowska, J. (1989). The role of predation in zooplankton succession. In: Sommer,

U. (eds) Plankton Ecology. Brock/Springer Series in Contemporary Bioscience. Springer, Berlin, Heidelberg.

https://doi.org/10.1007/978-3-642-74890-5_7

Guerra-García, J. M., Navarro-Barranco, C., Martínez-Laiz, G., Moreira, J., Giráldez, I., Morales, E.,

Fernández-Romero, A. et al., (2021). Assessing environmental pollution levels in marinas. Science of the Total

Environment, 762. https://doi.org/10.1016/j.scitotenv.2020.144169

Heery, E. C., Bishop, M. J., Critchley, L. P., Bugnot, A. B., Airoldi, L., Mayer-Pinto, M., Sheehan, E. V.,

Coleman, R. A. et al., (2017). Identifying the consequences of ocean sprawl for sedimentary habitats. In

Journal of Experimental Marine Biology and Ecology, 492, 31–48). Elsevier B.V.

https://doi.org/10.1016/j.jembe.2017.01.020

Jouffray, J. B., Blasiak, R., Norström, A. V., Österblom, H., & Nyström, M. (2020). The Blue

acceleration: the trajectory of human expansion into the ocean. one earth, 2(1), 43–54.

https://doi.org/10.1016/j.oneear.2019.12.016

554   Karakassis, I., Pitta, P., & Krom, M. D. (2005). Contribution of fish farming to the nutrient loading of

555 the Mediterranean. Scientia Marina, 69, 313–321. https://doi.org/10.3989/scimar.2005.69n2313

556   Keck, F., Blackman, R. C., Bossart, R., Brantschen, J., Couton, M., Hürlemann, S., Kirschner, D et al.,

557 (2022). Meta-analysis shows both congruence and complementarity of DNA and eDNA metabarcoding to

558 traditional methods for biological community assessment. Molecular Ecology, 31(6), 1820–1835.

559 https://doi.org/10.1111/mec.16364

560   Lapinski, M., Perrot, M., Le Direach, L., & Rouanet, E. (2014). Support to nursery functionalities of the

561 port of Marseilles-Fos given by artificial micro-habitats. https://doi.org/10.13140/2.1.1493.5843

562   Leclerc, J. C., Figueroa, N. N., Viard, F., & Brante, A. (2023). Distribution of functionally distinct native

563 and non-indigenous species within marine urban habitats. Diversity and Distributions, 29, 1445–1457.

564 https://doi.org/10.1111/ddi.13769

565   Lehtiniemi, M., Hakala, T., Saesmaa, S., & Viitasalo, M. (2007). Prey selection by the larvae of three

566 species of littoral fishes on natural zooplankton assemblages. Aquatic Ecology, 41, 85–94.

567 https://doi.org/10.1007/s10452-006-9042-6

568   Leslie, H. M., Breck, E. N., Chan, F., Lubchenco, J., & Menge, B. A. (2005). Barnacle reproductive

569 hotspots linked to nearshore ocean conditions. Proceedings of the National Academy of Sciences of the

570 United States of America, 102, 10534–10539. https://doi.org/10.1073/pnas.0503874102

571   Lokatis, S., & Jeschke, J. M. (2022). Urban biotic homogenization: Approaches and knowledge gaps.

572 Ecological Applications, 32. https://doi.org/10.1002/eap.2703

573   Luna, G. M., Manini, E., Turk, V., Tinta, T., D'Errico, G., Baldrighi, E., Baljak, V et al., (2019). Status of

574 faecal pollution in ports: a basin-wide investigation in the Adriatic Sea. Marine Pollution Bulletin, 147, 219–

575 228. https://doi.org/10.1016/j.marpolbul.2018.03.050

Macé, B., Mouillot, D., Dalongeville, A., Bruno, M., Deter, J., Varenne, A., Gudefin, A. et al., (2024).

The Tree of Life eDNA metabarcoding reveals a similar taxonomic richness but dissimilar evolutionary lineages

between seaports and marine reserves. Molecular Ecology, 33(12). https://doi.org/10.1111/mec.17373

Madon, B., David, R., Torralba, A., Jung, A., Marengo, M., & Thomas, H. (2023). A review of

biodiversity research in ports: Let's not overlook everyday nature! In Ocean and Coastal Management (Vol.

242). Elsevier Ltd. https://doi.org/10.1016/j.ocecoaman.2023.106623

Mahé, F., Rognes, T., Quince, C., de Vargas, C., & Dunthorn, M. (2014). Swarm: robust and fast

clustering method for amplicon-based studies. PeerJ, 2, e593. https://doi.org/10.7717/peerj.593

Mahé, F. (2023). "mumu: post-clustering curation tool for metabarcoding data"

https://github.com/frederic-mahe/mumu

Manel, S., Mathon, L., Mouillot, D., Bruno, M., Valentini, A., Lecaillon, G., Gudefin, A. et al., (2024).

Benchmarking fish biodiversity of seaports with eDNA and nearby marine reserves. Conservation Letters,

17(2). https://doi.org/10.1111/conl.13001

Mannino, A. M., Parasporo, M., Crocetta, F., & Balistreri, P. (2017). An updated overview of the

marine alien and cryptogenic species from the Egadi Islands Marine Protected Area (Italy). Marine

Biodiversity, 47, 469–480. https://doi.org/10.1007/s12526-016-0496-z

Martínez, M. L., Intralawan, A., Vázquez, G., Pérez-Maqueo, O., Sutton, P., & Landgrave, R. (2007).

The coasts of our world: Ecological, economic and social importance. Ecological Economics, 63(2–3), 254–

272. https://doi.org/10.1016/j.ecolecon.2006.10.022

Massé, C., Viard, F., Humbert, S., Antajan, E., Auby, I., Bachelet, G., Bernard, G et al., (2023). An

Overview of Marine Non-Indigenous Species Found in Three Contrasting Biogeographic Metropolitan French

Regions: Insights on Distribution, Origins and Pathways of Introduction. Diversity, 15(2), 161.

https://doi.org/10.3390/D15020161/S1

Mayer-Pinto, M., Cole, V. J., Johnston, E. L., Bugnot, A., Hurst, H., Airoldi, L., Glasby, T. M., & Dafforn,

K. A. (2018). Functional and structural responses to marine urbanisation. Environmental Research Letters,

13(1). https://doi.org/10.1088/1748-9326/aa98a5

McKinney, M. L. (2002). Urbanization, biodiversity, and conservation. BioScience, 52, 883–890.

https://doi.org/10.1641/0006-3568(2002)052[0883:ubac]2.0.co;2

McKinney, M. L., & Lockwood, J. L. (1999). Biotic homogenization: A few winners replacing many

losers in the next mass extinction. Trends in Ecology and Evolution, 14, 450–453.

https://doi.org/10.1016/s0169-5347(99)01679-1

Ojaveer, H., Galil, B. S., Carlton, J. T., Alleway, H., Goulletquer, P., Lehtiniemi, M., Marchini, A. t al.,

(2018). Historical baselines in marine bioinvasions: Implications for policy and management. PLOS ONE, 13(8),

e0202383. https://doi.org/10.1371/journal.pone.0202383

Oksanen, J., Blanchet, F., Kindt, R., Legendre, P., Minchin, P., & O'Hara, R. (2013). vegan: R package

for community ecologists: popular ordination methods, ecological null models and diversity analysis.

O'Shaughnessy, K. A., Hawkins, S. J., Yunnie, A. L. E., Hanley, M. E., Lunt, P., Thompson, R. C., & Firth,

L. B. (2020). Occurrence and assemblage composition of intertidal non-native species may be influenced by

shipping patterns and artificial structures. Marine Pollution Bulletin, 154, 111082.

https://doi.org/10.1016/j.marpolbul.2020.111082

Pérez-Ruzafa, A., García-Charton, J. A., Barcala, E., & Marcos, C. (2006). Changes in benthic fish

assemblages as a consequence of coastal works in a coastal lagoon: The Mar Menor (Spain, Western

Mediterranean). Marine Pollution Bulletin, 53, 107–120. https://doi.org/10.1016/j.marpolbul.2005.09.014

Rega-Brodsky, C. C., Aronson, M. F. J., Piana, M. R., Carpenter, E. S., Hahs, A. K., Herrera-Montes, A.,

Knapp, S. et al., (2022). Urban biodiversity: State of the science and future directions. Urban Ecosystems, 25,

1083–1096. https://doi.org/10.1007/s11252-022-01207-w

Rey, A., Viard, F., Lizé, A., Corre, E., Valentini, A., & Thiriet, P. (2023). Coastal rocky reef fish

monitoring in the context of the Marine Strategy Framework Directive: Environmental DNA metabarcoding

complements underwater visual census. Ocean & Coastal Management, 241, 106625.

https://doi.org/10.1016/j.ocecoaman.2023.106625

Rivero, N. K., Dafforn, K. A., Coleman, M. A., & Johnston, E. L. (2013). Environmental and ecological

changes associated with a marina. Biofouling, 29, 803–815. https://doi.org/10.1080/08927014.2013.805751

Rivkin, L. R., Santangelo, J. S., Alberti, M., Aronson, M. F. J., de Keyzer, C. W., Diamond, S. E., Fortin,

M. J. et al., (2019). A roadmap for urban evolutionary ecology. Evolutionary Applications, 12(3), 384–398.

https://doi.org/10.1111/eva.12734

Romoth, K., Darr, A., Papenmeier, S., Zettler, M. L., & Gogina, M. (2023). Substrate Heterogeneity as

a Trigger for Species Diversity in Marine Benthic Assemblages. Biology, 12(6).

https://doi.org/10.3390/biology12060825

Santos, L. N., Franco, A. C. S., de Souza, J. S., Miyahira, I. C., Rodrigues, A. J. S., Gonçalves, I. C. B.,

Krepsky, N. et al.,(2021). Using richness of native and non-native aquatic species along a climatic gradient to

test the intermediate disturbance hypothesis. Hydrobiologia, 848, 2055–2075.

https://doi.org/10.1007/s10750-021-04525-w

Schaefer, N., Bishop, M. J., Herbert, B., Hoey, A. S., Mayer-Pinto, M., Sherman, C. D. H. et al., (2025).

Major global ports alter light regimes for marine biofouling communities. Journal of Environmental

Management, 374. https://doi.org/10.1016/j.jenvman.2025.124119

Scott, B. D. (1978). Nutrient cycling and primary production in port hacking, new south wales. Marine

and Freshwater Research, 29, 803–815. https://doi.org/10.1071/mf9780803

Sebastiá, M. T., & Rodilla, M. (2013). Nutrient and phytoplankton analysis of a Mediterranean coastal

area. Environmental Management, 51, 225–240. https://doi.org/10.1007/s00267-012-9986-3

Seitz, R. D., Wennhage, H., Bergström, U., Lipcius, R. N., Ysebaert, T., & Seitz, R. D. (2014). Ecological

value of coastal habitats for commercially and ecologically important species. ICES Journal OfMarine Science,

3(71), 648–665. https://doi.org/10.1093/icesjms/fst152

Sempere-Valverde, J., Guerra-García, J. M., García-Gómez, J. C., & Espinosa, F. (2022). Coastal

urbanization, an issue for marine conservation. In Coastal Habitat Conservation: New Perspectives and

Sustainable Development of Biodiversity in the Anthropocene (pp. 41–79). Elsevier.

https://doi.org/10.1016/B978-0-323-85613-3.00007-4

Sghaier, Y. R., Zakhama-Sraieb, R., Hmida, A. Ben, & Charfi, F. (2019). An inventory of non-indigenous

species (NIS) inside and outside three tourist marinas from the southern Mediterranean coast.

https://www.researchgate.net/publication/332141163

Soares, M. de O., Campos, C. C., Santos, N. M. O., Barroso, H. de S., Mota, E. M. T., Menezes, M. O. B.

de, Rossi, S., & Garcia, T. M. (2018). Marine bioinvasions: Differences in tropical copepod communities

between inside and outside a port. Journal of Sea Research, 134, 42–48.

https://doi.org/10.1016/j.seares.2018.01.002

Stat, M., Huggett, M. J., Bernasconi, R., Dibattista, J. D., Berry, T. E., Newman, S. J., Harvey, E. S., &

Bunce, M. (2017). Ecosystem biomonitoring with eDNA: Metabarcoding across the tree of life in a tropical

marine environment. Scientific Reports, 7. https://doi.org/10.1038/s41598-017-12501-5

Stoch, F., Marrone, F., & Bruno, M. C. (2023). Classes Copepoda and Ichthyostraca. In Identification

and Ecology of Freshwater Arthropods in the Mediterranean Basin (pp. 131–156). Elsevier.

https://doi.org/10.1016/b978-0-12-821844-0.00012-0

Szulkin, M., Munshi-South, J., & Charmantier, A. (2020). Urban Evolutionary Biology. Urban

Evolutionary Biology. https://doi.org/10.1093/oso/9780198836841.001.0001

Tanasovici, R. M., Gibran, F. Z., & Dias, G. M. (2025). The proximity to marine infrastructure affects

fish diversity, the occurrence of non-indigenous species, and the dynamic of the sessile communities. Marine

Environmental Research, 207, 107086. https://doi.org/10.1016/J.MARENVRES.2025.107086

Tempesti, J., Langeneck, J., Maltagliati, F., & Castelli, A. (2020). Macrobenthic fouling assemblages

and NIS success in a Mediterranean port: The role of use destination. Marine Pollution Bulletin, 150.

https://doi.org/10.1016/j.marpolbul.2019.110768

Todd, P. A., Heery, E. C., Loke, L. H. L., Thurstan, R. H., Kotze, D. J., & Swan, C. (2019). Towards an

urban marine ecology: characterizing the drivers, patterns and processes of marine ecosystems in coastal

cities. Oikos, 128, 1215–1242. https://doi.org/10.1111/oik.05946

Touchard, F., Simon, A., Bierne, N., & Viard, F. (2023). Urban rendezvous along the seashore: Ports as

Darwinian field labs for studying marine evolution in the Anthropocene. Evolutionary Applications, 16(2),

560–579. https://doi.org/10.1111/eva.13443

Turon, M., Nygaard, M., Guri, G., Wangensteen, O. S., & Præbel, K. (2022). Fine-scale differences in

eukaryotic communities inside and outside salmon aquaculture cages revealed by eDNA metabarcoding.

Frontiers in Genetics, 13. https://doi.org/10.3389/fgene.2022.957251

Ulman, A., Ferrario, J., Occhpinti-Ambrogi, A., Arvanitidis, C., Bandi, A., Bertolino, M., Bogi, C. et al.,

(2017). A massive update of non-indigenous species records in Mediterranean marinas. PeerJ, 2017.

https://doi.org/10.7717/peerj.3954

Ulman, A., Ferrario, J., Forcada, A., Arvanitidis, C., Occhipinti-Ambrogi, A., & Marchini, A. (2019a). A

Hitchhiker's guide to Mediterranean marina travel for alien species. Journal of Environmental Management,

241, 328–339. https://doi.org/10.1016/j.jenvman.2019.04.011

Ulman, A., Ferrario, J., Forcada, A., Seebens, H., Arvanitidis, C., Occhipinti-Ambrogi, A., & Marchini, A.

(2019b). Alien species spreading via biofouling on recreational vessels in the Mediterranean Sea. Journal of

Applied Ecology, 56(12), 2620–2629. https://doi.org/10.1111/1365-2664.13502

Violet, Clément, Mathieu Chevalier, Amelia Curd, and Frédérique Viard. 2025. "ANIS-E: An Atlas of

Marine Non-Indigenous Species in Europe." *Submitted*.

Wangensteen, O. S., Cebrian, E., Palacin, C., & Turon, X. (2018). Under the canopy: Community-wide

effects of invasive algae in Marine Protected Areas revealed by metabarcoding. Marine Pollution Bulletin,

127(March 2017), 54–66. https://doi.org/10.1016/j.marpolbul.2017.11.033

Whittaker, R. J., Heegaard, E., Mittelbach, G. G., Scheiner, S. M., & Steiner, C. F. (2003). What is the

observed relationship between species richness and productivity? Ecology, 84, 3384–3395.

https://doi.org/10.1890/02-3128

Witalis, B., Iglikowska, A., Ronowicz, M., Weydmann-Zwolicka, A., & Kukliński, P. (2024). The

zooplankton community of Baltic Sea ports: diversity and seasonal dynamics. Oceanologia, 66, 353–364.

https://doi.org/10.1016/j.oceano.2024.02.001

Zarcero, J., Antich, A., Rius, M., Wangensteen, O. S., & Turon, X. (2024). A new sampling device for

metabarcoding surveillance of port communities and detection of non-indigenous species. IScience, 27(1),

108588. https://doi.org/10.1016/j.isci.2023.108588

Zenetos, A., & Galanidi, M. (2020). Mediterranean non indigenous species at the start of the 2020s:

Recent changes. Marine Biodiversity Records, 13(10). https://doi.org/10.1186/s41200-020-00191-4

Author contributions

GL: Methodology (equal); Data curation (lead); Formal analysis (lead); Visualization (lead); Writing – original draft (lead); Writing – review and editing (lead). **ACJ**: Methodology (equal); Investigation (equal); Resources (equal); Formal analysis (supporting); Software (equal); Writing – review and editing (Supporting). **CV**: Data curation (supporting); Resources (supporting); Writing – review and editing (Supporting). **MB**: Methodology (supporting); Resources (equal); Investigation (supporting); Writing – review and editing (Supporting). **XT**: Resources (equal); Formal analysis (supporting); Writing – original draft (supporting); Writing – review and editing (equal). **SAH**: Conceptualization (equal); Methodology (equal); Resources (equal); Data curation (equal); Formal analysis (supporting); Funding acquisition (equal); Writing – original draft (equal); Writing – original draft (supporting); Writing – review and editing (equal). **FV**: Conceptualization (equal); Methodology (equal); Investigation (supporting); Resources (equal); Formal analysis (supporting); Data curation (supporting); Funding acquisition (equal); Project administration (lead); Writing – original draft (equal); Writing – review and editing (equal).

Acknowledgements

The authors wish to thank all partners involved in the 2021 SUCHIMED campaign (RV L'Europe, https://doi.org/10.17600/1800 1619), and especially, Françoise Miralles, Fabienne Chavanon and Christophe Ravel. The authors are grateful to Pierre Taberlet for sharing the unpublished sequences for the primers of the teleo4 marker, and to Argyro Zenetos for her thorough review of the NIS list presented in this study and for offering insightful feedback. This work benefited from funding through the French National Research Agency (ANR) under the "Investissements d'Avenir" program with the reference ANR-16-IDEX-0006 (i-siteMUSE) supporting the MarEEE project. Data used in this work were partly produced through the GenSeq technical facilities of the « Institut des Sciences de l'Evolution de Montpellier » with the support of LabEx CeMEB, an ANR "Investissements d'avenir" program (ANR-10-LABX-04-01). This research was co-funded by the European Union (GA#101059915) supporting the BIOcean5D project through the funding of GL post-doctoral fellow. Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union. Neither the European Union nor the granting authority can be held responsible for them.

Data archiving statement

The codes and datasets used to perform the analyses will be provided upon acceptance of the paper.