Directed evolution of compact RNA-guided nucleases for enhanced activity in mammalian cells

Fedor Gorbenko[1*], Irene Sala[1*], Young-Yoon Lee[1], Lilly van de Venn[1], Charles D Yeh[1], András Tálas[3], Tautvydas Karvelis[2], Gytis Druteika[2], Luca V. Bechter[1], Iryna Vykhlyantseva[1], Markus S. Schröder[1], Ana Gvozdenovic[1], Gerald Schwank[3], Virginijus Siksnys[2] & Jacob E. Corn[1, **]

[1] Institute of Molecular Health Sciences, Department of Biology, Swiss Federal Institute of Technology (ETH) Zurich, Zurich, Switzerland

[2] Institute of Biotechnology, Life Sciences Center, Vilnius University, Vilnius, Lithuania

[3] Institute of Pharmacology and Toxicology, University of Zurich, Zurich, Switzerland

* Equal contribution

** Corresponding author. E-mail: jacob.corn@biol.ethz.ch

Abstract

RNA-guided nucleases enable DNA editing and offer promise for treating genetic diseases, particularly when used for precise sequence replacement. However, many of the most effective enzymes, such as *S. pyogenes* Cas9, are too large for delivery using vectors like adeno-associated virus (AAV). This has prompted interest in smaller alternatives from the Cas12f and TnpB families. Yet, these nucleases often show low activity in mammalian cells, limiting their utility. To address this, we used directed evolution in human cells to select variants with greatly improved activity. The resulting variants, Cas12f1Super and TnpBSuper, exhibited up to 11-fold increase in editing efficiency without increased off-target effects. When tested as a base editor, Cas12f1Super showed up to 10-fold improvement relative to the previously engineered CasMINI, suggesting utility beyond nuclease-related activities. These compact and efficient genome editors expand the current toolkit and hold promise for both research and therapeutic use in mammalian systems.

Introduction

CRISPR-Cas technology has revolutionized genome editing by enabling precise, RNA-guided manipulation of genomic DNA in a wide range of contexts. This advancement has unlocked powerful applications in biomedical research and holds great promise for treating genetic diseases through targeted genome correction. However, most existing highly active CRISPR systems, such as Cas9 and Cas12a, are large proteins, limiting their compatibility with clinically relevant size-restricted delivery vectors like adeno-associated viruses (AAVs)[1].

This constraint has driven growing interest in the development of compact genome editing systems that are more compatible with AAVs and other delivery platforms. Several miniature nucleases, including Cas12f and TnpB-based systems, have recently been explored as alternatives[2]. However, these proteins exhibit substantially lower editing efficiencies in mammalian systems than larger Cas enzymes, limiting their applicability and stimulating efforts to improve their activity[3–9].

Efforts to enhance enzyme activity typically rely on rational design or directed evolution. While rational design leverages structural and mechanistic insights to introduce beneficial mutations, it is inherently limited by our incomplete understanding of protein function and may miss unexpected improvements. Directed evolution allows broader exploration through functional selection from diverse variant pools. It is often performed in non-human systems, such as phage or bacteria[9,10], which offers several technical advantages such as the ability to interrogate large libraries necessary to cover large Cas proteins. However, these hosts fail to recapitulate many aspects of human cell biology, ranging from intracellular folding propensity to chromatin structure. Their DNA repair responses can also differ greatly from human cells, which for genome editing purposes necessitates evolution towards proxy targets such as targeting binding or catalytic activity. This may yield variants with high activity in the evolved context but modest improvement when applied to genome editing in human cells.

The effectiveness of directed evolution in mammalian systems is constrained by the scale of cell culture. For example, screening 2.5×10^7 variants in mammalian cells would allow testing of only about 7% of all possible single and double amino acid mutants for SpCas9 (1368 amino acids). This consideration underscores another advantage of compact editors.

With only 528 and 408 amino acids, respectively, Un1Cas12f1 and ISDra2 TnpB require far

fewer variants to cover a meaningful fraction of their sequence space[11,12,13]. Screening $2.5 \times$

10^7 variants covers approximately 50% of all single and double mutants for Un1Cas12f1 and

83% for ISDra2 TnpB. This reduced search space allows for effective directed evolution of

desired genome editing activities in mammalian cells. While all possible single point

mutations of AsCas12f1 have recently been tested by deep mutational scanning (DMS),

potentially more active variants with multiple mutations have never been explicitly selected

in human cells via directed evolution[8].

For many genome editing applications, including therapeutic gene correction, nuclease

activity alone is not sufficient. Editing is the result of host cell repair processes, including

high-fidelity homology-directed repair (HDR) that enables precise sequence changes when a

donor template is present. In mammalian cells, HDR is usually outstripped by the faster,

error-prone non-homologous end joining (NHEJ) pathway, leading to several workaround

technologies such as base and prime editing[14,15]. But base and prime editing add further size

to the editing architecture, again emphasizing the importance of compact editors. Improving

editing efficiency, particularly precise sequence manipulation such as HDR, remains a major

obstacle to expanding the utility of small genome editors for clinical use. In this regard,

mammalian-based selection offers a key advantage: it allows identification of variants that

specifically enhance editing performance in therapeutic-relevant settings rather than catalytic

activity.

Here we performed mammalian cell directed evolution on two small genome editing proteins:

Un1Cas12f1, a miniature Cas12f1-family nuclease derived from archaea, and ISDra2 TnpB, a

TnpB-family nuclease derived from bacteria. We applied directed evolution in a fluorescent

reporter system to isolate variants with increased HDR efficiency. Through iterative rounds

of mutagenesis and screening, we assembled combinatorial variants that show robust HDR

improvements achieving up to 11-fold increase in editing efficiency. The resulting variants

also exhibit increased NHEJ activity. We also demonstrated the portability of the best

Un1Cas12f1 variant by turning it into an adenine base editor with up to a 10-fold higher

editing efficiency relative to the engineered Un1Cas12f1 baseline. The engineered

"Cas12f1Super" and "TnpBSuper" enzymes reported here represent promising tools for

efficient genome editing in mammalian systems for both fundamental research and

therapeutic applications.

Results

Directed evolution of Un1Cas12f1 and ISDra2 TnpB for enhanced activity in mammalian cells

To enable directed evolution of compact nucleases, we first developed a fluorescent HDR reporter system for selecting, screening, and quantifying HDR-active variants in mammalian cells (**Fig. 1a**). This reporter contained a disrupted start codon in the *EGFP* gene, making *EGFP* expression dependent on successful HDR-mediated repair. A customizable protospacer and PAM/TAM directly upstream of *EGFP* allow precise targeting by various single guide RNAs (sgRNAs) or ωRNAs and assessment of HDR activity across different target sequences. These reporters (one for Un1Cas12f1 and one for ISDra2 TnpB) were polyclonally integrated into HEK293T cells at moderate multiplicity of infection, selecting for blasticidin resistance present in the cassette. For ease of recognition, we refer to these protospacer/PAM/TAM combinations as "Target CM1" for Un1Cas12f1 and "Target TB1" for ISDra2 TnpB (see Methods).

To evaluate the baseline HDR activity of Un1Cas12f1 and ISDra2 TnpB proteins in this format, we delivered each nuclease by lentiviral transduction with puromycin selection and subsequently electroporated both a plasmid expressing guide RNA (sgRNA for Un1Cas12f1 and ωRNA for TnpB) and a single-stranded oligodeoxynucleotide (ssODN) donor template for *EGFP* sequence repair. The percentage of EGFP-positive cells served as a readout of HDR efficiency. ISDra2 TnpB was tested in its wild-type (WT) form, hereafter referred to as TnpB. For Un1Cas12f1, we tested a recently engineered version that combines CasMINI V3.1[3] with the ge4.1 guide scaffold[6], both of which were previously shown to improve activity. For simplicity, we hereafter refer to this combination as CasMINI. Both CasMINI and TnpB exhibited detectable HDR at their respective targets (Target CM1 or Target TB1) as measured by EGFP-positive cell percentage (CasMINI: 16.1±3.6% and TnpB: 6.3±0.5%) only once both sg/ωRNA and ssODN were delivered (**Fig. S1a**). HDR alleles were verified for both systems by amplicon next-generation sequencing (ampliconNGS) (**Fig. S1b**). This established a foundation for using this reporter system for the selection of activity-enhancing mutations.

To perform directed evolution, we first performed error-prone polymerase chain reaction

(epPCR) to generate libraries of 1.5×10^7 variants for CasMINI and 4.5×10^7 variants for

TnpB, which is an average of ~2 amino acid–altering mutations per enzyme. We deliberately

chose a low mutagenesis rate to avoid the accumulation of deleterious changes that could

mask beneficial effects. As a result, about half of each library remained unmutated WT

sequence (**Fig. S2a**). Each library was delivered to cells harboring its corresponding Target

CM1 or Target TB1 within the *EGFP* reporter system. Cells were electroporated with a

plasmid expressing sg/ωRNA targeting the reporter along with ssODN donor to restore

EGFP expression. EGFP-positive cells were isolated by fluorescence-activated cell sorting

(FACS), and their genomic DNA was used for targeted PCR amplification of active variants

for cloning into a new plasmid library for the next round of selection (**Fig. 1b**).

We performed two initial rounds of selection with intervening error-prone PCR to increase

library diversity (**Fig. 1b**). Bulk library activity dropped during these rounds, consistent with

early rounds of selection that had not yet enriched highly active variants (**Fig. 1c**). We then

carried out two additional rounds of purifying selection without further mutagenesis, during

which EGFP levels gradually increased, indicating an enrichment for more active variants.

After rounds 1, 2, and 4 of selection we amplified and tagmented the nuclease construct and

used NGS to determine the unphased mutation spectrum of the bulk pool. For both CasMINI

and TnpB we found multiple amino acid changes that were steadily enriched compared to the

pre-selection libraries (**Fig. 1d**). In case of CasMINI we identified three clusters of enriched

mutations (clusters 1-3). Among them, CasMINI cluster 1 exhibited enrichment of mutations

at spatially proximal residues 52, 55, 66 (**Fig. 1d,e, S2b**), which are located near or within

hydrogen-bonding distance to the sgRNA. Additionally, we observed two clusters (residues

201, 206, 227, 244 and 499, 504, 506) containing at least one highly enriched mutation (**Fig.**

1d, S2b). Among the 39 most enriched mutations in CasMINI, 28 involved a change in

charge. Surprisingly for a DNA-binding protein, 24 of these selected mutations lost positive

charge or gained negative charge, including the most abundantly selected K52E, K55E, and

K506E mutations. E206K was also a highly enriched mutation, which is similar to the

previously reported E206R[16].

The selected TnpB pool exhibited a broader distribution of enriched mutations with only one

obvious cluster at positions 64, 66, 67, 73 (TnpB cluster 1), which the structure of TnpB

revealed to be in proximity to DNA (**Fig. 1d,e, S2c**). For TnpB, 17/34 most enriched

mutations involved a change in charge. In contrast to what we observed for CasMINI, only

one of these showed a loss of positive charge. The rest exhibited a loss of negative charge or

a gain of positive charge. This included mutations from TnpB cluster 1: E64K, E66K, E67K,

and E73K. Overall, enrichment of multiple mutations for both nucleases after selection for

HDR prompted us to investigate these mutations as potential drivers of HDR activity.

Generation of CasMINI and TnpB with enhanced activity

The unphased short-read sequencing we used to measure mutation abundance in the selected

pools cannot resolve co-occurring beneficial mutations. To develop high-performing

CasMINI and TnpB variants, we isolated and tested promising multi-mutation variants from

in cellulo evolution. Mutations from the best variants were combined and tested at multiple

reporter and endogenous targets (**Fig. 2a**). This strategy provided a structured path from a

diverse variant pool to functionally robust and generalizable genome editing configurations.

For CasMINI, we tested a total of 246 post-round four variants in HEK293T cells (88 tested

with stable expression using lentiviral delivery and 158 transiently expressed from plasmids,

Fig. S3a). We first tested these variants against the same target site used for selection (Target

CM1), resulting in eleven multi-mutation variants with a 1.2 to 1.6-fold increase in HDR

efficiency relative to CasMINI (**Fig. 2b, Supplementary Table 1**). To isolate activity drivers

from passengers, we cloned and tested 22 mutations from the promising variants as single

mutants (SMs) in the CasMINI background. We also tested 18 additional candidate SMs

identified as enriched by NGS analysis of the selected pool (**Fig. 1d**), resulting in a total of 40

tested mutations (**Fig. S3b**). We retested the top-performing 19 mutations against an

orthogonal target site (hereafter "Target CM2", see Methods) and identified eight mutations

that consistently enhanced HDR at both targets (**Fig. S3c**). We combined eight selected

mutations with high activity at both target sites into five distinct combinatorial variants (CVs)

(**Fig. 2c, Supplementary Table 1**). Rather than performing an exhaustive combinatorial

screen, we employed a structure-guided reduction strategy in which we cloned a "master"

variant with all selected mutations that individually enhanced activity and then derived sub-

variants by selectively reverting subsets of mutations to the parental amino acid. The design

was guided by structural proximity of mutations, with the aim of reducing the likelihood of

negative epistatic interactions between closely positioned residues. In individual testing using

plasmid expression, all CVs significantly outperformed both CasMINI and the individual
mutations at two independent target sites. Comparison of CV16, CV17 and CV18 showed
that having K52E alone (CV18) is more beneficial than having either K52E, K55E, R66E
altogether (CV16) or K52E paired with R66E (CV17). CV18 (six mutations) and CV20 (five
mutations) showed the highest levels of HDR, with CV20 achieving 5.3-fold improvement
relative to CasMINI at Target CM2 (**Fig. 2d**).

We selected CV20 for further studies (**Fig. 2d**) based on its high expression in HEK293T
cells (**Fig. S3d**) and target-independent activity. CV20 contains five additional mutations on
top of CasMINI (K52E, K129E, E206K, K227E, K506E), with at least one mutation from
each of the three clusters that we previously identified that map to the REC, WED and NUC
domains of CasMINI (**Fig. 2e**). We attempted to further improve CasMINI_CV20 using the
recently described EVOLVEpro machine learning framework[17], training on activity data
collected during single mutant testing (**Fig. S3c**). Eight different EVOLVEpro-suggested
point mutations introduced in the CV20 revealed no further improvements in activity (**Fig.**
S3e), and so we continued with CV20.

To assess the portability and contribution of the evolved mutations, we incorporated the
CV20 mutations into the WT Un1Cas12f1 background (Un1Cas12f1_CV20) instead of the
CasMINI background. We then compared the HDR activities of WT Un1Cas12f1,
Un1Cas12f1_CV20, CasMINI and CasMINI_CV20 at both Target CM1 and Target CM2.
We observed that the CV20 mutations greatly increased the activity of both WT Un1Cas12f1
and CasMINI (**Fig. S3f**). The HDR activity of CasMINI_CV20 was up to 11.4±1.2-fold
greater than that of WT Un1Cas12f1, so we selected it for later testing at endogenous sites.
We termed this final system with five coding mutations relative to CasMINI as Cas12f1Super
(**Fig. 2e**).

For TnpB, we tested a total of 311 post-round four variants in HEK293T cells (**Fig. S4a**, all
tested using plasmid expression). This yielded 12 variants with at least 1.4-fold improvement
(**Fig. 2f, Supplementary Table 1**) at the Target TB1 site used for selection. One variant,
designated V322 (five amino acid altering mutations) consistently exhibited the highest HDR
activity, achieving 2.4-fold improvement over TnpB. Notably, V322 contained E73K, which
was the most enriched mutation within the TnpB cluster 1 after four rounds of selection (**Fig.**
1d). To expand our search for stronger contributors, we cloned and tested 17 mutations from
the 12 TnpB variants, as well as 19 additional mutations identified by NGS of the selected

pool resulting in total of 36 SMs. In total, we identified 16 constructs that improved HDR by
at least a 1.9-fold over TnpB (**Fig. S4b**). We retested the top-performing variant and single
mutations at an orthogonal target (hereafter "Target TB2", see Methods) (**Fig S4c**). Several
SMs performed well at both target sites, and we selected a subset of these broadly effective
substitutions to incorporate on top of V322, yielding five CVs (**Fig. 2g, Supplementary**
Table 1). As with CasMINI, we sought to limit combining proximal mutations within the
same variant, based on the assumption that their effects might be redundant or negatively
epistatic. All CVs exhibited at least two-fold greater HDR than WT TnpB at both Target TB1
and Target TB2. We chose to move forward with the high-performing CV2 (**Fig. 2h**), which
carried nine mutations (**Fig. 2i**). We tested eight additional EVOLVEpro-suggested mutations
in the TnpB_CV2 background (**Fig. S4d**)[17], but none strongly increased activity.

Two final refinements of TnpB_CV2 and ωRNA further improved activity and were
incorporated into the final design. While CasMINI (and hence Cas12f1Super) was already
codon-optimized for mammalian cells, we performed selection with TnpB cloned with its
original bacterial codon usage. Mammalian codon optimization slightly improved the activity
of both TnpB and TnpB_CV2 across two targets (**Fig. S5a,b**). However, we observed no
great increase in the expression of TnpB_CV2 compared to TnpB in HEK293T cells **(Fig.**
S5c). A recently reported shorter version of the ωRNA (Trim2)[18] also marginally improved
activity for TnpB_CV2 at both target sites (**Fig. S5d,e**). We next explored an additional
refinement by combining the TnpBSuper mutations with the previously developed TnpBmax
construct design[19] to generate TnpBmaxSuper. However, the addition of TnpBmax
architecture did not improve activity at either of the two HDR reporters nor at two
endogenous sites (**Fig. S5f,g**), which is an unexpected result in light of prior observations
with TnpBmax as a nuclease[19]. We therefore did not incorporate this optimization in
subsequent experiments. We termed the final codon-optimized system with nine coding
mutations relative to TnpB and the Trim2 ωRNA as TnpBSuper **(Fig. 2i)**.

Cas12f1Super and TnpBSuper efficiently edit endogenous genomic loci

To evaluate whether the observed improvement in HDR for Cas12f1Super and TnpBSuper
translated to endogenous gene targets, we edited 13 distinct genomic sites in HEK293T cells
using plasmid-based delivery of nuclease and sg/ωRNA (**Fig. 3a**). Editing outcomes were

measured using ampliconNGS after enriching for both guide RNA and editor-positive cells
by FACS.

For Cas12f1Super we observed consistent and significant increases in HDR efficiency across
all seven sites (**Fig. 3b**), with improvements up to 4.0±0.8-fold relative to CasMINI at
TAOK1 (**Fig. S6a**) At one target site, *NLRC4*, absolute HDR efficiency reached up to 30±2%
(**Fig. 3b**). NHEJ rates were increased at five of seven targeted sites, with 2.5±1.0%-fold
improvement at *TAOK1* (**Fig S6b**) and reaching up to 56±1.4% at *TTR* (**Fig. 3c**). Next, we
evaluated the activity of Cas12f1Super in the absence of an ssODN donor to determine
whether enhanced activity would also be observed under these conditions. Across all three
endogenous sites tested, Cas12f1Super exhibited increased editing efficiency (**Fig. 3d**), with
improvements of up to 4.2±0.3-fold at *P2RX5* (**Fig. S6c**). At the *COL8A1* site, absolute
NHEJ editing efficiency reached 80±0.8% (**Fig. 3d**). Together, these results show that
Cas12f1Super is generally more active than CasMINI in mammalian cells for both HDR and
NHEJ outcomes.

Intrigued by the enhancement in activity of the evolved proteins, we compared the *in vitro*
activity of CasMINI and Cas12f1Super. We expressed and purified both nucleases and tested
cleavage at six different temperatures (ranging from 37°C to 55°C) at two targets (*KRT1* and
Target CM1) (**Fig. S7a-f**). We observed no significant differences in cleavage activity
between CasMINI and Cas12f1Super at any of the temperatures. The cleavage pattern also
remained the same in the evolved variant (**Fig. S7g,h**). The improvement in HDR and NHEJ
for CasMINI may therefore be unrelated to catalytic activity in the tube. The high activity of
these variants could instead reflect interactions with the mammalian cell environment, though
this remains to be tested and may be difficult to determine.

TnpBSuper demonstrated significant increases in HDR at four of six endogenous sites (**Fig.**
3e), with HDR improvements up to 5.0±0.5-fold over TnpB at *AGBL1* (Fig. S8a). NHEJ
efficiency was increased relative to TnpB at the same four sites (**Fig. 3f, S8b**). We next
assessed the activity of TnpBSuper in the absence of an ssODN donor, analogous to the
analysis performed for CasMINI and Cas12f1Super. Across all three endogenous sites tested,
TnpBSuper exhibited increased editing efficiency (**Fig. 3g**), with improvements of up to
2.1±0.1-fold at *TTR* (**Fig. S8c**) and absolute editing efficiency reaching 96±0.5% at the same
site (**Fig. 3g**). Overall, these results show that TnpBSuper outperforms TnpB in mammalian
cells across both HDR and NHEJ.

In order to further explore the potential of enhanced cutting efficiency of Cas12f1Super and
TnpBSuper, we tested all possible guides targeting exon 1 of the *TRAC* locus in HEK293T
cells, including nine guides for CasMINI and one guide for TnpB. For the majority of guides,
we observed efficient editing, with higher activity for the evolved variants compared to their
respective wild-type enzymes (**Fig. S8d**). Together, these results further support the enhanced
activity of the evolved systems across diverse guide sequences.

We next tested activity in the clinically relevant setting of primary human T cells. We
produced mRNAs for TnpB and TnpBSuper and co-electroporated them with synthetic
ωRNA and an appropriate ssODNs targeting three endogenous loci in activated primary T
cells. TnpBSuper exhibited consistent increases in HDR relative to TnpB at all three sites,
with improvements reaching up to 6.9±1.6-fold at *TTR* (**Fig. S8e**), and absolute HDR
efficiencies up to 5.1±0.1% at *TTLL11* (**Fig. 3h**). Enhancements in NHEJ were also
significant, with up to 7.9±0.6-fold improvement and levels reaching 52±4% at the *TTR* site
(**Fig. S8f, 3i**).

To benchmark our evolved systems, we compared their performance to that of the widely
used but much larger *S. pyogenes* Cas9 (*Sp*Cas9) nuclease. We selected two loci for CasMINI
system (*P2RX5* and *NLRC4*) and three loci for TnpB system (*EMX1*, *RUNX1*, and *TTR*) and
identified up to three SpCas9 guides per locus based on close proximity to CasMINI/TnpB
target sites and high Rule Set1 activity scores[20]. These guides were first evaluated for NHEJ
activity in mammalian cells (**Fig. S8g**), and the best-performing guide at each locus was
selected for direct comparison. We then compared SpCas9 side by side with
CasMINI/Cas12f1Super or TnpB/TnpBSuper for HDR efficiency (**Fig. 3j**). SpCas9
outperformed Cas12f1Super and TnpBSuper at most loci, consistent with over ten years of
development as a workhorse genome editing system. However, at the *TTR* site, TnpBSuper
achieved slightly higher HDR (19±0.4%) compared to SpCas9 (16±0.7%). At the *P2RX5*
locus, Cas12f1Super approached SpCas9 performance, with HDR efficiencies of 19±0.6%
and 21±0.5%, respectively. Cas12f1Super and TnpBSuper outperformed their parental
constructs in all cases.

The enhanced genome editing activity of Cas12f1Super and TnpBSuper could potentially
come at the cost of increased off-target effects. To evaluate this possibility, we employed
DISCOVER-Seq, which enables nuclease-agnostic off-target identification in mammalian
cells by monitoring the recruitment of DNA repair factors[21]. For each nuclease pair

(CasMINI/Cas12f1Super and TnpB/TnpBSuper), we tested two target sites: *KRT1* and *CLIC4*

for CasMINI/Cas12f1Super (**Fig. 3b**), and *TTR* and *RUNX1* for TnpB/TnpBSuper (**Fig. 3e**).

Consistent with the previously reported high fidelity of both CasMINI and TnpB, we

observed only a single off-target site for CasMINI at the *CLIC4* locus. NHEJ efficiency at

this off-target was comparable between CasMINI and Cas12f1Super (**Fig. 3k,l, Fig. S9a,b**).

Importantly, despite their enhanced on-target activity, neither Cas12f1Super nor TnpBSuper

exhibited additional off-target effects. These results demonstrate that the evolved, highly

active variants retain the high specificity of their parental nucleases.

Efficient in vivo editing using AAV delivery of Cas12f1Super and TnpBSuper

Encouraged by the performance of Cas12f1Super and TnpBSuper at endogenous loci in

HEK293T and T cells, we further tested these variants *in vivo* in the context of a

therapeutically relevant target gene. We selected *PCSK9*, which is an attractive target for *in*

vivo AAV delivery that would be enabled by the hyper-active miniature genome editors

developed here[22,23]. *PCSK9* encodes a liver-expressed protein that promotes degradation of

the low-density lipoprotein receptor (LDLR). Disrupting PCSK9's function leads to reduced

levels of circulating low-density lipoprotein cholesterol (LDL-C), a strategy that can be

exploited for treating familial hypercholesterolemia[24]. Since PCSK9 loss-of-function is

sufficient to reduce LDL-C, we leveraged the increased NHEJ activity we observed for

Cas12f1Super and TnpBSuper. We first screened for sgRNAs and ωRNAs targeting exons 2-

9 of *Pcsk9* in murine NIH-3T3 cells and identified five top-performing guides: three for

Cas12f1Super and two for TnpBSuper (**Fig. 4a,b**). These guides were then used to compare

the editing efficiency of *Pcsk9* by CasMINI, Cas12f1Super, TnpB, and TnpBSuper in the

murine hepatoma cell line Hepa 1-6. We found that Cas12f1Super outperformed CasMINI at

all three sites and by up to 2.0±0.13-fold at site 3 (**Fig. S10a**) and reaching 51±1.6% absolute

efficiency at site 7 (**Fig. 4c**). TnpBSuper outperformed TnpB at site 4 by 3.2±0.5-fold (**Fig.**

S10b), reaching 43±6% absolute efficiency (**Fig. 4d**).

For mouse *in vivo* experiments, we selected site 7 for CasMINI/Cas12f1Super and site 4 for

TnpBSuper. CasMINI/Cas12f1Super were delivered using a single-stranded AAV9

(ssAAV9) design, whereas TnpB/TnpBSuper were packaged in a self-complementary AAV9

(scAAV9) format, both based on previously optimized TnpBmax architectures[19] (**Fig. 4e**).

AAV vectors were administered via tail vein injection to healthy C57BL/6J mice and
circulating PCSK9 levels were quantified by ELISA at 3- and 6-week timepoints post-
injection. Editing efficiency in the liver was assessed at the 6-week timepoint (**Fig. 4f**). For
both systems, we observed significantly increased editing with Cas12f1Super and TnpBSuper
compared to their respective wild-type counterparts (**Fig. 4g,h**). Editing efficiencies in whole
liver reached 51±8% for Cas12f1Super and 61±6% for TnpBSuper, reflecting marked
improvements over the parental constructs. This improved activity was reflected in more
rapid drops in plasma PCSK9 protein levels for both Cas12f1Super and TnpBSuper, though
after 6 weeks all systems approached a consistent minimum amount of PCSK9 (**Fig. 4i,j**).

Together, these results demonstrate efficient in vivo editing of a therapeutically relevant
target by Cas12f1Super and TnpBSuper.

Improved variant dCas12f1Super outperforms dCasMINI as adenine base
editor

Encouraged by the high activity of our editing enzymes, we explored their potential
applications in base editing. The additional enzymatic machinery appended to Cas proteins
that enables base editing makes miniature editors particularly attractive in this context.
Following the example of an existing dCasMINI-based adenine base editor[3], we introduced
two inactivating mutations (D326A, D510A) to generate dCasMINI and dCas12f1Super and
fused TadA* from ABE8e to their N-termini[25]. We individually co-delivered these base
editors to HEK293T cells as plasmids together with sgRNA-encoding plasmids targeting the
genomically integrated Target CM1 or three endogenous loci (*TAOK1*, *P2RX5* and *TTR*)
(**Fig. 5a**). We measured base editing using ampliconNGS. dCas12f1Super consistently
outperformed dCasMINI fusion at all four sites, with up to a 10±2.5-fold increase at *TAOK1*
(**Fig. S11a**) and reaching 17±0.1% absolute base editing at *TTR* (**Fig. 5b**). The per-base
editing window remained similar between dCasMINI and dCas12f1Super, but with markedly
increased activity for the variant (**Fig. 5c-f**). We observed almost exclusively A-to-G
conversion, even at the most efficient *TTR* site (**Fig. 5g**). Notably, we observed zero indels
even at the most efficient *TTR* target site, consistent with the use of catalytically inactive
editor (**Fig. 5h, Fig. S11b**) This activity and product purity is encouraging for the potential
use of Cas12f1Super beyond its evolved HDR and NHEJ context.

Discussion

Progress in genome editing relies on the development of more active tools for mammalian (human) systems. While rational design allows direct testing in human cells, it is constrained by limited variant throughput and the challenge of predicting beneficial mutations from structure alone. Directed evolution in bacterial systems offers high throughput, but nucleases evolved in bacteria can underperform in human cells. Several studies have reported false positives, where mutations that appeared promising in bacterial cells failed to translate to human cells[26–28]. This may be due to differences in cellular context, including chromatin structure, DNA repair pathways and post-translational modifications.

These differences have motivated the development of eukaryotic platforms for directed evolution. Yeast-based systems partially address this gap[29], but the biological differences between yeast and human cells still make functional divergence likely. DMS in human cells, such as a recent screen of AsCas12f1, illustrates the power of selection directly in the relevant context[8]. However, DMS in the context of human cell culture is only feasible for single point mutations and does not capture the potential of synergistic combinations that may emerge during selection. In this work, we developed a mammalian-based selection strategy to enrich for highly active multi-mutant variants and applied it to CasMINI and TnpB, yielding Cas12f1Super and TnpBSuper with strongly enhanced activity in human cells.

For Cas12f1Super, four out of five mutations involve a switch from positively to negatively charged residues. One example, K52E, lies near the sgRNA scaffold and improves activity despite losing a positive charge which is an unexpected result given the presumed importance of electrostatic interactions in that region. Given K52's proximity to the sgRNA-interacting R66, its beneficial effect may result from stabilizing sgRNA binding by buttressing R66. The other charge-altering mutations could also reconfigure the protein to enhance nucleic acid engagement, though such effects are not readily apparent from the nuclease structure. E206K adds a positive charge near DNA, and a similar substitution (E206R) was previously proposed in a rational design study to enhance DNA binding[16]. In TnpBSuper, five of nine mutations originate from a single variant straight from selection (V322). Among the remaining mutations, E73K is positioned near DNA and may participate in protein-DNA interactions. Others include conservative changes such as L155V, L172V, V192L, and

G299A. L172 is involved in stabilizing the TAM-proximal end of the heteroduplex[30],
suggesting that L172V may support more efficient duplex formation. While this manuscript
was in revision, deep mutational scanning of ISDra2 TnpB individual point mutations in
yeast identified some overlapping beneficial mutations (e.g. V192L) and some partially
convergent mutations (e.g. L172G vs. L172V)[31]. However, the majority of mutations reported
here are distinct from those identified by yeast-based selection. These differences likely
reflect distinct selection environments (yeast vs. mammalian cell biology), non-saturating but
combinatorial mutagenesis in our approach, and locus-dependent effects.

Our mammalian selection strategy focused on promoting HDR without directly suppressing
NHEJ and the resulting variants showed higher HDR and NHEJ outcomes. Explicitly
including HDR during selection and screening yielded TnpB and Cas12f1 variants with high
overall activity, retaining the ability to perform both error-prone and precise editing.
Cas12f1Super exhibited identical *in vitro* cleavage patterns and enzymatic parameters to
CasMINI, and suggesting that its improved activity might stem from mammalian cell-specific
factors such as increased expression, greater protein stability, or even interaction with the
DNA repair machinery. Additionally, altered trans ssODN-cleavage activity of these
nucleases could affect the availability of the donor template during repair. Nevertheless, these
mechanisms remain speculative and require additional investigation.

In summary, our work demonstrates the power of mammalian-based selection to evolve
compact, highly active genome editors tailored for function in complex cellular
environments. The functional improvements observed in CasMINI and TnpB systems after
their engineering in mammalian cells, including enhanced HDR, NHEJ, and base editing
activity, together with their compatibility with AAV delivery, highlight their utility for broad
applications in therapeutic genome engineering.

Methods

Cell culture

HEK293T, NIH-3T3, Hepa 1-6, and HEK293T reporter cell lines were cultured in
Dulbecco's Modified Eagle Medium (DMEM) with GlutaMAX (Thermo Fisher Scientific),
supplemented with 10% fetal bovine serum (FBS, Thermo Fisher Scientific) and 100 U/mL
penicillin-streptomycin (Gibco). HEK293T and NIH-3T3 cells were obtained from certified

vendors, and Hepa 1-6 cells were provided by the laboratory of Prof. Dr. Markus Stoffel

(ETH Zürich). These cell lines were maintained at 37°C in a humidified incubator with 5%

CO_2 and routinely tested for mycoplasma contamination using the MycoAlert kit (Lonza).

Cells were never used beyond passage 20. Primary human T cells were isolated from

Leukopaks (StemCell Technologies) using the EasySep™ Human T Cell Enrichment Kit, and

activated by culturing in T25 flasks coated overnight at 4 °C with anti-CD3 (clone OKT3,

functional grade; Thermo Fisher Scientific) and anti-CD28 (clone CD28.2, BioLegend)

antibodies (1 µg/mL each in ultrapure H_2O). Prior to electroporation, activated T cells were

maintained for 48 hours in CTS medium (Advanced RPMI 1640 (44.5%), Click's Medium

(44.5%), FBS (10%), and 200 mM L-glutamine (1%)) supplemented with IL-7 (10 ng/mL)

and IL-15 (5 ng/mL).

HDR reporter cell line generation

Lentiviral packaging of HDR reporter constructs was performed in HEK293T cells using

polyethyleneimine (PEI) as the transfection reagent. The HDR construct plasmid was co-

transfected with the envelope plasmid (pCMV-VSV-G) and the packaging plasmid (pCMV-

dR8.2 dvpr) at a 1:1:1 molar ratio. Viral supernatant was collected 48 hours post-transfection,

filtered through a 0.45 µm filter, and used to transduce HEK293T cells with different HDR

reporters. Reporter construct with Target CM1 was transduced with MOI of 6, while all other

reporter constructs (Target CM2, Target TB1, Target TB2) were transduced with MOI of 0.2.

Twenty-four hours after transduction media was replaced. Forty-eight hours after

transduction, blasticidin selection was initiated at concentration of 6 µg/mL and continued for

3 weeks to establish stable cell lines.

Cloning variant libraries

Libraries were generated by error-prone PCR (epPCR) using the GeneMorph II kit (Agilent)

on CasMINI or TnpB-containing amplicons. Conditions were optimized to introduce ~2

amino acid-changing mutations per open reading frame (ORF). PCR products and destination

vectors were digested with AscI and BamHI for 4 hours, followed by overnight ligation at

16°C. Ligated products were isopropanol-precipitated and electroporated into MegaX

DH10B cells (Thermo Fisher Scientific) using a Bio-Rad electroporator. After a 1-hour

recovery in SOC medium, transformed cells were plated on large square agar plates and

incubated overnight at 30°C. Colonies were harvested, and plasmid DNA was purified using

a Qiagen Midiprep kit.

Workflow for HDR-enriched variant selection

Variant libraries and corresponding wild-type (WT) plasmids were packaged into lentivirus using PEI-mediated co-transfection with the envelope plasmid (pCMV-VSV-G) and the packaging plasmid (pCMV-dR8.2 dvpr) in HEK293T cells at a 1:1:1 molar ratio. Lentiviral supernatants were used to transduce HEK293T HDR reporter cells. Forty-eight hours post-transduction, cells underwent two consecutive 2-day puromycin selection rounds (concentration 1.5 µg/ml). After 1 day of recovery, cells were electroporated with a guide RNA targeting the HDR reporter (Target CM1 for CasMINI or Target TB1 for TnpB) and an ssODN donor template to restore *EGFP* expression. Electroporations were performed using Lonza Amaxa 4D Nucleofector in 100 µL single Nucleocuvettes. Three days post-electroporation, EGFP-positive cells were isolated by FACS. Genomic DNA was extracted using the Gentra Puregene Cell Kit (Qiagen). Variants were PCR-amplified from genomic DNA, digested with AscI and BamHI, and cloned to generate the next library generation. The primers for amplifying variants from genomic DNA are listed in **Supplementary Table 2.**

Design of ssODN donors

All ssODNs used for HDR at reporter and endogenous targets were 180 nucleotides long, complementary to the targeted strand, and symmetric around the cut site. For CasMINI or TnpB HDR reporter targets, the ssODNs included the 2-nucleotide substitution required to restore the *EGFP* start codon, along with an additional modification within the protospacer (positions 11–15 relative to the PAM or TAM sequence for CasMINI or TnpB, respectively). For CasMINI or TnpB endogenous targets, ssODNs were designed to include: (1) a 2-nucleotide substitution near the cut site (positions 23 & 24 relative to the PAM for CasMINI, or positions 17 & 18 relative to the TAM for TnpB); (2) a substitution within the protospacer (positions 11–15 for CasMINI, or 11–16 for TnpB); and (3) a PAM/TAM mutation (TTTR→TGTR for CasMINI, or TTGAT→TTGTT for TnpB). The protospacer and PAM/TAM modifications were included to prevent re-targeting following successful HDR. Final editing efficiency was quantified as the percentage of reads containing the ATG restoration (when editing HDR reporter cell lines) or the 2-nucleotide substitution near the cut site (when editing endogenous targets in HEK293T cell line). For SpCas9 endogenous targets, ssODNs were designed to include (1) a 2-nucleotide substitution near the cut site (positions 3 & 4 relative to the PAM); and (2) a PAM mutation (NGG→NTG).

492   ssODNs were ordered as Ultramer DNA oligos from Integrated DNA Technologies (IDT).

493   For electroporating T cells, chemically protected ssODNs were used (ordered with four

494   phosphorothioate modifications at the 5' and 3' ends). The sequences of ssODNs are listed in

495   **Supplementary Table 3**.

Testing workflow for HDR-enriched variant selection

The workflow for HDR-enriched variant selection was tested at Target CM1 with CasMINI.

CasMINI was delivered via lentivirus, produced by PEI-mediated co-transfection of the

envelope plasmid (pCMV-VSV-G), the packaging plasmid (pCMV-dR8.2 dvpr) and the

CasMINI expression plasmid into HEK293T cells at a 1:1:1 molar ratio. The resulting

lentiviral supernatant was used to transduce HEK293T HDR reporter cells (with Target

CM1). Forty-eight hours post-transduction, cells underwent two consecutive rounds of 2-day

puromycin selection (concentration 1.5 µg/ml). After 1 day of recovery, cells were

electroporated with a guide RNA targeting the HDR reporter (Target CM1) and an ssODN

donor template to restore *EGFP* expression. After 3 days cells were collected for flow

cytometry analysis and genomic DNA extraction for subsequent ampliconNGS.

Flow cytometry and fluorescence-activated cell sorting (FACS)

For all flow cytometry and FACS experiments cells were harvested 3 days post-transfection

by dissociation with 0.05% Trypsin and resuspension in PBS supplemented with 10% FBS.

Analytical flow cytometry for quantifying editing at HDR reporter targets in HEK293T-

derived cell lines was performed using an Attune NxT Flow Cytometer (Thermo Fisher

Scientific), and data were analyzed using FlowJo software (version 10.10.0). For all

experiments in HEK293T cells at HDR reporter targets except for experiment involving

TnpBmax in **Fig. S5f** cells were sequentially gated for live cells (FSC/SSC), singlets (FSC-

A/FSC-H), BFP+, mCherry+, and finally EGFP+ populations **(Fig. S12a)**. For experiment

involving comparison with TnpBmax in HEK293T cells in **Fig. S5f** cells were sequentially

gated for live cells (FSC/SSC), singlets (FSC-A/FSC-H), mCherry+, and finally EGFP+

populations **(Fig. S12b)**. FACS for analysis of editing at endogenous sites in HEK293T,

NIH-3T3 and Hepa 1-6 cell lines and FACS during big scale selections in HEK293T cells

was performed using a BD FACSAria Fusion or BD FACSAria III. In case of FACS for

analysis of editing at endogenous sites in HEK293T involving comparison with TnpBmax

(**Fig. S5g**), editing at TRAC locus (**Fig. S8d**) and comparison with SpCas9 (**Fig. 3j**, **S8g**)

cells were sequentially gated for live (FSC/SSC), singlets (FSC-A/FSC-H), and finally

mCherry+ populations **(Fig. S12b),** then 1×10^5 cells were processed for genomic DNA

extraction and subsequent ampliconNGS**.** In case of FACS for analysis of editing at

endogenous sites in Hepa 1-6 cell line, cells were sequentially gated for live (FSC/SSC),

singlets (FSC-A/FSC-H), and finally mCherry+ populations **(Fig. S12b),** then 1.2×10^4

mCherry+ cells were sorted, cultured for 5 days, and 1×10^5 cells were processed for

genomic DNA extraction and subsequent ampliconNGS. In case of FACS for analysis of

editing at endogenous sites in HEK293T (except for comparison with TnpBmax in **Fig. S5g,**

editing at TRAC locus in **Fig. S8d** and comparison with SpCas9 in **Fig. 3j**, **S8g**) and in NIH-

3T3 cell lines, cells were sequentially gated for live (FSC/SSC), singlets (FSC-A/FSC-H),

and finally mCherry+/BFP+ population **(Fig. S13a)**, then 1×10^5 BFP$^+$/mCherry$^+$ double

positive cells were sorted and processed for genomic DNA extraction and subsequent

ampliconNGS. In case of FACS during big scale selections in HEK293T cell line, cells were

sequentially gated for live cells (FSC/SSC), singlets, SYTOX$^{\text{TM}}$-Red-negative, and finally

mCherry+/EGFP+ populations **(Fig. S13b)**, then all EGFP-positive cells were sorted from 1.1

$\times 10^8$ mCherry+ cells during round one and two of selection or from 2×10^7 mCherry+ cells

during rounds three and four of selection and processed for genomic DNA extraction.

mRNA synthesis for T cell electroporation

mRNA encoding either TnpB or TnpBSuper was synthesized using the mMESSAGE

mMACHINE™ T3 Transcription Kit (Thermo Fisher AM1348). The expression plasmid

containing the T3 promoter was linearized, purified, and used for *in vitro* transcription. The

resulting mRNA was purified with Monarch Spin RNA Cleanup Kit (New England Biolabs

T2050L) and quantified by Qubit using the Qubit RNA Broad Range Kit (Thermo Fisher

Q10210). The integrity of the mRNA constructs was verified with an automated

electrophoresis system (Agilent).

Electroporation of mammalian cells

To assess editing at endogenous sites in HEK293T, HEK293T-derived, and NIH-3T3 cell

lines, 1-2 $\times$ 10 cells were electroporated per reaction with a total of 600 ng plasmid

DNA at a 1:1 molar ratio of nuclease-expression and guide RNA-expression plasmids,

carrying mCherry and BFP markers, respectively. When the nuclease was stably expressed in

HEK293T-derived lines, 200 ng of guide RNA-expressing plasmid (without a fluorescent

marker) was used. 100 pmol of ssODN was co-electroporated when transfecting HEK293T

and HEK293T-derived lines. For editing in activated primary T cells, 1×10^6 cells were

electroporated with 150 pmol of synthetic ωRNA (modified with 2'-O-methyl analogs on

the first and last three bases and 3' phosphorothioate internucleotide linkages between the

first three and last two bases; Synthego), 4 µg of mRNA encoding the nuclease and 200 pmol

of chemically protected ssODN. Electroporations were performed using the Lonza Amaxa 4D

Nucleofector with 20 µL Nucleocuvette Strips. Cell lines were electroporated with SF

buffer using program DG-130 (HEK293T and derivatives) or SG buffer using EN-158 (NIH-

3T3). T cells were electroporated with P3 buffer using program EH-115. Following

electroporation, HEK293T, HEK293T-derived, and NIH-3T3 cells were cultured for 3 days

before flow cytometry or FACS. For T cells, the medium was exchanged 24 hours post-

electroporation, and genomic DNA was harvested 3 days after the media change.

Lipofection of mammalian cells

567 To assess editing at endogenous sites in the Hepa 1-6 cell line, 1.5×10^5 cells were seeded

568 per well in a 24-well plate and transfected with 1000 ng of an "all-in-one" plasmid expressing

569 both the protein and guide RNA, along with an mCherry marker, using Lipofectamine 3000

570 (Thermo Fisher). Cells were cultured for 3 days prior to FACS.

Base editing

Two inactivating mutations (D326A, D510A) were introduced to CasMINI or Cas12f1Super

to get catalytically inactive variants: dCasMINI and dCas12f1Super. All sites in HEK293T

cells were base-edited by co-electroporation of dCasMINI or dCas12f1Super-expressing

plasmid and sgRNA-expressing plasmid as described above in the "electroporation of

mammalian cells" section of these Methods. Cells were collected for ampliconNGS 3 days

after transfection.

Amplicon next-generation sequencing (ampliconNGS)

Three days post-transfection (4 days for T cells), cells were dissociated, resuspended in PBS

+ 10% FBS, and 1×10^5 cells were pelleted and resuspended in 50 µL QuickExtract solution

(Lucigen). Genomic DNA was extracted by incubation at 65°C for 30 min, followed by 98°C

for 15 min. Amplicon libraries were prepared using a two-step PCR protocol. In PCR1, the

target locus was amplified with primers containing Illumina adapter sequences. Products

were validated on agarose gel and purified using SPRI beads (0.9× volume). In PCR2,

indexing primers were added. Final libraries were bead-purified (0.7× volume), pooled, and

sequenced using paired-end 2×150 bp reads on an Illumina platform. NGS data were

analyzed using CRISPResso2 (v2.3.1)[32]. For analysis of base editing outcomes overall editing

was calculated as a percentage of A→G conversion in all reads that had had at least 0.1%

representation. The primers used for ampliconNGS are listed in **Supplementary Table 2**.

Tagmentation of libraries

Amplicons corresponding to either the original sequences of CasMINI or TnpB (referred to

as "Ctrl"), or variants from libraries collected before selection (termed "before_R1") and

after rounds 1, 2, or 4 of selection ("R1," "R2," and "R4," respectively), were generated by

PCR. Each of these amplicons (1 ng input) was used to prepare sequencing libraries using the

Nextera XT DNA Library Preparation Kit (Illumina), which includes enzymatic

fragmentation and adapter tagging. The resulting libraries were subjected to high-throughput

sequencing on an Illumina platform.

Tagmented library analysis

Quality of raw sequencing files was checked with fastqc v0.11.9[33] with default arguments

and reports were combined using multiqc v1.12[34]. Raw reads were aligned to the respective

custom reference with bowtie2 v2.4.4[35] and arguments -p 60 and -local. Samtools v1.16.1[36]

was used to convert the alignments to BAM format, sort by genomic location and generate

indices. The callvariants.sh tool from BBMap v38.95[37] was used to call variants from all

alignments from the same reference in multisample mode with the arguments multisample =

t, ploid = 1, clearfilters, trimq = 25, minqualitymax = 25 and minmapqmax = 5. Variants

were further filtered using bcftools v1.15.1[38] with argument -v snps to only include SNPs.

Variants were annotated using ensemble-vep v112.0[39] using its docker image with custom

references and default arguments. The annotated SNPs were further processed with a custom

script in R v4.3.1[40]. Only SNPs with an allele fraction greater than or equal to 0.003 after R4

and with a before_R1/R4 ratio greater than 3.77 were kept. Plots were generated with base R

plotting functions or with ComplexHeatmap v2.18[41].

Western blot analysis

Briefly, HEK293T cells were electroporated with HA-tagged CasMINI/CasMINI_CV20 or

TnpB/TnpB_CV2-expressing plasmid and sg/ωRNA-expressing plasmid targeting Target

CM1 (for CasMINI/CasMINI_CV20) or Target TB1 (for TnpB/TnpB_CV2) in triplicate.

Electroporations were carried out using the Lonza Amaxa 4D Nucleofector system with

20 µl Nucleocuvette Strips. After 3 days, cells were collected and lysed in RIPA buffer

(50 mM Tris-HCl, 150 mM NaCl, 0.25% deoxycholic acid, 1% NP-40, 1 mM EDTA, pH

7.4) supplemented with protease inhibitors (Halt protease inhibitor cocktail, Thermo Fisher

Scientific). Proteins were separated by polyacrylamide gel electrophoresis (PAGE) and

transferred onto 0.2 µm nitrocellulose membranes using standard wet transfer methods.

Membranes were blocked in TBS-T (150 mM NaCl, 20 mM Tris, 0.1% Tween-20, pH 7.4)

containing 5% skim milk, then incubated with primary antibodies diluted in TBS-T

supplemented with 5% BSA and 0.05% sodium azide. The primary antibodies used were

anti-HA (Cell Signaling Technology, 3724) and anti-GAPDH (Cell Signaling Technology,

2118), both at 1:1,000 dilution. Detection was performed using fluorescent secondary

antibody (LI-COR Biosciences, 926-32213) at 1:10,000 dilution. Blots were imaged using the

Odyssey CLx scanner (LI-COR Biosciences).

EVOLVEpro

The identities and relative activities of the best-performing single mutants of CasMINI (16

point mutants) and TnpB (7 point mutants) were used as input for EVOLVEpro

(https://github.com/mat10d/EvolvePro)[17]. The top eight predicted activity-enhancing

mutations for CasMINI and TnpB were individually cloned onto the Cas12f1Super and

TnpBSuper backgrounds. These new variants were tested in HEK293T cells at Target CM1

and Target CM2 (for CasMINI variants) or Target TB1 and Target TB2 (for TnpB variants)

via co-delivery of a nuclease-expressing plasmid, a sg/ωRNA-expressing plasmid and

ssODN.

DISCOVER-Seq

HEK293T cells (2×10^7) were transfected with 120 µg of each "all-in-one" construct using

Lipofectamine 3000 (Thermo Fisher). Fifteen hours post-transfection, cells were pelleted and

resuspended in room temperature RPMI (without supplements). Cells were crosslinked with

1% formaldehyde (Thermo Fisher) and incubated for 15 min at room temperature.

Formaldehyde was quenched with 125 mM glycine for 3 min on ice. Cells were then pelleted

at 4°C, washed twice with ice-cold PBS, and snap frozen in liquid nitrogen. Pellets were

stored at -80°C prior to processing. MRE11 ChIP-Seq was performed next as described in

Wienert et al[42]. Briefly, samples were thawed on ice and lysed using ice-cold buffers LB1,

LB2, and LB3. The isolated DNA was sonicated to obtain ~300 bp chromatin fragments

using a Covaris S2 with the following settings: 12 cycles of duty cycle 5 %, intensity 5, 200

cycles per burst for 60 s. For each chromatin immunoprecipitation (ChIP), 10 µl of MRE11

antibody (PA3-16527R; Thermo Fisher) per ChIP was prebound to protein A Dynabeads

(Invitrogen). MRE11-associated chromatin was immunoprecipitated with antibody-bound

beads, rotating overnight at 4°C. Dynabeads were washed with RIPA buffer and the DNA

was eluted by incubating overnight at 65°C in elution buffer containing SDS. For the final

clean-up, the samples were digested with Proteinase K and RNase A in TE buffer for 1 h at

55°C. DNA fragments were purified using the MinElute Kit (Qiagen). Sequencing libraries

were prepared using the NEBNext Ultra II workflow (New England BioLabs). Indexed

libraries were pooled and sequenced on a NovaSeqX (Illumina) with 2×150 paired-end reads

and a target depth of 40 million reads per sample. To identify genome editing target sites,

NGS reads were first trimmed for the Illumina adapter sequences using cutadapt[43] (v4.6),

followed by alignment to the human genome (hg38) using bowtie2[44] (v2.5.2). Resulting

alignments were then deduplicated and subsampled to 40 million reads per sample using

samtools[38] (v1.19). Off-target site calling was performed using BLENDER[21,42]

(https://github.com/cornlab/blender) with custom parameters followed by manual inspection.

Plasmids used for DISCOVER-Seq are listed in **Supplementary Table 4.**

CasMINI and Cas12f1Super purification

Both nucleases (His^6-GFP-2xNLS-CasMINI/Cas12f1Super) were expressed from plasmids

listed in **Supplementary Table 4**. Plasmids were transformed into *Escherichia coli* BL21

(DE3) cells, and protein expression was induced at an optical density (OD□□□) of 0.5 by

adding IPTG to a final concentration of 0.5 mM. Cultures were then incubated at 18°C for 18

h. Cells were harvested and resuspended in buffer A (25 mM Tris-HCl (pH 7.6), 1 M NaCl, 5

mM β-mercaptoethanol) supplemented with 5% glycerol, 1 mM PMSF, 1 µg/ml pepstatin A,

and 1 µg/ml leupeptin. Lysis was performed by sonication, and the lysates were clarified by

centrifugation at 40,000 × *g* for 30 min. The supernatant was applied to Ni-NTA Superflow

resin (Qiagen) pre-equilibrated with buffer A without supplements, washed with buffer A

containing 25 mM imidazole, and eluted with buffer A containing 250 mM imidazole. The

His[6]-GFP tag was removed by overnight digestion with TEV protease at 4°C. The resulting

protein solution was diluted to reduce NaCl concentration to 0.2 M and loaded onto a HiTrap

SP HP column (Cytiva) for ion exchange chromatography, with elution performed using a

linear NaCl gradient (0.2–2 M). Finally, the protein fractions containing 2xNLS-

CasMINI/Cas12f1Super were concentrated in a 30kDa molecular weight cut-off Amicon

Ultra-15 Centrifugal Filter Unit (Merck) and further purified by size exclusion

chromatography using a Superdex 200 column (GE Healthcare) pre-equilibrated with 25 mM

Tris (pH 7.6), 200 mM NaCl, 2 mM DTT, 1 mM $MgCl_2$. Fractions containing the target

protein were pooled, diluted to a final concentration of 0.24 mg/ml, and stored at –80°C.

sgRNA synthesis for plasmid DNA cleavage assay

The single guide RNA (sgRNA) templates were generated by two rounds of PCR using

overlapping oligonucleotides containing the T7 promoter, sgRNA scaffold, and 20 nt

targeting sequence. The obtained templates were then used for *in vitro* transcription (IVT)

using the TranscriptAid T7 High Yield Transcription Kit (Thermo Fisher Scientific). The

IVT reaction products were purified using the GeneJET RNA Purification Kit (Thermo

Fisher Scientific). Oligonucleotides used for the template generation and sequences of the

sgRNA templates, are listed in **Supplementary Tables 2** and **5, respectively**.

CasMINI/Cas12f1Super-sgRNA complex assembly

CasMINI and Cas12f1Super ribonucleoprotein (RNP) complexes were assembled by mixing

CasMINI or Cas12f1Super protein with sgRNA at 2:1 molar ratio and incubating at 37°C for

30 min in 10 mM Tris–HCl (pH 7.5 at 37°C), 100 mM NaCl, 1 mM DTT, and 1 mM EDTA

buffer.

Plasmid DNA cleavage assay

Plasmid DNA cleavage reactions were performed by mixing 100 nM of CasMINI-sgRNA

complex with 3 nM of plasmid DNA (pTK1910 or pTK1911) in 2.5 mM Tris-HCl (pH 7.5 at

37°C), 25 mM NaCl, 10 mM $MgCl_2$, 0.25 mM DTT, and 0.25 mM EDTA reaction buffer.

The samples were incubated at 37-55°C, and the aliquots, removed at timed intervals (0, 1, 2,

5, 10 15, and 30 min), were mixed with 3×loading dye solution containing 75 mM EDTA,

0.1% SDS, and 0.03% Bromophenol Blue in 50% (v/v) glycerol. The samples were analyzed

by running agarose gel electrophoresis and quantified using GelAnalyzer (v23.1.1). The

linear cleavage products from the 30-minute endpoint aliquots were purified and subjected to

run-off Sanger sequencing to determine the cleavage patterns on both the target and non-

target DNA strands.

Adeno-associated virus production

AAV9 vectors were produced by the Viral Vector Facility at the Neuroscience Center Zurich.

In brief, vectors were purified by ultracentrifugation followed by diafiltration. Physical titers

(vg⬜ml⬜¹) were quantified using a Qubit 3.0 fluorometer (Invitrogen).

CasMINI/Cas12f1Super were packaged in a single-stranded AAV9 (ssAAV9) format,

whereas TnpB/TnpBSuper constructs were packaged in a self-complementary AAV9

(scAAV9) format.

Mouse studies

Animal experiments were performed following protocols approved by the Kantonales

Veterinäramt Zürich and in compliance with the Swiss Animal Welfare Act and Protection

ordinance. Mice (C57BL/6J) were housed in a pathogen-free animal facility at the Institute of

Pharmacology and Toxicology at the University of Zurich and kept in a temperature- and

humidity-controlled room (21⬜°C, 50% RH) on a 12-hour light-dark cycle and fed a standard

laboratory chow (Kliba Nafag no. 3437 with 18.5% crude protein). AAV vectors were

administered via tail vein injection in 6-week-old mice at a dose of 5×10^{13} vg/kg (n=4 per

group, 2 males and 2 females). Mice were fasted for 3-4⬜hours before blood was collected

from the tail vein for the 3-week timepoint or the inferior vena cava after the endpoint of the

experiments.

Serum biochemistry

Mouse PCSK9 protein levels in mouse plasma collected at 3 weeks and 6 weeks (endpoint)

post-injection were determined using Mouse Proprotein Convertase 9/PCSK9 Quantikine

ELISA Kit (R&D Systems MPC900) according to the instructions provided by the

manufacturer. PCSK9 protein levels were normalized to those of untreated control mice. For

each control mouse the 3- and 6-week timepoints were first averaged, and the resulting per-

mouse values were then averaged across control mice to obtain a single normalization

reference. One control mouse was excluded due to highly variable measurements between

timepoints.

Plasmid cloning and mutagenesis

Lentiviral reporter constructs

EFα-TAM/PAM-protospacer-EGFP-PGK-BSD fragment was inserted in a lentiviral

backbone (Addgene ID 60954) using PCR and Gibson assembly. Protospacer is either Target

CM1 or Target CM2 (for CasMINI reporter constructs) or Target TB1 or Target TB2 (for

TnpB reporter constructs). *EGFP* gene contained a different triplet instead of a start codon:

GGG for CasMINI reporters, AGA for TnpB reporters. Sequences or Target CM1, Target

CM2, Target TB1, Target TB2 are provided in **Supplementary Table 2**.

Lentiviral CasMINI/TnpB expressing constructs

EF1α-2xNLS-CasMINI/TnpB-PGK-Puro-T2A-mCherry fragment was inserted in a lentiviral

backbone (Addgene ID 52962) using PCR and Gibson assembly. These constructs were used

as templates for epPCR for generating libraries of variants of CasMINI and TnpB. Mutations

for single mutants (SMs) and combinatorial variants (CVs) were cloned using PCR

mutagenesis and Gibson assembly.

sgRNA/ωRNA expressing constructs

Scaffolds of CasMINI or TnpB were cloned downstream from the human U6 promoter in a

pUC19 plasmid. ge4.1 scaffold (100 nt) was used for CasMINI. Either the original (230 nt) or

Trim2 scaffold (99 nt) was used for TnpB. The targeting part of the guide RNA sequence was

cloned downstream from the scaffold sequence. EF1α-BFP fragment was cloned downstream

from the guide. A version of these constructs without the EF1α-BFP part was used during

HDR selection. Sequences of guide RNA scaffolds are provided in **Supplementary Table 6.**

"All-in-one" construct for transient co-expression of TnpB and Trim2 ωRNA (also called

ωRNA*) was obtained from Addgene (ID 212967). "TnpBmax" design was cloned in in this

backbone using Gibson assembly. This backbone was modified to co-express CasMINI and

its sgRNA using Gibson cloning. "All-in-one" construct for transient co-expression of

SpCas9 and its sgRNA was obtained from Addgene (ID 48138).

AAV constructs

The ssAAV construct used for *in vivo* experiments was kindly provided by the laboratory

of Gerald Schwank. CasMINI or Cas12f1Super, together with the sgRNA, were cloned into a

"TnpBmax"[19] design using Gibson assembly.

The scAAV construct was obtained from the Viral Vector Facility of the University of Zurich

(UZH), Zurich. TnpB or TnpBSuper, along with the ωRNA, were cloned into a "TnpBmax"[19]

design using Gibson assembly.

mRNA expressing constructs

NLS-TnpB/TnpBSuper-NLS fragment was placed under T3 promoter for subsequent *in vitro*

transcription.

Base editor constructs for CasMINI

Two inactivating mutations (D326A, D510A) were introduced in CasMINI and

Cas12f1Super with PCR to get dCasMINI and dCas12f1Super. Next, in ABE8e construct

(Addgene ID 138489) dCas9 was replaced either with dCasMINI or dCas12f1Super.

Plasmid DNA substrates for cleavage assays

The plasmid DNA substrates (pTK1910 and pTK1911) for *in vitro* cleavage reactions were

obtained by cloning PAM and target containing oligoduplex into a pUC18-based plasmid

(pTK1909) through BpiI restriction endonuclease (Thermo Fisher Scientific) sites. The

sequences of oligonucleotides are available in **Supplementary Table 2**.

For all Gibson assemblies, the New England Biolabs (NEB) HiFi Assembly Master Mix was

used. All PCR reactions were performed using Q5 DNA Polymerase (NEB).

All plasmids used in this study are listed in **Supplementary Table 4.**

Statistical analysis

GraphPad Prism 10.5.0 software was used for statistical analysis. Results are presented as

mean ± standard deviation. All *P* values were calculated using an unpaired, two-sided *t*-test.

Holm–Šidák correction was used for multiple comparisons. Schemes were created with

Biorender.com. Protein structures were visualized with ChimeraX[45].

Data availability

Sequencing files for tagmented libraries and DISCOVER-Seq data are available in Sequence Read Archive BioProject PRJNA1286789.

Acknowledgements

We thank Dr. Florian Mair, Dr. Malgorzata Kisielow, Anette Schütz, Renan Antonialli, Dr. Irini Vgenopoulou of the Flow Cytometry Core Facility of the ETH Zurich for cell sorting and support with flow cytometric analysis; Dr. Susanne Kreutzer of the ETH Zürich Genome Engineering and Measurement Lab for performing next-generation sequencing; Dr. M. Muhar, M. Schlapansky, Dr. Z. Kontarakis, Dr. E. Karasu and all of the other members or the Corn laboratory for helpful discussions; Prof. Dr. Markus Stoffel and Svenja Godbersen for gifting Hepa 1-6 cell line; Dr. Kim Marquart and the Viral Vector Facility of UZH (Zurich) for providing AAV construct backbones. J.E.C. is supported by the NOMIS Foundation and the Lotte und Adolf Hotz-Sprenger Stiftung. funded by the NOMIS Foundation and by the Lotte and Adolf Hotz-Sprenger Stiftung. T.K. is supported by "Mission-driven Implementation of Science and Innovation Programmes" No. 02-002-P-0001, funded by the Economic Revitalization and Resilience Enhancement Plan "New Generation Lithuania". This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation funding programmes (grant agreement no. 855741-DDREAMM-ERC-2019-SyG; no. 101057659; no. 101070740; no. 101057438), Swiss State Secretariat for Education, Research and Innovation (SERI), SNSF Project Funding grants (310030_188858 and 320030-227979) and SNF Bridge-Discovery grant (40B2-0_226603). LV.V. is supported by the ETH Pioneer Fellowship.

Contributions

F.G. and J.E.C conceptualized the study. F.G. and I.S. performed most of the experiments. Y.-Y.L. purified the recombinant proteins. LV.V. and C.DY. performed DISCOVER-Seq and produced graphs in **Fig 3k,l**. A.T. performed the *in vivo* experiments. M.S.S. performed bioinformatic analyses and produced graphs in **Fig. 1d**. T.K. and G.D. performed *in vitro* biochemical studies and produced graphs in **Fig. S7**. L.V.B. and I.V. performed editing of primary T cells. F.G., A.G. and J.E.C. wrote the article. G.S. and V.S. reviewed and edited the article. J.E.C. supervised the study.

Declaration of interests

J.E.C. is a co-founder of Serac Biosciences and a scientific advisory board (SAB) member of Relation Therapeutics and Hornet Bio. The laboratory of J.E.C. has had funded collaborations with Allogene and Cimeio. J.E.C. and F.G. are inventors on a patent filed by the ETH related to this work. The remaining authors declare no competing interests.

References

1. Wu, Z., Yang, H. & Colosi, P. Effect of genome size on AAV vector packaging. *Mol. Ther. J. Am. Soc. Gene Ther.* **18**, 80–86 (2010).

2. Altae-Tran, H. *et al.* Diversity, evolution, and classification of the RNA-guided nucleases TnpB and Cas12. *Proc. Natl. Acad. Sci. U. S. A.* **120**, e2308224120 (2023).

3. Xu, X. *et al.* Engineered miniature CRISPR-Cas system for mammalian genome regulation and editing. *Mol. Cell* **81**, 4333-4345.e4 (2021).

4. Ma, S. *et al.* hpCasMINI: An engineered hypercompact CRISPR-Cas12f system with boosted gene editing activity. *Nat. Commun.* **16**, 5001 (2025).

5. Park, S.-J. *et al.* Robust genome editing activity and the applications of enhanced miniature CRISPR-Cas12f1. *Nat. Commun.* **16**, 677 (2025).

6. Kim, D. Y. *et al.* Efficient CRISPR editing with a hypercompact Cas12f1 and engineered guide RNAs delivered by adeno-associated virus. *Nat. Biotechnol.* **40**, 94–102 (2022).

7. Marquart, K. F. *et al.* Effective genome editing with an enhanced ISDra2 TnpB system and deep learning-predicted ωRNAs. *Nat. Methods* **21**, 2084–2093 (2024).

8. Hino, T. *et al.* An AsCas12f-based compact genome-editing tool derived by deep mutational scanning and structural analysis. *Cell* **186**, 4920-4935.e23 (2023).

9. Zhang, L. *et al.* AsCas12a ultra nuclease facilitates the rapid generation of

therapeutic cell medicines. *Nat. Commun.* **12**, 3908 (2021).

10.   Doman, J. L. *et al.* Phage-assisted evolution and protein engineering yield

compact, efficient prime editors. *Cell* **186**, 3983-4002.e26 (2023).

11.   Harrington, L. B. *et al.* Programmed DNA destruction by miniature CRISPR-

Cas14 enzymes. *Science* **362**, 839–842 (2018).

12.   Karvelis, T. *et al.* PAM recognition by miniature CRISPR-Cas12f nucleases

triggers programmable double-stranded DNA target cleavage. *Nucleic Acids Res.*

48, 5016–5023 (2020).

13.   Karvelis, T. *et al.* Transposon-associated TnpB is a programmable RNA-

guided DNA endonuclease. *Nature* **599**, 692–696 (2021).

14.   Porto, E. M., Komor, A. C., Slaymaker, I. M. & Yeo, G. W. Base editing:

advances and therapeutic opportunities. *Nat. Rev. Drug Discov.* **19**, 839–859

(2020).

15.   Murray, J. B., Harrison, P. T. & Scholefield, J. Prime editing: therapeutic

advances and mechanistic insights. *Gene Ther.* **32**, 83–92 (2025).

16.   Hu, Y. *et al.* Engineering miniature CRISPR-Cas Un1Cas12f1 for efficient

base editing. *Mol. Ther. Nucleic Acids* **35**, 102201 (2024).

17.   Jiang, K. *et al.* Rapid in silico directed evolution by a protein language model

with EVOLVEpro. *Science* **387**, eadr6006 (2025).

18.   Nakagawa, R. *et al.* Cryo-EM structure of the transposon-associated TnpB

enzyme. *Nature* **616**, 390–397 (2023).

19.   Marquart, K. F. *et al.* Effective genome editing with an enhanced ISDra2 TnpB

system and deep learning-predicted ωRNAs. *Nat. Methods* **21**, 2084–2093 (2024).

20.   Doench, J. G. *et al.* Rational design of highly active sgRNAs for CRISPR-

Cas9-mediated gene inactivation. *Nat. Biotechnol.* **32**, 1262–1267 (2014).

21.   Wienert, B. *et al.* Unbiased detection of CRISPR off-targets in vivo using

DISCOVER-Seq. *Science* **364**, 286–289 (2019).

22.   Ding, Q. *et al.* Permanent alteration of PCSK9 with in vivo CRISPR-Cas9

genome editing. *Circ. Res.* **115**, 488–492 (2014).

23.   Rothgangl, T. *et al.* In vivo adenine base editing of PCSK9 in macaques

reduces LDL cholesterol levels. *Nat. Biotechnol.* **39**, 949–957 (2021).

24.   Abifadel, M. & Boileau, C. Genetic and molecular architecture of familial

hypercholesterolemia. *J. Intern. Med.* **293**, 144–165 (2023).

25.   Richter, M. F. *et al.* Phage-assisted evolution of an adenine base editor with

improved Cas domain compatibility and activity. *Nat. Biotechnol.* **38**, 883–891

(2020).

26.   Spencer, J. M. & Zhang, X. Deep mutational scanning of S. pyogenes Cas9

reveals important functional domains. *Sci. Rep.* **7**, 16836 (2017).

27.   Hu, J. H. *et al.* Evolved Cas9 variants with broad PAM compatibility and high

DNA specificity. *Nature* **556**, 57–63 (2018).

28.   Zhang, L. *et al.* Boosting genome editing efficiency in human cells and plants

with novel LbCas12a variants. *Genome Biol.* **24**, 102 (2023).

29.   Ruta, G. V. *et al.* Eukaryotic-driven directed evolution of Cas9 nucleases.

Genome Biol. **25**, 79 (2024).

30.   Sasnauskas, G. *et al.* TnpB structure reveals minimal functional core of

Cas12 nuclease family. *Nature* **616**, 384–389 (2023).

31.   Thornton, B. W. *et al.* Engineered TnpB genome editors for plants and human

cells identified by ribonucleoprotein mutational scanning. *Nat. Biotechnol.*

https://doi.org/10.1038/s41587-026-03059-7 (2026) doi:10.1038/s41587-026-

03059-7.

32.   Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing

sequence analysis. *Nat. Biotechnol.* **37**, 224–226 (2019).

33.   Andrews, Simon. FastQC: A Quality Control Tool for High Throughput

Sequence Data. (2010).

34.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize

analysis results for multiple tools and samples in a single report. *Bioinforma. Oxf.*

Engl. **32**, 3047–3048 (2016).

35.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2.

Nat. Methods **9**, 357–359 (2012).

36.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**,

giab008 (2021).

37.   Bushnell, B. BBMap: A Fast, Accurate, Splice-Aware Aligner. (2014).

38.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**,

giab008 (2021).

39.   McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol.* **17**,

122 (2016).

40.   R Core Team. R: A Language and Environment for Statistical Computing.

(2023).

41.   Gu, Z. Complex heatmap visualization. *iMeta* **1**, e43 (2022).

42.   Wienert, B., Wyman, S. K., Yeh, C. D., Conklin, B. R. & Corn, J. E. CRISPR

off-target detection with DISCOVER-seq. *Nat. Protoc.* **15**, 1775–1799 (2020).

43.   Martin, M. Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.journal* **17**, 10 (2011).

44.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2.

Nat. Methods **9**, 357–359 (2012).

45.   Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for

researchers, educators, and developers. *Protein Sci. Publ. Protein Soc.* **30**, 70–82

(2021).

Figure legends

Figure 1. Directed evolution of CasMINI and TnpB for enhanced HDR activity in mammalian cells.

(**a**) Schematic of the homology-directed repair (HDR) fluorescent reporter. A protospacer with its protospacer-associated motif (Un1Cas12f1 PAM: TTTA) or transposon-associated motif (ISDra2 TnpB TAM: TTGAT) was cloned upstream of an *EGFP* gene lacking a start codon. Lentiviral delivery into HEK293T cells generated a non-fluorescent polyclonal cell line. *EGFP* expression can be restored upon successful HDR following delivery of a nuclease (CasMINI or TnpB), a guide RNA (sgRNA for CasMINI and ωRNA for TnpB), and a donor single-stranded oligodeoxynucleotide (ssODN) containing the start codon. An arrow indicates approximate cut position. *BSD*-gene for blasticidin resistance, LTR-long terminal repeat; LV-lentiviral.

(**b**) Schematic of the directed evolution selection workflow. CasMINI or TnpB variant libraries were screened in reporter cells by iterative cycles of HDR, fluorescence-activated cell sorting (FACS) enrichment, and library regeneration to select for improved activity.

(**c**) HDR efficiency measured as bulk EGFP signal across four rounds of selection of indicated proteins, normalized to the respective wild-type nuclease. Bars represent the value of a single experimental replicate.

(**d**) Relative frequency of enriched mutations of the original sequences (Ctrl) and after rounds 1, 2, and 4 ("R1", "R2", "R4", respectively) of selection for indicated nucleases. Mutations with >0.3% frequency and >3.75-fold enrichment over the pre-selection libraries are shown. The clusters with enriched mutations are marked. FC, fold-change.

(**e**) Enriched mutations mapped onto the structures of nuclease Un1Ca12f1 (PDB: 7l49),

highlighted in teal and nuclease TnpB (PDB: 8H1J) highlighted in orange. Close-up views for

mutation cluster 1 are presented for each nuclease. Protein structures were visualized with

ChimeraX[45].

Figure 2. Generation of CasMINI and TnpB variants with enhanced activity.

(**a**) Schematic of the strategy for generating combinatorial variants (CVs). Screen-identified
variants were ranked by homology-directed repair (HDR) activity and used to guide
combinatorial assembly. An additional HDR reporter target was used to exclude target-
specific effects. Final combinatorial variants were validated at endogenous genomic sites.

(**b**) HDR activity of post-round 4 variants measured as percentage of EGFP-positive cells
from BFP/mCherry double positive cells (BFP marks the guide-expressing construct;
mCherry marks the nuclease-expressing construct), normalized to CasMINI ($n = 2$ biological
replicates). Variants exceeding 1.2-fold, highlighted in teal, were used to identify active
mutations.

(**c**) A table showing mutations present in CasMINI combinatorial variants 16-20 (CVs 16-
20).

(**d**) HDR activity of CVs at two targets measured as percentage of EGFP-positive cells from
BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry
marks the nuclease-expressing construct), normalized to CasMINI ($n = 2$ biological
replicates). The CV selected for further testing (CV20) at endogenous sites is highlighted.

(**e**) Mutations present in Cas12f1Super (CV20) are mapped onto the domain organization of
Un1Cas12f1 (PDB: 7l49).

(**f**) HDR activity of post-round 4 variants measured as percentage of EGFP-positive cells
from BFP/mCherry double positive cells (BFP marks the guide-expressing construct;
mCherry marks the nuclease-expressing construct), normalized to TnpB ($n = 2$ biological
replicates). Variants exceeding 1.4-fold, highlighted in orange, were used to identify active
mutations.

(**g**) A table showing mutations present in V322 and combinatorial variants 1-5 (CVs 1-5) of
TnpB.

(**h**) HDR activity of CVs at two targets measured as percentage of EGFP-positive cells from
BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry

marks the nuclease-expressing construct), normalized to TnpB ($n = 2$ biological replicates).
The CV selected (CV2) for further testing at endogenous sites is highlighted.

(**i**) Mutations present in TnpBSuper (CV2) are mapped onto the domain organization of TnpB
(PDB: 8H1J).

Each dot represents an individual biological replicate, and bars represent the mean.

Figure 3. Improved variants Cas12f1Super and TnpBSuper edit endogenous genomic
loci.

(**a**) Schematic of the workflow of endogenous site editing in HEK293T and NIH-3T3 cells.
Nuclease-expressing plasmid (containing mCherry marker) was co-delivered with guide-
expressing plasmid (containing BFP marker). ssODN was used only for HDR-related
experiments in HEK293T cells. After enrichment for cells that had taken up both plasmids,
ampliconNGS was performed.

(**b,c**) Absolute HDR (**b**) or NHEJ efficiency of Cas12f1Super (**c**) in the presence of ssODN
measured with ampliconNGS at indicated endogenous loci in HEK293T cells compared to
CasMINI ($n = 3$ biological replicates).

(**d**) Absolute NHEJ efficiency in the absence of ssODN of Cas12f1Super measured with
ampliconNGS at indicated endogenous loci in HEK293T cells compared to CasMINI ($n = 3$
biological replicates).

(**e,f**) Absolute HDR (**e**) or NHEJ efficiency of TnpBSuper (**f**) in the presence of ssODN at
indicated endogenous loci in HEK293T cells compared to TnpB measured with
ampliconNGS ($n = 3$ biological replicates).

(**g**) Absolute NHEJ efficiency in the absence of ssODN of TnpBSuper measured with
ampliconNGS at indicated endogenous loci in HEK293T cells compared to TnpB ($n = 3$
biological replicates).

(**h,i**) Absolute HDR (**h**) or NHEJ efficiency (**i**) of TnpBSuper at indicated endogenous loci in
primary T cells compared to TnpB measured with ampliconNGS ($n = 3$ biological replicates).
TnpBSuper or TnpB were electroporated as mRNA, along with synthetic ωRNA and ssODN.

(**j**) Comparison of CasMINI/Cas12f1Super and TnpB/TnpBSuper with SpCas9 measured

with ampliconNGS at indicated endogenous loci in HEK293T cells ($n = 3$ biological

replicates).

(**k,l**) Off-target sites identified by DISCOVER-Seq. Sequences are presented along with

Discover-seq scores for on-target and off-target editing with CasMINI/Cas12f1Super (**k**) or

TnpB/TnpBSuper (**l**).

Each dot represents an individual biological replicate, and bars represent the mean ± standard

deviation. All *P* values were calculated using an unpaired, two-sided *t*-test. ns, not significant

($P \geq 0.05$), *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 4. *Efficient in vivo editing using AAV delivery of Cas12f1Super and TnpBSuper*

(**a,b**) Absolute NHEJ efficiency in NIH-3T3 cells at targets within the *Pcsk9* gene upon

editing with CasMINI/Cas12f1Super (**a**) or TnpB/TnpBSuper (**b**) measured with

ampliconNGS ($n = 3$ biological replicates).

(**c,d**) Absolute NHEJ efficiency in Hepa 1-6 cells at selected targets within the *Pcsk9* gene

upon editing with CasMINI/Cas12f1Super (**c**) or TnpB/TnpBSuper measured with

ampliconNGS (**d**) ($n = 3$ biological replicates). Each nuclease-guide RNA combination was

delivered on a "all-in-one" plasmid.

(**e**) Schematic representation of ssAAV9 vectors used for CasMINI/Cas12f1Super and

scAAV9 vectors used for TnpB/TnpBSuper in *in vivo* experiments. Both designs are based on

the TnpBmax architecture[19]. SRAD-SV40_NLS-GS-BP_NLS, linker from "TnpBmax"

construct design; WPRE, woodchuck hepatitis virus post-transcriptional regulatory element;

pA, polyadenylation signal sequence; s.pA, synthetic polyadenylation signal sequence.

(**f**) Schematic overview of the *in vivo* experimental setup. AAV9 capsids were administered

to mice via tail vein injection. PCSK9 protein levels were measured in blood serum at weeks

3 and 6. At week 6, murine livers were harvested and analyzed by amplicon ampliconNGS.

(**g,h**) Editing efficiencies in whole liver at the 6-week timepoint following treatment with
CasMINI/Cas12f1Super (**g**) or TnpB/TnpBSuper (**h**), measured by amplicon NGS ($n = 4$
animals).

(**i,j**) PCSK9 protein levels following treatment with CasMINI/Cas12f1Super (**i**) or
TnpB/TnpBSuper (**j**), measured in blood serum by ELISA normalized to untreated (control)
mice ($n = 3$.

Each dot represents an individual biological replicate or animal, and bars represent the mean
± standard deviation. All P values were calculated using an unpaired, two-sided t-test. ns, not
significant ($P \geq 0.05$), *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 5. Improved variant dCas12f1Super outperforms dCasMINI as an adenine base
editor.

(**a**) Schematic showing adenine base editing of Target CM1 and endogenous sites in
HEK293T using catalytically-dead CasMINI or Cas12f1Super (dCasMINI or dCas12f1Super,
respectively). Plasmid expressing catalytically inactive enzyme fused with TadA* is co-
delivered with guide-expressing plasmid. AmpliconNGS was performed 3 days post-
transfection.

(**b**) Overall adenine base editing activity of dCasMINI and dCas12f1Super at Target CM1
and indicated endogenous sites ($n = 3$ biological replicates).

(**c–f**) Per-site adenine editing efficiencies of adenine base editors dCasMINI and
dCas12f1Super at indicated endogenous genomic sites in HEK293T cells ($n = 3$ biological
replicates). Editing outcomes were quantified as A•T-to-G•C conversion rates.

(**g**) Product distributions at *TTR* endogenous site in HEK293T cells edited using a
combination of adenine base editor dCas12f1Super and the corresponding sgRNA.

(**h**) Allele plot at *TTR* endogenous site in HEK293T cells edited using a combination of
adenine base editor dCas12f1Super and the corresponding sgRNA

Each dot represents an individual biological replicate, and bars represent the mean ± standard

deviation. All *P* values were calculated using an unpaired, two-sided *t*-test. ns, not significant

(P ≥ 0.05), *P < 0.05, **P < 0.01, ***P < 0.001.

Supplementary figure legends

Figure S1. HDR reporter validation.

(**a, b**) Testing of CasMINI and TnpB genome editors at two targets (Target CM1 for

CasMINI and Target TB1 for TnpB) within HDR reporter's framework in HEK293T cells (*n*

= 3 biological replicates). EGFP percentages assessed by flow cytometry (**a**) and HDR and

NHEJ allele percentages obtained through ampliconNGS (**b**) were used to quantify editing

outcomes for indicated nucleases. NT, non-targeting.

Each dot represents an individual biological replicate, and bars represent the mean ± standard

deviation.

Figure S2. Localization of mutations enriched after selection for HDR with CasMINI

and TnpB.

(**a**) Pie chart showing proportion of unmutated (WT) and mutated (variants) sequences in

libraries used for the first round of selection for indicated nucleases. For purpose of

estimating this proportion 28 (for CasMINI) or 25 (for TnpB) randomly picked colonies from

respective libraries were picked and analyzed by Sanger sequencing.

(**b,c**) Mutations from **Fig. 1d** mapped on the domain organization of CasMINI (PDB: 7l49)

(**b**) and TnpB (PDB: 8H1J) (**c**). The mutations from clusters of enriched mutations after

round 4 of selection are labelled for each nuclease.

Figure S3. Selection of CasMINI mutations for generating combinatorial variants

(CVs).

(**a**) HDR efficiencies of 88 CasMINI variants were tested in a lentiviral (LV) format and 158

CasMINI variants tested with plasmid delivery at Target CM1 in HEK293T cells. HDR

values were measured as percentage of EGFP-positive cells from BFP/mCherry double

positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-

expressing construct) and normalized to CasMINI. Highlighted variants (teal) were selected

for further validation using plasmid delivery in duplicate.

(**b**) HDR efficiencies of single mutants (SMs) derived from selected high-performing

CasMINI variants or enriched in NGS after round four of selection, tested at Target CM1 in

HEK293T cells. HDR values were measured as percentage of EGFP-positive cells from

BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry

marks the nuclease-expressing construct) and normalized to CasMINI ($n = 2$ biological

replicates). Highlighted SMs (teal) were selected for further testing at the orthogonal HDR

reporter target (Target CM2). A378V was not selected due to the presence of A378T with

similar or slightly higher editing efficiency, which was chosen for further analysis.

Each dot represents an individual biological replicate, and bars represent the mean.

(**c**) HDR efficiencies of selected CasMINI SMs tested in parallel at two distinct targets

(Target CM1 and Target CM2) in HEK293T cells, measured as percentage of EGFP-positive

cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct;

mCherry marks the nuclease-expressing construct), with values normalized to CasMINI ($n = $

2 biological replicates). The mutations selected for making combinatorial variants are

highlighted.

Each dot represents an individual biological replicate, and bars represent the mean.

(**d**) Western blot analysis of CasMINI and CasMINI_CV20 protein expression levels.

Expression of each nuclease was performed in triplicate together with sgRNA for Target

CM1 in HEK293T cells. GAPDH was used as a loading control.

(**e**) HDR efficiencies of eight newly generated CasMINI variants, each containing one

mutation derived from EVOLVEpro, cloned onto the CasMINI_CV20 background, tested in

HEK293T cells ($n = 2$ biological replicates). HDR values were measured as percentage of

EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-

expressing construct; mCherry marks the nuclease-expressing construct) and normalized to

the previously best-performing variant, CasMINI_CV20.

Each dot represents an individual biological replicate, and bars represent the mean.

(**f**) Comparison of editing efficiencies of Un1Cas12f1, CasMINI, Un1Cas12f1_CV20,

CasMINI_CV20 at Target CM1 and Target CM2 in HEK293T cells. HDR values were

measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells

(BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing

construct) and normalized to Un1Cas12f1 ($n = 3$ biological replicates).

Each dot represents an individual biological replicate, and bars represent the mean ± standard

deviation. All P values were calculated using an unpaired, two-sided t-test. ns, not significant

($P \geq 0.05$), *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure S4. Selection of TnpB mutations for generating combinatorial variants (CVs).

(**a**) HDR efficiencies of 311 TnpB variants with plasmid delivery at Target TB1 in HEK293T

cells. HDR values were measured as a percentage of EGFP-positive cells from BFP/mCherry

double positive cells (BFP marks the guide-expressing construct; mCherry marks the

nuclease-expressing construct) and normalized to TnpB. Highlighted variants (orange) were

selected for further validation using plasmid delivery in duplicate. Top-performing is

indicated (V322).

(**b**) HDR efficiencies of all TnpB single mutants (SMs) derived from selected high-

performing TnpB variants or enriched in NGS after round four of selection, tested at Target

TB1 in HEK293T cells ($n = 2$ biological replicates). HDR values were measured as

percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the

guide-expressing construct; mCherry marks the nuclease-expressing construct) and

normalized to TnpB. Highlighted SMs were selected for further testing at the orthogonal

HDR reporter target (Target TB2).

Each dot represents an individual biological replicate, and bars represent the mean.

(**c**) HDR efficiencies of selected TnpB SMs tested in parallel at two distinct targets (Target

TB1 and Target TB2) in HEK293T cells measured as percentage of EGFP-positive cells from

BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry

marks the nuclease-expressing construct), with values normalized to TnpB ($n = 2$ biological

replicates). The mutations selected for making combinatorial variants are highlighted.

Each dot represents an individual biological replicate, and bars represent the mean.

(**d**) HDR efficiencies of seven newly generated TnpB variants, each containing one mutation

derived from EVOLVEpro, cloned onto the TnpB_CV2 background, tested in HEK293T

cells (*n* = 2 biological replicates). HDR values were measured as percentage of EGFP-

positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing

construct; mCherry marks the nuclease-expressing construct) and normalized to the

previously best-performing variant, TnpB_CV2.

Each dot represents an individual biological replicate, and bars represent the mean.

Figure S5. Mammalian codon-optimization and usage of shorter scaffold for TnpB

enhance editing at HDR reporter targets in HEK293T cells.

(**a,b**) HDR efficiencies with mammalian codon-optimized (+CO) and not codon-optimized (-

CO) versions of TnpB and TnpB_CV2 in HEK293T cells at HDR reporter Target TB1 (**a**)

and Target TB2 (**b**) (*n* = 3 biological replicates). HDR was measured as percentage of EGFP-

positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing

construct; mCherry marks the nuclease-expressing construct).

Each dot represents an individual biological replicate, and bars represent the mean ± standard

deviation.

(**c**) Western blot analysis of mammalian codon-optimized TnpB and TnpB_CV2 protein

expression levels (*n* = 3 biological replicates). Expression of each nuclease was performed

together with ωRNA for Target TB1 in HEK293T cells. GAPDH was used as a loading

control.

(**d,e**) HDR efficiencies of mammalian codon-optimized TnpB and TnpB_CV2 combined with

either Trim2 ωRNA scaffold or original ωRNA in HEK293T cells at HDR reporter Target

TB1 (**d**) and Target TB2 (**e**). HDR was measured as percentage of EGFP-positive cells from

BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry

marks the nuclease-expressing construct) (*n* = 2 biological replicates).

Each dot represents an individual biological replicate, and bars represent the mean.

(**f**) HDR efficiencies of mammalian codon-optimized TnpB, TnpBmax, TnpB_CV2 and

TnpBmax_CV2 combined with Trim2 ωRNA scaffold in HEK293T cells at HDR reporter

targets TB1 and TB2. HDR was measured as percentage of EGFP-positive cells from
mCherry+ cells (all-in-one construct was used for this experiment) (*n* = 3 biological
replicates).

(**g**) HDR and NHEJ efficiencies of mammalian codon-optimized TnpB, TnpBmax,
TnpB_CV2 and TnpBmax_CV2 combined with Trim2 ωRNA scaffold in HEK293T cells at
indicated endogenous sites. HDR and NHEJ were measured with ampliconNGS (*n* = 3
biological replicates).

Each dot represents an individual biological replicate, and bars represent the mean ± standard
deviation. All *P* values were calculated using an unpaired, two-sided *t*-test. ns, not significant
(P ≥ 0.05), *$P<$0.05, **$P<$0.01, ***$P<$0.001.

Figure S6. Improved variant Cas12f1Super edits endogenous genomic loci.

(**a,b**) Fold change of HDR efficiency (**a**) or NHEJ efficiency (**b**) of Cas12f1Super relative to
CasMINI in presence of ssODN measured with ampliconNGS at indicated endogenous loci in
HEK293T cells (*n* = 3 biological replicates).

(**c**) Fold change of NHEJ efficiency of Cas12f1Super relative to CasMINI in absence of
ssODN measured with ampliconNGS at indicated endogenous loci in HEK293T cells (*n* = 3
biological replicates).

Bars represent mean ± standard deviation. Fold change was calculated from group means,
with propagated standard deviation shown as error bars.

Figure S7. *In vitro* cleavage assays with CasMINI and Cas12f1Super.

(**a-f**) Percentages of fully cleaved substrates at two different targets at designated time points
by CasMINI or Cas12f1Super (*n* = 3 biological replicates).

Each dot represents the mean of three biological replicates ± standard deviation.

(**g,h**) Run-off Sanger sequencing of the plasmid products cleaved by CasMINI (**g**) or
Cas12f1Super (**h**). Cleavage positions at both the non-targeted strand (NTS) and the targeted
strand (TS) are marked with black triangles.

Figure S8. Improved variants Cas12f1Super and TnpBSuper edit endogenous genomic loci.

(**a,b**) Fold change of HDR efficiency (**a**) or NHEJ efficiency (**b**) of TnpBSuper relative to TnpB in presence of ssODN measured with ampliconNGS at indicated endogenous loci in HEK293T cells ($n = 3$ biological replicates).

(**c**) Fold change of NHEJ efficiency of TnpBSuper relative to TnpB in absence of ssODN measured with ampliconNGS at indicated endogenous loci in HEK293T cells ($n = 3$ biological replicates).

Bars represent mean ± standard deviation. Fold change was calculated from group means, with propagated standard deviation shown as error bars.

(**d**) Absolute NHEJ efficiency of Cas12f1Super or TnpBSuper measured with ampliconNGS at indicated guides targeting *TRAC* locus in HEK293T cells compared to CasMINI or TnpB ($n = 2$ biological replicates).

Each dot represents an individual biological replicate, and bars represent the mean.

(**e,f**) Fold change of HDR (**c**) or NHEJ efficiency (**d**) of TnpBSuper relative to TnpB in the presence of ssODN measured with ampliconNGS at indicated endogenous loci in primary T cells ($n = 3$ biological replicates). TnpBSuper or TnpB were electroporated as mRNA, along with synthetic ωRNA and ssODN.

(**g**) Absolute NHEJ efficiency of SpCas9 in the absence of ssODN measured with ampliconNGS at indicated at indicated loci in HEK293T cells ($n = 2$ biological replicates).

Each dot represents an individual biological replicate, and bars represent the mean.

Figure S9. CasMINI/Cas12f1Super and TnpB/TnpBSuper NHEJ efficiency at on-target and off-target sites.

(**a**) NHEJ efficiency of CasMINI and Cas12f1Super measured with ampliconNGS at the indicated on-target and off-target sites in HEK293T cells.

(**b**) NHEJ efficiency of TnpB and TnpBSuper measured with ampliconNGS at the indicated

on-target sites in HEK293T cells.

Bars represent the value of a single experimental replicate.

Figure S10. Improved variants Cas12f1Super and TnpBSuper edit endogenous genomic

loci.

(**a,b**) Fold change of NHEJ efficiency of Cas12f1Super relative to CasMINI (**a**) or of

TnpBSuper relative to TnpB (**b**) measured with ampliconNGS at indicated targets within the

Pcsk9 gene in Hepa 1-6 cells ($n = 3$ biological replicates). Each nuclease-guide RNA

combination was delivered on a "all-in-one" plasmid.

Bars represent mean ± standard deviation. Fold change was calculated from group means,

with propagated standard deviation shown as error bars.

Figure S11. Improved variant dCas12f1Super outperforms dCasMINI as an adenine

base editor.

(**a**) Fold change of overall adenine base editing activity of dCas12f1Super relative to

dCasMINI measured with ampliconNGS at Target CM1 and indicated endogenous sites in

HEK293T cells ($n = 3$ biological replicates).

Bars represent mean ± standard deviation. Fold change was calculated from group means,

with propagated standard deviation shown as error bars.

(**b**) Frequency or reads without and with indels upon editing *TTR* site with adenine base

editor based on dCasMINI or dCas12f1Super in HEK293T cells ($n = 3$ biological replicates).

Each dot represents an individual biological replicate, and bars represent the mean.

Figure S12. Gating strategies for flow cytometry and FACS experiments related to

testing editing at HDR reporter targets and endogenous sites in HEK293T and Hepa 1-6

cell lines.

(**a**) Representative plots showing the gating strategy for measuring HDR at HDR reporter

targets in HEK293T cell line. Cells were gated for live cells, singlets, BFP positivity,

mCherry positivity (BFP marks the guide-expressing construct; mCherry marks the nuclease-
expressing construct) and finally, for EGFP positivity. In case of "edited" sample one of the
CasMINI variants was co-delivered with sgRNA for Target CM1 together with ssODN.
"BFP" only and "not transfected" samples serve as controls for gating. In case of "BFP only"
sample only sgRNA targeting Target CM1 was delivered.

(b) Representative plots showing the gating strategy for FACS-sorting HEK293T or Hepa 1-6
cells for measuring editing. Cells were gated for live cells, singlets and mCherry positivity. In
case of "edited" sample CasMINI and sgRNA were delivered on the same "all-in-one"
plasmid. "Not transfected" sample serves as control for gating.

Figure S13. Gating strategy for FACS experiments related to testing editing at
endogenous sites in HEK293T and NIH-3T3 cell line and to big scale selection of
CasMINI and TnpB for HDR in HEK293T cell line.

(a) Representative plots showing the gating strategy for FACS-sorting HEK293T and NIH-
3T3 cell lines for measuring NHEJ at endogenous sites. Cells were gated for live cells,
singlets and mCherry/BFP positivity (BFP marks the guide-expressing construct; mCherry
marks the nuclease-expressing construct). In case of "edited" sample CasMINI was co-
delivered with sgRNA and ssODN in HEK293T cells. "BFP only", "mCherry only" and "not
transfected" samples serve as controls for gating.

(b) Representative plots showing the gating strategy for FACS-sorting during round 1 of
HDR selection of CasMINI in HEK293T-reporter cell line. Cells were gated for live cells,
singlets, SYTOX^{TM}-Red-negativity, and finally for mCherry/EGFP positivity. In case of
"edited" sample sgRNA for Target CM1 was delivered together with ssODN. "Not
transfected" and "HEK293T" samples serve as controls for gating. In case of "not
transfected" control sgRNA and ssODN were not delivered.

Supplementary Tables

1265

Supplementary Table 1. Table listing all CasMINI and TnpB evolved variants and
combinatorial variants including their mutations.

Supplementary Table 2. Table listing the sequences of CasMINI and TnpB reporter and
endogenous targets, and all primers and oligonucleotides used in this study.

Supplementary Table 3. Table listing all CasMINI and TnpB ssODNs used for reporter and
endogenous targets.

Supplementary Table 4. Table listing all plasmids used in this study.

Supplementary Table 5. Table listing sgRNA sequences used *in vitro* cleavage assays for
CasMINI and Cas12f1Super.

Supplementary Table 6. Table listing all CasMINI and TnpB guide RNA scaffolds used in
this study.

Figure 1. Directed evolution of CasMINI and TnpB for enhanced HDR activity in mammalian cells.
(**a**) Schematic of the homology-directed repair (HDR) fluorescent reporter. A protospacer with its protospacer-associated motif (Un1Cas12f1 PAM: TTTA) or transposon-associated motif (ISDra2 TnpB TAM: TTGAT) was cloned upstream of an EGFP gene lacking a start codon. Lentiviral delivery into HEK293T cells generated a non-fluorescent polyclonal cell line. EGFP expression can be restored upon successful HDR following delivery of a nuclease (CasMINI or TnpB), a guide RNA (sgRNA for CasMINI and ωRNA for TnpB), and a donor single-stranded oligodeoxynucleotide (ssODN) containing the start codon. An arrow indicates approximate cut position. BSD-gene for blasticidin resistance, LTR-long terminal repeat; LV- lentiviral.
(**b**) Schematic of the directed evolution selection workflow. CasMINI or TnpB variant libraries were screened in reporter cells by iterative cycles of HDR, fluorescence-activated cell sorting (FACS) enrichment, and library regeneration to select for improved activity.
(**c**) HDR efficiency measured as bulk EGFP signal across four rounds of selection of indicated proteins, normalized to the respective wild-type nuclease. Bars represent the value of a single experimental replicate.
(**d**) Relative frequency of enriched mutations of the original sequences (Ctrl) and after rounds 1, 2, and 4 ("R1", "R2", "R4", respectively) of selection for indicated nucleases. Mutations with >0.3% frequency and >3.75-fold enrichment over the pre-selection libraries are shown. The clusters with enriched mutations are marked. FC, fold-change.
(**e**) Enriched mutations mapped onto the structures of nuclease Un1Ca12f1 (PDB: 7l49), highlighted in teal and nuclease TnpB (PDB: 8H1J) highlighted in orange. Close-up views for mutation cluster 1 are presented for each nuclease. Protein structures were visualized with ChimeraX45.

a

Evolved variants (V) → Activity assay — HDR ratio (V/WT), Variants → Combinatorial variants (CV) → Activity assay — HDR ratio (V/WT), Combinatorial variants (CV) → Validation at endogenous sites

b CasMINI — HDR improvement (V/CasMINI)

Selected / Discarded

V4, V16, V9, V37, V20, V75, V1, V163, V45, V25, V21, V108, V123

c CasMINI

position	52	55	66	129	206	227	504	506
CasMINI	K	K	R	K	E	K	N	K
CV16	E	E	E	E	K	E	D	E
CV17	E	K	E	E	K	E	D	E
CV18	E	K	R	E	K	E	D	E
CV19	E	K	E	E	K	E	N	E
CV20	E	K	R	E	K	E	N	E

d CasMINI — HDR improvement (CV/CasMINI)

Target CM1 / Target CM2

CV16, CV17, CV18, CV19, CV20

e Cas12f1Super (CV20)

WED | REC1 | WED | RuvC | REC2 | LID | RuvC | Nuc | RuvC

1 18 189 315 347 415 442 469 508 529
424

K52E K129E E206K K227E K506E

f TnpB — HDR improvement (V/TnpB)

Selected / Discarded

V322, V285, V8, V129, V158, V192, V32, V163, V315, V231, V235, V201, V173, V199, V86, V156, V152

g TnpB

position	73	155	163	172	192	209	240	299	384
TnpB	E	L	H	L	V	E	G	G	A
V322	K	V	H	L	V	E	R	A	T
CV1	K	V	H	L	L	E	R	A	T
CV2	K	V	Y	V	L	V	R	A	T
CV3	K	L	Y	V	L	V	R	A	T
CV4	K	V	Y	L	L	V	R	A	T
CV5	K	L	Y	L	L	V	R	A	T

h TnpB — HDR improvement (CV/TnpB)

Target TB1 / Target TB2

CV1, CV2, CV3, CV4, CV5

i TnpBSuper (CV2)

WED | REC | WED | RuvC | TNB | RuvC | CTD

13 116 177 185 327 360 376 408

E73K L155V H163Y V192L L172V E209V G240R G299A A384T

Figure 2. Generation of CasMINI and TnpB variants with enhanced activity.

(**a**) Schematic of the strategy for generating combinatorial variants (CVs). Screen-identified variants were ranked by homology-directed repair (HDR) activity and used to guide combinatorial assembly. An additional HDR reporter target was used to exclude target-specific effects. Final combinatorial variants were validated at endogenous genomic sites.

(**b**) HDR activity of post-round 4 variants measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct), normalized to CasMINI (n = 2 biological replicates). Variants exceeding 1.2-fold, highlighted in teal, were used to identify active mutations.

(**c**) A table showing mutations present in CasMINI combinatorial variants 16-20 (CVs 16-20).

(**d**) HDR activity of CVs at two targets measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct), normalized to CasMINI (n = 2 biological replicates). The CV selected for further testing (CV20) at endogenous sites is highlighted.

(**e**) Mutations present in Cas12f1Super (CV20) are mapped onto the domain organization of Un1Cas12f1 (PDB: 7I49).

(**f**) HDR activity of post-round 4 variants measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct), normalized to TnpB (n = 2 biological replicates). Variants exceeding 1.4-fold, highlighted in orange, were used to identify active mutations.

(g) A table showing mutations present in V322 and combinatorial variants 1-5 (CVs 1-5) of TnpB.

(h) HDR activity of CVs at two targets measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct), normalized to TnpB (n = 2 biological replicates). The CV selected (CV2) for further testing at endogenous sites is highlighted.

(**i**) Mutations present in TnpBSuper (CV2) are mapped onto the domain organization of TnpB (PDB: 8H1J).

Each dot represents an individual biological replicate, and bars represent the mean.

Fig. 3

Figure 3. Improved variants Cas12f1Super and TnpBSuper edit endogenous genomic loci.

(**a**) Schematic of the workflow of endogenous site editing in HEK293T and NIH-3T3 cells. Nuclease-expressing plasmid (containing mCherry marker) was co-delivered with guide-expressing plasmid (containing BFP marker). ssODN was used only for HDR-related experiments in HEK293T cells. After enrichment for cells that had taken up both plasmids, ampliconNGS was performed.

(**b,c**) Absolute HDR (b) or NHEJ efficiency of Cas12f1Super (c) in the presence of ssODN measured with ampliconNGS at indicated endogenous loci in HEK293T cells compared to CasMINI (n = 3 biological replicates).

(**d**) Absolute NHEJ efficiency in the absence of ssODN of Cas12f1Super measured with ampliconNGS at indicated endogenous loci in HEK293T cells compared to CasMINI (n = 3 biological replicates).

(**e,f**) Absolute HDR (e) or NHEJ efficiency of TnpBSuper (f) in the presence of ssODN at indicated endogenous loci in HEK293T cells compared to TnpB measured with ampliconNGS (n = 3 biological replicates).

(**g**) Absolute NHEJ efficiency in the absence of ssODN of TnpBSuper measured with ampliconNGS at indicated endogenous loci in HEK293T cells compared to TnpB (n = 3 biological replicates).

(**h,i**) Absolute HDR (h) or NHEJ efficiency (i) of TnpBSuper at indicated endogenous loci in primary T cells compared to TnpB measured with ampliconNGS (n = 3 biological replicates). TnpBSuper or TnpB were electroporated as mRNA, along with synthetic   RNA and ssODN.

(**j**) Comparison of CasMINI/Cas12f1Super and TnpB/TnpBSuper with SpCas9 measured with ampliconNGS at indicated endogenous loci in HEK293T cells (n = 3 biological replicates).

(**k,l**) Off-target sites identified by DISCOVER-Seq. Sequences are presented along with Discover-seq scores for on-target and off-target editing with CasMINI/Cas12f1Super (k) or TnpB/TnpBSuper (l).

Each dot represents an individual biological replicate, and bars represent the mean ± standard deviation. All P values were calculated using an unpaired, two-sided t-test. ns, not significant (P ≥ 0.05), *P < 0.05, **P < 0.01, ***P < 0.001.

Figure 4. Efficient in vivo editing using AAV delivery of Cas12f1Super and TnpBSuper

(**a,b**) Absolute NHEJ efficiency in NIH-3T3 cells at targets within the Pcsk9 gene upon editing with CasMINI/Cas12f1Super (a) or TnpB/TnpBSuper (b) measured with ampliconNGS (n = 3 biological replicates).

(**c,d**) Absolute NHEJ efficiency in Hepa 1-6 cells at selected targets within the Pcsk9 gene upon editing with CasMINI/Cas12f1Super (c) or TnpB/TnpBSuper measured with ampliconNGS (d) (n = 3 biological replicates). Each nuclease-guide RNA combination was delivered on a "all-in-one" plasmid.

(**e**) Schematic representation of ssAAV9 vectors used for CasMINI/Cas12f1Super and scAAV9 vectors used for TnpB/TnpBSuper in in vivo experiments. Both designs are based on the TnpBmax architecture19. SRAD-SV40_NLS-GS-BP_NLS, linker from "TnpBmax" construct design; WPRE, woodchuck hepatitis virus post-transcriptional regulatory element; pA, polyadenylation signal sequence; s.pA, synthetic polyadenylation signal sequence.

(**f**) Schematic overview of the in vivo experimental setup. AAV9 capsids were administered to mice via tail vein injection. PCSK9 protein levels were measured in blood serum at weeks 3 and 6. At week 6, murine livers were harvested and analyzed by amplicon ampliconNGS.

(**g,h**) Editing efficiencies in whole liver at the 6-week timepoint following treatment with CasMINI/Cas12f1Super (g) or TnpB/TnpBSuper (h), measured by amplicon NGS (n = 4 animals).

(**i,j**) PCSK9 protein levels following treatment with CasMINI/Cas12f1Super (i) or TnpB/TnpBSuper (j), measured in blood serum by ELISA normalized to untreated (control) mice (n = 3.

Each dot represents an individual biological replicate or animal, and bars represent the mean ± standard deviation. All P values were calculated using an unpaired, two-sided t-test. ns, not significant (P ≥ 0.05), *P < 0.05, **P < 0.01, ***P < 0.001.

Figure 5. Improved variant dCas12f1Super outperforms dCasMINI as an adenine base editor.

(**a**) Schematic showing adenine base editing of Target CM1 and endogenous sites in HEK293T using catalytically-dead CasMINI or Cas12f1Super (dCasMINI or dCas12f1Super, respectively). Plasmid expressing catalytically inactive enzyme fused with TadA* is co-delivered with guide-expressing plasmid. AmpliconNGS was performed 3 days post-transfection.

(**b**) Overall adenine base editing activity of dCasMINI and dCas12f1Super at Target CM1 and indicated endogenous sites (n = 3 biological replicates).

(**c–f**) Per-site adenine editing efficiencies of adenine base editors dCasMINI and dCas12f1Super at indicated endogenous genomic sites in HEK293T cells (n = 3 biological replicates). Editing outcomes were quantified as A•T-to-G•C conversion rates.

(**g**) Product distributions at TTR endogenous site in HEK293T cells edited using a combination of adenine base editor dCas12f1Super and the corresponding sgRNA.

(**h**) Allele plot at TTR endogenous site in HEK293T cells edited using a combination of adenine base editor dCas12f1Super and the corresponding sgRNA Each dot represents an individual biological replicate, and bars represent the mean ± standard deviation. All P values were calculated using an unpaired, two-sided t-test. ns, not significant (P ≥ 0.05), *P < 0.05, **P < 0.01, ***P < 0.001.

Figure S1. HDR reporter validation.
(**a, b**) Testing of CasMINI and TnpB genome editors at two targets (Target CM1 for CasMINI and Target TB1 for TnpB) within HDR reporter's framework in HEK293T cells (n = 3 biological replicates). EGFP percentages assessed by flow cytometry (a) and HDR and NHEJ allele percentages obtained through ampliconNGS (b) were used to quantify editing outcomes for indicated nucleases. NT, non-targeting.
Each dot represents an individual biological replicate, and bars represent the mean ± standard deviation.

a

b

c

Figure S2. Localization of mutations enriched after selection for HDR with CasMINI and TnpB.
(**a**) Pie chart showing proportion of unmutated (WT) and mutated (variants) sequences in libraries used for the first round of selection for indicated nucleases. For purpose of estimating this proportion 28 (for CasMINI) or 25 (for TnpB) randomly picked colonies from respective libraries were picked and analyzed by Sanger sequencing.
(**b,c**) Mutations from Fig. 1d mapped on the domain organization of CasMINI (PDB: 7l49) (b) and TnpB (PDB: 8H1J) (c). The mutations from clusters of enriched mutations after round 4 of selection are labelled for each nuclease.

Figure S3. Selection of CasMINI mutations for generating combinatorial variants (CVs).
(**a**) HDR efficiencies of 88 CasMINI variants were tested in a lentiviral (LV) format and 158 CasMINI variants tested with plasmid delivery at Target CM1 in HEK293T cells. HDR values were measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-ex-pressing construct; mCherry marks the nuclease-expressing construct) and normalized to CasMINI. Highlighted variants (teal) were selected for further validation using plasmid delivery in duplicate.
(**b**) HDR efficiencies of single mutants (SMs) derived from selected high-performing CasMINI variants or enriched in NGS after round four of selection, tested at Target CM1 in HEK293T cells. HDR values were measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct) and normalized to CasMINI (n = 2 biological replicates). Highlighted SMs (teal) were selected for further testing at the orthogonal HDR reporter target (Target CM2). A378V was not selected due to the presence of A378T with similar or slightly higher editing efficiency, which was chosen for further analysis.
Each dot represents an individual biological replicate, and bars represent the mean.
(**c**) HDR efficiencies of selected CasMINI SMs tested in parallel at two distinct targets (Target CM1 and Target CM2) in HEK293T cells, measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-ex-pressing construct), with values normalized to CasMINI (n = 2 biological replicates). The mutations selected for making combinatorial variants are highlighted.
Each dot represents an individual biological replicate, and bars represent the mean.
(**d**) Western blot analysis of CasMINI and CasMINI_CV20 protein expression levels. Expression of each nuclease was performed in triplicate together with sgRNA for Target CM1 in HEK293T cells. GAPDH was used as a loading control.
(**e**) HDR efficiencies of eight newly generated CasMINI variants, each containing one mutation derived from EVOLVEpro, cloned onto the CasMINI_CV20 background, tested in HEK293T cells (n = 2 biological replicates). HDR values were measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct) and normalized to the previously best-performing variant, CasMINI_CV20.
Each dot represents an individual biological replicate, and bars represent the mean.
(**f**) Comparison of editing efficiencies of Un1Cas12f1, CasMINI, Un1Cas12f1_CV20, CasMINI_CV20 at Target CM1 and Target CM2 in HEK293T cells. HDR values were measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct) and normalized to Un1Cas12f1 (n = 3 biological replicates).
Each dot represents an individual biological replicate, and bars represent the mean ± standard deviation. All P values were calculated using an unpaired, two-sided t-test. ns, not significant (P ≥ 0.05), *P < 0.05, **P < 0.01, ***P < 0.001.

Figure S4. Selection of TnpB mutations for generating combinatorial variants (CVs).

(**a**) HDR efficiencies of 311 TnpB variants with plasmid delivery at Target TB1 in HEK293T cells. HDR values were measured as a percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct) and normalized to TnpB. Highlighted variants (orange) were selected for further validation using plasmid delivery in duplicate. Top-performing is indicated (V322).

(**b**) HDR efficiencies of all TnpB single mutants (SMs) derived from selected high-performing TnpB variants or enriched in NGS after round four of selection, tested at Target TB1 in HEK293T cells (n = 2 biological replicates). HDR values were measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct) and normalized to TnpB. Highlighted SMs were selected for further testing at the orthogonal HDR reporter target (Target TB2).
Each dot represents an individual biological replicate, and bars represent the mean.

(**c**) HDR efficiencies of selected TnpB SMs tested in parallel at two distinct targets (Target TB1 and Target TB2) in HEK293T cells measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct), with values normalized to TnpB (n = 2 biological replicates). The mutations selected for making combinatorial variants are highlighted.
Each dot represents an individual biological replicate, and bars represent the mean.

(**d**) HDR efficiencies of seven newly generated TnpB variants, each containing one mutation derived from EVOLVEpro, cloned onto the TnpB_CV2 background, tested in HEK293T cells (n = 2 biological replicates). HDR values were measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct) and normalized to the previously best-performing variant, TnpB_CV2.
Each dot represents an individual biological replicate, and bars represent the mean.

Figure S5. Mammalian codon-optimization and usage of shorter scaffold for TnpB enhance editing at HDR reporter targets in HEK293T cells.

(**a,b**) HDR efficiencies with mammalian codon-optimized (+CO) and not codon-optimized (-CO) versions of TnpB and TnpB_CV2 in HEK293T cells at HDR reporter Target TB1 (a) and Target TB2 (b) (n = 3 biological replicates). HDR was measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct). Each dot represents an individual biological replicate, and bars represent the mean ± standard deviation.

(**c**) Western blot analysis of mammalian codon-optimized TnpB and TnpB_CV2 protein expression levels (n = 3 biological replicates). Expression of each nuclease was performed together with   RNA for Target TB1 in HEK293T cells. GAPDH was used as a loading control.

(**d,e**) HDR efficiencies of mammalian codon-optimized TnpB and TnpB_CV2 combined with either Trim2   RNA scaffold or original   RNA in HEK293T cells at HDR reporter Target TB1 (d) and Target TB2 (e). HDR was measured as percentage of EGFP-positive cells from BFP/mCherry double positive cells (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct) (n = 2 biological replicates). Each dot represents an individual biological replicate, and bars represent the mean.

(**f**) HDR efficiencies of mammalian codon-optimized TnpB, TnpBmax, TnpB_CV2 and TnpBmax_CV2 combined with Trim2   RNA scaffold in HEK293T cells at HDR reporter targets TB1 and TB2. HDR was measured as percentage of EGFP-positive cells from mCherry+ cells (all-in-one construct was used for this experiment) (n = 3 biological replicates).

(**g**) HDR and NHEJ efficiencies of mammalian codon-optimized TnpB, TnpBmax, TnpB_CV2 and TnpBmax_CV2 combined with Trim2   RNA scaffold in HEK293T cells at indicated endogenous sites. HDR and NHEJ were measured with ampliconNGS (n = 3 biological replicates).

Each dot represents an individual biological replicate, and bars represent the mean ± standard deviation. All P values were calculated using an unpaired, two-sided t-test. ns, not significant (P ≥ 0.05), *P < 0.05, **P < 0.01, ***P < 0.001.

Figure S6. Improved variant Cas12f1Super edits endogenous genomic loci.
(**a,b**) Fold change of HDR efficiency (a) or NHEJ efficiency (b) of Cas12f1Super relative to CasMINI in presence of ssODN measured with ampliconNGS at indicated endogenous loci in HEK293T cells (n = 3 biological replicates).
(**c**) Fold change of NHEJ efficiency of Cas12f1Super relative to CasMINI in absence of ssODN measured with ampliconNGS at indicated endogenous loci in HEK293T cells (n = 3 biological replicates).
Bars represent mean ± standard deviation. Fold change was calculated from group means, with propagated standard deviation shown as error bars.

Figure S7. In vitro cleavage assays with CasMINI and Cas12f1Super.
(**a-f**) Percentages of fully cleaved substrates at two different targets at designated time points by CasMINI or Cas12f1Super (n = 3 biological replicates). Each dot represents the mean of three biological replicates ± standard deviation.
(**g,h**) Run-off Sanger sequencing of the plasmid products cleaved by CasMINI (g) or Cas12f1Super (h). Cleavage positions at both the non-targeted strand (NTS) and the targeted strand (TS) are marked with black triangles.

Figure S8. Improved variants Cas12f1Super and TnpBSuper edit endogenous genomic loci.
(**a,b**) Fold change of HDR efficiency (a) or NHEJ efficiency (b) of TnpBSuper relative to TnpB in presence of ssODN measured with ampliconNGS at indicated endogenous loci in HEK293T cells (n = 3 biological replicates).
(**c**) Fold change of NHEJ efficiency of TnpBSuper relative to TnpB in absence of ssODN measured with ampliconNGS at indicated endogenous loci in HEK293T cells (n = 3 biological replicates).
Bars represent mean ± standard deviation. Fold change was calculated from group means, with propagated standard deviation shown as error bars.
(**d**) Absolute NHEJ efficiency of Cas12f1Super or TnpBSuper measured with ampliconNGS at indicated guides targeting TRAC locus in HEK293T cells compared to CasMINI or TnpB (n = 2 biological replicates).
Each dot represents an individual biological replicate, and bars represent the mean.
(**e,f**) Fold change of HDR efficiency (c) or NHEJ efficiency (d) of TnpBSuper relative to TnpB in the presence of ssODN measured with ampliconNGS at indicated endogenous loci in primary T cells (n = 3 biological replicates). TnpBSuper or TnpB were electroporated as mRNA, along with synthetic   RNA and ssODN.
(**g**) Absolute NHEJ efficiency of SpCas9 in the absence of ssODN measured with ampliconNGS at indicated at indicated loci in HEK293T cells (n = 2 biological replicates).
Each dot represents an individual biological replicate, and bars represent the mean.

Figure S9. CasMINI/Cas12f1Super and TnpB/TnpBSuper NHEJ efficiency at on-target and off-target sites.
(**a**) NHEJ efficiency of CasMINI and Cas12f1Super measured with ampliconNGS at the indicated on-target and off-target sites in HEK293T cells.
(**b**) NHEJ efficiency of TnpB and TnpBSuper measured with ampliconNGS at the indicated on-target sites in HEK293T cells.
Bars represent the value of a single experimental replicate.

Figure S10. Improved variants Cas12f1Super and TnpBSuper edit endogenous genomic loci.
(**a,b**) Fold change of NHEJ efficiency of Cas12f1Super relative to CasMINI (a) or of TnpBSuper relative to TnpB (b) measured with ampliconNGS at indicated targets within the Pcsk9 gene in Hepa 1-6 cells (n = 3 biological replicates). Each nuclease-guide RNA combination was delivered on a "all-in-one" plasmid.
Bars represent mean ± standard deviation. Fold change was calculated from group means, with propagated standard deviation shown as error bars.

Figure S11. Improved variant dCas12f1Super outperforms dCasMINI as an adenine base editor.
(**a**) Fold change of overall adenine base editing activity of dCas12f1Super relative to dCasMINI measured with ampliconNGS at Target CM1 and indicated endogenous sites in HEK293T cells (n = 3 biological replicates).
Bars represent mean ± standard deviation. Fold change was calculated from group means, with propagated standard deviation shown as error bars.
(**b**) Frequency or reads without and with indels upon editing TTR site with adenine base editor based on dCasMINI or dCas12f1Super in HEK293T cells (n = 3 biological replicates).
Each dot represents an individual biological replicate, and bars represent the mean.

Figure S12. Gating strategies for flow cytometry and FACS experiments related to testing editing at HDR reporter targets and endogenous sites in HEK293T and Hepa 1-6 cell lines.
(**a**) Representative plots showing the gating strategy for measuring HDR at HDR reporter targets in HEK293T cell line. Cells were gated for live cells, singlets, BFP positivity, mCherry positivity (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct) and finally, for EGFP positivity. In case of "edited" sample one of the CasMINI variants was co-delivered with sgRNA for Target CM1 together with ssODN. "BFP" only and "not transfected" samples serve as controls for gating. In case of "BFP only" sample only sgRNA targeting Target CM1 was delivered.
(**b**) Representative plots showing the gating strategy for FACS-sorting HEK293T or Hepa 1-6 cells for measuring editing. Cells were gated for live cells, singlets and mCherry positivity. In case of "edited" sample CasMINI and sgRNA were delivered on the same "all-in-one" plasmid. "Not transfected" sample serves as control for gating.

a

edited

BFP only

mCherry only

not transfected

b

edited

not transfected

HEK293T

Figure S13. Gating strategy for FACS experiments related to testing editing at endogenous sites in HEK293T and NIH-3T3 cell line and to big scale selection of CasMINI and TnpB for HDR in HEK293T cell line.

(**a**) Representative plots showing the gating strategy for FACS-sorting HEK293T and NIH-3T3 cell lines for measuring NHEJ at endogenous sites. Cells were gated for live cells, singlets and mCherry/BFP positivity (BFP marks the guide-expressing construct; mCherry marks the nuclease-expressing construct). In case of "edited" sample CasMINI was co-delivered with sgRNA and ssODN in HEK293T cells. "BFP only", "mCherry only" and "not transfected" samples serve as controls for gating.

(**b**) Representative plots showing the gating strategy for FACS-sorting during round 1 of HDR selection of CasMINI in HEK293T-reporter cell line. Cells were gated for live cells, singlets, SYTOXTM-Red-negativity, and finally for mCherry/EGFP positivity. In case of "edited" sample sgRNA for Target CM1 was delivered together with ssODN. "Not transfected" and "HEK293T" samples serve as controls for gating. In case of "not transfected" control sgRNA and ssODN were not delivered.