Common marmosets (*Callithrix jacchus*) do not differentiate between familiar and
unfamiliar conspecifics based on pitch contour information

Julia Victoria Grabner[1*], Emma Pigmans[2], Monika Mircheva[3], Job Morgan Knoester[1], Robert
Baumgartner[4], Ulrich Pomper[5], Thomas Bugnyar[1], Michelle J Spierings[1,2]

[1]Department of Behavioral and Cognitive Biology, University of Vienna, Vienna, Austria
[2]Animal Sciences, Institute of Biology, Leiden University, Leiden, The Netherlands
[3]Department of Evolutionary Anthropology, University of Zurich, Zurich, Switzerland
[4]Acoustics Research Institute, Austrian Academy of Sciences, Vienna, Austria
[5]Department of Cognition, Emotion and Methods in Psychology, Faculty of Psychology,
University of Vienna, Austria
[*]Corresponding author: julia.victoria.grabner@univie.ac.at

ORCID
Julia Victoria Grabner: 0009-0006-2512-6322
Job Morgan Knoester: 0009-0008-9946-3887
Robert Baumgartner: 0000-0003-0899-4903
Ulrich Pomper: 0000-0002-6534-968X
Thomas Bugnyar: 0000-0002-6072-9667
Michelle J Spierings: 0009-0004-7458-7251

Funding information
This work was supported by a Young Independent Researchers Group Grant (project number
ZK66 [10.55776/ZK66]) funded by the Austrian Science Fund (FWF) to M.J.S., R.B. and U.P.
and a Veni grant (31 212.264) funded by the Dutch Research Council (NWO) to M.J.S.. J.V.G
and J.M.K. were supported by the project DK Cognition and Communication 2 (project number
W1262-B29 [10.55776/W1262]) funded by the Austrian Science Fund (FWF). J.V.G was
additionally supported by a Marietta Blau Grant (MPC-2024-03740) funded by the Austrian
Federal Ministry of Education, Science and Research (BMBWF) and administered by the
Austrian Agency for Education and Internationalisation (OeAD). For open access purposes, the
authors have applied a CC BY public copyright license to any author-accepted manuscript
version arising from this submission.

Acknowledgements

We thank Steffen Hage, University of Tübingen for providing phee call recordings used for the unfamiliar stimuli. Furthermore, we are grateful to Awani Bapat, Christian Blum and David Meijer for valuable assistance in data analysis.

Ethical approval

The housing conditions and the experimental design were in accordance with the Austrian legislation and the *European Association of Zoos and Aquaria's* (EAZA) husbandry guidelines for Callitrichidae. The research procedures were approved by the Animal Ethics and Experimentation Board of the Faculty of Life Sciences of the University of Vienna (No. 2023-014). Individuals were neither food- nor water deprived at any point. Participation in the experiment was voluntary, and animals were never directly handled. Individuals entered the research room through a custom tunnel system, and we immediately terminated experiments in case the individual showed signs of distress.

Author contribution

J.V.G.: conceptualization, data curation, formal analysis, investigation, methodology, visualization, writing – original draft, writing – review and editing; E.P.: data curation, formal analysis, investigation, writing – review and editing; M.M.: conceptualization, methodology, writing – review and editing; J.M.K.: data curation, formal analysis, writing – review and editing; R.B.: funding acquisition, methodology, writing – review and editing; U.P.: funding acquisition, methodology, writing – review and editing; T.B.: conceptualization, funding acquisition, methodology, project administration, resources, supervision, writing – review and editing; M.J.S.: conceptualization, funding acquisition, methodology, project administration, resources, supervision, writing – review and editing

Abstract

Previous studies suggest that vocal-learning species use variations of prosodic cues, such as changes in pitch, amplitude, or duration, in the conveying of meaning and emotions as well as individual recognition. Although it is known that the long-distance contact calls ("phee" calls) of common marmosets, a vocal non-learner, vary in prosodic information depending on individual characteristics, evidence of the species' use of this information only recently started to emerge. In this study we tested 18 captive common marmosets' (*Callithrix jacchus*) reactions to playbacks of familiar and unfamiliar individuals' phee calls and extracted pitch-contour

information. The playbacks consisted of i) natural phee calls from group members or unfamiliar
conspecifics, or ii) synthesized pitch contours of familiar or unfamiliar phee calls following a
natural or artificial syllable order as well as pure tones of similar frequency and duration. We
found that although individuals seemed to show different reactions towards the natural calls of
familiar and unfamiliar individuals', we did not find group level differences. Marmosets
furthermore show significantly reduced interest in synthesized pitch contours compared to
natural phee calls and did not differentiate between pitch contour playbacks and pure tone
control stimuli, indicating that they did not categorize the pitch contours as conspecific calls.
Our results suggest that extracted pitch contour information alone is not sufficient for
marmosets to recognise conspecifics, however further studies are required to investigate the
exact mechanism of individual recognition in marmosets.
80

81   **Keywords:** individual recognition, prosody, non-human primate, auditory perception

Introduction

While living in social groups offers advantages like predator defence and foraging success (e.g., Rubenstein, 1978), it also introduces complex cognitive challenges. Hence, social interactions within and outside of a group have been proposed as one of the main drivers of cognitive evolution (Ashton et al., 2020; Byrne & Whiten, 1988; Dunbar, 1998). One cognitive challenge is to individually identify conspecifics. Differentiating between group members and outsiders, as well as individually recognising them and remembering their previous interactions with them, is crucial to navigate social life, cooperate and compete with others. Indeed, studies in various social primate species found that individuals can differentiate between group members, familiar outgroup individuals (i.e. neighbouring individuals) and unfamiliar individuals and adjust their reactions accordingly, showing for example differences in aggression or retreat behaviour between neighbours and strangers (e.g., French et al., 1995; Herbinger et al., 2009; Wich et al., 2002). However, the exact characteristics that are used to differentiate between individuals vary between species and situations and often remain unclear.

Depending on their environment and species-specific traits, individuals may rely on visual (e.g., Parr et al., 2000; Pokorny & de Waal, 2009a; Pokorny & de Waal, 2009b), olfactory (e.g., Henkel & Setchell, 2018; Smith et al., 1997) or acoustic information (e.g., Cheney & Seyfarth, 1980; Herbinger et al., 2009; Pereira, 1986; Rendall et al., 1996) or a combination of those modalities to discriminate between individuals. Acoustic signals seem especially important in species living in dense habitat, big complex groups or groups spread over large territories (Fichtel & Manser, 2010; Xie et al., 2024). Not only the vocalisation type itself, but also prosodic cues – variations in syllable duration, pitch and amplitude – prove to be important to provide context (see for example Filippi (2016) for a review on emotional prosody). Studies show that animals adapt their calling rates and acoustic properties of their vocalisations according to their surroundings (e.g., Ey et al., 2009; Koda et al., 2008; Sugiura, 2007). Furthermore, acoustic signals allow the extraction of several characteristics of an animal, such as their emotional state (Briefer, 2012), approximate size, sex, age and dominance status (e.g., Ey et al., 2007; Fischer et al., 2004; Norcross et al., 1999) as well as individual identity (e.g., Marler & Hobbett, 1975). Despite these individual differences in vocalisations and the fact that several animal species can distinguish individuals based on their vocalisations, studies investigating which characteristics individuals use are sparse.

One study on greater false vampire bats (*Megaderma lyra*) using synthesized conspecific calls showed that they could differentiate calls with different prosodic cues and mainly used pitch to

do so (Janßen & Schmidt, 2009). Further evidence about the importance of different prosodic cues comes from human and animal studies using discrimination tasks based on human speech syllables investigating which information is used to distinguish between acoustic signals. While humans can distinguish between different made-up words using only amplitude or pitch alone, Budgerigars (*Melopsittacus undulatus*) require more than one prosodic cue to distinguish two different stimuli (Hoeschele & Fitch, 2016). In a similar study Zebra finches (*Taeniopygia guttata*) performed comparably to humans, being able to use a single prosodic cue for differentiation between syllables, with pitch being the most salient cue (Spierings & ten Cate, 2014). While it seems that vocal-learning species display advanced abilities, rats (*Rattus norvegicus*), a vocal non-learning species, were shown to only be able to discriminate between human speech syllables when pitch, duration, amplitude and vowel quality information were available (Toro & Hoeschele, 2017). Just removing one of those prosodic cues led to very poor performance, suggesting a potential link between vocal learning and the perceptive abilities of prosodic cues. However, whether the same prosodic cues are also important in intraspecific communication and whether a similar difference between vocal learners and vocal non-learners exists remains unknown.

Common marmosets' (*Callithrix jacchus*) long-distance contact calls ("phee" calls) exhibit differences in prosodic-like features that are context- and sex dependent (Norcross & Newman, 1993) as well as group- and individual- specific (Miller et al., 2010; Zürcher & Burkart, 2017; Zürcher et al., 2019). Playback studies furthermore show that marmosets differentiate calls from different conspecifics (Kato et al., 2018; Miller & Thomas, 2012; *Callithrix jacchus*) and seem to categorise calls depending on callers' sex (Norcross et al., 1994; *Callithrix jacchus*) or familiarity (Rukstalis & French, 2005; *Callithrix kuhlii*). This suggests that vocalisations encode characteristics that allow individuals to individually recognize conspecifics. Although it has been found that common marmosets' vocalisations show individual characteristics in prosodic information and that individuals can differentiate between conspecifics, it is not known if they are able to differentiate conspecifics from a single salient cue or if they, as a vocal non-learner, need the complete prosodic information.

In this study, we used a playback paradigm to investigate whether common marmosets use prosodic information to individually identify conspecifics. We first investigated if marmosets show a difference in vocal response and looking time towards i) familiar and unfamiliar conspecifics' natural unaltered contact ("phee") calls, ii) extracted pitch information of those phee calls, as well as iii) control sounds with similar acoustic properties. As common marmosets

are territorial and show strong reactions towards unfamiliar individual's phee calls (e.g., Caselli

et al., 2018) we predicted that individuals would show clear differences in looking times with

longer looking times during playbacks of unfamiliar phee calls compared to familiar calls and

controls. Furthermore, as marmosets are known to engage in vocal turn-taking (Takahashi et

al., 2013), we predicted that individuals would vocally respond to the playbacks. We expected

them to vocalize more often after familiar individuals' calls, compared to unfamiliar

individuals' calls and pure tone control stimuli. For the extracted pitch information, we had two

alternative expectations: if pitch is the prominent information used by individuals, we expected

individuals to show similar patterns of responses when presented with unaltered playbacks or

extracted pitch information. However, if animals need more acoustic information to distinguish

conspecifics, we would expect individuals to not show any differentiation between playbacks

of familiar and unfamiliar individuals' pitch contours.

Material and Methods

Subjects

For this study we tested 18 adult common marmosets (6 f/12 m; mean ± SD: 9.4 ± 4.9 years) at

the Animal Care Facility of the Faculty of Life Sciences, University of Vienna, Vienna, Austria.

Individuals were housed in family groups of three to seven individuals in three separate keeping

rooms. Enclosures were equipped with various items such as branches, wooden boards,

(sleeping) baskets, hammocks, foraging boxes, heating- and UV-lamps and had bark mulch

floor covering. Additionally, individuals had access to roofed outdoor enclosures when

temperatures allowed. Enclosures were regularly enriched with a variety of food- and non-food

items. The monkeys were fed three times a day with a rich selection of vegetables, animal

protein and fruits and had ad libitum access to water. Gummi arabicum as well as additional

vitamin- and mineral supplements were provided daily.

Experimental procedure and stimuli

We performed a playback experiment during which each tested marmoset was presented with

a set of stimuli including altered and unaltered vocalisations of familiar and unfamiliar

conspecifics. We then measured the reaction of the tested individual towards those stimuli by

scoring behavioural responses. The experiment was divided into two conditions which were

tested with the same individuals in consecutive weeks: the Natural call condition, and the Pitch

contour condition, that are explained in more detail below. Before, between and after the stimuli

pink noise of random durations (10-30 seconds) was played to avoid habituation. Familiar

stimuli were vocalisations from individuals of the same family group. Unfamiliar stimuli were

vocalisations from individuals from separate colonies in Zurich and Tübingen which the test
subjects have never heard or seen before. All vocalisation stimuli were recorded during
habituation sessions while a single individual was separated from its family group. Recordings
were edited in Praat (version 6.2.14; Boersma, 2001) to match in amplitude, and single stimuli
were cut and combined into one playback. In each test session the calls of a different familiar
and unfamiliar conspecific were presented, while within a test sessions different calls of the
same familiar and unfamiliar individual were used. Example stimuli can be accessed on our
Open Science Framework repository (https://osf.io/kn3pt).

Natural call condition
In the Natural call condition, we wanted to find out how individuals react to playbacks of natural
calls from familiar and unfamiliar individuals. To do this, recordings of phee calls of familiar
and unfamiliar conspecifics as well as control sounds were used. Playbacks consisted of natural
three-syllable loud phee calls or three pure tones with similar properties (1 second pure tones,
7 – 8.4 kHz, 10 ms ramp-up/ramp-down, 300ms silence) as control (see figure 1*a,b* for
spectrograms of example stimuli). Each test session contained four randomized sets of stimuli,
each set comprising one familiar individual's phee call, one unfamiliar individual's phee call
and one control sound, making a total of 12 stimuli per session.

Pitch contour condition
In the Pitch contour condition, we investigated marmosets' reactions to extracted pitch
information of natural phee calls. Here, the pitch contour of each syllable of the call was
extracted and saved separately. This was done using the built-in Praat function "extract pitch
tier". The extracted pitch contour, containing only the fundamental frequency, was then
synthesized to sound (sine). After resynthesizing, the single syllables were concatenated to a
three-syllable string with 300ms of silence between syllables. In this step two different stimulus
types were produced: the *natural order* stimuli, that consisted of all three syllables of the natural
call in original order (1-2-3), and the *first syllable only* stimuli, in which only the first syllable
was used and repeated three times to exclude further temporal information (1-1-1) (see figure
1*c,d* for spectrograms of example stimuli). In each session of the Pitch contour condition three
randomized sets, each containing one familiar natural order pitch contour, one familiar first
syllable only pitch contour, one unfamiliar natural order pitch contour, one unfamiliar first
syllable only pitch contour and one control sound were presented, resulting in a total of 15
stimuli per session.

Experimental setup and data collection

In both conditions subjects were tested individually in a separate research room. After entering

the cage compartment (1m x 1m x 1m) through a tunnel system, the individual sat on a small

perch facing a loudspeaker. The experimenter then hid behind a curtain and started the trial. The

stimuli were played back from a loudspeaker (Anchor Explorer Pro) placed 1.5m from the

experiment compartment and hidden behind a fake plant, using Audacity (version 3.2.5) on a

MacBook pro13. Volume levels on Audacity, the laptop and the speaker were kept constant

throughout the experiment, such that the sound had a mean sound pressure level of 50dB in the

middle of the experimental cage. All experimental sessions were filmed using two cameras: The

main camera (Canon legria HF G25) was placed on top of the speaker directly facing the

experimental cage to measure the time the individuals look in the direction of the sound source.

Camera 2 (Canon legria HF R806) was mounted above the cage for a top view (figure 2).

Recordings from camera 2 were only used when an individual was out of view from the main

camera. Fourteen individuals were tested in three test sessions for each condition (Natural call

and Pitch contour). Each individual was tested once a week over the course of 8 weeks (three

weeks data collection for Natural call condition, two weeks break, three weeks data collection

for Pitch contour condition). Due to changes in group composition four individuals were only

tested in one (Lokri) or two (Kobold, Luna and Jack) test sessions per condition over the course

of 4 or 6 weeks, respectively.

Video coding

The videos were analysed by a single observer (E.P.) in BORIS (version 7.13; Friard & Gamba,

2016). The observer coded each video twice. In the first round all videos were analysed without

sound to code looking time, the amount of time the individual looked in the direction of the

speaker, while being blind to the stimulus category and onset. In the second round, the

frequency of contact ("phee") call vocalisations and stimuli on- and offset were coded. To

measure vocal responses to the playback we coded all phee calls occurring between the end of

the stimulus and 10 seconds after. Due to different lengths of stimuli and individuals sometimes

being out of sight we used proportions of looking times for the analysis. 10% of the videos were

additionally coded by a second observer (J.V.G.) to ensure interobserver reliability (IOR).

Interobserver reliability was excellent for overall duration (two-way intraclass correlation

coefficient: ICC (A,1) = 0.998) and frequency (ICC(A,1) = 0.972) of coded variables, as well

as looking time durations (ICC(A,1) = 0.921). Reliability was good for frequency of phee call

vocalisations (ICC(A,1) = 0.897).

Statistical analysis

 General approach

Data extraction and structuring as well as statistical analysis were done in R (version 4.5.0; R

Core Team, 2025) using RStudio (version 2024.12.1+563; Posit team, 2025). As response

variables we used the proportion of time the individuals were looking at the speaker (looking

time proportion) as well as a binary response (yes/no) variable indicating whether the

individuals responded by looking at the speaker (looking response) or producing phee calls

within ten seconds of the stimulus (vocal response). For the analysis we used Generalised

Linear Mixed Models (GLMM) with the function glmmTMB (Brooks et al., 2017) and applied

the full-null model comparison framework, where we compared a full model including all test

predictors, representing the working hypothesis against a null model excluding all test

predictors, representing the null hypothesis, to control for cryptic multiple testing (Forstmeier

& Schielzeth, 2011). If the model comparison was significant, meaning the sum of test

predictors significantly improved the model fit, we continued with model diagnostics and

investigated the model results. If the model comparison was not significant, we halted analysis

and did not further investigate the model. If the model comparison showed trend-level effects

(*p* values over 0.05 but below 0.1) we decided to follow the approach proposed by Titchener et

al. (2023), to not ignore the possibility of an effect and report both the full model and the

reduced model as well as the resulting analysis. We do however advice the readers to treat these

results with caution.

Before the model comparison, we performed model diagnostics. Firstly, we evaluated the

variance inflation factors (VIFs) using the function vif (of the package "car", version 3.1-3) to

investigate the extent of collinearity of predictors. Secondly, we checked model dispersion

using the custom function overdisp.test (Mundry, 2024 personal communication). For

binominal models we additionally produced basic diagnostic plots using the package DHARMa

(version 0.4.7; Hartig, 2024). Lastly, we visually inspected the plots of the best linear unbiased

predictors (BLUPs). These were also used to aid in the decision which random slopes to first

exclude in the case of model convergence problems. If model diagnostics showed acceptable

model fit the model results were investigated using the summary() function to compare levels

within a predictor against the reference. The drop1() function (with the argument test = "Chisq")

was then used to assess the influence of each fixed effect by excluding one fixed effect at a time

and comparing this reduced model to the full model including the effect. Where applicable

pairwise post hoc tests were conducted using the emmeans package (version 1.11.0; Lenth,

2025) and adjusted using Tukey correction. As an additional step we investigated model

stability by dropping one level of a specified grouping factor (in our case "subject") and

comparing the resulting estimates against each other using the function influence_ mixed from
the package glmmTMB.

Looking response / - time

First, we investigated differences in looking time proportions between stimuli within the
Natural call condition as well as the Pitch contour condition. Because of substantial zero
inflation of the response variable looking time proportion, we opted for a two-step approach:
First, we transformed the response variable to the binominal variable looking response,
representing a binary yes/no response. We then ran a GLMM with a binominal distribution to
compare between reaction and no reaction. Then, we excluded zeros and ran an additional
GLMM with beta regression, to test whether looking time proportion differed between stimuli.

For both the binominal and the beta model, we included as fixed effects test predictors the
interactions of stimulus type with the subject's sex, and stimulus type with the subject's age
including their main effects. We included the session number and set number as fixed effects as
control predictors. As random effects we included individual identity, family group, session
number and set number, plus all theoretically identifiable random slopes using the function
fe.re.tab (Mundry, 2023). The resulting null models included the same fixed and random effects
control predictors, but not the fixed effects test predictors. In case the following full-null model
comparison was significant, but an interaction had no significant effect, we formulated a null
model lacking that interaction, but containing its main effects, to check for an effect of those
main effects in absence of an interaction. In the case of model convergence issues, we simplified
the model stepwise: First we removed correlations between random effects intercepts and
random slopes. Then we excluded random slopes based on their biological, ecological relevance
and their expected importance in the experimental design. Once a full model converged and we
had to further simplify the random slopes because of convergence issues in the null model or
drop1 sub-models, we also used the BLUPs to assist in our decision making.

Vocal response

To investigate differences in vocal responses between stimuli within the Natural call condition
and the Pitch contour condition we again first transformed the response variable to the
binominal variable vocal response, representing a binary yes/no response. We then used a
GLMM with a binominal distribution including the same fixed and random effects as for
looking responses. As multiple phee call vocalisations were nearly never observed (only 4 out
of 588 cases in the Natural call condition and only 3 out of 735 cases in the Pitch contour
condition) we did not run further analyses for the frequency of phee call vocalisations.

Comparison Natural call condition and Pitch contour condition

To compare looking responses and -times between the Natural call condition and Pitch contour

condition we again used a GLMM with binominal and a GLMM with beta regression, both

including the experimental condition as test predictor and including the same fixed and random

effects as for looking responses but not including interactions with the subject's sex or age as

we did not expect any interactive effects between the experimental condition and the

individuals' characteristics.

Results

Natural call condition

Common marmosets show interest in conspecific´s natural phee calls

In the Natural call condition, for the binominal model investigating looking responses

(responses versus no responses) the full model including stimulus performed significantly better

than the null model ($\chi^2(6) = 16.02$, $p = 0.014$). Looking responses differed significantly between

the stimulus categories ($\chi^2(2) = 13.09$, $p = 0.001$) (figure 3a). A Tukey-adjusted post-hoc

pairwise comparison revealed that individuals responded more often when a conspecific call

was played back compared to a control stimulus (control vs familiar: estimate = -0.99, *SE* =

0.24, $z = -4.14$, $p < 0.001$; control vs unfamiliar: estimate = -1.00, *SE* = 0.34, $z = -2.90$, $p =$

0.011). However, although density plots indicate more interest in unfamiliar calls over familiar

ones (figure 4a) we did not find a difference in looking responses between the familiar and

unfamiliar stimulus (estimate = -0.01, *SE* = 0.34, $z = -0.02$, $p = 0.999$). For the beta model

investigating looking time proportions, the full model including stimulus type did not perform

significantly better compared to the null model ($\chi^2(6) = 1.41$, $p = 0.965$).

Common marmosets react differently to familiar and unfamiliar individual´s natural phee

calls on an individual level

Looking at figure 5a, it seems that although on a group level no difference in reaction between

familiar and unfamiliar calls was found, on an individual level marmosets might react

differently to familiar and unfamiliar conspecifics' phee calls, but in different directions and to

different extents. To investigate a possible effect of individual characteristics on the looking

response and -times we included the individuals' age and sex in the analysis. We did not find a

significant effect of those characteristics on overall looking responses in the binominal model

(sex: $\chi^2(1) = 0.001$, $p = 0.978$; age: $\chi^2(1) = 0.04$, $p = 0.836$). Interactions between sex or age and

the stimulus did not significantly improve the binominal model fit (stimulus:sex: $\chi^2(2) = 2.11$,

$p = 0.348$; stimulus:age: $\chi^2(2) = 0.22.$, $p = 0.897$), and graphical inspection does not show any

clear pattern (figure 6).

Marmosets rarely vocally respond to familiar and unfamiliar individual´s playbacks

349 Overall vocal responses were rare, with individuals only producing phee calls in response to

350 138 of 588 total stimuli. For the binominal model investigating the effect of vocal responses the

351 full model including the stimulus type did not perform significantly better compared to the null

352 model ($\chi^2(6) = 8.11$, $p = 0.230$).

Pitch contour condition

Common marmosets do not react differently to familiar and unfamiliar individual´s pitch
contours on a group level

In the Pitch contour condition, the full binominal model including stimulus did not improve the

model fit compared to the null model for both looking response ($\chi^2(12) = 6.90$, $p = 0.864$) and

vocal response ($\chi^2(12) = 8.61$, $p = 0.736$). When excluding zero responses, the full beta model

for looking time proportions again did not perform significantly better than the null model

($\chi^2(12) = 19.25$, $p = 0.083$). However, as the model comparison showed a trend level effect, we

report the model results but advice the readers to carefully evaluate these findings. As

interactions did not significantly improve the model fit (stimulus:sex: $\chi^2(4) = 4.11$, $p = 0.391$;

stimulus:age: $\chi^2(4) = 3.99$, $p = 0.408$) we fitted a reduced model including only the main effects

without interactions. We found that looking time proportions significantly differed between

stimuli ($\chi^2(4) = 9.81$, $p = 0.044$). However, post-hoc pairwise comparisons did not yield

significant differences (figure 3b).

Common marmosets do not react differently to familiar and unfamiliar individual´s pitch
contours on an individual level

We could not observe clear individual preferences in looking time proportions in the Pitch

contour condition. Differences in reactions between familiar and unfamiliar stimuli seem to be

considerably less pronounced compared to the Natural call condition and preferences do not

match the ones observed in the Natural call condition (figure 5b). Similarly to the Natural call

condition, sex ($\chi^2(1) = 0.09$, $p = 0.769$) and age ($\chi^2(1) = 0.23$, $p = 0.632$) did not significantly

influence overall looking time proportions (figure 6).

Comparison Natural call condition and Pitch contour condition

Common marmosets show less interest in pitch contours compared to natural call playbacks

The observed lack of differentiation between stimuli in the Pitch contour condition could be attributed to a loss of interest or habituation effect. Indeed, for both the binominal model investigating looking responses and the beta model investigating looking time proportions the full model including the experimental condition performed significantly better than the null model (looking responses: $\chi^2(1) = 7.27$, $p = 0.007$; looking time proportions: $\chi^2(1) = 23.68$, $p < 0.001$). Looking responses ($\chi^2(1) = 7.27$, $p = 0.007$) and looking time proportions ($\chi^2(1) = 23.68$, $p < 0.001$) both significantly decreased between the Natural call condition and the Pitch contour condition (figure 4b). However, while individuals' interest decreased between testing sets within a session (looking response: $\chi^2(1) = 5.23$, $p = 0.022$; looking time proportion: $\chi^2(1) = 3.72$, $p = 0.054$) there was no significant decrease in looking over testing sessions (looking response: $\chi^2(1) = 0.02$, $p = 0.902$; looking time proportion: $\chi^2(1) = 1.03$, $p = 0.311$), indicating only short time habituation or loss of interest (figure 4c,d). Furthermore, we found a significant increase in looking time proportions ($\chi^2(1) = 4.08$, $p = 0.043$) but not looking responses ($\chi^2(1) = 0.02$, $p = 0.876$) with age.

Discussion

We found that individuals' reactions towards familiar and unfamiliar stimuli did not differ when presented with pitch contour information only. This suggests that pitch information alone is not sufficient for marmosets to differentiate between conspecifics. Interestingly however, they also did not show different reactions between conspecific pitch contours and control stimuli. The overall decrease of looking time proportions between the Natural call condition and the Pitch contour condition suggests that marmosets were not interested in the pitch contour stimuli.

The reduced interest suggests an inability to perceive pitch contours as conspecific calls

As we observed a considerable decrease in attention between the two conditions but no significant decrease between testing sessions within each condition, it is unlikely, that the decrease is due to a habituation effect. Rather it seems that the pitch contours were not perceived as conspecific calls, leading to individual's interest dropping to control stimulus levels. One possibility is that the manipulation to extract the pitch contours made them too artificial for the individuals to be interesting. However, studies show that marmosets react similarly to natural and synthesized phee calls (e.g., Norcross et al., 1994). On the other hand literature suggests, that while vocal learners can distinguish between sounds using only limited prosodic information (Hoeschele & Fitch, 2016; Janßen & Schmidt, 2009; Spierings & ten Cate, 2014), vocal non-learners need the full spectrum of prosodic cues for the same task (Toro & Hoeschele, 2017). If the same is true for intraspecific communication, excluding prosodic information

would make it impossible for vocal non-learning individuals such as marmosets to distinguish between calls and could possibly lead to the observed loss of interest in the Pitch contour condition. While the current study design does not allow to disentangle between a possible effect of artificial sound manipulation and the general ability to extract information from one prosodic cue alone, we show that the marmosets clearly respond less to phee calls when only the fundamental frequency is maintained.

Individual characteristics are likely to lead to differing responses

Surprisingly, in contrast to previous studies investigating marmoset´s reactions to conspecific calls (e.g. *Callithrix jacchus*: Miller & Wang, 2006; *Callithrix kuhlii*: Rukstalis & French, 2005) we did not find a categorical difference in reaction towards familiar and unfamiliar individuals' calls. However, our data indicates there might be various effects of individual differences. Based on the graphical representation it seems that in the Natural call condition most individuals preferably react to one of the two categories. This preference however seems to vary in direction and strength between individuals which might lead to no clear pattern emerging on the group level. The discrepancy in reaction compared to what Rukstalis and French (2005) found could be caused by the fact that in our study additional individual characteristics could play a role. While they only used playbacks of pair bond partners or unfamiliar individuals of the opposing sex, we used familiar and unfamiliar playbacks originating from both males and females of varying age (2-17 years) and each reproductive status (breeders, helpers and pair-kept individuals without breeding experience). Furthermore, relationships between the individuals and playbacks used in this study varied. Individuals heard familiar calls from their pair partners, siblings, parents or offspring. According to previous studies individuals' reactions depend both on their own characteristics as well as the characteristics of the played back individual and an interplay of both. For example, Caselli et al. (2018) found that, when being played back unfamiliar individuals' calls, breeding individuals reacted first to playbacks of the opposite sex and breeding females reacted with pilo-erection to playbacks of unfamiliar females. Another study from Norcross and colleagues (1994) also shows differences in reaction, with breeding individuals reacting significantly stronger than juveniles and differences in the exhibited behaviours between males and females. Although our sample size and the number of potentially influencing variables did not allow us to investigate demographic effects in more detail, the possible presence of individual preferences might suggest that marmosets were able to differentiate between familiar and unfamiliar individual´s calls.

The absence of vocal responses highlights the importance of playback structure

Another surprising result was the lack of vocal responses. The tested individuals rarely

vocalized within the set time window of 10 seconds after the playbacks. Although, to our

knowledge, no study directly investigated differences in vocal response behaviour towards

familiar and unfamiliar playbacks, we expected individuals to strongly respond with phee calls

to conspecific vocalisations as shown in previous studies. In a study by Norcross and colleagues

(1994) marmosets produced twice as many phee calls after than before the playback of an

unfamiliar individual´s phee call, but did not show significant differences in response rates

towards playbacks from different sexes or call variants. The way the playback was presented in

our study might have influenced the individual´s responses. Previous studies found, that

individuals do not react differently to playbacks of phee calls and silent playbacks when the

playback is not presented in an interactive way (Miller & Wang, 2006) and when latencies

between an individuals' phee call and playbacks are too long (Miller et al., 2009). Furthermore,

it was found, that individuals responded less often after the identity of the played back

individual changed, than when the identity of the counterpart stayed the same (Miller &

Thomas, 2012). In our study each playback session consisted of calls from two different

individuals played back in a semi-random order. Moreover, playbacks were not presented in an

interactive way but in a fixed timescale and order, resulting in individuals sometimes starting

to vocalize shortly before the start of a stimulus and leading to overlaps between the

vocalisations and the playbacks. The comparison between our findings and the literature shows

that the way playbacks are presented can have a strong influence on the exposed subjects`

response.

Age is unlikely to drive the observed effects

The overall low number of observed reactions raises the question whether the individuals

perceived the stimuli correctly. One potential factor influencing the marmosets' reactions could

be their age. Marmosets above 8 years of age are often classified as "aged" and exhibit diverse

symptoms of age-related pathologies (Abbott et al., 2003; Tardif et al., 2011). Studies show that

older age is related to hearing-loss in marmosets (Harada et al., 1999; Sun et al., 2021). Two

reasons however speak against this explanation: Firstly, age-related hearing loss in marmosets

is described to be only moderate (Harada et al., 1999) and primarily affects ultra-high

frequencies above 16kHz (Sun et al., 2021). It is therefore unlikely to influence the perception

of phee calls that have a substantially lower fundamental frequency (5.5-10.5 kHz; Norcross &

Newman, 1993). Secondly, if hearing loss would be the reason for the lack in response, we

would expect individuals to also show overall declined reactions and no differentiation in

reaction towards control stimuli and conspecific stimuli. We do however find that overall

proportional looking time increased with age and that individuals do differentiate between control and conspecific stimuli in the Natural call condition. Taken together, although age might be an influencing factor it cannot fully explain the observed results. Rather, although we cannot statistically show it, we think the effect might stem from individuals displaying individual differences in their reactions and them not classifying the pitch contours as conspecific calls.

Our results give limited insight into individual recognition in marmosets

Our results only partially shed light on the acoustic characteristics used in individual recognition in marmosets. While it is known from previous studies that marmosets can differentiate between conspecifics based on vocalisations alone (Kato et al., 2018; Miller & Thomas, 2012) we set out to investigate which acoustic cues are used to do so. Marmosets are vocal non-learners and one vocal non-learning species has been shown to perform worse when differentiating sounds based on only one prosodic cue at least when using human speech syllables (Toro & Hoeschele, 2017). However, it is not clear yet whether vocal learning and the ability to use certain acoustic cues is connected. We therefore expected them to either show similar or less pronounced differences in reactions towards familiar and unfamiliar conspecifics' calls, depending on whether the inability to use single prosodic cues expands to marmosets. In line with the latter predictions, we found that marmosets do not react differently towards extracted pitch contours of familiar and unfamiliar individuals. However, given that the individuals did not show clear differentiation between familiar and unfamiliar calls in the Natural call condition and did not differentiate between pitch contours and control sounds we cannot conclude whether marmosets need more information or even the full acoustic spectrum or whether the observed behaviour results from other factors such as habituation, loss of interest or too artificial stimuli.

Conclusion

Taken together our results suggest that marmosets react differently to playbacks of natural familiar and unfamiliar conspecifics' contact calls on an individual basis. On a group level though, overall effects are outweighed by these individual preferences. When being presented with only the extracted pitch contours however, individuals seem to not differentiate anymore. Additionally, individuals showed no differentiation between extracted pitch contours and pure tone control sounds. Further studies are needed to disentangle whether marmosets cannot use pitch alone to distinguish between individuals or whether simply extracting the pitch contour and playing back a resynthesized version is too artificial for the monkeys to be perceived as a conspecific call.

References

Abbott, D., Barnett, D., Ricki, J., Yamamoto, M. E., & Schultz-Darken, N. (2003). Aspects of common marmoset basic biology and life history important for biomedical research. *Comparative medicine*, *53*, 339-350.

Ashton, B. J., Kennedy, P., & Radford, A. N. (2020). Interactions with conspecific outsiders as drivers of cognitive evolution. *Nature Communications*, *11*(1), 4937. https://doi.org/10.1038/s41467-020-18780-3

Boersma, P. (2001). Praat, a system for doing phonetics by computer. *Glot. Int.*, *5*(9), 341-345.

Briefer, E. F. (2012). Vocal expression of emotions in mammals: mechanisms of production and evidence. *Journal of Zoology*, *288*(1), 1-20. https://doi.org/10.1111/j.1469-7998.2012.00920.x

Brooks, M. E., Kristensen, K., Benthem, K. J. v., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal*, *9*(2), 378-400. https://doi.org/10.32614/RJ-2017-066

Byrne, R. W., & Whiten, A. (1988). *Machiavellian Intelligence: Social Expertise and the Evolution of Intellect in Monkeys, Apes, and Humans*. Clarendon Press.

Caselli, C. B., Ayres, P. H. B., Castro, S. C. N., Souto, A., Schiel, N., & Miller, C. T. (2018). The role of extragroup encounters in a Neotropical, cooperative breeding primate, the common marmoset: a field playback experiment. *Animal Behaviour*, *136*, 137-146. https://doi.org/10.1016/j.anbehav.2017.12.009

Cheney, D. L., & Seyfarth, R. M. (1980). Vocal recognition in free-ranging vervet monkeys. *Animal Behaviour*, *28*(2), 362-367. https://doi.org/10.1016/S0003-3472(80)80044-3

Dunbar, R. I. (1998). The social brain hypothesis. *Evolutionary Anthropology: Issues, News, and Reviews*, *6*(5), 178-190. https://doi.org/10.1002/(SICI)1520-6505(1998)6:5<178::AID-EVAN5>3.0.CO;2-8

Ey, E., Pfefferle, D., & Fischer, J. (2007). Do age- and sex-related variations reliably reflect body size in non-human primate vocalizations? A review. *Primates*, *48*(4), 253-267. https://doi.org/10.1007/s10329-006-0033-y

Ey, E., Rahn, C., Hammerschmidt, K., & Fischer, J. (2009). Wild Female Olive Baboons Adapt their Grunt Vocalizations to Environmental Conditions. *Ethology*, *115*(5), 493-503. https://doi.org/10.1111/j.1439-0310.2009.01638.x

Fichtel, C., & Manser, M. (2010). Vocal communication in social groups. In P. Kappeler (Ed.), *Animal Behaviour: Evolution and Mechanisms* (pp. 29-54). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-02624-9_2

Filippi, P. (2016). Emotional and Interactional Prosody across Animal Communication Systems: A Comparative Approach to the Emergence of Language [Review]. *Frontiers in Psychology*, *Volume 7 - 2016*. https://doi.org/10.3389/fpsyg.2016.01393

Fischer, J., Kitchen, D. M., Seyfarth, R. M., & Cheney, D. L. (2004). Baboon loud calls advertise male quality: acoustic features and their relation to rank, age, and exhaustion. *Behavioral Ecology and Sociobiology*, *56*(2), 140-148. https://doi.org/10.1007/s00265-003-0739-4

Forstmeier, W., & Schielzeth, H. (2011). Cryptic multiple hypotheses testing in linear models: overestimated effect sizes and the winner's curse. *Behavioral Ecology and Sociobiology*, *65*(1), 47-55. https://doi.org/10.1007/s00265-010-1038-5

French, J. A., Schaffner, C. M., Shepherd, R. E., & Miller, M. E. (1995). Familiarity with Intruders Modulates Agonism towards Outgroup Conspecifics in Wied's Black-tufted-ear Marmoset (*Callithrix kuhli*: Primates, Callitrichidae). *Ethology*, *99*(1-2), 24-38. https://doi.org/10.1111/j.1439-0310.1995.tb01086.x

Friard, O., & Gamba, M. (2016). BORIS: a free, versatile open-source event-logging software for video/audio coding and live observations. *Methods in Ecology and Evolution*, *7*(11), 1325-1330. https://doi.org/10.1111/2041-210X.12584

Harada, T., Tokuriki, M., & Tanioka, Y. (1999). Age-related changes in the brainstem auditory evoked potentials of the marmoset. *Hearing Research*, *128*(1), 119-124. https://doi.org/10.1016/S0378-5955(98)00201-9

Hartig, F. (2024). *DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models*. https://doi.org/10.32614/CRAN.package.DHARMa

Henkel, S., & Setchell, J. M. (2018). Group and kin recognition via olfactory cues in chimpanzees (*Pan troglodytes*). *Proceedings of the Royal Society B: Biological Sciences*, *285*(1889), 20181527. https://doi.org/10.1098/rspb.2018.1527

Herbinger, I., Papworth, S., Boesch, C., & Zuberbühler, K. (2009). Vocal, gestural and locomotor responses of wild chimpanzees to familiar and unfamiliar intruders: a playback study. *Animal Behaviour*, *78*(6), 1389-1396. https://doi.org/10.1016/j.anbehav.2009.09.010

Hoeschele, M., & Fitch, W. T. (2016). Phonological perception by birds: budgerigars can perceive lexical stress. *Animal Cognition*, *19*(3), 643-654. https://doi.org/10.1007/s10071-016-0968-3

Janßen, S., & Schmidt, S. (2009). Evidence for a perception of prosodic cues in bat communication: contact call classification by *Megaderma lyra*. *Journal of Comparative Physiology A*, *195*(7), 663-672. https://doi.org/10.1007/s00359-009-0441-8

Kato, M., Yokoyama, C., Kawasaki, A., Takeda, C., Koike, T., Onoe, H., & Iriki, A. (2018). Individual identity and affective valence in marmoset calls: in vivo brain imaging with vocal sound playback. *Animal Cognition*, *21*(3), 331-343. https://doi.org/10.1007/s10071-018-1169-z

Koda, H., Shimooka, Y., & Sugiura, H. (2008). Effects of caller activity and habitat visibility on contact call rate of wild Japanese macaques (*Macaca fuscata*). *American Journal of Primatology*, *70*(11), 1055-1063. https://doi.org/10.1002/ajp.20597

Lenth, R. V. (2025). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. In https://CRAN.R-project.org/package=emmeans

Marler, P., & Hobbett, L. (1975). Individuality in a long-range vocalization of wild chimpanzees. *Z Tierpsychol*, *38*(1), 37-109.

Miller, C. T., Beck, K., Meade, B., & Wang, X. (2009). Antiphonal call timing in marmosets is behaviorally significant: interactive playback experiments. *Journal of Comparative Physiology A*, *195*(8), 783-789. https://doi.org/10.1007/s00359-009-0456-1

Miller, C. T., Mandel, K., & Wang, X. (2010). The communicative content of the common marmoset phee call during antiphonal calling. *Am J Primatol*, *72*(11), 974-980. https://doi.org/10.1002/ajp.20854

Miller, C. T., & Thomas, A. W. (2012). Individual recognition during bouts of antiphonal calling in common marmosets. *Journal of Comparative Physiology A*, *198*(5), 337-346. https://doi.org/10.1007/s00359-012-0712-7

Miller, C. T., & Wang, X. (2006). Sensory-motor interactions modulate a primate vocal behavior: antiphonal calling in common marmosets. *Journal of Comparative Physiology A*, *192*(1), 27-38. https://doi.org/10.1007/s00359-005-0043-z

Mundry, R. (2023). *some R functions*. In Zenodo. https://doi.org/10.5281/zenodo.7670524

Norcross, J. L., & Newman, J. D. (1993). Context and gender-specific differences in the acoustic structure of common marmoset (*Callithrix jacchus*) phee calls. *American Journal of Primatology*, *30*(1), 37-54. https://doi.org/10.1002/ajp.1350300104

Norcross, J. L., Newman, J. D., & Cofrancesco, L. M. (1999). Context and sex differences exist in the acoustic structure of phee calls by newly–paired common marmosets (*Callithrix*

jacchus). *American Journal of Primatology*, *49*(2), 165-181. https://doi.org/10.1002/(SICI)1098-2345(199910)49:2<165::AID-AJP7>3.0.CO;2-S

Norcross, J. L., Newman, J. D., & Fitch, W. (1994). Responses to natural and synthetic phee calls by common marmosets (*Callithrix jacchus*). *American Journal of Primatology*, *33*(1), 15-29. https://doi.org/10.1002/ajp.1350330103

Parr, L. A., Winslow, J. T., Hopkins, W. D., & de Waal, F. B. (2000). Recognizing facial cues: individual discrimination by chimpanzees (*Pan troglodytes)* and rhesus monkeys (*Macaca mulatta*). *J Comp Psychol*, *114*(1), 47-60. https://doi.org/10.1037/0735-7036.114.1.47

Pereira, M. E. (1986). Maternal recognition of juvenile offspring coo vocalizations in Japanese macaques. *Animal Behaviour*, *34*(3), 935-937. https://doi.org/10.1016/S0003-3472(86)80084-7

Pokorny, J. J., & de Waal, F. B. (2009a). Face recognition in capuchin monkeys (*Cebus apella*). *J Comp Psychol*, *123*(2), 151-160. https://doi.org/10.1037/a0014073

Pokorny, J. J., & de Waal, F. B. (2009b). Monkeys recognize the faces of group mates in photographs. *Proceedings of the National Academy of Sciences*, *106*(51), 21539-21543. https://doi.org/10.1073/pnas.0912174106

Posit team. (2025). *RStudio: Integrated Development Environment for R. Posit software, PBC*. In http://www.posit.co/

R Core Team. (2025). *R: A Language and Environment for Statistical Computing*. In https://www.R-project.org/

Rendall, D., Rodman, P. S., & Emond, R. E. (1996). Vocal recognition of individuals and kin in free-ranging rhesus monkeys. *Animal Behaviour*, *51*(5), 1007-1015. https://doi.org/10.1006/anbe.1996.0103

Rubenstein, D. I. (1978). On Predation, Competition, and the Advantages of Group Living. In P. P. G. Bateson & P. H. Klopfer (Eds.), *Social Behavior* (pp. 205-231). Springer US. https://doi.org/10.1007/978-1-4684-2901-5_9

Rukstalis, M., & French, J. A. (2005). Vocal buffering of the stress response: exposure to conspecific vocalizations moderates urinary cortisol excretion in isolated marmosets. *Hormones and Behavior*, *47*(1), 1-7. https://doi.org/10.1016/j.yhbeh.2004.09.004

Smith, T. E., Abbott, D. H., Tomlinson, A. J., & Mlotkiewicz, J. A. (1997). Differential Display of Investigative Behavior Permits Discrimination of Scent Signatures from Familiar and Unfamiliar Socially Dominant Female Marmoset Monkeys (*Callithrix jacchus*). *Journal of Chemical Ecology*, *23*(11), 2523-2546. https://doi.org/10.1023/B:JOEC.0000006664.38169.e5

Spierings, M. J., & ten Cate, C. (2014). Zebra finches are sensitive to prosodic features of human speech. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1787), 20140480. https://doi.org/10.1098/rspb.2014.0480

Sugiura, H. (2007). Effects of proximity and behavioral context on acoustic variation in the coo calls of Japanese macaques. *American Journal of Primatology*, *69*(12), 1412-1424. https://doi.org/10.1002/ajp.20447

Sun, Z., Cheng, Z., Gong, N., Xu, Z., Jin, C., Wu, H., & Tao, Y. (2021). Neural presbycusis at ultra-high frequency in aged common marmosets and rhesus monkeys. *Aging (Albany NY)*, *13*(9), 12587-12606. https://doi.org/10.18632/aging.202936

Takahashi, Daniel Y., Narayanan, Darshana Z., & Ghazanfar, Asif A. (2013). Coupled Oscillator Dynamics of Vocal Turn-Taking in Monkeys. *Current Biology*, *23*(21), 2162-2168. https://doi.org/10.1016/j.cub.2013.09.005

Tardif, S. D., Mansfield, K. G., Ratnam, R., Ross, C. N., & Ziegler, T. E. (2011). The Marmoset as a Model of Aging and Age-Related Diseases. *ILAR Journal*, *52*(1), 54-65. https://doi.org/10.1093/ilar.52.1.54

Titchener, R., Thiriau, C., Hüser, T., Scherberger, H., Fischer, J., & Keupp, S. (2023). Social
disappointment and partner presence affect long-tailed macaque refusal behaviour in an
'inequity aversion' experiment. *Royal Society Open Science*, *10*(3), 221225.
https://doi.org/doi:10.1098/rsos.221225

Toro, J. M., & Hoeschele, M. (2017). Generalizing prosodic patterns by a non-vocal learning
mammal. *Animal Cognition*, *20*(2), 179-185. https://doi.org/10.1007/s10071-016-1036-
8

Wich, S. A., Assink, P. R., Becher, F., & Sterck, E. H. M. (2002). Playbacks of Loud Calls to
Wild Thomas Langurs (Primates; Presbytis thomasi): The Effect of Familiarity.
Behaviour, *139*(1), 79-87. https://doi.org/10.1163/15685390252902292

Xie, B., Brask, J. B., Dabelsteen, T., & Briefer, E. F. (2024). Exploring the role of vocalizations
in regulating group dynamics. *Philosophical Transactions of the Royal Society B:*
Biological Sciences, *379*(1905), 20230183. https://doi.org/10.1098/rstb.2023.0183

Zürcher, Y., & Burkart, J. M. (2017). Evidence for Dialects in Three Captive Populations of
Common Marmosets (*Callithrix jacchus*). *International Journal of Primatology*, *38*(4),
780-793. https://doi.org/10.1007/s10764-017-9979-4

Zürcher, Y., Willems, E. P., & Burkart, J. M. (2019). Are dialects socially learned in marmoset
monkeys? Evidence from translocation experiments. *PLOS ONE*, *14*(10), e0222486.
https://doi.org/10.1371/journal.pone.0222486

Figures:

Fig. 1

Types of playback stimuli used in the study. (a) natural call composed of three phee call syllables in their natural order, (b) control sound composed of three pure tones of similar duration and frequency, (c) pitch contour natural order, composed of pitch contours of the three syllables of a phee call in the natural order, (d) pitch contour first syllable, composed of three times the pitch contour of the first syllable of a phee call.

Fig. 2

688
689 *Schematic of the experimental setup. The stimuli were played back from a speaker hidden behind an artificial plant. On top of*
690 *the loudspeaker the main camera was placed to record the focal individual's reactions. Over the experimental cage an*
691 *additional camera (camera 2) was placed.*

692
693 **Fig. 3**

694
695 *Proportion of time looking at the playback location per stimulus and experimental condition. (a) Natural call condition, (b)*
696 *Pitch contour condition. While in the Natural call condition significant differences between familiar and unfamiliar individual's*
697 *calls compared to the control sound can be observed, looking time proportions did not differ between stimuli in the Pitch*
698 *contour condition.*

699
700 **Fig. 4**

Density plots for the difference in proportion of looking between familiar and unfamiliar individuals and the total proportion of looking time. (a) shows the overall difference in the proportion of looking times between familiar and unfamiliar stimuli. The plot of the Natural call condition shows a more skewed distribution indicating a slight preference in looking at unfamiliar stimuli, while the plot in the Pitch contour condition is more uniformly distributed. (b), (c) and (d) show the proportion of time looking during the stimulus presentations over stimuli category (b), testing session (c) and set number (d) respectively. All three show higher densities around 0 in the Pitch contour condition, indicating less interest in the stimuli in the Pitch contour condition compared to the Natural call condition.

Fig. 5

Delta values comparing times looking at the playback location between familiar and unfamiliar stimuli per individual (familiar – unfamiliar). In blue (above 0) individuals that looked longer at familiar stimuli, in green (below 0) individuals that looked longer at unfamiliar stimuli. (a) Natural call condition, ordered by degree of interest from most distinctly interested in familiar stimuli to most distinctly interested in unfamiliar stimuli. (b) Pitch contour condition, in the same order as the Natural call condition. In the Pitch contour condition overall decreased differences between familiar and unfamiliar stimuli can be observed. Individuals furthermore don't show the same looking patterns as in the Natural call condition.

Fig. 6

Effects of (a) sex, (b) family group and (c) age on the mean proportion of looking time per stimulus (control, familiar, unfamiliar), both in Natural call and Pitch contour condition. No clear patterns could be found for either of the variables.