1 **Mapping long-term memory modulation with transcranial alternating current**
2 **stimulation**
3
4
5 Running title: Mapping long-term memory
6

7 Dima Chitic, BSc[1,2], Miles Wischnewski, PhD[1,2,*]
8
9 [1]Department of Psychology, University of Groningen, Groningen, The Netherlands
10 [2]Research School of Behavioral and Cognitive Neurosciences, University of Groningen,
11 Groningen, The Netherlands

13 **Funding**
14 This work was supported by the Dutch Research Council [406.XS.25.01.035].

15
16 **Author statement**
17 DC: Data curation, Formal analysis, Methodology, Writing - original draft. MW:
18 Conceptualization, Formal analysis, Funding acquisition, Methodology, Supervision,
19 Visualization, Writing - review & editing
20
21 **Conflicts of interest**
22 The authors report no conflict of interests.
23

*Corresponding author:
Miles Wischnewski
Grote Kruisstraat 2/1
9712TS, Groningen
m.wischnewski@rug.nl
+31 621262594

Abstract

Background: Long-term memory (LTM), which is the ability to retain information over a long period of time, is governed by neural oscillations in various parts of the cerebral cortex. To investigate the relationship between oscillations and memory performance, transcranial alternating current stimulation (tACS) has been used to modulate LTM. However, due to the variability in stimulation parameters and locations, no clear picture has arisen on where tACS needs to be applied to improve LTM.

Methods: We collected behavioral data from 20 studies (58 effect sizes) and computationally modeled the corresponding brain electric fields based on each study's unique tACS parameters. As a result, we generated a map of performance–electric field index (PEI) values using linear and quadratic fits.

Results: Theta-tACS showed no regions with a positive linear association with LTM. However, left lateral prefrontal and temporal areas exhibited negative associations, meaning that theta stimulation of that region led to decreased LTM performance. A quadratic analysis revealed an inverted-U relation in the right parietal cortex, suggesting improved memory at moderate, but not high, electric field strengths. Gamma-tACS showed positive linear associations in parieto-occipital cortex and negative ones in frontal regions. For gamma, we found no regions that expressed a quadratic relationship.

Conclusion: These findings indicate that LTM-relevant tACS effects are frequency- and site-specific and potentially nonlinear with respect to intensity. Integrating meta-analytic effect sizes with individualized field estimates provides a brain-wide framework for identifying targets of memory modulation and for guiding future fundamental and translational research.

Keywords: Transcranial Alternating Current Stimulation; Long-term Memory; Meta-analysis; Electric Field Modeling; Neural oscillations

Introduction

The ability to retain information from hours, days, weeks, or even months ago is one of the hallmarks of the brain[1]. Besides the importance of gaining a fundamental understanding of the underlying brain mechanisms, long-term memory (LTM) studies are critical for further expanding the treatment options for memory-related deficits, such as Alzheimer's disease or traumatic brain injury[2,3]. In rodent models, memory is most commonly studied in subcortical areas, including the hippocampus and the amygdala[4–6]. However, human studies tend to focus on various cortical regions involved in processing and retaining information, including prefrontal, parietal, and temporal structures [7–12]. Furthermore, within these regions, particular neural oscillations, such as theta (3-8 Hz) and gamma (30-200 Hz), have been highlighted through their association with encoding and retrieval of memory[13–15].

Previous studies have investigated the role of oscillations using neurophysiological recordings. Osipova et al[16] engaged their participants in a picture recognition task while magnetoencephalographic (MEG) data were recorded. They found higher theta and gamma power both during encoding when the picture was subsequently remembered (subsequent memory effect) and during retrieval, when distinguishing old pictures from new. Furthermore, intracranial recordings from epilepsy patients showed a similar pattern, with increases in theta and gamma power over several cortical regions for subsequent memory effects[17–19]. In contrast, Hanouneh et al[20] showed that theta power in prefrontal regions was negatively correlated with LTM retrieval. However, Scholz et al[21] showed that theta power is positively associated with memory encoding.

Given the correlational nature and variability in results of the studies using electrophysiological techniques, others have used neuromodulation methods to pinpoint regions that causally relate to memory performance[22–24]. One particular technique suited to study neural rhythms is transcranial alternating current stimulation (tACS) and various studies have used it to alter cortical oscillations during memory tasks[25–28]. By applying a sinusoidal low-intensity current to the scalp, tACS has been shown to entrain neural spiking, which can result in a modulation of endogenous oscillatory activity[29,30]. Besides these neurophysiological effects, various studies have shown that tACS is able to modulate human behavior and cognition[25,31,32]. The rationale is that the tACS-induced entrainment of neural oscillations is directly related to changes in behavioral performance. As an example, long-term face recognition was improved by applying theta tACS to the right posterior parietal region during encoding[33]. On the other hand, Manippa et al[26] found reduced LTM performance following gamma tACS over bilateral temporal areas.

In accordance with the breadth of results in MRI and EEG studies, tACS has been applied in LTM experiments at various cortical locations and frequencies. Given the variety in parameters and protocols, a classic meta-analytic approach is suboptimal to summarize the efficacy of tACS, nor to identify core functional regions of LTM. By integrating meta-analytic results with electric field modeling, this limitation can be overcome[34–37]. Using SimNIBS software tACS-induced electric fields can be estimated based on a particular stimulation location and intensity[38]. By correlating the electric field distributions of different studies with their corresponding behavioral effect sizes (on LTM in this case), a meta-analytic brain map is generated that displays where tACS leads to increased, decreased, or no change in task performance[35].

102 In the present study, we leverage the meta-modeling approach to summarize the effects of
103 tACS on LTM. Thereby, we identify regions where neural oscillations in a given frequency are
104 directly related to memory performance. Altogether, these results aim to complement previous
105 research on cortical oscillations in LTM and point out cortical regions of interest for LTM
106 research.
107
Methods
Sample
We collected effect sizes, expressed in Hedges' g from available literature. The data was
obtained from articles published in peer-reviewed journals, pre-prints, and published theses,
by searching through PubMed, Web of Science, ProQuest (Dissertations and Theses),
BioRxiv and PsycINFO databases. The search was conducted in general accordance with the
PRISMA guidelines[39]. First, articles were identified using the following search terms: ("tACS"
OR "transcranial alternating current stimulation") AND ("memory"). After removal of duplicates,
the texts underwent abstract and full-text screening. Inclusion of studies in our modeling is
based on the following criteria: I) Articles are written in English, II) Full-text is available to the
researchers, III) manuscripts allowed for extraction of effect size (based on mean and standard
deviation) from text, tables, or figures, IV) the study used a control condition (e.g., placebo) to
which active stimulation can be compared, V) the study was conducted on a healthy adult
population (>= 18 years old), VI) tACS applied online or offline, but not during sleep, VII) long-
term memory assessed explicitly. For the criteria, we rely on a broad definition of long-term
memory as information retrieved from the past that does not rely on active maintenance[2]. A
preliminary search suggested that 15-25 articles could be eligible, suggesting that enough
data can be collected to perform our meta-modeling analyses. In total, the final search yielded
20 articles and 58 unique effect sizes (note that one study typically has multiple outcome
measures). Across all studies n = 1210 healthy volunteers were included between the ages of
18-77. Since previously published data were used, no ethical approval was obtained.
129
Effect size calculation
The standardized effect sizes were expressed as Hedges' g values[40]. Hedges' g values
express the ratio between differences in effects and the variability, similar to Cohen's d
values[41], with the addition of a correction factor for small sample sizes. If Hedges' g values
were reported, they were directly taken from the article. Alternatively, if the article provided the
Cohen's d value, the correction factor was applied to calculate Hedges' g. If no effect size
measures were directly reported, the Hedges' g was calculated based on reported averages
and standard deviations or those estimated from graphs. All effect sizes used in this analysis
reflect the difference between an active stimulation condition and a sham control. As such,
positive g values reflect improved memory performance in active tACS conditions compared
to sham, while negative g values relate to worsened memory performance in active tACS
conditions compared to sham.
142
Electric field modeling
Electric field simulations were performed on 50 realistic head models (age range 22-35, 22
females). The head models were generated based on data from the Human Connectome
Project S1200[42]. Quality and suitability for electric field modeling were assessed previously[43].
Each head model consists of 6 biological tissues, namely the skin, skull, cerebrospinal fluid,
eyes, gray matter and white matter. For each tissue standard conductivity values were used
($\sigma_{skull} = 0.01$ S/m, $\sigma_{skin} = 0.465$ S/m, $\sigma_{CSF} = 1.654$ S/m, $\sigma_{graymatter} = 0.275$ S/m, and

σwhitematter = 0.126 S/m)[44,45]. Electrode location, orientation, and material were gathered
from the articles and used for electric field modeling in SimNIBS 4.1[38]. For the purpose of this
study, we extracted the induced electric fields from the gray matter, as this reflects the tissue
at which tACS most likely induces functional changes[46,47]. In order to compare and average
between head models, all electric field simulations were transformed into the normalized
FreeSurver average (fsavg) standard brain. Importantly, by first performing simulations in the
individual head models and then transforming them into fsavg space (rather than directly
performing electric-field simulations in fsavg models), individual variability is still accounted
for.
159
Calculation of the performance-electric field index
To relate different elements of fsavg model to functional brain regions, each element was
categorized following the Human Connectome Project multimodal parcellation (HCP-MP)
atlas[48,49]. This atlas parcellates the brain models into 360 function regions (180 per
hemisphere). Given that we found 36 and 22 outcome measures for theta and gamma,
respectively, and 360 functional regions, a 36 x 360 matrix and a 22 x 360 matrix of electric
field values are obtained. This matrix is subsequently correlated with a vector of Hedges' g
values for each row of the matrix (i.e., for each functional region), resulting in a 1 x 360 matrix
that represents the correlation between electric fields and tACS-induced performance
changes[34–37]. We refer to these values as the performance-electric field index (PEI). Besides
this we report the uncorrected p-values. Note that due to the nature of meta-analytic datasets,
no a priori power analysis was performed, and the number of data points depends on the
available literature. As such, the exact p-value, and hence the optimal way to correct for
multiple comparisons, is somewhat arbitrary. Therefore, we opted to report uncorrected p-
values and visually represent them as $p<0.05$ (*), $p<0.01$ (**), and $p<0.001$ (***). For
visualization, we categorized the 360 HCP-MP atlas region into 22 larger regions[50].
176
In addition to the linear correlation analysis, we performed an exploratory analysis testing for
a potential quadratic relationship between electric field strength and performance. This
analysis was performed as several studies have shown that the effects of tACS follow a U-
shape[51–53] or inverted U-shape [54,55]. For each cortical region, we fit a linear ($y = \beta_0 + \beta_1 x$) and
quadratic regression model ($y = \beta_0 + \beta_1 x + \beta_2 x^2$), with the behavioral effect size (Hedges' g)
as the dependent variable and regional E-field magnitude as the predictor. For both linear and
quadratic models, a correlation value was defined as the signed square root of R-squared,
with the sign determined by the regression values being positive or negative. For each region,
we determined the better-fitting model (linear or quadratic) based on adjusted R-squared.
Across 50 head models, results were averaged per region. If more than 50% of the head
models yielded a quadratic fit for a particular region, the quadratic correlation values were
averaged. We refer to this value as the PEI_QUAD, where positive and negative values
correspond to a U-shaped and an inverted U-shaped relationship between electric field
strength and performance, respectively.
191

Table 1. Overview of studies using theta tACS included in the meta-analysis.

Reference	Montage	Freq. (Hz)	Intensity (mA)	Task	Time between encoding and retrieval	Hedges' g
Alekseichuk et al., 2020	P4, T8/C2/CP1/Oz	4	3	face recognition	20 min	0.31
	P3, T7/C1/CP2/Oz	4	3	face recognition	20 min	0.06
Chantrel et al., 2021	F3, left wrist	6	1.5	word recognition	90 min	-0.32
	F3, left wrist	6	1.5	word recognition	7 days	-0.15
Ergo et al., 2020	FCz, neck	6	2	word association task	Immediate	-0.21
Grover et al., 2022	FP1/F1/FC3/F5/AF7/Afz, Fz/AF3/cheek	4	1.58	delayed recall, words	Immediate	0.01
	C3/T7/CP1/FC5/TP7FT7/PO7/cheek, CP5	4	1.58	delayed recall, words	Immediate	0.06
	C3/T7/CP1/FC5/TP7FT7/PO7/cheek, CP5	4	1.58	delayed recall, words	1 month	0.17
Klink et al., 2020	Intersection of T3/F3/C3/F7, rSOR	5	1	face-occupation task	20 min	-0.14
	Intersection of T3/F3/C3/F7, rSOR	5	1	face-occupation task	1 day	0.03
Komissarenko et al., 2025	F7, P7	5.5	1	word association recognition	1 min	-0.04
	F7, P7	5.5	1	word association recognition	4 days	-0.14
Lang et al., 2019	right fusiform 4x1	6	4	face-scene association	online	0.56
Meng et al., 2021	P3, ring surround	6	2	face-scene task, pictures	5 min	-0.42
Murray et al., 2023	T3, T4	6	1.5	delayed recall, words	20 sec	-0.47
Shtoots et al., 2024	Chin, P1/P2/P3/P4	3	1.5	delayed recall, pictures	2 hours	0.29
	Chin, P1/P2/P3/P4	3	1.5	delayed recall, pictures	1 day	0.39
	Chin, P1/P2/P3/P4	3	1.5	delayed recall, pictures	7 day	0.4
	Fz, AF4/AF3/FC2/FC1	3	1.5	delayed recall, pictures	2 hours	1.11
	Fz, AF4/AF3/FC2/FC1	3	1.5	delayed recall, pictures	1 day	1.27
	Fz, AF4/AF3/FC2/FC1	3	1.5	delayed recall, pictures	7 day	1.26
Shtoots et al., 2025	Chin, P1/P2/P3/P4	3	1.5	multiple choice exam, video	20 min	1.05
	Chin, P1/P2/P3/P4	3	1.5	multiple choice exam, video	1 day	0.73
	Chin, P1/P2/P3/P4	3	1.5	multiple choice exam, video	7 day	0.65
	P3, C3/TP7/Pz/O1	3	1.5	multiple choice exam, video	20 min	1.24
	P3, C3/TP7/Pz/O1	3	1.5	multiple choice exam, video	1 day	1.16
	P3, C3/TP7/Pz/O1	3	1.5	multiple choice exam, video	7 day	1.01
	Fz, AF4/AF3/FC2/FC1	3	1.5	multiple choice exam, video	20 min	0.67

	Fz, AF4/AF3/FC2/FC1	3	1.5	multiple choice exam, video	1 day	0.83
	Fz, AF4/AF3/FC2/FC1	3	1.5	multiple choice exam, video	7 day	0.91
Varastegan et al., 2022	FPz, Oz	6	2	delayed recall, words	20 min	1.06
Wynn et al., 2024	AF4/P5, Cz	4	2	word recognition	15 min	0.09
Wynn et al., 2020	P7/P8, Cz	3.5	2	word recognition	7 min	0.03
	P7/P8, Cz	3.5	2	screen location recognition	7 min	0.00
	P7/P8, Cz	8	2	word recognition	7 min	0.02
	P7/P8, Cz	8	2	screen location recognition	7 min	0.00

Table 2. Overview of studies using gamma tACS included in the meta-analysis.

Reference	Montage	Freq. (Hz)	Intensity (mA)	Task	Time between encoding and retrieval	Hedges' g
Ambrus et al., 2015	F3/F4, left/right mastoid	140	1	cued recall, words	10 min	-0.20
	F3/F4, left/right mastoid	140	1	cued recall, words	12 hours	0.00
Chantrel et al., 2021	F3, left wrist	60	1.5	word recognition	90 min	-0.21
	F3, left wrist	60	1.5	word recognition	7 days	-0.49
	C4, left wrist	60	1.5	word recognition	7 days	-0.12
Grover et al., 2022	FP1/F1/FC3/F5/AF7/Afz, Fz/AF3/cheek	60	1.58	delayed recall, words	Immediate	-0.01
	FP1/F1/FC3/F5/AF7/Afz, Fz/AF3/cheek	60	1.58	delayed recall, words	1 month	0.17
	C3/T7/CP1/FC5/TP7FT7/PO7/cheek, CP5	60	1.58	delayed recall, words	Immediate	-0.02
Javadi et al., 2017	F3, left wrist	60	1.5	word recognition	1.5 hours	0.38
	F3, left wrist	60	1.5	word recognition	1.5 hours	0.46
	F3, left wrist	60	1.5	word recognition	1.5 hours	0.02
	F3, left wrist	60	1.5	word recognition	1.5 hours	0.38
Luckey et al., 2022	Bilateral C2 dermatome	40	3	word association task	7 days	1.05
Manippa et al., 2024	T7, T8	40	2	delayed recall, words	4 min	0.08
	T7, T8	60	2	delayed recall, words	4 min	-0.13
Nomura et al., 2019	F3, left arm	60	1.5	word recognition	Immediate	0.21
	F3, left arm	60	1.5	word recognition	1 day	0.74
	F3, left arm	60	1.5	word recognition	6 days	1.17
Sun et al., 2025	P3, right deltoid	60	1.5	cued recall, words	4 min	0.35

	P3, right deltoid	60	1.5	cued recall, words	7 days	0.81
	P3, right deltoid	60	1.5	cued recall, words	28 days	1.09
Wynn et al., 2024	AF4/P5, Cz	50	2	word recognition	15 min	0.10

Figure 1. *A) Electric field strength averaged over all montages used in the selected articles, split for theta and gamma stimulation. Warm colors, where electric fields were strong on average, reflect regions commonly targeted across studies, while cold colors, where electric fields were weak on average, indicate regions less commonly targeted. B) A sunburst plot providing a global reflection of montages used across studies. FP: frontoparietal, OP: occipitoparietal, TO: Temporo-occipital. C) Raincloud distribution of effect sizes for theta and gamma tACS studies, showing an overall cumulative effect. Box plot limits correspond to the 25th and 75th percentiles, with the center line representing the median, and whiskers representing 1.5x interquartile range.*

Results

In total 20 articles were identified for inclusion in the present meta-modeling analysis[25–28,33,56–70]. The analysis was split for studies that targeted theta oscillations (3-8 Hz) and gamma oscillations (30-200 Hz). For theta tACS studies, N=36, and for gamma tACS studies, N=22 effect sizes were obtained (Figure 1C). In three papers, tACS frequencies other than theta and gamma were studied, which were not included in our analysis (Supplementary Table 1). Since studies used various montages, we first aimed to get a sense of the general distribution of electric fields (Figure 1B). As such, we averaged electric fields across all studies (Figure 1A). For both frequencies, the simulations indicated widespread distribution across the brain, which is in line with electrodes being placed on frontal, temporal and occipito-parietal regions. However, for theta the averaged electric field was stronger at the left parietal cortex, whereas for gamma it was stronger for left frontal, temporal and parietal regions. This reflects a bias in the literature of preferentially targeting the left hemisphere.

219
220 *Figure 2. Distribution of PEI values across the brain related to theta tACS. Exact PEI values for all 360*
221 *regions, categorized into 22 larger regions (Huang et al., 2022), are reflected on the left and right.*
222

Theta tACS on LTM

For the theta tACS we found no regions that were related to improvement of LTM, with a
maximum of PEI = 0.052 (p = 0.761; right area 8B). In contrast, left frontal and temporal
regions showed a negative linear association between electric field strength and LTM
performance (p<0.001: N=2; 0.001<p<0.01: N=34; 0.01<p<0.05: N=39). Results are shown in
Figure 2 and Table 1. Strongest associations were found in region 6r (PEI = -0.545, p = 5.7e-
4) and 44 (PEI = -0.526, p = 9.6e-4) of the HCP-MMP atlas.
230

Figure 3. *Distribution of PEI values across the brain related to gamma tACS. Exact PEI values for all 360 regions, categorized into 22 larger regions (Huang et al., 2022), are reflected on the left and right.*

Gamma tACS on LTM

For gamma tACS there was tentative evidence for bilateral occipital and parietal regions being related to improved LTM (p<0.05: N=24). The regions with the strongest associations were V1 (PEI = 0.514, p = 0.014) and V3A (PEI = 0.504, p = 0.016). No regions were found with a significant negative association between electric field magnitude and LTM performance, with a minimum of PEI = -0.232 (p = 0.297, left area anterior 9-46v). Results are shown in Figure 3 and Table 2.

244
245 ***Figure 4.*** *Distribution of PEI*_{QUAD} *values across the brain related to theta tACS, reflecting a quadratic*
246 *relationship between electric field strength and behavioral performance.*
247
Quadratic associations between electric field magnitude and LTM
In an exploratory analysis we investigated whether regions show a quadratic relationship
between electric field and effect size (Figure 4). For theta tACS, we found that a quadratic
relationship explained the variance better in 92 out of the 360 regions (i.e., a linear relationship
explained more variance in the remaining 268 regions). This quadratic effect was observed in
right occipito-parietal region. These 92 regions corresponded to a significant negative
quadratic PEI, ranging between PEI$_{QUAD}$ = -0.353 (p = 0.035) and PEI$_{QUAD}$ = -0.468 (p = 0.007).
This negative relationship indicates that mid-level electric field strengths had a positive effect
on LTM performance, whereas low and high electric field strengths did not. No significant
positive quadratic PEI values were found. For gamma stimulation, no regions were identified
where a quadratic fit was superior to a linear fit.
259
Discussion
We aimed to summarize published LTM-tACS studies to date by correlating tACS-induced
electrical fields in brain models with corresponding changes in LTM performance. As a result,
we found that within the theta frequency range, no regions showed a significant linear
relationship that improved memory performance, but modulation of the left frontal and
temporal lobes led to worse performance across studies. Exploring quadratic relationships,
we found that theta over the right parietal cortex improves memory at moderate (but not high
and low) electric field strengths. For gamma frequencies, parieto-occipital areas were
associated with better performance, whereas frontal areas were associated with worse
performance.
270
In contrast to our expectations, no clear linear positive relationship was found between theta
stimulation and LTM performance. Only the dorsomedial prefrontal cortex showed a positive
association. And while theta in the medial PFC has been linked to LTM in animal studies[12,71–]
[73], our results provide little support for this in humans. In contrast, and against our
expectations, we found a negative association between LTM performance and electric field
strength in left lateral prefrontal and temporal areas. The lateral prefrontal cortex is often
associated with general cognitive performance and control, and various studies have at least
suggested a potential role in LTM[25,74–77]. The temporal lobe includes and is connected to the
hippocampus, which is often referred to as the main hub for LTM[78–81]. Within the hippocampus
and medial temporal lobe, theta is a dominant oscillation[19,82–84]. Therefore, we hypothesized a

positive PEI for this region. One possible explanation is that the exogenous stimulation in the
collected studies did not entrain but rather disrupted endogenous theta frequency or phase
coupling. While tACS is generally thought to entrain regions and thereby strengthen
endogenous oscillations[22], various studies have emphasized that this may depend on specific
circumstances[32]. For instance, several studies investigate in-phase vs anti-phase tACS,
suggesting that in-phase stimulation is more likely to increase synchronization between
regions[85–89]. Phase of the endogenous oscillations also matters. For instance, by using an
amplitude-modulated tACS design and sophisticated artifact cleaning methods, Haslacher et
al[90] were able to record EEG while tACS is applied in a working memory task. They found that
working memory depended on the phase lag between exogenous and endogenous theta
oscillations. While this study looked at a different memory modality, it demonstrates the
importance of phase alignment when aiming to modulate cognitive performance. Besides
phase, recent studies have highlighted the importance of tACS intensity[52,53]. Specifically, lower
intensities are more likely to decrease naturally occurring regional synchronization, rather than
entrain a region. Furthermore, entrainment effects follow an Arnold's tongue pattern, with the
strongest entrainment occurring for stimulation frequencies that are close to endogenous
rhythms[91,92]. This emphasizes the importance of personalized stimulation frequencies, which
many of the here included studies did not apply[47].
299
While we did not find any linear relationship between theta and LTM performance, when
performing an additional analysis investigating a quadratic relationship, we found that this
better explained the variance in 92 regions. Specifically, we found a significant negative
quadratic effect in right parietal regions. In this case, a negative quadratic association reflects
that the relation between electric field strength and LTM performance follows an inverted U
pattern. This suggests that parietal theta stimulation with moderate, but not high, electric field
strength was related to improved LTM performance. Indeed, various studies have
demonstrated a nonlinear relationship between tACS intensity and neurophysiological and
behavioral changes[51,54,55]. Wansbrough et al[55] demonstrated frequency-specific increase in
EEG coherence after 0.5 mA tACS, while 1 or 1.5 mA tACS decreased coherence. On a
behavioral level, Hsu et al[54] showed that 1 mA, but not 2 mA theta-tACS improve cognitive
performance in patients with multiple sclerosis. Such inverted U relations have also been
observed for other brain stimulation applications[93]. For example, Jamil et al[94] showed stronger
after effects of motor cortical transcranial direct current stimulation after 1 mA, compared to
sham and 2 mA stimulation. Indeed, various studies have proposed the idea of the brain as a
homeostatic system, where modulation effects depend on the initial state of the brain[95–98].
316
For gamma tACS, we found that parieto-occipital regions were most strongly related to
improvements in LTM. Indeed, studies have suggested that visual areas may serve as a
storage or reactivation site for visual memory representations. For example, imaging work
shows that visual memory traces may be maintained in the visual cortex[99–102]. Furthermore,
gamma oscillations in particular have been related to sensory memory information[16,102,103]. On
the other hand, gamma oscillations in the parieto-occipital cortex may reflect domain-general
attentional mechanisms that support robust encoding and consolidation of memory by
enhancing the binding of information and coordinating large-scale networks[104–110]. Previously,
we found that gamma tACS of similar posterior regions enhances working memory
performance[35], suggesting that parieto-occipital gamma is not exclusively related to LTM.
Rather, it may reflect a broader mechanism supporting cognitive operations across memory
timescales[14]. In sensory cortices, gamma oscillations have been linked to attentional selection

and feature binding, coordinating local neuronal ensembles to integrate information into
coherent representations[105,111,112]. Thus, gamma tACS over parieto-occipital areas may
facilitate long-term memory formation indirectly, by enhancing attentional and integrative
processes in visual regions that support effective encoding and consolidation. Whether
gamma tACS over the parietal cortex improves the sensory-mnemonic trace itself or improves
the attentional processes cannot be deduced from our results. Nevertheless, our results
suggest that stimulation of these regions most likely improves LTM performance.
336
Some considerations should be made when interpreting the results of the present study. First,
the included studies used a variety of tasks, stimuli, and sensory modalities. As such, we
cannot make any conclusions about specific memory domains. Rather, the PEI maps could
reflect improvement in domain-general LTM. Furthermore, while the results were split by
frequency, we pooled across different retrieval processes (recognition or recall), task
modalities (e.g., words, faces, or images), and stimulation timings (stimulation during
encoding, during retrieval, during both or offline). While it would be potentially interesting to
split by these factors, the limited sample of 20 studies does not allow for reliable subfactor
analyses. As such, more studies are required to explore the role of theta and gamma tACS in
different methodological approaches. Second, given the reliance on existing data, not all
potential montages have been explored. The PEI estimates rely on variability in electric fields
and capture effects within these commonly stimulated areas, while regions rarely or never
targeted should be interpreted with caution. The regions affected by the collected montages
were rather evenly distributed across the brain. Nonetheless, there was some bias towards
the temporoparietal areas and toward left hemispheric stimulation. Although such spatial bias
could influence the results, the lack of overlap between the average electric field and PEI maps
hints that any such effects were likely limited. Finally, it is important to note that the results
identify regions of interest based on the LTM-tACS studies conducted to date, not an exclusive
map of regions essential for memory per se. Various regions might be crucial for LTM
processing that are not susceptible to tACS-related improvements. Relatedly, since tACS
targets primarily cortical regions, we can draw no conclusion about the causal influence of
subcortical oscillations on LTM, for instance, those originating from hippocampal regions.
359
In conclusion, we illustrate the potential of noninvasive brain stimulation to modulate LTM and
shed light on the causal role of theta and gamma oscillations. As such, we find that tACS-
related improvement to LTM is location and frequency-specific. Additionally, tACS-induced
cognitive effects may not always scale linearly with intensity. Importantly, more studies are
required to explore specific aspects of LTM, including different modalities and various methods
of probing LTM. Nevertheless, we hope that these results provide a guide for future
fundamental, translational, and therapeutic studies aimed at modulating memory.
367

368 **Conflicts of interest**
369 The authors report no conflict of interests.
370
371 **Funding**
372 This work was supported by the XS grant of the Dutch Research Council (NWO;
373 406.XS.25.01.035).
374

375 **Author contributions**

376   DC: Data curation, Formal analysis, Methodology, Writing - original draft. MW:
377   Conceptualization, Formal analysis, Funding acquisition, Methodology, Supervision,
378   Visualization, Writing - review & editing
379
References

1.   Squire LR, Wixted JT. The Cognitive Neuroscience of Human Memory Since H.M.
Annual Review of Neuroscience. 2011;34(Volume 34, 2011):259-288.
doi:10.1146/annurev-neuro-061010-113720

2.   Jeneson A, Squire LR. Working memory, long-term memory, and medial temporal lobe
function. *Learn Mem*. 2012;19(1):15-25. doi:10.1101/lm.024018.111

3.   Pittenger C. Disorders of memory and plasticity in psychiatric disease. *Dialogues in
Clinical Neuroscience*. 2013;15(4):455-463. doi:10.31887/DCNS.2013.15.4/cpittenger

4.   Yang S, Yi YG, Chang MC. The effect of transcranial alternating current stimulation on
functional recovery in patients with stroke: a narrative review. *Front Neurol*.
2023;14:1327383. doi:10.3389/fneur.2023.1327383

5.   Daume J, Kamiński J, Schjetnan AGP, et al. Control of working memory by phase–
amplitude coupling of human hippocampal neurons. *Nature*. 2024;629(8011):393-401.
doi:10.1038/s41586-024-07309-z

6.   Sawangjit A, Oyanedel CN, Niethard N, Salazar C, Born J, Inostroza M. The
hippocampus is crucial for forming non-hippocampal long-term memory during sleep.
Nature. 2018;564(7734):109-113. doi:10.1038/s41586-018-0716-8

7.   Jetter W, Poser U, Freeman RB, Markowitsch HJ. A Verbal Long Term Memory Deficit
in Frontal Lobe Damaged Patients. *Cortex*. 1986;22(2):229-242. doi:10.1016/S0010-
9452(86)80047-8

8.   Kesner RP. The posterior parietal cortex and long-term memory representation of
spatial information. *Neurobiology of Learning and Memory*. 2009;91(2):197-206.
doi:10.1016/j.nlm.2008.09.004

9.   Olson IR, Berryhill M. Some surprising findings on the involvement of the parietal lobe
in human memory. *Neurobiology of Learning and Memory*. 2009;91(2):155-165.
doi:10.1016/j.nlm.2008.09.006

10. Squire LR. Memory for relations in the short term and the long term after medial
temporal lobe damage. *Hippocampus*. 2017;27(5):608-612. doi:10.1002/hipo.22716

11. Miller JA, Tambini A, Kiyonaga A, D'Esposito M. Long-term learning transforms
prefrontal cortex representations during working memory. *Neuron*. 2022;110(22):3805-
3819.e6. doi:10.1016/j.neuron.2022.09.019

12. Sheynikhovich D, Otani S, Bai J, Arleo A. Long-term memory, synaptic plasticity and
dopamine in rodent medial prefrontal cortex: Role in executive functions. *Front Behav
Neurosci*. 2023;16. doi:10.3389/fnbeh.2022.1068271

13. Axmacher N, Mormann F, Fernández G, Elger CE, Fell J. Memory formation by
neuronal synchronization. *Brain Res Rev*. 2006;52(1):170-182.
doi:10.1016/j.brainresrev.2006.01.007

14.   Nyhus E, Curran T. Functional Role of Gamma and Theta Oscillations in Episodic
Memory. *Neurosci Biobehav Rev*. 2010;34(7):1023-1035.
doi:10.1016/j.neubiorev.2009.12.014

15.   Köster M. The theta-gamma code in predictive processing and mnemonic updating.
Neuroscience & Biobehavioral Reviews. 2024;158:105529.
doi:10.1016/j.neubiorev.2023.105529

16.   Osipova D, Takashima A, Oostenveld R, Fernández G, Maris E, Jensen O. Theta and
Gamma Oscillations Predict Encoding and Retrieval of Declarative Memory. *J
Neurosci*. 2006;26(28):7523-7531. doi:10.1523/JNEUROSCI.1948-06.2006

17.   Sederberg PB, Kahana MJ, Howard MW, Donner EJ, Madsen JR. Theta and Gamma
Oscillations during Encoding Predict Subsequent Recall. *J Neurosci*.
2003;23(34):10809-10814. doi:10.1523/JNEUROSCI.23-34-10809.2003

18.   Sederberg PB, Schulze-Bonhage A, Madsen JR, et al. Hippocampal and Neocortical
Gamma Oscillations Predict Memory Formation in Humans. *Cereb Cortex*.
2007;17(5):1190-1196. doi:10.1093/cercor/bhl030

19.   Vivekananda U, Bush D, Bisby JA, et al. Theta power and theta-gamma coupling
support long-term spatial memory retrieval. *Hippocampus*. 2021;31(2):213-220.
doi:10.1002/hipo.23284

20.   Hanouneh S, Amin HU, Saad NM, Malik AS. EEG Power and Functional Connectivity
Correlates with Semantic Long-Term Memory Retrieval. *IEEE Access*. 2018;6:8695-
8703. doi:10.1109/ACCESS.2017.2788859

21.   Scholz S, Schneider SL, Rose M. Differential effects of ongoing EEG beta and theta
power on memory formation. *PLOS ONE*. 2017;12(2):e0171913.
doi:10.1371/journal.pone.0171913

22.   Polanía R, Nitsche MA, Ruff CC. Studying and modifying brain function with non-
invasive brain stimulation. *Nat Neurosci*. 2018;21(2):174-187. doi:10.1038/s41593-017-
0054-4

23.   Elyamany O, Leicht G, Herrmann CS, Mulert C. Transcranial alternating current
stimulation (tACS): from basic mechanisms towards first applications in psychiatry. *Eur
Arch Psychiatry Clin Neurosci*. 2021;271(1):135-156. doi:10.1007/s00406-020-01209-9

24.   Agboada D, Zhao Z, Wischnewski M. Neuroplastic effects of transcranial alternating
current stimulation (tACS): from mechanisms to clinical trials. *Front Hum Neurosci*.
2025;19. doi:10.3389/fnhum.2025.1548478

25.   Grover S, Wen W, Viswanathan V, Gill CT, Reinhart RMG. Long-lasting, dissociable
improvements in working memory and long-term memory in older adults with repetitive
neuromodulation. *Nat Neurosci*. 2022;25:1237-1246. doi:10.1038/s41593-022-01132-3

26.   Manippa V, Nitsche MA, Filardi M, et al. Temporal gamma tACS and auditory
stimulation affect verbal memory in healthy adults. *Psychophysiology*.
2024;61(11):e14653. doi:10.1111/psyp.14653

27.   Shtoots L, Nadler A, Partouche R, et al. Frontal midline theta transcranial alternating
current stimulation enhances early consolidation of episodic memory. *npj Sci Learn*.
2024;9(1):8. doi:10.1038/s41539-024-00222-0

459 28. Wynn SC, Townsend CD, Nyhus E. The role of theta and gamma oscillations in item
460   memory, source memory, and memory confidence. *Psychophysiology*.
461   2024;61(9):e14602. doi:10.1111/psyp.14602

462 29. Krause MR, Vieira PG, Csorba BA, Pilly PK, Pack CC. Transcranial alternating current
463   stimulation entrains single-neuron activity in the primate brain. *Proc Natl Acad Sci U S*
464   *A*. 2019;116(12):5747-5755. doi:10.1073/pnas.1815958116

465 30. Wischnewski M, Tran H, Zhao Z, et al. Induced neural phase precession through
466   exogenous electric fields. *Nat Commun*. 2024;15(1):1687. doi:10.1038/s41467-024-
467   45898-5

468 31. Schutter DJLG, Wischnewski M. A meta-analytic study of exogenous oscillatory electric
469   potentials in neuroenhancement. *Neuropsychologia*. 2016;86:110-118.
470   doi:10.1016/j.neuropsychologia.2016.04.011

471 32. Wischnewski M, Alekseichuk I, Opitz A. Neurocognitive, physiological, and biophysical
472   effects of transcranial alternating current stimulation. *Trends in Cognitive Sciences*.
473   2023;27(2):189-205. doi:10.1016/j.tics.2022.11.013

474 33. Alekseichuk I, Turi Z, Veit S, Paulus W. Model-driven neuromodulation of the right
475   posterior region promotes encoding of long-term memories. *Brain Stimulation*.
476   2020;13(2):474-483. doi:10.1016/j.brs.2019.12.019

477 34. Wischnewski M, Mantell KE, Opitz A. Identifying regions in prefrontal cortex related to
478   working memory improvement: A novel meta-analytic method using electric field
479   modeling. *Neuroscience & Biobehavioral Reviews*. 2021;130:147-161.

480 35. Wischnewski M, Berger TA, Opitz A, Alekseichuk I. Causal functional maps of brain
481   rhythms in working memory. *Proc Natl Acad Sci USA*. 2024;121(14):e2318528121.
482   doi:10.1073/pnas.2318528121

483 36. Wischnewski M, Berger TA, Opitz A. Meta-modeling the effects of anodal left prefrontal
484   transcranial direct current stimulation on working memory performance. *Imaging*
485   *Neuroscience*. 2024;2:1-14. doi:10.1162/imag_a_00078

486 37. Yachou Y, Bouaziz N, Makdah G, et al. Transcranial direct current stimulation in
487   patients with depression: An electric field modeling meta-analysis. *Journal of Affective*
488   *Disorders*. 2025;374:540-552. doi:10.1016/j.jad.2025.01.001

489 38. Thielscher A, Antunes A, Saturnino GB. Field modeling for transcranial magnetic
490   stimulation: A useful tool to understand the physiological effects of TMS? In: *2015 37th*
491   *Annual International Conference of the IEEE Engineering in Medicine and Biology*
492   *Society (EMBC)*. 2015:222-225. doi:10.1109/EMBC.2015.7318340

493 39. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic
494   reviews and meta-analyses: the PRISMA statement. *BMJ*. 2009;339:b2535.
495   doi:10.1136/bmj.b2535

496 40. Hedges LV, Olkin I. *Statistical Methods for Meta-Analysis*. Academic Press; 1985.

497 41. Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Academic Press;
498   1977.

42.   Van Essen DC, Smith SM, Barch DM, et al. The WU-Minn human connectome project:
an overview. *Neuroimage*. 2013;80:62-79. Accessed October 24, 2025.
https://www.sciencedirect.com/science/article/pii/S1053811913005351

43.   Berger TA, Wischnewski M, Opitz A, Alekseichuk I. Human head models and
populational framework for simulating brain stimulations. *Sci Data*. 2025;12(1):516.
doi:10.1038/s41597-025-04886-0

44.   Windhoff M, Opitz A, Thielscher A. Electric field calculations in brain stimulation based
on finite elements: an optimized processing pipeline for the generation and usage of
accurate individual head models. *Hum Brain Mapp*. 2013;34(4):923-935.
doi:10.1002/hbm.21479

45.   McCann H, Pisano G, Beltrachini L. Variation in reported human head tissue electrical
conductivity values. *Brain Topogr*. 2019;32(5):825-858. doi:10.1007/s10548-019-
00710-2

46.   Katoch N, Kim Y, Choi BK, et al. Estimation of brain tissue response by electrical
stimulation in a subject-specific model implemented by conductivity tensor imaging.
Front Neurosci. 2023;17. doi:10.3389/fnins.2023.1197452

47.   Hoornweder SV, Stagg CJ, Wischnewski M. Personalizing transcranial electrical
stimulation. *Trends in Neurosciences*. 2025;0(0). doi:10.1016/j.tins.2025.07.007

48.   Glasser MF, Coalson TS, Robinson EC, et al. A multi-modal parcellation of human
cerebral cortex. *Nature*. 2016;536(7615):171-178. doi:10.1038/nature18933

49.   Glasser MF, Smith SM, Marcus DS, et al. The Human Connectome Project's
neuroimaging approach. *Nat Neurosci*. 2016;19(9):1175-1187. doi:10.1038/nn.4361

50.   Huang CC, Rolls ET, Feng J, Lin CP. An extended Human Connectome Project
multimodal parcellation atlas of the human cortex and subcortical areas. *Brain Struct
Funct*. 2022;227(3):763-778. doi:10.1007/s00429-021-02421-6

51.   Moliadze V, Atalay D, Antal A, Paulus W. Close to threshold transcranial electrical
stimulation preferentially activates inhibitory networks before switching to excitation with
higher intensities. *Brain Stimulation*. 2012;5(4):505-511. doi:10.1016/j.brs.2011.11.004

52.   Krause MR, Vieira PG, Thivierge JP, Pack CC. Brain stimulation competes with
ongoing oscillations for control of spike timing in the primate brain. *PLOS Biology*.
2022;20(5):e3001650. doi:10.1371/journal.pbio.3001650

53.   Zhao Z, Shirinpour S, Tran H, Wischnewski M, Opitz A. Intensity- and frequency-
specific effects of transcranial alternating current stimulation are explained by network
dynamics. *J Neural Eng*. 2024;21(2):026024. doi:10.1088/1741-2552/ad37d9

54.   Hsu WY, Zanto T, Park JE, Gazzaley A, Bove RM. Effects of transcranial alternating
current stimulation on cognitive function in people with multiple sclerosis: A randomized
controlled trial. *Mult Scler Relat Disord*. 2023;80:105090.
doi:10.1016/j.msard.2023.105090

55.   Wansbrough K, Marinovic W, Fujiyama H, Vallence AM. Beta tACS of varying
intensities differentially affect resting-state and movement-related M1-M1 connectivity.
Front Neurosci. 2024;18. doi:10.3389/fnins.2024.1425527

56. Ambrus GG, Pisoni A, Primaßin A, Turi Z, Paulus W, Antal A. Bi-frontal transcranial
alternating current stimulation in the ripple range reduced overnight forgetting. *Front*
Cell Neurosci. 2015;9. doi:10.3389/fncel.2015.00374

57. Javadi AH, Glen JC, Halkiopoulos S, Schulz M, Spiers HJ. Oscillatory Reinstatement
Enhances Declarative Memory. *J Neurosci*. 2017;37(41):9939-9944.
doi:10.1523/JNEUROSCI.0265-17.2017

58. Lang S, Gan LS, Alrazi T, Monchi O. Theta band high definition transcranial alternating
current stimulation, but not transcranial direct current stimulation, improves associative
memory performance. *Sci Rep*. 2019;9(1):8562. doi:10.1038/s41598-019-44680-8

59. Nomura T, Asao A, Kumasaka A. Transcranial alternating current stimulation over the
prefrontal cortex enhances episodic memory recognition. *Exp Brain Res*.
2019;237(7):1709-1715. doi:10.1007/s00221-019-05543-w

60. Ergo K, Loof ED, Debra G, Pastötter B, Verguts T. Failure to modulate reward
prediction errors in declarative learning with theta (6 Hz) frequency transcranial
alternating current stimulation. *PLOS ONE*. 2020;15(12):e0237829.
doi:10.1371/journal.pone.0237829

61. Klink K, Peter J, Wyss P, Klöppel S. Transcranial Electric Current Stimulation During
Associative Memory Encoding: Comparing tACS and tDCS Effects in Healthy Aging.
Front Aging Neurosci. 2020;12. doi:10.3389/fnagi.2020.00066

62. Wynn SC, Kessels RPC, Schutter DJLG. Effects of parietal exogenous oscillatory field
potentials on subjectively perceived memory confidence. *Neurobiology of Learning and*
Memory. 2020;168:107140. doi:10.1016/j.nlm.2019.107140

63. Chantrel Y, Trabattoni V, Orton L, Javadi AH. Intrinsic reinstatement of induced
oscillatory context. *bioRxiv*. Preprint posted online January 26,
2021:2021.01.25.428096. doi:10.1101/2021.01.25.428096

64. Meng A, Kaiser M, de Graaf TA, et al. Transcranial alternating current stimulation at
theta frequency to left parietal cortex impairs associative, but not perceptual, memory
encoding. *Neurobiology of Learning and Memory*. 2021;182:107444.
doi:10.1016/j.nlm.2021.107444

65. Luckey AM, McLeod SL, Mohan A, Vanneste S. Potential role for peripheral nerve
stimulation on learning and long-term memory: A comparison of alternating and direct
current stimulations. *Brain Stimulation: Basic, Translational, and Clinical Research in*
Neuromodulation. 2022;15(3):536-545. doi:10.1016/j.brs.2022.03.001

66. Murray NWG, Graham PL, Sowman PF, Savage G. Theta tACS impairs episodic
memory more than tDCS. *Sci Rep*. 2023;13(1):716. doi:10.1038/s41598-022-27190-y

67. Varastegan S, Kazemi R, Rostami R, Khomami S, Zandbagleh A, Hadipour AL.
Remember NIBS? tACS improves memory performance in elders with subjective
memory complaints. *GeroScience*. 2023;45(2):851-869. doi:10.1007/s11357-022-
00677-2

68. Shtoots L, Barzilay R, Gigi T, Kostovetsky V, Pollock A, Levy DA. Theta Stimulation
Enhances Early Consolidation of Semantic Memory. *J Cogn Neurosci*.
2025;37(9):1496-1510. doi:10.1162/jocn_a_02322

69. Komissarenko A, Stupina E, Malyutina S. Targeting the neural bases of novel word
acquisition using theta-band transcranial alternating current stimulation. *Language,*
Cognition and Neuroscience. 2025;40(4):487-507.
doi:10.1080/23273798.2025.2454001

70. Sun S, Annaka H, Nomura T. Gamma-frequency transcranial alternating current
stimulation over the left posterior parietal cortex enhances the long-term retention of
associative memory. *Exp Brain Res*. 2025;243(3):62. doi:10.1007/s00221-025-07009-8

71. Miller EK, Cohen JD. An integrative theory of prefrontal cortex function. *Annu Rev*
Neurosci. 2001;24:167-202. doi:10.1146/annurev.neuro.24.1.167

72. Euston DR, Gruber AJ, McNaughton BL. The Role of Medial Prefrontal Cortex in
Memory and Decision Making. *Neuron*. 2012;76(6):1057-1070.
doi:10.1016/j.neuron.2012.12.002

73. Martínez MC, Villar ME, Ballarini F, Viola H. Retroactive interference of object-in-
context long-term memory: Role of dorsal hippocampus and medial prefrontal cortex.
Hippocampus. 2014;24(12):1482-1492. doi:10.1002/hipo.22328

74. Blumenfeld RS, Lee TG, D'Esposito M. The effects of lateral prefrontal transcranial
magnetic stimulation on item memory encoding. *Neuropsychologia*. 2014;53:197-202.
doi:10.1016/j.neuropsychologia.2013.11.021

75. Ghazizadeh A, Hong S, Hikosaka O. Prefrontal Cortex Represents Long-Term Memory
of Object Values for Months. *Current Biology*. 2018;28(14):2206-2217.e5.
doi:10.1016/j.cub.2018.05.017

76. Ghazizadeh A, Griggs W, Leopold DA, Hikosaka O. Temporal–prefrontal cortical
network for discrimination of valuable objects in long-term memory. *Proceedings of the*
National Academy of Sciences. 2018;115(9):E2135-E2144.
doi:10.1073/pnas.1707695115

77. Blumenfeld RS, Ranganath C. Chapter 12 - The lateral prefrontal cortex and human
long-term memory. In: D'Esposito M, Grafman JH, eds. *Handbook of Clinical*
Neurology. Vol 163. The Frontal Lobes. Elsevier; 2019:221-235. doi:10.1016/B978-0-
12-804281-6.00012-4

78. Treves A, Rolls ET. Computational analysis of the role of the hippocampus in memory.
Hippocampus. 1994;4(3):374-391. doi:10.1002/hipo.450040319

79. Simons JS, Spiers HJ. Prefrontal and medial temporal lobe interactions in long-term
memory. *Nat Rev Neurosci*. 2003;4(8):637-648. doi:10.1038/nrn1178

80. Squire LR, Stark CEL, Clark RE. The medial temporal lobe. *Annual Review of*
Neuroscience. 2004;27(Volume 27, 2004):279-306.
doi:10.1146/annurev.neuro.27.070203.144130

81. Lisman JE, Grace AA. The Hippocampal-VTA Loop: Controlling the Entry of Information
into Long-Term Memory. *Neuron*. 2005;46(5):703-713.
doi:10.1016/j.neuron.2005.05.002

82. Guderian S, Schott BH, Richardson-Klavehn A, Düzel E. Medial temporal theta state
before an event predicts episodic encoding success in humans. *Proceedings of the*

National Academy of Sciences. 2009;106(13):5365-5370.
doi:10.1073/pnas.0900289106

83. Lega BC, Jacobs J, Kahana M. Human hippocampal theta oscillations and the
formation of episodic memories. *Hippocampus*. 2012;22(4):748-761.
doi:10.1002/hipo.20937

84. Solomon EA, Stein JM, Das S, et al. Dynamic Theta Networks in the Human Medial
Temporal Lobe Support Episodic Memory. *Current Biology*. 2019;29(7):1100-1111.e4.
doi:10.1016/j.cub.2019.02.020

85. Polanía R, Nitsche MA, Korman C, Batsikadze G, Paulus W. The importance of timing
in segregated theta phase-coupling for cognitive performance. *Current Biology*.
2012;22(14):1314-1318. doi:10.1016/j.cub.2012.05.021

86. Violante IR, Li LM, Carmichael DW, et al. Externally induced frontoparietal
synchronization modulates network dynamics and enhances working memory
performance. *eLife*. 2017;6:e22001. doi:10.7554/eLife.22001

87. Schwab BC, Misselhorn J, Engel AK. Modulation of large-scale cortical coupling by
transcranial alternating current stimulation. *Brain Stimulation*. 2019;12(5):1187-1196.
doi:10.1016/j.brs.2019.04.013

88. Schwab BC, König P, Engel AK. Spike-timing-dependent plasticity can account for
connectivity aftereffects of dual-site transcranial alternating current stimulation.
NeuroImage. 2021;237:118179. doi:10.1016/j.neuroimage.2021.118179

89. Barzegar S, Kakies CFM, Ciupercă D, Wischnewski M. Transcranial alternating current
stimulation for investigating complex oscillatory dynamics and interactions. *International
Journal of Psychophysiology*. 2025;212:112579. doi:10.1016/j.ijpsycho.2025.112579

90. Haslacher D, Cavallo A, Reber P, et al. Working memory enhancement using real-time
phase-tuned transcranial alternating current stimulation. *Brain Stimulation*.
2024;17(4):850-859. doi:10.1016/j.brs.2024.07.007

91. Herrmann CS, Rach S, Neuling T, Strüber D. Transcranial alternating current
stimulation: a review of the underlying mechanisms and modulation of cognitive
processes. *Frontiers in human neuroscience*. 2013;7:279. Accessed October 15, 2025.
https://www.frontiersin.org/journals/human-
neuroscience/articles/10.3389/fnhum.2013.00279/full

92. Antal A, Herrmann CS. Transcranial Alternating Current and Random Noise
Stimulation: Possible Mechanisms. *Neural Plasticity*. 2016;2016(1):3616807.
doi:10.1155/2016/3616807

93. Bonaiuto JJ, Bestmann S. Chapter 4 - Understanding the nonlinear physiological and
behavioral effects of tDCS through computational neurostimulation. In: Bestmann S, ed.
Progress in Brain Research. Vol 222. Computational Neurostimulation. Elsevier;
2015:75-103. doi:10.1016/bs.pbr.2015.06.013

94. Jamil A, Batsikadze G, Kuo HI, et al. Systematic evaluation of the impact of stimulation
intensity on neuroplastic after-effects induced by transcranial direct current stimulation.
The Journal of Physiology. 2017;595(4):1273-1288. doi:10.1113/JP272738

95. Siebner HR, Lang N, Rizzo V, et al. Preconditioning of low-frequency repetitive
transcranial magnetic stimulation with transcranial direct current stimulation: evidence
for homeostatic plasticity in the human motor cortex. *Journal of Neuroscience*.
2004;24(13):3379-3385. Accessed December 18, 2024.
https://www.jneurosci.org/content/24/13/3379.short

96. Brem AK, Fried PJ, Horvath JC, Robertson EM, Pascual-Leone A. Is
neuroenhancement by noninvasive brain stimulation a net zero-sum proposition?
NeuroImage. 2014;85:1058-1068. doi:10.1016/j.neuroimage.2013.07.038

97. Schutter DJLG, Wischnewski M, Bekkering H. Stability through variability: Homeostatic
plasticity and psychological resilience. *Behav Brain Sci*. 2015;38:e118.
doi:10.1017/S0140525X14001678

98. Peterson EJ, Voytek B. Homeostatic mechanisms may shape the type and duration of
oscillatory modulation. *Journal of Neurophysiology*. 2020;124(1):168-177.
doi:10.1152/jn.00119.2020

99. Wheeler ME, Petersen SE, Buckner RL. Memory's echo: Vivid remembering
reactivates sensory-specific cortex. *Proceedings of the National Academy of Sciences*.
2000;97(20):11125-11129. doi:10.1073/pnas.97.20.11125

100. Fedurco M. Long-Term Memory Search across the Visual Brain. *Neural Plast*.
2012;2012:392695. doi:10.1155/2012/392695

101. Favila SE, Kuhl BA, Winawer J. Perception and memory have distinct spatial tuning
properties in human visual cortex. *Nat Commun*. 2022;13(1):5864. doi:10.1038/s41467-
022-33161-8

102. Griffiths BJ, Jensen O. Gamma oscillations and episodic memory. *Trends in
Neurosciences*. 2023;46(10):832-846. doi:10.1016/j.tins.2023.07.003

103. Köster M, Gruber T. Rhythms of human attention and memory: An embedded process
perspective. *Front Hum Neurosci*. 2022;16. doi:10.3389/fnhum.2022.905837

104. Keizer AW, Verment RS, Hommel B. Enhancing cognitive control through
neurofeedback: A role of gamma-band activity in managing episodic retrieval.
NeuroImage. 2010;49(4):3404-3413. doi:10.1016/j.neuroimage.2009.11.023

105. Morgan HM, Muthukumaraswamy SD, Hibbs CS, et al. Feature integration in visual
working memory: parietal gamma activity is related to cognitive coordination. *Journal of
Neurophysiology*. 2011;106(6):3185-3194. doi:10.1152/jn.00246.2011

106. Rouhinen S, Panula J, Palva JM, Palva S. Load Dependence of β and γ Oscillations
Predicts Individual Capacity of Visual Attention. *J Neurosci*. 2013;33(48):19023-19033.
doi:10.1523/JNEUROSCI.1666-13.2013

107. Tseng P, Chang YT, Chang CF, Liang WK, Juan CH. The critical role of phase
difference in gamma oscillation within the temporoparietal network for binding visual
working memory. *Sci Rep*. 2016;6(1):32138. doi:10.1038/srep32138

108. Magazzini L, Singh KD. Spatial attention modulates visual gamma oscillations across
the human ventral stream. *NeuroImage*. 2018;166:219-229.
doi:10.1016/j.neuroimage.2017.10.069

109. Marshall TR, den Boer S, Cools R, Jensen O, Fallon SJ, Zumer JM. Occipital Alpha
and Gamma Oscillations Support Complementary Mechanisms for Processing Stimulus
Value Associations. *J Cogn Neurosci*. 2018;30(1):119-129. doi:10.1162/jocn_a_01185

110. Kasten FH, Wendeln T, Stecher HI, Herrmann CS. Hemisphere-specific, differential
effects of lateralized, occipital–parietal α- versus γ-tACS on endogenous but not
exogenous visual-spatial attention. *Sci Rep*. 2020;10(1):12270. doi:10.1038/s41598-
020-68992-2

111. Tiesinga P, Sejnowski TJ. Cortical Enlightenment: Are Attentional Gamma Oscillations
Driven by ING or PING? *Neuron*. 2009;63(6):727-732.
doi:10.1016/j.neuron.2009.09.009

112. Misselhorn J, Schwab BC, Schneider TR, Engel AK. Synchronization of Sensory
Gamma Oscillations Promotes Multisensory Communication. *eNeuro*. 2019;6(5).
doi:10.1523/ENEURO.0101-19.2019

718
719

Supplementary Table 1. Overview of studies using tACS at frequencies outside of the theta or gamma range (not included in the meta-analysis)

Reference	Montage	Freq. (Hz)	Intensity (mA)	Task	Time between encoding and retrieval	Hedges' g
Luckey et al. 2022	Bilateral C2 dermatome	40	3	word association task	7 days	0.17
Shtoots et al., 2024	Fz, AF4/AF3/FC2/FC1	16	1.5	delayed recall, pictures	2 hours	0.06
	Fz, AF4/AF3/FC2/FC1	16	1.5	delayed recall, pictures	1 day	0.08
	Fz, AF4/AF3/FC2/FC1	16	1.5	delayed recall, pictures	7 day	0.08
Shtoots et al., 2025	Chin, P1/P2/P3/P4	16	1.5	multiple choice exam, video	20 min	0.14
	Chin, P1/P2/P3/P4	16	1.5	multiple choice exam, video	1 day	0.12
	Chin, P1/P2/P3/P4	16	1.5	multiple choice exam, video	7 day	0.29
	P3, C3/TP7/Pz/O1	16	1.5	multiple choice exam, video	20 min	0.16
	P3, C3/TP7/Pz/O1	16	1.5	multiple choice exam, video	1 day	0.14
	P3, C3/TP7/Pz/O1	16	1.5	multiple choice exam, video	7 day	0.17
	Fz, AF4/AF3/FC2/FC1	16	1.5	multiple choice exam, video	20 min	0.10
	Fz, AF4/AF3/FC2/FC1	16	1.5	multiple choice exam, video	1 day	0.09
	Fz, AF4/AF3/FC2/FC1	16	1.5	multiple choice exam, video	7 day	-0.12