Airflow Constraints Limit Natural Airborne Transmission of Tuberculosis

Kubra F. Naqvi[1], Yuhui Guo[2]†, Yash Kulkarni[3]†, Deepak Sapkota[2], Pei Lu[1], Shibo
Wang[2], Beatriz R. S. Dias[1], Victoria A. Ektnitphong[1], Arabella E. Martin[1], Bret M. Evers[3],
Lenette L. Lu[1,4], Hui Ouyang[2*], Lydia Bourouiba[3*] and Michael U. Shiloh[1,6*]

[1] Department of Internal Medicine, University of Texas Southwestern Medical Center,
Dallas, TX

[2] Department of Mechanical Engineering, University of Texas at Dallas, Richardson, TX

[3] The Fluid Dynamics of Disease Transmission Laboratory, Fluids and Health Network,
Massachusetts Institute of Technology, Cambridge, MA

[4] Department of Pathology, University of Texas Southwestern Medical Center, Dallas, TX

[5] Department of Immunology, University of Texas Southwestern Medical Center, Dallas,
TX

[6] Department of Microbiology, University of Texas Southwestern Medical Center, Dallas,
TX

† These authors contributed equally to this work

* Correspondence: Michael.shiloh@utsouthwestern.edu; lbouro@mit.edu;
Hui.Ouyang@utdallas.edu

ABSTRACT

Tuberculosis (TB) is transmitted through the air, yet the determinants of natural airborne

transmission remain poorly defined. Early twentieth-century guinea pig studies

demonstrated efficient airborne transmission of *Mycobacterium tuberculosis* (Mtb), but

this paradigm has not been reestablished in contemporary containment facilities. Here,

we show that ventilation can impose airflow constraints that suppress transmission

under otherwise permissive conditions. Using a guinea pig model of animal-to-animal

exposure, we combined transmission experiments with quantitative airflow

measurements and particle transport modeling to explain why some housing

configurations fail to support effective exposure. Static environments and excessive

unidirectional airflow prevented transmission, whereas controlled low-velocity airflow

restored evidence of exposure, including tuberculin skin test conversion, antigen-

specific immune responses, and pulmonary inflammation consistent with early infection.

These findings identify airflow as a critical constraint on airborne TB transmission and

provide a reproducible framework to dissect host, microbial, and environmental

determinants of spread.

INTRODUCTION

Tuberculosis (TB), caused by infection with *Mycobacterium tuberculosis* (Mtb), remains one of the leading causes of infectious mortality worldwide (*1*). Interrupting the transmission cascade is essential to reducing global incidence, yet progress has been hindered by an incomplete understanding of the host, bacterial and exposure-related factors that govern airborne transmission. Experimental systems that faithfully recapitulate natural transmission are therefore critical for dissecting the biological determinants of infection and for accelerating the development of transmission-blocking interventions and vaccines.

Guinea pigs have played a central role in the history of TB research (*2*). In 1882, Robert Koch established their use as a model for TB pathogenesis, noting that naïve animals housed near infected ones occasionally developed spontaneous pulmonary disease (*3, 4*). In the 1920s and 1930s, David Perla and Max Lurie expanded on these observations, demonstrating that healthy guinea pigs co-housed with infected animals developed bronchogenic TB, and that transmission could occur even between animals housed in separate cages within the same room (*5-9*). These seminal studies provided the first direct experimental evidence for airborne transmission of Mtb. Decades later, Riley, Wells, and Mills extended this concept to humans by showing that air exhausted from TB wards could infect sentinel guinea pigs (*10-14*).

Despite the foundational impact of these studies, the experimental paradigm of natural airborne transmission was not widely carried forward into the modern biosafety era. Contemporary biosafety level 3 (BSL-3) facilities impose stringent containment and ventilation requirements that differ substantially from the environments in which early

transmission experiments were conducted. How these modern conditions influence the

biological feasibility of natural airborne transmission has received relatively limited

experimental attention.

Recent studies using alternative animal models, including minipigs (*15*) and ferrets (*16*),

have demonstrated animal-to-animal transmission of Mtb, but these systems require

high infectious doses and do not fully capture natural respiratory behaviors such as

coughing. The guinea pig remains well suited for studying transmission biology: it is

relatively low-cost compared to other large-animal models, develops necrotizing

granulomas that resemble human disease (*17*), harbors Mtb within its airways during

infection (*18*), and exhibits respiratory neurophysiology closely resembling that of

humans (*19, 20*). While guinea pigs have long been used as sentinels for human-to-

animal transmission in air sampling studies (*12, 13, 21-23*), a modern system that

supports natural guinea pig-to-guinea pig airborne transmission under BSL-3 conditions

has not been established. This gap raises the possibility that features of contemporary

cage-level airflow and ventilation may impose constraints on transmission that are

independent of host susceptibility or bacterial virulence.

Airborne transmission of Mtb depends on the generation, persistence, and inhalation of

pathogen-containing aerosols produced during normal respiratory behaviors, including

coughing, breathing, sneezing, and singing (*24-35*). Once expelled, these aerosols are

shaped by environmental factors such as temperature, humidity, and ventilation (*33*)

that influence particle desiccation, dispersal, and persistence (*36-38*). In confined

animal housing systems, effective biological exposure is therefore determined not only

by aerosol generation but also by airflow patterns within the exposure environment.

Ventilation at the level of the cage can influence whether aerosols are retained long

enough to reach a susceptible host or are removed before inhalation. At the same time,

under modern BSL-3 containment, negative-pressure ventilation imposed at the facility

level can interact with imperfectly sealed enclosures, potentially altering airflow within

cages in ways that are not apparent from cage design or nominal ventilation rates

alone. Together, these considerations raise the possibility that multiple, interacting

airflow constraints shape biological exposure during experimental transmission, yet their

combined impact has been difficult to define in living systems.

To establish a modern experimental system capable of supporting natural airborne

transmission, we revisited guinea pig-to-guinea pig spread of Mtb using housing

designs adapted for contemporary BSL-3 environments. Our initial goal was to establish

a functional animal-to-animal transmission model that preserved biological exposure

while meeting biosafety requirements, and to understand which factors proved limiting

as the system was developed. When early housing configurations failed to support

transmission, we iteratively modified cage design and airflow conditions and

incorporated quantitative particle measurements to assess how features of the

exposure environment influenced aerosol retention and inhalation. Computational

modeling was then used to interpret these observations and explain how airflow

patterns and enclosure design shape access to the recipient breathing zone. Together,

this stepwise biological and engineering approach revives a classical transmission

paradigm with modern tools and establishes a biologically grounded platform for

dissecting host, microbial, and exposure-related determinants of airborne tuberculosis

transmission.

RESULTS

Initial housing configurations do not support guinea pig-to-guinea pig airborne Mtb transmission under BSL-3 conditions

Guided by early transmission studies in guinea pigs (*8*), we first tested whether a simple co-housing design could support natural airborne transmission under modern BSL-3 conditions. We therefore tested the first iteration of our guinea pig transmission system (system 0) by co-housing naïve recipient guinea pigs with Mtb-infected donors, separated by a perforated plexiglass barrier for 12 months (Fig. 1A, B). Infection of naïve animals was monitored every other month by tuberculin skin testing (TST).

This initial design emphasized simplicity, low-cost and ease of access to system components. We modified an existing static cage by inserting a plexiglass divider containing large holes to separate donor and recipient animals (Fig. 1B). The cages were housed on metal shelving located in the most negatively pressurized room of our ABSL3 facility (relative negative pressure -0.06 WC). Because guinea pigs are highly susceptible to progressive tuberculosis following low-dose infection (50-100 CFU) (*17*), infected donors were replaced every 7-8 weeks to maintain a constant infectious source. After 12 months of exposure, none of the recipient animals converted their TSTs or showed clinical or histopathologic evidence of infection, and no Mtb CFU were recovered at necropsy. The absence of transmission in this initial configuration indicated that this simple co-housing design was insufficient to support guinea pig-to-guinea pig airborne transmission under modern BSL-3 conditions.

The absence of transmission under these conditions contrasted sharply with historical

reports from Perla and Lurie and suggested that modern biosafety engineering

constraints might fundamentally alter airflow compared to the open systems used a

century ago. In contemporary BSL-3 facilities, negative-pressure ventilation ensures

containment but may also disrupt airborne particle movement between animals. On this

basis, we next tested whether enforcing directional airflow from infected donors to naïve

recipients would be sufficient to permit transmission.

To test this, we modified a commercially available guinea pig housing system

(Allentown, Inc.) where cages are connected to both supply and exhaust air blowers to

create "system 1" (Fig. 1C, D). The new design incorporated a stainless-steel barrier

between donor and recipient animals perforated with two panels of holes positioned at

the typical nose and mouth height of an adult guinea pig. To establish unidirectional

airflow, we blocked the donor-side exhaust port (right) and the recipient-side supply

ports (left) with rubber caps (Fig. 1E). Smoke testing confirmed right-to-left airflow from

the donor to the recipient compartment (Fig. 1F, Supplemental Movie 1). Recipient

animals were then exposed to infected donors for 12 months under these unidirectional

airflow conditions. Despite these adjustments, no transmission occurred: all recipients

remained TST-negative and free of histopathologic or microbiologic evidence of

infection. These findings indicated that unidirectional airflow alone was insufficient to

overcome transmission constraints imposed by BSL-3 conditions.

Airflow and cage design determine aerosol particle retention

Because systems 0 and 1 did not support transmission, we next directly measured

aerosol particle transport across housing configurations to determine whether physical

features of the cages limited retention and exposure of airborne particles. We undertook

an iterative series of aerosol studies to define the parameters that would permit particle

transport from donor to recipient compartments.

To test aerosolized particle transport, we connected prototype cages to a supply air line

on the donor side and an exhaust vacuum on the recipient side, allowing controlled

regulation of airflow within the system. To simulate infectious particles, we used a

particle generator to produce aerosols ranging from 0.8-7 µm and sampled air at three

locations: upstream (before entering the cage), within the donor compartment, and

within the recipient compartment (Fig. 2A). Particles were loaded for 1 hour to establish

steady-state size distribution (Supplemental Fig. 1A) and sampled three times for 20

seconds totaling 1 minute at each position.

We first characterized particle behavior under ambient room pressure, beginning with

the earliest systems, 0 and 1. In system 0, which lacked directional airflow, we observed

marked particle loss from both donor and recipient compartments, particularly for

particles larger than 4 µm (Fig. 2B-C). System 1, operated with the manufacturer's

default airflow settings (8.8 L/min supply; 2.5 L/min exhaust), also showed significant

particle depletion (Fig. 2D-E). These findings indicated that the high ventilation rates

used for standard animal husbandry, intended to prevent excessive humidity, ammonia

and CO_2 accumulation, were incompatible with retention of aerosolized particles.

Successful transmission would likely require precise titration of airflow to maintain

particle dwell time without compromising animal welfare.

Guided by these results, we next developed system 2A, in which air movement was

driven solely by the exhaust fan (2.5 L/min). This modification improved particle

retention within both donor and recipient compartments under ambient conditions,
especially for particles less than 2 µm (Fig. 2F-G).

Inspection of system 1 also revealed a structural limitation: its double-walled stainless-
steel barrier created a wide and obstructed path for particle transfer between
compartments. To address this, we engineered a new single-walled stainless-steel
barrier with larger perforations to reduce resistance to airflow. We also sealed the cage
to minimize particle leakage by taping the lid with aluminum tape, adding a rubber
gasket along the lid-base junction, and securing the lid with four spring-loaded clamps.
The resulting configuration, designated system 2B, was designed to optimize both
airflow directionality and particle retention (Fig. 2H). Under ambient pressure, system
2B achieved particle retention comparable to system 2A, with stable concentrations of
particles <2 µm in both compartments (Fig. 2I). The similarity between the two systems
suggested that the improvements achieved by airflow titration and cage sealing were
robust across minor design variations. However, because all prior experiments had
occurred under negative-pressure BSL-3 conditions, further testing under those
laboratory containment conditions was essential to evaluate real-world performance.
Together, these measurements demonstrated that standard housing configurations and
ventilation rates used for animal care rapidly deplete aerosolized particles, providing a
physical explanation for the absence of transmission observed in earlier system
iterations.

Negative-pressure BSL-3 conditions alter aerosol particle retention across
housing configurations

Because all Mtb transmission studies must be performed under BSL-3 conditions, we

next evaluated how negative pressure, an essential safety feature, affects particle

retention within the cage transmission systems. To do this, we placed each system

inside a custom airtight chamber maintained at the negative-pressure setting used in

our ABSL-3 facility (relative negative pressure -0.03 WC) (Fig. 3A). Of note, the

negative pressure imposed on the system mimicked the settings of the room in which

the transmission studies were performed, -0.03 (system 1, 2A, 2B) or -0.06 (system 0).

Under these conditions, we observed a striking loss of aerosolized particles in both

donor and recipient compartments of system 0 (Fig. 3B, C). The unsealed cage design

allowed particles to escape through lid gaps and barrier edges, resulting in near-

complete depletion of particles across all measured size ranges. System 1 also

exhibited substantial particle loss under negative pressure, particularly in the recipient

compartment (Fig. 3D, E), indicating that unidirectional airflow alone could not

overcome the strong external flow imposed by facility ventilation. System 2A, which

incorporated reduced exhaust flow and a modified barrier, also showed lower particle

retention in the recipient compartment (Fig 3F, G). The most dramatic improvement was

achieved with system 2B, which combined controlled exhaust flow with cage sealing.

Under negative pressure-induced flow, system 2B maintained high concentrations of

aerosols within both donor and recipient compartments, demonstrating markedly

improved particle retention compared to all previous iterations (Fig. 3H, I).

Together, these results reveal that negative pressure-induced flow profoundly alters

internal cage mixing, often removing aerosols before they can traverse from donor to

recipient compartments. Through progressive modification of airflow, barrier design, and

cage sealing, system 2B successfully counteracted these losses, providing stable

particle retention even under the stringent airflow conditions required for BSL-3

containment. This optimized configuration was therefore selected for subsequent

transmission experiments.

Biological constraints define airflow regimes compatible with transmission

Because particle retention alone does not establish biological exposure, we next asked

whether airflow conditions that preserved aerosols also supported bacterial transport

and inhalation in vivo. To model movement of bacteria-containing particles, we

nebulized a suspension of *Mycobacterium smegmatis,* a non-pathogenic mycobacterial

species, in the donor compartment and measured bacterial recovery from air sampled

on the recipient side. Recovery of *M. smegmatis* showed a clear dependence on airflow

rate, with the highest recovery observed at the lowest exhaust setting of 0.15 L/min (Fig.

4A). This pattern suggested that reduced airflow prolongs particle dwell time within the

cage and enhances the likelihood of bacterial transport between compartments.

We then tested if aerosolized bacteria from the donor compartment could deposit into

the lungs of naïve guinea pigs housed on the recipient side. We placed individual

guinea pigs in the recipient compartment and nebulized a suspension of *M. smegmatis*

in the donor compartment for one minute to simulate a cough paroxysm. After 15

minutes of exposure to *M. smegmatis*, we collected and homogenized the lungs of

recipient guinea pigs and quantified colony forming units (CFU). All four recipient guinea

pigs had recoverable *M. smegmatis* CFU in their lungs (Fig. 4B), demonstrating that

aerosolized bacteria generated in the donor compartment were effectively transported

and inhaled by recipient animals.

Because humidity strongly influences aerosol dynamics by affecting particle desiccation

and settling (*33, 39-42*), we next quantified relative humidity (RH) in occupied cages.

We housed two guinea pigs (one per compartment) for six days in system 2B cages,

and measured RH daily at three exhaust flow rates (2.5, 1.5, and 0.15 L/min). At the

lowest airflow rate, humidity rose rapidly, reaching approximately 75 percent by day six

(Fig. 4C). Higher airflow rates maintained RH near 65% throughout the experiment,

consistent with enhanced air exchange and moisture removal (Fig. 4C). Although lower

exhaust flow improved particle transport, the associated increase in humidity was

expected to impair aerosol stability and potentially affect animal welfare during long

term studies. Based on these considerations, we selected an exhaust flow rate of 2.5

L/min for subsequent Mtb transmission experiments using the system 2B cage design.

Together, these experiments defined an airflow regime that balanced aerosol transport

with biological feasibility but did not explain why effective exposure differed across

housing designs.

Computational modeling explains particle retention differences across housing

systems

To address this gap, we used computational fluid dynamics simulations to examine how

airflow patterns and cage architecture influence particle retention and animal exposure

across housing systems. Steady-state incompressible Navier-Stokes equations coupled

to particle tracer advection-diffusion were solved to characterize airflow structure and

aerosol transport within each system under both ambient conditions and BSL-3

negative-pressure laboratory environments (Supplemental Methods). These simulations

enabled interpretation of experimental differences in particle retention and spatial

distribution across housing systems. Full details of the computational framework,

boundary conditions, and sensitivity analyses are provided in the Methods and

Supplemental Methods.

We first defined the computational geometry and boundary conditions corresponding to

the donor-recipient cage systems and surrounding laboratory environment (Fig. 5A-B).

Simulations were performed for each system under both ambient pressure and BSL-3

negative-pressure conditions, incorporating tracer injection and exhaust locations

matched to the experimental particle release and sampling setup. In the simulations,

particle concentrations were evaluated within the naïve guinea pig breathing zone (BZ),

defined as the cage volume below the animal's typical breathing height of approximately

7 cm.

We quantified particle transport using the ratio of particle concentration in the recipient

compartment (R) to that introduced upstream in the donor compartment (U). The

computed R/U values were compared with the corresponding experimentally measured

concentration ratios reported in Figs. 2B-I and 3B-I, summed over particle size. For the

simulations, R/U is shown as the average across four locations within the recipient-side

BZ (Fig. 5C-D), reflecting spatial variation in predicted particle concentrations (see

below), with individual location-specific contributions shown in Supplemental Fig.2 A-B.

Despite differences in sampling resolution, simulated and experimental R/U values

tracked closely across systems and environmental conditions (Fig. 5C-D). In particular,

simulations recapitulated the pronounced reduction in R/U observed experimentally

under BSL-3 negative-pressure conditions for systems 0 and 2A and confirmed that

system 1 remained particle-depleted due to its high ventilation rate.

To understand the spatial variation underlying these averaged R/U values, we next

examined the simulated particle concentration fields across each housing system.

These analyses revealed a clear breakdown of the well-mixed assumption, with strongly

inhomogeneous aerosol distributions in most systems under both ambient and BSL-3

conditions (Fig. 5E-H; Supplemental Movies 2, 3). Spatial heterogeneity was most

pronounced in system 2A, where sharp concentration gradients were observed within

the recipient compartment (Fig. 5G). This heterogeneity is reflected in increased

variability among the location-specific R/U values for system 2A and indicates that the

inferred transmission proxy can be sensitive to measurement probe position in systems

with non-uniform airflow patterns.

By contrast, system 1 exhibited relatively uniform but low particle concentrations due to

rapid clearance, while system 2B maintained higher and more stable particle

concentrations within the recipient-side BZ under both ambient and BSL-3 conditions

(Fig. 5E-H, Supplemental Movies 2, 3). Importantly, these analyses showed that air

changes per hour alone were insufficient to explain particle retention in this

experimental context. Although systems 2A and 2B exhibited higher ventilation rates

than system 0 under ambient conditions, localized airflow pathways and spatial

heterogeneity dominated effective exposure within the recipient compartment.

Together, these simulations provide a physics-based interpretation of the particle tracer

experiments and explain why only system 2B maintained effective aerosol retention

under BSL-3 conditions, motivating its selection for subsequent transmission studies.

Buoyancy from animal heat and respiration reshapes exposure patterns

Although particle tracer experiments and simulations captured key features of airflow-

driven transport, these analyses initially treated particles as passive tracers in an

isothermal flow. In living systems, however, animals introduce additional physical forces

through body heat and respiratory exhalations that could plausibly reshape local airflow

and aerosol transport. To account for these effects, we extended our simulations to

incorporate buoyancy arising from guinea pig body heat and exhaled air, using

physiologic parameters drawn from the literature, including animal size, body

temperature, respiratory rate, tidal volume, and characteristic animal dimensions (*43-*

46) (see Methods and Supplemental Methods for details of equations and parameters).

Guinea pigs generate continuous thermal plumes and episodic respiratory exhalations.

A thermal plume is the gentle upward airflow produced by body heat, as warm air near

the skin becomes buoyant and rises. In animals, this persistent flow can transport

exhaled aerosols away from the body and interact with ambient ventilation. These host-

derived flows interact with background ventilation and cage geometry in ways that might

alter aerosol mixing, persistence, and access to the recipient-side breathing zone (BZ).

We therefore used buoyancy-resolving simulations to assess whether host-generated

forces could meaningfully influence effective exposure under BSL-3 housing conditions.

To quantify the biological implications of these buoyancy effects, we calculated a

normalized, time-averaged particle concentration within the recipient guinea pig's BZ,

expressed relative to the concentration of particles emitted from the donor animal ($\mathcal{C}$;

Fig. 6A; see Methods). This metric captures cumulative exposure experienced by the

naïve animal under steady-state housing conditions and enables direct comparison

across system designs without reliance on absolute particle counts.

Using this exposure metric, simulations incorporating buoyancy predicted substantially

higher recipient-side exposure in system 2B compared to earlier designs (Fig. 6B,C).

Under otherwise equivalent conditions, system 2B produced approximately an order-of-

magnitude greater normalized exposure than system 1, consistent with its enhanced

particle retention. Visualization of the underlying flow fields revealed that buoyancy-

driven upward mixing in system 2B promoted transfer of particles from donor to

recipient compartments, whereas poorly sealed or highly ventilated systems diluted or

redirected aerosols away from the recipient breathing zone (Fig. 6C; Supplemental

Movie 5).

Enhanced buoyancy-induced mixing was further evident in mid-plane concentration

slices of pathogen tracer distribution (Fig. 6C) and in the higher buoyancy-driven flow

velocities observed across systems (Supplemental Fig. 3A-F and Supplemental Fig. 4A-

F). Comparisons of simulations performed with and without buoyancy demonstrated that

inclusion of animal body heat and respiratory exhalations substantially altered predicted

airflow structure and particle transport pathways (Supplemental Movies 4, 5),

highlighting the importance of host-derived forces in shaping exposure within enclosed

transmission systems.

Together, these analyses indicate that airflow patterns imposed by cage design and

ventilation interact with host-generated buoyancy to shape effective exposure in

transmission systems. While airflow rate alone was insufficient to predict transmission

potential, the combined effects of ventilation, sealing, and animal-derived buoyancy

provide a mechanistic explanation for why system 2B uniquely supported sustained

exposure. These insights informed selection of housing conditions for subsequent Mtb

transmission experiments.

Mtb transmission under optimized conditions

Having established airflow and environmental parameters that favored particle retention,

we next assessed whether these optimized conditions supported Mtb transmission

between guinea pigs. We infected guinea pigs with Mtb (Supplemental Fig. 5A-B) and

housed them in the donor compartment of system 2B cages, while naïve guinea pigs

were placed in the recipient compartment for a 17-week exposure period. Because

guinea pigs develop progressive disease following infection, donor animals were

replaced every 6 weeks to maintain a consistent infectious source.

We monitored transmission using monthly tuberculin skin testing (TST) with a two-stage

protocol (*47-49*) in which animals were tested twice with a 7-day interval between doses

(Fig. 7A). Donor animals developed robust TST responses by four weeks post-infection

(mean 10 mm; range 6-16 mm), consistent with active infection (Fig. 7B). To define a

threshold for positive conversion in recipients, we performed repeated TSTs in

unexposed control animals and found that none exceeded 5.5 mm (Supplemental Fig.

5D). Using this baseline, together with TST responses from infected donors (Fig. 7B)

and uninfected naïve animals (Supplemental Fig. 5E), we established a cutoff of 6.5 mm

to indicate infection. Within one month of exposure, one recipient exceeded this

threshold, and by week 17, nine of twelve recipient animals demonstrated positive TST

conversion (Fig. 7B; Supplemental Fig. 6).

Because Mtb exposure also elicits humoral immune responses (*50-55*), we next

quantified serum antibody responses to Mtb antigens using a bead-based multiplex

assay (Supplemental Fig. 7A) (*55*). Antigens included complex mixtures such as PPD

and defined immunodominant proteins such as antigen 85B. Area-under-the-curve

(AUC) values were calculated from dilution series, and receiver operating characteristic

(ROC) analyses were performed for individual and combined antigen-specific IgG

responses (Supplemental Fig. 7B-G). Infected donor animals exhibited strong IgG

responses, whereas unexposed controls remained negative. Using a predictive

probability cutoff of 0.5 for the combined IgG response to antigen 85B and PPD

(Supplemental Fig. 7H), eight of twelve recipient animals displayed serologic responses

consistent with Mtb exposure (Fig. 7C).

We next examined lung tissues from recipient guinea pigs by blinded histopathology

analysis. Compared to unexposed controls, exposed recipients exhibited increased

pulmonary inflammation characterized by stage 1 granulomas, intra-alveolar

hemorrhage, and small foci of epithelioid macrophages (Fig. 7D-E, Supplemental Fig.

5F-G). Although recipient animals did not exhibit weight loss (Supplemental Fig. 5C)

and no culturable Mtb was recovered at necropsy, the concordance of TST conversion,

antigen-specific serologic responses, and histopathologic changes indicates that

recipient animals were exposed to and mounted immune responses to Mtb.

Together, these findings demonstrate that the optimized system 2B configuration

supports natural airborne transmission of Mtb between infected and naïve guinea pigs.

By integrating controlled airflow, cage sealing, and structural refinement, this system

enables sufficient aerosol retention to permit biological transmission under BSL-3

containment, providing a reproducible experimental platform for studying host,

microbial, and environmental determinants of airborne tuberculosis spread.

DISCUSSION

Our current understanding of TB transmission arises largely from epidemiologic

studies of Mtb spread within human populations or from analysis of exhaled aerosols

(*24, 25, 28, 56-58*). Experimental validation of these observations has been constrained

by the absence of modern animal-to-animal transmission systems that operate under

contemporary biosafety conditions. Here, we establish a guinea pig model that supports

natural airborne transmission of Mtb under BSL-3 containment, enabling direct

experimental interrogation of the biological determinants of transmission.

Using iterative cage design and controlled transmission experiments, we

identified airflow conditions that permit sufficient aerosol persistence to support

biological exposure. Transmission required retention of particles in the 1-5 µm

aerodynamic diameter range capable of carrying Mtb (*24, 25, 28, 57*), while remaining

compatible with animal welfare and biosafety constraints. Under these optimized

conditions, recipient guinea pigs mounted antigen-specific immune responses and

developed pulmonary inflammation consistent with early infection, demonstrating that

natural airborne transmission can be reconstituted experimentally.

Importantly, without tight seal or flow pattern control, both extremes of ventilation

suppressed transmission. Static environments, despite low nominal air-change rates,

permitted particle loss through uncontrolled leakage and buoyancy-driven escape,

limiting delivery of aerosols to the recipient breathing zone. Conversely, excessive

unidirectional ventilation rapidly depleted aerosolized particles before effective exposure

could occur. These findings demonstrate that ventilation rate alone is insufficient to

predict transmission risk and highlight the importance of local airflow patterns, enclosure

integrity, and host-generated forces in determining biological exposure.

The influence of overall ventilation on infection risk has long been recognized

from the classic studies of Riley and Wells to modern public health strategies for

airborne disease control. Renewed attention to ventilation during the COVID-19

pandemic emphasized air exchange as a tool to mitigate transmission in real-world

settings (*59*). Our results provide experimental confirmation of this principle, showing

that sufficiently high ventilation suppresses transmission even in close-proximity animal

housing. At the same time, they underscore the limitations of the well-mixed air

assumption and demonstrate that local flow topology and enclosure design critically

shape exposure under biosafety-mandated ventilation conditions. In addition, our results

provide experimental confirmation that airborne TB transmission can be mitigated with

sufficient replenishment of fresh air (*48, 60-63*), as illustrated by our experimental

outcomes showing absence of animal-to-animal transmission in systems 0 and 1 under

BSL3-containment conditions.

By integrating particle tracer experiments with computational analyses, we show

that airflow heterogeneity and buoyancy generated by live animals substantially alter

aerosol transport within confined housing systems. These physical insights help explain

why only specific cage designs supported sustained exposure and transmission, despite

similar average ventilation metrics. Accounting for these effects provides a mechanistic

explanation for the historical difficulty of establishing animal-to-animal transmission

models in modern biosafety facilities.

The ability to reproduce natural transmission has important implications for

tuberculosis biology. Few experimental systems currently allow direct testing of how

bacterial, host, and environmental factors interact to shape the transmission cascade.

Existing approaches rely on artificial aerosolization of Mtb to naïve animals (*64*),

human-to-animal exposure systems (*23, 65*), or alternative animal models with

uncertain relevance to human disease (*15, 16*). The system described here enables

transmission to be studied directly between infected and naïve animals in a well-

established model of tuberculosis pathogenesis. For example, this platform can be

adapted to investigate microbial determinants of transmission such as strain lineage

(*66-72*), lipid composition (*73-75*), genetic background (*76*) or specific Mtb genes (*37*). It

also provides an opportunity to examine how donor physiology influences aerosol

generation and infection probability.

The relationship between respiratory behavior and transmission remains a particularly

important gap in our understanding of Mtb transmission biology. Although tidal breathing

alone can generate infectious bioaerosols (*32, 77-79*), coughing produces larger

quantities of particles and propels them over longer distances (*28, 33, 39, 59, 80, 81*).

Our prior work identified Mtb cell wall glycolipids sufficient to induce cough in naïve

guinea pigs (*73, 74*), and guinea pigs infected with Mtb strains lacking both lipids did not

cough, despite severe pulmonary pathology (*74*). Applying the current transmission

model to such strains will allow direct testing of how pathogen-encoded cough-inducing

factors influence transmission efficiency. In the present simulations, breathing and

coughing were assigned equivalent emitted pathogen concentrations to isolate the role

of airflow and host-generated forces in shaping exposure, independent of variation in

bacterial source. Although a single cough is expected to generate a larger number of

particles than a single breath, breathing occurs orders of magnitude more frequently

over time, making both processes biologically relevant contributors to cumulative

exposure. Our controlled framework can be extended in future studies to explicitly vary

source strength and respiratory behavior to define how cough frequency and intensity

interact with airflow to influence transmission efficiency.

This study has limitations. Although recipient animals showed consistent

immunologic and histologic evidence of exposure, culturable Mtb was not recovered

from lungs at necropsy. Similar findings have been reported in human-to-guinea pig

transmission studies, where latent infection was revealed only after immunosuppression

(*23*). In addition, replacement of moribund donor animals precluded correlation of donor

bacterial burden or cough frequency with transmission outcomes. Longer exposure

durations, recovery phases, or transient immunosuppression may increase bacterial

detection; future studies will allow donor disease burden to be correlated with

transmission efficiency. Our modeling approach employed a continuum particle tracer

rather than explicitly tracking discrete particles of varying size and deposition behavior.

While this simplification limits quantitative inference about individual particle fate, it

preserves the dominant airflow structures governing aerosol transport within the system.

Accordingly, our conclusions primarily reflect relative exposure patterns and

transmission-relevant airflow dynamics rather than absolute particle deposition.

In summary, this work revives a classical model of airborne tuberculosis

transmission and adapts it for modern biosafety conditions. By defining airflow regimes

that permit biologically effective exposure, we establish a reproducible experimental

system that supports natural Mtb transmission between animals. This platform enables

mechanistic investigation of the microbial, host, and environmental factors that govern

airborne spread and provides a foundation for developing transmission-blocking

interventions.

MATERIALS AND METHODS

Animal studies

Protocols for guinea pig infection and transmission studies were reviewed and approved

by the Institutional Animal Care and Use Committee at the University of Texas

Southwestern Medical Center (Protocol # 2018-102280-USDA).

Particle transmission

Aerosol particles were generated using a custom-built particle generation system to

model transport of respirable airborne particles under defined airflow conditions. A

syringe pump (Braintree Scientific, MA, USA) delivered EMEM (Suppl. Fig 1B)

suspension solution at a flow rate of 0.4 mL/min. Compressed air at 60 psi atomized

and dried the aerosols, with a total flow of 21.5 L/min. Relative humidity along the tubing

was monitored with a probe (Ahlborn, Germany) and maintained at 48 ± 2%.

Continuous aerosol loading into the cage was controlled at 0.3 L/min, while excess flow

was vented using a vacuum pump (Gast Manufacturing, MI, USA) regulated by a mass

flow controller (MFC; Aalborg Instruments and Controls, NY, USA).

Four case systems were evaluated: System 0, System 1, System 2A, and System 2B.

Each consisted of donor and recipient compartments separated by a divider, with 0.3

L/min entering the donor side. Flow conditions varied among tests as summarized in

Table 1. Compressed supply air (8.8 L/min) could be directed into the donor side, and

exhaust air (2.5 L/min) was regulated using a custom-built laminar flow element

connected to the facility vacuum.

Table 1: Structural and airflow characteristics of guinea pig transmission housing

systems

Testing case	System 0	System 1	System 2A	System 2B
Divider Modification	Plexiglass, double wall	Stainless steel, double wall	Stainless steel, double wall	Stainless steel, single wall
Lid sealing	Off	Off	Off	On
8.8 L/min supply	Off	On	Off	Off
2.5 L/min exhaust	Off	On	On	On
Estimated ACH from computations at ambient lab pressure	0.3	8	2	2
Estimated ACH for BSL-3-like negative pressure chamber used in particle release and sampling experiments	20	10	10	2
Effective ACH imposed by negative pressure within animal BSL3 facility	4	8	-	4
Inlet valve sealing on the donor side	On	Off (used as 8.8 L/min supply)	Off	Off
System leakage locations at lid-compartment junction	Front + back+ left + right	Front + back+ left + right	Front + back+ left + right	left + right

Aerosol particle size distributions were measured on both donor and recipient sides

using an Aerodynamic Particle Sizer (APS; Model 3321, TSI, MN, USA). The APS drew

0.3 L/min directly from each sampling site, while dilution air was added through

compressed air controlled by a MFC to achieve the required 5 L/min sampling flow.

Before aerosol introduction, clean air was flushed through the cage for 30 min to

confirm negligible background particle concentrations. Aerosol particles were then

loaded for approximately 1 h to reach steady-state size distributions (Supplemental Fig.

1A). After loading, three 20-second samples were collected per minute, alternating

between donor and recipient sides. The particle size boundary was set to 0.8-7µm.

Sampling was repeated every 5 min over a 25-min period under both ambient and

negative-pressure conditions.

To reproduce biosafety level 3 negative-pressure environments, a custom chamber (24″

× 24″ × 48″) was constructed. An inline duct fan (TerraBloom LLC, CA, USA) regulated

airflow, and chamber pressure was monitored continuously using a differential gauge

(Dwyer Instruments, Inc., IN, USA), maintained at –0.03 or -0.06 inches of water.

Mycobacterium smegmatis particle transmission

Mycobacterium smegmatis cultures were grown in Middlebrook 7H9 broth (Sigma

#M0178) supplemented with 10% OADC, 0.5% glycerol, and 0.05% Tween-80 at 37 °C

with shaking (100 rpm). Cultures were diluted to $1 \times 10\square$ CFU/mL in PBS and

aerosolized using a Blaustein Atomizing Module (CH Technologies). The atomizer was

placed in the donor compartment of the transmission cage, and aerosols were

generated for 1 min. After a 2-min delay, air from the recipient compartment was

sampled for 30 s using a button sampler (SKC, PA, USA) with a gelatin filter at 10 L/min.

The filter was dissolved in PBS and plated on Middlebrook 7H10 agar supplemented

with OADC and glycerol. Colonies were counted after overnight incubation to quantify

recovered bacteria. This procedure was repeated at exhaust flow rates of 2.5, 1.5, and

0.15 L/min.

For transmission of *M. smegmatis* to guinea pigs, the atomizer was placed in the donor

compartment containing 1×10^5 CFU/mL of *M. smegmatis* in PBS. The recipient

compartment contained 2 guinea pigs. Aerosols were generated for 1 min and allowed

to pass to the recipient compartment and inhaled by the recipient animals for 15

minutes. Following the 15-minute period, animals were sacrificed and lungs were

homogenized in PBS and plated on four 7H10 agar plates per animal to quantify

recovered bacteria. This process was repeated for a total of four guinea pigs.

Guinea pig aerosol infection

Donor guinea pigs were infected with *M. tuberculosis* Erdman in three groups on a 6-7

week cycle. Bacteria were cultured in Middlebrook 7H9 broth supplemented with 10%

OADC, 0.5% glycerol, and 0.05% Tween-80 at 37 °C using a roller apparatus. Cultures

were washed three times with PBS, then centrifuged at 500 rpm for 5 min to remove

large clumps. The supernatant was sonicated at 90% amplitude for 15-s pulses

(Branson Ultrasonics, CT, USA) to generate a single-cell suspension and diluted to $3 \times$

$10\square$ CFU/mL in PBS. The inoculum was aerosolized using a Glas-Col inhalation

exposure system (Glas-Col, IN, USA). Two animals per group were sacrificed

immediately post-exposure for day-0 CFU quantification by lung plating on 7H11 agar.

All animals were weighed weekly, and donors were euthanized seven weeks post-

infection.

Transmission experiments and cage modifications

For transmission experiments, male Hartley guinea pigs (Charles River) were used. For system 0, we used a polycarbonate static cage (Tecniplast # 2000P00SU), with a raised lid (Tecniplast #2000P224) covered by a filter top (Tecniplast #U1144B420RBOX-300) with a custom designed double walled plexiglass divider perforated with holes for particle movement between compartments. For the transmission experiment, cages were maintained on a stainless-steel rack in our ABSL3. For systems 1 and 2B, we housed animals in an Allentown 1800 15 cage rack (Allentown, NJ, PA). The standard cage bottom (Allentown #315351-1) and external bottle top (Allentown #230347) were used with the following modifications. For systems 1, 2A and 2B, the donor-side exhaust port (right) and recipient-side supply ports (left) were sealed with rubber caps. While for system 1 the supply air was "on", for systems 2A and 2B, the supply hose was disconnected from the main air inlet of the rack. For system 2B, a custom stainless-steel cage divider with 102 ventilation holes was fabricated locally to separate donor and recipient animals within each cage. To improve sealing, the underside of the lid was covered with aluminum tape, a 1.6 mm rubber gasket was applied along the lid-base junction, and four 2-inch spring clamps secured the lid once cages were positioned on the rack.

Tuberculin skin test

To determine Mtb exposure, guinea pigs were tested every four weeks using a two-step TST protocol. Animals were anesthetized under 3% isoflurane, and the right or left flank was shaved and cleaned with 70% ethanol. Tuberculin purified protein derivative (PPD; 100 µL of 5 TU, McGuff Medical Products #001079) was injected intradermally. Induration was measured at 24 h and 48 h post-injection. The procedure was repeated

589   one week later on the opposite flank. Control animals received a single dose of

590   tuberculin at baseline or the same two-step PPD dosing scheme as exposed animals to

591   control for effects of repeated testing.

Serologic analysis

Antigen-specific IgG levels were quantified as previously described with minor

modifications (*55, 82, 83*). Carboxylated microsphere beads (Luminex Corp., Austin, TX,

USA) were coupled to protein antigens via covalent NHS-ester linkages using 1-ethyl-3-

[3-dimethylaminopropyl]   carbodiimide   hydrochloride   (EDC)   and   *N*-

hydroxysulfosuccinimide (NHS) (Thermo Fisher Scientific, Waltham, MA, USA),

following the manufacturer's instructions. MagPlex® Avidin microspheres (Luminex)

were conjugated to biotinylated sulfolipid-1 (SL-1) using the manufacturer's protocol.

The following antigens were used: Ag85A (BEI Resources NR-14871), Ag85B (NR-

14870), ESAT-6 (NR-49424), CFP-10 (NR-49425), purified protein derivative (PPD;

Statens Serum Institute, Copenhagen, Denmark), *M. tuberculosis* H37Rv whole-cell

lysate (NR-14822), and *M. tuberculosis* H37Rv cell wall fraction (NR-14828).

Antigen-coupled microspheres (1,000 beads per well) were incubated with diluted

guinea pig serum (1:50, 1:250, and 1:1,250) in 96-well Bio-Plex Pro flat-bottom plates

(Bio-Rad Laboratories, Hercules, CA, USA) at 4 °C for 16 h. After washing to remove

unbound antibodies, antigen-specific IgG was detected using phycoerythrin (PE)-

conjugated goat anti-guinea pig IgG (Novus Biologicals, NBP1-74874PE). Beads were

incubated with the secondary antibody for 2 h at room temperature, washed with PBS

containing 0.05% Tween-20, and resuspended in PBS. Fluorescence intensity was

acquired using a MAGPIX® instrument running xPONENT 4.2 software (Luminex).

Relative antigen-specific IgG levels were quantified as the area under the curve (AUC)

calculated from the three serial serum dilutions for each sample.

Receiver operating characteristic (ROC) curves were generated using GraphPad Prism

version 10.5 (GraphPad Software, San Diego, CA, USA). Sensitivity and specificity at

selected cutoff points, along with 95% confidence intervals, were calculated using

logistic regression.

Histologic analysis

At study completion, animals were euthanized, and lungs were collected in 10% neutral-

buffered formalin for histopathologic analysis. Fixed tissues were transferred to 70%

ethanol and processed by the Histo Pathology Core Facility at UT Southwestern.

Paraffin-embedded blocks were sectioned at 5 μm and stained with hematoxylin and

eosin. Slides were scanned using a NanoZoomer S60 digital slide scanner, randomized,

and blinded prior to evaluation by a pathologist. The number and stage of granulomas

were recorded, and pulmonary inflammation was described for each animal. After

unblinding, data were compared between naïve and exposed animals using a Mann-

Whitney test (GraphPad Prism v10).

Computational modeling of airflow and particle transport within transmission housing

systems

To interpret experimentally observed differences in particle retention and exposure

across housing designs, we used computational fluid dynamics simulations to model

airflow and tracer particle transport within the transmission cages. Simulations were

performed under both ambient laboratory conditions and negative-pressure BSL-3

environments, using cage geometries and airflow boundary conditions matched to

experimental configurations.

Airflow was modeled as incompressible, steady-state flow, and aerosol particles were

treated as passive tracers subject to advection and diffusion. To account for host-

generated effects present in occupied cages, additional simulations incorporated

buoyancy arising from animal body heat and respiratory exhalations using physiologic

parameters drawn from the literature, including guinea pig size, body temperature,

respiratory rate, tidal volume, and characteristic body dimensions.

Full governing equations, numerical methods, boundary conditions, parameter values,

and sensitivity analyses are provided in the Supplemental Methods.

Metric of relative exposure used to compare systems.

To enable biologically meaningful comparison of exposure across housing designs, we

defined a normalized, time-averaged exposure metric, $\mathcal{C}$, representing the average

concentration of particles within the recipient guinea pig breathing zone relative to the

concentration emitted by the donor animal. The breathing zone was defined as the

region from which inhaled air is predominantly drawn during normal respiration.

This metric captures cumulative inhalation exposure over time and allows comparison

across systems independent of absolute particle emission rates. The mathematical

definition of $\mathcal{C}$, spatial averaging procedures, and parameter choices are provided in the

Supplemental Methods.

References

1. Global Tuberculosis Report 2024. *Geneva:World Health Organization* **Licence: CC**
BY-NC-SA 3.0 IGO, (2024).

2. P. R. Donald, A. H. Diacon, C. Lange, A. M. Demers, F. von Groote-Bidlingmaier, E.
Nardell, Droplets, dust and guinea pigs: an historical review of tuberculosis
transmission research, 1878-1940. *Int J Tuberc Lung Dis* **22**, 972-982 (2018).

3. R. Koch, The etiology of tuberculosis. *The germ theory of disease*, 109-115 (1882).

4. R. Koch, Gesammelte Werke von Robt. Koch. *Leipsic*, 512 (1912).

5. D. Perla, Experimental Epidemiology of Tuberculosis : The Elimination of Tubercle
Bacilli in the Feces, Bile, and Urine of Infected Guinea Pigs. *J Exp Med* **45**, 1025-
1035 (1927).

6. D. Perla, Experimental Epidemiology of Tuberculosis. *J Exp Med* **45**, 209-226
(1927).

7. M. B. Lurie, Experimental Epidemiology of Tuberculosis : The Route of Infection in
Naturally Acquired Tuberculosis of the Guinea Pig. *J Exp Med* **51**, 769-776 (1930).

8. M. B. Lurie, Experimental Epidemiology of Tuberculosis : The Effect of Eliminating
Exposure to Enteric Infection on the Incidence and Course of Tuberculosis Acquired
by Normal Guinea Pigs Confined with Tuberculous Cage Mates. *J Exp Med* **51**, 753-
768 (1930).

9. M. B. Lurie, Experimental Epidemiology of Tuberculosis : Air-Borne Contagion of
Tuberculosis in an Animal Room. *J Exp Med* **51**, 743-751 (1930).

10. R. L. Riley, What nobody needs to know about airborne infection. *Am J Respir Crit*
Care Med **163**, 7-8 (2001).

11. W. F. Wells, *Airborne Contagion and Air Hygiene: An Ecological Study of Droplet*
Infections (Harvard University Press, 1955), pp. 423.

12. R. L. Riley, C. C. Mills, F. O'Grady, L. U. Sultan, F. Wittstadt, D. N. Shivpuri,
Infectiousness of air from a tuberculosis ward. Ultraviolet irradiation of infected air:
comparative infectiousness of different patients. *Am Rev Respir Dis* **85**, 511-525
(1962).

13. R. L. Riley, W. F. Wells, C. C. Mills, W. Nyka, R. L. McLean, Air hygiene in
tuberculosis: quantitative studies of infectivity and control in a pilot ward. *Am Rev*
Tuberc **75**, 420-431 (1957).

14. W. F. Wells, H. L. Ratcliffe, C. Grumb, On the mechanics of droplet nuclei infection;
quantitative experimental air-borne tuberculosis in rabbits. *Am J Hyg* **47**, 11-28
(1948).

15. L. Ramos, A. Obregon-Henao, M. Henao-Tamayo, R. Bowen, J. K. Lunney, M.
Gonzalez-Juarrero, The minipig as an animal model to study Mycobacterium
tuberculosis infection and natural transmission. *Tuberculosis (Edinb)* **106**, 91-98
(2017).

16. T. Gupta, N. Somanna, T. Rowe, M. LaGatta, S. Helms, S. O. Owino, T. Jelesijevic,
S. Harvey, W. Jacobs, T. Voss, K. Sakamoto, C. Day, C. Whalen, R. Karls, B. He, S.
M. Tompkins, A. Bakre, T. Ross, F. D. Quinn, Ferrets as a model for tuberculosis
transmission. *Front Cell Infect Microbiol* **12**, 873416 (2022).

17. I. M. Orme, D. J. Ordway, Mouse and Guinea Pig Models of Tuberculosis. *Microbiol
Spectr* **4**, (2016).

18. B. K. Podell, D. F. Ackart, A. Obregon-Henao, S. P. Eck, M. Henao-Tamayo, M.
Richardson, I. M. Orme, D. J. Ordway, R. J. Basaraba, Increased severity of
tuberculosis in Guinea pigs with type 2 diabetes: a model of diabetes-tuberculosis
comorbidity. *Am J Pathol* **184**, 1104-1118 (2014).

19. B. J. Canning, The cough reflex in animals: relevance to human cough research.
Lung **186 Suppl 1**, S23-28 (2008).

20. S. B. Mazzone, N. Mori, B. J. Canning, Synergistic interactions between airway
afferent nerve subtypes regulating the cough reflex in guinea-pigs. *J Physiol* **569**,
559-573 (2005).

21. E. A. Nardell, Air sampling for tuberculosis- homage to the lowly guinea pig. *Chest*
116, 1143-1145 (1999).

22. A. Stoltz, R. R. Nathavitharana, E. de Kock, V. Ueckermann, P. Jensen, C. M.
Mendel, M. Spigelman, E. A. Nardell, Estimating the Early Transmission Inhibition of
New Treatment Regimens for Drug-Resistant Tuberculosis. *J Infect Dis* **232**, 143-
151 (2025).

23. A. S. Dharmadhikari, R. J. Basaraba, M. L. Van Der Walt, K. Weyer, M. Mphahlele,
K. Venter, P. A. Jensen, M. W. First, S. Parsons, D. N. McMurray, I. M. Orme, E. A.
Nardell, Natural infection of guinea pigs exposed to patients with highly drug-
resistant tuberculosis. *Tuberculosis (Edinb)* **91**, 329-338 (2011).

24. K. P. Fennelly, E. C. Jones-Lopez, I. Ayakaka, S. Kim, H. Menyha, B. Kirenga, C.
Muchwa, M. Joloba, S. Dryden-Peterson, N. Reilly, A. Okwera, A. M. Elliott, P. G.
Smith, R. D. Mugerwa, K. D. Eisenach, J. J. Ellner, Variability of infectious aerosols
produced during coughing by patients with pulmonary tuberculosis. *Am J Respir Crit
Care Med* **186**, 450-457 (2012).

25. K. P. Fennelly, J. W. Martyny, K. E. Fulton, I. M. Orme, D. M. Cave, L. B. Heifets,
Cough-generated aerosols of Mycobacterium tuberculosis: a new method to study
infectiousness. *Am J Respir Crit Care Med* **169**, 604-609 (2004).

26. R. D. Turner, G. H. Bothamley, Cough and the transmission of tuberculosis. *J Infect*
Dis **211**, 1367-1372 (2015).

27. R. G. Loudon, S. K. Spohn, Cough frequency and infectivity in patients with
pulmonary tuberculosis. *Am Rev Respir Dis* **99**, 109-111 (1969).

28. T. C. Tristao, M. A. M. Ferreira, P. S. de Almeida Junior, L. G. S. Castellani, M. N.
Brunetti, E. C. Jones-Lopez, K. P. Fennelly, M. R. Barer, C. H. Fantecelle, S. A.
Morellato, D. J. Hadad, J. J. Ellner, R. Dietze, M. Palaci, Dynamic of infectious
aerosols generated by cough from patients with pulmonary tuberculosis. *J Aerosol*
Sci **189**, (2025).

29. J. P. Duguid, The size and the duration of air-carriage of respiratory droplets and
droplet-nuclei. *J Hyg (Lond)* **44**, 471-479 (1946).

30. R. G. Loudon, R. M. Roberts, Singing and the dissemination of tuberculosis. *Am*
Rev Respir Dis **98**, 297-300 (1968).

31. B. T. Mangura, E. C. Napolitano, M. R. Passannante, R. J. McDonald, L. B.
Reichman, Mycobacterium tuberculosis miniepidemic in a church gospel choir.
Chest **113**, 234-237 (1998).

32. R. Dinkele, S. Gessner, A. McKerry, B. Leonard, J. Leukes, R. Seldon, D. F. Warner,
R. Wood, Aerosolization of Mycobacterium tuberculosis by Tidal Breathing. *Am J*
Respir Crit Care Med **206**, 206-216 (2022).

33. L. Bourouiba, Fluid Dynamics of Respiratory Infectious Diseases. *Annu Rev Biomed*
Eng **23**, 547-577 (2021).

34. L. Bourouiba, IMAGES IN CLINICAL MEDICINE. A Sneeze. *N Engl J Med* **375**, e15
(2016).

35. B. E. Scharfman, A. H. Techet, J. W. M. Bush, L. Bourouiba, Visualization of sneeze
ejecta: steps of fluid fragmentation leading to respiratory droplets. *Exp Fluids* **57**, 24
(2016).

36. C. D. Brown, B. M. Lee, H. Liu, A. Wu, A. Tellez, H. Zou, P. Singh, K. Saito, S.
Mishra, M. Brown, A. Saleh, N. Odjourian, M. Cristaldo, M. Gan, Q. Liu, M.
Gengenbacher, S. Darst, E. Campbell, C. Nathan, K. Y. Rhee, Desiccation-induced
DNA damage facilitates drug resistance in Mycobacterium tuberculosis. *bioRxiv*,
(2025).

37. S. Mishra, P. R. Singh, X. Hu, L. Lopez-Quezada, A. Jinich, R. Jahn, L. Geurts, N.
Shen, M. A. DeJesus, T. Hartman, K. Rhee, M. Zimmerman, V. Dartois, R. M. Jones,

X. Jiang, R. Almada-Monter, L. Bourouiba, C. Nathan, Candidate transmission
survival genome of Mycobacterium tuberculosis. *Proc Natl Acad Sci U S A* **122**,
e2425981122 (2025).

38. P. R. Singh, S. Mishra, A. Jinich, X. Jiang, F. Tsang, M. DeJesus, K. Y. Rhee, R. J.
Kaner, P. L. Leopold, R. G. Crystal, C. Nathan, Genes required for Mycobacterium
tuberculosis to survive the transition from aerosol to pulmonary alveolar lining fluid
and early infection in a model of transmission. *bioRxiv*, (2025).

39. H. Li, F. Y. Leong, G. Xu, C. W. Kang, K. H. Lim, B. H. Tan, C. M. Loo, Airborne
dispersion of droplets during coughing: a physical model of viral transmission. *Sci*
Rep **11**, 4617 (2021).

40. C. A. Verheyen, L. Bourouiba, Associations between indoor relative humidity and
global COVID-19 outcomes. *Journal of The Royal Society Interface* **19**, (2022).

41. W. F. WELLS, ON AIR-BORNE INFECTION*: STUDY II. DROPLETS AND
DROPLET NUCLEI. *American Journal of Epidemiology* **20**, 611-618 (1934).

42. A. Skanata, F. Spagnolo, M. Metz, D. S. Smyth, J. J. Dennehy, Humidity Reduces
Rapid and Distant Airborne Dispersal of Viable Viral Particles in Classroom Settings.
Environ Sci Technol Lett **9**, 632-637 (2022).

43. M. O. Amdur, J. Mead, Mechanics of respiration in unanesthetized guinea pigs. *Am*
J Physiol **192**, 364-368 (1958).

44. H. H. Kibler, S. Brody, D. Worstell, Surface area and metabolism of growing guinea
pigs. *J Nutr* **33**, 331-338 (1947).

45. M. A. Sucknow, S. A. Karla, R. P. Wilson, *The Laboratory Rabbit, Guinea Pig,*
Hamster and other Rodents (Academic Press, 2012).

46. B. Wiggs, A. Churg, J. Wright, Influence of weight on pulmonary function in the adult
guinea pig. *Respiration* **58**, 37-41 (1991).

47. Targeted tuberculin testing and treatment of latent tuberculosis infection. This official
statement of the American Thoracic Society was adopted by the ATS Board of
Directors, July 1999. This is a Joint Statement of the American Thoracic Society
(ATS) and the Centers for Disease Control and Prevention (CDC). This statement
was endorsed by the Council of the Infectious Diseases Society of America. (IDSA),
September 1999, and the sections of this statement. *Am J Respir Crit Care Med*
161, S221-247 (2000).

48. P. A. Jensen, L. A. Lambert, M. F. Iademarco, R. Ridzon, Cdc, Guidelines for
preventing the transmission of Mycobacterium tuberculosis in health-care settings,
2005. *MMWR Recomm Rep* **54**, 1-141 (2005).

49. Z. Taylor, C. M. Nolan, H. M. Blumberg, S. American Thoracic, C. Centers for
Disease, Prevention, A. Infectious Diseases Society of, Controlling tuberculosis in
the United States. Recommendations from the American Thoracic Society, CDC,
and the Infectious Diseases Society of America. *MMWR Recomm Rep* **54**, 1-81
(2005).

50. L. R. L. Davies, M. T. Smith, D. Cizmeci, S. Fischinger, J. Shih-Lu Lee, L. L. Lu, E.
D. Layton, A. D. Grant, K. Fielding, C. M. Stein, W. H. Boom, T. R. Hawn, S. M.
Fortune, R. S. Wallis, G. J. Churchyard, G. Alter, C. Seshadri, IFN-gamma
independent markers of Mycobacterium tuberculosis exposure among male South
African gold miners. *EBioMedicine* **93**, 104678 (2023).

51. E. E. Kroon, C. J. Kinnear, M. Orlova, S. Fischinger, S. Shin, S. Boolay, G. Walzl, A.
Jacobs, R. J. Wilkinson, G. Alter, E. Schurr, E. G. Hoal, M. Moller, An observational
study identifying highly tuberculosis-exposed, HIV-1-positive but persistently TB,
tuberculin and IGRA negative persons with M. tuberculosis specific antibodies in
Cape Town, South Africa. *EBioMedicine* **61**, 103053 (2020).

52. L. R. L. Davies, C. Wang, P. Steigler, K. A. Bowman, S. Fischinger, M. Hatherill, M.
Fisher, S. K. Mbandi, M. Rodo, T. H. M. Ottenhoff, H. M. Dockrell, J. S. Sutherland,
H. Mayanja-Kizza, W. H. Boom, G. Walzl, S. H. E. Kaufmann, E. Nemes, T. J.
Scriba, D. Lauffenburger, G. Alter, S. M. Fortune, Age and sex influence antibody
profiles associated with tuberculosis progression. *Nat Microbiol* **9**, 1513-1525
(2024).

53. L. L. Lu, A. W. Chung, T. R. Rosebrock, M. Ghebremichael, W. H. Yu, P. S. Grace,
M. K. Schoen, F. Tafesse, C. Martin, V. Leung, A. E. Mahan, M. Sips, M. P. Kumar,
J. Tedesco, H. Robinson, E. Tkachenko, M. Draghi, K. J. Freedberg, H. Streeck, T.
J. Suscovich, D. A. Lauffenburger, B. I. Restrepo, C. Day, S. M. Fortune, G. Alter, A
Functional Role for Antibodies in Tuberculosis. *Cell* **167**, 433-443 e414 (2016).

54. L. L. Lu, J. Das, P. S. Grace, S. M. Fortune, B. I. Restrepo, G. Alter, Antibody Fc
Glycosylation Discriminates Between Latent and Active Tuberculosis. *J Infect Dis*
222, 2093-2102 (2020).

55. L. L. Lu, M. T. Smith, K. K. Q. Yu, C. Luedemann, T. J. Suscovich, P. S. Grace, A.
Cain, W. H. Yu, T. R. McKitrick, D. Lauffenburger, R. D. Cummings, H. Mayanja-
Kizza, T. R. Hawn, W. H. Boom, C. M. Stein, S. M. Fortune, C. Seshadri, G. Alter,
IFN-gamma-independent immune markers of Mycobacterium tuberculosis
exposure. *Nat Med* **25**, 977-987 (2019).

56. Y. Lan, I. Rancu, M. H. Chitwood, B. Sobkowiak, K. Nyhan, H. H. Lin, C. Y. Wu, B.
Mathema, T. S. Brown, C. Colijn, J. L. Warren, T. Cohen, Integrating genomic and
spatial analyses to describe tuberculosis transmission: a scoping review. *Lancet*
Microbe **6**, 101094 (2025).

57. K. P. Fennelly, Particle sizes of infectious aerosols: implications for infection control.
Lancet Respir Med **8**, 914-924 (2020).

58. B. Patterson, S. Hermans, R. Wood, F. Cobelens, Quantification of tuberculosis
exposure in a high-burdened setting. *Sci Rep* **15**, 22687 (2025).

59. L. Bourouiba, Turbulent Gas Clouds and Respiratory Pathogen Emissions: Potential
Implications for Reducing Transmission of COVID-19. *JAMA* **323**, 1837-1838
(2020).

60. W.H.O., "Global technical consultation report on proposed terminology for
pathogens that transmit through the air." (2024;
https://www.who.int/publications/m/item/global-technical-consultation-report-on-
proposed-terminology-for-pathogens-that-transmit-through-the-air).

61. C. B. Beggs, C. J. Noakes, P. A. Sleigh, L. A. Fletcher, K. Siddiqi, The transmission
of tuberculosis in confined spaces: an analytical review of alternative
epidemiological models. *Int J Tuberc Lung Dis* **7**, 1015-1026 (2003).

62. A. R. Escombe, C. C. Oeser, R. H. Gilman, M. Navincopa, E. Ticona, W. Pan, C.
Martinez, J. Chacaltana, R. Rodriguez, D. A. Moore, J. S. Friedland, C. A. Evans,
Natural ventilation for the prevention of airborne contagion. *PLoS Med* **4**, e68
(2007).

63. R. Granich, N. J. Binkin, W. R. Jarvis, P. M. Simone, H. L. Rieder, M. A. Espinal, J.
Kumaresan, *Guidelines for the prevention of tuberculosis in health care facilities in*
resource-limited settings (World health organization (WHO), 1999).

64. F. Nuritdinov, J. Woo, M. J. Schmidt, N. M. Odjourian, M. Cristaldo, M. Dougher, R.
Antilus-Sainte, T. Heldt, K. Rhee, L. Bourouiba, M. Gengenbacher, Experimental
system enables studies of Mycobacterium tuberculosis during aerogenic
transmission. *mBio* **16**, e0095825 (2025).

65. A. R. Escombe, C. Oeser, R. H. Gilman, M. Navincopa, E. Ticona, C. Martinez, L.
Caviedes, P. Sheen, A. Gonzalez, C. Noakes, D. A. Moore, J. S. Friedland, C. A.
Evans, The detection of airborne transmission of tuberculosis from HIV-infected
patients, using an in vivo air sampling model. *Clin Infect Dis* **44**, 1349-1357 (2007).

66. M. Coscolla, S. Gagneux, Consequences of genomic diversity in Mycobacterium
tuberculosis. *Semin Immunol* **26**, 431-444 (2014).

67. Z. Cui, D. Lin, V. Chongsuvivatwong, E. A. Graviss, A. Chaiprasert, P.
Palittapongarnpim, M. Lin, J. Ou, J. Zhao, Hot and Cold Spot Areas of Household
Tuberculosis Transmission in Southern China: Effects of Socio-Economic Status
and Mycobacterium tuberculosis Genotypes. *Int J Environ Res Public Health* **16**,
(2019).

68. C. C. Huang, A. L. Chu, M. C. Becerra, J. T. Galea, R. Calderon, C. Contreras, R.
Yataco, Z. Zhang, L. Lecca, M. B. Murray, Mycobacterium tuberculosis Beijing
Lineage and Risk for Tuberculosis in Child Household Contacts, Peru. *Emerg Infect*
Dis **26**, 568-578 (2020).

69. K. E. Holt, P. McAdam, P. V. K. Thai, N. T. T. Thuong, D. T. M. Ha, N. N. Lan, N. H.
Lan, N. T. Q. Nhu, H. T. Hai, V. T. N. Ha, G. Thwaites, D. J. Edwards, A. P. Nath, K.
Pham, D. B. Ascher, J. Farrar, C. C. Khor, Y. Y. Teo, M. Inouye, M. Caws, S. J.
Dunstan, Frequent transmission of the Mycobacterium tuberculosis Beijing lineage
and positive selection for the EsxW Beijing variant in Vietnam. *Nat Genet* **50**, 849-
856 (2018).

70. C. Yang, T. Luo, G. Sun, K. Qiao, G. Sun, K. DeRiemer, J. Mei, Q. Gao,
Mycobacterium tuberculosis Beijing strains favor transmission but not drug
resistance in China. *Clin Infect Dis* **55**, 1179-1187 (2012).

71. S. Niemann, R. Diel, G. Khechinashvili, M. Gegia, N. Mdivani, Y. W. Tang,
Mycobacterium tuberculosis Beijing lineage favors the spread of multidrug-resistant
tuberculosis in the Republic of Georgia. *J Clin Microbiol* **48**, 3544-3550 (2010).

72. B. Koleske, C. Schill, S. Rajagopalan, S. Shee, Y. B. Martinez-Martinez, M. Gupta,
J. Shen, W. R. Jacobs, Jr., W. R. Bishai, Loss of the PPE71-esxX-esxY-PPE38
locus drives adaptive transcriptional responses and hypervirulence of
Mycobacterium tuberculosis lineage 2. *Sci Adv* **11**, eadw5194 (2025).

73. K. F. Naqvi, L. L. Warden, J. H. M. van Dijk, D. K. Naik, F. de Jesus Espinosa-
Becerra, I. M. Okolie, V. V. F. Verhoeks, K. C. Rahlwes, C. P. Xavier, J. D. C. Codee,
T. J. Price, M. U. Shiloh, Mycobacterial Phenolic Glycolipid Triggers ATP-Mediated
Neuronal P2X3 Signaling and Cough. *bioRxiv*, (2025).

74. C. R. Ruhl, B. L. Pasko, H. S. Khan, L. M. Kindt, C. E. Stamm, L. H. Franco, C. C.
Hsia, M. Zhou, C. R. Davis, T. Qin, L. Gautron, M. D. Burton, G. L. Mejia, D. K. Naik,
G. Dussor, T. J. Price, M. U. Shiloh, Mycobacterium tuberculosis Sulfolipid-1
Activates Nociceptive Neurons and Induces Cough. *Cell* **181**, 293-305 e211 (2020).

75. K. Moopanar, N. E. Mvubu, Lineage-specific differences in lipid metabolism and its
impact on clinical strains of Mycobacterium tuberculosis. *Microb Pathog* **146**,
104250 (2020).

76. S. Verma, K. Bhatt, A. Lovey, R. Ribeiro-Rodrigues, J. Durbin, E. C. Jones-Lopez,
M. Palaci, S. A. Vinhas, D. Alland, R. Dietze, J. J. Ellner, P. Salgame, Transmission
phenotype of Mycobacterium tuberculosis strains is mechanistically linked to
induction of distinct pulmonary pathology. *PLoS Pathog* **15**, e1007613 (2019).

77. R. Dinkele, S. Gessner, B. Patterson, A. McKerry, Z. Hoosen, A. Vazi, R. Seldon, A.
Koch, D. F. Warner, R. Wood, Persistent Mycobacterium tuberculosis bioaerosol
release in a tuberculosis-endemic setting. *iScience* **27**, 110731 (2024).

78. R. Dinkele, P. Y. Khan, D. F. Warner, Mycobacterium tuberculosis transmission: the
importance of precision. *Lancet Infect Dis* **24**, 679-681 (2024).

79. B. Patterson, R. Wood, Is cough really necessary for TB transmission? *Tuberculosis*
(Edinb) **117**, 31-35 (2019).

80. L. Bourouiba, E. Dehandschoewercker, J. W. M. Bush, Violent expiratory events: on
coughing and sneezing. *J Fluid Mech* **745**, 537-563 (2014).

81. C. Wang, J. Xu, H. Zhai, L. K. So, H. Guo, Mapping full-range infection transmission
from speaking, coughing, and sneezing in indoor environments and its impact on
social distancing. *J Hazard Mater* **490**, 137782 (2025).

82. E. H. Adhikari, P. Lu, Y. J. Kang, A. R. McDonald, J. E. Pruszynski, T. A. Bates, S. K.
McBride, M. Trank-Greene, F. G. Tafesse, L. L. Lu, Diverging Maternal and Cord
Antibody Functions From SARS-CoV-2 Infection and Vaccination in Pregnancy. *J
Infect Dis* **229**, 462-472 (2024).

83. T. A. Bates, P. Lu, Y. J. Kang, D. Schoen, M. Thornton, S. K. McBride, C. Park, D.
Kim, W. B. Messer, M. E. Curlin, F. G. Tafesse, L. L. Lu, BNT162b2-induced
neutralizing and non-neutralizing antibody functions against SARS-CoV-2 diminish
with age. *Cell Rep* **41**, 111544 (2022).

84. R. R. Schmidt, K. C. Kailash, S. V. Patankar, in *Ninth Intersociety Conference on
Thermal and Thermomechanical Phenomena in Electronic Systems*. (IEEE, Las
Vagas, 2004), vol. 1, pp. 571-578.

85. Y. Tominaga, M. Shirzadi, RANS CFD modeling of the flow around a thin windbreak
fence with various porosities: Validation using wind tunnel measurements. *Journal
of Wind Engineering and Industrial Aerodynamics* **230**, 105176 (2022).

86. T. J. R. Hughes, *The Finite Element Method: Linear Static and Dynamic Finite
Element Analysis* (Dover Civil and Mechanical Engineering, Prentice Hall, 1987),
pp. 768.

87. P. J. Roache, QUANTIFICATION OF UNCERTAINTY IN COMPUTATIONAL FLUID
DYNAMICS. *Annual Review of Fluid Mechanics* **29**, 123-160 (1997).

88. J. Boussinesq, *Théorie analytique de la chaleur mise en harmonie avec la
thermodynamique et avec la théorie mecanique de la lumière* (Gauthier-Villars,
Paris, 1903), vol. II.

89. N. M. Leunberger, R. Yang, L. Münzel, R. Verzicco, D. Lohse, L. Bourouiba, P.
Jenny, Fast monotonically integrated large eddy simulation solver: validation of a
new scalable tool to study and optimize indoor ventilation. *FLOW* **4**, E23 (2024).

90. W. Hayes, *CRC Handbook of Chemistry and Physics* (CRC Press, Boca-Raton, FL,
ed. 99th, 1998).

91. A. C. Guyton, Measurement of the respiratory volumes of laboratory animals. *Am J
Physiol* **150**, 70-77 (1947).

92. G. K. Batchelor, *An Introduction to Fluid Dynamics* (Cambridge Mathematical
Library, Cambridge University Press, Cambridge, 2000).

93. *Prandtl-Essentials of Fluid Mechanics* H. Oertel Ed (Applied Mathematical Sciences,
Springer, ed. 3, 2009).

94. "2025 ASHRAE Handbook—Fundamentals" (American Society of Heating,
Refrigerating and Air-Conditioning Engineers, 2025), chap. Airflow Around Buildings

95. I. E. Idelchik, *Handbook of Hydraulic Resistance* (Begell House, New York, ed. 4th,
2007).

96. H. Oertel, *Prandtl – Essentials of Fluid Mechanics* (Springer, New York, NY, 2020).

97. E. Torricelli, *De motu gravium naturaliter descendentium, et projectorum* (Florence,
1644).

98. National Research Council, in *Guide for the Care and Use of Laboratory Animals*
(The National Academies Collection: Reports funded by National Institutes of
Health, National Academies Press, Washington (DC), ed. 8th, 2011).

99. University of California, "Biosafety Level 3 (BSL-3) Laboratory Design Standards"
(2015; https://www.ucop.edu/safety-and-loss-prevention/_files/hcloc/uc-bsl3-design-
standards-final.pdf).

ACKNOWLEDGMENTS

The authors would like to thank Cody Ruhl for preliminary experiments, and all

members of the Shiloh lab for their support and constructive feedback. We also thank

members of the Animal Resources Center at UT Southwestern for providing support

with animal husbandry and the engineering team at Allentown, Inc. for their feedback

throughout this project.

FUNDING:

NIH T32 AI007520 (K.F.N), NIH R21AI181258 (H.O.), NIH R21AI188518 (H.O.), NIH

R21AI137545 (M.U.S.), NIH R01 AI158688 (M.U.S), NIH R01AI158858 (L.L.L.) , NIH

P01 AI159402 (L.B., M.U.S), Burroughs Wellcome Fund 1017894 (M.U.S), Michael

Shiloh and Lenette Lu would also like to acknowledge generous support from the

Disease Oriented Clinical Scholar program at UT Southwestern Medical Center.

AUTHOR CONTRIBUTIONS

Conceptualization, K.F.N, L.B. and M.U.S. Formal Analysis, K.F.N, Y.G., Y.K., D.S., P.L.,

S.W., A.M., B.M.E., L.L., H.O., L.B., and M.U.S. Funding Acquisition, L.B. and M.U.S.

Investigation, K.F.N, Y.G., Y.K., D.S., P.L., S.W., B.R.S.D., V.A.E., A.E.M., B.M.E., L.L.,

and L.B. Project Administration, K.F.N and M.U.S.Resources, H.O., M.U.S. Validation,

K.F.N, Y.G., D.S., P.L., A.E.M., Y.K. Supervision, L.L.L., H.O., L.B. and M.U.S. Writing-

original draft, K.F.N, Y.G., Y.K., H.O., L.B., and M.U.S. Writing-Review and Editing, all

authors.

COMPETING INTERESTS

All authors declare that they have no competing interests.

Figure legends:

Figure 1: Airflow Modifications of Preliminary Transmission Systems. (A)

Experimental scheme for 12-month transmission studies. Blue ticks indicate when new

donor groups were added to cages and stars denote when TSTs were performed. (B)

System 0 cage set up with static airflow and plexiglass divider between donor and

recipient compartments. (C) 15 cage rack as available by Allentown, Inc. (D) System 1

cage set up. Donor side is connected to supply hose which delivers 8.8 L/min into the

cage. Air then passes through a stainless-steel divider to recipient compartment and is

exhausted at 2.5 L/min. (E) Modifications to cage rack include plugging the exhaust port

on the donor side and supply port on the recipient side to drive unidirectional airflow. (F)

Smoke test to confirm unidirectional movement of air within system 1. See also

Supplemental Movie 1.

Figure 2: Particle movement driven by airflow under ambient pressure. (A)

Experimental overview of particle transport. Cage systems are connected to supply air

(where applicable) and exhaust vacuum to control airflow within the cages. Particles are

introduced to the donor compartment using a particle generator. An APS particle

sampler collects particles upstream before entering the cage, in the donor and recipient

compartments. (B-C) Schematic and particle sampling of System 0 under static flow

conditions. (D-E) Schematic and particle sampling of System 1 with supply air set to 8.8

L/min and exhaust to 2.5 L/min. Dashed line indicates unsealed cage lid. (F) System 2A

schematic with air supply removed (0 L/min) and exhaust set to 2.5 L/min. (G) Particle

transport within system 2A. (H-I) Schematic and particle transport of system 2B

modified to include cage lid sealing (solid line) and a single stainless-steel barrier. All

particle transport results are represented as number of particles per cubic centimeter of

air (#/cc) in the size range of 0.8-7 μm. Sampling of particles includes upstream (red)

before entering the cage, in the donor compartment (blue) and the recipient

compartment (black).

Figure 3: Negative pressure influences retention of particles within cage systems.

(A) Experimental set-up of particle transport where cage systems are placed inside a

custom chamber under negative pressure . (B-C) System 0 cage design and particle

movement within cage system while under negative pressure (-0.06 WC). (D-E) Addition

of unidirectional airflow in system 1 design and particle transport within system 1 under

negative pressure (-0.03 WC). Dashed line indicates unsealed cage lid. (F-G)

Schematic of system 2A with airflow driven by the exhaust port (2.5 L/min) and particle

transport under negative pressure (-0.03 WC) conditions. (H-I) System 2B schematic

with modified cage lid sealing (solid line) and single cage barrier and particle transport

within system 2B. All particle transport results are represented as number of particles

per cubic centimeter of air (#/cc) in the size range of 0.8-7 μm. Sampling includes

upstream (red) before entering the cage, in the donor compartment (blue) and the

recipient compartment (black).

Figure 4. Optimization of environmental factors which influence transmission

efficiency. (A) Recovery of particles containing *M. smegmatis* from the recipient

compartment using a low-flow button sampler at varying exhaust flow rates (0.15, 1.5

and 2.5 L/min). (B) Nebulization of *M. smegmatis* in the donor compartment and

recovery of bacteria from lungs of recipient guinea pigs in system 2B cages. Colony

forming units (CFU) quantified from lung homogenates plated across four agar plates

from four different guinea pigs. Each point represents CFU on an individual plate. (C)

Humidity accumulation during guinea pig housing in system 2B cages. Relative humidity

(%) in three system 2B cages with guinea pigs housed in each compartment (two per

cage) across 6 days. Three different exhaust flowrates tested include 0.15, 1.5, and 2.5

L/min. Cages were changed at day 4 for all flowrates.

Figure 5. Particle tracer simulations, validated against experiments, reveal

heterogeneous aerosol distributions and the influence of BSL-3 containment on

particle retention. (A-B) Schematics of the donor-recipient cage system under ambient

and negative-pressure surrounding laboratory conditions, showing airflow pathways and

tracer injection and exhaust locations. The naïve guinea pig breathing zone (BZ) is

defined as the cage volume below the animal's typical breathing height of approximately

7 cm. (C-D) Comparison between computed and experimentally measured proxy for

particle transport, quantified as the ratio (R/U) of tracer concentration in the recipient

compartment (R) to upstream injected tracer concentration (U), shown for ambient (C)

and negative-pressure *=negative pressure-induced effective ACH (D) laboratory

conditions. Simulation values represent averages across four locations within the

recipient-side BZ, capturing spatial variation in predicted particle concentrations;

individual location-specific contributions are shown in Supplemental Fig. 2A-B.

Experimentally measured R/U values are derived from single-location particle

measurements and summed over particle size. Air-change rates per hour (ACH) for

each configuration are indicated for ambient conditions. (E-H) Complementary

animations for ambient and negative-pressure conditions are shown in Supplemental

Movie 2 and 3

Figure 6. Buoyancy-resolving simulations quantify recipient exposure. (A)

Definition of the recipient inhalation layer used to compute mean exposure ($\mathcal{C}$) (see

Supplemental Methods). (B) Typical normalized time averaged exposure, $\mathcal{C}$, of the

recipient guinea pig for systems 0, 1 and 2B under negative pressure. (C)

Representative mid-plane normalized tracer concentration field (c/c_0), where c_0 is the

donor emission concentration (darkest red).

Figure 7: Transmission of Mtb in an optimized cage system. (A) Experimental

timeline for 17-week transmission study. Blue ticks indicate when new donor groups

were added to the cages. Monthly TSTs using a two-step strategy are noted as filled

stars (first dose) and open stars (second dose). (B) TST measurements for recipient

animals throughout the exposure period. Any responses above 6.5 (dashed line) are

considered positive. (C) Predicted probability of Mtb exposure based on serum antigen

85B and PPD IgG levels. Values above 0.5 (dashed line) are predicted positive in the

donor, naïve, recipient and TST control groups. (D) Quantification of stage 1

granulomata in naïve and recipient lung sections. (E) Representative hematoxylin and

eosin staining of lung section from naïve, donor and recipient groups. Inflammatory

collection (naïve) and stage 1 granuloma (recipient) at 20x. p-values calculated by one-

way ANOVA for multiple comparisons or Mann-Whitney for comparison between two

groups. ns=not significant, *=p<0.05, **= p<0.01, ***=p<0.001, ****=p<0.0001.

1079

1080

Figure 1

A

Aerosol infection

Donor GPs

Recipient GPs

Month 0 1 2 3 4 5 6 7 8 9 10 11 12

TST ★ ★ ★ ★ ★ ★ ★

B

System 0

C

D

System 1

Exhaust 2.5LPM   Supply 8.8 LPM

E

F

Figure 2

A Exhaust vacuum — Room Pressure — Supply air

System 0-2B

Recipient — Donor

APS Particle sampler — Particle generator

B System 0

C Upstream / Donor / Recipient
Particles counted #/cc vs Particle size (µm)

D System 1
Exhaust 2.5LPM / Supply 8.8 LPM

E Upstream / Donor / Recipient
Particles counted #/cc vs Particle size (µm)

F System 2A
Exhaust 2.5LPM / Supply 0 LPM

G Upstream / Donor / Recipient
Particles counted #/cc vs Particle size (µm)

H System 2B
Exhaust 2.5LPM / Supply 0 LPM

I Upstream / Donor / Recipient
Particles counted #/cc vs Particle size (µm)

Figure 3

Figure 4

Figure 5

A

Sys 1/2A/2B: 2.5 L/min
Outlet

Sys 1: 8.8 L/min
Inlet

Vertical height of the inlet tube and counter probe

Varying location of sampling

Breathing Zone (BZ)

APS Particle Counter Probe 0.3 L/min Suction

Donor

Tracer Inlet Tube 0.3 L/min Injection

B

Sys 1/2A/2B: 2.5 L/min
Outlet

Sys 1: 8.8 L/min
Inlet

Recipient

Donor

C

Room Pressure

Concentration ratio R/U

● Simulation at avg tube location
◇ Experiment

Sys0 ACH 0.3 · Sys1 ACH 8 · Sys2A ACH 2 · Sys2B ACH 2

D

Negative Pressure

Concentration ratio R/U

● Simulation at avg tube location
◇ Experiment

Sys0 ACH 20* · Sys1 ACH 10* · Sys2A ACH 10* · Sys2B ACH 2*

E System 0

F System 1

G System 2A

H System 2B

0.005   0.002   0.05   0.1   0.2   0.5   1

Concentration field normalized by the donor exhalation concentration
c/c_0

Figure 6

A Time-averaged normalized tracer concentration $\mathscr{C}$ in the guinea pig inhalation layer

$$\mathscr{C} = \frac{1}{T} \int_0^T \bar{c}(t)/c_0 \; dt$$

T: Exposure time
t: time
c_0: Donor breath concentration

B $\mathscr{C}$ over a typical hour of exhalation

Simulation based ACH values

C

Concentration field normalized by the donor exhalation concentration
c/c_0

Figure 7

A

Donor GPs
Recipient GPs
Week   0   2   4   6   8   9   10   12 13   14   16 17   18
TST   ★   ★   ☆   ★   ☆   ★   ☆

Aerosol infection

B

TST diameter (mm)

****
****
****
****
**

Donors   Naive   Recipients   TST Control

C   Ag85B and PPD IgG

Predicted probability

****
****
****
ns
*
**

Donors   Naive   Recipients   TST control

D

Stage 1 Granulomata

**

Naive   Recipients

E

| Naive | Donor | Recipient |

10 mm   10 mm   10 mm   10 mm

100 µm   100 µm   100 µm   100 µm