TITLE

Bumblebee queens differ in brain morphology but not learning performance across life stages

AUTHORS

Leeah I Richardson[1], Shalene Jha[1,2], *Christopher M Jernigan[3,4], and *Felicity Muth[1,5]

[1] University of Texas at Austin, Department of Integrative Biology

[2] University of Texas at Austin, Ladybird Johnson Wildflower Center

[3] Wake Forest University, Department of Biology

[4] Cornell University, Department of Neurobiology and Behavior

[5] University of California Davis, Department of Neurobiology Physiology & Behavior

*co-last authors

ABSTRACT

Animal cognition and morphological investment across brain regions can vary between
individuals and across a lifetime as a function of individual social and ecological requirements.
Bumblebees have distinct ecological pressures acting upon individuals at different times: young
queens (gynes) and workers share similar challenges as they both forage as part of the colony,
but only queens overwinter and start a colony the following season, switching to a reproductive
stage once established in a (dark) colony. Past work shows that gynes are better at visual
learning than workers, but we know little about this varies between a foraging (gyne) stage and
reproductive (queen) stage. Here we compared bumblebee (*Bombus impatiens*) visual learning
and brain morphology across age-matched naive workers and gynes, and older reproductive
queens. We expected foraging-stage gynes to be better at visual learning than reproductive
queens and visual regions to be reduced in the latter stage, in line with past work in other social
insects. However, we found that gynes and queens performed similarly, while both performed
better than workers. We also found that reproductive queens had larger olfactory regions
(antennal lobes) than gynes, while visual regions (medulla and lobula) did not differ, indicating a
greater investment in olfactory regions in the reproductive stage. Brain regions scaled differently
across gynes and workers with body size, suggesting that gynes are not simply 'large workers'
in terms of their neuromorphology. Overall, our results provide behavioral and morphological
evidence that social and ecological roles in a colony shape cognition and neural investment.
30
31

INTRODUCTION

An animal's cognition reflects both environmental selection pressures and individual experience, and can vary both between individuals and within individuals as their needs change over time (Shettleworth, 2009; Smid and Vet, 2016). For example, desert locusts (*Schistocerca gregaria*) undergo a distinct change in phenotype from a solitary to a gregarious phase, associated with changes in morphology and physiology but also their cognition: solitary forms are capable of aversive taste learning while gregarious forms are not, despite both being capable of appetitive learning (Simões et al., 2013; Simões et al., 2016). These differences in cognition align with differences in observed feeding behavior: gregarious forms readily consume chemically-defended plants while solitary forms do not. Changes in behavior across an animal's lifetime can also be associated with neural changes: food-storing black-capped chickadees (*Poecile atricapillus*) show seasonal changes in spatial memory performance and hippocampal volume, a brain region critical for spatial cognition, in line with foraging demands (Sherry and Hoshooley, 2010). Such changes are expected because neural structures are functionally specialized (Mantini et al., 2013; Tanaka et al., 2012) and metabolically costly to maintain (Niven and Laughlin, 2008). As a result, cognition and its neuromorphological underpinnings are expected to dynamically reflect ecological demands both between and within individuals (Gronenberg and Riveros, 2009) coined as 'neuroecological theory' (O'Donnell et al., 2014).

Foraging can be a cognitively demanding behavior and can drive cognitive specialization and associated neuromorphology. For example, frugivorous primates have larger brains than folivores, which is thought to be related to their more cognitively complex foraging requirements (DeCasien et al., 2017), and seed-caching birds that are dependent on spatial memory for retrieving caches have better spatial cognition and larger hippocampi compared to non-seed caching species (Clayton, 1998). Even within a seed-caching species (*Poecile gambeli*), birds from populations that seed-cache in harsher environments have better spatial memory and larger hippocampi compared to lower elevation populations (Freas et al., 2012). Honeybees and bumblebees have long been used to study foraging behavior because foraging dominates much of foraging workers' ('foragers'') lives (Heinrich, 1979). As generalist nectarivores visiting a wide variety of flowers, they are excellent at learning visual and olfactory associations with floral rewards (Chittka and Thomson, 2001) and much is known about both their learning and its neural underpinnings (Gronenberg and Riveros, 2009; Menzel, 2021; Menzel and Giurfa, 2001), making them an ideal system for understanding how foraging relates to brain morphology. For example, the medulla and lobula process visual information peripherally (Kenyon, 1897; Paulk and Gronenberg, 2008; Paulk et al., 2009), while the antennal lobe processes olfactory cues peripherally including floral volatiles (Krofczik, 2008) and socially relevant cuticular hydrocarbons (CHCs) (Sharma et al., 2015). A major brain region that integrates both visual and olfactory information in the central brain and is known to be involved in learning and memory is the mushroom bodies (MBs), with the MB collar receiving primarily visual inputs (Strausfeld et al., 1998) and the MB lip receiving primarily olfactory inputs (Ehmer and Gronenberg, 2002; Gronenberg, 1999; Mobbs, 1982). While past work on workers suggests that the bumblebee brain shows plasticity dependent on age and foraging experience (Jones et al., 2013; Riveros and Gronenberg, 2010), we know much less about one of the most important castes: queens. Queens offer the opportunity to study how cognition and the brain can change with behavioral

76 requirements, since they transition from a foraging stage to a reproductive stage after starting a
77 colony.
78
Gynes are produced at the end of the colony lifecycle, where they forage as part of the colony
(Alford, 1969; Goulson, 2010); they then mate and are the sole colony members that overwinter.
Gynes then emerge in the spring and forage as solitary individuals during nest-searching and
initiation. Foraging gynes face similar challenges to foraging workers in terms of needing to
learn which flowers offer the best rewards, although gynes are better at learning color
associations than workers (Muth, 2021). After they find a nest site and start a colony, the
queens cease foraging and enter a reproductive phase once their first batch of foraging workers
are produced. In their reproductive phase, queens develop their ovaries, produce offspring for
the colony, and remain colony-bound in the dark for the rest of their lives (Goulson, 2010).
However, it is unclear whether queen cognition and/ or brain morphology changes between the
foraging phase and reproductive phase of their lifecycle. More broadly, ontogenic shifts from
free-living to colony-bound reproduction in insects have been shown to coincide with changes in
brain investment: in ant queens (*Messor pergandei* and *Pogonomyrmex rugosus*), peripheral
visual regions (medulla) shrink once individuals begin nest-founding (Julian and Gronenberg,
2002), and in another species, *Harpegnathos saltator* colony-bound reproductives had smaller
visual regions compared to non-reproductive foragers (Gronenberg and Liebig, 1999; Penick et
al., 2021). Similarly, in paper wasps (Polistinae), nest-bound foundress queens invest less in
visual regions relative to olfactory regions compared with workers (O'Donnell et al., 2014), but it
is not known if similar differences in investment exist for bumblebee workers, gynes, and
reproductive queens.
In this study we addressed inter- and intra-caste variation in cognition and neuromorphology in
bumblebees. Specifically, we asked whether queen visual learning and neuromorphology
differed between a foraging ('gyne') and reproductive ('queen') stage in line with ecological
requirements, i.e. in line with 'neuroecological theory' (Gronenberg and Riveros, 2009;
O'Donnell et al., 2014). In addition, we compared age- and experience-matched gynes and
workers to ask whether these groups differ in learning performance and in the relative size of
sensory (peripheral) and sensory integrating (central) brain regions. While gynes have been
found to be better at learning than workers (Muth, 2021), it is unclear how queen learning
abilities change across their lifetime. It is also currently unknown how gyne and queen brains
scale relative to each other and to those of workers. If queen cognition and neuromorphology
are associated with their ecological role, then we predict that foraging-stage gynes will perform
better at a visual learning task than older, reproductive-stage queens. In addition, we predicted
that gynes would have greater investment in visual regions of the brain relative to reproductive
queens, specifically the medulla, lobula, and mushroom body collar relative to brain regions that
primarily process olfactory information (antennal lobes and mushroom body lip). To ask if
queens differ from workers, we compared age- and experience-matched gynes to workers. We
expected gynes to learn better based on past findings (Muth, 2021). We then compared
investment in the sensory and sensory-integrating regions of the brain. Specifically, we
measured the volume of three regions involved in sensory processing in the peripheral brain
across both vision (medulla, lobula) and olfaction (antennal lobe) and two subregions involved in
sensory integration and learning in the central brain, one primarily receiving visual inputs (MB

collar) and the other olfactory (MB lip), to determine if they scaled similarly relative to body size
across groups (Table 1).

METHODS

Colony maintenance and bee selection

We tested 70 bees across 16 commercial bumblebee (*Bombus impatiens*) colonies (Koppert,
USA) (Table S1 with sample sizes per colony per assay). Colonies were kept in the boxes that
they were received in from Koppert, elevated to allow for airflow and maintained on
approximately 3g of honeybee-collected pollen (Koppert, USA) and 30% (w/w) sucrose,
replenished 3 times per week. With the exception of when bees were marked for identification,
colonies were also kept in dim-light conditions by placing a cardboard lid on top of each colony.
We periodically identified all newly eclosed adult bees (workers or gynes) by their whitish
unsclerotized appearance and put a small water-based paint mark (Posca, USA) on their
thoraces for later identification. Once workers and gynes were 7-10 days old they were removed
from the colony to be tested in a color learning assay (total n = 63; workers = 35; gynes = 28;
Table S1). We age-matched workers and gynes to control for effects of age-related changes
(i.e. as in Jernigan et al., 2019). Older queens in a reproductive stage (hereafter 'queens') were
removed from the colony and underwent the color learning assay once we had finished testing
gynes and workers from the same colony, such that all tested bees were taken from a
queenright colony (i.e., we did not sample from colonies after the original queen had been
removed). This resulted in a total of 7 queens tested; given that we were targeting queens from
late-stage colonies that were producing gynes, we did not test queens from the 9 remaining
colonies because they died before we were able to test them. As commercially reared queens
are unlikely to have free-flying foraging experience including visual foraging experience (Huang
et al., 2015), we ensured that all bees did not forage and instead remained in the colony for the
duration of their lives. As such, we believe that all bees tested (workers, gynes, and queens)
were naïve to visual stimuli and did not have foraging experience. We imaged brains from a
subset of bees that underwent the color learning assay (n = 26: workers = 10; gynes = 10;
queens = 6; Table S1).

Color learning

We conducted a free-moving proboscis extension response (FMPER) color discrimination assay
to assess differences in visual learning performance between lab-reared workers, gynes, and
queens (first described in Muth et al., 2018, updated in (Muth, 2021). We placed bees in testing
tubes (2.5cm x 2.5cm x 15cm) for 2hr 40min to acclimate and become sufficiently hunger-
motivated before training them to discriminate between two shades of human-blue, blue and
aqua (same colors as in Muth, 2021, training color counterbalanced across groups; Fig. S2A).
Training consisted of three phases: 1) two "no choice" learning trials, 2) four "choice" learning
trials, and 3) six unrewarded "test" trials (Fig. 1A). For the no choice trials, we first presented the
rewarding color (rewarded conditioned stimulus: CS+) dipped in 50% (w/w) sucrose solution to
the bee and allowed her to drink for 3s, and then presented the unrewarding color (unrewarded
conditioned stimulus: CS-) dipped in water and let her drink or antennate for 3s (bees frequently

did not consume the water) (Muth, 2021; Muth et al., 2018). For the choice trials we presented
the CS+ and CS- simultaneously such that the bee had to approach and choose one of the
stimuli (by extending her proboscis or by antennating it, given that bees can taste via their
antennae, Bitterman et al. 1983)). After making her first choice (denoted as 'correct' or
'incorrect'), we then allowed the bee to sample the other option, so that for each learning trial all
bees experienced both the rewarding stimulus and the unrewarding stimulus (Muth, 2021; Muth
et al., 2018). We planned the inter-trial interval (ITI) for all six learning trials (2 no choice and 4
choice trials) to be approximately 10 minutes (8-12 min) for all groups, however since the bee
ultimately controlled the timing, we measured the actual ITI and found that queens took longer
than workers (mean of 12.8 mins vs. 9.2 mins), but that gynes (mean of 10.3 mins) did not differ
from workers or queens ($X^2(2) = 7.319$, $p = 0.025$; Tukey post-hoc comparisons between
groups: Queens vs. workers $t = 2.566$, $p = 0.033$; queens vs gynes: $t = -1.717$, $p = 0.207$; gynes
vs workers: $t = 1.472$, $p = 0.311$).

For the unrewarded test trials, the two colors were again presented simultaneously for the bee
to choose between, but both strips were dipped in water, and were removed just before the bee
contacted the chosen strip; we did this to motivate bees to make multiple choices and to limit
learning that strips were unrewarded in the test phase (i.e. extinction learning). These 6 trials
were presented such that the bee could make successive choices within a few seconds (i.e. the
total test trial lasted an average +/- SD of 5.90 +/- 6.84 min; mean per group: queens = 8.00
min, gynes = 7.45 min, workers = 4.50 min). If a bee did not move for 12 minutes during the
assay, she was presented with a gray strip of paper dipped in sucrose solution and was allowed
to drink for 3s to re-motivate her to participate; we excluded bees due to lack of participation if
they continued to remain stationary (for 12 mins) after gray strips were presented twice (*sensu*
Muth et al., 2021); bees across the three groups did not differ in their motivation to participate in
the task ($X^2(2) = -4.425$, $p = 0.109$), with 5/7 queens, 25/28 gynes, and 34/35 workers
participating. The two blue colors were more chromatically similar to each other than to the gray
strips when plotted in bee color space (Fig. S2B), modeled from *B. impatiens* photoreceptor
sensitivity (Gawryszewski, 2018; Skorupski and Chittka, 2010) and presentation of gray strips
did not affect test trial performance across bee types (Fig. S1C).

For each bee we measured intertegular distance (ITD) (Cane, 1987; Hagen and Dupont, 2013),
marginal cell length (Medler, 1962), and hind tibia length (Jernigan et al., 2021) as proxies for
body size, by taking photographs under magnification (AmLite, USA) and then measuring each
in FIJI (Schindelin et al., 2012). We calculated a PC1 score for each individual from these
metrics (Fig. S2) as our measure of body size (Jernigan et al., 2021) to account for body size
differences when making comparisons between brain region volumes. Body size did not differ
between queens and gynes either for those used in the behavioral assays ($t(58) = -0.331$, $p =$
0.941), or in the subset that underwent brain imaging ($t(23) = 0.097$, $p = 0.995$). As expected,
workers were smaller than gynes and queens in both the full data set (queens vs workers: $t(58)$
$= 9.226$, $p < 0.001$; gynes vs workers: $t(58) = 16.157$, $p < 0.001$) and in the subset used for
brain imaging (queens vs workers: $t(23) = 5.798$, $p < 0.001$; gynes vs workers: $t(23) = 9.307$, $p <$
0.001).

Dissections and staining

Following participation in the color learning assay, bees that were part of the subset for brain imaging were immediately cold-anaesthetized at 4.4C (workers: n = 10, although due to damage during processing one of these brains could only be used for medulla and lobula measurements so is only included in those analyses; gynes: n = 10; queens: n = 6). Once immobilized, bees were decapitated and brains were dissected out of the head capsule in cold PBS (Fig. S3A), before being stored overnight in a 4% paraformaldehyde solution at 4°C. We then conducted a staining protocol consisting of three fluorophores in order to visualize neuropils of interest (modified from Jernigan et al., 2019 to include DAPI as a fluorophore in addition to phalloidin TRITC and anti-synapsin Alexa488 and increased incubation times to 5 days to ensure fluorophore penetration in the larger brains; Fig. S3B).

After sitting overnight in the paraformaldehyde fixative, we rinsed the brains twice in PBS, then 6x in PBSTX (1X PBS with 1% Triton X-100 v/v) for 20 min per rinse, and added 1uL of our primary anti-synapsin antibody synorf (mouse anti-fruit fly synapsin, 3C11 DSHB 1:800 dilution) with 10uL of normal donkey serum, and left them to incubate on a shaker for 5 days. We then rinsed the brains 6 times in PBSTX for 20min each rinse, and added 2.5uL of our secondary antibody donkey anti-mouse Alexa488 (Jackson Immuno, USA), along with 10uL of phalloidin conjugated with TRITC (Thermo Fisher, USA), and 1.5uL of the nuclear stain DAPI (Millipore Sigma, USA). Brains were left on the shaker to incubate for an additional 5 days. We then rinsed brains with PBS 6x for 10 min per rinse, followed by a 10 min rinse with 4% paraformaldehyde, and then two more 10 min rinses with PBS. We then conducted an alcohol dehydration, spending 20min at each concentration in ascending order (30%, 50%, 70%, and 95% ethanol). We kept the brains in 100% ethanol at 4°C for 24 hours, and replaced this with another aliquot of 100% ethanol to store in a -20°C freezer prior to imaging.

Imaging and reconstructions

We cleared the brains by submerging them in methyl salicylate for >15min (up to 1 hour) and conducted whole mount z-stack imaging of one hemisphere with a Zeiss 710 laser scanning confocal microscope with 10X air objective and 12 microns between slices with 16x bidirectional averaging to reduce noise. Imaging was performed at the Center for Biomedical Research Support Microscopy and Imaging Facility at UT Austin (RRID:SCR_021756). If one hemisphere did not fit into the frame, two z-stacks were imaged and then tiled together using the 3-D stitching plugin in FIJI (Preibisch et al., 2009; Schindelin et al., 2012). Brain reconstructions were completed by manually tracing neuropils for each region of interest (*sensu* Jernigan et al., 2021) using Amira software (Thermo Fisher, USA).

We reconstructed three peripheral sensory processing brain areas: two visual processing brain regions (medulla and lobula) and the primary olfactory processing region of the brain (antennal lobe). We also subdivided two regions of the mushroom body calyces known to integrate information in the central brain, one integrating visual information (mushroom body collar) and one olfactory information (mushroom body lip) (Fig. 2; Table 1).

Data analysis

Performance across learning trials: To test for the effect of bee type (worker, gyne, or queen) on color discrimination performance during the four learning 'choice' trials, we used a generalized linear mixed model (*glmer*) with a binomial distribution (Bates et al., 2015) with the interaction between bee type (queen, gyne, or worker) and trial number (1-4: continuous) as fixed factors, color the bee was trained to (blue or aqua) as a fixed factor, and individual bee ID as a random factor to account for multiple measures per individual.

Learning test performance: To test for the effect of bee type (queen, gyne, or worker) on performance in the test phase, we included bee type and the color the bee was trained to as fixed factors and individual bee ID as a random factor. Test trial number was not included in models (as in Muth, 2021; Muth et al., 2018) since test trials were designed to be equivalent and bees' behavior was not expected to change over time as we restricted sampling during these trials. We determined fixed factor significance with *Anova* in the *car* package (Fox and Weisberg, 2019). For significant terms we used the *emmeans* function in the package *emmeans* (Lenth, 2017) to conduct Tukey posthoc pairwise comparisons within factors. We also confirmed that each group had learned their trained discrimination (performance above chance at 50%) with a one tailed one sample t-test for each group, with the alpha value adjusted to 0.017 with a Bonferroni correction due to the three comparisons (Bonferroni, 1936).

Brain morphology analyses: To determine if bees (queen, gyne, or worker) varied in a number of brain region volumes (total constructed volume, medulla, lobula, antennal lobe, mushroom body collar, and mushroom body lip), while accounting for body size scaling (bumblebee worker brain volume scales allometrically (Mares et al., 2005) we used standard major axis regressions (*sma*) which are commonly used to understand scaling relationships between measures (including work with bees, Cariveau et al., 2016; Yamhure-Ramírez et al., 2025) with bee type (categorical with 3 levels: queen, gyne, or worker) and PC1 of body size (continuous) (similar to Jernigan et al., 2021) as predictors of region volume. These analyses had two parts; first: we tested if the way each region scaled with body size varied by bee type by comparing slopes. Second: for bee types that had the same slope, we then tested for differences in intercept (Table 1) to assess if their investment in that brain region was predicted by their body size or if they invested relatively more or less (Warton et al., 2012). We used PC1 as our primary measure of body size since it accounts for multiple metrics of body size variation (Jernigan et al., 2021), but our main results are the same even when we use only ITD (Table S2), another commonly used body size metric in bee biology (Hagen and Dupont, 2013).

To test for differences in relative investment in visual versus olfactory region brain volumes, we ran two linear models, one for peripheral processing and one for central processing. For peripheral processing we used the ratio between the two relevant visual regions (medulla and lobula) and the antennal lobe. For central processing we measured the ratio between the mushroom body collar and mushroom body lip. Relative measures between regions are common in these types of comparisons, and body size was not included as a factor in the models testing differences in relative investment (Jernigan et al., 2021; O'Donnell et al., 2014).

Because the absolute size of brain regions, irrespective of body size, can also be informative of behavioral/cognitive differences (Marino, 2006), we also ran all analyses without accounting for

body size to understand absolute differences in brain region volume across our bee types
(results shown in Supplementary Material; Table S3).

RESULTS

Color learning

All groups of bees learned the color they were trained to, performing above 50% chance level:
queens chose the rewarding color 90.0% of the time in the test trials (one-sample t-tests: t(4) =
6.000, α = 0.017, p = 0.001), gynes 79.3% of the time (t(24) = 9.507, α = 0.017, p < 0.001), and
workers 65.2% of the time (t(33) = 5.182, α = 0.017, p < 0.001). Queens and gynes made
similar numbers of errors in the test phase, but both made fewer errors than workers in the test
phase (*glmer*; full model, type: $X^2(2)$ = 12.576, p = 0.001; color: $X^2(1)$ = 0.046, p = 0.829; post-
hoc comparisons: queens vs workers: z = 2.486, p = 0.034; gynes vs workers: z = 2.500, p =
0.011; queens vs gynes: z = -1.318, p = 0.384; Fig. 1B). The three groups of bees did not differ
from each other in performance across the learning choice trials (*glmer*; type: $X^2(2)$ = 0.004, p =
0.997), nor did they improve over the trials (*glmer*; trial: $X^2(1)$ = 0.191, p = 0.661; type*trial:
$X^2(2)$ = 0.866, p = 0.648; Fig. S4). Bees' performance also did not differ based on the color they
were trained to (*glmer*; color: $X^2(1)$ = 2.837, p = 0.092; Fig. S1A).

Neuromorphology
301
In order to compare body size-corrected volumes of each brain region across our groups, we
first compared the slopes across our three groups to evaluate if the scaling between body size
and brain region volume varied by bee type. The total reconstructed volume scaling differed by
type (*sma*; LRS(2) = 6.860, p = 0.032); where gynes' total reconstructed volume scaled more
positively with body size than queens' (t(2) = 4.22, p = 0.039) and workers' (t(2) = 5.252, p =
0.021), while queens' did not scale differently with body size compared to workers' (t(2) = 0.061,
p = 0.804). Scaling also differed between bee types for the peripheral brain regions (*sma*;
medulla: LRS(2) = 7.704, p = 0.021; lobula: LRS(2) = 10.590, p = 0.005; and antennal lobe:
LRS(2) = 10.53, p = 0.005). However, scaling did not differ between bee types for the central
brain regions (*sma*; MB collar: LRS(2) = 2.177, p = 0.336; MB lip: LRS(2) = 1.900, p = 0.386).
312

For the peripheral visual regions, gyne medullas scaled more positively with body size than
worker medullas did (t(2) = 7.484, p = 0.006), but there were no other differences in scaling
relationships (queens vs workers: t(2) = 0.242, p = 0.622; queens vs gynes: t(2) = 2.463, p =
0.116). The lobula scaled more positively with body size for gynes than for workers (t(2) =
10.677, p = 0.001), but there were no other differences in scaling with body size for the other
comparisons (queens vs workers: t(2) = 3.483, p = 0.062; queens vs gynes: t(2) = 1.652, p =
0.198). Independent of body size scaling (comparing the intercepts), there were no differences
in medulla size between queens and gynes (*sma*; intercept: Wald $X^2(1)$ = 2.96, p = 0.085; Table
1), or between queens and workers (*sma*; intercept: Wald $X^2(1)$ = 0.633, p = 0.425; Table 1).
Independent of scaling, there was no difference in lobula size for the queens and gynes (*sma*;
intercept: Wald $X^2(1)$ = 1.177, p = 0.278; Table 1), but queens had relatively larger lobulas than
workers (*sma*; intercept: Wald $X^2(1)$ = 41.910, p < 0.001; Table 1).

For the peripheral olfactory region, we found that gyne antennal lobe volume scaled more
positively with body size compared to workers ($t(2) = 10.528$, $p = 0.001$), but there were no
other scaling differences between groups (queens vs gynes: $t(2) = 1.840$, $p = 0.174$; queens vs
workers: $t(2) = 2.006$, $p = 0.156$). Independent of body size scaling, queens had larger antennal
lobes than both gynes (*sma*; intercept: Wald $X^2(1) = 8.103$, $p = 0.004$; Fig. 2D; Table 1) and
workers (*sma*; intercept: Wald $X^2(1) = 83.530$, $p < 0.001$; Fig. 2D; Table 1).

For the central processing regions, we found that queens and gynes had larger mushroom body
collars, which receives visual input, than workers (*sma*; queens vs workers intercept: Wald $X^2(1)$
$= 19.048$, $p < 0.001$; gynes vs workers intercept: Wald $X^2(1) = 5.292$, $p = 0.021$; Table 1).
Independent of body scaling, queens, gynes, and workers did not different in volume (*sma*;
queens vs gynes intercept: Wald $X^2(1) = 0.677$, $p = 0.172$). Queens and workers did not differ in
size of the mushroom body lip, which receives olfactory input (*sma*; queens vs workers
intercept: Wald $X^2(1) = 0.968$, $p = 0.325$; Table 1), but gynes had a smaller mushroom body lip
than workers and marginally lower than queens (*sma*; queens vs gynes intercept: Wald $X^2(1) =$
3.595, $p = 0.057$; gynes vs workers intercept: Wald $X^2(1) = 4.574$, $p = 0.032$; Table 1) when
accounting for body size scaling.

Comparing visual vs olfactory regions, we found that the relative investment in peripheral visual
vs. peripheral olfactory regions varied by bee type (*lm*; $F(2) = 15.175$, $p < 0.001$; Fig. 2C) with
queens investing relatively less in peripheral visual regions (medulla + lobula) relative to the
peripheral olfactory region (antennal lobe) compared to gynes and workers (queens vs gynes:
$t(2) = 7.340$, $p < 0.001$; queens vs workers: $t(2) = -6.092$, $p < 0.001$), but there was no difference
in relative investment in peripheral brain regions between gynes and workers ($t(2) = 1.261$, $p =$
0.431; Fig 2C). Bee types did not differ in their relative investment of vision and olfaction in
central brain areas (*lm*; MB collar / MB lip: $F(2) = 2.304$, $p = 0.123$).

<u>DISCUSSION</u>

Neuroecological theory predicts that greater investment in specific brain regions and associated
cognitive abilities reflect ecological demands (Gronenberg and Riveros, 2009; O'Donnell et al.,
2014). Here we show that queens retain their visual learning capacities between a foraging and
older reproductive stage, while performing better than workers across both stages (as has been
found previously (Muth, 2021)). Despite this similarity in learning performance across stages,
differences in brain morphology were evident: reproductive queens showed a shift in modality-
specific peripheral investment in relative investment of vision vs olfaction, as expected.
However, while we expected visual regions would be smaller in reproductive queens relative to
gynes, instead we found evidence for increased investment in the peripheral olfactory region.

In the current study, we did not find that queen visual learning performance declined compared
to gynes despite the behavioral transition to a nest-bound reproductive role. One explanation is
that learning capacities are maintained in bumblebee queens to serve a function; colony
initiation is turbulent and can be disrupted by abiotic factors such as the length of the winter
(Ogilvie and CaraDonna, 2022) or due to other nest-founding queens usurping existing nests
(Richards, 1978). The ability to re-initiate nests after early stage disruptions varies across
species (Tripodi and Strange, 2019) and to our knowledge it is not known if queens are able to
re-initiate nests later in colony development, but future work could explore this possibility. While

we expected information processing to be modality-specific (Bitterman, 1965; Frost et al., 2015),
and thus visual learning to decline, queens' learning capacities could also be maintained
through domain general processes (Deaner et al., 2006). For example, queens need to maintain
dominance hierarchies with aggression in response to worker behavior (van Doorn and Heringa,
1986), monitor brood to make reproductive decisions (Peto et al., 2025), and process social
information broadly, as queens are the colony's social hub, interacting more with daughters than
they do with each other (Ruttenberg et al., 2025). It is also possible that older reproductive
queens decline in more subtle visual learning capacities that could be captured with a more
difficult color discrimination or more cognitively complex tasks, such as reversal learning. To
control for the lack of foraging experience the older reproductive queens had (since they were
commercially reared), we also restricted foraging experience in the gyne and worker groups.
However, since foraging experience is related to learning performance (i.e. see Riveros &
Gronenberg 2009), it is possible that with foraging experience, differences in learning capacity
between queens and gynes would become evident. Nevertheless, previous findings that gynes
are better at learning than workers (Muth, 2021) held in our experiment, indicating that at least
between castes, baseline differences in learning may play a larger role than experience.

Beyond differences in learning, we also found that queens, in particular, took longer to make
choices than gynes and workers, consistent with prior work showing that bumblebee female
reproductives are slower decision-makers (Evans and Raine, 2014a). This likely reflects
motivational differences, since reproductive queens would not be motivated to forage beyond
feeding themselves. It is also possible that these differences reflect a difference in foraging
strategy prioritizing accuracy over speed (Chittka et al., 2003), whereas workers may be more
willing to take risks in terms of sampling previously unrewarding options that could result in
finding new floral resources (Evans and Raine, 2014b) as a part of a social colony. Indeed, past
work shows that honeybee foragers have two distinct foraging phenotypes which differ in gene
expression (Liang et al., 2012) and the concentrations of tyramine and octopamine found in the
brain (Cook et al., 2019): exploratory "scouts" that quickly abandon unrewarding stimuli and
tend to prefer novel resources (Carr-Markell and Robinson, 2014; Cook et al., 2020) and the
persistent "recruits" that may be more risk-averse (Anselme, 2018). While these types of
differences in terms of distinct phenotypes that vary in neuromodulator concentrations have not
been explored in bumblebees, it is possible that similar distinct foraging behavioral repertoires
could be relevant to behavioral differences across bumblebee castes. Our findings suggest that
reproductive individuals (gynes and queens) may employ different foraging strategies than
workers, potentially of importance to their solitary role during colony initiation.

In accordance with our color learning results, we did not detect a reduction in queens' visual
region volume, despite previous work finding this pattern when gynes transition to colony-bound
reproductives in ants (*M. pergandei* and *P. rugosus*) (Julian and Gronenberg, 2002), and paper
wasps (O'Donnell et al., 2014), and reversibly when ants (*H. saltator*) are experimentally
manipulated to transition back and forth between colony-bound reproductive and foraging
stages (Penick et al., 2021). It is possible that queen brains do not change between the foraging
and reproductive stage – however, we found that the olfactory lobe was larger in reproductive
queens. As such, one possibility for the difference between our visual region results and past
studies is that we used naïve individuals. Bumblebee foraging experience is known to increase
the size of brain regions relevant to foraging (mushroom body volume) (Riveros & Gronenberg

2010) so it is possible that as gynes begin actively foraging, their brain regions may expand and
then later decrease in volume as they transition to reproduction, dependent on experience.
Alternatively, there could be general constraints on plasticity of the visual regions within *Bombus*
impatiens, as plasticity is known to vary across species (Murren et al., 2015; Touchon et al.,
2024) and even between populations of the same species (Gonda et al., 2011; Gonda et al.,
2013). For example, in honeybees, mushroom body size is known to increase as workers
transition from nurses to foragers, but this change is not reversed if a forager transitions back to
a nurse (Fahrbach et al., 2003). It is also possible that the cessation of foraging activity/ flight is
energy-saving enough to mitigate the costs of maintaining these visual regions since flight is
known to increase insect metabolic rate up to 100 fold (Kammer & Heinrich 1978). Past work
shows that when queens are required to learn associations before initiating colonies, this
experience does not impair nest initiation success even when resources are limited (Watrobska
et al., 2024) and our data suggests that maintaining the ability to learn is not costly for queens
but it is possible that resource limitation could impact the cost of maintaining neural tissue.

In contrast to the results for visual brain regions, we found that older reproductive queens
invested more in peripheral olfaction (antennal lobe volume) than the other two groups, even
when body size was accounted for in our models. Olfaction is important for kin recognition in
social insects (Van Zweden and d'Ettorre, 2010), and may play a role in how queens detect
worker-laid eggs (Endler et al., 2004), which queens will eat to prevent worker reproduction
(Zanette et al., 2012). Olfactory information is likely especially relevant while residing in a dark
colony with limited visual information; and visual deprivation is known to relate to expansion of
the antennal lobe (Jones et al., 2013). In this study, we assessed differences in learning and
differences in brain morphology, but we note that synaptic brain volume is not the only measure
of neural investment and organization (Logan et al., 2018) and future work could further explore
metrics that may be more sensitive like synaptic density (Kraft et al., 2019), dendritic complexity
(Dobrin et al., 2011), or gene expression (Santos et al., 2022).

We also found that queens and gynes showed greater investment in the mushroom body collar
(associated with central visual processing, Strausfeld et al., 1998) than workers when controlling
for body size. The mushroom body expands as honeybee workers transition into a foraging role
(Fahrbach et al., 2003) which is an experience-expectant change that occurs as bees age even
in the absence of foraging experience (Fahrbach et al., 1998), although in bumblebee (*B.
occidentalis*) workers this expansion appears to be more dependent on experience than age
(Riveros and Gronenberg, 2010). The mushroom body is involved in learning and multi-sensory
integration (Davis, 1993; Menzel, 2014), and more broadly its volume correlates with learning
and sensory sensitivity (Serway et al., 2020). As such, queens' proportionally larger mushroom
body collar region may relate to their higher learning performance.

Overall, we found that queens and gynes performed better in a color discrimination task, and
invested relatively more in most brain regions (aside from the medulla and lobula) than would be
predicted purely by body size. We would also expect gynes and queens to be more sensitive to
visual and olfactory information because of their larger size, correlated with eye and antennal
size and features (eyes: Spaethe and Chittka, 2003; Taylor et al., 2019; antennae: Spaethe et
al., 2007), which in turn are correlated with visual and olfactory brain region volumes,
respectively (visual: Gronenberg and Couvillon, 2010; olfactory: Beltz et al., 2003). Our data

support the idea that, in terms of brain morphology, queens and gynes are not simply larger
workers but instead that brain development is distinct across castes (Cnaani and Hefetz, 2001).
Our finding that queens' and gynes' mushroom body collars were disproportionately larger than
those of workers' agrees with broader work on the distributed cognition hypothesis. This
hypothesis suggests that larger social insect groups allow for individuals to specialize on tasks,
and thus the larger the group size, the smaller we would expect individuals' brains to be
(Kverková et al., 2018; O'Donnell et al., 2015). This hypothesis is supported by fungus-farming
ants with group size being inversely related to brain size across fungus-farming ant species
(tribe: Attini) (Riveros et al., 2012), and solitary wasps having larger mushroom bodies (involved
in learning and memory) than closely related social species (family: Vespidae) (O'Donnell et al.,
2015). Similarly, workers within a bumblebee colony may be able to divide tasks and specialize
more than afforded by queens which have a solitary phase. In contrast to the distributed
cognition hypothesis, the social brain hypothesis suggests that species in larger groups have an
increased necessity for larger brains to track interactions and relationships (Dunbar, 1998).
While there is evidence for this hypothesis in taxa like primates (Dunbar and Shultz, 2007) there
is more limited support in insects (Lihoreau et al., 2012), including in primitively eusocial bees
(Pahlke et al., 2021; Smith et al., 2010) and locusts (Ott and Rogers, 2010). As bumblebee
castes are developmentally distinct and cannot be reversed in adulthood (Cnaani et al., 1997)
social hierarchies may be less important to track than in species with flexible hierarchies which
could explain why distributed cognition is able to be be relevant to their colony structure.

In summary, our results provide nuanced support for the brain reflecting ecological requirements
(i.e. the neuroecological hypothesis); rather than a reduction in visual brain regions, we found
an increase in antennal lobe volume in older reproductive queens, suggesting a shift toward
enhanced olfactory processing. This finding is potentially tied to queens' role within the dark,
socially complex nest environment, but could also be related to the lack of visual input all bees
experienced. Interestingly, we do not find evidence for increased investment in olfaction
necessitating a trade-off with maintenance of visual regions as we do not see a reduction to
visual regions in queens compared to gynes. Additionally, the higher learning performance of
gynes and queens relative to workers suggests the possibility of selective pressures on
reproductive females to forage efficiently and with minimal risk during the solitary phase.
Together, these findings point to a complex relationship between sensory investment and
ecological function between bumblebee castes and throughout the lifespan of the queen. Going
forward, experimental work could investigate free-foraging bumblebees, sampling at multiple
time points throughout their lifecycle to quantify the shifting role of experience (social and
foraging) on learning and brain morphology.

Figure 1 A: Color learning methods depicting a bee trained to aqua as the rewarding conditioned stimulus (CS+) in three parts. Rewarded stimuli were paired with 50% (w/w) sucrose solution, and unrewarded stimuli (CS-) were paired with water. First, two no choice training trials; second, four rewarded choice trials; finally, 6 unrewarded test trials in which the bee does not make contact with either stimulus. **B:** Performance by each group during the 6 unrewarded test trials, depicting the significant differences between workers and the other two groups with letters. Each grey point represents the proportion of correct choices a single bee made in the test trials, points are jittered for clarity with the group mean +/- stand error depicted in color. Correct choices are defined as choices to the CS+. The grey dotted line represents performance at chance (50%).

Figure 2 A: Brain regions traced on one slice laid over a bee head to display anatomic relationships **B:** Schematic showing 3D reconstructions - blue tints label olfactory regions (dark blue is antennal lobe; light blue is mushroom body lip) and red tints label visual regions (dark red is medulla and red is lobula; light red is mushroom body collar). **C:** Relative investment between peripheral visual regions (medulla and lobula) and the peripheral olfactory region (antennal lobe) with group means and standard errors. **D:** Natural log of antennal lobe volume (cubic microns) across groups, showing that queens have larger antennal lobes even when accounting for how volume may scale with body size with 95% confidence intervals about the slopes.

Table 1: Reconstructed brain regions describing which modality (visual vs olfactory) and type of processing (peripheral vs central) they are associated with. Names (including abbreviations) and a 3-D reconstruction of each region are displayed. We also show a summary of the differences in region volume (accounting for body size scaling; significant differences in intercept are shown) we find across groups.

Modality	Processing	Region	Reconstruction	Volume Differences
Olfaction	Peripheral	Antennal lobe (AL)		Queens > Gynes Queens > Workers
	Central	Mushroom body (MB) lip		Workers > Gynes
Vision	Peripheral	Medulla (MED)		No differences independent of body size scaling
	Peripheral	Lobula (LOB)		Queens > Workers
	Central	Mushroom body (MB) collar		Queens > Workers Gynes > Workers
Whole reconstructed volume				Queens > Workers

REFERENCES

Alford, D. V. (1969). A Study of the Hibernation of Bumblebees (Hymenoptera:
Bombidae) in Southern England. *J. Anim. Ecol.* **38**, 149.

Anselme, P. (2018). Uncertainty processing in bees exposed to free choices: Lessons
from vertebrates. *Psychon. Bull. Rev.* **25**, 2024–2036.

Bates, D., Maechler, M., Bolker, B. and Walker, S. (2015). Fitting Linear Mixed-Effects
Models Using lme4. *J. Stat. Softw.* **67**, 1–48.

Beltz, B. S., Kordas, K., Lee, M. M., Long, J. B., Benton, J. L. and Sandeman, D. C.
(2003). Ecological, evolutionary, and functional correlates of sensilla number and
glomerular density in the olfactory system of decapod crustaceans. *J. Comp. Neurol.*
455, 260–269.

Bitterman, M. E. (1965). Phyletic differences in learning. *Am. Psychol.* **20**, 396–410.

Bonferroni, C. (1936). Teoria statistica delle classi e calcolo delle probabilit.
Pubblicazioni R Ist. Super. Sci. Econ. E Commer. Firenze **8**, 3–36.

Cane, J. H. (1987). Estimation of Bee Size Using Intertegular Span (Apoidea). *J. Kans.*
Entomol. Soc. **60**, 145–147.

Cariveau, D. P., Nayak, G. K., Bartomeus, I., Zientek, J., Ascher, J. S., Gibbs, J. and
Winfree, R. (2016). The Allometry of Bee Proboscis Length and Its Uses in Ecology.
PLOS ONE **11**, e0151482.

Carr-Markell, M. K. and Robinson, G. E. (2014). Comparing Reversal-Learning
Abilities, Sucrose Responsiveness, and Foraging Experience Between Scout and Non-
Scout Honey bee (*Apis mellifera*) Foragers. *J. Insect Behav.* **27**, 736–752.

Chittka, L. and Thomson, J. D. (2001). *Cognitive ecology of pollination*. Cambridge
University Press, UK.

Chittka, L., Dyer, A. G., Bock, F. and Dornhaus, A. (2003). Bees trade off foraging
speed for accuracy. *Nature* **424**, 388–388.

Clayton, N. S. (1998). Memory and the hippocampus in food-storing birds: a
comparative approach. *Neuropharmacology* **37**, 441–452.

Cnaani, J. and Hefetz, A. (2001). Are queen *Bombus terrestris* giant workers or are
workers dwarf queens? Solving the "chicken and egg" problem in a bumblebee species.
Naturwissenschaften **88**, 85–87.

Cnaani, J., Borst, D. W., Huang, Z.-Y., Robinson, G. E. and Hefetz, A. (1997). Caste
Determination in *Bombus terrestris*: Differences in Development and Rates of JH
Biosynthesis between Queen and Worker Larvae. *J. Insect Physiol.* **43**, 373–381.

Cook, C. N., Mosqueiro, T., Brent, C. S., Ozturk, C., Gadau, J., Pinter-Wollman, N.
and Smith, B. H. (2019). Individual differences in learning and biogenic amine levels

influence the behavioural division between foraging honeybee scouts and recruits. *J.*
Anim. Ecol. **88**, 236–246.

Cook, C. N., Lemanski, N. J., Mosqueiro, T., Ozturk, C., Gadau, J., Pinter-Wollman,
N. and Smith, B. H. (2020). Individual learning phenotypes drive collective behavior.
Proc. Natl. Acad. Sci. **117**, 17949–17956.

Davis, R. L. (1993). Mushroom bodies and drosophila learning. *Neuron* **11**, 1–14.

Deaner, R. O., van Schaik, C. P. and Johnson, V. (2006). Do Some Taxa Have Better
Domain-General Cognition than others? A Meta-Analysis of Nonhuman Primate Studies.
Evol. Psychol. **4**, 147470490600400114.

DeCasien, A. R., Williams, S. A. and Higham, J. P. (2017). Primate brain size is
predicted by diet but not sociality. *Nat. Ecol. Evol.* **1**, 0112.

Dobrin, S. E., Herlihy, J. D., Robinson, G. E. and Fahrbach, S. E. (2011). Muscarinic
regulation of Kenyon cell dendritic arborizations in adult worker honey bees. *Arthropod*
Struct. Dev. **40**, 409–419.

Dunbar, R. I. M. (1998). The social brain hypothesis. *Evol. Anthropol. Issues News Rev.*
6, 178–190.

Dunbar, R. I. M. and Shultz, S. (2007). Evolution in the Social Brain. *Science* **317**,
1344–1347.

Ehmer, B. and Gronenberg, W. (2002). Segregation of visual input to the mushroom
bodies in the honeybee (*Apis mellifera*). *J. Comp. Neurol.* **451**, 362–373.

Endler, A., Liebig, J., Schmitt, T., Parker, J. E., Jones, G. R., Schreier, P. and
Hölldobler, B. (2004). Surface hydrocarbons of queen eggs regulate worker
reproduction in a social insect. *Proc. Natl. Acad. Sci.* **101**, 2945–2950.

Evans, L. J. and Raine, N. E. (2014a). Changes in Learning and Foraging Behaviour
within Developing Bumble Bee (*Bombus terrestris*) Colonies. *PLoS ONE* **9**, e90556.

Evans, L. J. and Raine, N. E. (2014b). Foraging errors play a role in resource
exploration by bumble bees (*Bombus terrrestris*). *J. Comp. Physiol. A* **200**, 475–484.

Fahrbach, S. E., Moore, D., Capaldi, E. A., Farris, S. M. and Robinson, G. E. (1998).
Experience-Expectant Plasticity in the Mushroom Bodies of the Honeybee. *Learn. Mem.*
5, 115–123.

Fahrbach, S. E., Farris, S. M., Sullivan, J. P. and Robinson, G. E. (2003). Limits on
volume changes in the mushroom bodies of the honey bee brain. *J. Neurobiol.* **57**, 141–
151.

Fox, J. and Weisberg, S. (2019). An R companion to applied regression. *Sage*.

Freas, C. A., LaDage, L. D., Roth, T. C. and Pravosudov, V. V. (2012). Elevation-
related differences in memory and the hippocampus in mountain chickadees, *Poecile*
gambeli. Anim. Behav. **84**, 121–127.

Frost, R., Armstrong, B. C., Siegelman, N. and Christiansen, M. H. (2015). Domain
generality versus modality specificity: the paradox of statistical learning. *Trends Cogn.*
Sci. **19**, 117–125.

Gawryszewski, F. M. (2018). Color Vision Models: Some simulations, a general n-
dimensional model, and the {colourvision} R package. *Ecol. Evol.* **8**(16), 8159-8170.

Godfrey, R. K., Oberski, J. T., Allmark, T., Givens, C., Hernandez-Rivera, J. and
Gronenberg, W. (2021). Olfactory System Morphology Suggests Colony Size Drives
Trait Evolution in Odorous Ants (Formicidae: Dolichoderinae). *Front. Ecol. Evol.* **9**,
733023.

Gonda, A., Herczeg, G. and Merilä, J. (2011). Population variation in brain size of nine-
spined sticklebacks (*Pungitius pungitius*) - local adaptation or environmentally induced
variation? *BMC Evol. Biol.* **11**, 1-11.

Gonda, A., Herczeg, G. and Merilä, J. (2013). Evolutionary ecology of intraspecific
brain size variation: a review. *Ecol. Evol.* **3**, 2751–2764.

Goulson, D. (2010). *Bumblebees: behaviour, ecology, and conservation.* Oxford
University Press.

Gronenberg, W. (1999). Modality-Specific Segregation of Input to Ant Mushroom
Bodies. *Brain. Behav. Evol.* **54**, 85–95.

Gronenberg, W. and Couvillon, M. J. (2010). Brain composition and olfactory learning
in honey bees. *Neurobiol. Learn. Mem.* **93**, 435–443.

Gronenberg, W. and Liebig, J. (1999). Smaller Brains and Optic Lobes in Reproductive
Workers of the Ant *Harpegnathos. Naturwissenschaften* **86**, 343–345.

Gronenberg, W. and Riveros, A. J. (2009). Social Brains and Behavior—Past and
Present. In *Organization of insect societies: from genome to sociocomplexity*, pp. 377–
401. Harvard University Press.

Hagen, M. and Dupont, Y. L. (2013). Inter-tegular span and head width as estimators of
fresh and dry body mass in bumblebees (*Bombus* spp.). *Insectes Sociaux* **60**, 251–257.

Heinrich, B. (1979). *Bumblebee economics.* Harvard University Press.

Huang, W.-F., Skyrm, K., Ruiter, R. and Solter, L. (2015). Disease management in
commercial bumble bee mass rearing, using production methods, multiplex PCR
detection techniques, and regulatory assessment. *J. Apic. Res.* **54**, 516–524.

Jernigan, C. M., Halby, R., Gerkin, R. C., Sinakevitch, I., Locatelli, F. and Smith, B.
H. (2019). Experience-dependent tuning of early olfactory processing in the adult honey
bee, *Apis mellifera. J. Exp. Biol.* jeb.206748.

Jernigan, C. M., Zaba, N. C. and Sheehan, M. J. (2021). Age and social experience
induced plasticity across brain regions of the paper wasp *Polistes fuscatus. Biol. Lett.*
17, 20210073.

Jones, B. M., Leonard, A. S., Papaj, D. R. and Gronenberg, W. (2013). Plasticity of
the Worker Bumblebee Brain in Relation to Age and Rearing Environment. *Brain. Behav.*
Evol. **82**, 250–261.

Julian, G. E. and Gronenberg, W. (2002). Reduction of Brain Volume Correlates with
Behavioral Changes in Queen Ants. *Brain. Behav. Evol.* **60**, 152–164.

Kenyon, F. C. (1897). The Optic Lobes of the Bee's Brain in the Light of Recent
Neurological Methods. *Am. Nat.* **31**, 369–376.

Kraft, N., Spaethe, J., Rössler, W. and Groh, C. (2019). Neuronal Plasticity in the
Mushroom Body Calyx of Bumble Bee Workers During Early Adult Development. *Dev.*
Neurobiol. **79**, 287–302.

Krofczik, S. (2008). Rapid odor processing in the honeybee antennal lobe network.
Front. Comput. Neurosci. **2**, 1-13.

Kverková, K., Bělíková, T., Olkowicz, S., Pavelková, Z., O'Riain, M. J., Šumbera, R.,
Burda, H., Bennett, N. C. and Němec, P. (2018). Sociality does not drive the evolution
of large brains in eusocial African mole-rats. *Sci. Rep.* **8**, 1-14.

Lenth, R. V. (2017). emmeans: Estimated Marginal Means, aka Least-Squares Means.
1.10.3.

Liang, Z. S., Nguyen, T., Mattila, H. R., Rodriguez-Zas, S. L., Seeley, T. D. and
Robinson, G. E. (2012). Molecular Determinants of Scouting Behavior in Honey Bees.
Science **335**, 1225–1228.

Lihoreau, M., Latty, T. and Chittka, L. (2012). An Exploration of the Social Brain
Hypothesis in Insects. *Front. Physiol.* **3**, 1-7.

Logan, C. J., Avin, S., Boogert, N., Buskell, A., Cross, F. R., Currie, A., Jelbert, S.,
Lukas, D., Mares, R., Navarrete, A. F., et al. (2018). Beyond brain size: Uncovering the
neural correlates of behavioral and cognitive specialization. *Comp. Cogn. Behav. Rev.*
13, 55–89.

Mantini, D., Corbetta, M., Romani, G. L., Orban, G. A. and Vanduffel, W. (2013).
Evolutionarily Novel Functional Networks in the Human Brain? *J. Neurosci.* **33**, 3259–
3275.

Mares, S., Ash, L. and Gronenberg, W. (2005). Brain Allometry in Bumblebee and
Honey Bee Workers. *Brain. Behav. Evol.* **66**, 50–61.

Marino, L. (2006). Absolute brain size: Did we throw the baby out with the bathwater?
Proc. Natl. Acad. Sci. **103**, 13563–13564.

Medler, J. T. (1962). Measurements of the Labium and Radial Cell of *Psithyrus*
(Hymenoptera: Apidae). *Can. Entomol.* **94**, 444–447.

Menzel, R. (2014). The insect mushroom body, an experience-dependent recoding
device. *J. Physiol.-Paris* **108**, 84–95.

Menzel, R. (2021). A short history of studies on intelligence and brain in honeybees.
Apidologie **52**, 23–34.

Menzel, R. and Giurfa, M. (2001). Cognitive architecture of a mini-brain: the honeybee.
Trends Cogn. Sci. **5**, 62–71.

Mobbs, P. (1982). The brain of the honeybee *Apis mellifera.* The connections and
spatial organization of the mushroom bodies. *Philos. Trans. R. Soc. Lond.* **298**, 309–
354.

Murren, C. J., Auld, J. R., Callahan, H., Ghalambor, C. K., Handelsman, C. A.,
Heskel, M. A., Kingsolver, J. G., Maclean, H. J., Masel, J., Maughan, H., et al.
(2015). Constraints on the evolution of phenotypic plasticity: limits and costs of
phenotype and plasticity. *Heredity* **115**, 293–301.

Muth, F. (2021). Intra-specific differences in cognition: bumblebee queens learn better
than workers. *Biol. Lett.* **17**, 20210280.

Muth, F., Cooper, T. R., Bonilla, R. F. and Leonard, A. S. (2018). A novel protocol for
studying bee cognition in the wild. *Methods Ecol. Evol.* **9**, 78–87.

Muth, F., Tripodi, A. D., Bonilla, R., Strange, J. P. and Leonard, A. S. (2021). No sex
differences in learning in wild bumblebees. *Behav. Ecol.* **32**, 638–645.

Niven, J. E. and Laughlin, S. B. (2008). Energy limitation as a selective pressure on
the evolution of sensory systems. *J. Exp. Biol.* **211**, 1792–1804.

O'Donnell, S., Clifford, M. R., Bulova, S. J., DeLeon, S., Papa, C. and Zahedi, N.
(2014). A test of neuroecological predictions using paperwasp caste differences in brain
structure (Hymenoptera: Vespidae). *Behav. Ecol. Sociobiol.* **68**, 529–536.

O'Donnell, S., Bulova, S. J., DeLeon, S., Khodak, P., Miller, S. and Sulger, E. (2015).
Distributed cognition and social brains: reductions in mushroom body investment
accompanied the origins of sociality in wasps (Hymenoptera: Vespidae). *Proc. R. Soc. B*
Biol. Sci. **282**, 20150791.

Ogilvie, J. E. and CaraDonna, P. J. (2022). The shifting importance of abiotic and biotic
factors across the life cycles of wild pollinators. *J. Anim. Ecol.* **91**, 2412–2423.

Ott, S. R. and Rogers, S. M. (2010). Gregarious desert locusts have substantially larger
brains with altered proportions compared with the solitarious phase. *Proc. R. Soc. B Biol.*
Sci. **277**, 3087–3096.

Pahlke, S., Seid, M. A., Jaumann, S. and Smith, A. (2021). The Loss of Sociality Is Accompanied by Reduced Neural Investment in Mushroom Body Volume in the Sweat Bee *Augochlora Pura* (Hymenoptera: Halictidae). *Ann. Entomol. Soc. Am.* **114**, 637–642.

Paulk, A. C. and Gronenberg, W. (2008). Higher order visual input to the mushroom bodies in the bee, *Bombus impatiens*. *Arthropod Struct. Dev.* **37**, 443–458.

Paulk, A. C., Dacks, A. M., Phillips-Portillo, J., Fellous, J.-M. and Gronenberg, W. (2009). Visual Processing in the Central Bee Brain. *J. Neurosci.* **29**, 9987–9999.

Penick, C. A., Ghaninia, M., Haight, K. L., Opachaloemphan, C., Yan, H., Reinberg, D. and Liebig, J. (2021). Reversible plasticity in brain size, behaviour and physiology characterizes caste transitions in a socially flexible ant (*Harpegnathos saltator*). *Proc. R. Soc. B Biol. Sci.* **288**, rspb.2021.0141, 20210141.

Peto, B. R., Costa, C. P., Moore, M. E. and Woodard, S. H. (2025). Social control of egg-laying in independently nest-founding bumble bee queens. *BMC Ecol. Evol.* **25**, 30.

Preibisch, S., Saalfeld, S. and Tomancak, P. (2009). Globally optimal stitching of tiled 3D microscopic image acquisitions. *Bioinformatics* **25**, 1463–1465.

Richards, K. W. (1978). Nest site selection by bumble bees (Hymenoptera: Apidae) in Southern Alberta. *Can. Entomol.* **110**, 301–318.

Riveros, A. J. and Gronenberg, W. (2010). Brain Allometry and Neural Plasticity in the Bumblebee *Bombus occidentalis. Brain. Behav. Evol.* **75**, 138–148.

Riveros, A. J., Seid, M. A. and Wcislo, W. T. (2012). Evolution of brain size in class-based societies of fungus-growing ants (Attini). *Anim. Behav.* **83**, 1043–1049.

Ruttenberg, D. M., Wolf, S. W., Webb, A. E., Wyman, E. S., White, M. L., Melo, D., Traniello, I. M. and Kocher, S. D. (2025). Bumble bee workers adopt novel behavioral roles and reshape their social networks in the absence of a queen. 2025.01.07.630106. *bioRxiv.*

Santos, P. K. F., Galbraith, D. A., Starkey, J. and Amsalem, E. (2022). The effect of the brood and the queen on early gene expression in bumble bee workers' brains. *Sci. Rep.* **12**, 3018.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682.

Serway, C. N., Dunkelberger, B. S., Del Padre, D., Nolan, N. W. C., Georges, S., Freer, S., Andres, A. J. and De Belle, J. S. (2020). Importin-α2 mediates brain development, learning and memory consolidation in *Drosophila. J. Neurogenet.* **34**, 69–82.

Sharma, K. R., Enzmann, B. L., Schmidt, Y., Moore, D., Jones, G. R., Parker, J., Berger, S. L., Reinberg, D., Zwiebel, L. J., Breit, B., et al. (2015). Cuticular

Hydrocarbon Pheromones for Social Behavior and Their Coding in the Ant Antenna. *Cell Rep.* **12**, 1261–1271.

Sherry, D. F. and Hoshooley, J. S. (2010). Seasonal hippocampal plasticity in food-storing birds. *Philos. Trans. R. Soc. B Biol. Sci.* **365**, 933–943.

Shettleworth, S. (2009). *Cognition, evolution, and behavior*. Oxford University Press.

Simões, P. M. V., Niven, J. E. and Ott, S. R. (2013). Phenotypic Transformation Affects Associative Learning in the Desert Locust. *Curr. Biol.* **23**, 2407–2412.

Simões, P. M. V., Ott, S. R. and Niven, J. E. (2016). Environmental Adaptation, Phenotypic Plasticity, and Associative Learning in Insects: The Desert Locust as a Case Study. *Integr. Comp. Biol.* **56**, 914–924.

Skorupski, P. and Chittka, L. (2010). Photoreceptor Spectral Sensitivity in the Bumblebee, *Bombus impatiens* (Hymenoptera: Apidae). *PLOS ONE* **5**, e12049.

Smid, H. M. and Vet, L. E. (2016). The complexity of learning, memory and neural processes in an evolutionary ecological context. *Curr. Opin. Insect Sci.* **15**, 61–69.

Smith, A. R., Seid, M. A., Jiménez, L. C. and Wcislo, W. T. (2010). Socially induced brain development in a facultatively eusocial sweat bee *Megalopta genalis* (Halictidae). *Proc. R. Soc. B Biol. Sci.* **277**, 2157–2163.

Spaethe, J. and Chittka, L. (2003). Interindividual variation of eye optics and single object resolution in bumblebees. *J. Exp. Biol.* **206**, 3447–3453.

Spaethe, J., Brockmann, A., Halbig, C. and Tautz, J. (2007). Size determines antennal sensitivity and behavioral threshold to odors in bumblebee workers. *Naturwissenschaften* **94**, 733–739.

Strausfeld, N. J., Hansen, L., Li, Y., Gomez, R. S. and Ito, K. (1998). Evolution, Discovery, and Interpretations of Arthropod Mushroom Bodies. *Learn. Mem.* **5**, 11–37.

Tanaka, N. K., Endo, K. and Ito, K. (2012). Organization of antennal lobe associated neurons in adult *Drosophila melanogaster* brain. *J. Comp. Neurol.* **520**, 4067–4130.

Taylor, G. J., Tichit, P., Schmidt, M. D., Bodey, A. J., Rau, C. and Baird, E. (2019). Bumblebee visual allometry results in locally improved resolution and globally improved sensitivity. *eLife* **8**,.

Touchon, J. C., McMillan, W. O., Ibáñez, R. and Lessios, H. A. (2024). Flexible oviposition behavior enabled the evolution of terrestrial reproduction. *Proc. Natl. Acad. Sci.* **121**, e2312371121.

Tripodi, A. D. and Strange, J. P. (2019). No second chances for pollen-laden queens? *Insectes Sociaux* **66**, 299–304.

van Doorn, A. and Heringa, J. (1986). The ontogeny of a dominance hierarchy in
colonies of the Bumblebee *Bombus terrestris* (Hymenoptera, Apidae). *Insectes Sociaux*
33, 3–25.

Van Zweden, J. S. and d'Ettorre, P. (2010). Nestmate recognition in social insects and
the role of hydrocarbons. In *Insect Hydrocarbons*, pp. 222–243. Cambridge University
Press.

Warton, D. I., Duursma, R. A., Falster, D. S. and Taskinen, S. (2012). smatr 3 - an R
package for estimation and inference about allometric lines. *Methods Ecol. Evol.* **3**, 257–
259.

Watrobska, C. M., Šima, P., Ramos Rodrigues, A. and Leadbeater, E. (2024).
Potential costs of learning have no detectable impact on reproductive success for
bumble bees. *Anim. Behav.* **214**, 173–185.

Yamhure-Ramírez, D., Wainwright, P. C. and Ramírez, S. R. (2025). Sexual
dimorphism and morphological integration in the orchid bee brain. *Sci. Rep.* **15**, 8915.

Zanette, L. R. S., Miller, S. D. L., Faria, C. M. A., Almond, E. J., Huggins, T. J.,
Jordan, W. C. and Bourke, A. F. G. (2012). Reproductive conflict in bumblebees and
the evolution of worker policing: Reproductive conflict and policing in bumblebees.
Evolution **66**, 3765–3777.