An unusual ring pattern in the Rosβ lanthipeptide of the two-component lantibiotic roseocin

Emily K. Desormeaux[1], Lingyang Zhu,[2] Youran Luo,[1] Dipti Sareen,[3] and Wilfred A. van der Donk[1,2,4,*]

[1] Department of Chemistry and Howard Hughes Medical Institute, 600 South Mathews Avenue, University of Illinois at Urbana−Champaign, Urbana, Illinois 61801, USA.

[2] School of Chemical Sciences NMR Laboratory, University of Illinois at Urbana-Champaign, Urbana, 61801, IL, USA.

[3] Department of Biochemistry, Hargobind Khorana block, Sector-25, South Campus, Panjab University, Chandigarh-160014, India.

[4] Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, 1206 West Gregory Drive, Urbana, Illinois, 61801, USA.

[*] vddonk@illinois.edu

Abstract

Two-component lantibiotics comprise of two post-translationally modified peptides that synergistically exert antimicrobial activity. Most known two-component lantibiotics are made up of a structurally conserved α-lanthipeptide that binds to the peptidoglycan precursor lipid II and a β-lanthipeptide that interacts with the α-peptide-lipid II complex to disrupt the membrane of susceptible bacteria. A few two-component lantibiotics including roseocin produced by Streptomycetes do not follow this general scheme and act by different, currently unresolved mechanisms. An important first step in studying this latter group of lanthipeptides is determination of their chemical structure. Roseocin's β-peptide (Rosβ) is formed by the RosM lanthipeptide synthetase from the RosA1 precursor peptide. RosM carries out nine dehydrations of Ser and Thr residues to the corresponding dehydroamino acids followed by six Michael-type additions of the thiols of Cys residues to a subset of the dehydroamino acids to form six thioether crosslinks. Sequence alignment with other structurally characterized lanthipeptides does not allow prediction of the ring pattern of Rosβ. In this study, Rosβ was produced in *Escherichia coli* and its ring pattern was established by multi-dimensional NMR spectroscopy. The stereochemistry of the lanthionine and methyllanthionine residues was determined by Marfey's analysis with authentic standards. Rosβ is shown to have a unique ring pattern amongst previously characterized lanthipeptides.

Lanthipeptides are a group of peptidic natural products containing thioether crosslinks that are introduced by post-translational modification.[1] Two-component lantibiotics form a subgroup of these natural products that comprise two different post-translationally modified lanthipeptides that display synergistic antimicrobial activity.[2,3] Thus, far, all reported two-component lantibiotics have been members of class II lanthipeptides, for which a single enzyme (generically termed LanM) introduces multiple thioether macrocycles by first dehydrating selective Ser and Thr residues in their precursor peptides by phosphorylation of the alcohol side chains followed by phosphate elimination.[4] This process results in the formation of the dehydroamino acids dehydroalanine (Dha) from Ser and dehydrobutyrine (Dhb) from Thr (Figure 1A). Subsequently, in a different active site, the same enzyme catalyzes the Michael-type addition of Cys residues to the Dha and Dhb residues to form lanthionine (Lan) and methyllanthionine (MeLan) crosslinks, respectively.[5] These modifications take place in the C-terminal part of the precursor peptide called the core peptide; the N-terminal part of the precursor peptide, termed the leader peptide, serves as a recognition element for the post-translational modification enzymes and is removed by a protease in usually the last step of maturation.[6] The biosynthesis of several two-component lantibiotics has been studied demonstrating that the two individual subunits are produced either by a singular LanM enzyme, capable of modifying both peptides,[7-9] or by two distinct LanM enzymes, each modifying one of the two precursor peptides.[10-12]

The mechanism of action of most of the studied two-component lantibiotics involves a1:1 ratio of α- and β-peptides.[13-15] The α-peptides of most of these natural products are structurally highly conserved (Figure 1B) and bind to the peptidoglycan biosynthetic precursor lipid II using a sequence motif containing a highly conserved glutamate residue, analogous to that found in the single-component lanthipeptide mersacidin (Figure 1A).[16-19] The partner β-peptides have less

sequence similarity than their α-counterparts, but still often have conserved ring patterns in characterized peptides (Figure 1C). Whereas the constituent peptides of most reported two-component lantibiotics contain similar ring patterns as those shown in Figure 1, a handful of two-component lantibiotics have been reported that do not have this general architecture. These are cytolysin produced by *Enterococcus faecalis* and the structurally related bibacillins,[7,20] bicereucin produced by *Bacillus cereus* SJ1,[9] and roseocin encoded by *Streptomyces roseosporus* NRRL11379/15998 and *Streptomyces* sp. ADI96-02 (Figure S1).[8,21,22] Roseocin is the first two-component lantibiotic found in the genomes of a member of Actinobacteria and the ring patterns of its two constituents (Rosα and Rosβ) have not yet been determined.

Figure 1. (Me)Lanthionine biosynthesis and structures of select two-component lantibiotics. **A)** Post-translational modification process for class II lanthipeptides that results in Lan and MeLan. **B)** Sequence alignment of the predicted α-peptides of a group of structurally similar two-component lantibiotics. The glutamate residue, absent in RosA2, critical for lipid-II binding and antimicrobial activity in mersacidin-like peptides, is shown in purple. The additional Cys residues in RosA2 that are not conserved in the other peptides are highlighted in red. **C)** Sequence alignment

of the partner β-peptides. The additional Cys residues in RosA1 that are not conserved in the other peptides are highlighted in red. Conserved rings based on the other β-peptides are indicated above the respective alignments, and residues are colored as follows: green, conserved residues involved in ring formation or dehydration; blue, Pro generally conserved in β-peptides. **D)** Alternative manual alignment of the RosA1 core peptide with the CylL$_L$ core peptide. Hal, haloduracin from *Bacillus halodurans*;[11,23] Bht, peptides from *Streptococcus rattus* strain BHT;[24] Enw, enterocin W from *Enterococcus faecalis* NKR-4-1;[25] Lch, lichenicidin from *Bacillus licheniformis* DSM 13 and VK21;[26-28] Ltn, lacticin 3147 from *Lactococcus lactis* DPC3147;[29] Sac, staphylococcin C55 from *Staphylococcus aureus* C55;[30] Smb, peptides from *Bacillus licheniformis* DSM 13.[31]

Figure 2. Biosynthetic gene cluster of roseocin. A) BGC diagram with genes involved in the biosynthesis of roseocin. **B)** Precursor sequences for Rosα and Rosβ. The predicted cleavage sites between leader and core peptides are indicated by a line break. Dehydratable residues (Ser/Thr) in the putative core peptide are indicated in blue and Cys residues in purple.

The roseocin biosynthetic gene cluster (BGC) encodes the two precursor peptides (RosA1 and RosA2), one LanM enzyme (RosM), a transporter (RosT$_P$), likely immunity proteins (RosEFG), and a protein of unknown function (Figure 2).[8] A previous investigation showed that co-expression of RosM with each of the precursor peptides in *E. coli* resulted in modified peptides that displayed strong synergistic antibacterial activity after leader peptide removal.[8] Furthermore, a study on the native producer *S. roseosporus* NRLL15998 detected nine-fold dehydrated Rosβ, consistent with the product made in *E. coli,* but not Rosα.[21] Rosα is generated from the RosA2 precursor peptide, and Rosβ is produced from the RosA1 precursor peptide. Whereas the core peptide of RosA2 has some sequence homology to the precursor peptides of other α-peptides (but lacks the lipid II

binding motif, Figure 1B), the core peptide of the RosA1 precursor is divergent from other β-peptides. The constellation of Cys, Ser, and Thr residues at the N- and C-termini may result in three rings that are at similar positions as other β-peptides, but that would still leave three more Cys residues that are not present in those peptides (Figure 1C). In this study, we determined the ring pattern and stereochemistry of the Rosβ peptide using a combination of NMR spectroscopy, Marfey's analysis, and mass spectrometry (MS). In parallel, we also attempted structural elucidation of the Rosα peptide, but poor solubility in a variety of solvents and poor signal dispersity suggesting aggregation, precluded NMR characterization.

RESULTS

Sequence alignment of roseocin precursor peptides

Multiple sequence alignment of both the α and β core peptides of select two-component lantibiotics was performed (Figure 1). The Rosα peptide aligned well with other α-peptides with the exception of two regions, one located within the first large ring of other α-peptides and another located at the C-terminus of the peptide. These regions of Rosα also contain two additional Cys residues that were identified to be involved in disulfide formation.[8] Thus, Rosα contains four thioether rings as opposed to three in known α-peptides. Most notably, this alignment shows that Rosα is lacking the highly conserved glutamate residue that is found within the lipid II binding domain of mersacidin that is essential for its bioactivity.[18,32,33] In fact, the Rosα core peptide sequence does not contain any Glu or Asp residues, suggesting a potential different molecular target or mechanism of action than that of other α-peptides.

The β-counterparts of two-component lantibiotics associated with the known α-peptides aligned less well with Rosβ, with three of the four conserved rings in other β-peptides potentially also present in Rosβ (Figure 1C). An additional manual alignment was also performed with CylL$_L$" indicating a potentially alternative partially conserved ring pattern (Figure 1D). Cytolysin was the first studied two-component lanthipeptide and is composed of a large subunit, CylL$_L$", and a small subunit, CylL$_S$".[7,34,35] Cytolysin has lytic activity against both bacterial and eukaryotic cell lines, a rare observation for lanthipeptides. Its activity against eukaryotic cells is directly responsible for poor disease prognosis for patients infected with cytolysin-producing *E. faecalis*.[36] The various possible sequence alignments in Figure 1 highlight the need for a fully characterized structure of Rosβ.

Production of Rosβ

We co-expressed a His-tagged version of the RosA1 precursor with untagged RosM in *Escherichia coli* as reported previously.[8] After Ni^{2+}-affinity purification, analysis by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) indicated nine dehydrations (Figure S2). Several peptidases were utilized to attempt leader peptide removal including LahT150,[37] the C39 peptidase domain of the peptidase-containing ATP-binding cassette transporter (PCAT) LahT, and endoproteinase AspN. Leader peptide removal with LahT150 was unsuccessful, potentially because the cleavage site is predicted to be right next to a methyllanthionine ring. To obtain a singular Rosβ product, AspN was utilized to remove a portion of the leader peptide (Figure S2). We then removed the remaining residues of the leader peptide using proteinase K.[8] Using a similar procedure, we also produced Rosα, but its solubility properties were unsuitable for NMR analysis. Therefore, we concentrated on Rosβ for structural elucidation. We refer to a study on Rosα that used a different approach to determine its structure.[38]

Stereochemical analysis of Rosβ

After Ni^{2+}-affinity purification, stereochemical analysis was performed by a modified Marfey's analysis using liquid-chromatography coupled to mass spectrometry detection (LC-MS).[39] This analysis with authentic standards as described previously indicated that the roseocin β-peptide contains DL-Lan, and both LL- and DL-MeLan (Figure 3). While Marfey's analysis of these crosslinked bis-amino acids by LC-MS is not quantitative, the ratios of the peaks within each chromatogram are indicative of the ratios of rings within the peptide.[39] The MeLan analysis indicates LL- and DL-products in an approximately one-to-three ratio, suggesting one LL-ring and three DL-rings. Using this information along with the precursor sequence, it was inferred that Rosβ contains four MeLan rings, and two Lan rings in the RosM-modified peptide.

Figure 3. LCMS analysis of Marfey's assay with RosM-modified RosA1β. A) Extracted ion chromatograms (EICs) of RosM-modified RosA1β (top), Lan standards (middle), and co-injections (bottom) analyzed by LC-MS for $[M − H]^−$ m/z 795.2373. **B)** EICs of RosM-modified RosA1β (top), MeLan standards (middle), and co-injections (bottom) analyzed by LC-MS for $[M − H]^−$ m/z 809.2530. Coloring of each peak: purple, DL stereochemistry; blue, LL stereochemistry; green, D-*allo*-L-MeLan stereochemistry.

NMR analysis of Rosβ

NMR data on Rosβ was obtained in a mixture of acetonitrile and water (1:1) using peptides containing the natural abundance isotopic distributions. All peptide spin systems were assigned using a TOCSY spectrum (Figure 4). Positions of the amino acid assignments within the Rosβ sequence were determined by nuclear Overhauser effect spectroscopy (NOESY) using sequential amide proton assignments, as well as nuclear Overhauser effects (NOEs) between NH residues

and the αH or side chain proton signals of their neighboring residues. These spectra were used collectively to assign all spin systems (Table 1). Amino acid assignments indicated that Thr1, Thr2, Thr7, Thr9, Thr11, Ser16, Thr18, Ser24, and Thr29 were all dehydrated by RosM, while residues Thr13, Thr20, Ser26 escaped dehydration in the RosM-modified Rosβ. These observations were consistent with previously published tandem MS data.[8] The NMR data also indicated that thioether crosslinks were formed at Dhb1, Dhb11, Dha16, Dhb18, Dha24, and Dhb29. We will use the notations Ala16-Cys22 to indicate a lanthionine linkage between the former Ser16 and Cys22; for MeLan we will use the notation Abu1-Cys5 (*vide intra*) to indicate a linkage between aminobutyric acid (Abu) 1 derived from Thr and Cys5.

Two assignments were made for residues eight through eighteen because of two conformations containing either a cis or trans isomer at Pro17, which were present in a ratio of 2:3 (cis:trans). Because the spectra were obtained in 50% acetonitrile, thus likely representing non-natural conformations, no three-dimensional modeling was performed. All reported assignments below correspond to the isomer with the trans conformation at Pro17.

Figure 4. TOCSY spectrum of Rosβ in 50% ACN-d₃/50%H₂O showing the spin system of its component residues. The spectrum was taken with water suppression and a 70 ms relaxation time. Full spectrum without any axis breaks is provided in the SI (Figure S3). An overlay of TOCSY and NOESY spectra is shown in Figure S4.

The NOESY spectrum was used to determine the location of (Me)Lan within Rosβ, revealing six

rings, two Lan and four MeLan. The (Me)Lan structures identified within the peptide will be

termed rings A-F and were shown to be formed between the following residue pairs: Abu1-Cys5

(ring A, Figure S5), Abu11-Cys15 (ring B, Figure S6), Ala16-Cys22 (ring C, Figure S7), Abu18-

Cys25 (ring D, Figure S8), Ala24-Cys32 (ring E, Figure S9), and Abu29-Cys33 (ring F, Figure

S10). These rings were assigned through NOE interactions between the α, β, and γ (in the case of

Abu) protons in the ring forming residues (indicated in red within Figures S5-S10). In some cases,

weak NOE peaks were also observed between these side chain protons and the amide proton of a

residue involved in ring formation; an example of this being the C ring (Figure S7C), wherein

NOE peaks are observed between the former Cys22 amide and the β protons of the former Ser16. The Dhb residues at positions 7 and 9 both had the *Z* stereochemistry, as expected for a LanM catalyzed *anti*-elimination of phosphate (Figure 1A).

The described ring pattern observed in Rosβ (Figure 5A) has not been observed in other lanthipeptides thus far reported, and while the stereochemistry has not been determined for each individual ring, we hypothesize that ring A is the LL-MeLan ring, as it aligns well with previously characterized rings containing LL-stereochemistry, such as CylL$_L$",[7,40] haloduracin β,[40] geobacillin II,[41] and carnolysin.[42] All of these LL rings are formed from a precursor with the sequence DhxDhxXxxXxxCys (Dhx = Dha or Dhb; Xxx = other amino acids), which was shown to favor formation of the LL stereochemistry.[7,40]

Table 1. Chemical shift assignments for Rosβ in 50% ACN-d₃/50%H₂O. Cys and Ser/Thr residues involved in a ring formation are represented in matching colors.

#	AA ID	N-H	αH	βH	γH	δH	other
1	Abu	8.25	4.70	3.55	1.24		
2	Dhb	9.11	N/A	5.87	1.63		
3	Trp	8.18	4.31	3.23, 3.32	2H:7.26 (s), N(1)H:10.00 (br, s), 4H:7.53 (d), 5H:7.03 (t), 6H:7.13 (d), 7H:7.41 (t)		
4	Ala	7.74	4.09	1.37			
5	Cys	7.40	4.83	2.44, 3.39			
6	Ala	7.45	4.10	1.40			
7	Dhb	8.96	N/A	6.66	1.57		
8	Val	7.33	4.03	2.15	0.98		
9	Dhb	9.03	N/A	6.95	1.68		
10	Leu	7.68	4.51	1.59, 1.65	1.53	0.82, 0.90	
11	Abu	8.25	4.43	3.75	1.32		
12	Val	7.78	4.04	1.88	0.89, 0.74		
13	Thr	7.94	4.25	4.19	1.16		
14	Val	7.92	4.04	2.08	0.83		

15	Cys	7.73	4.44	2.78, 3.09			
16	Ala	8.38	4.46	2.80, 3.20			
17	Pro	N/A	4.66	2.30, 1.92	2.01	3.82, 3.64	
18	Abu	7.72	4.63	3.70	1.36		
19	Gly	8.40	3.91				
20	Thr	7.48	4.25	4.18	1.19		
21	Leu	7.73	4.65	1.66, 1.54	1.54	0.86, 0.82	
22	Cys	7.55	4.77	3.13, 2.87			
23	Gly	8.19	3.65 3.85				
24	Ala	8.26	4.90	3.24, 2.83			
25	Cys	7.63	4.63	2.93, 3.01			
26	Ser	9.15	4.08	3.90, 3.99			
27	Met	9.08	3.98	1.72, 2.06	2.35, 2.20	εCH_3: 1.87 (s)	
28	Gly	8.39	3.83				
29	Abu	7.29	4.91	3.51	1.28		
30	Arg	7.28	4.38	1.78, 1.61	1.97, 1.61	3.12	εNH:7.13
31	Gly	7.89	3.37				
32	Cys	8.04	4.26	2.95, 2.88			
33	Cys	7.62	4.47	2.93, 3.01			

Figure 5. Structure of Rosβ. A) Chemical structure of Rosβ. **B)** Shorthand structure of Rosβ. The red MeLan (labeled with an asterisk) most likely has the LL configuration based on sequence similarity to the ring at the N-terminus in other lanthipeptides for which that has been shown conclusively. For explanation of the shorthand notation, see Figure S1. **C)** Lowest energy of an AlphaFold model of unmodified RosA1 core peptide suggesting an alpha helix in the N-terminal region (residues 1-15). Residues involved in ring formation are colored, matching those in Table 1 and panels A and B.

Discussion

Structural characterization of lanthipeptides is imperative for understanding the link between structure and bioactivity. Roseocin, which has antimicrobial activity against several clinically relevant bacterial strains,[8] has novel precursor peptide sequences compared to other two-

component lanthipeptides, suggesting a potential novel structure and mode of action. In this study, Rosβ was structurally characterized using advanced Marfey's analysis and two-dimensional NMR spectroscopy. Attempts to also use NMR spectroscopy to characterize Rosα were unsuccessful. Roseocin's β-peptide has a unique lanthipeptide structure containing six (Me)Lan rings, with the last four rings highly intertwined. The overall ring pattern resembles the large subunit of cytolysin (CylL$_L$") more than the ring patterns of other well-studied two-component lanthipeptides, with similar N-terminal A and B rings and the C-terminal F ring. However, the B ring in CylL$_L$" has the LL-configuration whereas the B-ring in Rosβ has the DL-configuration.

The N-terminal two rings of Rosβ (A and B ring) show similarity in pattern and stereochemistry to both haloduracin β and lichenicidin β (Figure 6).[11,12,27,28] For lichenicidin, the three dimensional structure was determined by NMR spectroscopy in acidic methanol (pH 3.5),[28] showing an alpha helical structure from the start of the A-ring up until a Pro residue that is highly conserved in a number of lanthipeptides including Rosβ.[43] The NMR structure of CylL$_L$" in methanol also showed an α-helical structure for the N-terminus up to a stretch of three Gly residues. While the three-dimensional structure was not modeled for Rosβ, an AlphaFold model[44] of the linear RosA1 precursor core peptide also predicts an alpha helix in the N-terminal region (Figure 5C), which is supported by the observation of (*i*, *i*+3) NOEs in the spectra of Rosβ. This observation, combined with Rosα lacking the Glu residue crucial for bioactivity described for other mersacidin-like peptides, suggests a potential novel mode of action compared to other examined two-component lantibiotics that may not be lipid II dependent. The determination of the structure of Rosα,[38] and Rosβ (this study) sets the stage for mechanistic studies on its mode of action.

Figure 6. Structural similarity of Rosβ with other lanthipeptides. Created with BioRender (https://BioRender.com).

Materials and Methods

General materials and methods

Materials and reagents were purchased from Gold Biotechnology, Fisher Scientific, or Sigma-Aldrich unless otherwise noted. Molecular biology reagents for cloning (Q5 polymerase, and deoxynucleotides) were purchased from NEB. Oligonucleotide primers and genes were obtained from Integrated DNA Technologies and Twist Bioscience. DNA plasmids were prepared using MiniPrep kits (Qiagen) using the manufacturer's instructions. Sanger sequencing was performed by the Roy J. Carver Biotechnology Center (University of Illinois at Urbana-Champaign), ACGT, and SNPsaurus. MALDI-TOF-MS analysis was performed using a Bruker UltrafleXtreme MALDI-TOF mass spectrometer (Bruker Daltonics) in reflectron positive mode at the University of Illinois School of Chemical Sciences Mass Spectrometry Laboratory. All samples were analyzed using Super-DHB MALDI matrix. Multiple sequence alignements were performed using ClustalW.

Plasmid construction and generation

All cloning and mutagenesis experiments were performed using *E. coli* DH5α cells. The RosA1:RosM:pET28a vector to co-express RosA1 and RosM was reported previously.[22] A previous gene construct for the expression of LahT$_{150}$ in a pETDuet-1 vector with a His$_6$-tag was also utilized for this study.[37] Information for plasmids used or generated can be found in Table S1.

Protein and peptide expression

All expressions were performed using *E. coli* BL21 Rosetta2 (DE3) unless otherwise specified. All constructs were overexpressed using TB medium. All cells transformed with the pET28a vector were grown in the presence of 50 µg mL^{-1} kanamycin, and those transformed with pETDuet-1 were grown in the presence of 100 µg mL^{-1} ampicillin.

Starter cultures were made for each expression using TB medium and were supplied with antibiotics indicated above and inoculated with a single colony. Starter cultures were incubated at 37 °C and shaken at 220 rpm overnight. Expression cultures were inoculated with 10 mL of starter culture per L of expression, supplied with antibiotic, and incubated at 37 °C with shaking (200 rpm) until an OD of 0.8-1.0 was reached. The culture was then cooled at 4 °C for 30 min, and IPTG was added to a final concentration of 1 mM to initiate over-expression. The cultures were incubated at 18 °C with shaking (200 rpm). Cells were harvested after 18 h and stored at −70 °C until purification.

Protein and peptide purification

The modified RosA1 peptide was purified by resuspending cell paste in 20 mL of LanA B1 buffer (6.0 M guanidine HCl, 20 mM NaH$_2$PO$_4$, 0.5 mM imidazole, 1 mM TCEP, pH 7.5) for 1 L of cell

culture and lysis via sonication at 60% amplitude for 5 min with a 2.0 s on and 6.0 s off pulse. The lysate was clarified via centrifugation at 50 000×g for 1 h and the supernatant was filtered using 0.45 μm syringe filters. The filtered lysate was loaded onto a gravity flow Ni-NTA column with 1 mL of resin preequilibrated with 6 column volumes (CV) of LanA B1 buffer (6.0 M guanidine HCl, 20 mM NaH_2PO_4, 0.5 mM imidazole, 1 mM TCEP, pH 7.5). The column was washed with 10 CV of LanA B1 buffer, 5 CV of LanA B2 buffer (6.0 M guanidine HCl, 20 mM NaH_2PO_4, 30 mM imidazole, 1 mM TCEP, pH 7.5), and 5 CV of wash buffer (20 mM NaH_2PO_4, 30 mM imidazole, 300 mM NaCl, 1 mM TCEP, pH 7.5). The peptide was then eluted using 10 CV of elution buffer (20 mM NaH_2PO_4, 1 M imidazole, 100 mM NaCl, 1 mM TCEP, pH 7.5). After purification of RosA1 that was modified by RosM *in vivo*, endoproteinase AspN was used to remove a large portion of the leader peptide. The buffer of the purified peptide was first exchanged into AspN digestion buffer (50 mM Tris HCl pH 8.0, and 2.5 mM $ZnSO_4$; 2X reaction buffer provided by NEB) using a 3 kDa Amicon centrifugation filter and the sample was concentrated to approximately 5 mL. Digestion reactions contained a mass-to-mass ratio of 50:1 of peptide: AspN. Reactions were incubated for 2 h at 37 °C and AspN-digested peptides were purified via reversed phase HPLC using a Phenomenex Luna C5 semiprep column (250 × 10 mm, 10 mm, 100 Å) at a flow rate of 8 mL min^{-1} and with the following gradient over 32 min: 2% B for 10 min, 2– 30% B over 2 min, 30–60% B over 15 min, 60–100% B over 2 min, and hold at 100% B for 3 min (A: 0.1% TFA in H_2O, B: 100% MeCN, 0.1% TFA). HPLC-purified peptide was collected and lyophilized, and the peptides were stored as a powder at −20 °C until use. Then additional residues of the predicted leader peptide were removed utilizing proteinase K as previously described to obtain Rosβ peptide.[8]

LahT150 was expressed and purified as previously reported. Its activity was confirmed with ProcA2.8 as substrate.[37]

Structural characterization

Stereochemical characterization of the roseocin peptides was performed as previously described.[39] For NMR studies, the Rosβ peptide was dissolved in 530-560 µL of 50% acetonitile-d$_3$ in H$_2$O and transferred to a 5 mm thin wall precision NMR sample tube (7" L, 535-PP, purchased from Wilmad-LabGlass). An ^{1}H double pulsed field gradient spin-echo (DPFGSE) TOCSY with a 70 ms mixing time and DPFGSE NOESY with a 350 ms mixing time were acquired at 25 °C. The data were acquired on an Agilent VNMRS 750 MHz spectrometer using VNMRJ 4.2A software with the BioPack suite. The acetonitrile signal was referenced as 1.97 ppm prior to manual data analysis.

Determination of stereochemistry

Stereochemistry determination was adapted from the previously reported advanced Marfey's method.[39] The peptide sample was hydrolyzed in 6 M DCl for 20 h at 120 ℃. The resulting hydrolysate was derivatized with 1-fluoro-2,4-dinitrophenyl-5-L-leucinamide (L-FDLA) in 0.8 M aqueous NaHCO$_3$ for 3 h at 67 ℃. Following the reaction, the mixture was neutralized with 6 M HCl and lyophilized. The dried residue was then extracted with MeCN to remove residual inorganic salts. The extract containing the diastereomeric derivatives was analyzed by LC-MS using an Agilent 6545 LC/Q-TOF instrument with a Kinetex F5 Core−Shell HPLC column (1.7 µm, F5 100 Å, 100 mm× 2.1 mm). mCylL$_s$ (plasmid available at Addgene ID #208760) and mCoiA1 (Addgene ID #208761 and #208762) were used for lanthionine and methyllanthionine standards, respectively, for comparison and co-injection.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at https://pubs.acs.org/doi/
Figures S1-S10 showing spectroscopic data, and Table S1 with gene sequences.

AUTHOR INFORMATION

Corresponding Author

Wilfred A. van der Donk −Department of Chemistry and Howard Hughes Medical Institute, University of Illinois at Urbana-Champaign, Urbana, IL, United States; orcid.org/0000-0002-5467-7071;
Email: vddonk@illinois.edu.

Authors

Emily K. Desormeaux –Department of Chemistry, University of Illinois at Urbana-Champaign, Urbana, IL, United States. orcid: 0000-0001-7400-8046

Lingyang Zhu – School of Chemical Sciences NMR Laboratory, University of Illinois at Urbana-Champaign, Urbana, IL, United States; orcid: 0000-0002-6657-271X.

Youran Luo –Department of Chemistry, University of Illinois at Urbana-Champaign, Urbana, IL, United States. orcid: 0000-0002-8033-2472

Dipti Sareen - Department of Biochemistry, Hargobind Khorana block, Sector-25, South Campus, Panjab University, Chandigarh-160014, India. orcid: 0000-0001-9432-0870.

Funding

This work was supported by the Howard Hughes Medical Institute. A Bruker UltrafleXtreme mass spectrometer used was purchased with support from the National Institutes of Health (S10 RR027109). The grant received from ANRF-SERB, India (Grant Number: CRG/2023/003532-G) is duly acknowledged.

Notes
The authors declare no competing financial interests.

References

(1) Repka, L. M.; Chekan, J. R.; Nair, S. K.; van der Donk, W. A. Mechanistic understanding of lanthipeptide biosynthetic enzymes. *Chem. Rev.* **2017**, *117*, 5457-520.

(2) Garneau, S.; Martin, N. I.; Vederas, J. C. Two-peptide bacteriocins produced by lactic acid bacteria. *Biochimie* **2002**, *84*, 577-92.

(3) Lawton, E. M.; Ross, R. P.; Hill, C.; Cotter, P. D. Two-peptide lantibiotics: a medical perspective. *Mini Rev. Med. Chem.* **2007**, *7*, 1236-47.

(4) Chatterjee, C.; Miller, L. M.; Leung, Y. L.; Xie, L.; Yi, M.; Kelleher, N. L.; van der Donk, W. A. Lacticin 481 synthetase phosphorylates its substrate during lantibiotic production. *J. Am. Chem. Soc.* **2005**, *127*, 15332-3.

(5) Paul, M.; Patton, G. C.; van der Donk, W. A. Mutants of the zinc ligands of lacticin 481 synthetase retain dehydration activity but have impaired cyclization activity. *Biochemistry* **2007**, *46*, 6268-76.

(6) Eslami, S. M.; van der Donk, W. A. Proteases involved in leader peptide removal during RiPP biosynthesis. *ACS Bio. Med. Chem. Au* **2023**, *4*, 20-36.

(7) Tang, W.; van der Donk, W. A. The sequence of the enterococcal cytolysin imparts unusual lanthionine stereochemistry. *Nat. Chem. Biol.* **2013**, *9*, 157-9.

(8) Singh, M.; Chaudhary, S.; Sareen, D. Roseocin, a novel two-component lantibiotic from an actinomycete. *Mol. Microbiol.* **2020**, *113*, 326-37.

(9) Huo, L.; van der Donk, W. A. Discovery and characterization of bicereucin, an unusual D-amino acid-containing mixed two-component lantibiotic. *J. Am. Chem. Soc.* **2016**, *138*, 5254-7.

(10) McAuliffe, O.; Hill, C.; Ross, R. P. Each peptide of the two-component lantibiotic lacticin 3147 requires a separate modification enzyme for activity. *Microbiology* **2000**, *146*, 2147-54.

(11) McClerren, A. L.; Cooper, L. E.; Quan, C.; Thomas, P. M.; Kelleher, N. L.; van der Donk, W. A. Discovery and in vitro biosynthesis of haloduracin, a two-component lantibiotic. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 17243-8.

(12) Caetano, T.; Krawczyk, J. M.; Mosker, E.; Süssmuth, R. D.; Mendo, S. Heterologous expression, biosynthesis, and mutagenesis of type II lantibiotics from *Bacillus licheniformis* in *Escherichia coli*. *Chem. Biol.* **2011**, *18*, 90-100.

(13) Wiedemann, I.; Bottiger, T.; Bonelli, R. R.; Wiese, A.; Hagge, S. O.; Gutsmann, T.; Seydel, U.; Deegan, L.; Hill, C.; Ross, P.; Sahl, H. G. The mode of action of the lantibiotic lacticin 3147--a complex mechanism involving specific interaction of two peptides and the cell wall precursor lipid II. *Mol. Microbiol.* **2006**, *61*, 285-96.

(14) Oman, T. J.; van der Donk, W. A. Insights into the mode of action of the two-peptide lantibiotic haloduracin. *ACS Chem. Biol.* **2009**, *4*, 865-74.

(15) Mukherjee, S.; Huo, L.; Thibodeaux, G. N.; van der Donk, W. A. Synthesis and bioactivity of diastereomers of the virulence lanthipeptide cytolysin. *Org. Lett.* **2016**, *18*, 6188-91.

(16) Panina, I. S.; Balandin, S. V.; Tsarev, A. V.; Chugunov, A. O.; Tagaev, A. A.; Finkina, E. I.; Antoshina, D. V.; Sheremeteva, E. V.; Paramonov, A. S.; Rickmeyer, J.; Bierbaum, G.; Efremov, R. G.; Shenkarev, Z. O.; Ovchinnikova, T. V. Specific binding of the α-component of the lantibiotic lichenicidin to the peptidoglycan precursor lipid II predetermines its antimicrobial activity. *Int. J. Mol. Sci.* **2023**, *24*, 1332.

(17) Brötz, H.; Bierbaum, G.; Leopold, K.; Reynolds, P. E.; Sahl, H. G. The lantibiotic mersacidin inhibits peptidoglycan synthesis by targeting lipid II. *Antimicrob. Agents Chemother.* **1998**, *42*, 154-60.

(18) Szekat, C.; Jack, R. W.; Skutlarek, D.; Farber, H.; Bierbaum, G. Construction of an expression system for site-directed mutagenesis of the lantibiotic mersacidin. *Appl. Environ. Microbiol.* **2003**, *69*, 3777-83.

(19) Oman, T. J.; Lupoli, T. J.; Wang, T.-S. A.; Kahne, D.; Walker, S.; van der Donk, W. A. Haloduracin α binds the peptidoglycan precursor lipid II with 2:1 stoichiometry. *J. Am. Chem. Soc.* **2011**, *133*, 17544-7.

(20) Moreira, R.; Yang, Y.; Luo, Y.; Gilmore, M. S.; van der Donk, W. A. Bibacillin 1: a two-component lantibiotic from *Bacillus thuringiensis*. *RSC Chem. Biol.* **2024**.

(21) Kersten, R. D.; Yang, Y. L.; Xu, Y.; Cimermancic, P.; Nam, S. J.; Fenical, W.; Fischbach, M. A.; Moore, B. S.; Dorrestein, P. C. A mass spectrometry-guided genome mining approach for natural product peptidogenomics. *Nat. Chem. Biol.* **2011**, *7*, 794-802.

(22) Ayikpoe, R. S.; Shi, C.; Battiste, A. J.; Eslami, S. M.; Ramesh, S.; Simon, M. A.; Bothwell, I. R.; Lee, H.; Rice, A. J.; Ren, H.; Tian, Q.; Harris, L. A.; Sarksian, R.; Zhu, L.; Frerk, A. M.; Precord, T. W.; van der Donk, W. A.; Mitchell, D. A.; Zhao, H. A scalable platform to discover antimicrobials of ribosomal origin. *Nat. Commun.* **2022**, *13*, 6135.

(23) Lawton, E. M.; Cotter, P. D.; Hill, C.; Ross, R. P. Identification of a novel two-peptide lantibiotic, Haloduracin, produced by the alkaliphile *Bacillus halodurans* C-125. *FEMS Microbiol. Lett.* **2007**, *267*, 64-71.

(24) Hyink, O.; Balakrishnan, M.; Tagg, J. R. Streptococcus rattus strain BHT produces both a class I two-component lantibiotic and a class II bacteriocin. *FEMS Microbiol. Lett.* **2005**, *252*, 235-41.

(25) Sawa, N.; Wilaipun, P.; Kinoshita, S.; Zendo, T.; Leelawatcharamas, V.; Nakayama, J.; Sonomoto, K. Isolation and characterization of enterocin W, a novel two-peptide lantibiotic produced by *Enterococcus faecalis* NKR-4-1. *Appl. Environ. Microbiol.* **2012**, *78*, 900-3.

(26) Begley, M.; Cotter, P. D.; Hill, C.; Ross, R. P. Identification of a novel two-peptide lantibiotic, lichenicidin, following rational genome mining for LanM proteins. *Appl. Environ. Microbiol.* **2009**, *75*, 5451-60.

(27) Dischinger, J.; Josten, M.; Szekat, C.; Sahl, H. G.; Bierbaum, G. Production of the novel two-peptide lantibiotic lichenicidin by *Bacillus licheniformis* DSM 13. *PLoS One* **2009**, *4*, e6788.

(28) Shenkarev, Z. O.; Finkina, E. I.; Nurmukhamedova, E. K.; Balandin, S. V.; Mineev, K. S.; Nadezhdin, K. D.; Yakimenko, Z. A.; Tagaev, A. A.; Temirov, Y. V.; Arseniev, A. S.; Ovchinnikova, T. V. Isolation, structure elucidation, and synergistic antibacterial activity of a novel two-component lantibiotic lichenicidin from *Bacillus licheniformis* VK21. *Biochemistry* **2010**, *49*, 6462-72.

(29) Ryan, M. P.; Jack, R. W.; Josten, M.; Sahl, H. G.; Jung, G.; Ross, R. P.; Hill, C. Extensive post-translational modification, including serine to D-alanine conversion, in the two-component lantibiotic, lacticin 3147. *J. Biol. Chem.* **1999**, *274*, 37544-50.

(30) Navaratna, M. A.; Sahl, H. G.; Tagg, J. R. Identification of genes encoding two-component lantibiotic production in *Staphylococcus aureus* C55 and other phage group II *S. aureus* strains and demonstration of an association with the exfoliative toxin B gene. *Infect. Immun.* **1999**, *67*, 4268-71.

(31) Yonezawa, H.; Kuramitsu, H. K. Genetic analysis of a unique bacteriocin, Smb, produced by Streptococcus mutans GS5. *Antimicrob. Agents Chemother.* **2005**, *49*, 541-8.

(32) Deegan, L. H.; Suda, S.; Lawton, E. M.; Draper, L. A.; Hugenholtz, F.; Peschel, A.; Hill, C.; Cotter, P. D.; Ross, R. P. Manipulation of charged residues within the two-peptide lantibiotic lacticin 3147. *Microb. Biotechnol.* **2010**, *3*, 222-34.

(33) Cooper, L. E.; McClerren, A. L.; Chary, A.; van der Donk, W. A. Structure-activity relationship studies of the two-component lantibiotic haloduracin. *Chem. Biol.* **2008**, *15*, 1035-45.

(34) Booth, M. C.; Bogie, C. P.; Sahl, H. G.; Siezen, R. J.; Hatter, K. L.; Gilmore, M. S. Structural analysis and proteolytic activation of *Enterococcus faecalis* cytolysin, a novel lantibiotic. *Mol. Microbiol.* **1996**, *21*, 1175-84.

(35) Cox, C. R.; Coburn, P. S.; Gilmore, M. S. Enterococcal cytolysin: a novel two component peptide system that serves as a bacterial defense against eukaryotic and prokaryotic cells. *Curr. Protein Pept. Sci.* **2005**, *6*, 77-84.

(36) Duan, Y.; Llorente, C.; Lang, S.; Brandl, K.; Chu, H.; Jiang, L.; White, R. C.; Clarke, T. H.; Nguyen, K.; Torralba, M.; Shao, Y.; Liu, J.; Hernandez-Morales, A.; Lessor, L.; Rahman, I. R.; Miyamoto, Y.; Ly, M.; Gao, B.; Sun, W.; Kiesel, R.; Hutmacher, F.; Lee, S.; Ventura-Cots, M.; Bosques-Padilla, F.; Verna, E. C.; Abraldes, J. G.; Brown, R. S., Jr.; Vargas, V.; Altamirano, J.; Caballeria, J.; Shawcross, D. L.; Ho, S. B.; Louvet, A.; Lucey, M. R.; Mathurin, P.; Garcia-Tsao, G.; Bataller, R.; Tu, X. M.; Eckmann, L.; van der Donk, W. A.; Young, R.; Lawley, T. D.; Starkel, P.; Pride, D.; Fouts, D. E.; Schnabl, B. Bacteriophage targeting of gut bacterium attenuates alcoholic liver disease. *Nature* **2019**, *575*, 505-11.

(37) Bobeica, S. C.; Dong, S. H.; Huo, L.; Mazo, N.; McLaughlin, M. I.; Jimenéz-Osés, G.; Nair, S. K.; van der Donk, W. A. Insights into AMS/PCAT transporters from biochemical and structural characterization of a double glycine motif protease. *eLife* **2019**, *8*, e42305.

(38) Kishen, S.; Dhiman, G.; Luo, Y.; Martini, R. M.; Manasa, K. M.; Samson, A. S.; van der Donk, W. A.; Sareen, D. Structure-activity relationship of thioether cross-links in roseocin's alpha component. **2025**, manuscript under review.

(39) Luo, Y.; Xu, S.; Frerk, A. M.; van der Donk, W. A. Facile method for determining lanthipeptide stereochemistry. *Anal. Chem.* **2024**, *96*, 1767-73.

(40) Tang, W.; Jiménez-Osés, G.; Houk, K. N.; van der Donk, W. A. Substrate control in stereoselective lanthionine biosynthesis. *Nat. Chem.* **2015**, *7*, 57-64.

(41) Garg, N.; Goto, Y.; Chen, T.; van der Donk, W. A. Characterization of the stereochemical configuration of lanthionines formed by the lanthipeptide synthetase GeoM. *Biopolymers* **2016**, *106*, 834–42.

(42) Lohans, C. T.; Li, J. L.; Vederas, J. C. Structure and biosynthesis of carnolysin, a homologue of enterococcal cytolysin with D-amino acids. *J. Am. Chem. Soc.* **2014**, *136*, 13150-3.

(43) Zhao, X.; van der Donk, W. A. Structural characterization and bioactivity analysis of the two-component lantibiotic Flv system from a ruminant bacterium. *Cell Chem. Biol.* **2016**, *23*, 246-56.

(44) Jumper, J.; Evans, R.; Pritzel, A.; Green, T.; Figurnov, M.; Ronneberger, O.; Tunyasuvunakool, K.; Bates, R.; Žídek, A.; Potapenko, A.; Bridgland, A.; Meyer, C.; Kohl, S. A. A.; Ballard, A. J.; Cowie, A.; Romera-Paredes, B.; Nikolov, S.; Jain, R.; Adler, J.; Back, T.; Petersen, S.; Reiman, D.; Clancy, E.; Zielinski, M.; Steinegger, M.; Pacholska, M.; Berghammer, T.; Bodenstein, S.; Silver, D.; Vinyals, O.; Senior, A. W.; Kavukcuoglu, K.; Kohli, P.; Hassabis, D. Highly accurate protein structure prediction with AlphaFold. *Nature* **2021**, *596*, 583-9.

TOC graphic

Rosβ